The effects of caloric restriction on adipose tissue and metabolic health are sex- and age-dependent

Karla J. Suchacki[1#] & Ben J. Thomas[1#], Claire Fyfe[3], Adriana A.S. Tavares[1,2], Richard J. Sulston[1], Andrea Lovdel[1], Holly J. Woodward[1], Xuan Han[1], Domenico Mattiucci[1], Eleanor J. Brain[1], Carlos J. Alcaide-Corral[1,2], Gillian A. Gray[1], Phillip D. Whitfield[4,5], Roland H. Stimson[1], Nicholas M. Morton[1], Alexandra M. Johnstone[3] and William P. Cawthorn[1*].

[1]University/BHF Centre for Cardiovascular Science and [2]Edinburgh Imaging, University of Edinburgh, The Queen's Medical Research Institute, Edinburgh BioQuarter, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK. [3]Rowett Institute, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD, UK. [4]Division of Biomedical Sciences, University of the Highlands and Islands, Centre for Health Sciences, Inverness, UK. [5]Institute of Infection, Immunity & Inflammation, University of Glasgow, Glasgow, G12 8TA, UK.

[#]These authors contributed equally to this manuscript.

__Correspondence to__: William Cawthorn, University/BHF Centre for Cardiovascular Science, The Queen's Medical Research Institute, Edinburgh BioQuarter, 47 Little France Crescent, Edinburgh, EH16 4TJ. W.Cawthorn@ed.ac.uk

Keywords: Sex differences, caloric restriction, ageing, adipose tissue, white adipose tissue, glucose homeostasis, insulin sensitivity, PET/CT, lipidomics.

HIGHLIGHTS

- Caloric restriction (CR) decreases fat mass and improves glucose homeostasis in young male mice, but young females resist these effects.
- These effects are associated with improved hepatic insulin sensitivity and suppressed gluconeogenesis, without marked differences in peripheral glucose uptake.
- CR's effects on fat loss and glucose homeostasis are comparable in aged male and female mice, implicating oestrogen as the driver of the sexually dimorphic effects in young mice.
- In humans, females resist CR-induced fat loss in an age-dependent manner, further supporting the role of oestrogen in the sexually dimorphic effects of CR.

SUMMARY

Caloric restriction (CR) is a nutritional intervention that reduces the risk of age-related diseases in numerous species, including humans. CR's metabolic effects, including decreased fat mass and improved insulin sensitivity, play an important role in its broader health benefits. However, the extent and basis of sex differences in CR's health benefits are unknown. We found that 30% CR in young (3-month-old) male mice decreased fat mass and improved glucose tolerance and insulin sensitivity, whereas these effects were blunted or absent in young female mice. Resistance to fat and weight loss in females was not explained by decreased energy expenditure compared to males. Positron emission tomography-computed tomography (PET/CT) with ^{18}F-fluorodeoxyglucose (^{18}F-FDG) showed that peripheral glucose uptake was comparable between sexes. Instead, CR caused a pronounced suppression of hepatic gluconeogenesis and altered liver ceramide content in males but not in females. In aged mice (18-months old), when females are anoestrus, CR decreased fat mass and improved glucose homeostasis to a similar extent in both sexes. CR-induced fat loss in humans was also sex- and age-dependent: younger females (<45 years) resisted fat loss compared to younger males while in older subjects (>45 years) this sex difference was absent. Collectively, these studies identify age-dependent sex differences in the metabolic effects of CR and highlight adipose tissue, the liver and oestrogen as key determinants of CR's metabolic benefits. This has important implications for understanding the interplay between diet and health and for maximising the benefits of CR in humans.

INTRODUCTION

Caloric restriction (CR) is a therapeutic nutritional intervention involving a sustained decrease in calorie intake whilst maintaining adequate nutrition. CR extends lifespan and reduces the risk of age-related diseases in numerous species, ranging from yeast to primates (1-3). Trials of CR in humans show numerous health benefits, including the prevention of cardiovascular disease, hypertension, obesity, type 2 diabetes, chronic inflammation, and risk of certain cancers (4). Thus, the ability of CR to promote healthy ageing is now also recognised in humans. In addition to these health benefits, many effects of CR reflect evolutionary adaptations that confer a survival advantage during periods of food scarcity (5). Establishing the extent and basis of CR's effects may thereby reveal new knowledge of healthy ageing and the interplay between food, nutrition, and health.

A key contributor to the health benefits of CR is its impact on metabolic function. Ageing is characterised by hepatic insulin resistance, hyperinsulinaemia and excessive accumulation of white adipose tissue (WAT), particularly visceral WAT (6-8). The latter is coupled with adipose dysfunction, whereby WAT becomes unable to meet the demands for safe lipid storage. This results in ectopic lipid accumulation in the liver and other tissues, contributing to insulin resistance and metabolic dysregulation (6, 8). CR counteracts these effects, decreasing WAT mass, increasing fatty acid oxidation, preventing ectopic lipid deposition and enhancing insulin sensitivity and glucose tolerance (5). Notably, studies in rodents show that removal of visceral WAT, independent of CR, is sufficient to prevent insulin resistance, improve glucose tolerance and increase lifespan (9-11). Visceral WAT loss in humans is also strongly associated with CR's metabolic benefits (12). Thus, the ability of CR to decrease visceral WAT, and the resultant improvements in hepatic insulin sensitivity, are likely central to CR's effects on metabolic function and healthy ageing.

This importance of adiposity and metabolic function raises the possibility of sex differences in the CR response. Indeed, it is now well established that metabolic homeostasis and adipose biology differ substantially between males and females (13-16); however, most preclinical rodent studies have used males only (17), suggesting that much of the CR literature may have overlooked sex as a potential determinant of CR's effects. Nevertheless, some clinical and preclinical CR studies from our lab and others have identified sexually dimorphic responses (18-21), whereby males lose fat mass to a greater extent than females. Oestrogens underlie many metabolic sex differences (13, 22), suggesting that oestrogen may contribute to females' resistance to fat loss during CR. However, the extent and basis of sexual dimorphism in CR's metabolic effects remains to be firmly established.

Herein, we first systematically review the CR literature to establish the degree to which sex has been overlooked a determinant of the CR response. To further elucidate the extent and basis of sex differences, we studied CR in male and female mice and humans at ages where oestrogen is physiologically active or absent. Together, our findings show that the CR field has largely overlooked sex differences and reveal that both mice and humans display age-dependent sexual dimorphism in the metabolic effects of CR.

RESULTS

Sex is routinely overlooked as a biological variable in the CR literature.

One review of the recent CR literature found that most rodent studies use males only, with females typically used only to address female-specific experimental questions (18); however, the extent of this bias in earlier CR studies, and whether it applies to human CR research, has not been assessed. To address these issues, we first systematically quantified the use of males and females, and the consideration of sex as a biological variable, in mouse and human CR studies (Fig. 1, S1). We focussed on research published since 2003, when the European Commission first highlighted the importance of addressing sex as a biological variable (23). We also excluded studies that necessarily focussed on a single sex, such as those addressing effects of CR on female reproductive function, leaving only studies in which there is no scientific rationale for ignoring potential sexual dimorphism. We found that male-only studies predominated for mice, accounting for ~64% of all mouse CR studies since 2003 (Fig. 1A, S1B). This is consistent with previous analyses of the more-recent CR literature (18). Fewer than 20% of studies used females only, while around 7.3% combined males and females, suggesting an assumption that sex would not influence experimental outcomes. In contrast, ~60% of human studies combined males and females, while ~23% used only females and ~12% only males (Fig. 1C, S1B). Strikingly, by the end of 2021, studies that included both sexes and analysed data with sex as a variable were in a minority for both mice (~3.4%) and humans (~4.5%) (Fig. 1A, 1C). Moreover, the proportion of studies using each sex or combination of sexes has remained relatively constant since 2003 (Fig. 1A, 1C). Thus, efforts to increase the consideration of sex in experimental design, as promoted by the European Commission (23), Canadian Institutes of Health Research (CIHR) and National Institutes of Health (NIH) (24), appear to have had little impact on the field of CR research.

We next focussed on the minority of studies that did consider sex in their experiments ("M&F separate") and that included analysis of metabolic parameters. We found that results in ~74% of mouse studies (Fig. 1B) and ~62% of human studies (Fig. 1D) indicated a sex difference in the CR response. Thus, sex differences have been described in the CR literature, but the continuing dearth of studies that include both sexes suggests that this issue continues to be overlooked in the CR field.

Females resist CR-induced weight loss, fat loss, and adipocyte hypertrophy

To explore sex differences in the CR response, we assessed the effects of 30% CR, implemented from 9-15 weeks of age, in C57BL/6J and C57BL/6N mice. As expected, CR decreased body mass in both males and females (Fig. 2A) but this effect was greater in males, with ANOVA confirming a significant sex-diet interaction. This sex difference in response to CR is particularly clear when body mass is presented relative to baseline, pre-CR levels (Fig. 2B).

To determine how fat and lean mass contribute to these diet and sex effects, body composition was assessed weekly using time-domain nuclear magnetic resonance (TD-NMR; Fig. 2C-E, S2A-B). CR decreased fat and lean mass in males, whereas females maintained fat mass and lost only lean mass (Fig. 2C-E). A significant sex-diet interaction was apparent for absolute fat mass (Fig. 2C) and for fat mass relative

to baseline (Fig. 2D); the latter showed that CR decreased fat mass in males, whereas
AL males and AL or CR females increased fat mass to a similar extent over the six-
week duration (Fig. 2D). For lean mass, a significant sex-diet interaction occurred for
absolute mass, with losses being greater in males than in females (Fig. 2E). However,
when compared to baseline lean mass, the CR vs AL effect was similar between the
sexes, in part because AL females continued to increase lean mass over time (Fig.
2F).

Given that these body composition changes coincide with changes in overall body
mass, we also assessed fat and lean mass as % body mass. Males preferentially lost
fat mass and preserved lean mass in response to CR, with % lean mass being greater
in CR vs AL-fed males (Fig. S2). In contrast, diet did not alter % fat or % lean mass in
females, indicating that changes in body composition were proportionate to the overall
changes in body mass (Fig. S2).

After 6 weeks of AL or CR diet, tissue-specific changes in response to CR were
assessed (Fig. 2G; Fig. S3). We found that in males but not females CR decreased
the absolute mass of gonadal (gWAT), inguinal (iWAT), mesenteric (mWAT) and
perirenal (pWAT) WAT depots, as well as brown adipose tissue (BAT; Fig. S3A). CR
also tended to decrease the absolute mass of pericardial WAT (pcWAT) in males only
(Fig. S3A). To determine if these changes were proportionate to changes in overall
body mass, we further analysed the mass of each adipose depot as % body mass
(Fig. 2G). This showed that the significant sex-diet interaction persisted for gWAT,
iWAT, mWAT and pWAT. CR did not affect % BAT mass in either sex, though % BAT
mass was significantly greater in males than in females (Fig. 2G). Thus, both in
absolute terms and as % body mass, CR decreases WAT mass in males but not in
females.

We next analysed the masses of other tissues to determine if this sex-dependent effect
of CR was unique to WAT. CR significantly decreased the absolute mass of the liver,
pancreas, kidneys, gastrocnemius muscle (gastroc), heart, spleen and thymus (Fig
S3B). Significant sex-diet effects were detected for the liver and kidney, with CR
causing greater decreases in males than in females; however, for each of the other
tissues the CR effect was similar between the sexes (Fig. S3B). In contrast to these
effects on absolute mass, CR did not significantly affect the relative masses of each
of these tissues, nor did sex influence the CR effect (Fig. S3C). This indicates that the
absolute masses of these tissues decreased in proportion to the changes in overall
body mass. One notable exception is the adrenal glands, the mass of which was
increased with CR to a greater extent in males than in females (Fig. S3B-C). Together,
these data show that the sex differences in CR-induced loss of body mass and fat
mass were driven primarily by decreased WAT mass in males, which females robustly
resist.

Differences in WAT mass can be driven by changes in adipocyte size and/or adipocyte
number. Thus, given the marked sex differences in the effect of CR on WAT mass, we
next investigated this effect at the level of adipocyte size (Fig. 2H-I). In gWAT, CR
significantly decreased average adipocyte area in males but not in females (Fig. 2H&I).
The same relationships were observed in mWAT, another visceral adipose depot (Fig.
S4). These observations suggest that differences in lipolysis and/or lipogenesis
underpin the sex differences in CR's effects on WAT mass.

208
209
Females resist CR-induced improvements in glucose homeostasis
We next investigated if other metabolic effects of CR also differ between the sexes. We found that CR decreased blood glucose to a greater extent in males than in females (Fig. 3A). Consistent with this, oral glucose tolerance tests (OGTT) revealed that CR improved glucose tolerance in both sexes but this effect was greater in males (Fig. 3B). These diet and sex effects were reflected by the OGTT total area under the curve (tAUC), calculated against 0 mM blood glucose; however, they were not apparent for the incremental AUC (iAUC), calculated against fasting blood glucose for each mouse. This suggests that differences in fasting glucose are the main driver of CR-induced improvements in glucose tolerance and the sex differences therein.

To further assess the basis for these effects on glucose tolerance, we analysed plasma insulin during the OGTT. CR decreased plasma insulin in males but not in females, largely because AL females were already relatively hypoinsulinaemic (Fig. 3D). Based on these blood glucose and insulin concentrations we calculated the homeostasis model assessment for insulin resistance (HOMA-IR) (25) and the Matsuda Index of insulin sensitivity (26). Across both sexes CR significantly decreased HOMA-IR and increased the Matsuda Index; however, within each sex the CR effect was significant for males only, indicating that females resist the CR-induced improvements in insulin sensitivity (Fig. 3E).

Time of CR feeding does not alter the sex differences in CR's metabolic effects
Time of feeding can influence the metabolic effects of CR (5, 27). Therefore, to test if our feeding regime influenced the observed sex differences we studied a cohort of mice that received their daily ration of CR diet in the evening instead of the morning (Fig. S5-6). Consistently, we found that the sex-dependent effect of CR on body mass, fat mass, lean mass and WAT mass was conserved (Fig. S5A-H). Similarly, an OGTT revealed that the diet and sex effects on glucose homeostasis persisted (Fig. S6A-B).

CR decreases overall energy expenditure to a similar extent in male and female mice
We next investigated whether altered energy expenditure explained the sex differences in the CR response. As expected, CR significantly decreased absolute energy expenditure, with a greater decrease at night than in the day (Fig. 4A-B). The daytime diet differences disappeared when normalised to lean body mass (Fig. 4C), suggesting that they are driven primarily by the loss of lean mass and a consequent reduction in basal metabolic rate. In contrast, CR-induced decreases in nighttime and total energy expenditure persisted even when normalised to lean body mass (Fig. 4C), indicating that factors beyond altered lean mass contribute to this CR effect. The relationship between lean mass and energy expenditure (*P,* Slope) did not differ among the groups; however, the intercepts of the best-fit lines (*P,* Intercept) did differ significantly (Fig. 4D). Thus, for a given lean body mass, CR-fed males and females had significantly lower energy expenditure than their AL-fed counterparts (Fig. 4D). Within the CR-fed groups, females also had significantly lower total energy expenditure than males for a given lean mass (Fig. 4D). CR females also tended to have lower nighttime energy expenditure than CR males, but this did not reach

statistical significance (Fig. 4B). Indeed, within each diet there were no sex differences in absolute energy expenditure and no significant sex-diet interactions. Importantly, total energy expenditure did not differ between CR-fed males and females (Fig. 4B), indicating that females' resistance to overall weight loss is not a result of decreased energy expenditure during CR.

Sex differences in glucose homeostasis are not explained by differential glucose uptake
The basis for the sex differences in CR-improved glucose homeostasis was next investigated. We first used positron emission tomography-computed tomography (PET/CT) with ^{18}F-fluorodeoxyglucose (^{18}F-FDG) to assess glucose uptake *in vivo* (Fig. 5, S7). We focussed on tissues with high basal and/or insulin-stimulated glucose uptake, including the heart, brain, skeletal muscle, BAT, WAT, bone, and the bone marrow (BM) (28); the latter was also included because CR increases bone marrow adipose tissue (BMAT), another site of high basal glucose uptake (29).

Consistent with previous observations (29), we found that glucose uptake was highest in BAT, followed by the brain, bones, BM, the heart and skeletal muscle (Fig. 5A-B, S7A). However, CR did not stimulate glucose uptake in any of these tissues and decreased tibial glucose uptake (Fig. 5A), including in the BMAT-enriched distal tibia (Fig. S7A). Gamma counting further revealed no significant effect of CR in the gastrocnemius muscle, arm bones, liver, pancreas, spleen or thymus (Fig. 5C, S7B); however, across both sexes, CR tended to increase glucose uptake in iWAT and gWAT and this effect was significant when assessed across all WAT depots (Fig. 5C). CR also significantly increased glucose uptake in the kidneys (Fig. 5C). Several sex differences were further detected, independent of diet. Thus, females had lower glucose uptake into gastrocnemius muscle and higher glucose uptake into BAT, the heart, gWAT, the humerus, the forearm and the kidneys (Fig. 5A-C). Notably, however, sex did not significantly alter the effect of CR in any of the tissues assessed. Thus, the sexually dimorphic effects of CR on glucose homeostasis are not a result of CR increasing glucose disposal to a greater extent in males than in females.

CR exerts sexually dimorphic effects on hepatic gene expression
The lack of CR-driven differences in peripheral uptake suggested that reduced hepatic glucose production may underpin the sex differences in glucose homeostasis. To test this, we analysed transcript expression in the livers of AL and CR mice, focussing on factors important for the regulation of gluconeogenesis, lipid metabolism and insulin sensitivity. We found that CR significantly increased mRNA levels of the rate-determining gluconeogenic enzyme phosphoenolpyruvate carboxykinase (*Pck1)* in females but not in males (Fig. 6A). A similar profile occurred for peroxisome proliferator-activated receptor-alpha (*Ppara*) (Fig. 6A), a key regulator of hepatic lipid metabolism that prevents hypoglycaemia during fasting (30). Compared to CR males, CR females also had higher mRNA levels for peroxisome proliferator-activated receptor-gamma (*Pparg*), carbohydrate response element-binding protein (*Chrebp*) and *Mt-nd1,* which encodes a subunit of mitochondrial complex I (Fig. 6A). In contrast,

mRNA for *Mt-co1*, encoding a subunit of mitochondrial complex IV, did not differ between males and females but was increased with CR in both sexes. Regardless of diet, mRNA levels for fatty acid-binding protein 5 (*Fabp5*) and glycerol-3-phosphate dehydrogenase 1 (*Gpd1*) were higher in females than males, with the latter being significantly decreased by CR in each sex (Fig. 6A). Finally, principal component analysis of these qPCR data identified CR females as the most transcriptionally distinct group, further highlighting the overall sex differences in the CR response (Fig. 6B).

CR exerts sexually dimorphic effects on hepatic sphingolipid content

We next tested whether altered lipid storage, implicated in sex-dependent differences in hepatic glucose production (22), might contribute to the sexually dimorphic metabolic effects of CR. Gross hepatic triglyceride content was not affected by diet or sex (Fig. 6C). Therefore, we undertook a more-detailed molecular investigation of lipid species that may underpin altered hepatic glucose regulation, focussing on sphingolipids (31, 32). Targeted lipidomics revealed that total ceramide content was significantly lower in females than in males but was not influenced by CR (Fig. 6D); however, individual ceramide species were modulated by CR in a sex-dependent manner. Thus, 22:0 and 22:1 ceramides were lower in females than in males and were decreased by CR, with this diet effect being stronger in males (Fig. 6D, S8A). Conversely, 23:0 ceramides were greater in AL females than in AL males and were increased by CR in males only (Fig. 6D); a similar pattern occurred for 14:0, 16:0, 18:0, 18:1, 20:1, 26:0 and 26:1 ceramides (Fig. S8A).

Ceramides are generated from dihydroceramides (DHCs) and an elevated hepatic ceramide:DHC ratio promotes metabolic dysfunction and insulin resistance (32, 33). Thus, we further analysed DHCs and the ceramide:DHC ratio as potential mediators of CR's sexually dimorphic metabolic effects. As for total ceramides, total DHCs were lower in females than in males; however, unlike for total ceramides, CR increased total DHCs in males only (Fig. 6E). Differential effects of sex and/or diet also occurred for the specific DHC species. A common pattern was for CR to increase DHC concentrations in males only, such that these were lowest in AL males but similar in CR males, CR females and AL females; this pattern occurred for 16:0, 18:0, 18:1, 20:0, 23:0, 24:0, and 24:1 DHC species (Fig. 6E, S8B). CR increased 14:0 DHCs in both sexes, albeit more strongly in males, and decreased 22:0 DHCs across both sexes (Fig. 6E, S8B). The overall ceramide:DHC ratio was decreased by CR in males but not in females (Fig. 6E) and this also occurred for many of the individual species, including 16:0, 18:0, 20:0, 22:0, 22:1, 23:0, 24:0 and 24:1 (Fig. S8B); however, unlike for the total ceramide:DHC ratio, the ratio for each of these species was highest in the AL males and significantly lower in females, such that CR males had a similar ratio to females on either diet (Fig. S8B). Consistent with this, principal component analysis of these ceramide data identified AL males as the most distinct group, with CR males being more similar to females on either diet (Fig. 6F).

Together, these data indicate that CR suppresses ceramide synthesis and gluconeogenesis to a greater extent in males than in females, highlighting these as potential mechanisms for CR's sexually dimorphic effects on glucose homeostasis.

Sex differences in CR's metabolic effects are absent when CR is initiated in aged mice

To test if ageing influences sex differences in the CR response, we next investigated the effects of CR initiated in 18-month-old mice, when females are anoestrous (34). Unlike in young mice, CR decreased body mass and fat mass to a similar extent in aged males and females, while the decreases in lean mass were greater in aged females (Fig. 7A-C). Consistent with this, CR's effects on the masses of individual WAT depots did not differ between the sexes (Fig. 7D, S9A). Among non-adipose tissues CR significantly decreased pancreas mass only in aged females, but there were no sex differences in the effects of CR on the masses of other tissues (Fig. S9B-C). Thus, unlike in young mice, aged males and females show a similar effect of CR on overall adiposity and this is driven predominantly by loss of WAT.

We then investigated CR's effects on glucose homeostasis and insulin sensitivity. As shown in Figure 8A, in aged mice the ability of CR to decrease fasting blood glucose was not significantly influenced by sex. Similarly, CR improved glucose tolerance to a similar extent in males and females, with tAUC but not iAUC being significantly decreased with CR (Fig. 8B-C). The latter indicates that, as for young mice, CR improves glucose tolerance in aged mice primarily by decreasing fasting glucose. Despite no sex differences in glucose tolerance, both sex and diet decreased glucose-stimulated insulin secretion, with this being greatest in aged AL males and decreased by CR more so in males than in females (Fig. 8D). Interestingly, a sex-diet effect was observed for HOMA-IR but not for the Matsuda Index (Fig. 8E); this is similar to the effects observed in young mice (Fig. 3E) and suggests that in aged mice there remain some sex differences in the relationship between plasma insulin and gluconeogenesis in the fasted state. However, overall, these studies demonstrate that sex differences in the metabolic effects of CR are highly age dependent, with CR being equally effective in aged males and females whereas young females resist these aspects of the CR response.

In humans CR-induced fat loss is sex- and age-dependent

We next investigated if the metabolic effects of CR in humans are also sex- and age-dependent. To do so we analysed data from forty-two men and women who participated in a weight loss study involving a 4-week CR intervention. As in young mice, body mass and fat-free (lean) mass decreased more so in males than in females, with males also showing a trend for greater fold-decreases in total fat mass (Fig. 9A-F). When adjusted for age (Fig. 9G), fold-change in body mass differed significantly between males and females (*P,* Intercept = 0.025). This demonstrates that overall weight loss was greater in males than females, independent of age. In contrast, the decreases in fat mass and fat-free mass were sex- and age-dependent. Thus, fat loss in younger individuals was greater in males than females but, with age, fat loss increased in females but diminished in males (Fig. 9H) (*P,* Slope = 0.007). Conversely, the loss of fat-free mass in younger individuals was greater in females than in males and this relationship reversed in older individuals (Fig. 9I). As for fat mass, the relationship between age and loss of fat-free mass differed significantly between the sexes (*P,* Slope = 0.0004).

These age-sex interactions were particularly apparent when the cohort was divided into younger and older groups (<45 vs >45 years of age). Total weight loss was greater

in males than females regardless of age (Fig. 9J), but only in the younger age group did males show greater fat loss than females (Fig. 9K). Conversely, males' greater loss of fat-free mass occurred only in the older age group (Fig. 9L). Notably, a similar pattern of age-dependent fat loss was apparent for CR in young vs aged mice (Fig. S10). Thus, our data demonstrate that the ability of CR to decrease body mass, fat mass and lean mass is sex- and age-dependent in both mice and humans.

DISCUSSION

Herein we aimed to determine the extent and basis of sex differences in the metabolic effects of CR. We reveal that 96.6% (mouse) and 95.7% (human) of published CR research overlooks potential sex differences, with most mouse studies using males only whereas most human studies combine data from males and females. We show that, in young mice, CR decreases fat mass and improves glucose homeostasis to a greater extent in males than in females, whereas these sex differences are blunted or absent when CR is initiated in old mice. The effects of CR on hepatic gene expression and sphingolipid content also differ between young male and female mice, highlighting the liver as a key mediator of CR's sexually dimorphic metabolic effects. Finally, we reveal that humans also exhibit age-dependent sex differences in the CR response, with younger age groups showing greater fat loss in males than females whereas, with increasing age, CR-induced fat loss becomes similar between the sexes. Together, these data identify sex and age as factors that have a major influence on the metabolic health outcomes of CR.

Age-dependent sex differences in CR's metabolic benefits
Our observation that young females resist overall weight loss during CR is consistent with some previous findings in mice (20, 21, 35), monkeys (3), and humans (36-41). Young females' resistance to CR-induced fat loss has also been reported in mice (20, 21, 42-44) and humans (19, 38, 45-47); however, fewer studies have directly assessed sex differences in the effect of CR on glucose homeostasis and insulin sensitivity. Many studies have included males and females but did not sex-stratify their analyses, thereby overlooking the potential impact of sex on these important metabolic outcomes (12, 48). Nevertheless, CR in monkeys and humans has been found to improve glucose homeostasis and insulin sensitivity more so in males than in females (3, 39), consistent with our findings in young mice.

Our observations fundamentally extend this previous research by demonstrating, for the first time, that these sex differences are age dependent. The novelty of this finding highlights that relatively few studies have investigated CR when initiated in aged subjects. In 2004, Dhahbi *et al* showed that CR's health benefits persist even when initiated in aged mice (49), and subsequent research has compared the metabolic effects of CR when implemented in young vs aged animals (50, 51). Unfortunately, these studies did not include females, an oversight that has persisted in the CR field (Figs. 1,S1). Other notable research has comprehensively compared aged males and females, but only in the context of lifelong CR that was initiated in younger mice (52). Two recent studies have also highlighted age-dependent sex differences in the effects of methionine restriction or time-restricted feeding, but these dietary interventions were limited to young vs middle-aged mice (53, 54). The most relevant previous research is perhaps from Brandhorst *et al,* who observed significant visceral fat loss in female mice fed a 'fasting-mimicking diet' from 16-20.5 months of age (55). While this is consistent with our observations in aged females (Fig. 7,S9), it is limited by having not assessed glucose homeostasis nor including male counterparts for direct comparison. Thus, our study is the first to compare CR when initiated in young or old males and females and to thereby identify age-dependent sex differences in CR's metabolic effects.

Our findings have several implications for interpreting previous CR research. Although most studies show that young females resist overall weight loss and fat loss during CR, such sex differences have not been universally observed. For example, in young female rats some studies report that CR decreases body mass and/or fat mass to a similar extent in both sexes (56-58), while others find fat loss to be even greater in females than in males (59, 60). CR-induced fat loss has also been noted in young female mice (61-64), albeit in studies that lack males for direct comparison. Several human studies also show no differences in total fat loss with CR, even in younger women who are unlikely to be menopausal (39, 41, 65). What are the reasons for these inconsistent results, and can they further inform the basis for sex differences in the CR response?

Considering our present findings, one possibility is that, even in younger subjects, age differences might have influenced the outcomes of these CR studies. Indeed, both Janssen *et al* and Das *et al* studied women who, while pre-menopausal, were significantly younger than their male counterparts (41, 65). Given that males' fat loss decreases as they get older (Figs. 9H,K), this disparity could have undermined the ability of these studies to detect sex differences in fat loss. A related, broader issue is that much human CR research has studied females who are over 45 years old (36, 38, 39, 45, 47, 66-68). This is important because it is the age at which oestrogen levels typically begin to decline (69) and after which we demonstrate no sex differences in fat loss (Fig. 9K). Thus, our findings underscore the importance of properly age-matching males and females, and ideally confirming menopausal status, if the response to CR is to be reliably assessed.

While age is clearly an important variable, many animal studies still find no resistance to fat loss in young females (61-64), even when appropriately age-matched to male counterparts (56-60). Therefore, other factors must also be influencing CR's sexually dimorphic effects. One possibility is that males show a greater CR response because they are usually heavier and fatter than females at the onset of CR, and thus have more fat available to lose. Consistent with this 'baseline adiposity' hypothesis, during CR in rats Hill *et al* found total fat loss to be greater in young females than young males and, unlike in mice, pre-CR adiposity was greater in the females (59, 60). Moreover, we show that sex differences in baseline adiposity become less pronounced with ageing, such that % fat mass is similar in aged male and female mice (Fig. 7D). Therefore, this hypothesis could explain why CR decreases fat mass in the fatter, aged females, whereas young, leaner females resist this effect. However, two key points argue against this hypothesis. Firstly, we show that young females' resistance to fat loss still occurs in obese and overweight humans, among whom % fat mass is greater in females than in males, irrespective of age (Fig. S11). This is consistent with numerous previous studies of overweight or obese humans, in which CR causes a greater degree of weight loss and/or fat loss in males than females (19, 36-40, 45, 47, 66). Secondly, in young, non-obese humans, sex differences in fat loss persist even after controlling for variation in baseline fat and body mass (41). Thus, sex differences in the CR response are a not simply a consequence of baseline differences in these parameters.

A third possibility is that young females' resistance to weight and fat loss can be overcome if the extent and/or duration of CR is great enough. Our CR protocol involves a 30% decrease in daily caloric intake over six weeks. In contrast, those studies

reporting fat loss in young females typically used CR of a greater extent and/or
duration. For example, Hill *et al* studied 60% CR for 11 weeks (59, 60) while Valle *et*
al used 40% CR for 14 weeks (57, 58). Others have tested CR of a similar extent to
our studies (20-30%) but for longer durations, ranging from ten weeks to six months
(56, 61, 63). It is now well established that the CR response is dose and duration
dependent (5, 21, 43, 52, 70, 71), with the former having been systematically
investigated in male mice (72-75). Thus, a key goal for future studies will be to
determine how sex and age influence these dose- and duration-dependent effects.
519
A final important consideration is the distinction between total versus regional
adiposity. Indeed, the CALERIE trial finds no sex differences in the loss of total fat
mass but does show that, compared to men, women resist loss of trunk fat during CR
(41). This is consistent with our data showing that fat loss in young male mice is
greatest for the visceral depots (Figs. 2G, S3A, S5G), and with previous observations
for CR in obese humans (19, 45, 47, 66). Some CR studies may therefore have failed
to detect sex differences because they measured only total adiposity, missing out on
possible site-specific differences (39, 76). Thus, any studies seeking to fully
understand sex differences in CR's metabolic effects, whether in preclinical models or
in humans, should always aim to assess adiposity on both whole-body and regional
levels.
531
532
Basis for young females' resistance to weight loss and fat loss
Our studies also shed light on the mechanistic basis for these sex differences. The
ability of young females to resist weight loss and fat loss suggests that, compared to
males, females are better able to mitigate negative energy balance during CR.
Consistent with this, previous murine CR studies show that females decrease total
energy expenditure more than males, possibly resulting from greater suppression of
BAT activity (20, 57, 58). In contrast, we find that CR suppresses energy expenditure
similarly in young males and females (Fig. 4) without suppressing BAT activity (Fig.
5). This suggests that our females resist weight loss through other means. However,
a key consideration is the timing of these analyses: our mice underwent indirect
calorimetry from weeks 4 to 6 of CR, when body masses and adiposity are no longer
declining; instead, the sexually dimorphic decreases in body mass and composition
occur during the first two weeks of CR (Fig. 2A,C). Hence, we propose that females
do suppress energy expenditure more than males during this initial period, thereby
helping them to resist weight loss. Future studies should therefore ensure that any
calorimetry analyses coincide with relevant metabolic responses of CR.
549
Despite this limitation, our calorimetry data, together with our histological analyses,
provide further insights about why young females resist fat loss specifically. We show
that, compared to CR males, CR females tend to have greater decreases in energy
expenditure during the night (Fig. 4B), a time when, with our CR regimen, energy is
derived predominantly from increased fat oxidation (77). Thus, one possibility is that
young CR females maintain fat mass because of decreased fat oxidation. Previous
studies have also shown that CR in young female mice enhances WAT lipogenesis
without increasing lipolysis (62), consistent with our finding that CR does not decrease
adipocyte size in young females (Figs. 2H-I, S4). The situation in humans differs
slightly: CR enhances WAT lipolysis in both sexes but females resist this effect (38).
Together, these differences in lipid metabolism likely explain why fat loss in young

females is absent or blunted during CR. Whether ageing alters these sex differences
in lipid metabolism remains to be determined.
563
564
Importance of visceral fat loss for CR's metabolic effects
These effects on fat mass and lipid metabolism likely contribute to young females'
resistance to improved glucose homeostasis during CR. Human studies have shown
that inefficient WAT lipolysis is associated with impaired glucose metabolism in
women (78) and that, during CR, decreased fat mass is linked to the improvements
in insulin sensitivity (12). Similarly, studies of CR or lipectomy in animal models
suggest that visceral fat loss contributes to CR's broader metabolic benefits (6, 9-
12). However, no previous research has investigated if this relationship between fat
loss and insulin sensitisation differs between the sexes. Our work therefore supports
and extends these previous studies by demonstrating that CR-induced fat loss
coincides with improved glucose homeostasis and insulin sensitivity in males and
aged females, whereas young females' resist these metabolic effects.
577
578
The liver as a key driver of sex differences in glucose tolerance
Several of our findings highlight the importance of the liver as a key mediator of CR's
sexually dimorphic metabolic effects. Firstly, our PET/CT data show that CR improves
glucose tolerance without any marked increases in peripheral glucose uptake (Fig. 5).
This includes the BMAT-rich distal tibia, suggesting that, although BMAT has high
basal glucose uptake (29), increased BMAT glucose uptake is not required for
improved glucose tolerance during CR. Moreover, although CR does increase glucose
uptake in WAT and the kidneys this effect does not differ between males and females
and therefore is unlikely to explain the sex differences in glucose tolerance (Fig. 5B).
Together our data are consistent with human studies showing that, in the fasted state,
CR does not affect overall glucose disposal (79, 80). Secondly, during an OGTT we
show that CR decreases the tAUC but not the iAUC, highlighting decreases in fasting
glucose as the main driver of the improvements in glucose tolerance. Thirdly, we show
that CR's effects on hepatic gene expression differ between young male and female
mice. One notable finding is that CR increases *Pck1* expression in females only (Fig.
6A), consistent with CR females having greater fasting glucose than their male
counterparts. Diet and/or sex differences in hepatic *Ppara, Pparg, Chrebp, Mt-nd1,*
Fabp5 and *Gpd1* expression are also notable because these encode proteins that
coordinate lipid metabolism and gluconeogenesis during fasting (81-85); hence, their
differential expression supports the conclusion that CR in females enhances hepatic
gluconeogenesis, lipogenesis and lipid oxidation, whereas in males these effects of
CR are blunted and/or absent. This is similar to the results of Della Torre *et al,* who
show that fasting for 6 hours inhibits hepatic gluconeogenesis and lipogenesis in male
mice but not in females (22).
603
How is loss of visceral fat linked with these hepatic effects? Visceral adiposity is
strongly associated with increased hepatic fat storage (86), and decreased liver fat
coincides with improved insulin sensitivity during CR in rats and humans (87, 88).
Moreover, young female mice resist the decreases in hepatic triglycerides following
short-term fasting (22) and prolonged time-restricted feeding (35). Therefore, we were
surprised to find that CR does not affect hepatic triglyceride content in young mice of
either sex (Fig. 6C). This may be because our young AL controls are lean and healthy,

and therefore already have livers that are not excessively fatty. In contrast, many of the previous studies investigated CR in animals or humans that are older and/or with type 2 diabetes (35, 87, 88), thereby increasing the baseline/control hepatic fat content. This would not explain why our findings differ to those of Della Torre *et al,* who did study young, healthy mice (22); however, it is perhaps unsurprising that our 6-week CR elicits different effects to those induced during shorter-term fasting.

Our above observations are important because they show that CR can exert metabolic benefits without decreasing hepatic triglyceride content. In contrast, we find that effects on sphingolipids may play a role, with CR altering hepatic sphingolipid content more so in young males than in young females. Here, the effect of diet and/or sex depends on the sphingolipid species concerned (Fig. 6F; S8). This is reminiscent of the findings of Norheim *et al,* who report sex differences in hepatic ceramide content in mice fed obesogenic diets (89). They further showed that the relationship between ceramides, body fat and insulin sensitivity also differs between males and females, focussing on 16:0 ceramide (C16:0). Indeed, ceramides' influence on insulin sensitivity depends on the ceramide species. For example, hepatic increases in C16:0 have been implicated with the development of obesity-associated insulin resistance (32, 90-92). This is seemingly at odds with our finding that CR increases hepatic C16:0 in young males but not young females (Fig. S8). Thus, unlike in obesity, increased hepatic C16:0 during CR appears to be associated with improvements in insulin sensitivity.

The rate of conversion of ceramides to DHCs has also been suggested to be a root cause of insulin resistance (93, 94). This underscores the concept that the ratio between each ceramide and DHC species may have a larger influence on metabolic function than the absolute values of either, a concept supported by others (33). More research is needed to determine how specific ceramide and DHC species, and the ratios thereof, impact sex differences in the health benefits of CR. Despite these remaining questions, our work extends previous observations by revealing that CR's hepatic effects are sexually dimorphic and that this contributes to the sex differences in glucose tolerance and insulin sensitivity.

Oestrogen as a modulator of the CR response
In many cases young females' resistance to the metabolic effects of CR is associated with them already being relatively metabolically healthy on an AL diet. For example, young AL females have lower visceral fat mass and are more glucose tolerant than AL males (Figs. 2G, 3B-C), and are similar to CR males and females in terms of fasting insulin and HOMA-IR (Fig. 3D-E). We also show that, for most sphingolipid species, the hepatic ceramide:DHC ratio is similar between AL females and CR mice of either sex (Fig. S8C). These patterns suggest that the mechanisms through which CR decreases visceral adiposity, improves glucose tolerance and modulates sphingolipid metabolism are already at least partially active in young females, even on an AL diet.

This observation helps to clarify potential mechanisms for CR's age-dependent sex differences. Sexual dimorphism in metabolic function typically results from sex hormones, with genetic factors and/or developmental programming also playing a role. For example, X-chromosome-linked genetic factors influence sex differences in adiposity independently of sex hormones or other secondary sexual characteristics (95, 96), while the developmental and neonatal environment causes sexually

dimorphic metabolic programming (97). However, it is unlikely that these developmental or genetic factors explain the sex differences we observe. Developmental programming is unlikely because the early life environment of our mice was the same for both sexes and was devoid of any adverse or stressful stimuli. Similarly, if X-linked genetic factors were responsible then the sex differences would be expected to persist in aged females, but they mostly do not. This leaves sex steroids as the likely mediator. Consistent with this, oestrogen is now well known to influence metabolic function and the fasting response (13, 22), and many actions of oestrogens overlap with those central to the effects of CR (98-101). Herein, we show that the sex differences in CR's metabolic effects are absent or greatly diminished when CR is initiated at 18 months of age, when C57BL/6 females are typically anoestrus (34, 69). Similarly, CR does decrease fat mass in young ovariectomised females (102), while oestrogen treatment protects male mice from obesity-induced insulin resistance (103, 104). Therefore, we propose that young females resist the metabolic effects of CR because, through endogenous oestrogen action, they already have high activation of pathways that mediate CR's metabolic effects.

On a molecular level we speculate that these age and sex differences relate to oestrogen receptor alpha (ERα) activity in adipose tissue and the liver, a hypothesis based on numerous lines of evidence. For example, in mice ERα contributes to lower visceral adiposity in females than males (105), while in males pharmacological ERα activation decreases fat mass (106). Oestrogens also act via ERα to prevent insulin resistance and suppress hepatic gluconeogenesis (107, 108), whereas global ERα knockout mice have impaired hepatic insulin sensitivity (109). ERα has also been implicated in sex differences in hepatic metabolism during fasting (22). Thus, a key goal of future studies should be to determine if oestrogen and ERα contribute to CR's age-dependent, sexually dimorphic metabolic effects.

Limitations and further considerations
Our study represents a significant advance in knowledge by revealing that age and sex have a substantial influence over the metabolic benefits of CR. This highlights the need for preclinical and human CR research to address these variables as key determinants of the CR response, a consideration that we show has been lacking in most CR research. We show that these sex differences occur on a systemic, physiological level and that they extend to cellular and molecular effects within adipose tissue and the liver, observations that identify potential mechanisms for the age-dependent sex differences; however, a general limitation of our study is that these mechanisms remain to be directly addressed. Specific limitations are as follows.

Firstly, regarding young females' resistance to weight and fat loss, our calorimetry analyses did not find any significant sex differences in energy expenditure. As explained above, this may be because these analyses were done later in the CR protocol when dynamic changes in fat and body mass have stabilised. Our calorimetry system was also unable to reliably assess the respiratory exchange ratio of our CR mice because they have increased torpor and a lower metabolic rate that is at the limit of detection for our system. Therefore, future studies should analyse energy homeostasis within the first two weeks of onset of CR, both in young and old mice, using a more-sensitive indirect calorimetry system.

Secondly, regarding insulin sensitivity and glucose homeostasis, it would be informative to use hyperinsulinaemic-euglycaemic clamps to comprehensively assess systemic and hepatic insulin sensitivity and determine if, as we propose, the differences in glucose homeostasis relate primarily to effects on gluconeogenesis.

Thirdly, the role of oestrogen remains to be directly tested. This could be done in mice through ovariectomy studies and/or manipulation of ERα activity. We have also tried to determine oestradiol concentrations in our mouse plasma samples using mass spectrometry, a method required for reliable assessment of sex steroid concentrations (110). However, despite extensive optimisation efforts in our world-leading mass spectrometry facility, we have not yet been able to reliably measure oestradiol using the relatively small plasma volumes available. Related to this, in our human studies we were unable to confirm the menopausal status of the female participants. Other limitations of our human studies are that they included relatively few younger participants and lacked analysis of regional adiposity (e.g. visceral versus subcutaneous adipose tissue). Future human research must therefore prioritise assessment of menopausal status and regional adiposity if the outcomes of CR are to be fully interpreted.

A final limitation regards the underlying molecular mechanisms. Pathways implicated in mediating CR's effects include activation of AMPK, SIRT, IGF-1 and FGF21, modulation of mitochondrial function and inhibition of mTOR (5, 18). As discussed above, we speculate that ERα activation may impinge on these pathways and thereby contribute to sex differences in the CR response. Thus, genetic, pharmacological and/or other approaches to manipulate these candidate pathways could be used to further dissect the molecular mechanisms underlying CR's age-dependent sexually dimorphic metabolic effects. Such approaches are a promising basis for future research.

Beyond these specific limitations, a broader question is whether other effects of CR also differ between males and females in this age-dependent manner. Some research suggests that CR's effects on cardiovascular function and healthy ageing differ between males and females (111-113); however, given the dearth of CR studies that directly assess sex differences (Fig. 1, S1) there remains a critical need to determine how age and sex influence the many other outcomes of CR. This is important not only for understanding fundamental biology, but also if we are to translate the benefits of CR, or CR mimetics, to improve human health. For example, if younger women resist many of CR's health benefits, should any CR-based intervention focus only on older women, who may be more responsive?

Finally, while it will be useful to understand the extent and basis for these sex differences, it is also important to consider their broader purpose. Adipose tissue profoundly influences female reproductive function, with adipose-secreted factors acting within the central nervous system to indicate if the body's energy stores are sufficient to support reproduction (114). The hormone leptin is the key adipose-derived factor that impacts the hypothalamic-pituitary-gonadal axis: if fat mass is too low, for example in anorexia nervosa or lipodystrophy, then leptin levels are also insufficient, resulting in decreased gonadotropin secretion and impaired reproductive capacity. Normal menses resumes in anorexic individuals who regain weight and fat mass, or in lipodystrophic individuals following leptin treatment (114, 115). Notably, we

previously demonstrated that female mice resist hypoleptinaemia during CR (21). Thus, we speculate that females resist fat loss during CR to preserve reproductive function. Given the tight interplay between adipose tissue and systemic metabolic homeostasis, such resistance to fat loss then has consequences for CR's other metabolic effects. This evolutionary pressure to overcome periods of food scarcity would thereby have driven the divergence of the CR response in males and females, with implications for current efforts to exploit CR therapeutically to improve human health.

ACKNOWLEDGEMENTS

This work was supported by grants from the Medical Research Council (MR/M021394/1 to W.P.C), the British Heart Foundation (RG/16/10/32375 and FS/19/34/34354 to A.A.S.T.; 4-year BHF PhD Studentship to B.J.T and R.J.S.), the University of Edinburgh (Chancellor's Fellowship to W.P.C.; PhD Studentship to A.L.), the Carnegie Trust (RIG007416 to W.P.C.), the Wellcome Trust (Technology Development Award 221295/Z/20/Z to A.A.S.T.) and the Chief Scientist Office of the Scottish Government (SCAF/17/02 to R.H.S.). C.F. and A.M.J. gratefully acknowledge financial support from the Scottish Government as part of the RESAS Strategic Research Programme at the Rowett Institute, University of Aberdeen. We are also grateful to the British Heart Foundation for providing funding towards establishment of the Edinburgh Preclinical PET/CT laboratory (RE/13/3/30183) and support of the 2018 Very Important Project (VIP) PET prize. We also thank Dr Hai-Bin Ruan (University of Minnesota Medical School) and Dr Eleonora Pagano (Pontifical Catholic University of Argentina) for kindly providing a full text of Ballor and Poehlman's 1994 study (76), and all staff at Edinburgh Bioresearch & Veterinary Services for their superb technical support.

AUTHOR CONTRIBUTIONS

Contributions are based on the CRediT (Contributor Roles Taxonomy) and are as follows: *Conceptualisation*, K.J.S., B.J.T. G.G., R.J.S. A.M.J. and W.P.C.; *Methodology*, K.J.S., B.J.T., A.A.S.T., P.D.W., R.H.S. A.M.J. and W.P.C.; *Investigation*, K.J.S., B.J.T., R.J.S., A.L., H.J.W., X.H., D.M., C.J.A-C., P.D.W., C.F., R.H.S. A.M.J. and W.P.C.; *Formal Analysis*, K.J.S., B.J.T., R.J.S., A.L., P.D.W., C.F., N.M.M., A.M.J. and W.P.C.; *Resources*, A.A.S.T., G.G., N.M.M., R.H.S., A.M.J. and W.P.C.; *Writing – Original Draft*, K.J.S., B.J.T. and W.P.C.; *Writing – Review & Editing*, K.J.S., B.J.T., G.G., R.H.S, N.M.M W.P.C.; *Visualisation*, K.J.S., B.J.T., and W.P.C.; *Supervision,* G.G. N.M.M., R.H.S., A.M.J. and W.P.C.; *Funding Acquisition*, K.J.S, A.A.S.T., G.G., N.M.M., R.H.S., A.M.J. and W.P.C.

DECLARATION OF INTERESTS

No competing interests are declared.

METHODS

Table 1: Key Resources

REAGENT or RESOURCE	SOURCE	IDENTIFIER
CHEMICALS		
18F-Fluorodeoxyglucose	Edinburgh Clinical Research Imaging Centre (Edinburgh, UK)	N/A
Ribozol[TM]	Amresco (USA)	N580
OLIGONUCLEOTIDES		
See Table 4	This paper	N/A
EXPERIMENTAL MODELS: Organisms/Strains		
Mouse: C57BL/6J or C57BL/6NCrl	Charles River	027
SOFTWARE		
Prism	GraphPad Software, LLC	v9.2.0
PMOD	PMOD Technologies LLC (Zurich, Switzerland)	v3.806
OTHER		
TSE Phenomaster	TSE	PhenoMaster 1.0 with software version 6.1.9
OneTouch Verio Glucometer	OneTouch	User's manual here
Gamma counter	PerkinElmer (USA)	Wizzard[2]
Preclinical PET/CT scanner	Mediso (Budapest, Hungary)	nanoScan PET/CT 122S
Phenomenex Kinetex C8 HPLC system	Phenomenex	00D-4499-AN

Animals

Studies in C57BL/6NCrl and C57BL/6JCrl mice were approved by the University of Edinburgh Animal Welfare and Ethical Review Board and were done under project licenses granted by the UK Home Office. Mice were originally obtained from Charles River (UK) and then bred in-house. They were housed on a 12 h light/dark cycle in a specific-pathogen-free facility with free access to water and food, as indicated. Differences in mouse cohorts are summarised in Table 2, along with the groups being compared and the experimental unit. The exact number of mice used is stated in the figure legends. Sample sizes for each cohort were determined by power calculations using G*Power software, with effect sizes based on pilot studies and/or previously published data for fat loss and improved glucose tolerance (the primary outcomes being tested) (21, 42); to increase power, data for young mice are reported across multiple cohorts from separate studies, each of which revealed the same sex differences in CR's metabolic effects. Mice were randomly assigned to each diet. To minimise potential confounders, such as the order of treatments or measurements, mice were treated or analysed based on the order of their ID number, a variable randomly distributed across the experimental variables (diet and sex). Blinding was not possible during the *in vivo* experiments because the experimenter(s) can see,

based on presence or absence of diet, which mice are AL and which are CR; however, data analysis was done in a blinded manner. Mice were excluded from the final analysis only if there were confounding technical issues or pathologies discovered at necropsy. Ten mice from the young study were excluded from the analysis of OGTT and necropsy tissue masses because of a technical issue during gavaging for the OGTT, which compromised their subsequent food intake. One mouse was excluded from PET/CT data analysis because it accidentally had two doses of FDG administered. One aged mouse was excluded because of developing confounding liver tumours. For some cohorts, specific tissues were not weighed at necropsy and therefore these mice have no data for these readouts. For further details, please see the Source Data file.

Mouse CR studies
Before assigning to AL or CR diets, mice were housed individually between 8-9 weeks of age (for young mice) or 77-78 weeks of age (for old mice). During this time, mice were fed a control diet (Research Diets, D12450B) *ad libitum* and daily food intake was determined by weighing the food remaining in each cage each day. Following one week of single housing, all mice were randomly assigned to control diet (Research Diets D12450B, provided AL) or the CR diet (Research Diets D10012703, administered at 70% of the average daily AL diet consumption). This CR diet is formulated for iso-nutrition at 70% CR, ensuring that malnutrition is avoided. The evening-fed CR mice (Fig. S5-S6) were fed daily between 1800-1900, while all other CR mice were fed daily between 0900-1000. For the first two weeks of CR mice were monitored and weighed daily to ensure they did not lose more than 30% of their initial body mass. For the following 4 weeks of CR, body mass was recorded weekly. Mice were assessed non-invasively for body fat, lean mass, and free fluid weekly from weeks 0 to 6 using TD-NMR (Minispec LF90II; Bruker Optics, Billerica, MA, USA). At necropsy, mice were humanely sacrificed via cervical dislocation and decapitation. Tissues were harvested, weighed, and stored either in 4% formalin or on dry ice. Tissues in formalin were fixed at 4ºC for 2-3 days before being washed and stored in DPBS (Life Technologies). Tissues on dry ice were stored at -80ºC prior to downstream analysis.

	Young	Aged	Evening-fed
Age at single housing	8 weeks	77 weeks	8 weeks
Age at start of CR	9 weeks	78 weeks	9 weeks
Time of feeding	0900-1000	0900-1000	1800-1900
Duration of CR	6 weeks	6 weeks	4 weeks
Related data	Fig. 2-6, S2-S4, S7-S8, S10	Fig. 7-8, S9-S10	Fig. S5-S6
Experimental unit	Single mouse		
Groups compared	AL vs CR (within sex); Male vs Female (within diet); and diet*sex interactions		

Table 2. Summary of CR protocol in each group of mice. Because mice are singly housed, each mouse represents an independent experimental unit.

Human weight loss study – Participants
Forty-five overweight and obese males and females; age 21-61y, Body Mass Index (BMI) 26-42 kg/m^2, were recruited by newspaper advertisement to participate in a

weight loss (WL) dietary intervention study. Thus, the participants were non-randomly selected individuals who were sufficiently motivated to actively respond to the request for volunteers. During recruitment, participants underwent a medical examination and their general practitioners were contacted to confirm medical suitability to participate in the study. Written, informed consent was obtained, and the study was reviewed by the NHS North of Scotland Research Ethics Service (ethics number 09/S0801/80). Participants were informed that the study was funded by an external commercial sponsor, but the company was not named, and all packaging of the food range was removed before distribution. Three participants withdrew for personal reasons therefore data for n = 42 is presented throughout; participant characteristics are summarised in Table 3.

Sex	Number of subjects	Age (years)	BMI (kg/m^2)	Glucose (mM)	Diabetic
Male	20	48.61 $\pm$ 10.7	34.7 $\pm$ 4.2	5.6 $\pm$ 0.7	0%
Female	22	44.7 $\pm$ 9.7	32.8 $\pm$ 3.7	5.2 $\pm$ 0.3	0%

Table 3. Human subject characteristics at baseline. Age, BMI and glucose are mean $\pm$ SD. *ND* = not determined.

Human weight loss study – Protocol
Participants were asked to attend the Human Nutrition Unit (HNU) at the Rowett Institute of Nutrition and Health (Aberdeen, UK) twice weekly to pick up food supplies. All required food, drinks and snacks were provided and the study dietician was present to answer any questions. Participants were provided with a diet guide whereby women were encouraged to consume no more than 6.3 MJ/d (1,500 kcal) and men 8.4 MJ/d (2,000 kcal). They were encouraged to consume 3 meals a day (breakfast, lunch and dinner) with a portion of protein at each meal, and 5 portions of fruit and vegetables, 2 portions of fats, 3 portions of milk and dairy and 4 portions of wholegrain bread, potatoes and cereals each day. Participants completed a descriptive food diary every day to record what and when they ate. Participants ate *ad libitum*, choosing when and what to eat from their supplies. They completed a detailed 7d weighed dietary record during week 4 of the WL phase of the study, to compare eating habits with a baseline 7d weighed record they completed before the study started. Energy and nutrient intakes were calculated using the Windiets programme (Univation Ltd; The Robert Gordon University, Aberdeen, UK).

Human weight loss study – Measurement of body composition
Before and after WL, at weeks 0 and 4 respectively, measurements of body composition were conducted under standardized conditions as described previously (116), with participants instructed to fast overnight (at least 10 h). Height was measured at the beginning of the study to the nearest 0.5 cm using a stadiometer (Holtain Ltd, Crymych, Dyfed, Wales). The body weight of the participants was measured at screening, during food provision days and on all test days, wearing a previously weighed dressing gown, to the nearest 100 g on a digital scale (DIGI DS-410, C.M.S. Weighing Equipment Ltd, London, UK). Fat mass (FM) and fat free mass (FFM) were measured with the use of air displacement whole-body plethysmography (BodPod® Gold - Body Composition System, Cosmed, Italy) (117).

Indirect calorimetry
Male and female AL and CR C57BL/6J mice aged ~13 weeks were housed individually in TSE PhenoMaster cages for indirect calorimetry, monitoring of physical activity, and measurement of *ad libitum* food and water consumption. Mice in each group were first housed in these cages for 24 h for acclimation and then housed for a further 72 h for data collection.

Oral glucose tolerance tests (OGTT)
Mice were given a half portion of food the night (6pm) prior to the metabolic challenge. At 12:00 pm on the day of the OGTT a basal blood sample was collected by venesection of the lateral tail vein into EDTA-coated capillary tubes (Microvette 16.444, Sarstedt, Leicester, UK) and the basal glucose levels was measured using a glucometer (OneTouch Verio IQ, LifeScan Inc., Zug, Switzerland). D-glucose (Sigma, G8270,Poole, UK) was then administered at 2 mg per g of body mass by oral gavage of a 25% (w/v) solution. At 15, 30, 60 and 120 minutes post-gavage, blood glucose was measured by a glucometer and tail vein blood was sampled in EDTA tubes, as for the basal (0 min) timepoint. Blood samples were kept on ice prior to isolation of plasma. Insulin was then measured in plasma samples by ELISA, following the manufacturer's instructions (cat. #90080, ChrystalChem, Chicago, IL, USA). Insulin was converted from ng/ml to µU/mL by multiplying the ng/mL value by 28.7174 (based on mouse insulin molar mass of 5803.69). Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) and Matsuda index were calculated as previously described (25, 26)).

PET/CT scanning
Immediately before and 15- and 60-min post [18]F-FDG intraperitoneal injection, blood glucose was measured by tail venesection and blood was sampled directly into EDTA-microtubes (Sarstedt, Leicester, UK). At 60-minutes post [18]F-FDG injection, mice were anaesthetised and [18]F-FDG distribution was assessed by PET/CT. Following the PET/CT scan, mice were sacrificed by overdose of anaesthetic (isoflurane), death was confirmed by cervical dislocation and tissues of interest were then dissected in a lead-shielded area. Half of each of the dissected tissues were snap frozen on dry ice or placed into 10% formalin. [18]F-FDG uptake was then determined using a gamma counter (PerkinElmer) as previously described (29)

PET/CT analysis
PET/CT images were reconstructed and data analysed using PMOD version 3.806 (PMOD, Zurich, Switzerland). SUVs were calculated for regions of interest, namely BAT, iWAT, gWAT; heart; bone tissue (without BM) from tibiae, femurs, and humeri; To distinguish bone tissue from BM, a calibration curve was generated using HU obtained from the acquisition of a CT tissue equivalent material (TEM) phantom (CIRS, model 091) and mouse CT scans, as previously described (29).

Histology and Adipocyte size quantification
Dissected mouse tissue was fixed for 48 h and paraffin embedded by the histology core at The University of Edinburgh's Shared University Research Facilities (SuRF). Paraffin-embedded tissue sections were then sectioned at 100 µm intervals using a Leica RM2125 RTS microtome and collected onto 76 x 26 mm StarFrost 624 slides (VWR, UK). The slides were baked at 37ºC overnight before Haematoxylin and Eosin (H&E) staining. Quantification of adipocyte number and size was per formed on H&E

955 stained adipose tissue using the ImageJ plug-in Adiposoft, an automated software for
956 the analysis of adipose tissue cellularity in histological sections (118).
957
RNA isolation and reverse transcription
RNA was isolated from tissues using Ribozol™ solution (cat. No. N580, Amresco,
USA,) according to the manufacturer's protocol. RNA quantity and quality were
quantified using a NanoDrop spectrophotometer (Thermo Scientific,USA), and cDNA
was synthesised from 1 µg RNA using the Taqman® High Capacity cDNA Reverse
Transcriptase Kit (Applied Biosystems, USA, cat. no. N8080234) in accordance with
manufacturer's guidelines and as described previously (29).
965
qPCR
Transcript expression was analysed by qPCR in duplicate reactions using qPCRBIO
SyGreen Blue Mix (part number PB20.17-20; PCR Biosystems, UK). Reactions were
loaded into 384-well qPCR plates (part number 72.1985.202; Sarstedt, UK) and run
on a Light Cycler 480 (Roche). Primers (Table 4) were designed using NCBI Primer
Blast and validated using cDNA titrations, with melting analysis and gel
electrophoresis used to confirm specificity for the intended target. Transcript
expression was calculated based on a cDNA titration loaded on each plate and was
normalised to the geometric mean of *Tbp, Rn18s* and *Ppia* expression (119).
975

Transcript	Forward primer (5'-3')	Reverse primer (5'-3')
Chrebp	CCAGCCTCAAGGTGAGCAAA	CATGTCCCGCATCTGGTCA
Fabp5	TGAAAGAGCTAGGAGTAGGACTG	CTCTCGGTTTTGACCGTGATG
Gpd1	ATGGCTGGCAAGAAAGTCTG	CGTGCTGAGTGTTGATGATCT
Mt-co1	GAGAGGCCTTTGCTTCAAAAA	AGGTTGGTTCCTCGAATGTG
Mt-nd1	CACTCCTCGTCCCCATTCTA	ATGCCGTATGGACCAACAAT
Pck1	**GTTTGATGCCCAAGGCAACT**	**CCTTCCCAGTAAACACCCCC**
Ppara	CCTGAACATCGAGTGTCGAATAT	TCTTCTTCTGAATCTTGCAGCT
Pparg	GGAAAGACAACGGACAAATCAC	TACGGATCGAAACTGGCAC
Ppia	CACCGTGTTCTTCGACATCA	CAGTGCTCAGAGCTCGAAAGT
Rn18s	CGATGCTCTTAGCTGAGTGT	GGTCCAAGAATTTCACCTCT
Tbp	ACCTTATGCTCAGGGCTTGG	GCCGTAAGGCATCATTGGAC

976 **Table 4.** Sequences of primers used for qPCR
977
Liver ceramide quantification
Liver ceramide and dihydroceramide content was determined using LC-MS
(Lipidomics Core Facility; University of the Highlands and the Islands, Inverness, UK).
As per Folch *et al.* 1957 (120), 100 mg of frozen liver tissue was homogenised in ice-
cold MeOH using a mechanical tissue homogeniser. Fifty microlitres of each sample
was combined with 100 µL ceramide 17:0 (1 µM in MeOH) and 25 µL dihydroceramide
12:0 (2 µM in MeOH) (Avanti Polar Lipids, Alabaster, AL, USA) as internal standards.
MeOH (1.875 mL) and $CHCl_3$ (4 mL) were added. Samples were then stored on ice
for 1 hour with intermittent vortex mixing, centrifuged at 2,000 rpm for 5 minutes at 4°C
and the supernatant was collected. KCl (0.9%; 1.45 mL per sample) was added to the
supernatant and the samples were centrifuged at 2,000 rpm for 5 minutes at 4°C. The
lower phase was collected and evaporated under nitrogen and reconstituted in 1 mL
$CHCl_3$. Solid phase extraction was performed via a 100 mg/3 mL Isolute silica column
(Biotage, Uppsala, Sweden). The column was conditioned twice with 3 mL $CHCl_3$ and
then loaded with 1 mL of the sample. The bound sample was washed with 2 washes

of 3 mL CHCl$_3$ then eluted with 2 additions of ethylacetate CHCl$_3$ (1:1 v/v). The
collected eluate was evaporated and dried under vacuum and reconstituted in 200 μL
MeOH. LC-MS/MS analyses were performed in positive ion mode on a Thermo TSQ
Quantum Ultra triple quadrupole mass spectrometer equipped with a HESI probe and
coupled to a Thermo Accela 1250 UHPLC system. The ceramides and
dihydroceramides were separated on a Kinetex C8 column (1.7 μm; 100 × 2.1 mm)
(Phenomenex, Macclesfield, UK). Mobile phase A was 90% H$_2$O, 9.9% acetonitrile
and 0.1% formic acid. Mobile phase B was 99.9% acetonitrile and 0.1% formic acid.
The gradient was held at 80% mobile phase B for 1 minute then gradually increased
to 100% mobile phase B over 15 minutes. Mobile phase B at 100% was maintained
for 1 minute and then equilibrated to storage conditions. The flow rate used was 500
μL per minute at 40°C. Two μL of each sample was injected into the LC-MS/MS for
ceramide analysis and 10 μL for dihydroceramide. Total ceramide and
dihydroceramide concentrations were calculated from the summed concentrations of
all the monitored molecular species. All values were normalised to wet weight of liver.
1008
Statistical analysis
Data were analysed for normal distribution using the Shapiro-Wilk normality test.
Normally distributed data were analysed by ANOVA, mixed models or t-tests, as
appropriate. Where data were not normally distributed, non-parametric tests were
used. When appropriate, *P* values were adjusted for multiple comparisons. Data are
presented as mean ± SEM or as Violin plots, as indicated in the figure legends. All
statistical analyses were performed using Prism software (GraphPad, USA). A *P*-value
<0.05 (after adjustment for multiple comparisons) was considered statistically
significant.
1018
1019 **Data availability**
1020 All source data from which the figures are based is available through University of
1021 Edinburgh DataShare at https://doi.org/10.7488/ds/3277. All other relevant data are
1022 available from the authors upon reasonable request.

FIGURE TITLES AND LEGENDS

Figure 1 – Summary of sex differences in mouse and human CR research.
PubMed was searched using MeSH terms to identify research articles that studied caloric restriction *in vivo* in mice (**A,B**) or humans (**C,D**), published between 2003 and 2021. Search results were then classified into the following groups depending on the sexes included in each study: 'Male only' = male subjects used exclusively; 'Female only' = female subjects used exclusively; 'Male & Female separate' = male and female subjects used with data reported respective to each sex, allowing sex differences to be addressed; 'Male & female combined' = male and female subjects used with data from both sexes combined either in part or in full; 'N/A' = no sex data available. **(A,C)** Cumulative publications for studies within each group. The boxes between (A) and (C) highlight the dates of funders' policies highlighting the importance of sex as a biological variable. (**B,D**) Pie charts of those studies in the 'Male & Female separate' group that considered sex a biological variable, and the proportion of these that did or did not detect sex differences. Source data are provided as a Source Data file. See also Supplementary Figure 1.

Figure 2 – Female mice resist CR-induced weight loss, fat loss and adipocyte hypotrophy.
Male and female mice on a C57BL/6NCrl or C57BL/6J background were fed *ad libitum* (AL) or a 30% CR diet from 9-15 weeks of age (0-6 weeks of CR). **(A-F)** Each week mice were weighed (A,B) and body composition was determined by TD-NMR (C-F). Body mass, fat mass and lean mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). **(G)** Masses of gWAT (gonadal WAT), iWAT (inguinal WAT), mWAT (mesenteric WAT), pWAT (perirenal WAT), pcWAT (pericardial WAT) and BAT (brown adipose tissue) were recorded at necropsy and are shown as % body mass. **(H,I)** Micrographs of H&E-stained sections of gWAT (H) were used for histomorphometric analysis of adipocyte area (I); in (H), scale bar = 100 µm. Data in (A-F) are shown as mean ±SEM of the following numbers of mice per group: *male AL*, n=42; *female AL,* n=43; *male CR,* n=44; *female CR,* n=52. Data in (G) are shown as violin plots of the following numbers of mice per group and tissue type: *male AL,* n=29 (iWAT, gWAT, mWAT, BAT), 15 (pWAT) or 14 (pcWAT); *female AL,* n=28 (iWAT, gWAT, mWAT, BAT), 14 (pWAT) or 13 (pcWAT); *male CR,* n=33 (iWAT, gWAT, mWAT, BAT), 16 (pWAT) or 15 (pcWAT); *female CR,* n=34 (iWAT, gWAT, mWAT, BAT), 15 (pWAT) or 14 (pcWAT). Data in (H-I) are shown as representative micrographs or violin plots from 5 (*male AL)* or 6 (*female AL, male CR, female CR)* mice per group. For (A-F), significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA or a mixed-effects model. For (G), significant effects of diet and/or sex within each tissue were determined by 2-way ANOVA with Tukey's multiple comparisons tests. For (I), significant effects of diet and/or sex on adipocyte area (Ad.Ar) were determined using a mixed-effects model, while significant differences in median Ad.Ar between AL and CR mice were determined 2-way ANOVA with Šídák's multiple comparisons test. *P values* from ANOVA or mixed models are shown beneath the graphs, as indicated. For (G) and (I), significant differences between comparable groups are indicated by * ($P<0.05$), ** ($P<0.01$) or *** ($P<0.001$). Source data are provided as a Source Data file. See also Supplementary Figures 2-5 and 10.

Figure 3 – The effects of CR on glucose homeostasis differ between young male and female mice.

Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. **(A)** Random-fed blood glucose was recorded each week. **(B-D)** At 13 weeks of age mice underwent an oral glucose tolerance test (OGTT). (B) Blood glucose readings during the OGTT. (C) Area under the curve (AUC) during the OGTT was determined relative to 0 mmol/L (total AUC: tAUC) and relative to baseline (incremental AUC: iAUC). (D) Glucose-stimulated insulin secretion in mice during OGTT was assessed using an insulin ELISA. **(E)** HOMA-IR and Matsuda indices of mice calculated from glucose and insulin concentrations during the OGTT. Data in (A), (B), and (D) are presented as mean ± SEM. Data in (C) and (E) are presented as violin plots overlaid with individual data points. For each group and timepoint, the following numbers of mice were used: **(A)**: *male AL,* n=42 (Wk 0), 36 (Wk2, Wk 2), 34 (Wk 3), 29 (Wk 4), 26 (Wk 5), or 22 (Wk 6); *female AL,* n=43 (Wk 0), 28 (Wk 1), 35 (Wk 2), 31 (Wk 3), 27 (Wk 4), 26 (Wk 5), or 21 (Wk 6); *male CR,* n=44 (Wk 0), 40 (Wk 1), 38 (Wk 2), 35 (Wk 3), 31 (Wk 4), 29 (Wk 5), or 26 (Wk 6); *female CR,* n= 51 (Wk 0), 44 (Wk 1, Wk 2), 41 (Wk 4), 35 (Wk 4), 33 (Wk 5), or 27 (Wk 6). **(B)**: *male AL,* n=27 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female AL,* n=26 (T_{15}, T_{30}, T_{120}), or 25 (T_0, T_{60}); *male CR,* n=27 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female CR,* n=28 (T_0, T_{15}, T_{30}, T_{60}, T_{120}). **(C)**: *male AL,* n=27; *female AL,* n=26; *male CR,* n=27; *female CR,* n=28. **(D)**: *male AL,* n=9 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female AL,* n=11 (T_0, T_{15}), 10 (T_{30}, T_{60}), or 8 (T_{120}); *male CR,* n=17 (T_0), 16 (T_{30}), 15 (T_{60}), 13 (T_{15}), or 11 (T_{120}); *female CR,* n=11 (T_{15}, T_{30}, T_{60},), or 10 (T_0, T_{120}). **(E)**: *male AL,* n=9; *female AL,* n=9; *male CR,* n=16; *female CR,* n=9. In (A), (B), and (D), significant effects of diet, sex and/or time, and interactions thereof, were determined using a mixed-effects model. In (C) and (E), significant effects of sex, diet, and sex-diet interaction for tAUC, iAUC, HOMA-IR and Matsuda Index were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown with each graph. For (C) and (E), statistically significant differences between comparable groups are indicated by *** ($P<0.001$) or with *P* values shown. Source data are provided as a Source Data file. See also Supplementary Figure 6.

Figure 4 – CR decreases energy expenditure in male and female mice, with a trend for greater nighttime decreases in females.

Male and female mice were fed AL or CR diets as described for Figure 2. At 12-13 or 14-15 weeks of age mice were housed individually in TSE PhenoMaster cages for 3 days for indirect calorimetry. **(A,B)** Absolute energy expenditure (watts, W) was recorded every 15 minutes throughout the 3 days. **(A)** Average energy expenditure per hour over the 24 h light (Day) and dark (Night) period, spanning days 2-3 of PhenoMaster housing. **(B)** Energy expenditure (kJ) during the Day, Night, or Day + Night (Total). **(C)** Energy expenditure normalised to lean body mass (kJ/g) during the Day, Night, or Day + Night (Total). **(D)** Linear regression of lean mass vs Total (Day + Night) energy expenditure (kJ/24 h). Data in (A-D) are from 13 (*male AL*) or 14 (*male CR, female AL, female CR)* mice per group. In (A), data are shown as mean ± SEM; significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA, with *P* values shown beneath the graph. In (B-C), data are shown as violin plots overlaid with individual data points; within each time period (Day, Night, or Total), significant effects of sex, diet, and diet*sex interaction were assessed using 2-way ANOVA, with *P* values shown beneath each graph. Statistically significant differences between comparable groups were further assessed by Tukey's multiple

comparisons tests and are indicated by ** (*P*<0.01) or *** (*P*<0.001). For linear regression in (D), ANCOVA was used to test if the relationship between lean mass and total energy expenditure differs significantly across all of the individual diet-sex groups (*i. All mice*); between AL and CR mice, irrespective of sex (*ii. AL vs CR (M+F)*); and between males and females fed AL diet (*iii*) or CR diet (*iv*) only. ANCOVA *P* values for differences in slope and intercept are reported beneath the graph.

Figure 5 – The effects of CR on glucose uptake do not differ between male and female mice. Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. At 15 weeks of age, glucose uptake was assessed by PET/CT with [18]F-fluorodeoxyglucose ([18]F-FDG). **(A-C)** [18]F-FDG uptake in the indicated tissues was determined as SUVs from PET/CT scans (A) or as % injected dose per gram of tissue (%ID/g) from gamma counting of dissected whole tissues (C). **(B)** PET/CT images confirm that [18]F-FDG uptake in interscapular BAT is greater in females than in males, regardless of diet. Data are shown as violin plots (A,C) or representative images (B) of 7 (*male AL, female AL, male CR*) or 6 (*female CR*) mice per group. For each tissue in (A) and (C), significant effects of sex, diet, and sex-diet interaction were assessed by 2-way ANOVA; overall *P* values for each variable, and their interactions, are shown beneath each graph. Significant differences between each group, as determined using Tukey's multiple comparisons test, are indicated by * (*P* <0.05). Source data are provided as a Source Data file. See also Supplementary Figure 7.

Figure 6 – CR exerts sexually dimorphic effects on hepatic gene expression and sphingolipid content.
Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. Livers were sampled at necropsy (15 weeks) and used to assess mRNA transcripts, triglyceride content, and ceramide and dihydroceramide content. **(A)** Expression of the indicated transcripts was determined by qPCR. Transcript expression was normalised to the geometric mean of *Tbp, Rn18s* and *Ppia* expression and is presented relative to the highest group average. **(B)** Principal component analysis of transcript expression in (A). **(C)** Hepatic triglyceride content. **(D,E)** LC-MS analysis of total, 22:0 and 23:0 ceramides (D), dihydroceramides (E, left side) and the ceramide:DHC ratio (E, right side). **(F)** Principal component analysis of ceramide and dihydroceramide content based on data in (D-E) and Supplementary Figure 8. Data in (A), (C), (D), and (E) are presented as truncated violin plots overlaid with individual data points. Data represent the following numbers of mice per group. **(A-B)**: *male AL,* n=6; *female AL,* n=6; *male CR,* n=6; *female CR,* n=6. **(C)**: *male AL,* n=9; *female AL,* n=8; *male CR,* n=9; *female CR,* n=8. **(D-F):** 6 mice per group. For (A), (C), (D) and (E), significant effects of sex, diet, and sex-diet interaction were assessed using 2-way ANOVA; overall *P* values for each variable, and their interactions, are shown beneath each graph. Significant differences between comparable groups were assess using Tukey's multiple comparison test and are indicated by * (*P*<0.05), ** (*P*<0.01) or *** (*P*<0.001). Source data are provided as a Source Data file. See also Supplementary Figure 8.

Figure 7 – Sex differences in CR-induced weight loss and fat loss are absent when CR is initiated in aged mice.

Male and female C57BL6/NCrl mice were single housed and fed AL or a CR diet (70% of daily AL intake) from 78-84 weeks of age. **(A-F)** Each week mice were weighed (A,B) and body composition was determined by TD-NMR (C-F). Body mass, fat mass and lean mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). **(G)** Masses of gWAT, iWAT, mWAT, pWAT, pcWAT and BAT were recorded at necropsy and are shown as % body mass. Data in (A-F) are shown as mean ±SEM. Data in (G) are shown as violin plots overlaid with individual data points. For each group and timepoint, the following numbers of mice were used: **(A,B)**: *male AL*, n=9 (Wk 0, Wk 2, Wk3, Wk 4), 8 (Wk 1), or 6 (Wk 5, Wk 6); *female AL,* n=13 (Wk 0, Wk 2), 12 (Wk 4, Wk 5), 11 (Wk 6), 9 (Wk 1, or 8 (Wk 3); *male CR,* n=10 (Wk 0, Wk 2-4), 9 (Wk 1), 7 (Wk 5), or 6 (Wk 6); *female CR,* n=14 (Wk 0, Wk 4, Wk 5), 13 (Wk 2, Wk 6), 10 (Wk 3), or 9 (Wk 1). **(C-F)**: *male AL,* n=9 (Wk 0, Wk 2), 8 (Wk 3, Wk 4, Wk 6), or 7 (Wk 1, Wk 5); *female AL,* n=13 (Wk 0, Wk 2), 9 (Wk 4, Wk 6), 8 (Wk 5), 6 (Wk 1), or 5 (Wk 3); *male CR,* n=10 (Wk 0, Wk 2), 9 (Wk3-6), or 8 (Wk 1); *female CR,* n=14 (Wk 1, Wk 2), 11 (Wk 4-6), 7 (Wk 3), or 6 (Wk 1). **(G)**: *male AL,* n=9; *female AL,* n=12 (iWAT, mWAT) or 11 (gWAT, pWAT); *male CR,* n=10 (iWAT, pWAT), or 9 (gWAT, mWAT); *female CR,* n=14 (mWAT, pWAT), 13 (iWAT), or 12 (gWAT). For (A-F), significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA or a mixed-effects model. For (G), significant effects of sex, diet, and sex-diet interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown beneath each graph. For (G), significant differences between comparable groups are indicated by ** (*P*<0.01) or *** (*P*<0.001). Source data are provided as a Source Data file. See also Supplementary Figures 9 and 10.

Figure 8 – Sex differences in CR's effects on glucose homeostasis are largely absent when CR is initiated in aged mice.

Male and female C57BL6/NCrl mice were fed AL or CR diet from 78-84 weeks of age, as described for Figure 7. **(A)** Random-fed blood glucose was recorded each week. **(B-D)** At 82 weeks of age mice underwent OGTTs. Blood glucose was recorded at each timepoint (B) and the tAUC and iAUC was determined (C). (D) Glucose-stimulated insulin secretion in mice during OGTT was assessed using an insulin ELISA. **(E)** HOMA-IR and Matsuda indices of mice calculated from glucose and insulin concentrations during the OGTT. Data in (A), (B), and (D) are presented as mean ± SEM. Data in (C) and (E) are presented as violin plots overlaid with individual data points. For each group and timepoint the following numbers of mice were used: **(A)**: *male AL,* n=9 (Wk 0, Wk 2-4, Wk 6), 8 (Wk 1), or 5 (Wk 5); *female AL,* n=12 (Wk 0, Wk2, Wk4), 8 (Wk 1, Wk 3), or 11 (Wk 5, Wk 6); *male CR,* n=10 (Wk 0, Wk 2-4, Wk 6), 9 (Wk 1), or 7 (Wk 5); *female CR,* n= 14 (Wk 0, Wk 2, Wk 4-6), 9 (Wk 1) or 10 (Wk 3). **(B-E)**: *male AL,* n=9 (B-C) or 5 (D-E); *female AL,* n=12; *male CR,* n=10; *female CR,* n=14. In (A), (B), and (D), significant effects of diet, sex and/or time, and interactions thereof, were determined using a mixed-effects model or 3-way ANOVA. In (C) and (E), significant effects of sex, diet, and sex-diet interaction for tAUC, iAUC, HOMA-IR and Matsuda Index were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown with each graph. For (C) and (E), statistically significant differences

1221   between comparable groups are indicated by * ($P<0.05$) or *** ($P<0.001$), or with P
1222   values shown. Source data are provided as a Source Data file.
1223
Figure 9 – Effects of CR on body mass and body composition in humans are
sex- and age-dependent.
Twenty male and twenty-two female volunteers participated in a weight loss study
involving a 4-week dietary intervention. **(A-F)** Body mass was recorded each week
(A,B)**.** Fat mass (FM) and fat-free mass (FFM) were measured by air-displacement
whole-body plethysmography at weeks 0 and 4 (C-F). Body mass, fat mass and fat-
free mass are shown as absolute masses (A,C,E) or fold-change relative to baseline
(B,D,F). Data are presented as mean ± SEM**.** For (A-C) and (E), significant effects of
time, sex, and time*sex interaction were assessed using 2-way ANOVA. In (B), (D)
and (F), significant differences between males and females at each time point were
determined by Šídák's multiple comparisons test (B) or unpaired T-test (D,F) and are
indicated by * ($P<0.05$) or ** ($P<0.01$). **(G-I)** Simple linear regression of age vs fold-
change (*week 4 vs week 0)* in body mass (G), fat mass (H) and fat-free mass (I). For
each sex, significant associations between age and outcome (fold-change) are
indicated beneath each graph as '*P,* Slope'. ANCOVA was further used to test if the
age-outcome relationship differs significantly between males and females. ANCOVA
results are reported beneath each graph as '*P,* Slope' and '*P,* Intercept' for males vs
females (M vs F). In (G), similar slopes but different intercepts show that sex
significantly influences weight loss, but the influence of age does not differ between
the sexes. In (H,I) the slopes differ significantly, indicating that the age-outcome
relationship differs between the sexes. **(J-L)** Fold-change (week 4 vs week 0) in body
mass (J), fat mass (K) and fat-free mass (L) for males vs females separated into
younger (<45 years) and older (>45 years) groups. Data are presented as violin plots
overlaid with individual data points. Significant effects of age, sex, and age*sex
interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons
test. Overall *P* values for each variable, and their interactions, are shown beneath each
graph. Significant differences between comparable groups are indicated by * ($P<0.05$)
or *** ($P<0.001$). Source data are provided as a Source Data file. See also Table 3
and Supplementary Figures 10 and 11.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1, Related to Figure 1 – Methods used and additional outcomes relating to the comprehensive review of the CR literature.
The NIH PubMed database was searched using MeSH terms to identify research articles that studied CR *in vivo* in mice or humans, published between 2003 and 2021. **(A)** Method of database search presented as a flow diagram, including search terms, reasons for exclusion, and number of results in each category. **(B)** Percentage of mouse or human publications for each year, with publications classified into each group as described for Figure 1. Source data for (B) are provided as a Source Data file.

Supplementary Figure 2, Related to Figure 2 – The effect of CR on percent fat and lean mass differs between male and female mice.
Male and female mice were fed AL or CR diets and body composition was assessed weekly, as described for Figure 2. **(A,B)** Fat mass (A) and lean mass (B) were determined by TD-NMR and calculated as % body mass. Data are shown as mean ±SEM of the following numbers of mice per group: *male AL*, n=42; *female AL,* n=43; *male CR,* n=44; *female CR,* n=52. Significant effects of diet, sex and/or duration were determined by 3-way ANOVA (A) or a mixed-effects model (B); *P* values are shown beneath each graph. Source data are provided as a Source Data file.

Supplementary Figure 3, Related to Figure 2 – CR and sex effects on the masses of adipose and non-adipose tissues.
Male and female mice were fed AL or CR diets, as described for Figure 2. After six weeks of CR (15 weeks of age) tissues were sampled and weighed. **(A-C)** Violin plots, overlaid with individual data points, showing the absolute mass (A,B) or % body mass (C) of each tissue; Gastroc = gastrocnemius muscle. For (A), the numbers of mice per group and tissue type are as reported for Figure 2G. For (B) and (C), the following numbers of mice were used for each group and tissue type: *male AL,* n=29 (liver, spleen), 23 (pancreas, kidneys, heart, thymus), 22 (adrenals) or 16 (gastroc.); *female AL,* n=33 (liver, spleen), 27 (pancreas, kidneys, heart, thymus, adrenals) or 16 (gastroc.); *male CR,* n=28 (liver, spleen), 22 (pancreas, kidneys, heart, thymus, adrenals) or 14 (gastroc.); *female CR,* n=34 (liver, spleen), 28 (pancreas, kidneys, heart, thymus, adrenals) or 15 (gastroc.). For each tissue, significant effects of diet and/or sex were determined by 2-way ANOVA with Tukey's multiple comparisons tests and are reported as described for Figure 2. Source data are provided as a Source Data file.

Supplementary Figure 4, Related to Figure 2 – CR decreases mWAT adipocyte size in male but not in female mice.
Male and female mice were fed AL or CR diets as described for Figure 2. After six weeks of CR (15 weeks of age) mWAT was sampled and processed for histological analysis. **(A,B)** Micrographs of H&E-stained sections of mWAT (A) were used for histomorphometric analysis of adipocyte area (B); in (A), scale bar = 100 μm. Data in are shown as representative micrographs (A) or violin plots (B) from 5 (*male AL, male CR)* or 6 (*female AL, female CR)* mice per group. For (B), significant effects of diet

and/or sex on adipocyte area (Ad.Ar) were determined by 3-way ANOVA, with *P* values shown beneath the graph. Significant differences in median Ad.Ar between AL and CR mice were further determined 2-way ANOVA with Šídák's multiple comparisons test; * = *P*<0.05. Source data are provided as a Source Data file.

Supplementary Figure 5, Related to Figure 2 – Sex differences in the effects of CR on body mass and composition persist when CR mice are fed in the evening. Male and female C57BL6/NCrl mice were single housed and fed AL or CR diet (70% of daily AL intake) from 9-13 weeks of age, with CR mice receiving their daily diet ration between 18:00-19:00. **(A-F)** Each week mice were weighed (A,B) and body composition was determined by TD-NMR (C-F). Body mass, fat mass and lean mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). **(G,H)** Masses of the indicated tissues were recorded at necropsy and are shown as % body mass. Data in (A-H) represent 5 (*male AL, female AL*) or 6 (*male CR, female CR*) mice per group and are shown as mean ±SEM (A-F) or as truncated volcano plots overlaid with individual data points (G,H). For (A-F), significant effects of diet, sex or time, and interactions thereof, were determined by 3-way ANOVA. For (G-H), significant effects of sex, diet, and sex-diet interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown beneath each graph. For (G-H), statistically significant differences between comparable groups are indicated by ** (*P*<0.01). Source data are provided as a Source Data file.

Supplementary Figure 6, Related to Figure 3 – Sex differences in the effects of CR on glucose homeostasis persist when CR mice are fed in the evening. Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Supplementary Figure 5. **(A,B)** At 13 weeks of age mice underwent OGTTs. Blood glucose was recorded at each timepoint (A) and the tAUC and iAUC was determined (B). Data represent 5 (*male AL, female AL*) or 6 (*male CR, female CR*) mice per group and are shown as mean ±SEM (A) or as truncated volcano plots overlaid with individual data points (B). In (A), significant effects of diet, sex or time, and interactions thereof, were determined by 3-way ANOVA. In (B), significant effects of sex, diet, and sex-diet interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown beneath each graph. For (B), statistically significant differences between comparable groups are indicated by ** (*P*<0.01). Source data are provided as a Source Data file.

Supplementary Figure 7, Related to Figure 5 – The effects of CR on glucose uptake in peripheral tissues. Male and female C57BL6/NCrl mice were fed AL or CR diet and glucose uptake was assessed by ^{18}F-FDG PET/CT, as described for Figure 5. **(A,B)** ^{18}F-FDG uptake in the indicated tissues was determined by PMOD analysis of PET/CT scans (A) or gamma counting of dissected whole tissues (B). Data in (A) and (B) are presented as truncated violin plots, overlaid with individual data points, from 7 (*male AL, female AL, male CR*) or 6 (*female CR*) mice per group; in (B), data for spleen are from 6 mice per group

1352 only. Statistical significance was determined as described for Figure 5. Source data
1353 are provided as a Source Data file.
1354
1355
Supplementary Figure 8, Related to Figure 6 – CR exerts sexually dimorphic
effects on hepatic sphingolipid content.
Male and female mice were fed AL or CR diets as described for Figure 2. **(A-C)** At
necropsy, livers were sampled for analysis of ceramide and dihydroceramide content
by LC-MS. Data show the concentration of each species of ceramide (A),
dihydroceramide (B), or dihydroceramide:ceramide ratio (C) and are presented as
truncated violin plots showing all points. The following numbers of mice were used per
group and species: *male AL*, n=6 (all except: DHC 20:1 & 26:0, Cer:DHC 20:1 & 26:0),
1 (DHC & Cer:DHC 20:1), or 0 (DHC & Cer:DHC 26:0); *female AL*, n=6; *male CR*, n=6
(all except DHC & Cer:DHC 26:0), or 1 (DHC & Cer:DHC 26:0); *female CR*, n=6. For
each species, significant effects of sex, diet, and sex-diet interaction were assessed
using 2-way ANOVA; overall *P* values for each variable, and their interactions, are
shown beneath each graph. Significant differences between comparable groups were
assess using Tukey's multiple comparison test and are indicated by * ($P<0.05$), **
($P<0.01$) or *** ($P<0.001$). Source data are provided as a Source Data file.
1371
1372
Supplementary Figure 9, Related to Figure 7 – CR and sex effects on the masses
of adipose and non-adipose tissues in aged mice.
Male and female mice were fed AL or CR diets from 78-84 weeks of age, as described
for Figure 7. After six weeks of CR (84 weeks of age) tissues were sampled and
weighed. **(A-C)** Violin plots, overlaid with individual data points, showing the absolute
mass (A,B) or % body mass (C) of each tissue; Gastroc = gastrocnemius muscle. The
numbers of mice per group and tissue type are as reported for Figure 7G. For each
tissue, significant effects of diet and/or sex were determined by 2-way ANOVA with
Tukey's multiple comparisons test and are reported as described for Figure 7. Source
data are provided as a Source Data file.
1383
1384
Supplementary Figure 10, Related to Figures 2, 7 and 9 – CR-induced fat loss in
mice is age- and sex-dependent.
Male and female mice were fed AL or CR diets from 9-15 or 78-84 weeks of age and
fat mass was determined each week, as described for Figures 2 and 7. For each
mouse (CR and AL diets) the ratio of absolute fat masses (g) was calculated for week
4 vs week 0 (i.e. 13 vs 9 weeks of age for young mice; 82 vs 78 weeks for aged mice);
week 4 was selected for consistency with the 4-week duration of our human CR study
(Fig. 9). For the CR mice, the fold-change in fat mass, relative to AL mice, was then
determined as follows: i) the mean week 4:week 0 ratio was calculated for AL mice in
each age group; ii) within each sex and age group, the week 4:week 0 ratio for each
CR mouse was calculated relative to the mean week 4:week 0 ratio for the
corresponding AL group. The rationale for showing these "fold changes of fold
changes" is that the AL mice gain fat mass over the course of the study and therefore
we must determine how much the fat mass of each CR mouse has changed compared
to where we would expect it to be if it had continued on AL diet. Data in are shown as
violin plots of the following numbers of mice per group: *young female*, n=31; *young*
male, n=26; *old female*, n=11; *old male*, n=9. Significant effects of diet and/or sex were

determined by 2-way ANOVA with Tukey's multiple comparisons test and are reported
as described for Figure S3. Source data are provided as a Source Data file.
1404
1405
Supplementary Figure 11, Related to Figure 9 – Body fat percentage in human
CR participants.
Twenty male and twenty-two female volunteers participated in a weight loss study
involving a 4-week dietary intervention, as described for Figure 9. Total % fat mass at
week 0 is shown for males vs females separated into younger (<45 years) and older
(>45 years) groups. Data are presented as violin plots overlaid with individual data
points. Significant effects of age, sex, and age*sex interaction were assessed using
2-way ANOVA with Tukey's multiple comparisons test. Overall P values for each
variable, and their interactions, are shown beneath the graph. Significant differences
between comparable groups are indicated by ** ($P<0.01$) or *** ($P<0.001$). See also
Table 3. Source data are provided as a Source Data file.

REFERENCES

1. Lin SJ, Kaeberlein M, Andalis AA, Sturtz LA, Defossez PA, Culotta VC, et al. Calorie restriction extends Saccharomyces cerevisiae lifespan by increasing respiration. *Nature.* 2002;418(6895):344-8.
2. Weindruch R, Walford RL, Fligiel S, and Guthrie D. The retardation of aging in mice by dietary restriction: longevity, cancer, immunity and lifetime energy intake. *J Nutr.* 1986;116(4):641-54.
3. Mattison JA, Colman RJ, Beasley TM, Allison DB, Kemnitz JW, Roth GS, et al. Caloric restriction improves health and survival of rhesus monkeys. *Nat Commun.* 2017;8:14063.
4. Most J, Tosti V, Redman LM, and Fontana L. Calorie restriction in humans: An update. *Ageing Res Rev.* 2017;39:36-45.
5. Speakman JR, and Mitchell SE. Caloric restriction. *Mol Aspects Med.* 2011;32(3):159-221.
6. Barzilai N, Banerjee S, Hawkins M, Chen W, and Rossetti L. Caloric restriction reverses hepatic insulin resistance in aging rats by decreasing visceral fat. *J Clin Invest.* 1998;101(7):1353-61.
7. López-Otín C, Galluzzi L, Freije JMP, Madeo F, and Kroemer G. Metabolic Control of Longevity. *Cell.* 2016;166(4):802-21.
8. Mancuso P, and Bouchard B. The Impact of Aging on Adipose Function and Adipokine Synthesis. *Front Endocrinol (Lausanne).* 2019;10:137-.
9. Gabriely I, Ma XH, Yang XM, Atzmon G, Rajala MW, Berg AH, et al. Removal of visceral fat prevents insulin resistance and glucose intolerance of aging: an adipokine-mediated process? *Diabetes.* 2002;51(10):2951-8.
10. Muzumdar R, Allison DB, Huffman DM, Ma X, Atzmon G, Einstein FH, et al. Visceral adipose tissue modulates mammalian longevity. *Aging cell.* 2008;7(3):438-40.
11. Tran TT, Yamamoto Y, Gesta S, and Kahn CR. Beneficial effects of subcutaneous fat transplantation on metabolism. *Cell Metab.* 2008;7(5):410-20.
12. Larson-Meyer DE, Heilbronn LK, Redman LM, Newcomer BR, Frisard MI, Anton S, et al. Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care.* 2006;29(6):1337-44.
13. Mauvais-Jarvis F. Gender differences in glucose homeostasis and diabetes. *Physiol Behav.* 2017.
14. Oliva M, Muñoz-Aguirre M, Kim-Hellmuth S, Wucher V, Gewirtz ADH, Cotter DJ, et al. The impact of sex on gene expression across human tissues. *Science.* 2020;369(6509):eaba3066.
15. Valencak TG, Osterrieder A, and Schulz TJ. Sex matters: The effects of biological sex on adipose tissue biology and energy metabolism. *Redox Biology.* 2017;12:806-13.
16. Maggi A, and Della Torre S. Sex, metabolism and health. *Mol Metabol.* 2018;15:3-7.
17. Prendergast BJ, Onishi KG, and Zucker I. Female mice liberated for inclusion in neuroscience and biomedical research. *Neurosci Biobehav Rev.* 2014;40:1-5.
18. Kane AE, Sinclair DA, Mitchell JR, and Mitchell SJ. Sex differences in the response to dietary restriction in rodents. *Current Opinion in Physiology.* 2018;6:28-34.

19. Redman LM, Heilbronn LK, Martin CK, Alfonso A, Smith SR, Ravussin E, et al. Effect of calorie restriction with or without exercise on body composition and fat distribution. *J Clin Endocrinol Metab.* 2007;92(3):865-72.

20. Shi H, Strader AD, Woods SC, and Seeley RJ. Sexually dimorphic responses to fat loss after caloric restriction or surgical lipectomy. *Am J Physiol Endocrinol Metab.* 2007;293(1):E316-26.

21. Cawthorn WP, Scheller EL, Parlee SD, Pham HA, Learman BS, Redshaw CM, et al. Expansion of Bone Marrow Adipose Tissue During Caloric Restriction Is Associated With Increased Circulating Glucocorticoids and Not With Hypoleptinemia. *Endocrinology.* 2016;157(2):508-21.

22. Della Torre S, Mitro N, Meda C, Lolli F, Pedretti S, Barcella M, et al. Short-Term Fasting Reveals Amino Acid Metabolism as a Major Sex-Discriminating Factor in the Liver. *Cell Metab.* 2018;28(2):256-67 e5.

23. Lee SK. Sex as an important biological variable in biomedical research. *BMB reports.* 2018;51(4):167-73.

24. Johnson J, Sharman Z, Vissandjée B, and Stewart DE. Does a change in health research funding policy related to the integration of sex and gender have an impact? *PLoS One.* 2014;9(6):e99900-e.

25. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, and Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985;28(7):412-9.

26. Matsuda M, and DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care.* 1999;22(9):1462.

27. Zhang L, Xue X, Zhai R, Yang X, Li H, Zhao L, et al. Timing of Calorie Restriction in Mice Impacts Host Metabolic Phenotype with Correlative Changes in Gut Microbiota. *mSystems.* 2019;4(6):e00348-19.

28. Bartelt A, Koehne T, Todter K, Reimer R, Muller B, Behler-Janbeck F, et al. Quantification of Bone Fatty Acid Metabolism and Its Regulation by Adipocyte Lipoprotein Lipase. *Int J Mol Sci.* 2017;18(6).

29. Suchacki KJ, Tavares AAS, Mattiucci D, Scheller EL, Papanastasiou G, Gray C, et al. Bone marrow adipose tissue is a unique adipose subtype with distinct roles in glucose homeostasis. *Nat Commun.* 2020;11(1):3097.

30. Kersten S. Integrated physiology and systems biology of PPARα. *Mol Metabol.* 2014;3(4):354-71.

31. Petersen MC, and Shulman GI. Roles of Diacylglycerols and Ceramides in Hepatic Insulin Resistance. *Trends Pharmacol Sci.* 2017;38(7):649-65.

32. Chaurasia B, Tippetts TS, Mayoral Monibas R, Liu J, Li Y, Wang L, et al. Targeting a ceramide double bond improves insulin resistance and hepatic steatosis. *Science.* 2019;365(6451):386-92.

33. Apostolopoulou M, Gordillo R, Gancheva S, Strassburger K, Herder C, Esposito I, et al. Role of ceramide-to-dihydroceramide ratios for insulin resistance and non-alcoholic fatty liver disease in humans. *BMJ Open Diabetes Research & Care.* 2020;8(2):e001860.

34. Gee DM, Flurkey K, and Finch CE. Aging and the Regulation of Luteinizing Hormone in C57BL/6J Mice: Impaired Elevations after Ovariectomy and Spontaneous Elevations at Advanced Ages1. *Biol Reprod.* 1983;28(3):598-607.

35. Piotrowska K, Tarnowski M, Zgutka K, and Pawlik A. Gender Differences in Response to Prolonged Every-Other-Day Feeding on the Proliferation and Apoptosis of Hepatocytes in Mice. *Nutrients.* 2016;8(3):176-.

36. Musante GJ. The dietary rehabilitation clinic: Evaluative report of a behavioral and dietary treatment of obesity. *Behav Ther.* 1976;7(2):198-204.

37. O'Neil PM, Currey HS, Hirsch AA, Riddle FE, Taylor CI, Malcolm RJ, et al. Effects of sex of subject and spouse involvement on weight loss in a behavioral treatment program: a retrospective investigation. *Addict Behav.* 1979;4(2):167-77.

38. Mauriege P, Imbeault P, Langin D, Lacaille M, Almeras N, Tremblay A, et al. Regional and gender variations in adipose tissue lipolysis in response to weight loss. *J Lipid Res.* 1999;40(9):1559-71.

39. Wong MHT, Holst C, Astrup A, Handjieva-Darlenska T, Jebb SA, Kafatos A, et al. Caloric restriction induces changes in insulin and body weight measurements that are inversely associated with subsequent weight regain. *PLoS One.* 2012;7(8):e42858-e.

40. Azar M, Nikpay M, Harper M-E, McPherson R, and Dent R. Can response to dietary restriction predict weight loss after Roux-en-Y gastroplasty? *Obesity.* 2016;24(4):805-11.

41. Das SK, Roberts SB, Bhapkar MV, Villareal DT, Fontana L, Martin CK, et al. Body-composition changes in the Comprehensive Assessment of Long-term Effects of Reducing Intake of Energy (CALERIE)-2 study: a 2-y randomized controlled trial of calorie restriction in nonobese humans. *The American Journal of Clinical Nutrition.* 2017;105(4):913-27.

42. Cawthorn WP, Scheller EL, Learman BS, Parlee SD, Simon BR, Mori H, et al. Bone Marrow Adipose Tissue Is an Endocrine Organ that Contributes to Increased Circulating Adiponectin during Caloric Restriction. *Cell Metab.* 2014;20(2):368-75.

43. Dionne DA, Skovso S, Templeman NM, Clee SM, and Johnson JD. Caloric Restriction Paradoxically Increases Adiposity in Mice With Genetically Reduced Insulin. *Endocrinology.* 2016;157(7):2724-34.

44. Li X, Cope MB, Johnson MS, Smith DL, Jr., and Nagy TR. Mild calorie restriction induces fat accumulation in female C57BL/6J mice. *Obesity.* 2010;18(3):456-62.

45. Leenen R, van der Kooy K, Droop A, Seidell JC, Deurenberg P, Weststrate JA, et al. Visceral fat loss measured by magnetic resonance imaging in relation to changes in serum lipid levels of obese men and women. *Arterioscler Thromb.* 1993;13(4):487-94.

46. Fontana L, Klein S, and Holloszy JO. Effects of long-term calorie restriction and endurance exercise on glucose tolerance, insulin action, and adipokine production. *Age.* 2010;32(1):97-108.

47. Evans EM, Mojtahedi MC, Thorpe MP, Valentine RJ, Kris-Etherton PM, and Layman DK. Effects of protein intake and gender on body composition changes: a randomized clinical weight loss trial. *Nutr Metab (Lond).* 2012;9(1):55.

48. Kraus WE, Bhapkar M, Huffman KM, Pieper CF, Krupa Das S, Redman LM, et al. 2 years of calorie restriction and cardiometabolic risk (CALERIE): exploratory outcomes of a multicentre, phase 2, randomised controlled trial. *The Lancet Diabetes & Endocrinology.* 2019;7(9):673-83.

49. Dhahbi JM, Kim HJ, Mote PL, Beaver RJ, and Spindler SR. Temporal linkage between the phenotypic and genomic responses to caloric restriction. *Proc Natl Acad Sci U S A.* 2004;101(15):5524-9.

50. Sheng Y, Xia F, Chen L, Lv Y, Lv S, Yu J, et al. Differential Responses of White
Adipose Tissue and Brown Adipose Tissue to Calorie Restriction During Aging.
The Journals of Gerontology: Series A. 2020;76(3):393-9.
51. Rohrbach S, Aurich AC, Li L, and Niemann B. Age-associated loss in adiponectin-
activation by caloric restriction: lack of compensation by enhanced inducibility of
adiponectin paralogs CTRP2 and CTRP7. *Mol Cell Endocrinol.* 2007;277(1-2):26-
34.
52. Mitchell SJ, Madrigal-Matute J, Scheibye-Knudsen M, Fang E, Aon M, González-
Reyes JA, et al. Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging
in Mice. *Cell Metab.* 2016;23(6):1093-112.
53. Forney LA, Stone KP, Gibson AN, Vick AM, Sims LC, Fang H, et al. Sexually
Dimorphic Effects of Dietary Methionine Restriction are Dependent on Age when
the Diet is Introduced. *Obesity.* 2020;28(3):581-9.
54. Chaix A, Deota S, Bhardwaj R, Lin T, and Panda S. Sex- and age-dependent
outcomes of 9-hour time-restricted feeding of a Western high-fat high-sucrose
diet in C57BL/6J mice. *Cell reports.* 2021;36(7).
55. Brandhorst S, Choi In Y, Wei M, Cheng Chia W, Sedrakyan S, Navarrete G, et al.
A Periodic Diet that Mimics Fasting Promotes Multi-System Regeneration,
Enhanced Cognitive Performance, and Healthspan. *Cell Metab.* 2015;22(1):86-
99.
56. Porter MH, Fine JB, Cutchins AG, Bai Y, and DiGirolamo M. Sexual dimorphism
in the response of adipose mass and cellularity to graded caloric restriction. *Obes
Res.* 2004;12(1):131-40.
57. Valle A, Catala-Niell A, Colom B, Garcia-Palmer FJ, Oliver J, and Roca P. Sex-
related differences in energy balance in response to caloric restriction. *Am J
Physiol Endocrinol Metab.* 2005;289(1):E15-22.
58. Valle A, Garcia-Palmer FJ, Oliver J, and Roca P. Sex differences in brown
adipose tissue thermogenic features during caloric restriction. *Cellular physiology
and biochemistry : international journal of experimental cellular physiology,
biochemistry, and pharmacology.* 2007;19(1-4):195-204.
59. Hill JO, Talano CM, Nickel M, and DiGirolamo M. Energy utilization in food-
restricted female rats. *J Nutr.* 1986;116(10):2000-12.
60. Hill JO, Latiff A, and DiGirolamo M. Effects of variable caloric restriction on
utilization of ingested energy in rats. *Am J Physiol.* 1985;248(5 Pt 2):R549-59.
61. Fenton JI, Nunez NP, Yakar S, Perkins SN, Hord NG, and Hursting SD. Diet-
induced adiposity alters the serum profile of inflammation in C57BL/6N mice as
measured by antibody array. *Diabetes, obesity & metabolism.* 2009;11(4):343-
54.
62. Varady KA, Allister CA, Roohk DJ, and Hellerstein MK. Improvements in body fat
distribution and circulating adiponectin by alternate-day fasting versus calorie
restriction. *The Journal of Nutritional Biochemistry.* 2010;21(3):188-95.
63. Harvey AE, Lashinger LM, Otto G, Nunez NP, and Hursting SD. Decreased
systemic IGF-1 in response to calorie restriction modulates murine tumor cell
growth, nuclear factor-kappaB activation, and inflammation-related gene
expression. *Mol Carcinog.* 2013;52(12):997-1006.
64. Zgheib S, Mequinion M, Lucas S, Leterme D, Ghali O, Tolle V, et al. Long-term
physiological alterations and recovery in a mouse model of separation associated
with time-restricted feeding: a tool to study anorexia nervosa related
consequences. *PLoS One.* 2014;9(8):e103775.

65. Janssen I, and Ross R. Effects of sex on the change in visceral, subcutaneous
adipose tissue and skeletal muscle in response to weight loss. *Int J Obes Relat*
Metab Disord. 1999;23(10):1035-46.
66. Wirth A, and Steinmetz B. Gender differences in changes in subcutaneous and
intra-abdominal fat during weight reduction: an ultrasound study. *Obes Res.*
1998;6(6):393-9.
67. Parker B, Noakes M, Luscombe N, and Clifton P. Effect of a high-protein, high-
monounsaturated fat weight loss diet on glycemic control and lipid levels in type
2 diabetes. *Diabetes Care.* 2002;25(3):425-30.
68. Weiss EP, Racette SB, Villareal DT, Fontana L, Steger-May K, Schechtman KB,
et al. Improvements in glucose tolerance and insulin action induced by increasing
energy expenditure or decreasing energy intake: a randomized controlled trial.
Am J Clin Nutr. 2006;84(5):1033-42.
69. Diaz Brinton R. Minireview: translational animal models of human menopause:
challenges and emerging opportunities. *Endocrinology.* 2012;153(8):3571-8.
70. Chen JH, Ouyang C, Ding Q, Song J, Cao W, and Mao L. A Moderate Low-
Carbohydrate Low-Calorie Diet Improves Lipid Profile, Insulin Sensitivity and
Adiponectin Expression in Rats. *Nutrients.* 2015;7(6):4724-38.
71. Zhu M, Miura J, Lu LX, Bernier M, DeCabo R, Lane MA, et al. Circulating
adiponectin levels increase in rats on caloric restriction: the potential for insulin
sensitization. *Exp Gerontol.* 2004;39(7):1049-59.
72. Mitchell SE, Tang Z, Kerbois C, Delville C, Konstantopedos P, Bruel A, et al. The
effects of graded levels of calorie restriction: I. impact of short term calorie and
protein restriction on body composition in the C57BL/6 mouse. *Oncotarget.*
2015;6(18):15902-30.
73. Mitchell SE, Delville C, Konstantopedos P, Hurst J, Derous D, Green C, et al. The
effects of graded levels of calorie restriction: II. Impact of short term calorie and
protein restriction on circulating hormone levels, glucose homeostasis and
oxidative stress in male C57BL/6 mice. *Oncotarget.* 2015;6(27):23213-37.
74. Mitchell SE, Tang Z, Kerbois C, Delville C, Derous D, Green CL, et al. The effects
of graded levels of calorie restriction: VIII. Impact of short term calorie and protein
restriction on basal metabolic rate in the C57BL/6 mouse. *Oncotarget.*
2017;8(11):17453-74.
75. Derous D, Mitchell SE, Green CL, Wang Y, Han JDJ, Chen L, et al. The Effects
of Graded Levels of Calorie Restriction: X. Transcriptomic Responses of
Epididymal Adipose Tissue. *J Gerontol A Biol Sci Med Sci.* 2017.
76. Ballor DL, and Poehlman ET. Exercise-training enhances fat-free mass
preservation during diet-induced weight loss: a meta-analytical finding. *Int J Obes*
Relat Metab Disord. 1994;18(1):35-40.
77. Bruss MD, Khambatta CF, Ruby MA, Aggarwal I, and Hellerstein MK. Calorie
restriction increases fatty acid synthesis and whole body fat oxidation rates. *Am*
J Physiol Endocrinol Metab. 2010;298(1):E108-16.
78. Arner P, Andersson DP, Backdahl J, Dahlman I, and Ryden M. Weight Gain and
Impaired Glucose Metabolism in Women Are Predicted by Inefficient
Subcutaneous Fat Cell Lipolysis. *Cell Metab.* 2018;28(1):45-54 e3.
79. Gazdag AC, Dumke CL, Kahn CR, and Cartee GD. Calorie restriction increases
insulin-stimulated glucose transport in skeletal muscle from IRS-1 knockout mice.
Diabetes. 1999;48(10):1930-6.

80. Johnson ML, Distelmaier K, Lanza IR, Irving BA, Robinson MM, Konopka AR, et
al. Mechanism by Which Caloric Restriction Improves Insulin Sensitivity in
Sedentary Obese Adults. *Diabetes.* 2016;65(1):74-84.

81. Lee YK, Park JE, Lee M, and Hardwick JP. Hepatic lipid homeostasis by
peroxisome proliferator-activated receptor gamma 2. *Liver Research.*
2018;2(4):209-15.

82. Abdul-Wahed A, Guilmeau S, and Postic C. Sweet Sixteenth for ChREBP:
Established Roles and Future Goals. *Cell Metab.* 2017;26(2):324-41.

83. Hou W-L, Yin J, Alimujiang M, Yu X-Y, Ai L-G, Bao Y-q, et al. Inhibition of
mitochondrial complex I improves glucose metabolism independently of AMPK
activation. *J Cell Mol Med.* 2018;22(2):1316-28.

84. Syamsunarno MRAA, Iso T, Hanaoka H, Yamaguchi A, Obokata M, Koitabashi
N, et al. A Critical Role of Fatty Acid Binding Protein 4 and 5 (FABP4/5) in the
Systemic Response to Fasting. *PLoS One.* 2013;8(11):e79386.

85. Sato T, Morita A, Mori N, and Miura S. The Roles of GPD1 in Maintaining Blood
Glucose Levels under Fasting Conditions. *The FASEB Journal.*
2016;30(S1):628.6-.6.

86. Liu Y, Basty N, Whitcher B, Bell JD, Sorokin EP, van Bruggen N, et al. Genetic
architecture of 11 organ traits derived from abdominal MRI using deep learning.
eLife. 2021;10:e65554.

87. Perry RJ, Peng L, Cline GW, Wang Y, Rabin-Court A, Song JD, et al. Mechanisms
by which a Very-Low-Calorie Diet Reverses Hyperglycemia in a Rat Model of
Type 2 Diabetes. *Cell Metab.* 2018;27(1):210-7.e3.

88. Taylor R, Al-Mrabeh A, Zhyzhneuskaya S, Peters C, Barnes AC, Aribisala BS, et
al. Remission of Human Type 2 Diabetes Requires Decrease in Liver and
Pancreas Fat Content but Is Dependent upon Capacity for β Cell Recovery. *Cell*
Metab. 2018;28(4):547-56.e3.

89. Norheim F, Bjellaas T, Hui ST, Chella Krishnan K, Lee J, Gupta S, et al. Genetic,
dietary, and sex-specific regulation of hepatic ceramides and the relationship
between hepatic ceramides and IR [S]. *J Lipid Res.* 2018;59(7):1164-74.

90. Turpin SM, Nicholls HT, Willmes DM, Mourier A, Brodesser S, Wunderlich CM, et
al. Obesity-induced CerS6-dependent C16:0 ceramide production promotes
weight gain and glucose intolerance. *Cell Metab.* 2014;20(4):678-86.

91. Raichur S, Wang ST, Chan PW, Li Y, Ching J, Chaurasia B, et al. CerS2
haploinsufficiency inhibits β-oxidation and confers susceptibility to diet-induced
steatohepatitis and insulin resistance. *Cell Metab.* 2014;20(4):687-95.

92. Raichur S, Brunner B, Bielohuby M, Hansen G, Pfenninger A, Wang B, et al. The
role of C16:0 ceramide in the development of obesity and type 2 diabetes: CerS6
inhibition as a novel therapeutic approach. *Molecular metabolism.* 2019;21:36-
50.

93. Holland WL, Brozinick JT, Wang LP, Hawkins ED, Sargent KM, Liu Y, et al.
Inhibition of ceramide synthesis ameliorates glucocorticoid-, saturated-fat-, and
obesity-induced insulin resistance. *Cell Metab.* 2007;5(3):167-79.

94. Zhang Q-J, Holland WL, Wilson L, Tanner JM, Kearns D, Cahoon JM, et al.
Ceramide Mediates Vascular Dysfunction in Diet-Induced Obesity by PP2A-
Mediated Dephosphorylation of the eNOS-Akt Complex. *Diabetes.*
2012;61(7):1848-59.

95. Mauvais-Jarvis F, Arnold AP, and Reue K. A Guide for the Design of Pre-clinical
Studies on Sex Differences in Metabolism. *Cell Metab.* 2017;25(6):1216-30.

96. Link JC, Wiese CB, Chen X, Avetisyan R, Ronquillo E, Ma F, et al. X chromosome dosage of histone demethylase KDM5C determines sex differences in adiposity. *The Journal of Clinical Investigation.* 2020;130(11):5688-702.

97. Dearden L, Bouret SG, and Ozanne SE. Sex and gender differences in developmental programming of metabolism. *Mol Metabol.* 2018;15:8-19.

98. Shen T, Ding L, Ruan Y, Qin W, Lin Y, Xi C, et al. SIRT1 Functions as an Important Regulator of Estrogen-Mediated Cardiomyocyte Protection in Angiotensin II-Induced Heart Hypertrophy. *Oxid Med Cell Longev.* 2014;2014:713894.

99. Wei Y, and Huang J. Role of estrogen and its receptors mediated-autophagy in cell fate and human diseases. *J Steroid Biochem Mol Biol.* 2019;191:105380.

100. Gupte AA, Pownall HJ, and Hamilton DJ. Estrogen: An Emerging Regulator of Insulin Action and Mitochondrial Function. *J Diabetes Res.* 2015;2015:916585.

101. Klinge CM. Estrogenic control of mitochondrial function. *Redox Biol.* 2020;31:101435.

102. Wheatley KE, Nogueira LM, Perkins SN, and Hursting SD. Differential effects of calorie restriction and exercise on the adipose transcriptome in diet-induced obese mice. *J Obes.* 2011;2011:265417.

103. Dakin RS, Walker BR, Seckl JR, Hadoke PW, and Drake AJ. Estrogens protect male mice from obesity complications and influence glucocorticoid metabolism. *Int J Obes (Lond).* 2015;39(10):1539-47.

104. Zhu L, Martinez MN, Emfinger CH, Palmisano BT, and Stafford JM. Estrogen signaling prevents diet-induced hepatic insulin resistance in male mice with obesity. *Am J Physiol Endocrinol Metab.* 2014;306(10):E1188-E97.

105. Tao Z, Zheng LD, Smith C, Luo J, Robinson A, Almeida FA, et al. Estradiol signaling mediates gender difference in visceral adiposity via autophagy. *Cell death & disease.* 2018;9(3):309.

106. Xie B, Pan D, Liu H, Liu M, Shi X, Chu X, et al. Diosmetin Protects Against Obesity and Metabolic Dysfunctions Through Activation of Adipose Estrogen Receptors in Mice. *Molecular Nutrition & Food Research.* 2021;n/a(n/a):2100070.

107. Guillaume M, Handgraaf S, Fabre A, Raymond-Letron I, Riant E, Montagner A, et al. Selective Activation of Estrogen Receptor α Activation Function-1 Is Sufficient to Prevent Obesity, Steatosis, and Insulin Resistance in Mouse. *Am J Pathol.* 2017;187(6):1273-87.

108. Yan H, Yang W, Zhou F, Li X, Pan Q, Shen Z, et al. Estrogen Improves Insulin Sensitivity and Suppresses Gluconeogenesis via the Transcription Factor Foxo1. *Diabetes.* 2019;68(2):291-304.

109. Bryzgalova G, Gao H, Ahren B, Zierath JR, Galuska D, Steiler TL, et al. Evidence that oestrogen receptor-alpha plays an important role in the regulation of glucose homeostasis in mice: insulin sensitivity in the liver. *Diabetologia.* 2006;49(3):588-97.

110. Handelsman DJ, and Wartofsky L. Requirement for Mass Spectrometry Sex Steroid Assays in the Journal of Clinical Endocrinology and Metabolism. *The Journal of Clinical Endocrinology & Metabolism.* 2013;98(10):3971-3.

111. Kane AE, Hilmer SN, Boyer D, Gavin K, Nines D, Howlett SE, et al. Impact of Longevity Interventions on a Validated Mouse Clinical Frailty Index. *The Journals of Gerontology: Series A.* 2016;71(3):333-9.

112. Ferland G, Doucet I, and Mainville D. Phylloquinone and Menaquinone-4 Tissue Distribution at Different Life Stages in Male and Female Sprague-Dawley Rats

Fed Different VK Levels Since Weaning or Subjected to a 40% Calorie Restriction
since Adulthood. *Nutrients.* 2016;8(3):141.
113. Austad SN, and Bartke A. Sex Differences in Longevity and in Responses to Anti-
Aging Interventions: A Mini-Review. *Gerontology.* 2016;62(1):40-6.
114. Bohler H, Jr., Mokshagundam S, and Winters SJ. Adipose tissue and
reproduction in women. *Fertil Steril.* 2010;94(3):795-825.
115. Misra M, and Klibanski A. Endocrine consequences of anorexia nervosa. *The
lancet Diabetes & endocrinology.* 2014;2(7):581-92.
116. Johnstone AM, Murison SD, Duncan JS, Rance KA, and Speakman JR. Factors
influencing variation in basal metabolic rate include fat-free mass, fat mass, age,
and circulating thyroxine but not sex, circulating leptin, or triiodothyronine. *Am J
Clin Nutr.* 2005;82(5):941-8.
117. McCrory MA, Gomez TD, Bernauer EM, and Molé PA. Evaluation of a new air
displacement plethysmograph for measuring human body composition. *Med Sci
Sports Exerc.* 1995;27(12):1686-91.
118. Galarraga M, Campión J, Muñoz-Barrutia A, Boqué N, Moreno H, Martínez JA, et
al. Adiposoft: automated software for the analysis of white adipose tissue
cellularity in histological sections. *J Lipid Res.* 2012;53(12):2791-6.
119. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al.
Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. *Genome biology.*
2002;3(7):RESEARCH0034.
120. Folch J, Lees M, and Sloane Stanley GH. A simple method for the isolation and
purification of total lipides from animal tissues. *J Biol Chem.* 1957;226(1):497-
509.
1783

Figure 1

A *Mouse CR studies*

Publications (% cumulative total)

Legend: Male only, Female only, M&F separate, M&F combined, N/A

2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021

2003: European Comission highlights addressing relevance of sex in research methodology

2010: CIHR introduces measures to improve the integration of sex in health research

2014: NIH introduces guidelines to include sex as a biological variable in preclinical research

B *Mouse CR studies*

Total = 57 publications

74% Sex difference(s) found
14% Sex difference(s) not found
7% Not comparable

C *Human CR studies*

Publications (% cumulative total)

Legend: Male only, Female only, M&F separate, M&F combined, N/A

2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021

D *Human CR studies*

Total = 50 publications

62% Sex difference(s) found
24% Sex difference(s) not found
14% Not comparable

Figure 1 – Summary of sex differences in mouse and human CR research.
PubMed was searched using keywords to identify research articles that studied caloric restriction *in vivo* in mice (**A,B**) or humans (**C,D**), published between 2003 and 2021. Search results were then classified into the following groups depending on the sexes included in each study: 'Male only' = male subjects used exclusively; 'Female only' = female subjects used exclusively; 'Male & Female separate' = male and female subjects used with data reported respective to each sex, allowing sex differences to be addressed; 'Male & female combined' = male and female subjects used with data from both sexes combined either in part or in full; 'N/A' = no sex data available. **(A,C)** Cumulative publications for studies within each group. The boxes between (A) and (C) highlight the dates of funders' policies highlighting the importance of sex as a biological variable. (**B,D**) Pie charts of those studies in the 'Male & Female separate' group that considered sex a biological variable, and the proportion of these that did or did not detect sex differences. Source data are provided as a Source Data file. See also Supplementary Figure 1.

Figure 2

Figure 2 – Female mice resist CR-induced weight loss, fat loss and adipocyte hypotrophy.

Male and female mice on a C57BL/6NCrl or C57BL/6J background were fed *ad libitum* (AL) or a 30% CR diet from 9-15 weeks of age (0-6 weeks of CR). **(A-F)** Each week mice were weighed (A,B) and body composition was determined by TD-NMR (C-F). Body mass, fat mass and lean mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). **(G)** Masses of gWAT (gonadal WAT), iWAT (inguinal WAT), mWAT (mesenteric WAT), pWAT (perirenal WAT), pcWAT (pericardial WAT) and BAT (brown adipose tissue) were recorded at necropsy and are shown as % body mass. **(H,I)** Micrographs of H&E-stained sections of gWAT (H) were used for histomorphometric analysis of adipocyte area (I); in (H), scale bar = 100 µm. Data in (A-F) are shown as mean ±SEM of the following numbers of mice per group: *male AL,* n=42; *female AL,* n=43; *male CR,* n=44; *female CR,* n=52. Data in (G) are shown as violin plots of the following numbers of mice per group and tissue type: *male AL,* n=29 (iWAT, gWAT, mWAT, BAT), 15 (pWAT) or 14 (pcWAT); *female AL,* n=28 (iWAT, gWAT, mWAT, BAT), 14 (pWAT) or 13 (pcWAT); *male CR,* n=33 (iWAT, gWAT, mWAT, BAT), 16 (pWAT) or 15 (pcWAT); *female CR,* n=34 (iWAT, gWAT, mWAT, BAT), 15 (pWAT) or 14 (pcWAT). Data in (H-I) are shown as representative micrographs or violin plots from 5 (*male AL)* or 6 (*female AL, male CR, female CR)* mice per group. For (A-F), significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA or a mixed-effects model. For (G), significant effects of diet and/or sex within each tissue were determined by 2-way ANOVA with Tukey's multiple comparisons tests. For (I), significant effects of diet and/or sex on adipocyte area (Ad.Ar) were determined using a mixed-effects model, while significant differences in median Ad.Ar between AL and CR mice were determined 2-way ANOVA with Šídák's multiple comparisons test. *P values* from ANOVA or mixed models are shown beneath the graphs, as indicated. For (G) and (I), significant differences between comparable groups are indicated by * (*P*<0.05), ** (*P*<0.01) or *** (*P*<0.001). Source data are provided as a Source Data file. See also Supplementary Figures 2-5.

Figure 3

Figure 3 – The effects of CR on glucose homeostasis differ between young male and female mice.

Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. **(A)** Random-fed blood glucose was recorded each week. **(B-D)** At 13 weeks of age mice underwent an oral glucose tolerance test (OGTT). (B) Blood glucose readings during the OGTT. (C) Area under the curve (AUC) during the OGTT was determined relative to 0 mmol/L (total AUC: tAUC) and relative to baseline (incremental AUC: iAUC). (D) Glucose-stimulated insulin secretion in mice during OGTT was assessed using an insulin ELISA. **(E)** HOMA-IR and Matsuda indices of mice calculated from glucose and insulin concentrations during the OGTT. Data in (A), (B), and (D) are presented as mean ± SEM. Data in (C) and (E) are presented as violin plots overlaid with individual data points. For each group and timepoint, the following numbers of mice were used: **(A)**: *male AL,* n=42 (Wk 0), 36 (Wk2, Wk 2), 34 (Wk 3), 29 (Wk 4), 26 (Wk 5), or 22 (Wk 6); *female AL,* n=43 (Wk 0), 28 (Wk 1), 35 (Wk 2), 31 (Wk 3), 27 (Wk 4), 26 (Wk 5), or 21 (Wk 6); *male CR,* n=44 (Wk 0), 40 (Wk 1), 38 (Wk 2), 35 (Wk 3), 31 (Wk 4), 29 (Wk 5), or 26 (Wk 6); *female CR,* n= 51 (Wk 0), 44 (Wk 1, Wk 2), 41 (Wk 4), 35 (Wk 4), 33 (Wk 5), or 27 (Wk 6). **(B)**: *male AL,* n=27 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female AL,* n=26 (T_{15}, T_{30}, T_{120}), or 25 (T_0, T_{60}); *male CR,* n=27 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female CR,* n=28 (T_0, T_{15}, T_{30}, T_{60}, T_{120}). **(C)**: *male AL,* n=27; *female AL,* n=26; *male CR,* n=27; *female CR,* n=28. **(D)**: *male AL,* n=9 (T_0, T_{15}, T_{30}, T_{60}, T_{120}); *female AL,* n=11 (T_0, T_{15}), 10 (T_{30}, T_{60}), or 8 (T_{120}); *male CR,* n=17 (T_0), 16 (T_{30}), 15 (T_{60}), 13 (T_{15}), or 11 (T_{120}); *female CR,* n=11 (T_{15}, T_{30}, T_{60},), or 10 (T_0, T_{120}). **(E)**: *male AL,* n=9; *female AL,* n=9; *male CR,* n=16; *female CR,* n=9. In (A), (B), and (D), significant effects of diet, sex and/or time, and interactions thereof, were determined using a mixed-effects model. In (C) and (E), significant effects of sex, diet, and sex*diet interaction for tAUC, iAUC, HOMA-IR and Matsuda Index were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown with each graph. For (C) and (E), statistically significant differences between comparable groups are indicated by *** (*P*<0.001) or with *P* values shown. Source data are provided as a Source Data file. See also Supplementary Figures 6-7.

Figure 4

Figure 4 – CR decreases energy expenditure in male and female mice, with a trend for greater nighttime decreases in females.

Male and female mice were fed AL or CR diets as described for Figure 2. At 12-13 or 14-15 weeks of age mice were housed individually in TSE PhenoMaster cages for 3 days for indirect calorimetry. **(A,B)** Absolute energy expenditure (watts, W) was recorded every 15 minutes throughout the 3 days. **(A)** Average energy expenditure per hour over the 24 h light (Day) and dark (Night) period, spanning days 2-3 of PhenoMaster housing. **(B)** Energy expenditure (kJ) during the Day, Night, or Day + Night (Total). **(C)** Energy expenditure normalised to lean body mass (kJ/g) during the Day, Night, or Day + Night (Total). **(D)** Linear regression of lean mass vs Total (Day + Night) energy expenditure (kJ/24 h). Data in (A-D) are from 13 (*male AL*) or 14 (*male CR, female AL, female CR)* mice per group. In (A), data are shown as mean ± SEM; significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA, with P values shown beneath the graph. In (B-C), data are shown as violin plots overlaid with individual data points; within each time period (Day, Night, or Total), significant effects of sex, diet, and diet*sex interaction were assessed using 2-way ANOVA, with P values shown beneath each graph. Statistically significant differences between comparable groups were further assessed by Tukey's multiple comparisons tests and are indicated by ** ($P<0.01$) or *** ($P<0.001$). For linear regression in (D), ANCOVA was used to test if the relationship between lean mass and total energy expenditure differs significantly across all of the individual diet-sex groups (*i. All mice*); between AL and CR mice, irrespective of sex (*ii. AL vs CR (M+F)*); and between males and females fed AL diet (*iii*) or CR diet (*iv*) only. ANCOVA P values for differences in slope and intercept are reported beneath the graph.

Figure 5

A — FDG uptake (SUV, g/mL)

Legend: Male AL, Male CR, Female AL, Female CR

	BAT	Brain	Tibia	Tibia BM	Heart
Diet:	0.1512	0.1585	**0.0342**	0.1733	0.0517
Sex:	**0.0035**	0.5617	0.0829	0.0928	**0.0010**
Sex*Diet:	0.4190	0.9282	0.5143	0.8955	0.6854

B

Male AL   Male CR   Female AL   Female CR

SUV (g/mL)   0 — 2

C — FDG uptake (gamma counts, %ID/g)

Legend: Male AL, Male CR, Female AL, Female CR

	Gastroc	iWAT	gWAT	mWAT	pWAT	WAT (all)	Humerus	Forearm	Bones (all)	Kidneys
Diet:	0.4443	0.0889	0.0553	0.1807	0.1220	**0.0200**	0.8700	0.8362	0.2682	**0.0058**
Sex:	**0.0447**	0.5962	**0.0373**	0.8837	0.5290	0.1747	**0.0232**	**0.0005**	0.1470	**0.0085**
Sex*Diet:	0.9959	0.7461	0.3961	0.6896	0.2926	0.5137	0.7253	0.8818	0.7469	0.2069

Figure 5 – The effects of CR on glucose uptake do not differ between male and female mice. Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. At 15 weeks of age, glucose uptake was assessed by PET/CT with [18]F-fluorodeoxyglucose ([18]F-FDG). **(A-C)** [18]F-FDG uptake in the indicated tissues was determined as SUVs from PET/CT scans (A) or as % injected dose per gram of tissue (%ID/g) from gamma counting of dissected whole tissues (C). **(B)** PET/CT images confirm that [18]F-FDG uptake in interscapular BAT is greater in females than in males, regardless of diet. Data are shown as violin plots (A,C) or representative images (B) of 7 (*male AL, female AL, male CR*) or 6 (*female CR*) mice per group. For each tissue in (A) and (C), significant effects of sex, diet, and sex*diet interaction were assessed by 2-way ANOVA; overall *P* values for each variable, and their interactions, are shown beneath each graph. Significant differences between each group, as determined using Tukey's multiple comparisons test, are indicated by * (*P* <0.05). Source data are provided as a Source Data file. See also Supplementary Figure 7.

Figure 6

Figure 6 – CR exerts sexually dimorphic effects on hepatic gene expression and sphingolipid content.

Male and female C57BL6/NCrl mice were fed AL or CR diet from 9-15 weeks of age, as described for Figure 2. Livers were sampled at necropsy (15 weeks) and used to assess mRNA transcripts, triglyceride content, and ceramide and dihydroceramide content. **(A)** Expression of the indicated transcripts was determined by qPCR. Transcript expression was normalised to the geometric mean of *Tbp, Rn18s* and *Ppia* expression and is presented relative to the highest group average. **(B)** Principle component analysis of transcript expression in (A). **(C)** Hepatic triglyceride content. **(D,E)** LC-MS analysis of total, 22:0 and 23:0 ceramides (D), dihydroceramides (E, left side) and the ceramide:DHC ratio (E, right side). **(F)** Principle component analysis of ceramide and dihydroceramide content based on data in (D-E) and Supplementary Figure 8. Data in (A), (C), (D), and (E) are presented as truncated violin plots overlaid with individual data points. Data represent the following numbers of mice per group. **(A-B)**: *male AL,* n=6; *female AL,* n=6; *male CR,* n=6; *female CR,* n=6. **(C)**: *male AL,* n=9; *female AL,* n=8; *male CR,* n=9; *female CR,* n=8. **(D-F):** 6 mice per group. For (A), (C), (D) and (E), significant effects of sex, diet, and sex*diet interaction were assessed using 2-way ANOVA; overall *P* values for each variable, and their interactions, are shown beneath each graph. Significant differences between comparable groups were assess using Tukey's multiple comparison test and are indicated by * ($P<0.05$), ** ($P<0.01$) or *** ($P<0.001$). Source data are provided as a Source Data file. See also Supplementary Figure 8.

Figure 7

Figure 7 – Sex differences in CR-induced weight loss and fat loss are absent when CR is initiated in aged mice.

Male and female C57BL6/NCrl mice were single housed and fed AL or a CR diet (70% of daily AL intake) from 78-84 weeks of age. **(A-F)** Each week mice were weighed (A,B) and body composition was determined by TD-NMR (C-F). Body mass, fat mass and lean mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). **(G)** Masses of gWAT, iWAT, mWAT, pWAT, pcWAT and BAT were recorded at necropsy and are shown as % body mass. Data in (A-F) are shown as mean ±SEM. Data in (G) are shown as violin plots overlaid with individual data points. For each group and timepoint, the following numbers of mice were used: **(A,B)**: *male AL*, n=9 (Wk 0, Wk 2, Wk3, Wk 4), 8 (Wk 1), or 6 (Wk 5, Wk 6); *female AL,* n=13 (Wk 0, Wk 2), 12 (Wk 4, Wk 5), 11 (Wk 6), 9 (Wk 1, or 8 (Wk 3); *male CR,* n=10 (Wk 0, Wk 2-4), 9 (Wk 1), 7 (Wk 5), or 6 (Wk 6); *female CR,* n=14 (Wk 0, Wk 4, Wk 5), 13 (Wk 2, Wk 6), 10 (Wk 3), or 9 (Wk 1). **(C-F)**: *male AL,* n=9 (Wk 0, Wk 2), 8 (Wk 3, Wk 4, Wk 6), or 7 (Wk 1, Wk 5); *female AL,* n=13 (Wk 0, Wk 2), 9 (Wk 4, Wk 6), 8 (Wk 5), 6 (Wk 1), or 5 (Wk 3); *male CR,* n=10 (Wk 0, Wk 2), 9 (Wk3-6), or 8 (Wk 1); *female CR,* n=14 (Wk 1, Wk 2), 11 (Wk 4-6), 7 (Wk 3), or 6 (Wk 1). **(G)**: *male AL,* n=9; *female AL,* n=12 (iWAT, mWAT) or 11 (gWAT, pWAT); *male CR,* n=10 (iWAT, pWAT), or 9 (gWAT, mWAT); *female CR,* n=14 (mWAT, pWAT), 13 (iWAT), or 12 (gWAT). For (A-F), significant effects of diet, sex and/or time, and interactions thereof, were determined by 3-way ANOVA or a mixed-effects model. For (G), significant effects of sex, diet, and sex*diet interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown beneath each graph. For (G), significant differences between comparable groups are indicated by ** (*P*<0.01) or *** (*P*<0.001). Source data are provided as a Source Data file. See also Supplementary Figure 9.

Figure 8

Figure 8 – Sex differences in CR's effects on glucose homeostasis are largely absent when CR is initiated in aged mice.

Male and female C57BL6/NCrl mice were fed AL or CR diet from 78-84 weeks of age, as described for Figure 7. **(A)** Random-fed blood glucose was recorded each week. **(B-D)** At 82 weeks of age mice underwent OGTTs. Blood glucose was recorded at each timepoint (B) and the tAUC and iAUC was determined (C). (D) Glucose-stimulated insulin secretion in mice during OGTT was assessed using an insulin ELISA. **(E)** HOMA-IR and Matsuda indices of mice calculated from glucose and insulin concentrations during the OGTT. Data in (A), (B), and (D) are presented as mean ± SEM. Data in (C) and (E) are presented as violin plots overlaid with individual data points. For each group and timepoint the following numbers of mice were used: **(A)**: *male AL,* n=9 (Wk 0, Wk 2-4, Wk 6), 8 (Wk 1), or 5 (Wk 5); *female AL,* n=12 (Wk 0, Wk2, Wk4), 8 (Wk 1, Wk 3), or 11 (Wk 5, Wk 6); *male CR,* n=10 (Wk 0, Wk 2-4, Wk 6), 9 (Wk 1), or 7 (Wk 5); *female CR,* n= 14 (Wk 0, Wk 2, Wk 4-6), 9 (Wk 1) or 10 (Wk 3). **(B-E)**: *male AL,* n=9 (B-C) or 5 (D-E); *female AL,* n=12; *male CR,* n=10; *female CR,* n=14. In (A), (B), and (D), significant effects of diet, sex and/or time, and interactions thereof, were determined using a mixed-effects model or 3-way ANOVA. In (C) and (E), significant effects of sex, diet, and sex*diet interaction for tAUC, iAUC, HOMA-IR and Matsuda Index were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown with each graph. For (C) and (E), statistically significant differences between comparable groups are indicated by * (*P*<0.05) or *** (*P*<0.001), or with *P* values shown. Source data are provided as a Source Data file.

Figure 9

Figure 9 – Effects of CR on body mass and body composition in humans are sex- and age-dependent.

Twenty male and twenty-two female volunteers participated in a weight loss study involving a 4-week dietary intervention. **(A-F)** Body mass was recorded each week (A,B)**.** Fat mass (FM) and fat-free mass (FFM) were measured by air-displacement whole-body plethysmography at weeks 0 and 4 (C-F). Body mass, fat mass and fat-free mass are shown as absolute masses (A,C,E) or fold-change relative to baseline (B,D,F). Data are presented as mean ± SEM**.** For (A-C) and (E), significant effects of time, sex, and time*sex interaction were assessed using 2-way ANOVA. In (B), (D) and (F), significant differences between males and females at each time point were determined by Šídák's multiple comparisons test (B) or unpaired T-test (D,F) and are indicated by * ($P<0.05$) or ** ($P<0.01$). **(G-I)** Simple linear regression of age vs fold-change (*week 4 vs week 0)* in body mass (G), fat mass (H) and fat-free mass (I). For each sex, significant associations between age and outcome (fold-change) are indicated beneath each graph as '*P,* Slope'. ANCOVA was further used to test if the age-outcome relationship differs significantly between males and females. ANCOVA results are reported beneath each graph as '*P,* Slope' and '*P,* Intercept' for males vs females (M vs F). In (G), similar slopes but different intercepts show that sex significantly influences weight loss, but the influence of age does not differ between the sexes. In (H,I) the slopes differ significantly, indicating that the age-outcome relationship differs between the sexes. **(J-L)** Fold-change (week 4 vs week 0) in body mass (J), fat mass (K) and fat-free mass (L) for males vs females separated into younger (<45 years) and older (>45 years) groups. Data are presented as violin plots overlaid with individual data points. Significant effects of age, sex, and age*sex interaction were assessed using 2-way ANOVA with Tukey's multiple comparisons test. Overall *P* values for each variable, and their interactions, are shown beneath each graph. Significant differences between comparable groups are indicated by * ($P<0.05$) or *** ($P<0.001$). See also Table 3. Source data are provided as a Source Data file.