Cobalamin riboswitches are broadly sensitive to corrinoid cofactors to enable an efficient
gene regulatory strategy

Kristopher J. Kennedy[1], Florian J. Widner[1], Olga M. Sokolovskaya[1,2], Lina V. Innocent[1],
Rebecca R. Procknow[1], Kenny C. Mok[1], and Michiko E. Taga[1,*]

[1]Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA
94720, USA

[2]Present address: Department of Infectious Diseases, Genentech Inc., 1 DNA Way, South San
Francisco, CA, 94080, USA

[*]Corresponding author: taga@berkeley.edu

Abstract

In bacteria, many essential metabolic processes are controlled by riboswitches, gene regulatory RNAs that directly bind and detect metabolites. Highly specific effector binding enables riboswitches to respond to a single biologically relevant metabolite. Cobalamin riboswitches are a potential exception because over a dozen chemically similar but functionally distinct cobalamin variants (corrinoid cofactors) exist in nature. Here, we measured cobalamin riboswitch activity *in vivo* using a *Bacillus subtilis* fluorescent reporter system and found that among 38 tested riboswitches, a subset responded to corrinoids promiscuously, while others were semi-selective. Analyses of chimeric riboswitches and structural models indicate that, unlike other riboswitch classes, cobalamin riboswitches indirectly differentiate among corrinoids by sensing differences in their structural conformation. This regulatory strategy aligns riboswitch-corrinoid specificity with cellular corrinoid requirements in a *B. subtilis* model. Thus, bacteria can employ broadly sensitive riboswitches to cope with the chemical diversity of essential metabolites.

Introduction

Controlling gene expression is an essential task that cells accomplish in a variety of ways. Non-coding RNAs are one such means of gene regulation, acting in parallel or in concert with historically better-studied protein-based mechanisms (1). In bacteria and archaea, riboswitches are a widespread type of gene regulatory RNA with the distinct ability to sense particular intracellular metabolites by direct binding (2). These RNAs are typically located in the 5'-untranslated region of mRNA transcripts and function as cis-regulators of downstream genes within their transcripts. A riboswitch is composed of an effector-binding aptamer domain and an expression platform. The aptamer domain adopts a three-dimensional structure that can bind its cognate effector molecule. The expression platform domain is a regulatory switch that interprets the effector-binding state of the upstream aptamer typically to promote or disrupt the transcription or translation of downstream genes (3). The diversity of riboswitch effectors and regulatory mechanisms has revealed fundamental insights into how bacteria sense and respond to dynamic environments and has also driven new approaches for precise control and manipulation of microbes for human purposes (4-7).

Cobalamin (Cbl) riboswitches (also called 'B$_{12}$ riboswitches' or 'adenosylcobalamin riboswitches') are among the most widespread and structurally diverse types of riboswitch in bacteria (8). They directly bind various forms of the enzyme cofactor Cbl as a cognate effector (Figure 1A, C) (9) to regulate genes involved in the biosynthesis, transport, and usage of Cbl. Cbl-dependent enzymes function in common metabolic pathways including methionine synthesis, deoxyribonucleotide synthesis, tRNA modification, and the degradation of certain amino acids, fatty acids, and biopolymers (10-23). Cbl is also required for rarer metabolic processes involved in antibiotic synthesis, mercury methylation, catabolism of steroids, and many others (24-37). Comparative genomic studies indicate that most bacteria perform Cbl-dependent metabolism and that Cbl-riboswitches often regulate these processes (8, 38, 39). However, an overlooked facet among most riboswitch studies is that Cbl is just one member of a class of enzyme cofactors known as corrinoids (Figure 1B) (40). In fact, Cbl-dependent enzymes in bacteria often function with corrinoids other than Cbl. Yet, it remains unclear whether the dozens of naturally occurring corrinoid cofactors are also Cbl-riboswitch effectors (41-47).

Corrinoid cofactors contain a highly substituted corrin ring with a central cobalt ion, a variable 'upper ligand' moiety coordinating the β axial face of the cobalt, and a tail structure extending from the corrin ring and terminating in a variable 'lower ligand' moiety that often coordinates the α axial face of the cobalt (Figure 1A, B). The molecular basis of selectivity of Cbl-riboswitches for upper ligand variants of Cbl has been relatively well studied, but selectivity for corrinoid tail variants remains mostly unexplored (9, 48, 49). To our knowledge, only one study has directly examined corrinoid tail specificity of a single Cbl-riboswitch. *In vitro* binding measurements showed that the aptamer of the *Escherichia coli btuB* Cbl-riboswitch binds the complete corrinoids Cbl and 2-methyladeninylcobamide ([2-MeAde]Cba) with a 3.2-fold difference in affinity (K_D = 89 and 290 nM, respectively). Furthermore, Cbi, an incomplete corrinoid with a truncated tail, binds the aptamer with roughly 8,000-fold lower affinity than Cbl (K_D = 753 μM), suggesting that this aptamer binds corrinoids in a selective manner (50). In light

of these previous studies of corrinoid-specific metabolisms and Cbl-riboswitches, we
hypothesize that Cbl-riboswitches harbor a range of distinct corrinoid tail-specific activities.

Here, we examined how a panel of 38 Cbl-riboswitches derived from 12 bacterial species
responds to the distinct tail structures of four corrinoids: Cbl, pCbl, and CreCba, representatives
of the benzimidazolyl, purinyl, and phenolyl cobamides, respectively, and Cbi, an incomplete
corrinoid (Figure 1B). To compare activities among several dozen Cbl-riboswitches, we devised
a live cell fluorescence-based reporter system in *Bacillus subtilis*. In contrast to conventional *in*
vitro biochemical approaches, this riboswitch reporter system captures the complete corrinoid-
responsive gene regulatory process and provides rapid functional measurements with multiple
effectors in parallel. Our results obtained from experiments in the reporter system in conjunction
with comparative structural analyses of Cbl-riboswitches and corrinoid effectors allowed us to
develop a mechanistic model for how corrinoid tail-specific gene regulation is achieved.
Additionally, we examine a gene regulatory strategy for the corrinoid specificity of a Cbl-
riboswitch and discuss the conceptual and practical implications of these findings.

Results

Experimental strategy – Development of an in vivo *reporter system*

In order to compare corrinoid selectivity among several Cbl-riboswitches and corrinoids, we
constructed an *in vivo* GFP reporter system. We initially attempted to use an *E. coli* host for the
reporter system but found that most of the riboswitches we tested did not function in the *E. coli*
host. We chose *B. subtilis* as an alternative host organism because of the robust genome
engineering and gene expression toolsets available. Furthermore, the *B. subtilis* genome does not
contain any annotated corrinoid biosynthesis or remodeling genes that would potentially interfere
with a riboswitch reporter assay. We engineered the strain to overexpress the Cbl uptake and
adenosylation operon, *btuFCDR*, and deleted *queG*, which encodes the only Cbl-dependent
enzyme in the genome. We found that *btuFCDR* overexpression increased uptake of not only
Cbl, but also pCbl, CreCba and Cbi (Figure 2A-D). Additionally, each corrinoid was transformed
from the cyanated to adenosylated form, suggesting that the corrinoids are internalized to the
cytoplasm, and not simply accumulating on the outer cell surface. This strain is effective for
measuring the response of the *B. subtilis btuF* Cbl-riboswitch to a broad range of concentrations
of Cbl (Figure 2E-G). Notably, deletion of *btuR*, which encodes the adensosyltransferase that
installs the Ado upper ligand group, rendered the *B. subtilis btuF* Cbl-riboswitch reporter
insensitive to exogenously supplied CNCbl, MeCbl, and OHCbl, while retaining dose-dependent
repression in response to AdoCbl (Figure S1). This strongly suggests that the *B. subtilis btuF*
Cbl-riboswitch only responds to Cbl containing the 5'-deoxyadenosine upper ligand, in contrast
with a report suggesting that this riboswitch aptamer can also bind MeCbl and OHCbl (51).

Comparison of corrinoid specificity among Cbl-riboswitches

In the strain background described above, we constructed 86 reporter strains to examine
riboswitches from 20 bacterial species including 10 species known to produce or require specific

corrinoids. Of the 86 reporters, 38 repressed GFP expression 0.5-fold or greater in response to
one or more corrinoids. 37 of these 38 functional riboswitch expression platforms contain a
predicted intrinsic transcriptional terminator suggesting that they are transcriptional riboswitches.
We observed extensive variation in sequence length and nucleotide composition throughout the
aptamers and expression platforms of the 38 riboswitches that were functional in *B. subtilis*
(Figure 1D). Nine of the functional riboswitches are from *Priestia* (formerly *Bacillus*)
megaterium, which produces Cbl (41), and 12 are from *Sporomusa ovata* and *Veillonella parvula*
which both produce CreCba (44, 45, 52, 53). These results show that Cbl-riboswitches of diverse
sequence composition and origin can be examined with the *in vivo* reporter system. To address
whether Cbl-riboswitches are corrinoid selective, and how corrinoid selectivity varies among
Cbl-riboswitches, we measured the dose-responses of the 38 functional Cbl-riboswitches to four
corrinoids, Cbl, pCbl, CreCba and Cbi. Our results show that all of the riboswitches responded to
more than one corrinoid, and a subset responded to all four (Figure 3). Strikingly, all of the tested
riboswitches are either semi-selective (responding to more than one corrinoid) or promiscuous
(responding to all four corrinoids) (Figure 3A-B). We did not find any highly selective
riboswitches that respond to only one corrinoid. The semi-selective and promiscuous
riboswitches all respond to Cbl and pCbl (Figure 3C), and the promiscuous riboswitches
additionally respond to Cbi and CreCba (Figure 3D-E, points above the horizontal dashed line).
Furthermore, the semi-selective riboswitches are generally more sensitive to Cbl than to pCbl,
while the promiscuous riboswitches respond similarly to these two corrinoids. Almost all of the
riboswitches respond weakly to CreCba compared to the other three corrinoids (Figures S2A-C).
In general, corrinoid selectivity of a riboswitch appears to be associated with its taxonomic
origin (Figure 3C-E). The riboswitches from the Bacilli class are exclusively semi-selective
(Figure S2A), whereas those from Negativicutes are predominantly promiscuous (Figure S2B).
The *S. ovata cobT* riboswitch is a notable exception that will be discussed later. In contrast to
taxonomy, corrinoid selectivity of a riboswitch is not strongly associated with the function of its
regulatory target genes (Figure S3).

Next, we attempted to identify the RNA sequence features that underlie corrinoid selectivity.
Chimeric fusions of the semi-selective *P. megaterium metE* riboswitch and the promiscuous *V.*
parvula mutA riboswitch enabled us to examine the effects of specific domain and subdomain
sequences on corrinoid selectivity (Figure S4). Fusing the *P. megaterium metE* aptamer domain
to the expression platform of the *V. parvula mutA* riboswitch produced a semi-selective
riboswitch chimera, while the reciprocal chimera was promiscuous, suggesting that the aptamer
domain is a major determinant of corrinoid selectivity (Figure S4A). However, results from
aptamer subdomain swaps of stem P1, stem-loop P2-L2, stem-loop P4-L4, and P6 accessory
region were less conclusive. In the context of the *P. megaterium metE* riboswitch scaffold,
swapping stem-loop P2-L2 or the P6 accessory region with the corresponding structures of the *V.*
parvula mutA riboswitch produced chimeras that are less selective by gaining sensitivity to Cbi
and CreCba (Figure S4B). This suggests that these subdomains confer corrinoid promiscuity.
Yet, within the *V. parvula mutA* riboswitch scaffold, swapping stem P1 increased corrinoid
selectivity by retaining sensitivity to Cbl and pCbl, but losing sensitivity to CreCba and Cbi
(Figure S4C). The remaining chimeras partially or completely lost overall activity. These results
demonstrate that subdomains distributed throughout the aptamer domain may impact corrinoid

selectivity; no single conserved substructure completely controls corrinoid selectivity, nor did
any single structure fully convert a riboswitch's corrinoid selectivity. Thus, the source of the
corrinoid selectivity phenotype appears to be complex and requires inputs from multiple
subdomains of the aptamer.

Corrinoid tail structure impacts selectivity of Cbl-riboswitches

We next sought to identify how structural differences in the corrinoid tail affect the response of
semi-selective Cbl-riboswitches to corrinoids. There are no predicted hydrogen bond interactions
between the lower ligand and the RNA in the X-ray crystal structures of Cbl-bound riboswitches,
making it difficult to surmise how a Cbl-riboswitch might distinguish between corrinoids (51,
54, 55). Could the overall structural conformation of the corrinoid, rather than specific
interactions between the RNA and the corrinoid tail, influence Cbl-riboswitch activity?

Corrinoids undergo major conformational changes when spontaneously switching between two
distinct states known as 'base-on' and 'base-off' (56, 57). In the base-on state, a nitrogen atom in
the lower ligand base is coordinated to the central cobalt atom of the corrin ring (as shown in
Figure 1A). In the base-off state, the lower ligand base is de-coordinated, allowing the tail to
move more freely (58, 59). Benzimidazolyl and purinyl cobamides (e.g., Cbl, pCbl) can switch
between base-on and base-off states . However, the tail moieties of phenolyl cobamides (e.g.,
CreCba) and Cbi cannot coordinate cobalt and so these corrinoids exist exclusively in a de-
coordinated state (52). Interestingly, we noticed that the semi-selective riboswitches respond
strongly to Cbl and pCbl, but respond weakly to Cbi and CreCba. We also observe that semi-
selective Cbl-riboswitches are most sensitive to Cbl, which forms the base-on state more readily
than pCbl (Figures 3C, S2) (58, 59). In line with these results, the *E. coli btuB* riboswitch
aptamer was previously shown to bind Cbl with higher affinity than Cbi and [2-MeAde]Cba (a
primarily base-off corrinoid) (50). Also, all six X-ray crystal structures of Cbl-riboswitches
contain Cbl in the base-on state (51, 54, 55). Based on these observations, we hypothesized that
semi-selective riboswitches distinguish between base-on and base-off states of corrinoids. We
therefore used a range of corrinoids with diverse lower ligand structures to test whether the
activity of Cbl-riboswitches quantitatively correlates with the base-on tendency of corrinoids.

We selected a panel of sixteen corrinoids for this analysis, including both natural and synthetic
benzimidazolyl, purinyl, and aza-benzimidazolyl corrinoids. These corrinoids span a range of
base-on tendency between that of Cbl and pCbl, which we measured as the ratio of spectral
absorbance at 525 and 458 nm in the adenosylated form (Figures 4A, S5). Base-on/base-off
equilibrium constants for AdoCbl, Ado[2-MeAde]Cba, and pCbl in aqueous conditions have
been reported as 76, 0.48, and 0.30, respectively, and are consistent with our measurements of
corrinoid base-on tendency (58, 59). We observed a strong association between base-on tendency
and riboswitch response in semi-selective riboswitches (Figure 4B). Even among promiscuous
Cbl-riboswitches, we observe measurable sensitivity to base-on tendency, albeit to a much
smaller degree (Figure 4C). These results support the hypothesis that Cbl-riboswitches
selectively respond to corrinoids by distinguishing between the base-on and base-off states of
corrinoids.

Structural comparisons between base-on and base-off tail orientations

The results presented above led us to speculate about how a Cbl-riboswitch could detect the base-on and base-off state of a corrinoid. In all published X-ray crystal structures of Cbl-riboswitches, aptamer-effector binding is achieved mainly through van der Waals forces and shape complementarity between the binding site and base-on Cbl. Only a few hydrogen bonds between the RNA and corrinoid are observed, none of which occur with the lower ligand group of Cbl (51, 54, 55). Thus, it appears unlikely that the riboswitch is directly detecting the specific chemical differences among corrinoid lower ligands. Instead, we considered whether the riboswitch discriminates base-on and base-off forms of a corrinoid by sensing corrinoid conformation. In the base-on state, the tail is spatially constrained due to the Co-N coordinate bond, whereas in the base-off form it is able to sample a wider range of spatial positions (60). To develop mechanistic insight into how the base-on and base-off states a corrinoid could impact Cbl-riboswitch activity, we leveraged the plethora of publicly accessible X-ray crystal structures of macromolecule-bound Cbl (37, 51, 54, 55, 61-70). We first assessed the range of structural conformations that are potentially sampled by corrinoids as they dynamically switch between base-on and base-off states by aligning and visually comparing various structural models of Cbl. Six base-on Cbl models were obtained from structural studies of Cbl-riboswitches and synthetic Cbl RNA aptamers, whereas base-off/His-on Cbl models were obtained from X-ray crystal structures of ten Cbl-dependent enzymes (Table S2). After aligning and superimposing these molecular models by their central cobalt and coordinating nitrogen atoms, we observed that the corrin rings and their amide and methyl substituents occupy similar spatial positions, but the tails of base-on and base-off Cbl structures occupy distinct positions (Figure 5A-B). Moreover, the base-on Cbl tails appear in very similar positions with lower ligands in close proximity to the central cobalt ion, whereas the base-off Cbl tails appear more scattered with lower ligands more distal to the cobalt ion. These structural alignments visually convey the degree to which the conformations of base-on and base-off corrinoids can vary among biomolecular complexes.

Next, we compared the positions of the aligned base-on and base-off Cbl models in the context of 3D Cbl-riboswitch models. We analyzed X-ray crystal structures of the two Cbl-riboswitches that contain resolved kissing loop structures: one from *Thermoanaerobacter tengcongensis* (Figure 5C-D) and one identified from a marine metagenome sequence (Figure S6A-B) (54). These structural models show that the base-on tails are contained within the binding site, whereas the tails of the base-off Cbl structures protrude away from the binding site and clash with the L5-L13 kissing loop. Although the *B. subtilis btuF* (Figure S6C-D) and *Symbiobacterium thermophilum cblT* (Figure S6E-F) riboswitch models do not contain the L13 structure of the kissing loop, some of the modeled base-off tails clash with L5 of the aptamer in these structures (51, 55). The kissing loop has been shown to play a key mechanistic role of sensing the corrinoid-binding state of the aptamer domain to influence downstream regulatory structures in the expression platform (71, 72). If kissing loop formation in semi-selective Cbl-riboswitches is sensitive to the base-on and base-off states of the corrinoid, then corrinoid selectivity may be mediated by either selective binding by the aptamer or selective formation of downstream regulatory structures.

To determine whether the expression platform structures can impact corrinoid selectivity, we
examined corrinoid-selective binding separately from subsequent corrinoid-selective regulation.
We tested for promiscuous binding by comparing the Cbl dose-response of the *P. megaterium*
metE Cbl-riboswitch in the presence and absence of competing 100 nM Cbi and found that the
response to Cbl is unaffected by Cbi (Figure S7A-B). This indicates that Cbi does not compete
with Cbl for riboswitch binding, supporting corrinoid-selective binding as the mechanism of
semi-selectivity. However, when the aptamer of this semi-selective riboswitch is replaced with
the aptamer of the promiscuous *S. ovata nikA* riboswitch, it retains semi-selectivity, which
suggests that the *P. megaterium* expression platform also plays a role in corrinoid selectivity
(Figure S7A). Interestingly, the Cbl dose-response of the *S. ovata nikA* / *P. megaterium metE*
chimeric riboswitch does become sensitized to competing Cbi addition, confirming that the *S.*
ovata nikA aptamer retains sensitivity to base-off corrinoid in the context of this chimera (Figure
S7C). Taken together, these results show that base-off corrinoids may impede both Cbl-
riboswitch binding and formation of regulatory structures, explaining the link between corrinoid
base-on tendency and riboswitch activity observed in Figure 4.

Gene regulatory strategy of corrinoid selectivity

While the prior experiments clearly demonstrate that Cbl-riboswitches are capable of
distinguishing between corrinoids, we wondered what purpose Cbl-riboswitch corrinoid
selectivity might serve in the organisms containing these regulatory systems. We posit that Cbl-
riboswitch selectivity reflects a regulatory strategy that complements the corrinoid-specific
requirements of the cell and avoids gene mis-regulation. As a specific example, we hypothesize
that only corrinoids that are functionally compatible with a Cbl-dependent enzyme should cause
riboswitch-mediated repression of the expression of its Cbl-independent counterpart (Figure 6A,
B). We tested this hypothesis directly in *B. subtilis* by examining the function and regulation of
methionine synthase isozymes MetE (Cbl-independent) and MetH (Cbl-dependent). In bacterial
genomes with both *metE* and *metH*, a tandem SAM-Cbl riboswitch is commonly found upstream
of the *metE* gene (73, 74). The *B. subtilis* genome contains *metE* but lacks *metH*, and no Cbl-
riboswitch is located upstream of *metE*. We therefore constructed strains of *B. subtilis* that
heterologously express the *metE* or *metH* locus from the Cbl-producing species *P. megaterium*,
in a Δ*metE* background with overexpressed corrinoid uptake genes (Figure 6C) (75-77). In each
strain, the *P. megaterium* genes are constitutively transcribed from the promoter P_{Veg} and also
contain a transcriptionally fused *gfp* to measure expression levels. Growth of the *B. subtilis* strain
expressing *P. megaterium metH* in a medium lacking methionine was supported to varying
extents by most benzimidazolyl and both phenolyl cobamides, but not by [5-OHBza]Cba, the
purinyl cobamides, or Cbi (Figure 6D, red squares). This result indicates that MetH-dependent
growth is influenced by the corrinoid tail structure, as observed previously in other bacteria (78-
82). In the *B. subtilis* strain containing the *P. megaterium metE* locus which includes the
repressing SAM-Cbl-riboswitch, growth was suppressed by benzimidazolyl cobamides to
different extents (Figure 6D, blue circles). This growth pattern coincides with the GFP repression
measured for the *metE* riboswitch (Figure 6E), except that the repression of *metE* by purinyl
cobamides is apparently insufficient to suppress growth in this context. Comparison of the two
strains in response to a suite of corrinoids reveals a striking correspondence between riboswitch-

mediated suppression of growth in the *metE*-containing strain and growth promotion by
corrinoids in the *metH*-containing strain (Figure 6C). [5-OHBza]Cba and the phenolyl cobamides
are exceptions to the trend, though in neither case is MetE-dependent growth completely
suppressed by a corrinoid incompatible with MetH. This result demonstrates that riboswitch-
based repression and cobalamin-dependent isozyme function are largely aligned for the *P.*
megaterium metE-metH pair and suggests that Cbl-riboswitch specificity may generally adhere
to a regulatory strategy reflecting the cell's corrinoid preference.

Discussion

Riboswitches are key regulators of microbial gene expression. The Cbl-riboswitch was the first
type discovered and is among the most widely distributed riboswitch classes in bacteria and
archaea (9, 83). Previous biochemical and structural studies have uncovered the major molecular
features of the Cbl-riboswitch response to Cbl, including how upper ligand variants of Cbl
impact their function (48, 49, 51, 54, 55, 71, 72, 74, 84). Yet few studies have examined how
other naturally occurring corrinoids containing diverse lower ligand structures impact gene
regulation by Cbl-riboswitches (50). Here, we found that Cbl-riboswitches vary in their ability to
discriminate between corrinoids, with some being semi-selective on the basis of corrinoid base-
on/off state, and others being promiscuous. These results were enabled by a carefully designed
fluorescent reporter system capable of measuring the responses of dozens of Cbl-riboswitches to
multiple corrinoids *in vivo* (Figures 2, S1). Since several naturally occurring corrinoids other
than Cbl appear to be potent effectors for Cbl-riboswitches, we propose that the term 'corrinoid
riboswitch' be adopted to describe this broad class of RNAs more accurately. This may also
mitigate the inconsistent and overlapping terminology used in the literature (i.e. cobalamin
riboswitch, adenosylcobalamin riboswitch, B_{12} riboswitch, vitamin B_{12} riboswitch, etc.).

We can roughly estimate the intracellular corrinoid concentrations in our corrinoid dose-response
experiments to assess the physiological relevance the *in vivo* Cbl-riboswitch data. With the
observation that most of the corrinoid in the medium is imported by the *B. subtilis* riboswitch
reporter cells (Figure 2A-D), and assuming a cell volume of 10^{-15} L and a culture cell titer of 10^{12}
cells/L at $OD_{600} = 1$, we calculate the cytoplasmic corrinoid concentrations to range from 0.01 to
100 μM in the corrinoid dose-response experiments – a 1,000-fold increase from the cell culture
medium to the cytoplasm. Reports of binding affinity (K_D) for riboswitch aptamers to AdoCbl
range from 0.026 to 90 μM, which is within the range of our estimated intracellular corrinoid
concentrations (48, 50, 54, 72, 85). Additionally, Cbl uptake has been measured in some
bacterial species. In *E. coli*, the minimum cytoplasmic Cbl concentration to support MetH-
dependent growth is roughly 0.03 μM (86). In studies of Cbl uptake across several bacterial
species, saturating Cbl uptake can result in cytoplasmic Cbl concentrations in the low μM to low
mM range, depending on the species (87, 88). Taken together, these data and calculations
support the physiological relevance of the Cbl-riboswitch responses measured in this study.

We observed that Cbl-riboswitches display different degrees of corrinoid selectivity, with some
that respond to a subset of corrinoids (semi-selective) and others that respond to all tested

corrinoids (promiscuous) (Figures 3, S2). Our chimeric riboswitch results suggest that sequence
and structural determinants of corrinoid selectivity are dispersed throughout the Cbl-riboswitch
aptamer scaffold rather than being confined to a single conserved region (Figure S4). This
finding contrasts with other studies of riboswitch specificity. For example, in a study of Cbl
upper ligand specificity, a few key residues in the Cbl binding site were sufficient to fully
convert a Cbl-riboswitch from MeCbl-specific to AdoCbl-specific (48). In purine riboswitches,
effector specificity is achieved by positioning of a critical conserved uracil or cytosine residue in
the binding site of the aptamer, which forms a base-pair with the adenine or guanine effector,
respectively. Among the various families of SAM riboswitches, highly specific binding of SAM
and exclusion of SAH is achieved by RNA structures that discriminate the charged sulfonium ion
of SAM from the uncharged sulfoether of SAH (89). In contrast, the SAM/SAH riboswitch class
attains effector promiscuity for SAM and SAH by a general lack of interaction between the RNA
and the aminocarboxypropyl side chains of these effectors (90, 91). This is reminiscent of Cbl-
riboswitches which similarly have few molecular contacts between the RNA and corrinoid tail
(50, 54, 55, 84).

Overall, our structural model analyses support a mechanism in which semi-selective Cbl-
riboswitches primarily sense the distinct corrinoid tail orientations of the base-on and base-off
forms. In this case, Cbl-riboswitches only indirectly sense the chemical composition of the
variable lower ligand group, with differential binding largely determined by steric effects and
shape complementarity (Figures 4-5). Recent molecular dynamics simulations of the *T.
tengcongensis* Cbl-riboswitch suggest that the kissing loop structure may form prior to effector
binding, which would place even greater constraints on the corrinoid tail orientation to achieve
shape complementarity with its binding site (92). Additionally, some Cbl-riboswitches may in
fact bind base-off corrinoids but disrupt subsequent formation of downstream regulatory
structures of the expression platform, perhaps by interfering with the kissing loop (Figure S7). A
similar feature has been observed in tetrahydrofolate (THF) riboswitches where chemical
variations in the para-aminobenzoic acid moiety of THF analogs differentially perturb expression
platform structures without affecting aptamer binding (93). The mechanisms of corrinoid
selectivity of Cbl-riboswitches could be directly tested in future structural or biochemical studies
of promiscuous Cbl-riboswitches with base-on and base-off corrinoids.

In regard to gene regulatory strategies, it seems sensible that Cbl-riboswitches are not highly
effector-specific because bacteria are often flexible in their corrinoid usage. A variety of
corrinoids have been shown to support growth of *C. difficile*, *S. ovata, and Ensifer meliloti*
despite each of these organisms displaying highly specific corrinoid production (82, 94, 95).
Furthermore, since corrinoid auxotrophy is prevalent among corrinoid-dependent bacteria, many
organisms may need to take advantage of the wide range of corrinoids that may be available in
their environment (96). Thus, the range of effector selectivity that we observe among Cbl-
riboswitches may reflect a coevolution between corrinoid-responsive gene regulation and
corrinoid-dependent physiology. Our result demonstrating complementary corrinoid selectivity
between *P. megaterium* MetH-dependent growth and Cbl-riboswitch-dependent expression of
MetE is consistent with this notion (Figure 6).

Alternatively, the preference for base-on corrinoids among Cbl-riboswitches may function as a
proxy to discriminate complete corrinoid coenzymes from incomplete corrinoids such as Cbi,
which often function poorly as coenzymes. This idea has been proposed as an explanation for the
remarkably high selectivity of the corrinoid uptake system in mammals (97). Interestingly, we
found that all but one of the *S. ovata* riboswitches tested are promiscuous types that can respond
to its natively produced CreCba. The notable exception is a semi-selective riboswitch upstream
of the gene *cobT* (Figure S2B), which functions in a late step of corrinoid biosynthesis that
occurs after synthesis of Cbi (53, 98-103). Thus, this Cbl-riboswitch that discriminates against
Cbi may allow homeostatic regulation of *cobT* in response to complete corrinoids like Cbl, while
preventing unproductive repression of *cobT* in the presence of incomplete corrinoids like Cbi.
Future studies examining corrinoid-specific gene regulation of riboswitches in the context of
their native organisms may help clarify which regulatory strategies are generally at play in
corrinoid-related bacterial physiology.

Our findings fit into a broader discussion of how corrinoids impact complex microbial
communities (104). In future studies, it will be worth examining how the interplay between
corrinoid-specific gene regulation and corrinoid-specific metabolic pathways influence microbial
interactions. There is also significant interest in the fields of bioengineering and synthetic
biology to use riboswitches as gene regulatory devices because they act more rapidly and
efficiently than protein-based regulatory systems (6, 7, 105, 106). Riboswitches have also
become desirable therapeutic drug targets because they often control essential metabolic
pathways in pathogenic microbes (4, 5, 107). Broader consideration of the conformational
dynamics of larger types of effector molecules including organic cofactors, antibiotics, and their
analogs could aid efforts to engineer synthetic RNA-based regulatory platforms that function
robustly *in vivo*. This could also inform future efforts to develop synthetic antimetabolites that,
for example, elicit gene mis-regulation, instead of a more common strategy of creating inhibitors
for enzymes (108, 109).

The chemical diversity of corrinoids is intrinsically linked to a vast array of metabolic processes
and microbial interactions. Yet it remains unclear how microbes have evolved to cope with and
thrive on the assortment of natural corrinoid analogs, especially when compared to other primary
metabolites including organic cofactors, nucleotides and amino acids which typically require one
specific structural form for precise biological functions. We have gained new appreciation for the
impacts of chemical diversity on biological function by focusing on the Cbl-riboswitch with its
distinctively complex structure and regulatory mechanism, and by accounting for the often
overlooked biological and ecological roles of corrinoid analogs. Future studies into the evolution
of microbial molecular specificity for corrinoids may yield further insight into the nature of these
exceptionally versatile coenzymes.

Materials and methods

Cbl-riboswitch sequence analysis

Cbl-riboswitch sequences, chromosomal coordinates and regulon information were downloaded
from the RiboD online database (73) (Table S1). The genome of *Sporomusa ovata* was not
included in the RiboD database, so we used the RiboswitchScanner webserver to search for Cbl-
riboswitches in this organism (110, 111). Cbl-riboswitch aptamer sequences were manually
aligned by conserved secondary structures bounded by the 5' and 3' ends of the P1 stem (8, 85)
(Data S1). The P13-L13 stem loop and potential intrinsic transcriptional termination hairpin
structures of the expression platform were identified using secondary structure prediction tools in
RNAstructure 6.2 (112, 113). Intrinsic terminators were identified as stem loops directly
preceding a sequence of five or more consecutive uracil residues (114, 115). Sequence alignment
and annotation was carried out in JalView 2.11.1.4 (116). Cartoons of riboswitch secondary
structures were constructed using the StructureEditor program of RNAstructure 6.2 (113).

Corrinoid production, extraction, purification, and analysis

Cyanocobalamin, adenosylcobalamin, methylcobalamin, hydroxocobalamin, and
dicyanocobinamide were purchased from MilliporeSigma. All other corrinoids used in this study
were produced in bacterial cultures and purified in cyanated form as previously described (82,
95, 117, 118). For the experiments in Figures 4 and S5, corrinoids other than cobalamin were
chemically adenosylated to obtain the coenzyme (5'-deoxyadenosylated) form as previously
described (82, 117).

UV/Vis spectra were collected from corrinoid samples in UV/Vis-transparent 96-well microtiter
plates (greiner bio-one UV-STAR® 675801) using a BioTek Synergy 2 or Tecan Infinite M1000
Pro plate reader. To measure concentrations of corrinoid stock solutions, corrinoid samples were
diluted 10-fold in 10 mM sodium cyanide to obtain the dicyanated base-off form of the corrinoid.
The concentration of the dicyanated corrinoid was calculated using the extinction coefficient
$\varepsilon 580 = 10.1 \square mM^{-1} \square cm^{-1}$ (52, 119). For adenosylated corrinoids used in Figures 4 and S5, base-
on tendency at neutral pH was measured as the ratio of spectral absorbance at 525 nm and 458
nm in phosphate buffered saline solution pH 7.3 at 37°C (47, 59).

Plasmid and strain construction

Plasmids generated in this study were constructed with one-step isothermal assembly (120) and
introduced into *E. coli* strain XL1-Blue by heat shock transformation. Riboswitch reporter
plasmids were constructed in the shuttle vector pSG29 for single copy integration at the amyE
locus of the *B. subtilis* chromosome (121). Riboswitch DNA sequences were inserted between
the transcriptional start site of the constitutive P$_{Veg}$ promoter and the *gfp* translational start site of
pSG29. For riboswitch sequences that resulted in no detectable GFP signal under any conditions,
a synthetic ribosome binding site (RBS) sequence R0 (5'-
GATTAACTAATAAGGAGGACAAAC-3') from pSG29 was placed between the riboswitch
sequence and *gfp* translational start site (Table S1B-C).

Strains used in this study are listed in Table S1A. All *B. subtilis* riboswitch fluorescent reporter
strains and *B. subtilis* strains expressing *P. megaterium metE* and *metH* used in this study are
derived from the high-efficiency transformation strain SCK6, which has a xylose-inducible
competence gene cassette (122). Preparation of competent cells and transformations of all SCK6-

derived strains were performed as previously described (122). The strain KK642, which
constitutively overexpresses the corrinoid uptake genes, was constructed by deletion of gene
queG and replacement of the promoter and 5' untranslated region of the *btuFCDR* operon with
the P_{Veg} promoter and R0 RBS (121). *B. subtilis* genes *queG*, *btuR*, and *metE* were targeted for
deletion by recombination with kanamycin resistance cassettes containing flanking sequence
homology to each respective locus. Kanamycin resistance cassettes were PCR-amplified from
genomic DNA of *B. subtilis* strains BKK08910 ($\Delta queG::kan^R$), BKK33150 ($\Delta btuR::kan^R$), and
BKK13180 ($\Delta metE::kan^R$) (123). Kanamycin resistance cassettes were removed by Cre-Lox
recombination using plasmid pDR244 as previously described (123).

B. subtilis strains heterologously expressing *metE* and *metH* from *P. megaterium* were
constructed as follows. The *metE* and *metH* genes were PCR-amplified from genomic DNA of *P.*
megaterium DSM319 and cloned between the transcriptional start site and the *gfp* translational
start site of pSG29. The *P. megaterium metE* amplified fragment starts at the SAM-Cbl tandem
riboswitch in the 5' UTR and ends at the *metE* stop codon, whereas the *P. megaterium metH*
fragment starts at the *metH* RBS, which is composed of 20 nucleotides preceding the *metH*
translational start site and ends at the *metH* stop codon.

Riboswitch reporter plasmids and plasmids containing *P. megaterium metE* and *metH* were
linearized by restriction enzyme digest with ScaI-HF (New England Biolabs) and selected for
integration at the *amyE* locus of *B. subtilis* by plating on lysogeny broth (LB) agar with 100
µg/mL spectinomycin. Colonies were screened for integration at *amyE* by colony PCR. All
stocks of bacterial strains were stored in 15% glycerol at -80 □C.

Intracellular corrinoid accumulation experiments

B. subtilis strains were inoculated from single colonies into 50 mL LB and grown with aeration
at 37 □C in a shaking incubator (Gyromax 737R, Amerex Instruments, Inc.) for 5-6 hours until
reaching an optical density at 600 nm (OD_{600}) of 1.0 to 1.5. Each culture was diluted 10-fold in
LB and split into 13 25 mL cultures containing 0, 25, 250, or 2500 picomoles of a cyanated
corrinoid (Cbl, pCbl, CreCba and Cbi). These cultures were incubated at 37 □C with aeration for
3-4 hours to a final OD_{600} of 1.5 to 2.0. The cells were pelleted by centrifugation at 4,000 g for
10 min. Cell pellets were rinsed three times by resuspension in 10 mL phosphate buffered saline
solution pH 7.3 followed by centrifugation. After the final centrifugation, tubes were wrapped in
aluminum foil to protect adenosylated corrinoids from exposure to light.

To extract intracellular corrinoids, cell pellets were resuspended in 5 mL of 100% methanol by
vigorous vortexing for 30 secs. Samples were stored at -80 □C until the next day. Frozen lysates
were heated in an 80 □C water bath for 1.5 hours, with 15 seconds of vortexing every 30
minutes. Methanol concentration of each sample was diluted to 10% by adding 45 mL of water,
and cell debris was pelleted by centrifugation at 4,000 g for 10 mins. The supernatants were used
for the subsequent steps.

All of the following steps were carried out in darkened rooms illuminated with red light to
preserve light-sensitive adenosylated corrinoid samples. Solid-phase extraction of adenosylated
corrinoids with Sep-Pak C18 cartridges (Waters) was performed as previously described (82).

Solvents were evaporated in a vacuum concentrator centrifuge (Savant SPD1010, Thermo
Scientific) at 45 □C and the samples were resuspended in 500 µL deionized water and passed
through 0.45 µm pore-size filters (Millex-HV, MilliporeSigma).

Corrinoids were analyzed on an Agilent 1200 series (high-performance liquid chromatography
system equipped with a diode array detector (Agilent Technologies). Samples were injected into
an Agilent Zorbax SB-Aq column (5-µm pore size, 4.6 by 150□mm). The following HPLC
method was used: solvent A, 0.1% formic acid–deionized water; solvent B, 0.1% formic acid–
methanol; flow rate of 1□mL/minute at 30°C; 25% to 34% solvent B for 11□minutes, followed
by a linear gradient of 34% to 50% solvent B over 2□minutes, followed by a linear gradient of
50% to 75% solvent B over 8 minutes.

Riboswitch fluorescent reporter assays

Corrinoid dose-response assays of riboswitch reporter strains were set up as follows. Saturated
cultures of the riboswitch reporter strain in LB were diluted 200-fold in LB and dispensed into
96-well microtiter plates (Corning Costar Assay Plate 3904) containing a range of concentrations
of various corrinoids. The plates were sealed with gas diffusible membranes (Breathe-Easy,
MilliporeSigma) and incubated at 37 □C for 4 to 5 hours in a benchtop heated plate shaker
(Southwest Science) at 1,200 revolution per minute (rpm). GFP fluorescence
(excitation/emission/bandwidth = 485/525/10 nm) and absorbance at 600 nm (A_{600}) were
measured on a Tecan Infinite M1000 Pro plate reader. The A_{600} measurements of uninoculated
medium and fluorescence measurements of the parental control strains lacking *gfp* were
subtracted from all readings. Data were plotted and analyzed in GraphPad Prism 9.

3D structural analysis of corrinoids and macromolecular models

Molecular models of cobalamin in the base-on and base-off/His-on state in complex with various
proteins and RNAs were downloaded from the Protein DataBank (PDB) (Table S2) (124). PDB
files were analyzed in UCSF Chimera 1.14 (125). Corrinoid molecular models were aligned with
each other by the central cobalt atom and coordinating nitrogen atoms of the corrin ring, using
the PDB ID 4GMA Cbl model as a reference. Corrinoid models were aligned within the binding
sites of riboswitch structures PDB IDs 4GMA, 4FRN, 4GXY, and 6VMY (51, 54, 55).

Methionine-dependent growth of B. subtilis *strains*

B. subtilis strains were streaked from frozen stocks onto LB agar plates and incubated overnight
at 37 □C for 14-18 hrs. Single colonies were used to inoculate 3 mL liquid starter cultures
containing Spizizen minimal medium supplemented with 0.02% D-glucose and 0.2% L-Histidine
(SMM) (126). Starter cultures of the *metH*-expressing strain were supplemented with 1 nM
CNCbl to support growth, whereas the *metE*-expressing strains were cultured in SMM without
CNCbl. Starter cultures were incubated overnight shaking (250 rpm, 37 □C) for 20 hours,
reaching cell density of about $OD_{600} = 1.0$. Starter cultures were diluted 500-fold by transferring
50 µL of starter culture to 25 mL of SMM. Then 75 µL of the diluted culture were dispensed into
wells of a 96 well microtiter plate containing 75 µL of SMM supplemented with 40 nM of
various corrinoids. Plates were sealed with gas diffusible membranes (Breathe-Easy,

MilliporeSigma) and incubated at 37 □C on the 'high shaking' setting of a BioTek Synergy2
plate reader. Growth kinetics and *metE* and *metH* expression were measured by A_{600} and GFP
fluorescence every 15 minutes for 72 hours. Data were plotted and analyzed in GraphPad Prism
9.

Acknowledgments

We thank past and present members of the Taga lab for support. In particular, we thank
Sebastian Gude, Zachary Hallberg, and Gordon Pherribo for critical reading of the manuscript;
Alexa Nicolas and Amanda Shelton for helpful discussions; Anna Grimaldo and Jong-Duk Park
for reagents. We thank Amrita Hazra, Arash Komeili, Michael Marletta, Kathleen Ryan, and
Matthew Traxler for helpful discussions. We especially thank Ming Hammond for critical
reading of the manuscript and helpful advice. We are grateful to the labs of Benjamin Blackman,
John Deuber, Krishna Niyogi, and Matthew Traxler for use of their equipment. This work was
supported by NIH grants R01GM114535 and R35GM139633 to M.E.T. We acknowledge that
this work was conducted on the ancestral and unceded land of the Ohlone people.

Author contributions

525 Conceptualization, K.J.K., F.J.W., K.C.M., and M.E.T.; Methodology, K.J.K., F.J.W., O.M.S.,
526 K.C.M., and M.E.T.; Validation: R.R.P.; Investigation, K.J.K. and L.V.I.; Writing – Original
527 Draft, K.J.K. and M.E.T.; Writing – Review & Editing, K.J.K., O.M.S., K.C.M., and M.E.T.;
528 Funding Acquisition, M.E.T.; Resources, O.M.S. and K.C.M.; Supervision, M.E.T.

Declaration of interests

531 The authors declare no competing interests.

References

1. Cech Thomas R & Steitz Joan A (2014) The Noncoding RNA Revolution - Trashing Old
Rules to Forge New Ones. *Cell* 157(1):77-94.
2. Roth A & Breaker RR (2009) The structural and functional diversity of metabolite-
binding riboswitches. *Annu Rev Biochem* 78:305-334.
3. Winkler WC & Breaker RR (2005) Regulation of bacterial gene expression by
riboswitches. *Annu Rev Microbiol* 59:487-517.
4. Panchal V & Brenk R (2021) Riboswitches as Drug Targets for Antibiotics. *Antibiotics*
(Basel, Switzerland) 10(1).
5. Quereda JJ & Cossart P (2017) Regulating Bacterial Virulence with RNA. *Annu Rev*
Microbiol 71:263-280.
6. Wrist A, Sun W, & Summers RM (2020) The theophylline aptamer: 25 years as an
important tool in cellular engineering research. *ACS Synth Biol* 9(4):682-697.
7. Hallberg ZF, Su Y, Kitto RZ, & Hammond MC (2017) Engineering and In Vivo
Applications of Riboswitches. *Annu Rev Biochem* 86:515-539.
8. McCown PJ, Corbino KA, Stav S, Sherlock ME, & Breaker RR (2017) Riboswitch
diversity and distribution. *RNA* 23(7):995-1011.
9. Nahvi A*, et al.* (2002) Genetic control by a metabolite binding mRNA. *Chem Biol*
9(9):1043.
10. Licht S, Gerfen GJ, & Stubbe J (1996) Thiyl radicals in ribonucleotide reductases.
Science 271(5248):477-481.
11. Berg IA, Kockelkorn D, Buckel W, & Fuchs G (2007) A 3-hydroxypropionate/4-
hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea. *Science*
318(5857):1782-1786.
12. Ljungdahl LG (1986) The autotrophic pathway of acetate synthesis in acetogenic
bacteria. *Annu Rev Microbiol* 40:415-450.
13. Krasotkina J, Walters T, Maruya KA, & Ragsdale SW (2001) Characterization of the
B12- and iron-sulfur-containing reductive dehalogenase from Desulfitobacterium
chlororespirans. *J Biol Chem* 276(44):40991-40997.
14. Scarlett FA & Turner JM (1976) Microbial metabolism of amino alcohols. Ethanolamine
catabolism mediated by coenzyme B12-dependent ethanolamine ammonia-lyase in
Escherichia coli and Klebsiella aerogenes. *J Gen Microbiol* 95(1):173-176.
15. Stupperich E & Konle R (1993) Corrinoid-Dependent Methyl Transfer Reactions Are
Involved in Methanol and 3,4-Dimethoxybenzoate Metabolism by Sporomusa ovata.
Appl Environ Microbiol 59(9):3110-3116.
16. Jeter RM (1990) Cobalamin-dependent 1,2-propanediol utilization by Salmonella
typhimurium. *J Gen Microbiol* 136(5):887-896.
17. Forage RG & Foster MA (1982) Glycerol fermentation in Klebsiella pneumoniae:
functions of the coenzyme B12-dependent glycerol and diol dehydratases. *J Bacteriol*
149(2):413-419.
18. Erb TJ, Rétey J, Fuchs G, & Alber BE (2008) Ethylmalonyl-CoA mutase from
Rhodobacter sphaeroides defines a new subclade of coenzyme B12-dependent acyl-CoA
mutases. *J Biol Chem* 283(47):32283-32293.
19. Korotkova N, Chistoserdova L, Kuksa V, & Lidstrom ME (2002) Glyoxylate
regeneration pathway in the methylotroph Methylobacterium extorquens AM1. *J*
Bacteriol 184(6):1750-1758.

20. Ferguson DJ, Jr. & Krzycki JA (1997) Reconstitution of trimethylamine-dependent
coenzyme M methylation with the trimethylamine corrinoid protein and the isozymes of
methyltransferase II from Methanosarcina barkeri. *J Bacteriol* 179(3):846-852.
21. Barker HA (1985) beta-Methylaspartate-glutamate mutase from Clostridium
tetanomorphum. *Methods Enzymol* 113:121-133.
22. Chen HP, Wu SH, Lin YL, Chen CM, & Tsay SS (2001) Cloning, sequencing,
heterologous expression, purification, and characterization of adenosylcobalamin-
dependent D-ornithine aminomutase from Clostridium sticklandii. *J Biol Chem*
276(48):44744-44750.
23. Chang CH & Frey PA (2000) Cloning, sequencing, heterologous expression, purification,
and characterization of adenosylcobalamin-dependent D-lysine 5, 6-aminomutase from
Clostridium sticklandii. *J Biol Chem* 275(1):106-114.
24. Kim HJ, *et al.* (2013) GenK-catalyzed C-6' methylation in the biosynthesis of
gentamicin: isolation and characterization of a cobalamin-dependent radical SAM
enzyme. *J Am Chem Soc* 135(22):8093-8096.
25. Knox HL, *et al.* (2021) Structural basis for non-radical catalysis by TsrM, a radical SAM
methylase. *Nat Chem Biol* 17(4):485-491.
26. Choi SC & Bartha R (1993) Cobalamin-mediated mercury methylation by *Desulfovibrio*
desulfuricans LS. *Appl Environ Microbiol* 59(1):290-295.
27. Choi SC, Chase T, Jr., & Bartha R (1994) Enzymatic catalysis of mercury methylation by
Desulfovibrio desulfuricans LS. *Appl Environ Microbiol* 60(4):1342-1346.
28. Parks JM, *et al.* (2013) The genetic basis for bacterial mercury methylation. *Science*
339(6125):1332-1335.
29. Wang PH, *et al.* (2020) Retroconversion of estrogens into androgens by bacteria via a
cobalamin-mediated methylation. *Proc Natl Acad Sci U S A* 117(3):1395-1403.
30. Romine MF, *et al.* (2017) Elucidation of roles for vitamin B12 in regulation of folate,
ubiquinone, and methionine metabolism. *Proc Natl Acad Sci U S A* 114(7):E1205-e1214.
31. Bridwell-Rabb J, Zhong A, Sun HG, Drennan CL, & Liu HW (2017) A B12-dependent
radical SAM enzyme involved in oxetanocin A biosynthesis. *Nature* 544(7650):322-326.
32. Allen KD & Wang SC (2014) Initial characterization of Fom3 from Streptomyces
wedmorensis: The methyltransferase in fosfomycin biosynthesis. *Arch Biochem Biophys*
543:67-73.
33. Pierre S, *et al.* (2012) Thiostrepton tryptophan methyltransferase expands the chemistry
of radical SAM enzymes. *Nat Chem Biol* 8(12):957-959.
34. Marous DR, *et al.* (2015) Consecutive radical S-adenosylmethionine methylations form
the ethyl side chain in thienamycin biosynthesis. *Proc Natl Acad Sci U S A*
112(33):10354-10358.
35. Takano H, *et al.* (2015) Role and Function of LitR, an Adenosyl B12-Bound Light-
Sensitive Regulator of Bacillus megaterium QM B1551, in Regulation of Carotenoid
Production. *J Bacteriol* 197(14):2301-2315.
36. Takano H, *et al.* (2011) Involvement of CarA/LitR and CRP/FNR family transcriptional
regulators in light-induced carotenoid production in Thermus thermophilus. *J Bacteriol*
193(10):2451-2459.
37. Jost M, *et al.* (2015) Structural basis for gene regulation by a B12-dependent
photoreceptor. *Nature* 526(7574):536-541.

38. Rodionov DA, Vitreschak AG, Mironov AA, & Gelfand MS (2003) Comparative
genomics of the vitamin B12 metabolism and regulation in prokaryotes. *J Biol Chem*
278(42):41148-41159.
39. Pavlova N, Kaloudas D, & Penchovsky R (2019) Riboswitch distribution, structure, and
function in bacteria. *Gene* 708:38-48.
40. Kennedy KJ & Taga ME (2020) Cobamides. *Curr Biol* 30(2):R55-r56.
41. Moore SJ, *et al.* (2013) Elucidation of the anaerobic pathway for the corrin component of
cobalamin (vitamin B12). *Proc Natl Acad Sci U S A* 110(37):14906-14911.
42. Battersby AR (2000) Tetrapyrroles: the pigments of life. *Nat Prod Rep* 17(6):507-526.
43. Hazra AB, *et al.* (2015) Anaerobic biosynthesis of the lower ligand of vitamin B12. *Proc*
Natl Acad Sci U S A 112(34):10792-10797.
44. Chan CH & Escalante-Semerena JC (2012) ArsAB, a novel enzyme from *Sporomusa*
ovata activates phenolic bases for adenosylcobamide biosynthesis. *Mol Microbiol*
81(4):952-967.
45. Newmister SA, Chan CH, Escalante-Semerena JC, & Rayment I (2012) Structural
insights into the function of the nicotinate mononucleotide:phenol/p-cresol
phosphoribosyltransferase (ArsAB) enzyme from *Sporomusa ovata*. *Biochemistry*
51(43):8571-8582.
46. Yan J, *et al.* (2018) Purinyl-cobamide is a native prosthetic group of reductive
dehalogenases. *Nat Chem Biol* 14(1):8-14.
47. Hoffmann B, *et al.* (2000) Native corrinoids from *Clostridium cochlearium* are
adeninylcobamides: spectroscopic analysis and identification of pseudovitamin B12 and
factor A. *J Bacteriol* 182(17):4773-4782.
48. Polaski JT, Webster SM, Johnson JE, Jr., & Batey RT (2017) Cobalamin riboswitches
exhibit a broad range of ability to discriminate between methylcobalamin and
adenosylcobalamin. *J Biol Chem* 292(28):11650-11658.
49. Polaski JT, Kletzien OA, Drogalis LK, & Batey RT (2018) A functional genetic screen
reveals sequence preferences within a key tertiary interaction in cobalamin riboswitches
required for ligand selectivity. *Nucleic Acids Res* 46(17):9094-9105.
50. Gallo S, Oberhuber M, Sigel RK, & Kräutler B (2008) The corrin moiety of coenzyme
B12 is the determinant for switching the *btuB* riboswitch of *E. coli*. *Chembiochem*
9(9):1408-1414.
51. Chan CW & Mondragón A (2020) Crystal structure of an atypical cobalamin riboswitch
reveals RNA structural adaptability as basis for promiscuous ligand binding. *Nucleic*
Acids Res 48(13):7569-7583.
52. Stupperich E, Eisinger HJ, & Kräutler B (1988) Diversity of corrinoids in acetogenic
bacteria. P-cresolylcobamide from *Sporomusa ovata*, 5-methoxy-6-
methylbenzimidazolylcobamide from *Clostridium formicoaceticum* and vitamin B12
from *Acetobacterium woodii*. *Eur J Biochem* 172(2):459-464.
53. Crofts TS, Seth EC, Hazra AB, & Taga ME (2013) Cobamide structure depends on both
lower ligand availability and CobT substrate specificity. *Chem Biol* 20(10):1265-1274.
54. Johnson JE, Jr., Reyes FE, Polaski JT, & Batey RT (2012) B12 cofactors directly
stabilize an mRNA regulatory switch. *Nature* 492(7427):133-137.
55. Peselis A & Serganov A (2013) Structural insights into ligand binding and gene
expression control by an adenosylcobalamin riboswitch. *Nat Struct Mol Biol*
19(11):1182-1184.

56.   Barker HA*, et al.* (1960) Isolation and properties of crystalline cobamide coenzymes
670   containing benzimidazole or 5,6-dimethylbenzimidazole. *J Biol Chem* 235:480-488.
671   57.   Ladd JN, Hogenkamp HP, & Barker HA (1961) Structure of cobamide coenzymes:
672   influence of pH on absorption spectra and ionophoretic mobilities. *J Biol Chem*
673   236:2114-2118.
674   58.   Brown KL & Hakimi JM (1986) Heteronuclear NMR studies of cobalamins. 4. alpha-
675   Ribazole-3'-phosphate and the nucleotide loop of base-on cobalamins. *J Am Chem Soc*
676   108(3):496-503.
677   59.   Fieber W*, et al.* (2002) Pseudocoenzyme B12 and adenosyl-Factor A: Electrochemical
678   synthesis and spectroscopic analysis of two natural B12 coenzymes with predominantly
679   'base-off' constitution. *Helvetica Chimica Acta* 85(3):927-944.
680   60.   Marques HM, Zou X, & Brown KL (2000) The solution structure of adenosylcobalamin
681   and adenosylcobinamide determined by nOe-restrained molecular dynamics simulations.
682   *Journal of Molecular Structure* 520(1):75-95.
683   61.   Sussman D & Wilson C (2000) A water channel in the core of the vitamin B12 RNA
684   aptamer. *Structure* 8(7):719-727.
685   62.   Mancia F*, et al.* (1996) How coenzyme B12 radicals are generated: the crystal structure
686   of methylmalonyl-coenzyme A mutase at 2 A resolution. *Structure* 4(3):339-350.
687   63.   Froese DS*, et al.* (2010) Structures of the human GTPase MMAA and vitamin B12-
688   dependent methylmalonyl-CoA mutase and insight into their complex formation. *J Biol*
689   *Chem* 285(49):38204-38213.
690   64.   Drennan CL, Huang S, Drummond JT, Matthews RG, & Ludwig ML (1994) How a
691   protein binds B12: A 3.0 A X-ray structure of B12-binding domains of methionine
692   synthase. *Science* 266(5191):1669-1674.
693   65.   Dowling DP*, et al.* (2016) Molecular basis of cobalamin-dependent RNA modification.
694   *Nucleic Acids Res* 44(20):9965-9976.
695   66.   Payne KA*, et al.* (2015) Epoxyqueuosine Reductase Structure Suggests a Mechanism for
696   Cobalamin-dependent tRNA Modification. *J Biol Chem* 290(46):27572-27581.
697   67.   Payne KA*, et al.* (2015) Reductive dehalogenase structure suggests a mechanism for
698   B12-dependent dehalogenation. *Nature* 517(7535):513-516.
699   68.   Jost M, Born DA, Cracan V, Banerjee R, & Drennan CL (2015) Structural Basis for
700   Substrate Specificity in Adenosylcobalamin-dependent Isobutyryl-CoA Mutase and
701   Related Acyl-CoA Mutases. *J Biol Chem* 290(45):26882-26898.
702   69.   Gruber K, Reitzer R, & Kratky C (2001) Radical shuttling in a protein: Ribose
703   pseudorotation controls alkyl-radical transfer in the coenzyme B12-dependent enzyme
704   glutamate mutase. *Angewandte Chemie International Edition* 40(18):3377-3380.
705   70.   Berkovitch F*, et al.* (2004) A locking mechanism preventing radical damage in the
706   absence of substrate, as revealed by the x-ray structure of lysine 5,6-aminomutase. *Proc*
707   *Natl Acad Sci U S A* 101(45):15870-15875.
708   71.   Lussier A, Bastet L, Chauvier A, & Lafontaine DA (2015) A kissing loop is important for
709   btuB riboswitch ligand sensing and regulatory control. *J Biol Chem* 290(44):26739-
710   26751.
711   72.   Polaski JT, Holmstrom ED, Nesbitt DJ, & Batey RT (2016) Mechanistic Insights into
712   Cofactor-Dependent Coupling of RNA Folding and mRNA Transcription/Translation by
713   a Cobalamin Riboswitch. *Cell Rep* 15(5):1100-1110.

73. Mukherjee S, *et al.* (2019) RiboD: a comprehensive database for prokaryotic
riboswitches. *Bioinformatics* 35(18):3541-3543.
74. Sudarsan N, *et al.* (2006) Tandem riboswitch architectures exhibit complex gene control
functions. *Science* 314(5797):300-304.
75. Moore SJ, Mayer MJ, Biedendieck R, Deery E, & Warren MJ (2014) Towards a cell
factory for vitamin B12 production in Bacillus megaterium: bypassing of the cobalamin
riboswitch control elements. *New biotechnology* 31(6):553-561.
76. Collins HF, *et al.* (2013) *Bacillus megaterium* has both a functional BluB protein
required for DMB synthesis and a related flavoprotein that forms a stable radical species.
PLoS One 8(2):e55708.
77. Biedendieck R, *et al.* (2010) Metabolic engineering of cobalamin (vitamin B12)
production in Bacillus megaterium. *Microb Biotechnol* 3(1):24-37.
78. Ma AT, Tyrell B, & Beld J (2020) Specificity of cobamide remodeling, uptake and
utilization in Vibrio cholerae. *Molecular Microbiology* 113(1):89-102.
79. Degnan PH, Barry NA, Mok KC, Taga ME, & Goodman AL (2014) Human gut microbes
use multiple transporters to distinguish vitamin B12 analogs and compete in the gut. *Cell
Host Microbe* 15(1):47-57.
80. Tanioka Y, *et al.* (2010) Methyladeninylcobamide functions as the cofactor of
methionine synthase in a Cyanobacterium, *Spirulina platensis* NIES-39. *FEBS Lett*
584(14):3223-3226.
81. Helliwell KE, *et al.* (2016) Cyanobacteria and eukaryotic algae use different chemical
variants of vitamin B12. *Curr Biol* 26(8):999-1008.
82. Sokolovskaya OM, *et al.* (2019) Cofactor Selectivity in Methylmalonyl Coenzyme A
Mutase, a Model Cobamide-Dependent Enzyme. *mBio* 10(5):e01303-01319.
83. Nahvi A, Barrick JE, & Breaker RR (2004) Coenzyme B12 riboswitches are widespread
genetic control elements in prokaryotes. *Nucleic Acids Res* 32(1):143-150.
84. Gallo S, Mundwiler S, Alberto R, & Sigel RK (2011) The change of corrin-amides to
carboxylates leads to altered structures of the B12-responding *btuB* riboswitch. *Chem
Commun (Camb)* 47(1):403-405.
85. Choudhary PK, *et al.* (2013) Diversity of cobalamin riboswitches in the corrinoid-
producing organohalide respirer *Desulfitobacterium hafniense*. *J Bacteriol* 195(22):5186-
5195.
86. Di Girolamo PM, Kadner RJ, & Bradbeer C (1971) Isolation of vitamin B 12 transport
mutants of Escherichia coli. *J Bacteriol* 106(3):751-757.
87. Kashket S, Kaufman JT, & Beck WS (1962) The metabolic functions of vitamin B12. III.
Vitamin B12 binding in Lactobacillus leichmannii and other lactobacilli. *Biochim
Biophys Acta* 64:447-457.
88. Giannella RA, Broitman SA, & Zamcheck N (1971) Vitamin B12 uptake by intestinal
microorganisms: mechanism and relevance to syndromes of intestinal bacterial
overgrowth. *J Clin Invest* 50(5):1100-1107.
89. Price IR, Grigg JC, & Ke A (2014) Common themes and differences in SAM recognition
among SAM riboswitches. *Biochim Biophys Acta* 1839(10):931-938.
90. Huang L, Liao TW, Wang J, Ha T, & Lilley DMJ (2020) Crystal structure and ligand-
induced folding of the SAM/SAH riboswitch. *Nucleic Acids Res* 48(13):7545-7556.
91. Weickhmann AK, *et al.* (2019) The structure of the SAM/SAH-binding riboswitch.
Nucleic Acids Res 47(5):2654-2665.

92. Ma B, Bai G, Nussinov R, Ding J, & Wang YX (2021) Conformational Ensemble of
TteAdoCbl Riboswitch Provides Stable Structural Elements for Conformation Selection
and Population Shift in Cobalamin Recognition. *The journal of physical chemistry. B*
125(10):2589-2596.
93. Trausch JJ & Batey RT (2014) A disconnect between high-affinity binding and efficient
regulation by antifolates and purines in the tetrahydrofolate riboswitch. *Chem Biol*
21(2):205-216.
94. Shelton AN, Lyu X, & Taga ME (2020) Flexible cobamide metabolism in *Clostridioides*
(*Clostridium*) *difficile* 630 Δerm. *J Bacteriol* 202(2).
95. Mok KC & Taga ME (2013) Growth Inhibition of *Sporomusa ovata* by Incorporation of
Benzimidazole Bases into Cobamides. *J Bacteriol* 195(9):1902-1911.
96. Shelton AN*, et al.* (2019) Uneven distribution of cobamide biosynthesis and dependence
in bacteria predicted by comparative genomics. *ISME J* 13(3):789-804.
97. Fedosov SN, Fedosova NU, Kräutler B, Nexo E, & Petersen TE (2007) Mechanisms of
discrimination between cobalamins and their natural analogues during their binding to the
specific B12-transporting proteins. *Biochemistry* 46(21):6446-6458.
98. Chan CH*, et al.* (2014) Dissecting cobamide diversity through structural and functional
analyses of the base-activating CobT enzyme of *Salmonella enterica*. *Biochim Biophys*
Acta 1840(1):464-475.
99. Hazra AB, Tran JL, Crofts TS, & Taga ME (2013) Analysis of substrate specificity in
CobT homologs reveals widespread preference for DMB, the lower axial ligand of
vitamin B12. *Chem Biol* 20(10):1275-1285.
100. Cheong CG, Escalante-Semerena JC, & Rayment I (2001) Structural investigation of the
biosynthesis of alternative lower ligands for cobamides by nicotinate
mononucleotide:5,6-dimethylbenzimidazole phosphoribosyltransferase from *Salmonella*
enterica. *J Biol Chem* 276(40):37612-37620.
101. Cheong CG, Escalante-Semerena JC, & Rayment I (1999) The three-dimensional
structures of nicotinate mononucleotide:5,6-dimethylbenzimidazole
phosphoribosyltransferase (CobT) from *Salmonella typhimurium* complexed with 5,6-
dimethybenzimidazole and its reaction products determined to 1.9 A resolution.
Biochemistry 38(49):16125-16135.
102. Trzebiatowski JR & Escalante-Semerena JC (1997) Purification and characterization of
CobT, the nicotinate-mononucleotide:5,6-dimethylbenzimidazole
phosphoribosyltransferase enzyme from *Salmonella typhimurium* LT2. *J Biol Chem*
272(28):17662-17667.
103. Trzebiatowski JR, O'Toole GA, & Escalante-Semerena JC (1994) The *cobT* gene of
Salmonella typhimurium encodes the NaMN: 5,6-dimethylbenzimidazole
phosphoribosyltransferase responsible for the synthesis of N1-(5-phospho-alpha-D-
ribosyl)-5,6-dimethylbenzimidazole, an intermediate in the synthesis of the nucleotide
loop of cobalamin. *J Bacteriol* 176(12):3568-3575.
104. Sokolovskaya OM, Shelton AN, & Taga ME (2020) Sharing vitamins: Cobamides unveil
microbial interactions. *Science* 369(6499).
105. Villa JK, Su Y, Contreras LM, & Hammond MC (2018) Synthetic Biology of Small
RNAs and Riboswitches. *Microbiol Spectr* 6(3).
106. Galizi R & Jaramillo A (2019) Engineering CRISPR guide RNA riboswitches for in vivo
applications. *Curr Opin Biotechnol* 55:103-113.

806 107. Mansjö M & Johansson J (2011) The riboflavin analog roseoflavin targets an FMN-
807   riboswitch and blocks Listeria monocytogenes growth, but also stimulates virulence
808   gene-expression and infection. *RNA Biol* 8(4):674-680.

809 108. Kräutler B (2020) Antivitamins B12 - Some Inaugural Milestones. *Chemistry*
810   26(67):15438-15445.

811 109. Zelder F, Sonnay M, & Prieto L (2015) Antivitamins for Medicinal Applications.
812   *Chembiochem* 16(9):1264-1278.

813 110. Mukherjee S & Sengupta S (2016) Riboswitch Scanner: an efficient pHMM-based web-
814   server to detect riboswitches in genomic sequences. *Bioinformatics* 32(5):776-778.

815 111. Singh P, Bandyopadhyay P, Bhattacharya S, Krishnamachari A, & Sengupta S (2009)
816   Riboswitch detection using profile hidden Markov models. *BMC Bioinformatics* 10:325.

817 112. Mathews DH*, et al.* (2004) Incorporating chemical modification constraints into a
818   dynamic programming algorithm for prediction of RNA secondary structure. *Proc Natl*
819   *Acad Sci U S A* 101(19):7287-7292.

820 113. Reuter JS & Mathews DH (2010) RNAstructure: software for RNA secondary structure
821   prediction and analysis. *BMC Bioinformatics* 11:129.

822 114. Wilson KS & von Hippel PH (1995) Transcription termination at intrinsic terminators:
823   the role of the RNA hairpin. *Proc Natl Acad Sci U S A* 92(19):8793-8797.

824 115. Gusarov I & Nudler E (1999) The mechanism of intrinsic transcription termination. *Mol*
825   *Cell* 3(4):495-504.

826 116. Waterhouse AM, Procter JB, Martin DM, Clamp M, & Barton GJ (2009) Jalview Version
827   2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics*
828   25(9):1189-1191.

829 117. Sokolovskaya OM*, et al.* (2020) Naturally occurring cobalamin (B12) analogs can
830   function as cofactors for human methylmalonyl-CoA mutase. *Biochimie*.

831 118. Yi S*, et al.* (2012) Versatility in corrinoid salvaging and remodeling pathways supports
832   corrinoid-dependent metabolism in *Dehalococcoides mccartyi*. *Appl Environ Microbiol*
833   78(21):7745-7752.

834 119. Mok KC*, et al.* (2020) Identification of a novel cobamide remodeling enzyme in the
835   beneficial human gut bacterium *Akkermansia muciniphila*. *mBio* 11(6):e02507-02520.

836 120. Gibson DG*, et al.* (2009) Enzymatic assembly of DNA molecules up to several hundred
837   kilobases. *Nature methods* 6(5):343-345.

838 121. Guiziou S*, et al.* (2016) A part toolbox to tune genetic expression in Bacillus subtilis.
839   *Nucleic Acids Res* 44(15):7495-7508.

840 122. Zhang XZ & Zhang Y (2011) Simple, fast and high-efficiency transformation system for
841   directed evolution of cellulase in Bacillus subtilis. *Microb Biotechnol* 4(1):98-105.

842 123. Koo BM*, et al.* (2017) Construction and Analysis of Two Genome-Scale Deletion
843   Libraries for Bacillus subtilis. *Cell Syst* 4(3):291-305.e297.

844 124. Berman HM*, et al.* (2000) The Protein Data Bank. *Nucleic Acids Research* 28(1):235-
845   242.

846 125. Pettersen EF*, et al.* (2004) UCSF Chimera--a visualization system for exploratory
847   research and analysis. *Journal of computational chemistry* 25(13):1605-1612.

848 126. Anagnostopoulos C & Spizizen J (1961) Requirements for transformation in Bacillus
849   subtilis. *J Bacteriol* 81(5):741-746.

Figure 1 – Main corrinoids and riboswitches examined in this study. (A) Chemical structure
of cobalamin, also called vitamin B_{12}. The gray solid line delineates the corrinoid tail region,
which contains the variable lower ligand group in the gray dashed-line box. The R group is the
upper ligand. (B) Names, lower ligand structures and abbreviations of the corrinoids used
throughout this study. Cbl, pCbl and CreCba are 'complete corrinoids' (cobamides). Cbi is an
'incomplete corrinoid' that lacks the phosphoribosyl and lower ligand groups as indicated by the
bracket and wavy line in the corrinoid tail in panel A. (C) Secondary structural model of the Cbl-
riboswitch upstream of *B. subtilis btuF*. Base paired stems (P), loops (L), and junctions (J) are
labeled. The P6 accessory region is highly variable in length and number of stems across Cbl-
riboswitch sequences, often containing up to six paired regions. In this example, P6 accessory
consists of only two paired regions. A kissing loop interaction (KL, dashed line) occurs between
L5 of the aptamer and L13 of the expression platform. M1 indicates mutations C107U and
C108U, which are examined in Figure 2F. (D) Secondary structural alignments of seven
representative Cbl-riboswitches examined in this study. Row "2° str" indicates paired bases as
parentheses, loops and junctions as periods, and kissing loop base pairs by brackets. PDB
numbers or abbreviations are given for riboswitches from the following organisms and genes:
4GMA, *Thermanaerobacter tencongensis*; *S.the, Symbiobacterium thermophilum cblS*; *B.sub*, *B.
subtilis btuF; P.meg*, *Priestia megaterium metE*; *D.haf*, *Desulfitobacterium hafniense* DSY0087;
S.ova, *Sporomusa ovata btuB2*; *V.par*, *Veillonella parvula mutA*. Non-conserved sequences
between P13 and the start codon are indicated by nucleotide sequence length (nt), but the actual
sequences were omitted for clarity.

Figure 2 – Characterization of a live cell Cbl-riboswitch reporter system. (A-D) Intracellular
accumulation of corrinoids in *Bacillus subtilis* strains containing the wild type (diamonds) or
constitutively overexpressed (circles) corrinoid uptake genes *btuFCDR*. (E-G) Dose-responses of
B. subtilis btuF GFP riboswitch reporters. 'Mutant' riboswitch refers to the M1 mutant version of
the *B. subtilis btuF* riboswitch (Fig. 1C). Data points in panels A-D represent single
measurements from one representative experiment. Data points and error bars in panels E-G
represent mean and standard deviation of four independent replicates, and horizontal dotted lines
demarcate no change in expression.

Figure 3 – Corrinoid specificity among 38 Cbl-riboswitches. (A,B) Corrinoid dose-responses
of two riboswitch reporter strains with distinct corrinoid specificities, representing the semi-
selective (A) and promiscuous (B) types. (C-E) Pairwise comparisons of GFP fold repression
induced by 100 nM Cbl versus (C) pCbl, (D) Cbi, and (E) CreCba. Data points are colored by
taxonomic class with the number of riboswitches analyzed from each class indicated in
parentheses. Vertical and horizontal gray dashed lines demarcate a response to only one of the
two corrinoids. Diagonal line demarcates equal response to two corrinoids. Data points and error
bars in panels A and B represent mean and standard deviation of ten experiments for *P.
megaterium metE* and six experiments for *V. parvula mutA*. Data points and error bars in panels
C-E represent mean and range of at least two independent experiments for each riboswitch.

Figure 4 – Corrinoid base-on tendency correlates with corrinoid potency. (A) Sixteen adenosylated corrinoids used to test the relationship between corrinoid base-on tendency and corrinoid potency. (B) Semi-selective riboswitches *P. megaterium metE* and *D. hafniense* DSY0087 and (C) promiscuous riboswitches *V. parvula mutA* and *S. ovata btuB2* were used to measure the response to each corrinoid. Base-on tendency was measured as the ratio between spectral absorbance at 525 and 458 nm in a pH 7.3 solution. Absorbance spectra are displayed in Figure S5. The cumulative corrinoid response was measured as the area under the dose-response curve of the riboswitch reporter strain for that corrinoid ($AUC_{corrinoid}$) relative to its dose-response to Cbl (AUC_{Cbl}). Corrinoids with benzimidazole (blue circles), azabenzimidazole (purple circles), and purine (red circles) lower ligands can adopt the base-on conformation. The corrinoids that are unable to adopt the base-on state (CreCba, [Phe]Cba and Cbi) are represented by empty circles. Trendlines in B and C were fit to data points of corrinoids with benzimidazole, azabenzimidazole and purine lower ligands, with strictly base-off corrinoids excluded. Each data point represents a single measurement of riboswitch dose-response and corrinoid base-on tendency.

Figure 5 – Distinct tail positions among base-on and base-off corrinoids may impact binding of corrinoids to riboswitches. 3D alignments of Cbl structural models derived from published X-ray crystal structures of (A) base-on Cbl in complex with RNAs and (B) base-off Cbl in complex with proteins. The Cbl-binding site in the X-ray crystal structure of the *Thermoanaerobacter tengcongensis* Cbl-riboswitch (PDB ID 4GMA) is depicted with (C) base-on and (D) base-off Cbl alignments. Cbl models were aligned by the cobalt and coordinating nitrogen atoms in the corrin ring. Structures of the corrin ring, cobalt, and tail of Cbl are colored in black, blue, and green, respectively. Upper ligand structures of Cbl were omitted for clarity. Riboswitch RNA structures are depicted as space-filled models with the L5-L13 kissing loop in pink and the rest of the RNA in gray.

Figure 6 – Corrinoid specificities of the *P. megaterium metE* riboswitch and *P. megaterium* MetH enzyme are aligned. Expression of the Cbl-independent methionine synthase MetE and enzymatic activity of the Cbl-dependent methionine synthase MetH are potentially impacted by the presence of (A) MetH-compatible corrinoids and (B) MetH-incompatible corrinoids. (C) The *metH*-expressing strain (red) is a *B. subtilis* Δ*metE::loxP* Δ*queG::loxP* P_{Veg}-*btuFCDR* strain heterologously expressing *P. megaterium metH*. The *metE*-expressing strain (blue) is a *B. subtilis* Δ*metE::loxP* Δ*queG::loxP* P_{Veg}-*btuFCDR* strain heterologously expressing *P. megaterium metE* downstream from a SAM-Cbl tandem riboswitch (RS). Each methionine synthase gene is constitutively expressed from the P_{Veg} promoter and is transcriptionally fused to *gfp*. (D) Growth rates of the *metE*-expressing strain (blue circles) and *metH*-expressing strain (red squares) strains were measured in methionine-dependent culture conditions containing 20 nM corrinoid. (E) Expression of *metE* was measured as GFP fluorescence per OD_{600} in medium lacking methionine, supplemented with 20 nM corrinoids. Data points are individual measurements and black horizontal lines represent the mean of the four replicate measurements.

Legends for supplemental items

Figure S1 – The *B. subtilis btuF* Cbl-riboswitch is sensitive to Cbl with 5'-deoxyadenosyl,
but not methyl, hydroxyl, or cyanyl upper ligand moieties. Dose responses of *B. subtilis btuF*
Cbl-riboswitch reporter strains with (A) adenosylcobalamin (AdoCbl), (B) methylcobalamin
(MeCbl), (C) hydroxocobalamin (OHCbl), and (D) cyanocobalamin (CNCbl). Data points and
error bars represent mean and standard deviation of four independent replicates. Horizontal
dotted lines demarcate no change in expression. Note that the two dose-response curves in panel
A are overlapping.

Figure S2 – Corrinoid dose-responses of riboswitch reporter strains. Riboswitch names
(species and gene downstream gene) are indicated for each dose-response. Data points are
plotted for at least two independent experiments for each strain. Lines connect mean values. (A)
Riboswitches from the class Bacilli. Species include *Bacillus subtilis*, *Priestia megaterium*, and
Alkalihalobacillus halodurans. (B) Riboswitches from the class Negativicutes. Species include
Sporomusa ovata, *Veillonella parvula*, and *Selenomonas sputigena*. (C) Riboswitches from the
classes Clostridia, Deltaproteobacteria, and Bacteroidia. *Clostridioides difficile*, *Clostridium*
novyi, *Desfulitobacterium hafniense*, and *Symbiobacterium thermophilum* are members of
Clostridia. *Desulfobulbus propionicus* and *Pelobacter propoionicus* are members of
Deltaproteobacteria. *Bacteroides thetaiotaomicron* is a member of Bacteroidia.

Figure S3 – Cbl-riboswitch specificity does not cluster by regulatory gene target function.
Pairwise comparisons of GFP fold repression induced by 100 nM doses of Cbl versus Cbi (top
left) pCbl (top right), and CreCba (bottom left). Data points are colored by predicted function of
downstream regulatory target genes. The number of riboswitches analyzed from each group is
indicated in parentheses. Vertical and horizontal gray dashed lines demarcate a lack of response
to one corrinoid. Diagonal line indicates equal repression between corrinoids. The distance of a
point from the diagonal line indicates the bias in response towards one of the two corrinoids.

Figure S4 – Chimeric riboswitches show that multiple components contribute to corrinoid
specificity. Repression of GFP expression with 100 nM corrinoid is shown for (A) *P.*
megaterium metE and *V. parvula mutA* riboswitches and chimeric fusions of their aptamer and
expression platform domains. Aptamer subdomains swaps within the (B) *P. megaterium metE*
riboswitch scaffold and (C) *V. parvula mutA* riboswitch scaffold. Columns and error bars
represent mean and standard deviation of 3 replicates. Cartoon riboswitches depict *P.*
megaterium riboswitch sequence in cyan and *V. parvula* riboswitch sequence in orange.
Sequences in black are identical between riboswitches.

Figure S5 – Corrinoid lower ligand structure impacts base-on tendency. Absorbance spectra
of corrinoids were measured in a neutral buffered solution (pH 7.3). Absorbance peaks at 458 nm
and 525 nm are associated with base-off and base-on conformations, respectively. Base-on
tendency was measured as the ratio between absorbance at 525 nm and 458 nm (thin vertical
lines). Each panel is labeled with corrinoid name, base-on tendency value ($Abs_{525nm} / Abs_{458nm}$),
and lower ligand chemical structure. Note that the tail structures of PheCba, CreCba, and Cbi
cannot coordinate cobalt, and thus these corrinoids cannot be assigned a 'base-on tendency' per
se. The Abs_{525nm}/Abs_{468nm} ratios of PheCba, CreCba, and Cbi simply reflect an absence of any

Co-N coordination at the α axial face. For reference, the dashed line in the upper left panel is the
absorbance spectrum of nearly complete base-off Cbl measured in acidic solution (pH 1.57).

Figure S6 – Cbl-binding sites in X-ray crystal structures of various Cbl-riboswitches. The
marine metagenome derived riboswitch *env8* (PDB ID 4FRN) (A, B), the *B. subtilis btuF* Cbl-
riboswitch (PDB ID 6VMY) (C, D), and the *S. thermophilum cblT* Cbl-riboswitch (PDB ID
4GXY) are depicted with base-on (A, C, E) and base-off (B, D, F) Cbl alignments. Cbl models
were aligned by the cobalt and coordinating nitrogen atoms in the corrin ring. Structures of the
corrin ring, cobalt, and tail of Cbl are colored in black, blue and green, respectively. Upper
ligand structures of Cbl were omitted for clarity. Riboswitch RNA structures are depicted as
space-filled models with the L5-L13 kissing loop in pink in panels A and B, and just the L5 in
pink in panels C-F. The rest of the RNA structures in each panel are in gray. Arrows in panels D
and F point to clashes between RNA and the Cbl tails.

Figure S7 – The aptamer and expression platform of the *P. megaterium* riboswitch
contribute to corrinoid selectivity. (A) Repression of GFP expression with 100 nM corrinoid is
shown for the *P. megaterium metE* riboswitch, *S. ovata nikA* riboswitch, and a chimeric fusion of
the *S. ovata nikA* aptamer and *P. megaterium metE* expression platform domains. Cbl dose-
responses with or without 100 nM Cbi in (B) *P. megaterium metE* and (C) chimeric
riboswitches. Data points and error bars represent mean and standard deviation of four
independent replicates.

Table S1 – Strain list and riboswitch information. (A) List of strains used in this study. (B)
Genome and sequence information of wild type riboswitches analyzed in this study. (C)
Sequences of mutant riboswitches analyzed in this study.

Table S2 – X-ray crystal structural models used for 3D Cbl structural alignments.

Data S1 – Riboswitch sequence alignment in Stockholm sequence format.

A semi-selective riboswitch *P. megaterium metE*

B promiscuous riboswitch *V. parvula mutA*

Cbl
pCbl
CreCba
Cbi

Bacilli (12)
Negativicutes (16)
Deltaproteobacteria (5)
Clostridia (4)
Bacteroidia (1)

C GFP fold repression with 100 nM pCbl

D GFP fold repression with 100 nM Cbi

E GFP fold repression with 100 nM CreCba

GFP fold repression (F_0/F)

[corrinoid] (nM)

GFP fold repression with 100 nM Cbl

A MetH-compatible corrinoid

MetH: homocysteine → methionine

terminated *metE* mRNA transcript

B MetH-incompatible corrinoid

MetH inactive

MetE: homocysteine → methionine

full-length mRNA transcript

C Strain background:
B. subtilis Δ*metE::loxP* Δ*queG::loxP* P$_{Veg}$-*btuFCDR*

P$_{Veg}$ — *metH*-expressing strain — *P.m. metH* — *gfp*

P$_{Veg}$ — *metE*-expressing strain — RS — *P.m. metE* — *gfp*

D benzimidazolyl cobamides | purinyl cobamides | phenolyl cobamides

Growth rate (h^{-1})

Corrinoids: Cbl, [5-MeBza]Cba, [5-OMeBza]Cba, [5-OHBza]Cba, [Bza]Cba, [5-FBza]Cba, [2-MeAde]Cba, pCbl, CreCba, [Phe]Cba, Cbi, No corrinoid

E *metE* expression level (GFP/OD$_{600}$)

Corrinoid: Cbl, [5-MeBza]Cba, [5-MeOBza]Cba, [5-OHBza]Cba, [Bza]Cba, [5-FBza]Cba, [2-MeAde]Cba, pCbl, CreCba, [Phe]Cba, Cbi, No corrinoid

A

B

C

GFP fold repression (F_0/F)

[corrinoid] (nM)

- Cbl
- pCbl
- CreCba
- Cbi

C. difficile cbiM

C. novyi cobT

D. hafniense DSY0087

S. thermophilum cbiT (45°C)

D. propionicus mutA-1

D. propionicus mutA-2

D. propionicus cbiZ

D. propionicus exbB

P. propionicus cobU

B. thetaiotaomicron nrdD

Riboswitch reporter construct

A

B

Semi-selective riboswitch
P. megaterium metE

C

Chimeric riboswitch
So nikA aptamer
Pm metE exp. plat.

— Cbl + 0 nM Cbi
- - - Cbl + 100 nM Cbi