Defensive shimmering responses in *Apis dorsata* are triggered by dark stimuli moving against a bright background

Sajesh Vijayan[1]*, Eric J Warrant[2] and Hema Somanathan[1]

1: IISER TVM Centre for Research and Education in Ecology and Evolution (ICREEE), School of Biology, Indian Institute of Science Education and Research Thiruvananthapuram, Kerala, India
2: Lund Vision Group, Department of Biology, University of Lund, Sölvegatan 35, Lund, S-22362 Sweden

ORCID-ID: SV, 0000-0002-5110-1536; EJW, 0000-0001-7480-7016; HS, 0000-0002-4803-3894

*sajeshksv17@iisertvm.ac.in

Keywords
collective behaviour, defensive behaviour, honeybees, predator-prey interaction, visual ecology

Abstract

Giant honeybees, including the open-nesting Asian giant honeybee *Apis dorsata,* display a spectacular collective defence behaviour – known as "shimmering" – against predators, which is characterised by travelling waves generated by individual bees flipping their abdomens in a coordinated and sequential manner across the bee curtain. We examined if shimmering is visually-mediated by presenting moving stimuli of varying sizes and contrasts to the background (dark or light) in bright and dim ambient light conditions. Shimmering was strongest under bright ambient light, and its strength declined under dim-light in this facultatively nocturnal bee. *A. dorsata* shimmered only when presented with the darkest stimulus against a light background, but not when this condition was reversed (light stimulus against dark background). This response did not attenuate with repeated exposure to the stimuli, suggesting that shimmering behaviour does not undergo habituation. We suggest that this is an effective anti-predatory strategy in open-nesting *A. dorsata* colonies which are exposed to high ambient light, as flying predators are more easily detected when they appear as dark moving objects against a bright sky. Moreover, the stimulus detection threshold (smallest visual angular size) is much smaller in this anti-predatory context (1.6º - 3.4º) than in the context of foraging (5.7º), indicating that ecological context affects visual detection threshold.

Introduction

Honeybee colonies consist of several thousand individuals and brood cells, as well as abundant stores of honey and pollen that must be guarded from predators (Fuchs and Tautz, 2011; Seeley, 1995). The giant honeybee *Apis dorsata* is a tropical open-nesting species with colonies occupying cliff faces, tall trees or man-made structures such as water tanks and window ledges (Misra et al., 2016; Neupane et al., 2013; Roy et al., 2011; Tan et al., 1997; Thomas et al., 2009). Honeybee species exhibit a wide array of anti-predatory behaviours (Fuchs and Tautz, 2011; Phiancharoen et al., 2011; Seeley et al., 1982), and shimmering behaviour reported in the giant honeybees is the first line of defence against aerial flying predators such as birds and hornets which are their major predators (*A. dorsata*: (Kastberger et al., 2011), *A. laboriosa*: (Batra, 1996)). This behaviour is visualised as a dark travelling wave on the comb surface, and can precede a more aggressive collective stinging response (Seeley et al., 1982; Kastberger and Sharma, 2000; Kastberger et al., 2011; Wongsiri et al., 1997).

Shimmering in *A. dorsata* arises from 'flickering' behaviour, a form of dorsoventral abdomen-flipping that occurs stochastically (Weihmann et al., 2012). Once initiated at trigger centres on the bee curtain (Kastberger et al., 2010a; Schmelzer and Kastberger, 2009), the shimmering wave follows the path of the flying intruder (Kastberger et al., 2010a). More trigger centres can subsequently form if the intruder persists, as quiescent bees from other regions on the comb initiate shimmering, leading to progressively stronger and larger displays (Weihmann et al., 2012). The waves also accelerate at a few trigger centres that detect approaching waves from further away (Kastberger et al., 2012; Kastberger et al., 2014). Shimmering confuses potential predators, and serves as a multi-modal alerting signal for nestmates using mechanoreceptive cues (Kastberger et al., 2011), low-frequency comb vibrations (Kastberger et al., 2013), or by the release of Nasonov pheromone during wave generation (Kastberger et al., 2010b).

In this study we used artificial moving stimuli that simulate aerial predators to examine the precise relationships between the visual features of the stimuli and the strength of shimmering, by varying the relative contrast and visual angles subtended by the stimuli on the bee's eye at different ambient light levels during daylight and twilight in this facultatively nocturnal honeybee (Dyer, 1985; Somanathan et al., 2009). We predicted that the strength of shimmering will increase under the following conditions: (1) with increasing contrast between the object and the background, (2) with larger stimuli and (3) at higher ambient light levels. We also examine whether repeated exposure to potential threats (stimuli) leads to a progressive reduction in the strength of shimmering due to habituation. Since shimmering occurs in response to potential threats from predators/intruders with consequences for colony survival, we expected that habituation would not occur. We thus predicted that the shimmering response does not reduce or dampen with recurring exposure to the stimuli over trials.

Methods

A. dorsata colonies
The experiments were conducted in October - December 2020 and in June 2022 using two natural *A. dorsata* colonies located on the Biological Sciences building at the IISER Thiruvananthapuram campus, India (8.68ºN, 77.14ºE). Both colonies consisted of a single comb and nested under the roof rafters. The first colony, henceforth referred to as Hive A (1 m width, 1.2 m height), was situated 2 m above the floor level on the open terrace of the building (Fig 1A, Supp. Video) and the second colony, Hive B (1.8 m width, 0.7 m height), was 5-storeys above ground level (Fig 1B, Supp. Video). Though hives of *Apis dorsata* are often found hanging from rooftop rafters and window ledges of multi-storey buildings, accessing hives for performing controlled experiments is often risky or challenging. Both experimental hives were naturally located in their respective positions. Translocating hives to convenient locations for ease of experimentation has been done in other studies (Young et al., 2021). Since this could trigger absconding or change the baseline state of the colony, we refrained from doing so.

Experimental setup
The stimuli used in this study were made from circular cardboard cut-outs that varied in shade (light-grey, dark-grey or black) and size (diameter: 4, 8 or 16 cm; visual angle subtended on the eye of the bee: 1.6º, 3.3º and 6.5º, respectively). The relative contrasts of the stimuli were further varied by using two different backgrounds (grey or black). In the case of Hive A, a frame that consisted of two panels that could be swivelled along their vertical axes was placed 1.4 m from the colony (Fig. 1C), and the front and rear sides of the panels were covered with grey and black papers, respectively (papers

procured locally). The rear side of each stimulus was attached to a string and moved either vertically or
horizontally by the experimenter standing behind the frame and out of sight of the colony. To achieve
this, the string was strung around the setup vertically or horizontally. Moving the strings without a
stimulus was treated as the experimental control. The position of Hive B on an overhanging roof rafter
obstructed by walls and ledges prevented the use of the same arrangement. Instead, one of two
backgrounds consisting of cardboard sheets (1 m x 0.6 m) covered with the black or grey papers was
hung in front of the wall facing the colony during the experiments (Fig. 1D). The hive-to-background
distance of 1.4 m was the same as for Hive A. The experimenter stood behind the wall and moved the
stimuli horizontally with the help of a wire, and moving the wire alone acted as an experimental
control. Videos of the trials were recorded using a Sony Handycam HDR CX-405.
103
We carried out 799 trials on Hive A, randomising the order of presentation of the stimuli across the
following variables to prevent nestedness: stimulus shade, stimulus size, background colour, orientation
of motion (vertical/horizontal) and side of the setup on which the trial was done (left/right). We
conducted 240 trials on Hive B, randomised across stimulus shade, stimulus size, and background.
Trials for both hives were carried out at least 15 minutes apart to ensure that the colonies reached a
state of quiescence, and 9 – 45 trials were conducted each day between 06:00 h and 18:00 h.
110
The effect of ambient light intensity on shimmering was examined by moving the largest black
stimulus against the grey background under dim light conditions prevailing during astronomical,
nautical and civil twilight (n(Hive A) = 26, n(Hive B) = 20; illuminance between 0.002-45 lux), and the
responses were compared to shimmering elicited in bright daylight (illuminance > 1000 lux).
Illuminance levels were measured in lux using a Hagner Screenmaster (Hive A) and a Hagner E4-X
Digital Luxmeter (Hive B).
117
118 To quantify whether habituation occurred as a result of repeated exposure within experimental days, we
119 examined the strength of shimmering response over trials. We excluded the first data-point from each
120 day since more than 12 hours had elapsed after the last trial.
121
Relative contrasts of the stimuli against the backgrounds
The grey stimuli were made using laminated sheets of grey paper manufactured by Color-Aid Corp.
The dark and light-grey stimuli were made using Color-Aid Gray-5 and Color-Aid Gray-9 respectively.
Black stimuli were made from the same paper as the black background. All stimuli were pasted on to a
hard cardboard base to ensure that their shapes remained unchanged over the course of the experiment.
The reflectance spectra of the stimuli and backgrounds were recorded using an Ocean Insight Ocean-
HDX-UV-VIS spectrophotometer connected to an Ocean Optics PX-2 pulsed Xenon light-source (Fig.
S1A). The spectra were captured to a PC (Acer One 110-ICT) running the Ocean View software and
saved as a spreadsheet (see electronic supplementary materials). The contrasts of the stimuli against the
background were quantified using Weber contrast (O'Carroll and Wiederman, 2014)

132
$$C_{Weber} = \frac{Intensity_{object} - Intensity_{background}}{Intensity_{background}}.$$

133 The light-grey, dark-grey and black stimuli had contrasts of 16.4, 6.6 and 0, respectively, against the
134 black background, and 2.4, 0.5 and -0.8, respectively, against the grey background.
135
Quantification of shimmering

A trial lasted 15 s and consisted of the stimulus moving 15 times from one side to the other against the
front of the frame. The movement was timed using a metronome set to 60 beats per minute to maintain
a constant velocity of 1 m/s (40^o/s in the visual field of a bee). A shimmering response is characterised
by a distinct ascending phase, a climax and a descending phase (Fig. 1E-G, (Kastberger et al., 2014)) of
which only the area of the climax stage was considered for quantifying the strength of shimmering (Fig
1F). Videos were converted to AVI format using FFMPEG (at 25 fps) and analysed frame-by-frame
using ImageJ software (Schneider et al., 2012). The overall strength of shimmering in a trial S_{trial} was
quantified as

$$S_{trial} = \frac{\sum_{i=1}^{n} S_R}{n_{mov}}$$

146   where $S_R = Area_{climax} / Area_{hive}$ for the i_{th} climax event, and n_{mov} is the number of times the stimulus was
147   moved in a trial (fixed at 15 per trial). In other words, S_{trial} denotes the average proportion of the hive
148   that shimmered when the stimulus moved once.
149
Statistical analyses
We compared the strength of shimmering (S_{trial}) in relation to stimulus shade, size and background.
Since the values of S_{trial} belong to the interval [0,1], interactions were modelled using beta regression
with the hive identity as a random effect (Douma and Weedon, 2019). The side of the frame against
which the stimuli were presented (left side/right side) and the direction of motion (vertical/horizontal)
which were carried out only with Hive A were excluded as the shimmering response for these showed
no significant differences in pairwise comparisons with Bonferroni correction (Fig. S2).
157
158   Since the beta probability distribution is defined in the open interval (0,1), and 86% of the trials elicited
159   no shimmering (S_{trial}=0), we rescaled S_{trial} in the dataset using the following equation:
160
161     $x_i^o = \dfrac{x_i(n-1) + 0.5}{n}$
162
163
where x_i is the observed value, n is the number of trials (1039 in total) and x_i^o is the rescaled value for
that observation. All analyses were carried out in R version 3.6.3 (R Core Team, 2021), and the beta
regression analyses were done using the *glmmTMB* package (Brooks et al., 2017). We compared five
beta-regression models against two null models (one with pooled data and one with hive identity as a
random effect): a model with data pooled from both hives and single precision (ϕ) across all variables
(Model 1), a model with hive identity as a fixed effect (Model 2), a hierarchical (mixed effects) model
with hive identity as a random effect and single ϕ (Model 3), a hierarchical model with hive identity as
random effect and variable ϕ across stimulus size (Model 4), and a hierarchical model with hive
identity as random effect and variable ϕ across stimulus shade (Model 5). The best performing model
was chosen on the basis of the Aikake Information Criterion (AIC) (see Table S1). The effect of
ambient light intensity was estimated using hierarchical beta-regression modelling with light condition
(daylight/twilight) being the fixed effect and hive identity as random effect ($n_{daylight}$(Hive A) = 26),
$n_{daylight}$(Hive B) = 10, $n_{twilight}$(Hive A) = 25, $n_{twilight}$(Hive B) = 20), and the best performing model was
chosen on the basis of AIC (see Table S2). The dataset used for the daylight condition is a subset of the
data used in the previous analysis with the largest black stimulus against the grey background. Effect of

habituation was analysed using linear models for both hives by estimating the impact of duration
between consecutive trials on S_{trial}.
181
Results
A. dorsata shimmers strongly to a dark object moving against a light background
Shimmering responses were only observed in trials where the black stimulus was moved against the
grey background (Fig. 2). The hierarchical model with hive identity as random effect and variable
precision (ϕ) across stimulus size (Model 4) was found to be the most parsimonious compared to the
other models, including Model 2 where hive identity was considered as a fixed effect (Table S1, Fig.
S3). Hence, we did not carry out post-hoc statistical comparisons between hives. Shimmering in Hive A
was strongest with the largest black stimulus (subtending a visual angle of 6.5°, $S_{\text{trial}} = 0.25 \pm 0.12$) and
reduced in strength with the intermediate-sized black stimulus (3.3°, $S_{\text{trial}} = 0.09 \pm 0.10$), whereas both
stimuli elicited an equally strong response in the case of Hive B (S_{trial} (6.5°) $= 0.20 \pm 0.05$, S_{trial} (3.3°) $=$
0.15 ± 0.04). The smallest stimulus (1.6°) did not elicit shimmering in either hive (Fig. 2). Thus, the
detection threshold for a moving dark object is likely between 1.6° and 3.3°. The distinct behavioural
responses and the contrasts of the stimuli (Fig. S1) suggest that bees can distinguish and respond to the
stimuli on the basis of contrast and size, leading to strong shimmering responses as the visual angle
subtended by the intruding object/predator increases in the visual field of the bees.
197
Shimmering response is less pronounced in dim-light
31 trials out of the 45 carried out during dawn and dusk with the 16 cm black stimulus against the grey
background did not elicit shimmering ($n_{\text{Hive A}} = 13$ and $n_{\text{Hive B}} = 18$; illuminance between 0.1 - 3.9 lux),
and the rest showed varying strengths of shimmering (illuminance between 3.9 - 45 lux). The model in
which the data were pooled for both hives was the most parsimonious (Table S2). The mean strength of
shimmering under bright daylight conditions was an order of magnitude higher than the response
during twilight hours (S_{trial} (daylight) $= 0.243 \pm 0.101$, S_{trial} (twilight) $= 0.022 \pm 0.053$; Fig. 3).
205
A. dorsata does not habituate with repeated exposure to the stimuli
Since *A. dorsata* shimmered only to the 8 cm and 16 cm black stimuli moving against the grey
background, we excluded trials with other stimulus-background treatments from the analyses
examining if habituation occurred. Since the order of presentation of stimuli and backgrounds were
randomised (see Methods), the black-on-grey trials were interspersed with other treatments and the
trials were often not successive. 51 trials from Hive A and 14 trials from Hive B satisfied the condition
of black stimuli (8/16 cm diameter) moving against a grey background and repeating within a day
(06:00 AM – 06:00 PM). We did not observe any significant change in the strength of the shimmering
response with time elapsed between trials (range: 15 – 622 minutes, Fig. 4A), and there was no effect
of time of the day on the strength of the shimmering response (Fig. 4B).
216
Discussion
Defence behaviour in honeybees involves several sensory modalities (Fuchs and Tautz, 2011), and
shimmering is predominantly mediated by visual cues (Kastberger et al., 2010a; Kastberger et al.,
2011). When we simulated predator movement using artificial stimuli, the strongest shimmering
responses were obtained only in bright ambient light when the stimulus was much darker than the
background. This is unlike the role of contrast in the context of foraging (in *A. mellifera*), where the
stimulus can be either brighter or darker than the background and yet elicit similar responses (de Ibarra
et al., 2000). We also found that the strength of shimmering decreased as stimulus size reduced and

resulted in no response with the smallest stimulus. Repeated exposure to the experimental stimuli did
not lead to a change in the strength of the shimmering response in *A. dorsata,* although habituation or
sensitisation occurs in some invertebrates in response to repeated exposure to a threat (Evans et al.,
2019).
229
Detection of dark objects moving against bright backgrounds (e.g. the sky) has been reported in several
insects, including bees, although mostly in the context of mate-detection or predation (Bergman et al.,
2015; Sauseng et al., 2003). Male carpenter bees (*Xylocopa tenuiscapa*) detect and chase inanimate
projectiles, mistaking them to be potential mates or rivals, when these objects subtend very small visual
angles of 0.4º - 1º (which darken the visual field of a single ommatidium by as little as 2%)
(Somanathan et al., 2017). *A. mellifera* drones fly towards moving objects subtending as little as 0.4º
(which darkens the visual field by just 8%) (Vallet and Coles, 1993). *A. mellifera* workers can also
detect a small black stimulus (0.6º x0.6º) moving at 65º/s against a white background (Rigosi et al.,
2017), which is significantly smaller than their detection threshold during foraging (3.7º-5º) (Giurfa and
Vorobyev, 1998). Anatomical estimates from the compound eyes of *A. dorsata* suggest coarser visual
resolution compared to other honeybees (Somanathan et al., 2009), which agrees with recent estimates
of detection thresholds in *A. dorsata* in the context of foraging (achromatic: 5.7º, chromatic: 12.4º,
Balamurali et al. *in prep*).
243
Our experiments suggest that *A. dorsata* shimmers in response to dark objects moving against a bright
background that subtend a visual angle of at least 3.3º (and likely even smaller, between 1.6º and 3.3º),
which is smaller than their achromatic detection threshold while foraging. Using a conservative
detection threshold of 3.3º for shimmering behaviour, and assuming that a colony shimmers to a hornet
(a common predator) which is ~2.3-2.7 cm in length (Girish and Srinivasan, 2010), the detection-
distance threshold corresponds to ~40-47 cm, which agrees with results from real-world interactions
involving free-flying hornets near *A. dorsata* colonies (Kastberger et al., 2008). Besides predators, *A.*
dorsata also shimmer in response to intruders such as non-nestmate bees who display erratic flight
patterns and splay their legs outwards while doing so (Weihmann et al., 2014). Based on a length of ~2
cm for an *A. dorsata* worker (pers. obs.) and a detection threshld of 3.3º, we speculate that the distance
at which shimmering is triggered is around 35 cm. This suggests that shimmering could be a general
response to aerial threats.
256
Open-nesting honeybees such as *A. dorsata* are likely to perceive flying predators as dark objects
moving against the bright sky. Hornets are major predators of *A. dorsata* hives (Kastberger et al., 2013;
Seeley et al., 1982), and *Vespa tropica* (the greater banded hornet) is a common predator in our study
location (Balamurali et al., 2021). Moreover, hornets can act as strong evolutionary drivers in the
evolution of specialised anti-predatory responses in honeybees (Cappa et al., 2021; Ken et al., 2005;
Mattila et al., 2020; Mattila et al., 2021; Papachristoforou et al., 2007; Tan et al., 2013). Shimmering
resembles the anti-hornet 'I-see-you' display found in the sympatric cavity-nesting *A. cerana* in which
guard bees at the entrance shake their abdomens in a to-and-fro motion in the presence of hornets,
which supposedly warns the predator that the bees have detected its presence (Tan et al., 2013).
Simulating this threat with black discs resulted in strong abdomen-shaking responses in *A. cerana* to
stimuli of size 8º moving at 140º/s (Koeniger and Fuchs, 1973, as cited in Fuchs and Tautz, 2011)),
which agrees with the achromatic visual detection threshold of 7.7º in *A. cerana* (Meena et al., 2021).
In the context of an anti-predatory response such as shimmering, habituation may prove costly, and we
found that the shimmering response does not reduce in strength after repeated exposure to the stimuli,
which is advantageous to ward off repeated approaches/attacks by predators.

272
Although in our study we found that the shimmering response increased with larger stimulus size, up to
a maximum target size of 6.5° used in our experiments, we cannot rule out the possibility that the
response increases further for even larger targets, or simply saturates. Beyond a certain size threshold
(which is between 6.5° (current study) and 9.5° (Koeniger et al., 2017)), the colony responds to the
object with stinging attacks (Koeniger et al., 2017), which resembles the defensive response of *A.*
dorsata against much larger predators such as birds (Kastberger and Sharma, 2000). The findings from
our study using moving objects are consistent with the notion that the shimmering behaviour of the
giant honeybee is a response to an aerial predator moving in front of the hive against a bright sky
background.
282

Acknowledgements
284 SV was funded by the Council for Scientific and Industrial Research, India, SV and HS were funded by
285 IISER Thiruvananthapuram and EJW was funded by the Swedish Research Council (2016-04014). We
286 thank GS Balamurali for his valuable inputs, and PC Anand for helping with the pilot experiments for
287 this study. We also thank three anonymous reviewers for their valuable comments on the manuscript.

288
289

References
291

Balamurali, G. S., Reshnuraj, R. S., Johnson, J., Kodandaramaiah, U. and Somanathan, H.
(2021). Visual associative learning and olfactory preferences of the greater banded hornet,
Vespa tropica. *Insect. Soc.* **68**, 217–226.

Balamurali, G. S., Warrant, E. J. and Somanathan, H. Visual detection threshold and contrast
sensitivity in the Asian giant honey bee *Apis dorsata (in prep)*.

Batra, S. W. T. (1996). Biology of *Apis laboriosa* Smith, a Pollinator of Apples at High Altitude in the
Great Himalaya Range of Garhwal, India, (Hymenoptera: Apidae). *Journal of the Kansas
Entomological Society* **69**, 177–181.

Bergman, M., Lessios, N., Seymoure, B. M. and Rutowski, R. L. (2015). Mate detection in a
territorial butterfly—the effect of background and luminance contrast. *Behav Ecol* **26**, 851–860.

**Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen, A.,
Skaug, H., J., Mächler, M. and Bolker, B., M.** (2017). glmmTMB Balances Speed and
Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R
Journal* **9**, 378.

Cappa, F., Cini, A., Bortolotti, L., Poidatz, J. and Cervo, R. (2021). Hornets and Honey Bees: A
Coevolutionary Arms Race between Ancient Adaptations and New Invasive Threats. *Insects* **12**,
1037.

de Ibarra, N. H., Vorobyev, M., Brandt, R. and Giurfa, M. (2000). Detection of bright and dim
colours by honeybees. *J Exp Biol* **203**, 3289–3298.

Douma, J. C. and Weedon, J. T. (2019). Analysing continuous proportions in ecology and evolution: A practical introduction to beta and Dirichlet regression. *Methods Ecol Evol* **10**, 1412–1430.

Dyer, F. C. (1985). Nocturnal orientation by the Asian honey bee, *Apis dorsata*. *Animal Behaviour* **33**, 769–774.

Evans, D. A., Stempel, A. V., Vale, R. and Branco, T. (2019). Cognitive Control of Escape Behaviour. *Trends in Cognitive Sciences* **23**, 334–348.

Fuchs, S. and Tautz, J. (2011). Colony Defence and Natural Enemies. In *Honeybees of Asia* (ed. Hepburn, H. R.) and Radloff, S. E.), pp. 369–395. Berlin, Heidelberg: Springer Berlin Heidelberg.

Girish, P. and Srinivasan, G. (2010). Taxonomic studies of hornet wasps (Hymenoptera: Vespidae) *Vespa* Linnaeus of India. *Rec Zool Surv India* **110**, 57–80.

Giurfa, M. and Vorobyev, M. (1998). The angular range of achromatic target detection by honey bees. *J Comp Physiol A* **183**, 101–110.

Kastberger, G. and Sharma, D. K. (2000). The predator-prey interaction between blue-bearded bee eaters (*Nyctyornis athertoni* Jardine and Selby 1830) and giant honeybees (*Apis dorsata* Fabricius 1798). *Apidologie* **31**, 727–736.

Kastberger, G., Schmelzer, E. and Kranner, I. (2008). Social Waves in Giant Honeybees Repel Hornets. *PLoS ONE* **3**, e3141.

Kastberger, G., Weihmann, F. and Hoetzl, T. (2010a). Complex social waves of giant honeybees provoked by a dummy wasp support the special-agent hypothesis. *Commun. Integr. Biol.* **3**, 179–180.

Kastberger, G., Raspotnig, G., Biswas, S. and Winder, O. (2010b). Evidence of Nasonov Scenting in Colony Defence of the Giant Honeybee *Apis dorsata*. *Ethology* **104**, 27–37.

Kastberger, G., Weihmann, F. and Hoetzl, T. (2011). Self-Assembly Processes in Honeybees: The Phenomenon of Shimmering. In *Honeybees of Asia* (ed. Hepburn, H. R.) and Radloff, S. E.), pp. 397–443. Berlin, Heidelberg: Springer Berlin Heidelberg.

Kastberger, G., Weihmann, F., Hoetzl, T., Weiss, S. E., Maurer, M. and Kranner, I. (2012). How to Join a Wave: Decision-Making Processes in Shimmering Behavior of Giant Honeybees (*Apis dorsata*). *PLoS ONE* **7**, e36736.

Kastberger, G., Weihmann, F. and Hoetzl, T. (2013). Social waves in giant honeybees (*Apis dorsata*) elicit nest vibrations. *Naturwissenschaften* **100**, 595–609.

Kastberger, G., Hoetzl, T., Maurer, M., Kranner, I., Weiss, S. and Weihmann, F. (2014). Speeding Up Social Waves. Propagation Mechanisms of Shimmering in Giant Honeybees. *PLoS ONE* **9**, e86315.

Ken, T., Hepburn, H. R., Radloff, S. E., Yusheng, Y., Yiqiu, L., Danyin, Z. and Neumann, P. (2005). Heat-balling wasps by honeybees. *Naturwissenschaften* **92**, 492–495.

Koeniger, N. and Fuchs, S. (1973). Sound production as colony defence in *Apis cerana*. In *Proceedings of 7th International Congress, IUSSI*, p. London.

Koeniger, N., Kurze, C., Phiancharoen, M. and Koeniger, G. (2017). "Up" or "down" that makes the difference. How giant honeybees (*Apis dorsata*) see the world. *PLOS ONE* **12**, e0185325.

Mattila, H. R., Otis, G. W., Nguyen, L. T. P., Pham, H. D., Knight, O. M. and Phan, N. T. (2020). Honey bees (*Apis cerana*) use animal feces as a tool to defend colonies against group attack by giant hornets (*Vespa soror*). *PLoS ONE* **15**, e0242668.

Mattila, H. R., Kernen, H. G., Otis, G. W., Nguyen, L. T. P., Pham, H. D., Knight, O. M. and Phan, N. T. (2021). Giant hornet (*Vespa soror*) attacks trigger frenetic antipredator signalling in honeybee (*Apis cerana*) colonies. *R. Soc. Open Sci.* **8**, 211215.

Meena, A., Kumar, A. M. V., Balamurali, G. S. and Somanathan, H. (2021). Visual detection thresholds in the Asian honeybee, *Apis cerana*. *J Comp Physiol A* **207**, 553–560.

Misra, T. K., Pahari, S., Murmu, S. and Raut, S. K. (2016). Nesting Behaviour of the Giant Honeybees *Apis dorsata* Occurring in Jhargram, West Bengal, India. *Proc Zool Soc*.

Neupane, K. R., Woyke, J., Poudel, S. M., Zhang, S., Yang, H. and Singh, L. (2013). Nesting site preference and behavior of giant honey bee *Apis dorsata*. *Apimondia 2013* **1225**, 41–42.

O'Carroll, D. C. and Wiederman, S. D. (2014). Contrast sensitivity and the detection of moving patterns and features. *Phil. Trans. R. Soc. B* **369**, 20130043.

Papachristoforou, A., Rortais, A., Zafeiridou, G., Theophilidis, G., Garnery, L., Thrasyvoulou, A. and Arnold, G. (2007). Smothered to death: Hornets asphyxiated by honeybees. *Curr Biol* **17**, R795–R796.

Phiancharoen, M., Duangphakdee, O. and Hepburn, H. R. (2011). Biology of Nesting. In *Honeybees of Asia* (ed. Hepburn, H. R.) and Radloff, S. E.), pp. 109–131. Berlin, Heidelberg: Springer Berlin Heidelberg.

R Core Team (2021). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.

Rigosi, E., Wiederman, S. D. and O'Carroll, D. C. (2017). Visual acuity of the honey bee retina and the limits for feature detection. *Sci Rep* **7**,.

Roy, P., Leo, R., Thomas, S. G., Varghese, A., Sharma, K., Prasad, S., Bradbear, N., Roberts, S., Potts, S. G. and Davidar, P. (2011). Nesting requirements of the rock bee *Apis dorsata* in the Nilgiri Biosphere Reserve, India. *Trop Ecol* **52**, 285–291.

Sauseng, M., Pabst, M.-A. and Kral, K. (2003). The dragonfly *Libellula quadrimaculata* (Odonata: Libellulidae) makes optimal use of the dorsal fovea of the compound eyes during perching. *Eur. J. Entomol.* **100**, 475–479.

Schmelzer, E. and Kastberger, G. (2009). 'Special agents' trigger social waves in giant honeybees (*Apis dorsata*). *Naturwissenschaften* **96**, 1431–1441.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671–675.

Seeley, T. D. (1995). *The wisdom of the hive: the social physiology of honey bee colonies*. Cambridge, Mass: Harvard University Press.

Seeley, T. D., Seeley, R. H. and Akratanakul, P. (1982). Colony Defense Strategies of the Honeybees in Thailand. *Ecol Monogr* **52**, 43–63.

Somanathan, H., Warrant, E. J., Borges, R. M., Wallen, R. and Kelber, A. (2009). Resolution and sensitivity of the eyes of the Asian honeybees *Apis florea*, *Apis cerana* and *Apis dorsata*. *J Exp Biol* **212**, 2448–2453.

Somanathan, H., Borges, R. M., Warrant, E. J. and Kelber, A. (2017). Visual Adaptations for Mate Detection in the Male Carpenter Bee *Xylocopa tenuiscapa*. *PLOS ONE* **12**, e0168452.

Tan, N. Q., Chinh, P. H., Thai, P. H. and Mulder, V. (1997). Rafter beekeeping with *Apis dorsata:* some factors affecting the occupation of rafters by bees. *J Apicult Res* **36**, 49–54.

Tan, K., Wang, Z., Chen, W., Hu, Z. and Oldroyd, B. P. (2013). The 'I see you' prey–predator signal of *Apis cerana* is innate. *Naturwissenschaften* **100**, 245–248.

Thomas, S., Varghese, A., Roy, P., Bradbear, N., Potts, S. and Davidar, P. (2009). Characteristics of trees used as nest sites by *Apis dorsata* (Hymenoptera, Apidae) in the Nilgiri Biosphere Reserve, India. *J Trop Ecol* **25**, 559–562.

Vallet, A. M. and Coles, J. A. (1993). The perception of small objects by the drone honeybee. *J Comp Physiol A* **172**, 183–188.

Weihmann, F., Hoetzl, T. and Kastberger, G. (2012). Training for Defense? From Stochastic Traits to Synchrony in Giant Honey Bees (*Apis dorsata*). *Insects* **3**, 833–856.

Weihmann, F., Waddoup, D., Hötzl, T. and Kastberger, G. (2014). Intraspecific Aggression in Giant Honey Bees (Apis dorsata). *Insects* **5**, 689–704.

Wongsiri, S., Lekprayoon, C., Thapa, R., Thirakupt, K., Rinderer, T. E., Sylvester, H. A., Oldroyd, B. P. and Booncham, U. (1997). Comparative biology of *Apis andreniformis* and *Apis florea* in Thailand. *Bee World* **78**, 23–35.

Young, A. M., Kodabalagi, S., Brockmann, A. and Dyer, F. C. (2021). A hard day's night: Patterns in the diurnal and nocturnal foraging behavior of Apis dorsata across lunar cycles and seasons. *PLoS ONE* **16**, e0258604.

Figure 1. Experimental setup and quantification of shimmering response. A-B) Experimental colonies Hive A (A) and Hive B (B) used in the experiment. **C)** Experimental setup for Hive A. Hive-facing side of the panels were covered with the grey background, while the experimenter-facing side had a black background which could be changed by swivelling the panels. **D)** Experimental setup for Hive B. The cardboard panel covered with grey/black sheets was fastened in place from the

297 experimenter-facing side of the wall. **E-G)** An occurrence of shimmering in Hive B. The shimmering
298 wave (*outlined in yellow dots*) has an *onset/ascending* phase (E), *peak/climax* phase (F), and an
299 *offset/descending* phase (G). The strength of shimmering S_R is the area covered by the climax-stage
300 wave as a proportion of the area of the hive (F).

Figure 2. Shimmering responses in *A. dorsata* to moving stimuli against dark or light backgrounds. The relative strength of shimmering (S_{trial}) when presented with moving circular stimuli of varying diameters/visual subtenses (x-axes) and shades (light-grey, dark-grey and black (*insets*)) against either a black (*left*) or grey (*right*) background. The *hatched boxes* correspond to Hive A, the *cross-hatched boxes* correspond to Hive B, the *hinges* (horizontal bounds of the box) correspond to the interquartile range (IQR), the *bold horizontal lines* correspond to the medians, the *whiskers* denote the upper/lower hinge ± 1.5xIQR, and the *points* outside the *whiskers* represent outliers.

Figure 3. Effect of ambient light on strength of shimmering response, S_{trial}. The colonies showed only a weak response (Hive A) and no response (Hive B) in twilight (04:45-06:45 h and 18:00-20:00 h; 0.002-45 lux) to the largest black stimulus (diameter 16 cm, subtense 6.5°) when it moved against the grey background. During daylight (07:00-17:00 h; illuminance >1000 lux), the response to the same stimulus-background combination was much more pronounced. The *hatched boxes* correspond to Hive A, the *cross-hatched boxes* correspond to Hive B, the *hinges* (horizontal bounds of the box) correspond to the interquartile range (IQR), the *bold horizontal line* corresponds to the median, the *whiskers* denote the upper/lower hinge ± 1.5xIQR, and the *points* outside the *whiskers* represent outliers.

Figure 4. Effect of repeated exposure of stimuli on the shimmering response, S_{trial}. A) Effect of time elapsed on S_{trial}. Neither colony habituated to the black stimuli moving against the grey background. The *filled circles* (●) correspond to the 8 cm diameter black stimulus (visual subtense 3.3°), the *filled triangles* (▲) correspond to the 16 cm diameter black stimulus (visual subtense 6.5°), and the *solid line* is the linear regression of the strength of shimmering, S_{trial}, against the time elapsed between consecutive trials. The *inset text* indicated the Pearson correlation coefficient R, the p-value and the linear relation that best explains the data. **B)** Effect of time of the day on S_{trial}. Shimmering response did not exhibit any relation with the time of day when trials were conducted. The *filled circles* (●) correspond to the 8 cm diameter black stimulus (visual subtense 3.3°), the *filled triangles* (▲) correspond to the 16 cm diameter black stimulus (visual subtense 6.5°), and the *solid line* is the linear regression of the strength of shimmering, S_{trial}, against the time of the day. The *inset text* indicated the Pearson correlation coefficient R, the p-value and the linear relation that best explains the data.

16