Alternate response of *Bacillus anthracis atxA* and *acpA* to serum, HCO$_3^-$ and CO$_2$

[Ψ] Itai Glinert, [Ψ]Elad Bar-David, Amir Ben-Shmuel, Assa Sittner, Reut Puni, Shira Laredo, David Kobiler, Shay Weiss and Haim Levy[*]

Department of Infectious Diseases, Israel Institute for Biological Research, P.O. Box 19, Ness Ziona 74100, Israel.

[Ψ] equal contribution

*corresponding author

haiml@iibr.gov.il

Abstract

Bacillus anthracis overcomes host immune responses by producing capsule and secreting toxins. Production of these virulence factors in response to entering the host environment was shown to be regulated by *atxA*, the major virulence regulator, known to be activated by HCO_3^- and CO_2. While toxin production is regulated directly by *atxA*, capsule production is independently mediated by two regulators; *acpA* and *acpB*. In addition, it was demonstrated that *acpA* has at least two promotors, one of them shared with *atxA*. We used a genetic approach to study capsule and toxin production under different conditions. Unlike previous works utilizing NBY-HCO_3^- medium under CO_2 enriched conditions, we used a sDMEM-based medium. Thus, toxin and capsule production can be induced in ambient or CO_2 enriched atmosphere. Using this system, we could differentiate between induction by 10% NRS, 10% CO_2 or 0.75% HCO_3^-. In response to high CO_2, capsule production is induced by *acpA* in an *atxA*-independent manner, with little to no toxin (protective antigen PA) production. *atxA* is activated in response to serum independently of CO_2, inducing toxin and capsule production in an *acpA* or *acpB* dependent manner. HCO_3^- was also found to activate *atxA*, but in non-physiological concentrations. Our findings may help explain the first stages of inhalational infection, in which spores germinating in dendritic cells require protection (by encapsulation) without affecting cell migration to the draining lymph-node by toxin secretion.

Introduction

Bacillus anthracis, the causative agent of anthrax, is a gram positive, spore forming bacterium that infects humans via three major routes; inhalational (lung), cutaneous (skin) and gastrointestinal (gut) [1, 2]. The infectious form of *B. anthracis* is the spore, durable and environmentally stable for decades. In response to entering the host, spores germinate and overcome the immune system utilizing two major virulence factors; the poly-δ-D-glutamic antiphagocytic capsule and the tripartite toxin system [3]. Capsule producing genes are encoded as a polycistronic gene cluster on the pXO2 virulence plasmid, comprised of the *capB,C,A,D,E* genes. CapB and CapC are the D-glutamic acid polymerization enzymes, while CapA and CapE form the transport channel that exports the polymer from the cytoplasm through the cell wall and membrane to the cell surface [4-6]. Once there, CapD hydrolyzes the polymer to shorter chains and covalently links them to the cell wall [7]. CapD also regulates the length of the bound polymer chains through its hydrolysis activity [4, 8, 9]. The tripartite toxins are encoded by the *pag* (protective antigen-PA), *lef* (lethal factor-LF) and *cya* (edema factor-EF) genes, located on the pXO1 virulence plasmid [1]. LF is a metalloprotease that specifically cleaves members of the MAP kinase regulatory pathway of mammalian cells [10]. EF is a potent calmudolin dependent adenylate cyclase that interferes with cell regulation by elevating the internal cAMP levels [11]. Both LF and EF are driven into mammalian cells by PA, which binds specific receptors ANTXR1 (TEM8) and ANTXR2 (CMG2) [12]. Following binding to the receptor, PA is processed by cell-associated furin, activating oligomerization, forming a heptamer that binds LF and EF [12, 13]. The complex is then phagocyted and upon lysosomal fusion, the pH drop causes a conformational shift resulting in LF and EF injection into the cytoplasm. These toxin moieties then disrupt cell activity by disrupting intra-cellular signaling. Intoxication of immune cells results in their inactivation, disrupting host immune responses [12].

The AtxA was shown to be the major virulence regular of *B. anthracis* [2, 14]. This pXO1-located gene (encoding for the AtxA protein) activates a cascade of regulatory processes, resulting in up and down regulation of both chromosomal and plasmid encoded genes [15]. Capsule production is regulated by two pXO2 encoded genes – *acpA* [15] and *acpB* [16, 17]. Though it was demonstrated that AtxA regulates these two genes [18], it is not essential for capsule production since ΔpXO1 strains remain capable of capsule production [19, 20]. Since *B. anthracis* does not produce toxins or capsule under normal laboratory growth conditions, specific host-simulating growth conditions were developed. It was reported that growth of *B. anthracis* in NBY broth supplemented with glucose and bicarbonate in 5-15% CO_2 atmosphere induces toxin secretion and capsule production [16, 20-22]. These growth conditions were used to study different aspects of virulence regulation in *B. anthracis*.

Resulting findings indicated that AtxA in induced in response to HCO_3^- and CO_2, activating both toxin

and capsule production. This concept implies that toxins and capsule are produced simultaneously in the

host.

Previously we reported that growth of *B. anthracis* in supplemented DMEM (a high glucose cell culture

medium supplemented with pyruvate, glutamine, nonessential amino acids, (henceforth sDMEM) with the

addition of 10% serum, and incubated in a 10% CO_2 atmosphere, induces virulence factor production [17,

19]. These conditions were used to examine the effects of serum (here - normal rabbit serum, NRS),

HCO_3^- and/or CO_2 atmosphere, on the regulation of toxin and capsule production. This was coupled with a

systematic genetic approach, to dissect the regulation of the bacterial virulence response. Unlike previous

reports, we show that *AtxA* is induced by NRS or HCO_3^-, but not CO_2. The capsule regulator AcpA can be

induced by CO_2 in an *atxA*-independent manner, or by NRS in an *atxA*-dependent manner. AcpB was

activated only in an *atxA*-dependent manner, with *atxA* deletion resulting in complete abrogation of *acpB*

capsule regulation. Our results indicate that bacteria is capable of independent capsule production,

uncoupled from toxin production, but that toxins are always be co-induced with the capsule.

Materials and Methods

Bacterial strains, media and growth conditions.

Bacterial strains used in this study are listed in **Table 1**. For the induction of toxins and capsule production, we employed a modified DMEM (supplemented with 4mM L-glutamine, 1 mM Sodium pyruvate, 1% non-essential amino acid) that was supplemented with 10% normal rabbit serum or 0.75% NaHCO3 (Biological Industries – Israel) – hence sDMEM. Spores of the different mutants were seeded at a concentration of 5×10^5 CFU/ml and grown in 96 well tissue culture plates (100μl per well) for 5 or 24h at 37oC in ambient or 10% CO2 atmosphere.

Strain	Description/characteristics	Source
B. anthracis		
Vollum	ATCC 14578	IIBR collection
VollumΔpXO1	Complete deletion of the plasmid pXO1	IIBR collection
VollumΔ_atxA_	Complete deletion of the _atxA_ gene	This study
VollumΔ_acpA_	Complete deletion of the _acpA_ gene	[17]
VollumΔ_acpB_	Complete deletion of the _acpB_ gene	[17]
VollumΔ_acpA_Δ_acpB_	Complete deletion of the _acpA_ and _acpB_ genes	[17]
VollumΔ_pag_Δ_cya_Δ_lef_ (ΔTox)	Complete deletion of the _pag_, _lef_ and _cya_ genes	[23]
VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpA_	Complete deletion of the _acpA_ gene in the VollumΔ_pag_Δ_cya_Δ_lef_ mutant	[17]
VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpB_	Complete deletion of the _acpB_ gene in the VollumΔ_pag_Δ_cya_Δ_lef_ mutant	[17]
VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpA_Δ_acpB_	Complete deletion of the _acpA_ and _acpB_ genes in the VollumΔ_pag_Δ_cya_Δ_lef_ mutant	[17]
VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpA_Δ_atxA_	Complete deletion of the _atxA_ gene in the VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpA_ mutant	This study
VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpB_Δ_atxA_	Complete deletion of the _atxA_ gene in the VollumΔ_pag_Δ_cya_Δ_lef_Δ_acpB_ mutant	This study
VollumΔpXO2 Δ_bclA_:: _pag_prom::_capA-E_	Genomic insertion of the _pagA_ promotor in front of the CAP operon replacing the _bclA_ gene in the VollumΔpXO2	[17]
VollumΔpXO2 Δ_bclA_:: _pag_prom::_capA-E_Δ_atxA_	Deletion of the _atxA_ gene from the strain having Genomic insertion of the _pagA_ promotor in front of the CAP operon replacing the _bclA_ gene in the VollumΔpXO2	[17]

Table 1: Bacterial strains used in this study

Mutant strain construction.

Oligonucleotide primers used in this study were previously described [17, 19, 23]. The oligonucleotide primers were designed according to the genomic sequence of *B. anthracis* Ames strain. Genomic DNA (containing the chromosomal DNA and the native plasmids, pX01 and pX02) for cloning the target gene fragments was extracted from the Vollum strain as previously described [24]. Target genes were disrupted by homologous recombination, using a previously described method [25]. In general, gene deletion or insertion was accomplished by a marker-less allelic exchange technique that replaced the complete coding region with the SpeI restriction site or the desired sequence. At the end of the procedure the resulting mutants did not code for any foreign sequences and the only modification is the desired gene insertion or deletion. Deletion of the atxA gene was performed as previously described [25]. All the mutants were tested for their ability to produce capsule by incubation in sDMEM. The capsule was visualized by negative staining with India ink.

Toxin quantification

Protective antigen (PA) concentration was determine by capture ELISA using the combination of a polyclonal and monoclonal αPA antibodies previously described [23]

Results

The effect of supplements and growth condition on capsule production in sDMEM medium. The basic medium used to test capsule and toxin induction was high glucose DMEM supplanted with glutamine, pyruvate and non-essential amino acids, hence sDMEM. We first tested the effect of growth media volume on capsule production in a 96-well tissue culture microplate format. This was done to allow higher-throughput testing of growth conditions and genetic manipulations. We inoculated *B. anthracis* Vollum spores (5×10^5 CFU/ml) into 100µl, 200µl and 300µl sDMEM at $37^{\circ}C$ in ambient atmosphere for 24h. No capsule production could be detected in 100µl sDMEM (**Figure 1**). However, increasing the culture volume resulted in partial (200µl) or full (300µl) capsule production (**Figure 1**). These findings could be possibly explained by higher CO_2 concentrations reached in the well when bacteria are grown in larger volumes, due to a lower surface area to volume ratio, limiting gas exchange. This is coupled with a larger volume of multiplying and respiring bacteria, further increasing CO_2 concentrations. This hypothesis is supported by the finding that growth in 10% CO_2 atmosphere does induce capsule production in 100µl sDMEM (**Figure 1**). Having established non-capsule inducing growth conditions for our experiments, we next tested the effect of adding normal rabbit serum (NRS) to growth media. NRS was shown to induce capsule production, regardless of CO_2 concentrations in the incubator's atmosphere, with all the bacteria grown with NRS encapsulated in ambient or 10% CO_2 atmosphere (**Figure 1**).

The role of *atxA* on capsule production and toxin secretion under different growth conditions. To determine the role of *atxA* on capsule and toxin production, we compared capsule production of the wild type Vollum strain with the ΔpXO1 or Δ*atxA* mutants (**Table 1**) following growth in sDMEM under the different growth conditions (NRS/CO_2 presence). None of the strains produced capsule under ambient atmosphere in un-supplemented sDMEM (**Figure 2**). Addition of 10% NRS, under ambient atmosphere, induced capsule production by the wild type Vollum strain but not by the *atxA* null mutants (both ΔpXO1 or Δ*atxA*, **Figure 2**). Under 10% CO_2, all three strains produced capsule. We therefore conclude that the response to NRS is *atxA* dependent, as lacking either the atxA gene or the entire pXO1 plasmid abrogates

this response (**Figure 2**). These results also demonstrate the presence of another, atxA independent mechanism, responding to CO_2. However, the effect of NRS in the presence of CO_2 on capsule production under these conditions could not be resolved.

Figure 2. Capsule production of the Vollum wild type, ∆pXO1 or ∆atXA mutants under different growth conditions. Spores of the wild type and different mutants (top panel) were seeded into 100μl of sDMEM. The same strains were grown in sDMEM supplemented with 10% NRS (second row). The same conditions were applied in the presence of 10% CO_2 (lower two rows) for 24h. Capsule was imaged by India ink negative staining (capsule presence forms a typical bright outer layer).

As *atxA*, the major virulence regulator, regulates toxin expression (LF, EF and PA), we tested their secretion into growth medium after growth, both for the Vollum and the Vollum∆*atxA* mutant (the ∆pXO1 mutant lacks the entire set of toxin genes and is therefore irrelevant for such a test). Toxins secretion was determined by ELISA for the most abundant component, PA. For the Vollum∆*atxA* mutant, PA was undetectable, both under 10% CO_2, with 10% NRS and in the presence of both. These results confirm the role of atxA in toxin expression regulation (**Table 2**). For the wild type Vollum, no toxins were induced in sDMEM alone coupled with under ambient atmosphere. The addition of 10% NRS alone induced high PA

expression. sDMEM without NRS under 10% CO_2 again did not induce PA expression (**Table 2**). However, the addition of both NRS and CO2 resulted in reduced PA expression (about a 70% reduction) (**Table 2**). This finding was maintained throughout different, independent experiments. However, we cannot exclude the possibility that this difference resulted from growth rates variation under the tested growth conditions. No PA secretion could be detected in the *atxA* null mutant under any of the tested growth conditions. Our finding that 10% CO_2 atmosphere in itself does not activate *atxA*-dependent toxin expression is surprising, since it was repeatedly demonstrated such activation is achievable by HCO_3^- addition. We therefore tested both capsule production and toxin expression in sDMEM supplemented with 0.75% HCO_3^- under 10% CO_2 atmosphere. These atmospheric conditions were chosen since HCO_3^- in aqueous media is in equilibrium with CO_2 and H_2O, dramatically increasing the levels of the soluble CO_2 in the medium even under ambient atmosphere, and we sought to expose our experimental controls to conditions as similar as possible.

Table 2. Protective Antigen secretion of Vollum or VollumΔ*atxA* under different growth conditions.

	VollumΔ*atxA*	Vollum	Atmosphere	Supplements
PA µg/ml	<0.1	<0.1	Ambient	-
	<0.1	8.7	Ambient	10% NRS
	<0.1	<0.1	10% CO_2	-
	<0.1	2.6	10% CO_2	10% NRS

According to our previous findings, capsule production is CO_2 dependent. Therefore, we found that the above tested strains were encapsulated under the conditions applied (**Figure 3**). PA secretion was tested only for Vollum and VollumΔ*atxA*. Supplementing the sDMEM with 0.75% HCO_3^- induced PA secretion by the Vollum strain but not by the VollumΔ*atxA* mutant. This indicates that HCO_3^- induces toxin secretion in an *atxA* dependent manner (**Table 3**). To validate these findings, we used a previously reported VollumΔpXO2 chimera in which we substituted the genomic *bclA* gene (which is a redundant exosporium glycoprotein) with a CAP operon altered to be regulated by the PA$_{prom}$ (**Table 1**). Thus, this mutant serves here as an indicator for AtxA dependent activation of the PA promotor, which results in capsule production (assayable readout). This mutant was then used as the basis for another mutant strain, in which the *atxA* gene was deleted, on top of the existing genetic alterations (**Table 1**). Thus, the exact role *atxA* plays in the regulation on the PA-promotor under the tested conditions could be resolved.

Figure 3. Effect of 0.75% HCO_3^- on capsule production of Vollum, Vollum$\Delta atxA$ or VollumΔpXO1. Spores were seeded into 100μl of sDMEM as is or supplemented with 0.75% HCO_3^- or 10%NRS, and incubated at 37°C under 10% CO_2 for 24h. Capsule was imaged by India ink negative staining (capsule presence forms a typical bright outer layer).The different mutations are indicated in the top panel.

Table 3. PA secretion by Vollum and Vollum$\Delta atxA$ in sDMEM supplemented with HCO_3^-.

	Vollum$\Delta atxA$	Vollum	Atmosphere	Supplements
PA μg/ml	<0.1	<0.1	10% CO_2	-
	<0.1	3.12	10% CO_2	0.75% HCO_3-
	<0.1	5.3	10% CO_2	10% NRS

While 10% CO_2 atmosphere induced capsule production in the Vollum wild type strain, little or no capsule production could be detected in the VollumΔpXO2$\Delta bclA$::$pagA_{prom}$-capA-E chimeric strain (**Figure 4**). Supplementing sDMEM with 0.75% HCO_3^- resulted in capsule production in the Vollum and VollumΔpXO2$\Delta bclA$::$pagA_{prom}$-capA-E chimera. However, deletion of the *atxA* gene, in the background of the chimera mutant described, resulted in loss of capsule production (VollumΔpXO2$\Delta bclA$::$pagA_{prom}$-capA-E$\Delta atxA$, **Figure 4**). Similar results were obtained by supplementing sDMEM with 10% NRS. These results

complement the PA secretion experiment, indicating that HCO_3^- and NRS induce toxin secretion in an

atxA dependent manner.

Figure 4. Capsule production in response to HCO_3^- in a PA$_{prom}$- regulated genomic CAP operon. Spores were

seeded into 100μl of sDMEM as is or supplemented with 0.75% HCO_3^- or 10% NRS and incubated at 37°C under an

atmosphere of 10% CO_2 for 24h. Capsule was imaged by India ink negative staining (capsule presence forms a typical

bright outer layer).The different mutations are indicated (top panel).

The effect of short incubations on capsule and toxins production in response the different growth

conditions. Aerobic growth affects different parameters of the liquid medium such as pH and O_2/CO_2

concentrations, especially when the bacteria reaches high concentration (CFU/ml). As was shown

previously [26], capsule production and toxin secretion can be detected as early as 2-5h of growth in

sDMEM under 10% CO_2 atmosphere. To minimize changes in media conditions resulting from bacterial

growth, we examined capsule production and PA secretion after 5h growth in different growth conditions.

Vollum growth under ambient atmosphere did not result in any capsule accumulation or toxin secretion

following 24h incubation (Figure 2, Table 2). This was true also for a short (5h) incubation (Figure 5,

Table 4). Supplementing the media with 10% NRS induced capsule production and PA secretion following

24h incubation under ambient atmosphere (Figure 2, Table 2). A shorter (5h) incubation induced

measurable PA secretion (Table 4) but no capsule production (Figure 5). This PA secretion is AtxA

dependent, as deletion of the *atxA* gene resulted in no PA accumulation following 5h (Table 4) or 24h

incubation (**Table 2**). Incubating the bacteria in 10% CO_2 atmosphere for 5h resulted in capsule production (**Figure 5**), similarly to that observed following a 24h incubation (**Figures 1-3**). This capsule accumulation was AtxA independent and was not significantly affected by the addition of 10% NRS or HCO_3^- (**Figure 5**). PA secretion was not induced by 10% CO_2 in itself, but required addition of NRS or HCO_3^- (**Table 4**), however unlike the 24h incubation (**Table 3**), the amount of PA secreted in response to HCO_3^- was significantly less (~10%) than of that induced by NRS (**Table 4**). Under all conditions the PA secretion was AtxA dependent.

Figure 5. Effect of deleting *atxA* on capsule production after a short (5h) incubation under different growth conditions. Spores of the wild type and Δ*atxA* mutant (top panel) were seeded into 100μl of sDMEM as is

246   Table 4. PA secretion by Vollum and Vollum$\Delta atxA$ following 5h growth in different conditions

	Vollum$\Delta atxA$	Vollum	Atmosphere	Supplements
PA µg/ml	<0.1	<0.1	Ambient	-
	<0.1	1.89	Ambient	10% NRS
	<0.1	<0.1	10% CO_2	-
	-	4.25	10% CO_2	10% NRS
	-	0.29	10% CO_2	0.75% HCO_3-

Regulation of *acpA* and *acpB* in response to different growth conditions. Toxin production is induced in an *atxA* dependent manner in response to HCO_3^- or NRS, while capsule production is also induced by a CO_2 enriched (10%) atmosphere. Capsule production is known to be regulated by two regulatory proteins, AcpA and AcpB. We therefore tested the effect of deleting each of these genes on capsule production in response to different growth conditions. The complete coding region of *acpA* or *acpB* was deleted independently in the background of wild type Vollum or the toxin deficient mutant - VollumΔTox (Vollum$\Delta pag\Delta cya\Delta lef$ **Table 1**). As was previously shown for the wild type Vollum strain (**Figure 2**), none of these mutants produced capsule following growth in sDMEM under ambient atmosphere (**Figure 6**). The presence of either *acpA* or *acpB* is sufficient for capsule production in sDMEM supplemented with 10% NRS, regardless to the presence or absence of 10% CO_2 atmosphere (**Figure 6**). However, in the absence of NRS, only AcpA expressing mutants (lacking *acpB*) produce significant capsule when grown in 10% CO_2 atmosphere. Adding 0.75% HCO_3^- to sDMEM induced capsule production in the presence of either *acpA* or *acpB*. Mutants lacking *acpA* (expressing only *acpB*), did not produce significant capsule in 10% CO_2 atmosphere. To examine the role of *atxA* in these processes, we deleted the *atxA* gene in the background of our VollumΔTox$\Delta acpA$ or $\Delta acpB$ mutants. Deleting *atxA* in the VollumΔTox$\Delta acpA$, expressing *acpB*, abolished capsule production under all tested conditions (**Figure 7**). However, deleting *atxA* in the VollumΔTox$\Delta acpB$, expressing *acpA,* did not affect capsule accumulation, compared to the AtxA expressing mutant. This finding indicates that *acpA* operates in an AtxA independent manner (**Figure 7**). As we demonstrated (**Figure 2**), capsule production could be induced in ambient atmosphere by adding 10% NRS to sDMEM. This induction is AtxA dependent, since no capsule production was detected in the Vollum$\Delta atxA$ mutant under these conditions (**Figure 2**). Since AcpA dependent capsule production in 10% CO_2 atmosphere was AtxA independent, we tested the role of AtxA on AcpA dependent capsule production in response to 10% NRS in ambient atmosphere. As 10% NRS induced capsule production of Vollum$\Delta acpB$ under ambient atmosphere (**Figure 6, Figure 8**) we tested the effect of *atxA* deletion on

capsule production under these conditions. Unlike the CO_2 induction, under ambient atmosphere, AcpA

dependent capsule production in response to NRS is AtxA dependent (**Figure 8**).

Figure 6. The effect of absence of *acpA* or *acpB* on capsule production in response to 10% NRS under

ambient or 10% CO_2 atmosphere. Spores of the Δ*acpA* or *acpB* mutants (top panel) were seeded into 100μl of

sDMEM as is or supplemented with0.75% HCO_3^- or 10% NRS and incubated at 37°C in an ambient or 10% CO_2

atmosphere (as indicated on the right) for 24h. Capsule was imaged by India ink negative staining (capsule presence

forms a typical bright outer layer).

Figure 7. Effect of AtxA on capsule production in the presence of either AcpA or AcpB. Spores of the

different mutants (top panel) were seeded into 100μl of sDMEM as is or supplemented with 0.75% HCO$_3^-$ or 10% NRS

and incubated at 37°C in 10% CO$_2$ atmosphere (as indicated on the right) for 24h. Capsule was imaged by India ink

negative staining (capsule presence forms a typical bright outer layer).

Figure 8. Effect of AtxA on capsule production in response to 10% NRS in ambient atmosphere. Spores

of the different mutants (top panel) were seeded into 100μl of sDMEM as is or supplemented with 10% NRS and

incubated at 37°C in ambient atmosphere (as indicated on the right) for 24h. Capsule was imaged by India ink negative

staining (capsule presence forms a typical bright outer layer).

Gas content of the supplemented and basic DMEM media under the different growth conditions. Gas

content was determined by Abbott iSTAT blood gas analyzer. The different media were incubated at

37°C in ambient or 10% CO_2 atmosphere for 4 h and analyzed using the EC8+ cartridge. The results shown

in **Table 5** indicate that the main parameter that is affected by the 10% CO_2 is the soluble CO_2 (PCO_2)

rather than HCO_3^-. Unlike the HCO_3^- levels that are higher under all growth conditions from those

considered normal for human blood, the PCO_2 levels in the presence of 10%CO_2 are well within or very

close to normal values.

Table 5. Gas analysis of the sDMEM media under the different growth conditions

10% CO_2	-	+	+	-	+	Normal arterial	Normal venous
Supplement	-	-	10% FCS	10% FCS	0.75% HCO_3 1:1*		
pH	8.0	7.691	7.447	8.149	7.972	7.35-7.45	7.31-7.41
PCO_2 (mmHg)	18.4	34.9	58.1	14.2	34.3	35-45	41-51
PO_2 (mmHg)	230	148	126	173	183	80-100	35-40
HCO_3 (mmol/L)	48.7	42.2	40.1	49.5	79.2	22-26	22-26
sO_2 (%)	100	100	99	100	100	95-100	/

301 *at the end of the 4h incubation the media was diluted 1:1 in sDMEM prior to the analysis since the
302 undiluted media was off scale.

Discussion

Successful invasion requires the pathogen regulate its virulence factors in a way that will maximize their effect on host defense mechanisms. The trigger for such activation is usually host derived. This can be biological (such as proteins) of physical (pH or temperature for example). *B. anthracis* naturally infects humans following spore inhalation, contact with broken skin or ingestion of undercook contaminated meat. These routes present different environmental conditions [2]. It was previously demonstrated that toxin secretion and capsule production could be induced by growing the bacteria in culture media supplemented with HCO_3^- or serum (10-50%) in a CO_2 enriched (5-15%) atmosphere [5, 16, 18, 19, 27-29]. HCO_3^- /CO_2 condition were commonly used to study *atxA*, *acpA* and *acpB* regulation and their effect on toxin and capsule biosynthetic genes [16, 18, 30]. Since these conditions always included these two components, it was concluded that *atxA* was induced in response to CO_2, regulating the induction of *acpA* and *acpB*. Although in some reports, capsule production was shown to be AtxA dependent [16, 31], the fact that ΔpXO1 variants are encapsulated contradicts this finding, alluding to additional, AtxA independent regulation of the process [19, 27]. The use of sDMEM as growth media enabled the examination of the effect of CO_2, HCO_3^- and serum on these processes.

The parameter of soluble CO_2 is influenced by multiple parameters, such as surface area to volume ratio and aerobic bacterial growth. Therefore, normal growth conditions were set to 100μl media/well (96 well tissue culture plate) for 24h at 37°C under ambient atmosphere (**Figure 1**). This baseline enabled testing the effect of different supplements and/or growth conditions on capsule (**Figure 1**) or toxin (**Figure 2, Table 2**) induction. Capsule production is induced by the addition of 10% NRS or growth under a 10% CO_2 atmosphere (**Figure 1**). The serum capsule induction (under ambient atmosphere) is *atxA* dependent (**Figure 2**), since there was no significant capsule production in mutants that do not express AtxA (VollumΔ*atxA* and VollumΔpXO1). Alternatively, capsule production in response to CO_2 enriched atmosphere is AtxA independent, as there is no significant difference in capsule production, under these conditions, between AtxA expressing and *atxA* null mutants (**Figures 2, 3**). Toxin secretion, as determined by measuring PA media concentrations, is serum dependent (**Tables 2, 3**), as PA could be detected only in NRS supplemented sDMEM regardless of CO_2 enrichment. Adding HCO_3^- induced toxin secretion in an *atxA* dependent manner, similarly to serum (**Figure 3, Table 3**), with PA expressed by the wild type Vollum and not by the *atxA* null mutant. The same NRS/HCO_3^- dependence and CO_2 independence of PA induction was demonstrated using a previously described mutant strain, in which pXO2 is missing and a copy of the CAP operon was inserted chromosomally, regulated by a PA promotor [17]. This mutant strain produces capsule when grown in sDMEM supplemented with NRS or HCO_3^- but not in non-supplemented sDMEM under 10% CO_2 atmosphere.

We also found differences in the speed *B. anthracis* reacts to the different stimuli. This was done by examining toxin secretion and capsule production after a short incubation (5h). HCO_3^- was not as robust as NRS in inducing toxin secretion. Examining PA concertations after 5h incubation in sDMEM supplemented with 0.75% HCO_3^- revealed about 1/10 of the concentration compared to that measured after 24h incubation (**Figure 5, Table 4**). However, NRS induction yielded similar concentrations at these two timepoints. Testing the effect of NRS on capsule production, reveals that 5h incubation under ambient atmosphere, inducing significant PA secretion, does not result in significant capsule production. Growth under a 10% CO_2 atmosphere induced capsule production at 5h even in the absence of supplemented NRS or HCO_3^- (**Table 4**). Hence, Serum seemed more effective then HCO_3^- in inducing toxin secretion, with two process being AtxA dependent. The AtxA independent induction of capsule production by 10% CO_2 appeared more effective then AtxA dependent serum induction (**Figure 5**).

Two major regulators; AcpA and AcpB control capsule biosynthesis by promoting transcription of *acpB,C,A,D,E* operon. *acpA* was shown to be regulated by AtxA (activated by NRS), while also depicting an additional *atxA* independent activation capacity by CO_2 and HCO_3^-). This independent activation could be direct or mediated by an as yet unknown, additional promotor, in turn responding to an CO_2 ingredient. We found that deleting *acpA* causes the bacteria to produce significantly less capsule in response to CO_2, while maintaining its ability to respond to NRS or HCO_3^- (**Figure 6**). Deletion of *acpB* did not have any effect on capsule production under all tested conditions (**Figure 6**), supporting our previous *in vivo* data, which showed no effect on virulence [17]. A double deletion of *atxA* and *acpA* or *acpB* revealed that AcpB activity is strictly AtxA dependent under all the conditions tested (**Figure7**). AcpA activity is not affected by the absence of *atxA* in the presence of CO_2 (**Figure7**) but is nulled in response to NRS under ambient atmosphere (**Figure 8**).

Our findings delineate the following regulation cascade; CO_2 induces capsule production by activating of *acpA* in an AtxA independent manner. Serum activates the AtxA dependent cascade, inducing toxin secretion and eventually capsule production, by activating *acpA* and *acpB* (**Figure 9**). The order of the processes can be deduced from the lack of capsule production following 5h growth in NRS supplemented sDMEM under ambient atmosphere (**Figure 5**). HCO_3^- induces toxin secretion through the AtxA cascade, but in a less efficient manner (compared to NRS, **Table 4**). Direct activation of capsule production by HCO_3^- in an AtxA independent manner could not be eliminated, since even under ambient atmosphere, it modifies the levels of soluble CO_2 (PCO_2) and possibly induces capsule production via AcpA (**Table 5**). In terms of the initial infection stages of pathology, this differential regulation of toxins and capsule has a logical role. Inhalational and cutaneous infections involve spore phagocytosis by local innate immune cells and their migration to a draining lymph node. While en route (and within the phagocytic cell), the spore

needs to germinate and produce the protective capsule. During this stage, toxin production could prove

counterproductive, as it disrupts normal cellular function, possibly arresting the cell and preventing it

from reaching the lymph node. Once there, toxin production is desirable, possibly enhancing bacterial

release from the cells in the lymph node, to the blood stream, promoting pathogenesis. The pathway

sensing serum and CO_2 remains to be elucidated and requires more research. Such a pathway may prove

common to other pathogens as well as possibly providing additional therapeutic targets for intervention.

Figure 9. Proposed regulatory scheme for CO_2, Serum and HCO_3^- regulation of capsule production and

toxin secretion.

References

1. Dixon TC, Meselson M, Guillemin J, Hanna PC. Anthrax. New England Journal of Medicine. 1999;341(11):815-26. doi: 10.1056/nejm199909093411107. PubMed PMID: 10477781.

2. Hanna P. Anthrax pathogenesis and host response. Curr Top Microbiol Immunol. 1998;225:13-35. Epub 1997/12/05. PubMed PMID: 9386326.

3. Swartz MN. Recognition and management of anthrax--an update. N Engl J Med. 2001;345(22):1621-6. Epub 2001/11/13. doi: 10.1056/NEJMra012892. PubMed PMID: 11704686.

4. Candela T, Fouet A. Bacillus anthracis CapD, belonging to the gamma-glutamyltranspeptidase family, is required for the covalent anchoring of capsule to peptidoglycan. Mol Microbiol. 2005;57(3):717-26. Epub 2005/07/28. doi: 10.1111/j.1365-2958.2005.04718.x. PubMed PMID: 16045616.

5. Candela T, Mock M, Fouet A. CapE, a 47-amino-acid peptide, is necessary for Bacillus anthracis polyglutamate capsule synthesis. J Bacteriol. 2005;187(22):7765-72. Epub 2005/11/04. doi: 10.1128/JB.187.22.7765-7772.2005. PubMed PMID: 16267300; PubMed Central PMCID: PMCPMC1280324.

6. Heine HS, Shadomy SV, Boyer AE, Chuvala L, Riggins R, Kesterson A, et al. Evaluation of Combination Drug Therapy for Treatment of Antibiotic-Resistant Inhalation Anthrax in a Murine Model. Antimicrob Agents Chemother. 2017;61(9). Epub 2017/07/12. doi: 10.1128/AAC.00788-17. PubMed PMID: 28696235; PubMed Central PMCID: PMCPMC5571330.

7. Candela T, Fouet A. Poly-gamma-glutamate in bacteria. Molecular Microbiology. 2006;60(5):1091-8. doi: doi:10.1111/j.1365-2958.2006.05179.x.

8. Wu R, Richter S, Zhang RG, Anderson VJ, Missiakas D, Joachimiak A. Crystal structure of Bacillus anthracis transpeptidase enzyme CapD. J Biol Chem. 2009;284(36):24406-14. Epub 2009/06/19. doi: 10.1074/jbc.M109.019034. PubMed PMID: 19535342; PubMed Central PMCID: PMCPMC2782033.

9. Richter S, Anderson VJ, Garufi G, Lu L, Budzik JM, Joachimiak A, et al. Capsule anchoring in Bacillus anthracis occurs by a transpeptidation reaction that is inhibited by capsidin. Mol Microbiol. 2009;71(2):404-20. Epub 2008/11/20. doi: 10.1111/j.1365-2958.2008.06533.x. PubMed PMID: 19017271; PubMed Central PMCID: PMCPMC2770442.

10. Klimpel KR, Arora N, Leppla SH. Anthrax toxin lethal factor contains a zinc metalloprotease consensus sequence which is required for lethal toxin activity. Mol Microbiol. 1994;13(6):1093-100. Epub 1994/09/01. doi: 10.1111/j.1365-2958.1994.tb00500.x. PubMed PMID: 7854123.

11. Tang W-J, Guo Q. The adenylyl cyclase activity of anthrax edema factor. Molecular Aspects of Medicine. 2009;30(6):423-30.

12. Liu S, Moayeri M, Leppla SH. Anthrax lethal and edema toxins in anthrax pathogenesis. Trends Microbiol. 2014;22(6):317-25. Epub 2014/04/02. doi: 10.1016/j.tim.2014.02.012. PubMed PMID: 24684968; PubMed Central PMCID: PMCPMC4041834.

13. Ebrahimi CM, Kern JW, Sheen TR, Ebrahimi-Fardooee MA, van Sorge NM, Schneewind O, et al. Penetration of the blood-brain barrier by Bacillus anthracis requires the pXO1-encoded BslA protein. J Bacteriol. 2009;191(23):7165-73. Epub 2009/10/13. doi: 10.1128/JB.00903-09. PubMed PMID: 19820089; PubMed Central PMCID: PMCPMC2786561.

14. Uchida I, Hornung JM, Thorne CB, Klimpel KR, Leppla SH. Cloning and characterization of a gene whose product is a trans-activator of anthrax toxin synthesis. J Bacteriol. 1993;175(17):5329-38. Epub 1993/09/01. doi: 10.1128/jb.175.17.5329-5338.1993. PubMed PMID: 8366021; PubMed Central PMCID: PMCPMC206586.

15. Dai Z, Sirard JC, Mock M, Koehler TM. The atxA gene product activates transcription of the anthrax toxin genes and is essential for virulence. Mol Microbiol. 1995;16(6):1171-81. Epub 1995/06/01. doi: 10.1111/j.1365-2958.1995.tb02340.x. PubMed PMID: 8577251.

16. Drysdale M, Bourgogne A, Hilsenbeck SG, Koehler TM. atxA controls Bacillus anthracis capsule synthesis via acpA and a newly discovered regulator, acpB. J Bacteriol. 2004;186(2):307-15. Epub

428   2004/01/02. doi: 10.1128/jb.186.2.307-315.2004. PubMed PMID: 14702298; PubMed Central PMCID:
429   PMCPMC305762.

17.   Sittner A, Bar-David E, Glinert I, Ben-Shmuel A, Schlomovitz J, Levy H, et al. Role of acpA and
431   acpB in Bacillus anthracis capsule accumulation and toxin independent pathogenicity in rabbits. Microb
432   Pathog. 2021;155:104904. Epub 2021/05/01. doi: 10.1016/j.micpath.2021.104904. PubMed PMID:
433   33930422.

18.   Fouet A. AtxA, a Bacillus anthracis global virulence regulator. Res Microbiol. 2010;161(9):735-42.
435   Epub 2010/09/25. doi: 10.1016/j.resmic.2010.09.006. PubMed PMID: 20863885.

19.   Levy H, Glinert I, Weiss S, Sittner A, Schlomovitz J, Altboum Z, et al. Toxin-independent
437   virulence of Bacillus anthracis in rabbits. PLoS One. 2014;9(1):e84947. Epub 2014/01/15. doi:
438   10.1371/journal.pone.0084947. PubMed PMID: 24416317; PubMed Central PMCID: PMCPMC3885664.

20.   Fouet A, Mock M. Differential influence of the two Bacillus anthracis plasmids on regulation of
440   virulence   gene   expression.   Infect   Immun.   1996;64(12):4928-32.   Epub   1996/12/01.   doi:
441   10.1128/IAI.64.12.4928-4932.1996. PubMed PMID: 8945528; PubMed Central PMCID: PMCPMC174470.

21.   Bertin M, Chateau A, Fouet A. Full expression of Bacillus anthracis toxin gene in the presence of
443   bicarbonate requires a 2.7-kb-long atxA mRNA that contains a terminator structure. Res Microbiol.
444   2010;161(4):249-59. Epub 2010/04/03. doi: 10.1016/j.resmic.2010.03.003. PubMed PMID: 20359529.

22.   Vietri NJ, Marrero R, Hoover TA, Welkos SL. Identification and characterization of a trans-
446   activator involved in the regulation of encapsulation by Bacillus anthracis. Gene. 1995;152(1):1-9. Epub
447   1995/01/11. doi: 10.1016/0378-1119(94)00662-c. PubMed PMID: 7828912.

23.   Levy H, Weiss S, Altboum Z, Schlomovitz J, Glinert I, Sittner A, et al. Differential Contribution
449   of Bacillus anthracis Toxins to Pathogenicity in Two Animal Models. Infect Immun. 2012;80(8):2623-31.
450   Epub 2012/05/16. doi: IAI.00244-12 [pii]
451   10.1128/IAI.00244-12. PubMed PMID: 22585968.

24.   Levy H, Fisher M, Ariel N, Altboum Z, Kobiler D. Identification of strain specific markers in
453   Bacillus anthracis by random amplification of polymorphic DNA. FEMS Microbiol Lett. 2005;244(1):199-
454   205. PubMed PMID: 15727841.

25.   Levy H, Weiss S, Altboum Z, Schlomovitz J, Rothschild N, Glinert I, et al. The effect of deletion
456   of the edema factor on Bacillus anthracis pathogenicity in guinea pigs and rabbits. Microb Pathog.
457   2012;52(1):55-60. Epub 2011/10/25. doi: S0882-4010(11)00184-7 [pii]
458   10.1016/j.micpath.2011.10.002. PubMed PMID: 22020310.

26.   Makdasi E, Laskar O, Glinert I, Alcalay R, Mechaly A, Levy H. Rapid and Sensitive Multiplex Assay
460   for the Detection of B. anthracis Spores from Environmental Samples. Pathogens. 2020;9(3). Epub
461   2020/03/04. doi: 10.3390/pathogens9030164. PubMed PMID: 32120986; PubMed Central PMCID:
462   PMCPMC7157734.

27.   Welkos SL. Plasmid-associated virulence factors of non-toxigenic (pX01-) Bacillus anthracis.
464   Microb Pathog. 1991;10(3):183-98. Epub 1991/03/01. doi: 10.1016/0882-4010(91)90053-d. PubMed PMID:
465   1654495.

28.   Chiang C, Bongiorni C, Perego M. Glucose-dependent activation of Bacillus anthracis toxin gene
467   expression and virulence requires the carbon catabolite protein CcpA. J Bacteriol. 2011;193(1):52-62.
468   Epub 2010/10/26. doi: 10.1128/JB.01656-09. PubMed PMID: 20971911; PubMed Central PMCID:
469   PMCPMC3019961.

29.   McCall RM, Sievers ME, Fattah R, Ghirlando R, Pomerantsev AP, Leppla SH. Bacillus anthracis
471   Virulence Regulator AtxA Binds Specifically to the pagA Promoter Region. J Bacteriol. 2019;201(23). Epub
472   2019/10/02. doi: 10.1128/JB.00569-19. PubMed PMID: 31570528; PubMed Central PMCID:
473   PMCPMC6832065.

30.   Guignot J, Mock M, Fouet A. AtxA activates the transcription of genes harbored by both Bacillus
475   anthracis virulence plasmids. FEMS Microbiol Lett. 1997;147(2):203-7. Epub 1997/02/15. doi:
476   10.1111/j.1574-6968.1997.tb10242.x. PubMed PMID: 9119194.

477   31.   Drysdale M, Bourgogne A, Koehler TM. Transcriptional analysis of the Bacillus anthracis capsule
478   regulators. J Bacteriol. 2005;187(15):5108-14. Epub 2005/07/21. doi: 10.1128/JB.187.15.5108-
479   5114.2005. PubMed PMID: 16030203; PubMed Central PMCID: PMCPMC1196023.
480