1

An averaging model for analysis and interpretation of high-order genetic interactions

Fumiaki Katagiri

Department of Plant and Microbial Biology, Microbial and Plant Genomics Institute, University of

Minnesota, St. Paul, MN 55108, USA

katagiri@umn.edu

Author contribution: FK is the sole author and responsible for all the content

Competing Interest Statement: The author declares no competing interest

Keywords: multi-gene system; high-order genetic interaction; distributive law; general linear model;

averaging model

This PDF file includes:

Main Text

Figures 1 to 5

Supplemental Texts S1 and S2

ABSTRACT

While combinatorial genetic data collection from biological systems in which quantitative phenotypes are controlled by functional and non-functional alleles of multiple genes (multi-gene systems) is becoming common, a standard analysis method for such data has not been established. A common additive model of the non-functional allele effects contrasted against the functional alleles, based on ANOVA with interaction, has three issues. First, although it is a long tradition in genetics, modeling the effect of the non-functional allele (a null mutant allele) contrasted against that of the functional allele (the wild-type allele) is not suitable for mechanistic understanding of multi-gene systems. Second, an additive model fails in estimation of interactions among more than two genes when the phenotypic response is not linear. Third, interpretation of higher-order interactions defined by an additive model is not intuitive. I propose an averaging model, which is suitable for mechanistic understanding of multi-gene systems: the effect of the functional allele is contrasted against the effect of the non-functional allele for easier mechanistic interpretations; it is stable in estimation of higher-order interactions even when the phenotypic response is not linear; and the higher-order interactions it defines are highly intuitive. Yet, the averaging model is still a general linear model, so model fitting is easy and accurate using common statistical tools.

INTRODUCTION

Accumulation of genetic knowledge in many biological systems and technological advances that made combining multiple genetic loci easier have facilitated combinatorial genetic analysis among multiple genes, each of which has functional ("wild-type") and non-functional (null "mutant") allele states, involved in single quantitative traits, e.g., (1-4), which I here call multi-gene systems. However, conventional genetics is not well built for analysis and interpretation of high-order genetic interactions among multiple genes involved in a single quantitative trait. This is because conventional genetics is an extension of early objectives of analyzing functionally independent and/or qualitative genes. First, comparing multiple mutant phenotypes to the wild-type phenotype does not allow simple mechanistic interpretations. The phenotype of a particular genotype should be compared to the phenotype of the most disrupted mutant state (e.g., a quadruple null mutant in a 4-gene system) for simple mechanistic interpretations.

Second, a method to define and interpret interactions among multiple genes is not definitively integrated. The main topic of this paper concerns this second point. An additive model based on ANOVA with interaction is a simple implementation for analysis of high-order genetic interactions. However, such an additive model fails with a nonlinear system, such as a system with a saturating response, when it contains genetic interactions among more than two genes. This problem is caused by the fact that the additive model requires the conservation of the distributive law ((A + B):C = A:C + B:C, where ":" indicates the interaction defined in the additive model), whereas conservation of this law cannot be assumed in a nonlinear system. We previously proposed a network reconstitution (formerly called signaling allocation) general linear model (NR model), which does not assume conservation of the distributive law, to resolve this problem (2, 5). This was achieved by averaging the interaction terms at each order ("averaging principle"), instead of simply adding them (e.g., ABC = A + B + C + (A:B + B:C + C:A) / 3 +A:B:C in the NR model, while ABC = A + B + C + A:B + B:C + C:A + A:B:C in the additive model; the italicized upper-case letters denote the genotype carrying wild-type alleles A, B, and C; the intercept is omitted).

I recently realized that the interaction operator in the NR model is not consistent, e.g., in the above example of the ABC description, The ":" operators in A:B and A:B:C are not defined in a consistent manner. This was caused by the fact that the 2-gene interaction was defined in the same way in both the NR and additive models, AB = A + B + A:B. To resolve this issue, the averaging principle should be extended to the 1-gene effect terms as well: e.g., ABC = (A + B + C) / 3 + (A;B + B;C + C;A) / 3 +A;B;C. I call this an averaging model. It is a general linear model with the averaging principle applied consistently. I use a ";" to denote the interaction operator defined in the averaging model ("averaging interaction").

The averaging model does not assume distributivity. Consequently, it can estimate high-order genetic interactions in a stable manner with nonlinear multi-gene systems. This is because each averaging interaction can be defined using only observed values, e.g., A;B;C = ABC − (AB + AC + BC) / 3. Since all the interaction terms are defined simply and consistently, the interpretations of averaging interactions are intuitive and consistent: a genetic interaction is the average impact of adding the last gene; note that the above definition of A;B;C is the equivalent of A;B;C = { (ABC − AB) + (ABC − AC) + (ABC − BC) } / 3. I

83   further demonstrate that the additive and averaging models without interaction can be considered as
84   two extreme approximations in a simple 2-gene system: there is no mechanistic reason to favor one of
85   the models without additional mechanistic information. Since the averaging model does not fail in
86   estimations of high-order genetic interactions, unlike the additive model, I propose the averaging model
87   as a standard general linear model for descriptions of multi-gene system behaviors.
88
89
90   **RESULTS AND DISCUSSION**
91
92   *Objective of the study*
93
I define a multi-gene system as one in which multiple genes affect a single quantitative trait while each
gene can have two states, functional and non-functional, arising from the wild-type and null mutant
alleles, respectively. Such a system necessarily implies a gene network, in which the gene functions are
not organized in a series (i.e., not in a single pathway). This is because a series of genes, each of which
can only take a functional or non-functional state, can only generate an on-or-off, non-quantitative
output. Instead, a gene network must have a converging node(s) to generate a single trait. Converging
nodes are sources of complex system behaviors (6, 7). For a data set, I consider the measurement of the
quantitative trait as the phenotype and measurements made with exhaustively combinatorial genotypes
(i.e., for a *n*-gene system, the number of the exhaustively combinatorial genotypes is 2^n) as the data.
103
104   The objective of this study is to best describe the output of such a multi-gene system using the general
105   linear model framework to facilitate mechanistic interpretations of the system behavior. The approach
106   was limited to a general linear model framework due to assumed lack of mechanistic information for
107   more complex (nonlinear) modeling. Fitting a general linear model is computationally easy and accurate.
108   The general linear model could serve as a simple and versatile platform in many cases.
109
110
111   *General notation rules*
112
I assume that all the genes of interest are homozygous in diploid organisms. A single gene is denoted by
a single alphabetical letter in italics, with upper-case and lower-case letters denoting the wild-type and
null mutant alleles, respectively. For example, *ABc* represents the genotype with the wild-type alleles for
genes *A* and *B* and the mutant allele for gene *C*. I also use the genotype notation omitting the mutant
alleles, such as *AB* instead of *ABc*, for simplicity, clarity, and generalization. The notation of a genotype
can represent its phenotype as well. The non-italic lower-case letters, such as a, b, and c, represent the
mutant allele effects defined in comparison to the wild-type alleles. The wild-type allele effects,
represented by non-italic upper-case letters, such as A, B, and C, are defined in comparison with the
mutant allele. The additive effect of A and B is denoted A + B. The additive and averaging interactions
between A and B are denoted A:B and A;B, respectively. In a mechanistic network model underlying the
observation, the node corresponding to gene A and its output are denoted as nA. Although I often use a

3-gene system, *ABC*, with the intercept term omitted for simple examples, the points discussed
subsequently can be generalized to a system consisting of an arbitrary number of genes.
126
127
128   *Comparing to the most disrupted state instead of the intact state gives better interpretability*
129
A convention in genetics is to compare a mutant phenotype to the wild-type phenotype. Here I argue
that instead, comparing a phenotype of any genotype to the phenotype of the most disrupted state,
e.g., comparing to the triple mutant state in a 3-gene system, leads to much better mechanistic
interpretations. For a simple example, I use a system defined by an ANOVA-based, 3-gene additive
model, i.e., assuming a linear phenotypic response (although I will later discuss a linear assumption
issue).
136

Fig. 1. A simple network behavior can be well described by the wild-type allele effects of a multi-gene system but not by the mutant allele effects. (a) A mechanistic model of a network containing 3 nodes that can be mutationally manipulated (a 3-gene system). The network consists of 6 nodes, among which nA, nB, and nC are mutationally manipulable and nX, nY, and nZ are not. The output of each node is given either as a value or an equation. The output of nZ is the quantitative phenotype of the system. (b) The phenotype values of all 8 combinatorial genotypes. (c) The values for the mutant allele effects and interactions. (d) The values for the wild-type allele effects and interactions.

138
Fig. 1a shows a mechanistic network underlying a system with 6 nodes, in which three nodes (nA, nB,
and nC) can be manipulated by mutations and the other three (nX, nY, and nZ) cannot. Thus, for the
purpose of genetic analysis, this is a 3-gene system. nA, nB, nX, and nY are input nodes, and their values
are arbitrarily set at 5, -3, 4, and 2, respectively. nZ is the output node, and the output of nZ can be
measured as the quantitative trait of the system. Simple additive rules at nodes nC and nZ are assumed,
nC = nA + nB + nX and nZ = nC + nY, respectively. Fig. 1b shows the nZ output (i.e., phenotype) of 8
exhaustively combinatorial genotypes. Fig. 1c shows the effects and interactions of the mutant alleles
that are calculated according to an ANOVA model with interaction. Fig. 1d shows the effects and
interactions of the wild-type alleles that are calculated according to an ANOVA model with interaction.
With Fig. 1d, it is easy to reconstitute the mechanistic network shown in Fig. 1a: there is a basal activity

of 2 in the absence of A, B, and C; A and B are not active by themselves, while C has its own activity of 4
regardless of A and B; the connection between A and C is positive with a value of 5, and the connection
between B and C is negative with a value of -3; No A:B:C interaction means that additive effects up to
two-gene interactions can explain the system behavior completely. In comparison, mechanistic
interpretations based on Fig. 1c are not simple: e.g., it is not easy to decipher that B is completely
dependent on C from *ABC* = 8, b = 3, c = -6, and b:c = -3. I conclude that the behavior of a multi-gene
system should be interpreted using wild-type allele effects. I will subsequently model a system with
wild-type allele effects and their interactions.
157
158
159 *Derivation of the averaging model*
160
See Text S1 for details of this section. The additive model requires the distributive law but this law is
violated when the system behavior is nonlinear. A typical biological system has a limited phenotype
range. When the sizes of the effects and interactions of the genes in a multi-gene system are substantial
compared to the phenotype range, the system behavior is nonlinear, and the distributive law is violated
- consequently the additive model fails. The NR model (2, 5) allows non-distributivity. However, the
definitions of the 2-gene interaction and of the 3-or-more-genes interaction were not consistent in the
NR model. The averaging model was derived by extending the definition of the 3-or-more-genes
interaction in the NR model to the 2-gene interaction. Thus, the averaging model allows non-
distributivity, and its interaction definition is consistent regardless of the number of genes involved. The
three models and a 1-way ANOVA model for each genotype are all general linear models with different
ways to decompose the fitted values. When the models are fit to data with replication as statistical
models, their fitted values and residuals are identical. Therefore, subsequently I sometimes use only the
mean estimates of the models for model comparisons (such as Fig. 2).
174
175
176 *Unlike the additive model, the averaging model stably estimates high-order genetic interactions*
177
The situation in which the additive model fails due to a limited phenotype range was simulated using a
7-gene system. A system with many genes was used because it shows the problem more clearly. With 7
genes, the number of combinatorial genotypes is 2^7 = 128. A simulated mean phenotype value for each
genotype was randomly sampled from a uniform distribution ranging from 1 to 10, and each model was
solved using these randomly generated data values. This simulation procedure was repeated 10,000
times and the model estimate distributions, except for the model intercept (i.e., the septuple mutant
value), were visualized as a box plot (Fig. 2). Fig. 2a shows that in the additive model, the higher the
order of interactions is, the higher the representations of the interactions are. The length of the box (the
difference between the 75[th] and 25[th] percentiles) of the 7-gene interaction is about 7.5 times larger than
those of the 1-gene effects and about 5 times larger than the phenotype range. Therefore, if the
additive model is used, the absolute values of higher order additive interactions are grossly
overestimated in general. This problem was solved in the NR model, except for the 2-gene interactions,
which were still additive interactions (Fig. 2b). Note the *y*-axis scale difference between Fig. 2a and Figs.

2b and 2c: the distributions of the 1-gene effects are essentially the same across the models. The
distributions of estimates were very consistent across all the effects and averaging interactions in the
averaging model (Fig. 2c). These results strongly suggest that the averaging interactions that do not
assume distributivity can be stably estimated even when the system behavior is nonlinear.
195

Fig. 2. Distributions of the gene effects and
interaction values when the phenotype values
were randomly sampled from a uniform
distribution with (a) additive, (b) NR, or (c)
averaging models. Each order of interactions is
color-coded separately, and the color coding is
shown at the bottom of (a). Note that the scales in
the y-axes are much larger in (a) than in (b) or (c).

196
197
To investigate a situation with actual biological data, we reanalyzed our previously published data set
with a 4-gene system (2) using the additive, NR, and averaging models (Fig. 3). Here we focus on the
behaviors of the model estimates from the three models. Biological information about the data set is
described later. Note that the data set had replicated observations for each combinatorial mutant, the
models were fit to the data (instead of solving), and the confidence intervals of the model estimates
were obtained. The 95% confidence intervals of the full additive models (which include up to the 4-gene
interaction) were evidently wider with high-order interactions in the additive model (Fig. 3a, black bars).
This problem of uneven confidence interval sizes was moderated in the NR model (Fig. 3b) and not
noticeable in the averaging model (Fig. 3c). Furthermore, in the additive model, since the effect sizes of
higher order interactions were overestimated, the estimates for the 1-gene effect and lower order
interactions were substantially affected when higher-order interactions were omitted (model reduction;
Fig. 3a, compare red to black and blue to red). In addition, the 95% confidence interval sizes of the same

terms were affected by model reduction in the additive model. These problems in model reduction were
moderated in the NR model, and almost non-existent in the averaging model (Figs. 3b and 3c). In
summary, the problems associated with non-distributivity in this biological data set are evident in the
additive model while the averaging model appears free of these problems. These observations made
with a biological data set directly corroborate with the simulation results in Fig. 2
215

Fig. 3. Coefficient estimates for the contribution to immunity using the data from Tsuda et al. in (a) additive, (b) NR, and (c) averaging models. The 95% confidence interval is shown as a horizontal bar, with the mean as a point. Different levels of model reduction (omitting higher order interactions from the model) are color-coded according to the color code in (a). Different shades of gray background are used to show different orders of interactions. ":" and ";" indicate additive and averaging interactions, respectively.

216
217
218
219   *Why does the averaging model describe a multi-gene system better than the additive model?*
220
In the 2-gene additive interaction, A:B = $AB - (A + B)$, the 2-gene additive interaction A:B can be
estimated using only observed values, i.e., estimation of A:B can be independent of other model
estimates. Essentially, an error originated from the nonlinearity of data is absorbed by the 2-gene
additive interaction A:B. However, estimation of a 3-gene additive interaction, A:B:C = $ABC - (A + B + C)$
- (A:B + A:C + B:C), is not independent of other model estimates, A:B, A:C, and B:C. In a nonlinear system,
there is no guarantee that the value of A:B in the 3-gene model is the same as that of A:B estimated by
the 2-gene model. Thus, on average, higher-order interactions accumulate errors from lower-order
interactions in the additive model with a nonlinear system, which was demonstrated in simulation and
analysis of an actual data set (Figs. 2a and 3a).

On the other hand, any order of averaging interactions can be estimated using only observed values,

e.g., A;B;C = $ABC - (AB + AC + BC) / 3$, so their estimations can be independent of other model estimates.

Thus, nonlinearity-originated errors are confined to individual averaging interactions and do not

accumulate toward higher-order interactions (Figs. 2c and 3c). This is the reason the additive model

including 3-or-more-gene interactions fails with nonlinear systems while the averaging model does not.

Interpretation of the averaging model outcome

It should be emphasized that the definitions of the additive and averaging interactions are different, and

consequently their interpretations are different. The 2-gene additive interaction is understood as the

difference from the addition of the 1-gene effects, A:B = $AB - (A + B)$. When A, B, A:B > 0, A and B have a

synergistic effect. When A, B > 0, A:B < 0, A and B have a compensating effect (Fig. 4a). However,

synergistic or compensating interpretations of additive interactions become unclear when A and B have

opposite signs (Fig. 4b) or the additive interactions are higher-order, e.g., A:B:C = $ABC - (A + B + C + A:B +$

A:C + B:C) (Fig. 4c).

In contrast, the interpretation of averaging interactions is consistent and highly interpretable at all

orders of the averaging interactions, i.e., the averaging model is highly scalable to the number of genes

in the system. An averaging interaction is the deviation of the corresponding genotype from the average

of all involved genotypes that have one gene fewer. For example, in a 2-gene system, A;B = $AB - (A +$

$B)/2$ (Fig. 4d). Note that not just the values but also the signs of the interactions could be different

between the additive and averaging interactions (compare *AB* (case 1) in Figs. 4a and 4d). The

interpretations of averaging interactions are consistent even when A and B have opposite signs (Fig. 4e)

or in the case of higher-order averaging interactions, e.g., A;B;C = $ABC - (AB + AC + BC) / 3$ (Fig. 4f).

A 3-gene averaging interaction, A;B;C = $ABC - (AB + AC + BC) / 3$, (Fig. 4d) could be a 3-gene interaction

in a 7-gene system, A;B;C = $ABCdefg - (ABcdefg + AbCdefg + aBCdefg) / 3$. Thus, a genotype notation

that omits the mutant alleles is a generalized notation for the averaging model.

Fig. 4. Interpretations of interactions in (a-c) additive and (d-f) averaging models. (a, d) Two-gene interactions when both 1-gene effects A and B are positive. Two different cases (cases 1 and 2) of the *AB* phenotype values are used. (b, e) Two-gene interactions when the 1-gene effects have opposite signs. (c, f) Three-gene interactions. ":" and ";" indicate additive and averaging interactions, respectively.

An averaging model-based multi-gene analysis should only contain genes significantly involved in the phenotype.

Since the averaging interaction is the phenotypic deviation of the corresponding genotype from the average of all genotypes with one gene fewer, it is affected if the analysis includes unnecessary genes. Such unnecessary genes can be detected by comparing all the genotypes containing the gene in question to the corresponding genotypes without the gene. For example, in a 3-gene system with genes *A*, *B*, and *C*, the test for whether gene *C* should be included is whether any of $ABC - AB$, $AC - A$, $BC - B$, and $C - abc$ have values significantly different from 0. If none of them are significant, gene *C* should be removed from the averaging model.

Reinterpretation of previous results using the averaging model

Using the averaging model, I reinterpreted results from my laboratory of exhaustively combinatorial genotype analysis in a 4-gene system, which were originally analyzed using the NR model shown in Fig. 4 of (2). The study consisted of four cases of inducible immunity in the model plant Arabidopsis against strains of the bacterial pathogen *Pseudomonas syringae*, which are designated as the AvrRpt2-ETI, AvrRpm1-ETI, flg22-PTI, and elf18-PTI cases. ETI is Effector-Triggered Immunity, and AvrRpt2 and AvrRpm1 are triggering effectors (8-12). PTI is Pattern-Triggered Immunity, and flg22 and elf18 are triggering molecular patterns (13-15). The inhibition of bacterial growth in the plant leaf, in

285      $\log_{10}(\text{cfu/cm}^2)$, was the immunity phenotype measure. The hub genes of four major signaling sectors
286   (subnetworks) in the plant immune signaling network were subjected to mutational analysis. The
287   signaling sectors were the jasmonate, ethylene, PAD4, and salicylate sectors, which are indicated as J, E,
288   P, and S, respectively. I also call their hub genes *J*, *E*, *P*, and *S*, in this context of analysis of the 4-gene
289   system. Biological and experimental details are provided in (2).
290
Each of the AvrRpt2-ETI, AvrRpm1-ETI, flg22-PTI, and elf18-PTI cases was first tested to determine
whether all four genes were significantly involved in the phenotype variation. Except for the elf18-PTI
case, all four genes were significant, and the averaging model for the 4-gene system was used. In the
elf18-PTI case, the phenotype was not significantly affected by the *J* gene in any genotype context, so
the averaging model for the 3-gene system with the *E*, *P*, and *S* genes was used. The results of
applications of the averaging model to these four immunity cases are shown in Fig. 5.
297

Fig. 5. Coefficient estimates for the contribution to immunity from averaging model analysis of the data in Tsuda et al. (a) AvrRpt2-ETI, (b) AvrRpm1-ETI, (c) flg22-PTI, and (d) elf18-PTI. The 95% confidence intervals are shown as horizontal bars, with the means as points. The Holm-corrected *p*-values are shown in the left part of each plot: red, $p < 0.05$; blue, $p \geq 0.05$. The dataset used for AvrRpt2-ETI in Fig. 5a is the same as that in used in Fig. 3, and Fig. 5a is the same as the full model (black lines) in Fig. 3c. ";" indicates the averaging interaction.

298
299
300   In AvrRpt2-ETI with the averaging model (Fig. 5a), the values for the 1-gene effects were all positive. P
301   had the largest effect, which was close to the wild-type level. Among the 2-gene averaging interactions,
302   only J;S was significantly positive: while the J and S effects are both positive, combining these two genes
303   together (*JS* genotype) increases the immunity from the average of the *J* and *S* genotypes, which makes
304   the *JS* phenotype similar to wild type.
305
In AvrRpm1-ETI with the averaging model (Fig. 5b), most immunity was explained by the intercept (i.e.,
the immunity level in the *jeps* genotype), showing that the quadruple mutant still maintains most of the
immunity of wild-type plants. This observation can be explained by the fast kinetics of AvrRpm1-ETI
signaling compared to AvrRpt2-ETI, in respect to the gating timing of the ETI-Mediated and PTI-Inhibited
Sector (EMPIS) by PTI signaling (16). Although all the 1-gene effects and the averaging interactions had
lower amplitudes, their general up and down trends were similar to those of AvrRpt2-ETI, suggesting
that the 4-gene network apart from EMPIS behaves similarly in AvrRpm1-ETI and AvrRpt2 ETI. J;S was
the only significant averaging interaction with a positive contribution to immunity. In addition, the
positive P effect was marginally significant (Holm-corrected $p = 0.054$).

In flg22-PTI (Fig. 5c), all the 1-gene effects except S were significantly positive with E as the highest. The

2-gene averaging interactions were largely low and/or not significant, except P;S, which was significantly

and strongly positive. The 3-gene and 4-gene averaging interactions were significantly and strongly

positive, indicating that all the genes substantially increase the immunity level when added to the

system as the 3rd or 4th genes.

In elf18-PTI (Fig. 5d), only P;S was significant among all the averaging model terms, except the intercept.

The P;S averaging interaction was strongly positive, indicating that a single mutation in genes *P* or *S*

almost completely abolishes elf18-PTI. The difference in the importance of the *E* gene clearly separated

flg22-PTI and elf18-PTI. Another difference between flg22-PTI and elf18-PTI was the 3-gene and 4-gene

averaging interactions. All were strongly positive in flg22-PTI, and none were significant in elf18-PTI.

It is noteworthy that the roles of J;S and P;S were very different in ETI and PTI. A strongly positive P;S

averaging interaction was observed in PTI (Figs. 5c and 5d). Positive functional interactions between the

P and *S* genes have been well documented in many aspects of plant immunity (17). In contrast, this

averaging interaction was insignificant in ETI, except for their contributions through higher-order

averaging interactions, J;P;S, E;P;S, and J;E;P;S. On the other hand, a strongly positive J;S averaging

interaction was observed in ETI while it was insignificant in PTI (Fig. 5). Although negative functional

interactions between the *J* and *S* genes are often described in plant immunity (17), these two genes

interact positively in ETI (Figs. 5a and 5b). In addition, in flg22-PTI the 3-gene and 4-gene averaging

interactions were strongly positive while they were moderately positive in AvrRpt2-ETI. A disadvantage

of strongly positive 3-gene and 4-gene averaging interactions is that a mutation(s) in one or two genes

results in large loss of immunity. Relatively weak 3-gene and 4-gene averaging interactions in ETI

indicates that ETI is more resilient against damage to one or two of these major immune signaling

sectors, which could be caused by pathogen effectors (7). In summary, the averaging model analysis

highlighted that while the 4-gene system is important in both ETI and PTI (with flg22), the way they are

used in ETI and PTI is quite different, and ETI is more resilient than PTI against perturbations to the

signaling sectors. It also highlighted substantial differences, particularly in the roles of the *J* and *E* genes,

in regulation between flg22-PTI and elf18-PTI.

Mechanistic basis of the additive and averaging models

To provide a concrete idea about what the additive and averaging models without interaction represent

mechanistically, I use the following idealized 2-gene signaling network based on chemical reactions. Two

genes *A* and *B* qualitatively (i.e., on or off) control two input nodes nA and nB of an output node nC (i.e.,

directed links from nA and nB converging on nC). The activity of each node is represented by the

corresponding chemical, A, B, or C, and the chemical reactions involved are A + X → C and B + Y → C. We

assume an equilibrium (or a steady state) for the reactions, and the concentration relationships, [X], [Y]

>> [A], [B], [C], i.e., [X] and [Y] can be considered constant in the reactions. Under these conditions, we

can describe the relationships among [A], [B], and [C] as: k_A[A] = [C] and k_B[B] = [C], where k_A and k_B are

positive constants. Then the additive and averaging models without interaction can well approximate
the system behavior when k_A, $k_B \gg 1$ (i.e., [C] $\gg$ [A],[B]) and when k_A, $k_B \ll 1$ (i.e., [C] $\ll$ [A],[B]),
respectively (Text S2). This discussion of an idealized system shows that the additive and averaging
models without interaction could represent two opposing extreme approximations of the underlying
mechanistic state. Therefore, one of the models cannot be favored over the other unless mechanistic
information about the system is available in addition to the phenotypic values.
363
364
365 *Limitation in mechanistic interpretations with a general linear model*
366
The chemical reaction system discussion in the last section also shows that when the actual values of k_A
and k_B are between the extreme values, both models handle the deviation with the interaction, A:B or
A;B (with opposite signs). Note that the interaction is a consequence of shared [C] between two
reactions: one reaction affects the other, i.e., it is truly a mechanistic interaction. However, the
interaction value from a model cannot reveal the nature of the interaction – we can interpret it
mechanistically in this example only because we know the underlying system mechanism. This example
illustrates a limitation to describing a nonlinear system with a general linear model in terms of
mechanistic interpretations based on the model. However, it should also be reemphasized that the
genetic interpretations of higher-order interactions are always clear with the averaging model while this
is not the case with the additive model.
377
378
379 *Concluding Remarks*
380
I have demonstrated that multi-gene systems subjected to exhaustively combinatorial mutation analysis
typically violate the distributive law due to their nonlinearity and that therefore, the additive model that
requires the conservation of the distributive law is not appropriate for analysis of such systems. In
contrast, an averaging model allows non-distributivity and maintains consistency from the 1-gene
effects to the highest order of averaging interactions. Furthermore, averaging model results are
consistently and intuitively interpretable from the 1-gene effects to the highest order of averaging
interactions. I propose the averaging model as a standard general linear model for combinatorial
mutation analysis of multi-gene systems.
389
390
391
392 **METHODS**
393
394 *Data sets*
395
396 Biological data sets used in this study are the same data sets used in Figs. 4A and 4B in Tsuda et al.
397 (2009). Each data set consists of bacterial counts ($\log_{10}$(colony forming units/cm^2)) for 16 exhaustively

398 combinatorial genotypes for a 4-gene system, with or without treatment, with replication. Since the raw
399 bacterial count data were not published previously, they are provided as Supplemental Dataset 1.
400
401
402 *Random simulation with three models*
403
The simulation was performed with a 7-gene system. The phenotype values for $2^7 = 128$ genotypes were
randomly sampled from a uniform distribution ranging from 1 to 10. The 128 phenotype values were
solved for the coefficients (gene effects and interactions) in each of the additive, NR, and averaging
models. To solve the 128 equations per model, the 7-gene system matrix equivalent of the 3-gene
system matrix in Fig. TS1.1 (Text S1) was used (the matrices are provided in an R workspace file in
Supplemental Dataset 2). This procedure was repeated 10,000 times for each model, and the
distribution of each coefficient (except the intercept) across the repeats is shown by a box-and-whiskers
in Fig. 2.
412
413
414 *Fitting averaging models to the data*
415
A linear mixed-effect model (the lme function in the nlme R package (18)) was used. This was because (i)
each data set has factors regarding the experimental design, which were included as random effects in
the model and (ii) the numbers of replicates were not the same across the genotype x treatment
combinations. First, a linear mixed-effect model with the genotype x treatment interactions was fit to
each of the data sets for "AvrRpt2-ETI", "AvrRpm1-ETI", "flg22-PTI", and "elf18-PTI". The formula for the
fixed effects was "`~ genotype/treatment -1`". The random effects for the data sets were "`~
1|replicate/flat/pot`". The interaction coefficients of the linear mixed-effect model were used
to test whether each gene is significant. For example, to test the significance of the *J* gene, the estimate
differences, *JEPS – EPS*, *JEP – EP*, *JPS – PS*, *JES – ES*, *JE – E*, *JP – P*, *JS – S*, and *J – jeps* were subjected to *t*-
tests using the associated standard errors calculated from the variance/covariance matrix and the
residual degree of freedom. The *p*-values from all *t*-tests for a single data set were correct by the
Benjamini-Hochberg FDR. If none of the corrected *p*-values were smaller than 0.05 for the gene, the
gene was designated insignificant and omitted from the following averaging model analysis. Only the *J*
gene in "elf18-PTI" was insignificant. In this case, the data were bundled by ignoring the *J* gene. For
example, the *JEPS* data were considered as part of the *EPS* data.
431
Second, the averaging model using the significant genes was fit. The 4-gene system equivalent matrix of
the 3-gene system matrix in Fig. TS1.1c (Text S1) or the 3-gene system matrix was used (the matrices are
provided in an R workspace file in Supplemental Dataset 2). The rows were replicated according to the
genotypes of the observations (the design matrix for the averaging model coefficients, denoted as "m.").
Using the design matrix m., the fixed effects were, "`~ m. -1 + genotype`" and the random effects
were, "`~ 1|replicate/flat/pot`" in the lme function. The averaging model coefficient estimates,
their standard errors, and the *p*-values were extracted from the coefficient table of the lme model. The
estimates, the standard errors, and the residual degree of freedom of the lme model were used to

calculate the 95% confidence intervals. The *p*-values were subjected to the Holm multiple tests

correction. The R script used to generate Fig. 5 from the raw bacterial count data sets is provided as

Supplemental Dataset 3. Supplemental Datasets are available from

https://github.com/fumikatagiri/Averaging_Model.

ACKNOWLEDGEMENTS

I thank Dave Mackey and Alex Turo for exposing me to their unpublished data from a 7-gene system,

which led me to realization of the averaging model, Takashi Hirayama for his question, which led to the

discussion regarding the additive and averaging models as approximated behaviors of a chemical

signaling network, and Kenichi Tsuda for the raw data used in Figs. 3 and 5 and for critical reading of the

manuscript. I also thank Yaniv Brandvain, Ruth Shaw, and Linda Jeanguenin for critical reading of the

manuscript and Jane Glazebrook for editing. This work was supported by grants from National Science

Foundation (MCB-1518058 and IOS-1645460) and from US Department of Agriculture-National Institute

of Food and Agriculture (2020-67013-31187).

REFERENCES

1.   R. A. Hillmer *et al.*, The highly buffered Arabidopsis immune signaling network conceals the
functions of its components. *PLoS Genet* **13**, e1006639 (2017).

2.   K. Tsuda, M. Sato, T. Stoddard, J. Glazebrook, F. Katagiri, Network properties of robust immunity
in plants. *PLoS Genet* **5**, e1000772 (2009).

3.   A. Celaj *et al.*, Highly Combinatorial Genetic Interaction Analysis Reveals a Multi-Drug
Transporter Influence Network. *Cell Syst* **10**, 25-38.e10 (2020).

4.   E. Kuzmin *et al.*, Exploring whole-genome duplicate gene retention with complex genetic
interaction analysis. *Science* **368** (2020).

5.   F. Katagiri, Network Reconstitution for Quantitative Subnetwork Interaction Analysis. *Methods*
Mol Biol **1578**, 223-231 (2017).

6.   F. Katagiri, A global view of defense gene expression regulation - a highly interconnected
signaling network. *Current Opinion in Plant Biology* **7**, 506-511 (2004).

7.   F. Katagiri, Review: Plant immune signaling from a network perspective. *Plant Sci* **276**, 14-21
(2018).

8.   J. D. G. Jones, J. L. Dangl, The plant immune system. *Nature* **444**, 323-329 (2006).

9.   S. T. Chisholm, G. Coaker, B. Day, B. J. Staskawicz, Host-microbe interactions: Shaping the
evolution of the plant immune response. *Cell* **124**, 803-814 (2006).

10.   M. Mindrinos, F. Katagiri, G. L. Yu, F. M. Ausubel, The A. thaliana disease resistance gene RPS2
encodes a protein containing a nucleotide-binding site and leucine-rich repeats. *Cell* **78**, 1089-
1099 (1994).

11.   A. F. Bent *et al.*, RPS2 OF ARABIDOPSIS-THALIANA - A LEUCINE-RICH REPEAT CLASS OF PLANT-
DISEASE RESISTANCE GENES. *Science* **265**, 1856-1860 (1994).

12.   M. R. Grant *et al.*, STRUCTURE OF THE ARABIDOPSIS RPM1 GENE ENABLING DUAL-SPECIFICITY
DISEASE RESISTANCE. *Science* **269**, 843-846 (1995).

13.   C. Zipfel *et al.*, Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature*
428, 764-767 (2004).
14.   L. Gómez-Gómez, G. Felix, T. Boller, A single locus determines sensitivity to bacterial flagellin in
Arabidopsis thaliana. *Plant J* **18**, 277-284 (1999).
15.   C. Zipfel *et al.*, Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts
Agrobacterium-mediated transformation. *Cell* **125**, 749-760 (2006).
16.   N. Hatsugai *et al.*, A plant effector-triggered immunity signaling sector is inhibited by pattern-
triggered immunity. *EMBO J* **36**, 2758-2769 (2017).
17.   C. M. J. Pieterse, D. Van der Does, C. Zamioudis, A. Leon-Reyes, S. C. M. Van Wees, Hormonal
Modulation of Plant Immunity. *Annual Review of Cell and Developmental Biology, Vol 28* **28**,
489-521 (2012).
18.   J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar, R. C. Team) (2021) nlme: Linear and Nonlinear Mixed
Effects Models.

FIGURE LEGENDS

Fig. 1. A simple network behavior can be well described by the wild-type allele effects of a multi-gene
system but not by the mutant allele effects. (a) A mechanistic model of a network containing 3 nodes
that can be mutationally manipulated (a 3-gene system). The network consists of 6 nodes, among which
nA, nB, and nC are mutationally manipulable and nX, nY, and nZ are not. The output of each node is
given either as a value or an equation. The output of nZ is the quantitative phenotype of the system. (b)
The phenotype values of all 8 combinatorial genotypes. (c) The values for the mutant allele effects and
interactions. (d) The values for the wild-type allele effects and interactions.

Fig. 2. Distributions of the gene effects and interaction values when the phenotype values were
randomly sampled from a uniform distribution with (a) additive, (b) NR, or (c) averaging models. Each
order of interactions is color-coded separately, and the color coding is shown at the bottom of (a). Note
that the scales in the y-axes are much larger in (a) than in (b) or (c).

Fig. 3. Coefficient estimates for the contribution to immunity using the data from Tsuda et al. in (a)
additive, (b) NR, and (c) averaging models. The 95% confidence interval is shown as a horizontal bar,
with the mean as a point. Different levels of model reduction (omitting higher order interactions from
the model) are color-coded according to the color code in (a). Different shades of gray background are
used to show different orders of interactions. ":" and ";" indicate additive and averaging interactions,
respectively.

Fig. 4. Interpretations of interactions in (a-c) additive and (d-f) averaging models. (a, d) Two-gene
interactions when both 1-gene effects A and B are positive. Two different cases (cases 1 and 2) of the *AB*
phenotype values are used. (b, e) Two-gene interactions when the 1-gene effects have opposite signs.
(c, f) Three-gene interactions. ":" and ";" indicate additive and averaging interactions, respectively.

Fig. 5. Coefficient estimates for the contribution to immunity from averaging model analysis of the data
in Tsuda et al. (a) AvrRpt2-ETI, (b) AvrRpm1-ETI, (c) flg22-PTI, and (d) elf18-PTI. The 95% confidence
intervals are shown as horizontal bars, with the means as points. The Holm-corrected p-values are
shown in the left part of each plot: red, $p < 0.05$; blue, $p \geq 0.05$. The dataset used for AvrRpt2-ETI in Fig.
5a is the same as that in used in Fig. 3, and Fig. 5a is the same as the full model (black lines) in Fig. 3c. ";"
indicates the averaging interaction.
533
534
535   **SUPPLEMENTAL TEXTS**
536   Text S1. Derivation of the averaging model and non-distributivity of the data
537
538   Text S2. Averaging and additive models without interaction are two extreme approximations of the
539   behavior of a two-input chemical signaling system
540
541
542   **SUPPLEMENTAL DATASETS**
543   Supplemental Datasets are available from https://github.com/fumikatagiri/Averaging_Model.
544
Supplemental Dataset 1. A folder containing four bacterial count data files (tab-delimited text) for
"AvrRpt2_ETI", "AvrRpm1_ETI", "flg22_PTI", and "elf18_PTI". Each has columns of genotype, treatment,
replicate, flat, pot, and colony. The colony column has log10-transformed colony counts (colony forming
unit/cm2). Although the summarized data were originally reported in (2), these raw data were not
published.
Supplemental Dataset 2. A .RData file (R workspace file) containing a list object, "ave.model.mats",
which contains the matrices for the averaging model for 2- to 7-gene systems (equivalents of matrix in
Fig. TS1.1 in Text S1 for different number-gene systems).
Supplemental Dataset 3. An R script file (.r file), which was used to generate Fig. 5 from the data in
Supplemental Dataset 1 (the data should be in the subfolder "Supplemental.Dataset1" as in this folder).
In addition, an empty "outputs" subfolder is required for the outputs of this script (the "outputs"
subfolder in this folder is already populated with the actual outputs of the script). The outputs of the
script includes "reanalyzed.w.average.model....pdf", which is Fig. 5. The script includes algorithms for
selecting significant genes for the analysis and the averaging model. In the script, the object of a 3-gene
or 4-gene matrix included in Supplemental Dataset 2 is called "rec.mx", which is generated by a
function, "make.rec.mx".
561

Text S1. Derivation of the averaging model and non-distributivity of the data

Laws of algebra

An additive model of gene effects and interactions involves two operators: additive, "+", and interactive,
":". Different models can be derived if we assume different laws for these operations. Three types of
laws define algebra involving two operators: commutative, associative, and distributive laws. The
commutative laws are A + B = B + A and A:B = B:A. The associative laws are (A + B) + C = A + (B + C) and
(A:B):C = A:(B:C). The distributive law is (A + B):C = A:C + B:C.

I assume the commutative laws for both "+" and ":" because a single quantitative phenotype cannot
experimentally distinguish A + B from B + A or A:B from B:A. I also assume the associative law for "+"
since without this assumption the general linear model framework cannot be used.

The associative law for ":" is also required for the general linear model framework (see below).
However, as I show below, the impact of a violation of the associative law for ":" can be moderated
using an averaging principle, in which the arithmetic mean of multiple different expressions for the same
quantity is taken as the true value of the quantity. This moderation by the averaging principle is
important in applications of the general linear model framework to multi-gene systems because we
cannot generally assume that the associative law for ":" holds. For example, in the case of Fig. 1 in the
main text, A:B = 0, and thus, (A:B):C = 0. However, B:C ≠ 0, so, A:(B:C) may not be 0 particularly when A:C
≠ 0. Thus, (A:B):C ≠ A:(B:C), i.e., a violation of the associative law, could happen.

In the following sections, I will show that the distributive law is required in the additive model. I will also
show that a range-limiting nonlinearity of a system, such as a saturation response, would violate the
distributive law. Such responses are common in biological systems. I will also show that there is a
general linear model that allows violation of the distributive law under the assumption of the averaging
principle. I call this model an averaging model.

Derivation of the averaging model

The part of the following discussion describing derivation of the NR model was modified from Text S1 in
(1). In this section, for simplicity, the intercept value (i.e., the phenotype value for the most disrupted
state) is subtracted from all the measured values so that the intercept value is 0.

According to the additive model, I assume *AB* = A + B + A:B ... (1) as the starting point. In this case, the
interaction is the deviation of the corresponding genotype from arithmetic addition of the 1-gene
effects. Let's extend this to a system consisting of three genes *A*, *B*, and *C*. The phenotype *ABC* can be
considered as being expressed in three different ways: adding C to the genetic background of *AB*; adding
A to the genetic background of *BC*; or adding B to the genetic background of *CA*.
By adding C to *AB*, the genotype *ABC* is expressed as:

ABC = *AB* + C + *AB*:C = (A + B + A:B) + C + *AB*:C = A + B + C + A:B + *AB*:C … (2)

If the distributive law is not assumed, (2) cannot be simplified.

If the distributive law is assumed, (2) can be simplified to:

ABC = A + B + C + A:B + (A + B + A:B):C = A + B + C + A:B + B:C + C:A + (A:B):C   … (3)

Similarly, if the distributive law is not assumed:

By adding A to *BC*, *ABC* = A + B + C + B:C + *BC*:A   … (4)

By adding B to *CA*, *ABC* = A + B + C + C:A + *CA*:B   … (5)

If the distributive law is assumed:

By adding A to *BC*, *ABC* = A + B + C + A:B + B:C + C:A + A:(B:C) … (6)

By adding B to *CA*, *ABC* = A + B + C + A:B + B:C + C:A + B:(C:A) … (7)

Although the expressions are not the same, (3), (6), and (7) must be the same in a model to explain *ABC*.

Therefore, for this model framework to work exactly, the associative law for the interaction operator ":"

is necessary,

(A:B):C = A:(B:C) = B:(C:A) = A:B:C … (8)

If (8) is true, (3), (6), and (7) become the same expression:

ABC = A + B + C + A:B + B:C + C:A + A:B:C … (9)

(9) is the additive model for three genes. This can be extended to a system consisting of more genes. In

summary, the additive model is a good description of a multi-gene system if the associative law for ":"

and the distributive law are conserved.

However, as discussed above, the associative law cannot be generally assumed for the ":" operator in a

multi-gene system. This contradiction about associativity indicates a failure of the general linear model

as a general description of a multi-gene system. A compromise to maintain the general linear model

framework is to define *ABC* as the arithmetic mean of (3), (6), and (7):

ABC = [{A + B + C + A:B + B:C + C:A + (A:B):C} + {A + B + C + A:B + B:C + C:A + A:(B:C)}

630   + { A + B + C + A:B + B:C + C:A + B:(C:A)}] / 3

= A + B + C + A:B + B:C + C:A + {(A:B):C + A:(B:C) + B:(C:A)} / 3 … (10)

I call this practical approach to avoiding the contradiction in the general linear model by averaging all

possible cases the averaging principle.

Since {(A:B):C + A:(B:C) + B:(C:A)} cannot be expressed by the lower order terms, A, B, C, A:B, B:C, and

C:A, it is reasonable to define A:B:C = {(A:B):C + A:(B:C) + B:(C:A)} / 3 … (11). Then,

ABC = A + B + C + A:B + B:C + C:A + A:B:C … (9)

Thus, with the averaging principle, the additive model can conform to the assumption of no associativity

in the interaction operator ":". This can be extended to a system consisting of more genes. In summary,

the additive model should be a reasonable description of a multi-gene system if the distributive law

holds.

If the distributive law cannot be assumed, (2), (4), and (5) must still be the same to express *ABC*. Here

again we observe a failure of the general linear model as a general description of a multi-gene system. I

apply the averaging principle to (2), (4), and (5) to express *ABC*:

ABC = [{A + B + C + A:B + *AB*:C} + {A + B + C + B:C + *BC*:A} + {A + B + C + C:A + *CA*:B}] / 3

= A + B + C + (A:B + B:C + C:A) / 3 + (AB:C + BC:A + CA:B) / 3 … (12)
Since (AB:C + BC:A + CA:B) cannot be expressed by the lower order terms, A, B, C, A:B, B:C, and C:A, it is
reasonable to define A;B;C = (AB:C + BC:A + CA:B) / 3 … (13). I use the semicolon ";" to distinguish this
different definition of interaction from that of the interaction in the additive model and call ";" the
averaging interaction operator and ":" the additive interaction operator.
ABC = A + B + C + (A:B + B:C + C:A) / 3 + A;B;C … (14)
Therefore, if the distributive law is not assumed, the average of the 2-gene additive interactions should
be used to express the all wild-type allele state of ABC. In general, the rule that the terms in each order
of the interactions (2-gene additive interactions, 3-gene averaging interactions, 4-gene averaging
interactions, …) must be averaged can be derived by extending this to a system with more genes.
For example, with a system consisting of 4 genes, A, B, C, and D:
$ABCD$ = A + B + C + D + (A:B + A:C + A:D + B:C + B:D + C:D) / 6 + (A;B;C + A;B;D + A;C;D + B;C;D) /4 +
A;B;C;D … (15)
This is the NR model (2) (previously called the signaling allocation model (1)). Note that in the NR model,
2-gene interactions are additive interactions while 3- or higher-order interactions are averaging
interactions.
662
The assumption of non-distributivity does not require any more changes in the model. However, the
above derivation of the NR model began with an arbitrary definition of the 2-gene additive interaction,
AB = A + B + A:B … (1), which is the reason the NR model is a mixture of additive and averaging
interactions. The model would be more mathematically consistent if the averaging interaction definition
were extended to 1-gene effect terms to make the 2-gene interactions averaging interactions as well,
i.e., AB = (A + B) / 2 + A;B … (16). I demonstrate in Figs. 2 and 3 in the main text that (16) is indeed
required for mathematical consistency of the model.
670
By applying (16):
ABC = (A + B + C) / 3 + (A;B + B;C + C;A) / 3 + A;B;C … (17)
$ABCD$ = (A + B + C + D) / 4 + (A;B + A;C + A;D + B;C + B;D + C;D) / 6 + (A;B;C + A;B;D + A;C;D + B;C;D) /4 +
A;B;C;D … (18)
Now the rule is that the terms in each order of the interactions, including 1-gene effects (the first order),
must be averaged. (17) and (18) are equivalents of:
ABC = (AB + BC + CA) / 3 + A;B;C … (19)
A;B;C = ABC - (AB + BC + CA) / 3 … (19)' (variated from (19))
$ABCD$ = (ABC + ABD + ACD + BCD) /4 + A;B;C;D … (20)
A;B;C;D = $ABCD$ - (ABC + ABD + ACD + BCD) /4 …(20)' (variated from (20))
From (19)' and (20)', the highest order averaging interaction is defined as the deviation of the
corresponding genotype from the average of all genotypes with one gene fewer. This definition of the
averaging interaction is highly interpretable. I call this extended model with all averaging interactions an
averaging model. With the definitions of the averaging interactions of different orders in (16), (19), and
(20), it is clear that the averaging model does not require the distributive law because these definitions
do not include any terms that could be affected by whether the distributive law holds or not.
687

Note that the mean estimates from the additive model, NR model, averaging model, and 1-way ANOVA
for all genotypes are just different ways to linearly decompose the phenotype values (when the full
model terms are kept). Thus, when the models are fit to actual data with replication, all these models
yield the same fitted and residual values. The numbers of estimated values are the same, i.e., the
models have the same residual degree of freedom. Therefore, I sometimes use only the mean estimates
of the models for my arguments. The coefficient matrices to solve the linear equations for the means in
the three models using the genotype mean values in a 3-gene system are shown in Fig. TS1.1.
695

a Additive model

	Intercept	A	B	C	A:B	A:C	B:C	A:B:C
triple.mut	1	0	0	0	0	0	0	0
A	1	1	0	0	0	0	0	0
B	1	0	1	0	0	0	0	0
C	1	0	0	1	0	0	0	0
AB	1	1	1	0	1	0	0	0
AC	1	1	0	1	0	1	0	0
BC	1	0	1	1	0	0	1	0
ABC	1	1	1	1	1	1	1	1

b NR model

	Intercept	A	B	C	A:B	A:C	B:C	A;B;C
triple.mut	1	0	0	0	0	0	0	0
A	1	1	0	0	0	0	0	0
B	1	0	1	0	0	0	0	0
C	1	0	0	1	0	0	0	0
AB	1	1	1	0	1	0	0	0
AC	1	1	0	1	0	1	0	0
BC	1	0	1	1	0	0	1	0
ABC	1	1	1	1	1/3	1/3	1/3	1

c Averaging model

	Intercept	A	B	C	A;B	A;C	B;C	A;B;C
triple.mut	1	0	0	0	0	0	0	0
A	1	1	0	0	0	0	0	0
B	1	0	1	0	0	0	0	0
C	1	0	0	1	0	0	0	0
AB	1	1/2	1/2	0	1	0	0	0
AC	1	1/2	0	1/2	0	1	0	0
BC	1	0	1/2	1/2	0	0	1	0
ABC	1	1/3	1/3	1/3	1/3	1/3	1/3	1

Fig. TS1.1. Matrices for the linear equations to obtain model coefficients from the genotype values in a 3-gene system for (a) additive, (b) NR, and (c) averaging models. The rows are genotypes and the columns are model variables. ":" and ";" indicate the additive and averaging interactions, respectively.

696
697
698
699 *Violation of the distributive law is prevalent in multi-gene systems*
700
The averaging model does not assume the distributive law. Do we really need to consider non-
distributivity in a biological system? Let's consider a simple 3-gene system, in which nA and nB are input
nodes and nC is the output node (Fig. TS1.2). Mechanistically, signals from nA and nB are first summed,
and then modulated by a nonlinear function f_1 before the signal is output from nC. Thus,
(A + B):C = nC = f_1(nA + nB) … (21)
A:C + B:C = f_1 (nA) + f_1 (nB) … (22)
Therefore, if the distributive law holds,
f_1(nA + nB) = f_1 (nA) + f_1 (nB) … (23)
709
Let's make f_1 a Michaelis-Menten function for a saturating response (Fig. TS1.2b):
$f_1(x) = \dfrac{10}{1+\frac{7}{x}}$ … (24)
When nA = 5, nB = 2,
f_1(nA + nB) = f_1(5 + 2) = 5
f_1 (nA) + f_1 (nB) = f_1(5) + f_1(2) = 4.16… + 2.22… = 6.38…

Thus, $f_1(nA + nB) \neq f_1(nA) + f_1(nB)$ ... (25)

and the distributive law is violated.

Generally, nonlinearity in a system leads to violation of the distributive law.

Fig. TS1.2. Non-linearity in a system violates the distributive law. (a) a 3-gene system, in which signals from nA and nB feed into nC. The output of nC is defined as $f_1(nA + nB)$. (b) When $f_1(x) = \frac{10}{1+\frac{7}{x}}$, the input-output relationships at nC are shown. If the input is nA, the output is expressed as A:C in the additive model. This plot clearly shows that (A + B):C ≠ A:C + B:C (Y-axis values in orange and green, respectively), a violation of the distributive law.

A saturating response limits the output range. Without nonlinearity, the range of the system output is

not limited, and this is the condition the additive model requires. Thus, the additive model generally

cannot be used in a system consisting of multiple genes (when the model includes 3 -or-more-gene

interactions, strictly speaking) when the phenotype value range is limited compared to the ranges of the

gene effects and interactions.

References

1.   K. Tsuda, M. Sato, T. Stoddard, J. Glazebrook, F. Katagiri, Network properties of robust immunity
in plants. *PLoS Genet* **5**, e1000772 (2009).
2.   F. Katagiri, Network Reconstitution for Quantitative Subnetwork Interaction Analysis. *Methods*
Mol Biol **1578**, 223-231 (2017).

Text S2. Averaging and additive models without interaction are two extreme approximations of
the behavior of a two-input chemical signaling system

We consider a hypothetical system with two converging input signals. One input is signal molecule A,
the other input is signal molecule B, and the output is signal molecule C (Fig. S2.1).

Fig. S2.1. The structure of a two-input system.

The system is modeled assuming an equilibrium among the three signal molecules. It can also be
considered as a steady state of a dynamical system, which is more realistic for a signaling system. The
system can be considered as a system of chemical reactions (e.g., A + X → C and B + Y → C), in which
other molecules in the reactions (like X and Y) are not limiting (i.e., [X], [Y] >> [A], [B], [C]). We can only
observe the final output of [C], the concentration of C, under the conditions of presence or absence of A
and B. We cannot observe [A] and [B].

In the system, the sum of [A] and the fraction of [C] that derived from [A] is denoted as $[A]_0$, and the
sum of [B] and the fraction of [C] that derived from [B] is denoted as $[B]_0$. When the system lacks B
(genotype *Ab*), $[C]_A$, the concentration of C specific to this condition is expressed with $[A]_A$, the
concentration of A specific to this condition, and the conversion constant k_A as $[C]_A = k_A[A]_A$. Since $[C]_A =$
$[A]_0 − [A]_A$, by eliminating $[A]_A$,

$[C]_A = \{ k_A / (k_A + 1) \} [A]_0$ ... (1)

Similarly, when the system lacks A (genotype *aB*), with the conversion constant k_B,

$[C]_B = \{ k_B / (k_B + 1) \} [B]_0$ ... (2)

When the system has both A and B (genotype *AB*), with $[A]_{AB}$ and $[B]_{AB}$, the concentration of A and B
specific to this condition, $[C]_{AB}$, the concentration of C specific to this condition, can be expressed as:

$[C]_{AB} = k_A[A]_{AB} = k_B[B]_{AB}$ ... (3)

Also, $[C]_{AB} = ([A]_0 − [A]_{AB}) + ([B]_0 − [B]_{AB})$ ... (4)

By eliminating $[A]_{AB}$ and $[B]_{AB}$ from (3) and (4),

$[C]_{AB} = \{ 1 / (1 / k_A + 1/ k_B + 1) \} ([A]_0 + [B]_0)$ ... (5)

We now consider the situations with different k_A and k_B values.

I. When $0 < k_A, k_B << 1$,

(1), (2), and (5) can be approximated as:

$[C]_A \approx k_A[A]_0$ … (6)

$[C]_B \approx k_B[B]_0$ … (7)

$[C]_{AB} \approx \{ 1 / (1 / k_A + 1 / k_B) \} ([A]_0 + [B]_0)$ … (8)

k_A and $[A]_0$ cannot be distinguished as we only know $[C]_A$ as the phenotype, and similarly, k_B and $[B]_0$
cannot be distinguished as we only know $[C]_B$ as the phenotype. Thus, we assume $k \approx k_A \approx k_B$ to remove
unnecessary complexity from the model. Then,

$[C]_A \approx k[A]_0$ … (6)′

$[C]_B \approx k[B]_0$ … (7)′

$[C]_{AB} \approx k ([A]_0 + [B]_0) / 2$ … (8)′

From (6)′, (7)′, and (8)′,

$[C]_{AB} \approx ([C]_A + [C]_B) / 2$ … (9)

Therefore, under the conditions of $0 < k_A$, $k_B \ll 1$ (meaning [A],[B] $\gg$ [C]), the averaging model without
interaction describes the system well.

II. When k_A, $k_B \gg 1$,

(1), (2), and (5) can be approximated as:

$[C]_A \approx [A]_0$ … (10)

$[C]_B \approx [B]_0$ … (11)

$[C]_{AB} \approx [A]_0 + [B]_0$ … (12)

From (10), (11), and (12),

$[C]_{AB} \approx [C]_A + [C]_B$ … (13)

Therefore, under the conditions of k_A, $k_B \gg 1$ (meaning [A],[B] $\ll$ [C]), the additive model without
interaction describes the system well.

The discussion in this supplemental text demonstrates that whether the averaging model or the additive
model describes the no interaction state better in a two-input system depends on the system. Since we
can only observe the output of the system, [C], (but not [A] or [B]), we cannot determine which of the
averaging and additive models would be better for a particular system.