# Estimation of the firing behaviour of a complete motoneuron pool by combining electromyography signal decomposition and realistic motoneuron modelling

**Abbreviated title:** Estimation of the firing behaviour of a complete motoneuron pool

Arnault H Caillet[1], Andrew T.M. Phillips[1], Dario Farina[2],*, Luca Modenese[1,3]*

[1]Department of Civil and Environmental Engineering, Imperial College London, SW7 2AZ, UK
[2]Department of Bioengineering, Imperial College London, SW7 2AZ, UK
[3]Graduate School of Biomedical Engineering, University of New South Wales, Sydney 2052, Australia

Email: arnault.caillet17@imperial.ac.uk

* These authors have equal contributions and share the senior authorship

NUMBER OF PAGES: 54

NUMBER OF FIGURES: 14

NUMBER OF TABLES: 3

CONFLICT OF INTEREST STATEMENT: The authors declare no competing financial interests.

ACKNOWLEDGEMENTS: AC was supported by the Skempton Scholarship and LM by an Imperial College Research Fellowship granted by Imperial College London.

# Abstract

Our understanding of the firing behaviour of motoneuron (MN) pools during human voluntary muscle contractions is currently limited to electrophysiological findings from animal experiments extrapolated to humans, mathematical models of MN pools not validated for human data, and experimental results obtained from decomposition of electromyographical (EMG) signals. These approaches are limited in accuracy or provide information on only small partitions of the MN population. Here, we propose a method based on the combination of high-density EMG (HDEMG) data and realistic modelling for predicting the behaviour of entire pools of motoneurons in humans. The method builds on a physiologically realistic model of a MN pool which predicts, from the experimental spike trains of a smaller number of individual MNs identified from decomposed HDEMG signals, the unknown recruitment and firing activity of the remaining unidentified MNs in the complete MN pool. The MN pool model is described as a cohort of single-compartment leaky fire-and-integrate (LIF) models of MNs scaled by a physiologically realistic distribution of MN electrophysiological properties and driven by a spinal synaptic input, both derived from decomposed HDEMG data. The MN spike trains and effective neural drive to muscle, predicted with this method, have been successfully validated experimentally. A representative application of the method in MN-driven neuromuscular modelling is also presented. The proposed approach provides a validated tool for neuroscientists, experimentalists, and modelers to infer the firing activity of MNs that cannot be observed experimentally, investigate the neuromechanics of human MN pools, support future experimental investigations, and advance neuromuscular modelling for investigating the neural strategies controlling human voluntary contractions.

# Author Summary

Our experimental understanding of the firing behaviour of motoneuron (MN) pools during human voluntary muscle contractions is currently limited to the observation of small samples of active MNs obtained from EMG decomposition. EMG decomposition therefore provides an important but incomplete description of the role of individual MNs in the firing activity of the complete MN pool, which limits our understanding of the neural strategies of the whole MN pool and of how the firing activity of each MN contributes to the neural drive to muscle. Here, we combine decomposed high-density EMG (HDEMG) data and a physiologically realistic model of MN population to predict the unknown recruitment and firing activity of the remaining unidentified MNs in the complete MN pool.

33    In brief, an experimental estimation of the synaptic current is input to a cohort of MN models, which

34    are calibrated using the available decomposed HDEMG data, and predict the MN spike trains fired by

35    the entire MN population. This novel approach is experimentally validated and applied to muscle force

36    prediction from neuromuscular modelling.

# INTRODUCTION

During voluntary muscle contractions, pools of spinal alpha-motoneurons (MNs) convert the synaptic input they receive into a neural command that drives the contractile activity of the innervated muscle fibres, determining limb motion. Identifying the recruitment and firing dynamics of MNs is fundamental for understanding the neural strategies controlling human voluntary motion, with applications in sport sciences [1-3], and neurological and musculoskeletal rehabilitation [4-8]. Determining the MN-specific contributions to the MN population activity also allows more realistic control of neuromuscular models [9-12], investigation of muscle neuromechanics [13,14], prediction of limb motion from MN-specific behaviour [15], or improvement in human-machine interfacing and neuroprosthetics [16,17].

Our understanding of MN pool firing behaviour during human voluntary tasks is however currently limited. While the MN membrane afterhyperpolarization and axonal conduction velocity can be inferred from indirect specialized techniques [18,19], most of the other electro-chemical MN membrane properties and mechanisms that define the MN recruitment and discharge behaviour cannot be directly observed in humans in vivo. Analysis of commonly adopted bipolar surface EMG recordings, which often lump the motor unit (MU) trains of action potentials into a single signal assimilated as the neural drive to muscle, cannot advance our understanding of the MN pool activity at the level of single MNs. Our experimental knowledge on the remaining MN membrane properties in mammals is therefore obtained from in vitro and in situ experiments on animals [20]. The scalability of these mechanisms to humans is debated [21] due to a systematic inter-species variance in the MN electrophysiological properties in mammals [22]. Decomposition of high-density EMG (HDEMG) or intramuscular EMG (iEMG) signals [23,24] allows the in vivo decoding in human muscles of the firing activity of individual active motoneurons during voluntary contractions and provide a direct window on the internal dynamics of MN pools. Specifically, the non-invasive EMG approach to MN decoding has recently advanced our physiological understanding of the neurophysiology of human MU pools and of the interplay between the central nervous system and the muscle contractile machinery [25,26]

Yet, the activity of all the MNs constituting the complete innervating MN population of a muscle cannot be identified with this technique. High-yield decomposition typically detects at most 30-40 MNs [27], while MU pools typically contain hundreds of MUs in muscles of the hindlimb [20]. The small sample of recorded MNs is besides not representative of the continuous distribution of the MN electrophysiological properties in the complete MN pool with a bias towards the identification of mainly high-threshold MUs. The samples of spike trains obtained from signal decomposition therefore

provide a limited description of the role of individual MNs in the firing activity of the complete MN
pool.

To allow the investigation and description of specific neurophysiological mechanisms of the complete
MN population, some studies have developed mathematical frameworks and computational models
of pools of individual MNs. These MN pool models have provided relevant insights for interpreting
experimental data [28,29], investigating the MN pool properties and neuromechanics [30-32],
neuromuscular mechanisms [12,33], and the interplay between muscle machinery and spinal inputs
[34]. However, none of these MN pool models have been tested with experimental input data, instead
either receiving artificial gaussian noise [32], sinusoidal [31] or ramp [29] inputs, inputs from
interneurons [30] or feedback systems [28]. These MN pool models have therefore never been tested
in real conditions of voluntary muscle contraction. The forward predictions of MN spike trains or neural
drive to muscle obtained in these studies were consequently not or indirectly validated against
experimental recordings.

The MN-specific recruitment and firing dynamics of these MN pool models are usually described with
comprehensive or phenomenological models of MNs. The biophysical approaches [30-33], which rely
on a population of compartmental Hodgkin-Huxley-type MN models provide a comprehensive
description of the microscopic MN-specific membrane mechanisms of the MN pool and can capture
complex nonlinear MN dynamics [35]. However, these models are computationally expensive and
remain generic, involving numerous electrophysiological channel-related parameters for which
adequate values are difficult to obtain in mammalian experiments [36,37] and must be indirectly
calibrated or extrapolated from animal measurements in human models [38]. On the other hand,
phenomenological models of MNs [10,28,39] provide a simpler description of the MN pool dynamics
and rely on a few parameters that can be calibrated or inferred in mammals including humans. They
are inspired from the Fuglevand's formalism [29], where the output MN firing frequency is the
gaussian-randomized linear response to the synaptic drive with a MN-specific gain. However, these
phenomenological models cannot account for the MN-specific nonlinear mechanisms that dominate
the MN pool behaviour [20,35,40,41]. MN leaky integrate-and-fire (LIF) models, the parameters of
which can be defined by MN membrane electrophysiological properties for which mathematical
relationships are available [42], are an acceptable trade-off between Hodgkin-Huxley-type and
Fuglevand-type MN models with intermediate computational cost and complexity, and accurate
descriptions of the MN macroscopic discharge behaviour [43] without detailing the MN's underlying
neurophysiology [44]. While repeatedly used for the modelling and the investigation of individual MN
neural dynamics [45-47], MN LIF models are however not commonly used for the description of MN
pools.

To the authors' knowledge there is no systematic method to record the firing activity of all the MNs in

a MN pool, or to estimate from a sample of experimental MN spike trains obtained from signal

decomposition the firing behaviour of MNs that are not recorded in the MN pool. There is no

mathematical model of a MN pool that (1) was tested with experimental neural inputs and investigated

the neuromechanics of voluntary human muscle contraction, (2) involves a cohort of MN models that

relies on MN-specific profiles of inter-related MN electrophysiological properties, (3) is described by a

physiologically-realistic distribution of MN properties that is consistent with available experimental

data. This limits our understanding of the neural strategies of the whole MN pool and of how the firing

activity of each MN contributes to the neural drive to muscle.

In this study, a novel four-step approach is designed to predict, from the neural information of $N_r$ MN

spike trains obtained from HDEMG signal decomposition, the recruitment and firing dynamics of the

$N - N_r$ MNs that were not identified experimentally in the investigated pool of $N$ MNs. The model of

the MN pool was built upon a cohort of $N$ single-compartment LIF models of MNs. The LIF parameters

are derived from the available HDEMG data, are MN-specific, and account for the inter-relations

existing between mammalian MN properties [42]. The distribution of $N$ MN input resistances hence

obtained defines the recruitment dynamics of the MN pool. The MN pool model is driven by a common

synaptic current, which is estimated from the available experimental data as the filtered cumulative

summation of the $N_r$ identified spike trains. The MN-specific LIF models phenomenologically transform

this synaptic current into accurate discharge patterns for the $N_r$ MNs experimentally identified and

predict the MN firing dynamics of the $N - N_r$ unidentified MNs. The blind predictions of the spike

trains of the $N_r$ identified MNs and the effective neural drive to the muscle, computed from the firing

activity of the complete pool of $N$ virtual MNs, are both successfully validated against available

experimental data.

Neuroscientists can benefit from this proposed approach for inferring the neural activity of MNs that

cannot be observed experimentally and for investigating the neuromechanics of MN populations.

Experimenters can use this approach for better understanding their experimental dataset of $N_r$

identified discharging MNs. Moreover, this approach can be used by modelers to design and control

realistic neuromuscular models, useful for investigating the neural strategies in muscle voluntary

contractions. In this study, we provide an example for this application by using the simulated discharge

patterns of the complete MN pool as inputs to Hill-type models of muscle units to predict muscle force.

# METHODS

**<u>Overall approach</u>**

The spike trains $sp_j^{sim}(t)$ elicited by the entire pool of $N$ MNs were inferred from a sample of $N_r$ experimentally identified spike trains $sp_i^{exp}(t)$ with a 4-step approach displayed in Figure 1. The $N_r$ experimentally-identified MNs were allocated to the entire MN pool according to their recorded force recruitment thresholds $F_i^{th}$ (step 1). The common synaptic input to the MN pool was estimated from the experimental spike trains $sp_i^{exp}(t)$ of the $N_r$ MNs (step 2) and was linearly scaled for simplicity to the total postsynaptic membrane current $I(t)$ responsible for spike generation. A cohort of $N_r$ single-compartment leaky-and-fire (LIF) models of MNs, the electrophysiological parameters of which were mathematically determined by the unique MN size parameter $S_i$, transformed the input $I(t)$ to simulate the experimental spike trains $sp_i^{exp}(t)$ after calibration of the $S_i$ parameter (step 3). The distribution of the $N$ MN sizes $S(j)$ in the entire MN pool, which was extrapolated by regression from the $N_r$ calibrated $S_i$ quantities, scaled the electrophysiological parameters of a cohort of $N$ LIF models. The $N$ calibrated LIF models predicted from $I(t)$ the spike trains $sp_j^{sim}(t)$ of action potentials elicited by the entire pool of $N$ virtual MNs.

The following assumptions were made. (1) The MU pool is idealized as a collection of $N$ independent MUs that receive a common synaptic input and possibly MU-specific independent noise. (2) In a pool of $N$ MUs, $N$ MNs innervate $N$ muscle units (mUs). (3) In our notation, the pool of $N$ MUs is ranked from $j = 1$ to $j = N$, with increasing recruitment threshold. For the $N$ MNs to be recruited in increasing MN and mU size and recruitment thresholds according to Henneman's size principle [20,48-53], the distribution of morphometric, threshold and force properties in the MN pool follows

$$\forall j, k \in [1; N], (j < k) \Longrightarrow \left( S_j < S_k \Leftrightarrow I_j^{th} < I_k^{th} \Leftrightarrow IR_j < IR_k \Leftrightarrow F_j^{th} < F_k^{th} \Leftrightarrow f_{iso,j}^{max} < f_{iso,k}^{max} \right)$$

where $S$ is the MN surface area, $I^{th}$ the MN current threshold for recruitment, $IR$ the MU innervation ratio defining the MU size, $F^{th}$ is the MU force recruitment threshold, and $f_{iso}^{max}$ is the MU maximum isometric force. (4) The MN-specific electrophysiological properties are mathematically defined by the MN size $S$ [42]. This extends the Henneman's size principle to:

$$\forall j, k \in [1; N], (j < k) \Longrightarrow \left( S_j < S_k \Leftrightarrow C_j < C_k \Leftrightarrow R_j > R_k \Leftrightarrow \tau_j > \tau_k \right)$$

174 Where $C$ is the MN membrane capacitance, $R$ the MN input resistance and $\tau$ the MN membrane time
175 constant.

**Experimental data**

The four sets of experimental data used in this study, named as reported in the first column of Table
1, provide the time-histories of recorded MN spike trains $sp_i^{exp}(t)$ and whole muscle force trace $F(t)$
(left panel in Figure 1), and were obtained from the studies [54,55] and [26,27], as open-source
supplementary material and personal communication, respectively. In these studies, the HDEMG
signals were recorded with a sampling rate $f_s = 2048 Hz$ from the Tibialis Anterior (TA) and
Gastrocnemius Medialis (GM) human muscles during trapezoidal isometric contractions. As displayed
in Figure 2, the trapezoidal force trajectories are described in this study by the $t_{tr_{0 \to 5}}$ times reported in
Table 1, as a zero force in $[t_{tr0}; t_{tr1}]$, a ramp of linearly increasing force in $[t_{tr1}; t_{tr2}]$, a plateau of
constant force in $[t_{tr2}; t_{tr3}]$, a ramp of linearly decreasing force in $[t_{tr3}; t_{tr4}]$, and a zero force in
$[t_{tr4}; t_{tr5}]$.

*Figure 2: Definition of times $t_{tr_i}$ that describe the trapezoidal shape of the muscle isometric contraction.*

*Table 1:The four experimental datasets processed in this study. $N_r$ spike trains are identified per dataset during trapezoidal*
*contractions of the Tibialis Anterior (TA) or Gastrocnemius Medialis (GM) muscles. The trapezoidal force trace is described*
*by times $t_{tr_i}$ in seconds up to a dataset-specific level of maximum voluntary contraction (%MVC).*

| Dataset | Muscle | %MVC | $t_{tr0}$ | $t_{tr1}$ | $t_{tr2}$ | $t_{tr3}$ | $t_{tr4}$ | $t_{tr5}$ | $N_r$ | Reference paper |
|---------|--------|------|------|------|------|------|------|------|------|-----------------|
| $DTA_{35}$ | TA | 35 | 0 | 2.2 | 10.6 | 20.5 | 30 | 30 | 32 | [26,27] |
| $HTA_{35}$ | TA | 35 | 0 | 2.1 | 10.5 | 20.5 | 30 | 33 | 21 | |
| $HTA_{50}$ | TA | 50 | 0 | 1.6 | 12 | 21.8 | 34.5 | 35 | 14 | [54,55] |
| $HGM_{30}$ | GM | 30 | 0 | 3.1 | 9.1 | 28 | 33.5 | 107 | 27 | |

193 The HDEMG signals were decomposed with blind-source separation techniques and $N_r$ MN spike trains
194 $sp_i^{exp}(t)$ were identified. In this study, the experimental $N_r$ MNs were ranked from $i = 1$ to $i = N_r$ in
195 the order of increasing recorded force recruitment thresholds $F_i^{th}$, i.e. $\forall i \in [1; N_r], F_i^{th} < F_{i+1}^{th}$. The

196  sample time of the k$^{th}$ firing event of the i$^{th}$ identified MN is noted as $ft_i^k$, and the binary spike train of

197  the i$^{th}$ identified MN was mathematically defined as:

$$\begin{cases} sp_i^{exp}(t = ft_i^k) = 1 \\ sp_i^{exp}(t \neq ft_i^k) = 0 \end{cases} \qquad \text{(EQ. 1)}$$

198  The train of instantaneous discharge frequency $IDF_i(t)$ of the i$^{th}$ identified MN was computed

199  between firing times $ft_i^k$ and $ft_i^{k+1}$ as:

$$\begin{cases} IDF_i(t = ft_i^k) = \dfrac{1}{ft_i^{k+1} - ft_i^k} \\ \quad IDF_i(t \neq ft_i) = 0 \end{cases} \qquad \text{(EQ. 2)}$$

200  The IDFs were moving-average filtered by convolution with a Hanning window of length 400ms [56],

201  yielding the continuous filtered instantaneous discharge frequencies (FIDFs) for all $N_r$ identified MNs.

### Approximation of the TA and GM MU pool size

The typical number $N$ of MUs was estimated for the TA muscle from cadaveric studies [57], statistical

methods [58], decomposed-enhanced spike-triggered-averaging (DESTA) approaches [59-64], and

adapted multiple point stimulation methods [65] in 20-80-year-old human subjects. Because of

method-specific limitations [66], results across methods varied substantially, with estimates for $N$ of,

respectively, 445, 194, 190, 188 and 300 MUs for the TA muscle. DESTA methods systematically

underestimate the innervation ratio due to the limited muscle volume covered by the surface

electrodes. Cadaveric approaches rely on samples of small size and arbitrarily distinguish alpha from

gamma MNs. Twitch torque measurements are an indirect method for estimating $N$. Accounting for

these limitations, we estimated the potentially conservative $N_{TA} = 400$ MUs in a typical adult TA

muscle. Assuming 200,000 fibres in the TA muscle [67,68], $N_{TA} = 400$ yields a mean of 500 fibres per

TA MU, consistently with previous findings [68]. In two cadaveric studies [57,69], the estimate for the

GM was $N_{GM} = 550$ MUs, which is consistent with $N_{TA} = 400$ as the GM muscle volume is typically

larger than TA's [70]. A sensitivity analysis on the performance of the method presented in Figure 1

upon variations on the value of $N_{TA}$ was included in this study.

### Step (1): MN mapping

In the first step of the approach overviewed in Figure 1, the $N_r$ experimentally identified MNs were

allocated to the entire pool of $N$ MNs according to their recorded force recruitment thresholds $F_i^{th}$.

Three studies measured in the human TA muscle *in vivo* the force recruitment thresholds $F^{th}$ of MUs,

given as a percentage of the maximum voluntary contraction (%MVC) force, for 528 [64], 256 [71], and

302 [72] MUs. Other studies investigated TA MU pools but reported small population sizes [73] and/or

did not report the recruitment thresholds [74-76].

We digitized the scatter plot in Figure 3 in [64] using the online tool WebPlotDigitizer [77]. The

normalized MU population was partitioned into 10%-ranges of the values of $F^{th}$ (in %MVC), as

reported in [71,72]. The distributions obtained from these three studies were averaged. The

normalized frequency distribution by 10%MVC-ranges of the $F^{th}$ quantities hence obtained was

mapped to a pool of $N$ MUs, providing a step function relating each $j^{th}$ MU in the MU population to

its 10%-range in $F^{th}$. This step function was least-squares fitted by a linear-exponential trendline (EQ.

3) providing a continuous frequency distribution of TA MU recruitment thresholds in a MU pool that

reproduces the available literature data.

$$F^{th}(j) = k_1 \cdot \left( k_2 \cdot \frac{j}{N} + \Delta_{\mathrm{F}}^{\left(\frac{j}{N}\right)^{k_3}} \right), j \in [\![1; N]\!] \qquad \text{(EQ. 3)}$$

Simpler trendlines, such as $F^{th}(j) = k_1 \cdot \Delta_{\mathrm{F}}^{\left(\frac{j}{N}\right)^{k_3}}$ [29], returned fits of lower $r^2$ values. According to the

three studies and to [20], a $\Delta_{\mathrm{F}} = 120$-fold range in $F^{th}$ was set for the TA muscle, yielding $F^{th}(N) =$

90%MVC $= \Delta_{\mathrm{F}} \cdot F^{th}(1)$, with $F^{th}(1) = 0.75$%MVC. Finally, the equation $F^{th}(N_i) = k_1 \cdot \left( k_2 \cdot \frac{N_i}{N} + \right.$

$\left. \Delta_{\mathrm{F}}^{\left(\frac{N_i}{N}\right)^{k_3}} \right) = F_i^{th}$ was solved for the variable $N_i$ for all $N_r$ identified MUs for which the experimental

threshold $F_i^{th}$ was recorded. The $N_r$ identified MUs were thus assigned the $N_i^{th}$ locations of the

complete pool of $N$ MUs ranked in order of increasing $F^{th}$:

239                    $i \in [\![1; N_r]\!] \rightarrow N_i \in [\![1; N]\!]$

When considering a 100-ms electromechanical delay between MN recruitment time and onset of

muscle unit force, the mapping $i \in [\![1; N_r]\!] \rightarrow N_i \in [\![1; N]\!]$ did not substantially change. It was

therefore simplified that a MN and its innervated muscle unit were recruited at the same time, and

the $i \in [\![1; N_r]\!] \rightarrow N_i \in [\![1; N]\!]$ mapping derived for muscle units was extrapolated to MNs.

Considering that typically less than 30 MUs (5% of the GM MU pool) can be currently identified by

HDEMG decomposition in GM muscles [27], and that the few papers identifying GM MUs with

intramuscular electrodes either did not report the MU $F^{th}$ [78-80] or identified less than 24 MUs up

to 100% MVC [81,82], a GM-specific $F^{th}(j)$ distribution could not be obtained from the literature for

the GM muscle. The $F^{th}(j)$ distribution obtained for the TA muscle was therefore used for the

simulations performed with the GM muscle, which is acceptable as an initial approximation based on

visual comparison to the scattered data provided in these studies.

**Step (2): Current input $I(t)$**

In the second step of the approach in Figure 1, the common synaptic input to the MN pool was first

estimated from the experimental data. The cumulative spike train (CST) was obtained as the temporal

binary summation of the $N_r$ experimental spike trains $sp_i^{exp}(t)$.

$$CST = \sum_i sp_i^{exp}(t) \tag{EQ. 4}$$

256  The effective neural drive ($eND$) to the muscle was estimated by low-pass filtering the CST in the

257  bandwidth $[0; 10]Hz$ relevant for force generation [47]. As the pool of MNs filters the MN-specific

258  independent synaptic noise received by the individual MNs and linearly transmits the common

259  synaptic input ($CSI$) to the MN pool [31,83], the $CSI$ was equalled to the $eND$ in arbitrary units:

$$CSI(t) = k \cdot eND(t) \tag{EQ. 5}$$

The common synaptic control ($CSC$) signal was obtained by low pass filtering the $CSI$ in $[0; 4]Hz$.

This approach, which estimates the $CSI$ from the $N_r$ experimental spike trains is only valid if the sample

of $N_r$ MNs is 'large enough' and 'representative enough' of the complete MN pool for the linearity

properties of the population of $N_r$ MNs to apply [83]. To assess if this approximation holds with the $N_r$

MNs obtained experimentally, the following two validations were performed. (1) The coherence

$coher_{\frac{N_r}{2}} \in [0; 1]$, averaged in $[1; 10]Hz$, was calculated between two cumulative spike trains $CST_{\frac{N_r}{2},1}$

and $CST_{\frac{N_r}{2},2}$ computed from two complementary random subsets of $\frac{N_r}{2}$ MNs. This was repeated 20

times for random permutations of complementary subsets of $\frac{N_r}{2}$ MNs, and the $coher_{\frac{N_r}{2}}$ values were

average yielding $coher_{\frac{N_r}{2},mean} \in [0; 1]$. The coherence $coher_{N_r}$ between the complete experimental

sample of $N_r$ MNs and a virtual sample of $N_r$ non-identified MNs was finally estimated by reporting

the pair $\left(\frac{N_r}{2}; \; coher_{\frac{N_r}{2},mean}\right)$ similarly to Figure 2A in [47]. (2) The time-histories of the normalized

force $\bar{F}(t)$ and common synaptic control $\overline{CSC}$, which should superimpose if the linearity properties

apply (see Figure 6 in [83]), were compared with calculation of the normalized root-mean-square error

(nRMSE) and coefficient of determination $r^2$. If $coher_{N_r} > 0.7$, $r^2 > 0.7$ and $nRMSE < 30\%$, it was

assumed that the sample of $N_r$ MNs was large and representative enough of the MN pool for the

linearity properties to apply, and the $eND$ was confidently assimilated as the $CSI$ to the MN pool in

arbitrary units. It must be noted that if $coher_{N_r} < 1$, the linearity properties do not fully apply for the

sample of $N_r$ MNs, and the $CSI$ computed from the $N_r$ MNs is expected to relate to the true $CSI$ with

a coherence close but less than $coher_{N_r}$.

The $CSI$ hence obtained reflects the net excitatory synaptic influx common to the MN population,

which triggers the opening of voltage-dependent dendritic channels and the activation of intrinsic

membrane currents, such as persistent inward currents (PICs), which contribute to the total dendritic

membrane current $I(t)$ responsible for spike generation. As the single-compartment LIF model

considered in this study does not describe the dendritic activity, the PIC-related nonlinearity in the

$CSI - I$ transformation was excluded from the model and $I(t)$ was simplified to be linearly related to

$CSI$ with a constant gain $G$ across the MN pool. To avoid confusion, $I(t)$ is referred as 'current input'

in the following. The limitations of this simplification are investigated in the Limitations Section

(Limitation 3). To identify $G$, it must be noted that the $CSI$, which was computed from a subset of the

MN pool, does not capture the firing activity of the MNs that are recruited before the smallest

identified $N_1^{th}$ MN, which is recruited at time $ft_{N_1}^1$. It non-physiologically yields $CSI\left(t < ft_{N_1}^1\right) = 0$.

Accounting for this experimental limitation, $I(t)$ was defined to remain null until the first identified

MN starts firing at $ft_{N_1}^1$, and to non-continuously reach $I_{N_1}^{th}$ at $t = ft_{N_1}^1$:

$$I(t) = \begin{cases} 0 \; if \; t < ft_{N_1}^1 \\ I_{N_1}^{th} + G \cdot CSI(t) \; else \end{cases} \quad \text{with} \quad G = \frac{I_{N_r}^{th} - I_{N_1}^{th}}{CSI\left(ft_{N_r}^1\right) - CSI\left(ft_{N_1}^1\right)} \quad \text{(EQ. 6)}$$

In (EQ. 6), the rheobase currents of the first and last identified MNs $I_{N_1}^{th}$ and $I_{N_r}^{th}$ were estimated from

a typical distribution of rheobase in a MN pool $I^{th}(j) = k_1 \cdot \Delta_I^{\left(\frac{j}{N}\right)^{k_3}}$, obtained as for the distribution of

$F^{th}$, from normalized experimental data from populations of hindlimb alpha-MNs in adult rats and cats

in vivo [84-92]. It is worth noting that the experimental $I^{th}$ were obtained by injecting current pulses

directly into the soma of identified MNs, thus bypassing the dendritic activity and most of the

aforementioned PIC-generated nonlinear mechanisms [40]. A $\Delta_I = 9.1$-fold range in MN rheobase in

$[3.9; 35.5]nA$ was taken [21,42], while a larger $\Delta_I$ is also consistent with the literature ([93], Table 4)

and larger values of $I^{th}$ can be expected for humans [21].

**Step (3): LIF model – parameter tuning and distribution of electrophysiological properties in the**

**MN pool**

In the third step of the approach in Figure 1, single-compartment LIF MN models with current synapses

are calibrated to mimic the discharge behaviour of the $N_r$ experimentally identified MNs.

 MN LIF model: description

A variant of the LIF MN model was chosen in this study for its relative mathematical simplicity and low

computational cost and its adequacy in mimicking the firing behaviour of MNs. The LIF model described

in (EQ. 7) describes the discharge behaviour of a MN of rheobase $I^{th}$ and input resistance $R$ as a

capacitor charging with a time constant $\tau$ and instantaneously discharging at time $ft$ when the

membrane voltage $V_m$ meets the voltage threshold $V_{th}$, after which $V_m$ is reset and maintained to the

membrane resting potential $V_{rest}$ for a time duration called 'inert period' $IP$ in this study. For simplicity

without loss of generalisation, the relative voltage threshold was defined as $\Delta V_{th} = V_{th} - V_{rest} > 0$,

and $V_{rest}$ was set to 0.

The model is described by the following set of equations:

$$
\begin{cases}
\tau \dfrac{dV_m}{dt} = R \cdot I(t) - V_m \\
\tau = RC \\
V_m(ft) = \Delta V_{th} \\
\lim_{t \to ft^+} V_m = 0 \\
\forall t \in [ft; ft + IP], V_m = 0
\end{cases}
\tag{EQ. 7}
$$

315 The differential equation was solved with a time step $dt \leq 0.001s$ as:

$$
\begin{cases}
V_m(0) = 0 \\
\forall n \in \mathbb{N}^*, V_m[nT] = e^{-\frac{T}{\tau}} V_m[(n-1)T] \; + \; R\dfrac{T}{\tau} \cdot I(nT)
\end{cases}
\tag{EQ. 8}
$$

This model includes 5 electrophysiological parameters: $R, C, \tau, \Delta V_{th}$ and $IP$. The passive parameters

$R$ and $C$ were mathematically related to the MN surface area S as $R = \dfrac{k_R}{S^{2.43}}$ and $C = C_m \cdot S$ after an

extensive meta-analysis of published experimental data on hindlimb alpha-MNs in adult cats in vivo

[42], that sets the constant value of $C_m$ to $1.3\mu F \cdot cm^2$, supports the equality $\tau = RC$ and the validity

of Ohm's law in MNs as $I^{th} = \dfrac{2.7 \cdot 10^{-2}}{R} = \dfrac{\Delta V_{th}}{R}$, setting the constant value $\Delta V_{th} = 27mV$ in this study.

The model was thus reduced to the MN size parameter $S$ and to the $IP$ parameter.

MN LIF model: the IP parameter

324 Single-compartment LIF models with no active conductances receiving current inputs can predict non-

325 physiologically large values of MN firing frequency (FF) because of a linear FF-I gain at large but

326 physiological current inputs $I(t)$ [44]. The saturation in the FF-I relation typically observed in the

327 mammalian motor unit firing patterns is primarily mediated by the voltage-dependent activity of the

328 PIC-generating dendritic channels [40], that was overlooked in the $CSI - I$ transformation in Step (2),

and which decreases the driving force of the synaptic current flow as the dendritic membrane depolarizes. While a physiological modelling of this saturation could be achieved with a LIF model with conductance synapses, as discussed in Limitation 3 in the Limitations section, this nonlinear behaviour could also be captured in a simple approach by tuning the phenomenological IP parameter in (EQ. 7). Considering a constant supra-threshold current $I \gg I^{th}$ input to a LIF MN, the steady-state firing frequency FF predicted by the LIF model is:

$$FF(I) = \frac{1}{IP - RC \ln\left(1 - \frac{\Delta V_{th}}{RI}\right)} \underset{I \gg I^{th}}{\Longleftrightarrow} FF(I) \approx \frac{1}{IP + \frac{C \cdot \Delta V_{th}}{I}} \qquad \text{(EQ. 9)}$$

As $I$ and $C$ typically vary over a 10-fold and 2.4-fold range respectively [42], the FF predicted by the LIF model is dominantly determined by the value of $IP$ as the input current increasingly overcomes the MN current threshold: $FF(I \gg I^{th}) \approx \frac{1}{IP}$.

In the previous phenomenological models of MNs [9,10,29], a maximum firing rate $FF^{max}$ was defined and a non-derivable transition from $FF(I)$ to constant $FF^{max}$ was set for increasing values of $I(t)$. Here, $IP$ was integrated to the dynamics of the LIF model and was derived from experimental data to be MN-specific in the following manner. For each of the $N_r$ identified MNs, the time-course of the MN instantaneous discharge frequencies (IDFs) was first smoothed, as performed in [55], with a sixth-order polynomial trendline ($IDF_{trend}$) to neglect any unexplained random noise in the analysis. The mean $M$ of the trendline values during the plateau of force $[t_{r_2}; t_{r_3}]$ was then obtained. If $\exists t \in [t_{r_1}; t_{r_2} - 1], IDF_{trend}(t) > 0.9M$, i.e. if the MN reached during the ramp of $I(t)$ in $[t_{r_1}; t_{r_2} - 1]$ an IDF larger than 90% of the IDF reached one second before the plateau of force, the MN was identified to 'saturate'. Its $IP$ parameter was set to $IP = \frac{1}{\max(IDF_{trend})}$, which constrains the MN maximum firing frequency for high input currents to $FF^{max} \approx \frac{1}{IP} = max(IDF_{trend})$. A power trendline $IP(j) = a \cdot j^b$ was finally fitted to the pairs $\left(N_i; IP_{N_i}\right)$ of saturating MNs and the $IP_{N_i}$ values of the non-saturating MNs were predicted from this trendline. To account for the residual variation in FF observed to remain at high $I(t)$ due to random electrophysiological mechanisms, the $IP$ parameter was randomized at each firing time, taking the value $IP + o$, where $o$ was randomly obtained from a normal gaussian distribution of $\frac{IP}{10}$ standard deviation.

MN LIF model: MN size parameter calibration

The remaining unknown parameter - the MN size $S$ - defines the recruitment and firing dynamics of the LIF model. The size $S_i$ of the $i^{th}$ identified MN was calibrated by minimizing over the time range $\left[t_{tr_0}; \frac{t_{tr_2} + t_{tr_3}}{2}\right]$ (Table 1) the cost function $J(S_i)$ computed as the root-mean-square difference

between experimental $FIDF_i^{exp}(t)$ and LIF-predicted $FIDF_i^{sim}(t)$ filtered instantaneous discharge frequencies:

$$
\begin{cases}
J(S_i) = \sqrt{\dfrac{dt}{\dfrac{t_{tr_2}+t_{tr_3}}{2} - t_{r_0}} \cdot \sum_{t_k=t_{r_0}}^{\frac{t_{tr_2}+t_{tr_3}}{2}} \left( FIDF_i^{sim}(t_k) - FIDF_i^{exp}(t_k) \right)^2} \\
\min_{S_i} J(S_i), i \in [\![1;N_r]\!]
\end{cases}
\tag{EQ. 10}
$$

To assess how well the calibrated LIF models can replicate the available experimental data, the normalized RMS error (nRMSE) (%) and coefficient of determination $r^2$ between $FIDF_i^{exp}(t)$ and $FIDF_i^{sim}(t)$, and the error in seconds between experimental and predicted recruitment times $\Delta f t_i^1 = f t_i^{1,sim} - f t_i^{1,exp}$ were computed for the $N_r$ MNs. Finally, a power trendline in (EQ. 11) was fitted to the pairs $(N_i; S_{N_i})$, and the continuous distribution of MN sizes in the entire pool of $N$ MNs was obtained.

$$
S(j) = k_1 \cdot \Delta_S^{\left(\frac{j}{N}\right)^{k_3}}
\tag{EQ. 11}
$$

$\Delta_S = 2.4$ was taken in [42]. The $S(j)$ distribution defines the continuous distribution of the MN-specific electrophysiological properties across the MN pool ([42], Table 6).

<u>MN LIF model: parameter identification during the derecruitment phase</u>

The time-range $\left[t_{r_3}; t_{r_5}\right]$ over which the MNs are being derecruited was not considered in the $S$ calibration in (EQ. 10) because the MN's current-voltage relation presents a hysteresis triggered by long-lasting PICs, as discussed in [40,94,95]. This hysteresis, which leads MNs to be derecruited at lower current threshold than at recruitment, can be phenomenologically interpreted in the scope of a single-compartment LIF model with no active conductance as an increase in the 'apparent' MN resistance $R$ during derecruitment to a new value $R^d$. To determine $R^d$, a linear trendline $I^{dth} = k_{th}^{dth} \cdot I^{th}$ ($k_{th}^{dth} < 1$) was fitted to the association of experimental MN recruitment $I^{th}$ and recruitment $I^{dth}$ current thresholds for the $N_r$ recruited MNs, and the distribution of MN input resistance $R = \frac{k_R}{S^{2.43}}$ was increased over the derecruitment time range $\left[t_{r_3}; t_{r_5}\right]$ to:

$$
R^d = \frac{k_R^d}{S^{2.43}} \text{ with } k_R^d = \frac{k_R}{k_{th}^{dth}} > k_R
\tag{EQ. 12}
$$

As reviewed in [20,40], the current-voltage hysteresis also explains the typically lower MN discharge rate observed at derecruitment than at recruitment. For purely modelling perspectives, this

phenomenon was phenomenologically captured by increasing over the time-range $\left[t_{r_3}; t_{r_5}\right]$ the value

of the $C_m$ property, which is the main parameter influencing the precited MN FF in (EQ. 7) for current

inputs close to $I^{th}$. The simulated $FIDF_i^{sim}$ traces were iteratively simulated for the $N_r$ MNs over the

time-range $\left[t_{r_3}; t_{r_5}\right]$ with the $N_r$ $S_i - IP_i$ −calibrated LIF models obtained from Step (3),  for $0.1\mu F \cdot$

$cm^2$ incremental changes in the value of the membrane specific capacitance $C_m$. The $FIDF_i^{sim}$ results

were compared to the $FIDF_i^{exp}$ traces with nRMSE and $r^2$ values. The 'apparent' $C_m$ value returning

the lowest output for (EQ. 13)  was retained and was renamed $C_m^d$.

$$J(C_m) = \frac{\dfrac{\overline{nRMSE}}{100} - \overline{r^2}}{2} \tag{EQ. 13}$$

In the following, the individual spike trains $sp_i^{sim}(t)$ were predicted with $C_m$ over the $[t_{r_0}; t_{r_3}[$ time

range and $C_m^d$ over the $\left[t_{r_3}; t_{r_5}\right]$ time range.

It must be noted that, although this approach can accurately capture the observed MN nonlinear input-

output behaviour and is coherent with the modelling constraints imposed by the LIF formulation in

(EQ. 7), as the $C_m$ and $R$ parameters mainly affect the discharge and recruitment properties of the LIF

model respectively in this model, the $R$ and $C_m$ parameters are passive MN properties which remain

independent from the MN synaptic and membrane activity in actual MNs, while the $R$-to-$R^d$ and $C_m$-

to-$C_m^d$ transformations are merely phenomenological interpretations of complex neurophysiological

mechanisms, and do not provide any insights into the working principles underlying MN recruitment.

This limitation is further discussed in Limitation 3.

**Step (4): Simulating the MN pool firing behaviour**

The firing behaviour of the complete MN pool, i.e. the MN-specific spike trains $sp_j(t)$ of the $N$ virtual

MNs constituting the MN pool, was predicted with a cohort of $N$ LIF models receiving the common

synaptic current $I(t)$ as input. The inert period $IP_j$ and MN size $S_j$ parameters scaling each LIF model

were obtained from the distributions $IP(j)$ and $S(j)$ previously derived at Step (3).

**Validation**

Validation 1

We assessed whether the $N_r$ experimental spike trains $sp_i^{exp}(t)$ were accurately predicted by this 4-

step approach (Figure 1). Steps (2) and (3) (Figure 1) were iteratively repeated with samples of $N_r - 1$

spike trains, where one of the experimentally recorded MNs was not considered. At each $i^{th}$ iteration

of the validation, the $i^{th}$ identified spike train $sp_i^{exp}(t)$ was not used in the derivation of the synaptic

current $I(t)$ (step 2) and in the reconstruction of the MN size and $IP$ distributions $S(j)$ and $IP(j)$ in

the MN pool (step 3). As in Step (4), the $FIDF_i^{sim}$ of the $i^{th}$ MN was finally predicted with a LIF model,

which was scaled with the parameters $S(N_i)$ and $IP(N_i)$ predicted from the $S(j)$ and $IP(j)$

distributions. For validation, $FIDF_i^{sim}$ was compared to the experimental $FIDF_i^{exp}$ with calculation of

$\Delta ft_i^1$, $r^2$ and nRMSE values. This validation was iteratively performed for all the $N_r$ identified MN spike

trains.

<u>Validation 2</u>

We assessed whether the $N$ MN spike trains $sp_j^{sim}(t)$, predicted in Step 4 for the entire MN pool from

the $N_r$ identified trains $sp_i^{exp}(t)$, accurately predicted the effective neural drive (eND$_N$) to the muscle.

The eND$_N$ was computed as the $[0;4]$Hz low-pass filtered cumulative spike train of the $N$ predicted

spike trains, $CST_N = \sum_{j=1}^{N} sp_j^{sim}(t)$. As suggested for isometric contractions [83], the normalized

$eND_N$ was compared for validation against the normalized experimental force  trace $\bar{F}(t)$ with

calculation of the nRMSE and $r^2$ metrics. $\bar{F}(t)$ was also compared with nRMSE and $r^2$ to the

normalized effective neural drive $eND_{N_r}$ computed directly from the $N_r$ experimentally identified MN

spike trains. The added value of the presented workflow (Steps 1 to 4) in predicting the neural drive to

muscle was finally assessed by comparing the (nRMSE, $r^2$) pairs obtained for the $eND_{N_r}$ and $eND_N$

traces. Considering the uncertainty on the value of $N$ in the literature, as previously discussed, the

performance of the method with reconstructed populations of $N = \{N_r, 100, 200, 300, 400\}$ MNs was

investigated. This sensitivity analysis provides different scaling factors $N$ for the normalized MN

mapping (Step 1) and $IP$ and $S_{MN}$ parameter distributions (Step 3) and constrains the number of firing

MNs involved in the computation of eND$_N$ (Step 4).

**Application to MN-driven muscle modelling**

The $N$ MN-specific spike trains $sp_j^{sim}(t)$ predicted in step (4) were input to a phenomenological muscle

model to predict the whole muscle force trace $F_N^{sim}$ in a forward simulation of muscle voluntary

isometric contraction. As displayed in Figure 3, the muscle model was built as $N$ in-parallel Hill-type

models which were driven by the simulated spike trains $sp_j^{sim}(t)$ and replicated the excitation-

contraction coupling dynamics and the contraction dynamics of the $N$ MUs constituting the whole

muscle. In brief, the binary spike train $sp_j^{sim}(t)$ triggered for each $j^{th}$ MU the trains of nerve and

muscle fibre action potentials that drove the transients of calcium ion $Ca^{2+}$ and $Ca^{2+}$-troponin

complex concentrations in the MU sarcoplasm, yielding in a last step the time-history of the MU active

state $a_j(t)$. The MU contraction dynamics were reduced to a normalized force-length relationship that

scaled nonlinearly with the MU active state [96] and transformed $a_j(t)$ into a normalized MU force

trace $\overline{f_j}(t)$. The experiments being performed at constant ankle joint (100°, 90° being ankle

perpendicular to tibia) and muscle-tendon length, it was simplified, lacking additional experimental

insights, that tendon and fascicle length both remained constant during the whole contractile event at

optimal MU length $\overline{l_{opt,J}} = 1$. The dynamics of the passive muscle tissues and of the tendon and the

fascicle force-velocity relationships were therefore neglected. Finally, the MU-specific forces $f_j(t)$

were derived with a typical muscle-specific distribution across the MU pool of the MU isometric tetanic

forces $f_{iso}^{max}(j)$ [64,71,74]. The whole muscle force was obtained as the linear summation of the MU

forces $F_N^{sim}(t) = \sum_j f_j(t)$.

To validate $F_N^{sim}(t)$, the experimental muscle force $F^{exp}(t)$ was first approximated from the

experimental force trace $F(t)$, which was recorded at the foot with a force transducer [26]. The

transducer-ankle joint and ankle joint-tibialis anterior moment arms $L_1$ and $L_2$ were estimated using

OpenSim [97] and a generic lower limb model [98]. Using the model muscle maximum isometric forces,

it was then inferred the ratio $q$ of transducer force $F(t)$ that was taken at MVC by the non-TA muscles

spanning the ankle joint in MVC conditions. The experimental muscle force was estimated as

$F^{exp}(t) = (1 - q) \cdot \frac{L_1}{L_2} \cdot F(t)$ and was compared with calculation of normalized maximum error

(nME), nRMSE and $r^2$ values against the muscle force $F_N^{sim}(t)$ predicted by the MN-driven muscle

model from the $N$ neural inputs.

The whole muscle force $F_{N_r}^{sim}(t)$ was also predicted using the $N_r$ experimental spike trains $sp_i^{exp}(t)$

as inputs to the same muscle model of $N_r$ in-parallel Hill-type models (Figure 3). In this case, each

normalized MU force trace $\overline{f_j}(t)$ was scaled with the same $f_{iso}^{max}(j)$ distribution, however assuming

the $N_r$ MNs to be evenly spread in the MN pool. $F_{N_r}^{sim}(t)$ was similarly compared to $F^{exp}(t)$ with

calculation of $nME, nRMSE$ and $r^2$ values. To assess the added value of the step (1-4) approach in the

modelling of MN-driven muscle models, the $(nME, nRMSE, r^2)$ values obtained for the predicted

$F_{N_r}^{sim}(t)$ and $F_N^{sim}(t)$ were compared.

*Figure 3: MN-driven neuromuscular model. The $N$ in-parallel Hill-type models take as inputs the $N$ spike trains $sp_j^{sim}(t)$*

*predicted in steps (1-4) and output the predicted whole muscle force trace $F_m^{sim}(t)$. The MU-specific active states $a_j(t)$ are*

*obtained from the excitation-contraction coupling dynamics adapted from [99] and extended to model the concentration of*

*the Calcium-troponin system. The MU normalized forces $\overline{f}_j(t)$ are computed by the MU contractile elements ($CE_j$) at MU*

*optimal length and are scaled with values of MU maximum isometric forces $f_{iso,j}^{max}$ to yield the MU force traces $f_j(t)$. The*

*predicted whole muscle force is taken as $F_N^{sim}(t) = \sum_j f_j(t)$.*

# RESULTS

## Experimental data

As reported in Table 1, the experimental datasets $DTA_{35}$ and $HTA_{35}$ respectively identified 32 and 21

spike trains $sp_i^{exp}(t)$ from the trapezoidal isometric TA muscle contraction up to 35%MVC, $HTA_{50}$

identified 14 $sp_i^{exp}(t)$ up to 50%MVC, and $HGM_{30}$ identified 27 $sp_i^{exp}(t)$ from the GM muscle up to

30%MVC. The $N_r = 32$ MN spike trains, identified in this dataset across the complete TA pool of $N =$

400 MNs, are represented in Figure 4A in the order of increasing force recruitment thresholds $F_i^{th}$.

The $N_r$ MNs were globally derecruited at relatively lower force thresholds (Figure 4B) and generally

discharged at a relatively lower firing rate (Figure 4C) at derecruitment than at recruitment.

*Figure 4: Experimental data obtained from HDEMG signal decomposition in the dataset $DTA_{35}$. (A) Time-histories of the*

*transducer force trace in %MVC (green curve) and of the $N_r = 32$ MN spike trains identified from HDEMG decomposition*

*and ranked from bottom to top in the order of increasing force recruitment thresholds $F^{th}$. (B) Association between force*

*recruitment and derecruitment threshold, fitted by a linear trendline $y = 0.9 \cdot x$ ($r^2 = 0.85$). Identity is displayed as a black*

*dotted line. (C) Time-histories of the MN instantaneous discharge frequencies (IDFs, blue dots) smoothed by convolution*

*with a Hanning window of length 400ms (red curve) and with a sixth-order polynomial trendline (green curve) of the lowest-*

*threshold identified ($1^{st}$) MN.*

## Step (1): MN mapping

493 The $N_r = 32$ MNs identified in the dataset $DTA_{35}$ were allocated to the entire pool of $N = 400$ MNs

494 according to their recruitment thresholds $F_i^{th}$ (%MVC). The typical TA-specific frequency distribution

495 of the MN force recruitment thresholds $F^{th}$, which was obtained from the literature and reported in

496 the bar plot in Figure 5A, was approximated (Figure 5B) by the continuous relationship:

$$F^{th}(j) = 0.50 \cdot \left( 58.12 \cdot \frac{j}{N} + \Delta_F^{\left(\frac{j}{N}\right)^{1.83}} \right), j \in [\![1; N]\!] \qquad \text{(EQ. 14)}$$

With this distribution, 231 TA MNs, i.e. 58% of the MN pool is recruited below 20%MVC, which is

consistent with previous conclusions [20].

From the $F^{th}(j)$ distribution, the $N_r$ identified MNs were mapped to the complete MN pool (blue

crosses in Figure 5B) according to their recorded force recruitment thresholds $F_i^{th}$ (ordinates in Figure

4B). As shown in Figure 5C, the $N_r$ MNs identified experimentally were not homogeneously spread in

the entire MN pool ranked in the order of increasing force recruitment thresholds, as two MNs fell in

the first quarter of the MN pool, 5 in the second quarter, 18 in the third quarter and 5 in the fourth

quarter. Such observation was similarly made in the three other experimental datasets, where no MN

was identified in the first quarter and in the first half of the MN pool in the datasets $HGM_{30}$ and $HTA_{50}$

respectively (second column of Table 2). In all four datasets, mostly high-thresholds MNs were

identified experimentally.

*Figure 5: Distribution of force recruitment thresholds $F^{th}$ in the human Tibialis Anterior (TA) muscle and mapping of the $N_r$*

*identified MNs to the complete MN pool. (A) Typical partition obtained from the literature of the TA MN pool in 10%*

*increments in normalized f $F^{th}$. (B) Equivalent $F^{th}$ stepwise distribution (black dots) in a TA pool of $N = 400$ MNs,*

*approximated by the continuous relationship $F^{th}(j)$ (red curve). The mapping (blue crosses) of the $N_r = 32$ MNs identified in*

*the dataset $DTA_{35}$ was obtained from the recorded $F_i^{th}$ (Figure 4B). (C) $N_r = 32$ MNs (red dots) of unknown properties (Left)*

*are mapped (Right) to the complete MN pool from the $F^{th}(j)$ distribution, represented by the blue dots of increasing sizes.*

*The MNs represented here are numbered from left to right and bottom to top from $j = 1$ to $400$.*

**Step (2): Common current input $I(t)$**

To approximate the common synaptic input $CSI(t)$ to the MN pool, the CST and the eND to the MN

pool were obtained in Figure 6 from the $N_r$ MN spike trains identified experimentally using (EQ. 4) and

(EQ. 5). After 20 random permutations of $\frac{N_r}{2}$ complementary populations of MNs, an average

coherence of $coher_{\frac{N_r}{2},mean} = 0.56$ was obtained between $\frac{N_r}{2}$-sized CSTs of the $DTA_{35}$ dataset. From

Figure 2 in [47], a coherence of $coher_{N_r} = 0.8$ is therefore expected between the CST in Figure 6A and

a typical CST obtained with another virtual group of $N_r = 32$ MNs, and by extension with the true CST

obtained with the complete MN pool. The normalized $eND$ and force traces (black and green curves

respectively in Figure 6B) compared with $r^2 = 0.92$ and $nRMSE = 20.0\%$ for the $DTA_{35}$ dataset.

With this approach, we obtained $coher_{N_r} > 0.7$, $r^2 > 0.7$ and $nRMSE < 30\%$ for all four datasets,

with the exception of $HGM_{30}$ for which $coher_{N_r} < 0.7$ (third column of Table 2). For the TA datasets,

the sample of $N_r$ identified MNs was therefore concluded to be sufficiently representative of the

complete MN pool for its linearity property to apply, and the $eND$ (red curve in Figure 6B for the

dataset $DTA_{35}$) in the bandwidth $[0,10]Hz$ was confidently identified to be the common synaptic

input ($CSI$) to the MN pool. As observed in Figure 6B, a non-negligible discrepancy, which partially

explains why $coher_{N_r} \neq 1$, was however systematically obtained between the $eND$ and force traces

in the regions low forces, where mostly small low-threshold MNs are recruited. As discussed in the

Discussion Section, this discrepancy reflects the undersampling of small MUs identified from

decomposed HDEMG signals and a bias towards mainly identifying the large high-threshold units which

are recruited close to the plateau of force. From Figure 2 in [47], it is worth noting that the computed

$CSI$ in Figure 6B (red curve) accounts for 60% of the variance of the true synaptic input, which is linearly

transmitted by the MN pool, while the remaining variance is the MN-specific synaptic noise, which is

assumed to be filtered by the MN pool and is neglected in the computation of the $eND$ in this

workflow.

*Figure 6: Neural drive to the muscle derived from the $N_r$ identified MN spike trains in the dataset $DTA_{35}$. (A) Cumulative*

*spike train (CST) computed by temporal binary summation of the $N_r$ identified MN spike trains. (B) Effective neural drive –*

*Upon applicability of the linearity properties of subsets of the MN pool, the effective neural drive is assimilated as the*

*synaptic input to the MN pool. The normalized common synaptic input (red), control (black) and noise (blue) are obtained*

*from low-pass filtering the CST in the bandwidths relevant for muscle force generation. The normalized experimental force*

*trace (green curve) is displayed for visual purposes.*

As discussed in the Methods section, the normalized $CSI$ (red curve in Figure 6B) was simplified to be

linearly related to the total dendritic membrane current $I(t)$. To do so, the scaling factor was

determined with a typical distribution of the MN membrane rheobase $I^{th}(j)$ in a cat MN pool, which

was obtained (Figure 7A) from the literature [85,87,88,90,91,100-102] as:

$$I^{th}(j) = 3.9 \cdot 10^{-9} \cdot \Delta_I^{\left(\frac{j}{N}\right)^{1.18}}, j \in [\![1; N]\!] \qquad \text{(EQ. 15)}$$

552 Using (EQ. 6), the time-history of the current input $I(t)$ was obtained (Figure 7B):

$$I(t) = \begin{cases} 0 \ if \ t < ft_{N_1}^1 \\ 3.9 \cdot 10^{-9} + 6.0 \cdot 10^{-8} \cdot CSI(t) \ else \end{cases} \qquad \text{(EQ. 16)}$$

554 *Figure 7: (A) Typical distribution of MN current recruitment threshold $I^{th}(N_i)$ in a cat MN pool according to the literature.*

555 *(B) Current input $I(t)$, taken as a non-continuous linear transformation of the common synaptic input (red curve in Figure*

556 *6B) for the dataset $DTA_{35}$.*

### Step (3): LIF model – MN size calibration and distribution

Because of the modelling choices made for our MN LIF model, the MN inert period ($IP$) and the MN

size $S$ parameters entirely define the LIF-predictions of the MN firing behaviour. The $IP_i$ parameters

of the $N_r$ MNs in the dataset $DTA_{35}$ (Figure 8) were obtained from the maximum firing frequency of

the 20 MNs identified to 'saturate', from which the distribution of $IP$ values in the entire pool of $N$

MNs was obtained: $IP(j)[s] = 0.04 \cdot j^{0.05}$. With this approach, the maximum firing rate assigned to

the first recruited and unidentified MN is $\frac{1}{IP_1} = \frac{1}{0.04 \ s} = 25Hz$. The $IP$ distributions obtained with this

approach for the three other datasets are reported in the fourth column of Table 2 and yielded

physiological approximations of the maximum firing rate for the unidentified lowest -threshold MN for

all datasets, with the exception of the dataset $HTA_{50}$, which lacks the information of too large a

fraction of the MN pool for accurate extrapolations to be performed.

The size parameter $S_i$ of the $N_r$ LIF models was calibrated using the minimization function in (EQ. 14)

so that the LIF-predicted filtered discharge frequencies $FDIF_i^{sim}(t)$ of the $N_r$ MNs replicated the

experimental $FDIF_i^{exp}(t)$, displayed in Figure 9A and B (blue curves). As shown in Figure 9C, the

recruitment time $ft^1$ of three quarters of the $N_r$ identified MNs was predicted with an error less than

250ms. The calibrated LIF models were also able to accurately mimic the firing behaviour of the 27

lowest-threshold MNs as experimental and LIF-predicted FIDF traces compared with $r^2 > 0.8$ and

$nRMSE < 15\%$ (Figure 9D and E). The scaled LIF models reproduced the firing behaviour of the five

highest-threshold MNs ($Ni > 300$) with moderate accuracy, with $\Delta ft^1$ and $nRMSE$ values up to -1.5s

and 18.2%. Despite modelling limitations discussed in the Discussion section, the global results in

Figure 9 confirm that the two-parameters calibrated LIF models can accurately reproduce the firing

and recruitment behaviour of the $N_r$ experimental MNs. It must be noted that the $\Delta ft^1$ and $r^2$ metrics

were not included in the calibration procedure and the $(\Delta ft^1, r^2)$ values reported in Figure 9 were

therefore blindly predicted.

The cloud of $N_r$ pairs $\{N_i; S_{N_i}\}$ of data points (black crosses in Figure 10A), obtained from MN mapping (Figure 5B and C) and size calibration for the dataset $DTA_{35}$, were least-squares fitted (red curve in Figure 10A, $r^2 = 0.96$) by the power relationship in (EQ. *17*), with $\Delta_S = 2.4$:

$$S(j) = 1.49 \cdot 10^{-7} \cdot \Delta_S^{\left(\frac{j}{N}\right)^{1.47}} \qquad \text{(EQ. 17)}$$

As reported in the last column of Table 2, the minimum MN size (in $m^2$) obtained by extrapolation of this trendline was in the range $[1.20, 1.73] \cdot 10^{-1}mm^2$ for the datasets $DTA_{35}, HTA_{35}$ and $HGM_{30}$, with expected maximum MN size (in $m^2$) in the range $[2.88, 4.15] \cdot 10^{-1}mm^2$ which is consistent with typical cat data [42,103-108], even if slightly in the lower range. The low value for the $c$ coefficient observed in Table 2 for the $TA_{50}$ and $GM_{30}$ datasets suggests that the typical skewness in MN size distribution reported in the literature [42] was not captured for these two datasets, in which low-threshold MNs in the first quarter of the MN population were not identified from HDEMG signals.

606 *trains (CST), and the coefficients defining the distributions in the complete MN pool of (3) the inert period (IP) parameter and*

607 *of (4) the MN size (S). For the TA and GM muscles, $N = 400$ and $N = 550$ respectively.*

| | Identified population | | CST | $IP(j)[s] = a \cdot j^b$ | | $S(j)[m^2] = S_{min} \cdot \Delta_S^{\left(\frac{j}{N}\right)^c}$ | |
|---|---|---|---|---|---|---|---|
| Dataset | $N_r$ | $i \rightarrow N_i$ | $coher_{N_r}$ | $a$ | $b$ | $S_{min}[m^2]$ | $c$ |
| $DTA_{35}$ | 32 | 9-313 | 0.80 | 0.04 | 0.06 | $1.49 \cdot 10^{-7}$ | 1.47 |
| $HTA_{35}$ | 21 | 72-313 | 0.71 | 0.04 | 0.05 | $1.73 \cdot 10^{-7}$ | 2.76 |
| $HTA_{50}$ | 14 | 233-346 | 0.71 | 0.0006 | 0.80 | $1.35 \cdot 10^{-7}$ | 0.85 |
| $HGM_{30}$ | 27 | 162-412 | 0.65 | 0.03 | 0.20 | $1.20 \cdot 10^{-7}$ | 0.57 |

As displayed in Figure 9A and B, the phenomenological adjustment of the $R$ and $C_m$ parameters

described in (EQ. 12) and (EQ. 13) successfully captured the main effects of the hysteresis mechanisms

involved with MN derecruitment, where MNs discharge at lower rate and stop firing at lower current

input than at recruitment. In all four datasets, $J(C_m) = \frac{\overline{nRMSE} - \overline{r^2}}{2}$ was parabolic for incremental values

of $C_m$ and a global minimum was found for $C_m^d = 2.0, 2.0, 1.6$ and $2.2 \mu F \cdot cm^2$ for the datasets

$DTA_{35}, HTA_{35}, HTA_{50}$ and $HGM_{30}$, respectively. These values for the parameter $C_m^d$ were retained

for the final simulations (Figure 12 and Figure 13) over the $\left[t_{r_3}; t_{r_5}\right]$ time range.

**Step (4): Simulating the MN pool firing behaviour**

As displayed in Figure 10B for the dataset $DTA_{35}$, the $S(j)$ distribution determined the MN-specific

electrophysiological parameters (input resistance $R$ and membrane capacitance $C$) of a cohort of $N =$

400 LIF models, which predicted from $I(t)$ the spike trains of the entire pool of $N$ MNs (Figure 10C).

As displayed in the plot of smoothed FIDFs in Figure 10C, the output behaviour of the reconstructed

MN pool agrees with the Onion-skin scheme [109] where lower-threshold MNs start discharging at

higher frequencies and reach higher discharge rates than larger recruited units.

*Figure 10: Reconstruction of the firing behaviour and recruitment dynamics of the complete MN pool. (A) The $N_r$ calibrated*

*MN sizes (black crosses) are lest-squares fitted by the power trendline $S(j)[m^2] = 1.49 \cdot 10^{-7} \cdot \Delta_S^{\left(\frac{j}{N}\right)^{1.47}}$, which reconstructs*

*the distributions of MN sizes in the complete MN pool. (B) The $S(j)$ distribution determines the MN-specific R and C parameters*

*of a cohort of N LIF models, which is driven by the current input $I(t)$ and predicts (C) the spike trains $sp_j^{sim}(t)$ of the N virtual*

*MNs constituting the complete MN pool. (D) The MN instantaneous discharge frequencies were computed from the simulated*

*the spike trains $sp_j^{sim}(t)$ and were smoothed with a sixth-order polynomial [55]. One out of ten IDFs is displayed for clarity.*

*The blue-to-red gradient for small-to-large MNs shows that the output behaviour of the reconstructed MN population is in*

*agreement with the Onion-Skin scheme.*

*Figure 11: Detailed description of the 4-step workflow applied to the $DTA_{35}$ dataset. The firing activity of a fraction of the MN pool is obtained from decomposed HDEMG signals. These $N_r = 32$ experimental spike trains provide an estimate of the effective neural drive to muscle and explain most of the MN pool behaviour (coherence = 0.8). From a mapping of the $N_r$ identified MNs to the complete MN pool (Step (1)), literature knowledge on the typical distibution of $I^{th}$ in a mammalian MN pool, and using the linearity properties of the population of $N_r$ MNs, the current input $I(t)$ is estimated (Step (2)). $I(t)$ is input to $N_r$ LIF models of MN to derive, after a one-parameter calibration step minimizing the error between experimental and LIF-predicted FIDFs, the distribution of MN sizes across the complete MN pool (Step (3)). The distribution of MN sizes, which entirely describes the distribution of MN-specific electrophysiological parameters across the MN pool, scales a cohort of $N = 400$ LIF models which transforms $I(t)$ into the simulated spike trains of the $N$ MNs of the MN pool (Step (4)). The effective neural drive to muscle is estimated from the $N$ simulated spike trains.*

 **Validation**

The four-step approach summarized in Figure 1 is detailed in Figure 11 and was validated in two
ways.

<u>Validation 1</u>

The simulated spike trains were validated for the $N_r$ MNs by comparing experimental and LIF-predicted
FIDFs, where the experimental information of the investigated MN was removed from the
experimental dataset and not used in the derivation of the $IP(j)$ and $S(j)$ distributions of the synaptic
current $I(t)$. For the $N_r$ MNs of each dataset, Figure 12 reports the absolute error $\Delta f t^1$ in predicting
the MN recruitment time (1st row) and the comparison between experimental and LIF-predicted
filtered instantaneous discharge frequencies (FIDFs) with calculation of $nRMSE$ (%) (2nd row) and $r^2$
(3rd row) values over the $\left[t_{r_0}; t_{r_5}\right]$ time range. In all datasets, the recruitment time of 45-to-65% of the
$N_r$ identified MNs was predicted with an absolute error less than $\Delta f t^1 = 250 ms$. In all datasets, the
LIF-predicted and experimental FIDFs of more than 80% of the $N_r$ MNs compared with $nRMSE < 20\%$
and $r^2 > 0.8$, while 75% of the $N_r$ MN experimental and predicted FIDFs compared with $r^2 > 0.8$ in
the dataset $HGM_{30}$. These results confirm that the four-step approach summarized in Figure 1 is valid
for all four datasets for blindly predicting the recruitment time and the firing behaviour of the $N_r$ MNs
recorded experimentally. The identified MNs that are representative of a large fraction of the complete
MN pool, i.e. which are the only identified MN in the range $\left[N_i - \frac{N}{10}; N_i + \frac{N}{10}\right]$ of the entire MN pool,
such as the 1st MN in the dataset $DTA_{35}$ (Figure 5C), are some of the MNs returning the highest $\Delta f t^1$
and $nRMSE$ and lowest $r^2$ values. As observed in Figure 12, ignoring the spike trains $sp_i^{exp}(t)$ of those
'representative' MNs in the derivation of $I(t), IP(j)$ and $S(j)$ in steps (2) and (3) therefore affects the
quality of the predictions more than ignoring the information of MNs that are representative of a small
fraction of the MN population. In all datasets, $nRMSE > 20\%$ and $r^2 < 0.8$ was mainly obtained for
the last-recruited MNs (4th quarter of each plot in Figure 12) that exhibit recruitment thresholds close
to the value of the current input $I(t)$ during the plateau of constant force in the time range $\left[t_{tr_2}; t_{tr_3}\right]$.
In all datasets, the predictions obtained for all other MNs that have intermediate recruitment
thresholds and are the most identified MNs in the datasets, were similar and the best among the pool
of $N_r$ MNs.

*Figure 12: Validation of the MN recruitment and firing behaviour predicted with the 4-step approach (Figure 1) for the $N_r$ MNs*

*experimentally identified in the four datasets $DTA_{35}, HTA_{35}, HTA_{50}$ and $HGM_{30}$ described in Table 1. For the validation of*

*each $i^{th}$ predicted MN spike train, the experimental information of the $i^{th}$ identified MN was omitted when deriving the*

*current input $I(t)$ and the $IP(j)$ and $S(j)$ distributions (steps 2 and 3), from which the $IP_i$ and $S_i$ parameters are obtained*

*without bias for the $i^{th}$ MN. The spike train $sp_i^{sim}(t)$ of the $i^{th}$ MN is then predicted with an $IP_i, S_i$-scaled LIF model receiving*

*$I(t)$ as input. The absolute error in predicting the experimental recruitment time $\Delta ft_i^1$ (s) is reported for each of the $N_r$ MNs*

*in the first row of the figure. Experimental and filtered instantaneous discharge frequencies $FIDF_i^{exp}(t)$ and $FIDF_i^{sim}(t)$,*

*computed from $sp_i^{exp}(t)$ and $sp_i^{sim}(t)$ are compared with calculation of $nRMSE$ (%) and $r^2$ in the second and third rows of*

*the figure. The dashed lines represent the $\Delta ft^1 \in [-250; 250]ms$, $nRMSE \in [0; 15]\%MVC$ and $r^2 \in [0.8, 1.0]$ intervals of*

*interest respectively.*

<u>Validation 2</u>

The effective neural drive predicted by the 4-step approach summarized in Figure 1 was validated by

comparing for isometric contractions, the normalized $eND_N$ (orange traces in Figure 13) computed

from the $N$ predicted spike trains $sp_j^{sim}(t)$ to the normalized force trace $\bar{F}(t)$ (green trace in Figure

13). The $eND_N$ was accurately predicted for the datasets $DTA_{35}$ and $HTA_{35}$, with $r^2 = 0.97 - 0.98$

and $nRMSE < 8\%$ (Table 3). The results obtained from the dataset $HGM_{30}$ returned $r^2 = 0.89$ and

$nRMSE = 18.2\%$ (Table 3), accurately predicting the $eND_N$ for the positive ramp and first half of the

plateau of force and then underestimating the true neural drive ($HGM_{30}$, Figure 13). The $eND_N$

predicted for the dataset $HTA_{50}$ returned results of lower accuracy ($r^2 = 0.88$ and $nRMSE = 15.1\%$)

compared to the three other datasets. A null $eND$ was predicted for one third of the simulation where

a muscle force up to $20\%MVC$ is generated, and the $eND$ was overestimated for the rest of the

(de)recruitment phase ($HTA_{50}$, orange trace, Figure 12). As detailed in the discussion, the latter is

explained by an inadequate $IP(j)$ distribution (Table 2), due to a lack of experimental information in

the dataset $HTA_{50}$, which returns non-physiological maximum firing rates for the low-thresholds MNs.

Two data points (16; 0.032) and (118; 0.037), obtained from the experimental data in the dataset

$DTA_{35}$ with a scaling factor of $\frac{35}{50}$ applied to the $IP$ values, were appended to the list of $N_r$ data points

$(N_i; IP_{N_i})$ to describe the maximum firing behaviour of the first half of the MN pool for which no MN

was identified for the dataset $HTA_{50}$ (Table 2). A new $IP(j)$ distribution was obtained, returning an

improved estimation of the $eND_N$ ($HTA_{50}$, purple trace, Figure 12) with respectively lower and

higher $nRMSE$ and $r^2$ metrics (Table 3). With three times lower $nRMSE$ and higher $r^2$ values for all

four datasets (Table 3), the $eND_N$ predicted from the 4-step approach (orange dotted traces in Figure

13) was a more accurate representation of the real effective neural drive than the $eND_{N_r}$ (blue dotted

traces in Figure 13) computed from the $N_r$ experimental spike trains, especially in the phases of MN

(de)recruitment where the real effective neural drive was underestimated by $eND_{N_r}$. The predicted

$eND_N$ also produced less noise than the $eND_{N_r}$ trace during the plateau of force. With $r^2 > 0.85$ and

$nRMSE < 20\%$, this four-step approach is valid for accurately reconstructing the $eND$ to a muscle

produced by a collection of $N$ simulated MN spike trains, which were predicted from a sample of $N_r$

experimental spike trains. It is worth noting that the accuracy of the $eND_N$ predictions, performed in

Table 3 and Figure 13 for $N_{TA} = 400$ and $N_{GM} = 550$ MNs as discussed in the Methods Section, was

not sensitive to the size $N$ of the reconstructed MN population. For example, for the $DTA_{35}$ dataset,

the $eND_N$ computed with populations of $N = \{32, 100, 200, 300, 400\}$ MNs systematically compared

to the real effective neural drive with $r^2 = 0.98$, and respectively returned $nRMSE = 7.7, 6.9, 6.5, 6.1$

and $5.9\%$, with slightly more accurate predictions of $eND_N$ with an increasing $N$ at low force during

derecruitment.

*Table 3: For all four datasets, the $r^2$ and nRMSE values obtained for the comparison of the time-histories of the normalized*

*experimental force trace against the effective neural drive (1) $eND_{N_r}$ and (2) $eND_N$ computed from the spike trains of (1)*

*the $N_r$ identified MNs and (2) the N virtual MNs. The results for $HTA_{50}$ (1) were obtained with the standard approach,*

*while those for $HTA_{50}$ (2) were obtained with a revisited $IP(j)$ distribution (see text).*

| | $N_r$ MNs (experimental) | | $N$ MNs (simulated) | |
|---|---|---|---|---|
| Dataset | $r^2$ | $nRMSE$ (%) | $r^2$ | $nRMSE$ (%) |
| $DTA_{35}$ | 0.92 | 19.5 | 0.98 | 5.9 |
| $HTA_{35}$ | 0.85 | 26.8 | 0.97 | 7.9 |
| $HTA_{50}$ (1) | 0.83 | 26.9 | 0.88 | 15.1 |
| $HTA_{50}$ (2) | | | 0.92 | 10.6 |
| $HGM_{30}$ | 0.85 | 28.8 | 0.89 | 18.2 |

*Figure 13: For the 4 datasets, validation against the normalized force trace $\bar{F}(t)$ (green trace) of the normalized effective*

*neural drive to muscle $(eND_N)$ produced by the complete MN pool (orange trace) and computed from the N MN spike trains*

*$sp_j^{sim}(t)$ predicted with the 4-step approach. For comparison, the $eND_{N_r}$ directly computed from the $N_r$ identified MNs is*

*also reported (blue dashed trace). For the dataset $HTA_{50}$, an additional prediction was performed (purple trace), where the*

*$IP(j)$ distribution obtained from step (3) was updated to return physiological maximum firing rates for all N MNs.*

**Application to MN-driven modelling**

Figure 14 reports for the datasets $DTA_{35}$ and $HTA_{35}$, for which the $eND_N$ was the most accurately

predicted among the four datasets (Figure 13, Table 3), the time evolutions of the whole muscle force

$F(t)$ recorded experimentally (green curve) and the whole muscle forces predicted with the MN-

driven muscle model described in Figure 3 using $N_r$ experimental ($F_{N_r}(t)$, blue curves) and $N$ simulated

($F_N(t)$, red curves) spike trains as inputs. The muscle force trace $F_N(t)$ obtained from the $N$ simulated

spike trains derived in steps (1-4) compared with the experimental force $F(t)$ with $r^2 = 0.95$ and

$nRMSE = 11\%$ for the dataset $DTA_{35}$ and $r^2 = 0.97$ and $nRMSE = 11\%$ for the dataset $HTA_{35}$.

These results validate the approach provided in this study for predicting the whole muscle force as a

collection of MU force contributions from the MN-specific contributions to the effective neural drive.

The muscle force trace $F_N(t)$ obtained from the $N$ MN spike trains returned more accurate predictions

than $F_{N_r}(t)$ ($r^2 = 0.88$ and $nRMSE = 27\%$ for the dataset $DTA_{35}$ and $r^2 = 0.79$ and $nRMSE =$

$33\%$ for the dataset $DTA_{35}$), in which case the reconstruction of the MN pool in steps (1-4) was

disregarded and the $N_r$ experimental spike trains were directly used as inputs to the muscle model in

Figure 3.

*Figure 14: Prediction of the experimental force trace $F(t)$ (green curve) with the MN-driven muscle model described in*

*Figure 3. $F_{N_r}(t)$ (blue curve) was predicted with the $N_r$ experimental spike trains input to the muscle model. $F_N(t)$ (red*

*trace) was obtained from the cohort of $N$ experimental spike trains simulated by the four-step approach.*

# DISCUSSION

This study reports a novel four-step approach, summarized in Figure 1 and displayed in detail in Figure 11, to reconstruct the recruitment and firing behaviour of a complete human pool of $N$ MNs from a sample of $N_r$ experimental spike trains obtained from the decomposition of HDEMG or intramuscular recordings during voluntary contractions. This approach can help neuroscientists, experimentalists, and modelers to investigate MN pool neuromechanics, better understand experimental datasets, and control more detailed neuromuscular models to advance our understanding of the neural strategies taken by the human central nervous system to control voluntary muscle contractions.

The three first steps of our approach identify from a sample of $N_r$ experimental spike trains a distribution of the MN electrophysiological properties across the MN pool. The $N_r$ MN spike trains are used to approximate the common synaptic input $CSI(t)$ to the complete MN pool (Figure 7). For simplicity, the $CSI$ is linearly related to the post-synaptic total dendritic membrane current $I(t)$, which is input to a cohort of $N_r$ single-compartment LIF models with current synapses. The LIF firing behaviour is entirely described by the phenomenological MN Inert Period parameter $IP_i$, derived from the experimental data (Figure 8), and by the MN size $S_i$ parameter to which all the MN electrophysiological properties are related, according to the relations provided in [42]. After calibration of the $S_i$ parameter, the $N_r$ LIF models accurately mimic the filtered discharge behaviour and accurately predict the recruitment dynamics of the $N_r$ experimental MNs (Figure 9). The $N_r$ MNs are allocated into the complete MN pool (Figure 5B and C) according to their recorded force recruitment thresholds $F_i^{th}$ (Figure 4B) and a typical species- and muscle-specific $F^{th}(j)$ distribution (Figure 5). From the previous findings, the continuous distribution of MN sizes $S(j)$ (Figure 10A) is derived for the complete pool of $N$ MNs. $S(j)$ defines the electrophysiological properties [42] of the MNs constituting the complete MN pool. The neural behaviour of the complete pool of $N$ MNs is predicted in the 4th step with a cohort of $N$ $S_j$-$IP_j$-scaled LIF models and the application of the current drive $I(t)$ (Figure 10).

**Validation of the approach**

The approach was successfully validated both for individual MNs and for the complete pool of $N$ MNs. By blindly scaling the LIF models in steps (1-3) with permutations of $N_r - 1$ input experimental spike trains, the filtered discharge behaviour and the recruitment dynamics of the $N_r$ individual MNs were accurately predicted for the four investigated datasets (Figure 12). The effective neural drive ($eND_N$) to muscle elicited by the complete pool of $N$ MNs was also accurately predicted by the 4-step approach

for the $HTA_{35}$, $DTA_{35}$ and $HGM_{30}$ datasets (Figure 13, Table 3). The latter result suggests that the

recruitment dynamics and the normalized distribution of firing rates across the firing MN pool were

accurately predicted for the non-identified population of $N - N_r$ MNs. The accuracy of the $eND_N$

predictions was not sensitive to the size $N$ of the reconstructed population, with acceptable $eND_N$

predictions ($nRMSE < 20\%, r^2 > 0.9$) obtained with as few as ten distributed MNs. This suggests that

the MN mapping (Step 1) and the derivation of the MN properties distribution (Step 3) in Figure 8 and

Figure 10A are the key contributions for describing the complete MN pool behaviour from the input

experimental data.

Despite the modelling limitations, discussed in [44] and in the Limitations Section, inherent to single-

compartment LIF models with current synapses and related to the phenomenological $IP$, $C^d$ and $R^d$

parameters, the LIF models scaled following the methodology described in Step (3) can capture the

distribution of MN discharging characteristics across the MN population. For example, the firing

activity of the reconstructed MN population agrees with the onion-skin scheme (Figure 10D), as

expected for slow low-force trajectories in human muscle contraction. While the MN size parameter

calibration (Figure 9) only relies on the minimization of the root-mean square difference between

experimental and predicted FIDFs, the normalized time-history of the FIDFs (high $r^2$ in Figure 12) and

the MN recruitment times (low $\Delta ft^1$ values in Figure 12) of the $N_r$ MNs are both accurately mimicked

(Figure 9) and blindly predicted (Figure 12) for all four datasets. Some realistic aspects of the MN

neurophysiology are also maintained.  For example, the range of MN surface areas $[0.15; 0.36]mm^2$

and input resistances $[0.5; 4.1]M\Omega$ obtained from the $S(j)$ distribution falls into the classic range of

MN properties for cats [21,42], which is consistent with the distribution of rheobase values used in

Step (2) that was obtained from cat data. The distribution of MN input resistance in the MN pool,

obtained from the calibrated sizes $S(j)$, defines the individual MN rheobase thresholds and predicts

that 80% of the human TA MU pool is recruited at 35%MVC (Figure 10C), which is consistent with

previous findings in the literature [20,64].

**Benefits of the approach**

Benefit 1 – Better understanding the MN pool firing behaviour inferred from decomposed HDEMG

outputs

The workflow provides a method to better understand datasets of $N_r$ experimental spike trains

obtained from signal decomposition. HDEMG decomposition techniques commonly identify samples

of few tens of MNs [27] at most, i.e. typically less than 10-25% of the MU pool. The mapping in Figure

5 and Figure 10A demonstrates that these $N_r$-sized samples of MNs currently are not representative

of the complete MN population, as supported by the $[0.65; 0.80]$ $coher_{N_r}$ values in Table 2. EMG

decomposition shows a bias towards under-sampling small low-threshold MUs and identifying large

high-threshold MUs because of their larger electric signal amplitude detected by the HDEMG

electrodes. Consequently, the $CST_{N_r}$ computed from the $N_r$ identified MNs is not an accurate

representation of the true muscle neural drive (Figure 13, Table 3). The method in Figure 11 identifies

credible values for some MN-specific electrophysiological and morphometric properties, including MN

membrane surface area, input resistance, capacitance, rheobase and time constant, for the dataset of

$N_r$ experimental MNs and for the $N - N_r$ MNs that cannot be recorded. Using a realistic distribution

of these properties, the $eND_N$ produced by the reconstructed MN population is accurately estimated,

insensitive to the value of $N$, because it includes the firing activity of the complete fraction of the

smaller MNs that were underrepresented in the experimental dataset, especially during the force

ramps in $[t_1;\ t_2]$ and $[t_3;t_4]$ (Figure 2), where the $eND_{N_r}$ can reach 70% normalized maximum error.

The $eND_N$ besides includes the activity of the complete population of high-threshold MNs, which

reduces noise (orange vs blue trace in Figure 13) during the plateau of force in $[t_2;t_3]$. The $eND_N$ also

filters the local non-physiological variations displayed by the $eND_{N_r}$ such as the decrease in $eND_{N_r}$ in

$[10;15]s$ in $DTA_{35}$ and $HTA_{35}$ (top plots in Figure 13), the sudden rise and drop in $eND_{N_r}$ at $t = 10s$

and $t = 23s$ in $HTA_{50}$ (bottom left plot in Figure 13) or the steeply decreasing $eND_{N_r}$ from $t = 28s$

in $HGM_{30}$ (bottom right plot in Figure 13). Understanding and accounting for these experimental

limitations is important and has critical implications in HDEMG-driven neuromuscular modelling when

the $CST_{N_r}$ is input to a muscle model [110,111].

The workflow also provides a method to support future experimental investigations. The signal

decomposition process can be refined and more MNs can be identified by iteratively running this

workflow and using the $N - N_r$ simulated spike trains of the non-identified MNs as a model for

identifying more MNs from the recorded signals. The method moreover provides simple

phenomenological two-parameter-scaled LIF models described in step (3) to accurately replicate the

recruitment and firing behaviour of the $N_r$ identified MNs (Figure 9) for further investigation.

Benefit 2 – Advances in MN pool modelling

This four-step approach advances the state-of-the-art in MN pool modelling. As discussed in the

introduction, using a sample of $N_r$ experimental MN spike trains permits for the first time in human

MN pool modelling to (1) estimate the true time-course of the common synaptic input to the MN pool,

which cannot be measured experimentally, (2) approximate with a one-parameter calibration,

available HDEMG data and knowledge from the literature, the MN-specific electrophysiological

properties of all the MNs in the MN pool and a realistic distribution of these properties across the pool,

which could, to date, only be obtained in specialized experimental and MN pool modelling studies in

animals [39], and (3) validate the forward predictions of MN spike trains and effective neural drive to

muscle to human experimental data. The pool of LIF models, the maximum firing frequency of which

is obtained from the available experimental data, intrinsically accounts for the onion-skin theory [109]

(Figure 10D), and better replicates the MN membrane dynamics of the MN pool than Fuglevand-type

phenomenological models [10,29], where the MN-specific firing rates are predicted as a linear function

of the current input $I(t)$. Moreover, single-compartment current-input LIF models can credibly

replicate most of the MN membrane dynamics [43] and allowed accurate predictions of the MN pool

behaviour (Figure 12, Figure 13), while they provide a simpler modelling approach and a more

convenient framework for MN electrophysiological parameter assignment in the MN pool than

comprehensive Hodgkin-Huxley-type MN models [30,32,112,113]. This four-step approach

demonstrated to be robust to systematic differences in the input experimental datasets. For example,

it accurately predicted the individual MN firing behaviour of the $N_r$ MNs for all four datasets (Figure

12) despite the latter being obtained at different levels of muscle contractions, on different subjects

and muscles and in different experimental approaches. The approach accurately predicted the $eND_N$

of both $DTA_{35}$ and $HTA_{35}$ datasets from $N_r$ =32 and 21 identified MNs respectively (Table 3),

suggesting that the method is not sensitive to the number $N_r$ of identified MNs, providing that the $N_r$

identified MNs and their properties are homogeneously spread in the MN pool, as in datasets $DTA_{35}$

and $HTA_{35}$ where at least one MN is identified in the each 10%-range of the rheobase domain. The

accurate prediction of $eND_N$ for the dataset $HGM_{30}$ in the time range $[4.7; 7.2]\ s$ (Figure 13 bottom-

right plot) also suggests that the quality of the predictions is not sensitive to the hindlimb muscle

investigated, providing that the $F^{th}$ distribution is representative of the investigated muscle.

Benefit 3 – Relevance for neuromuscular modelling

A MN-driven neuromuscular model of in-parallel Hill-type actuators describing the MUs (Figure 3) is

described in the Methods section and is used to predict the experimental muscle isometric force (green

trace in Figure 14) from the $N_r$ experimental or $N$ reconstructed MN spike trains. The complete

reconstruction of the discharging MN population detailed in Figure 11 is a key step towards advancing

the state-of-the-art in neuromuscular modelling on several aspects.

Firstly, the neural input to the neuromuscular model in Figure 3 is a vector of experimental spike trains

that is a more easily interpretable and a more detailed description of the muscle neural drive than the

rectified-normalized-filtered envelopes of recorded bipolar EMGs (bEMGs) [96,114,115] or the $CST_{N_r}$

[110,111,116] typically used in single-input muscles models. Secondly, despite advanced multi-scale

approaches in modelling whole muscles as a single equivalent MUs [99,117-121], the multiple Hill-MU

model in Figure 3 provides a more convenient framework to model in detail the continuous distribution

of the MU excitation-contraction and force properties in the MU pool. Thirdly, a few other multi-MU

models were proposed in the literature [9,10,12,39,99,122,123], some of which  used the Fuglevand's

formalism [29] to model the MN firing behaviour and recruitment dynamics of complete pools of $N$

MNs, which are intrinsic to the experimental $sp_i^{exp}(t)$, to predict whole muscle forces with collections

of $N$ MU Hill-type [9,123] Hill-Huxley-type [10] or twitch-type [39] muscle models. However, these

studies lacked experimental neural control at the MN level and considered artificial synaptic inputs to

their model, the predicted force of which was indirectly validated against results from other models

and not against synchronously recorded experimental data, as performed in Figure 12 to Figure 14 in

this study. Finally, Figure 14 demonstrates that the reconstruction of the complete MN population

described in this study (steps 1 to 4 in Figure 1) is a key step for accurate MN-driven neuromuscular

predictions of muscle force. The $N_r$-MU model, that receives the $N_r$ individual MN spike trains

$sp_i^{exp}(t)$, intrinsically underestimates the whole muscle activity when dominantly low-threshold MUs

are recruited but are under-represented in the experimental sample (Figure 5, Figure 13). The $N$-MU

model, which receives as inputs the $N$ spike trains $sp_j^{sim}(t)$ generated by the four-step approach,

allows a more realistic assignment of distributed MU properties (MU type and maximum isometric

force) to the complete MU population, and returned more accurate force predictions than the $N_r$-MU

model (Figure 14). It is worth noting that the $N$-model did not require any parameter calibration except

the MN size in step 3. The detailed modelling of the distribution of the excitation-contraction

properties of the MUs makes the $N$-model more suitable for investigating the muscle neuromechanics

than typical EMG-driven models, the neural parameters of which do not have a direct physiological

correspondence and must be calibrated to match experimental joint torques [96,124,125].

**Limitations of the approach and potential improvements**

Despite the aforementioned good performance of the four-step workflow, the method presents 4

main limitations, for some of which potential improvements are proposed in the following.

Limitation 1 – Model validation

The two validations of the approach (Figure 12 and Figure 13) present some limitations. The local

validation in Figure 12 only ensures that the method accurately estimates the MN firing behaviour for

the fractions of the MN pool that were experimentally identified. This local validation alone does not

inform on the accuracy of the predictions for the non-identified regions of the MN pool and must be

coupled with a global validation of the MN pool behaviour by validating the predicted $eND_N$. While

the local validation was successful for the $HTA_{50}$ dataset (Figure 12), where less than 30% of the active

MN pool is represented, it is inferred in Figure 13C that the individual MN firing rates were
overestimated for the first half of the non-recorded MN pool. This local validation would be self-
sufficient for experimental samples that contain a large and homogeneously spread population of
identified MNs, obtained from decomposed fine-wire intramuscular electrode and HDEMG grid signals.
The validation of the $eND_N$ is performed in this study against an experimental force recorded by a
transducer (green traces in Figure 13), which accounts for the force-generating activity of both the
muscle of interest and the synergistic and antagonistic muscles crossing the ankle. The experimental
force trace measured in the TA datasets may be an acceptable validation metric as the TA muscle is
expected to explain most of the recorded ankle torque in dorsiflexion. The TA muscle is indeed the
main dorsiflexor muscle with a muscle volume and maximum isometric force larger than the flexor
hallucis longus muscle, which moment arm is besides not aligned with the dorsiflexion direction and
mainly acts for ankle inversion. However, the gastrocnemius lateralis, soleus, and peroneus muscles
acts synergistically with the GM muscle for ankle plantarflexion and the experimental torque recorded
in dataset $HGM_{30}$ may not be representative of the GM muscle force-generation activity and may not
be a suitable validation metric for $eND_N$. In Figure 13, the decreasing $eND_N$ and $eND_{N_r}$ (orange and
blue dotted traces) may accurately describe a gradually increasing sharing of the ankle torque between
synergistic muscles initially taken by the GM muscle, which the constant validation trace does not
capture. To answer such limitations, the predicted $eND_N$ should be validated against a direct measure
of muscle force, which can be performed as in other recent studies [13,14,117] with ultrasound
measurements of the muscle fascicle or of the muscle tendon concurrently obtained with (HD)EMG
recordings of the muscle activity. Finally, these two validations do not provide any indication whether
the parameters calibrated with a trapezoidal force profile would generalize, for the same subject, to
another force trajectory or a trapezoidal force profile up to another force level and provide accurate
predictions of the $N_r$ experimental spike trains and of the $eND_N$. To perform such validation, the
parameter identification in Step 3 (Figure 1) would be overlooked. The $N_r^{new}$ spike trains identified
from the HDEMG signals concurrently measured with the new force profile would be used to derive
the new current input $I^{new}(t)$ (Step 2) to drive a cohort of $N$ MN models (Step 4), the characteristics
of which would be defined by the $IP$ and $S$ distributions (Step 3) derived with the first contraction
profile, to predict the new effective neural drive $eND_N^{new}$. The MN mapping (Step 1) would serve to
identify the MNs in the reconstructed MN populations, the predicted MN spike trains of which should
be compared against the $N_r$ identified $sp_i^{exp}(t)$. Performing this final validation step was however
impossible in this study because there exists, to the authors' knowledge, no open-source datasets of
edited HDEMG signals recorded for the same subject for different force profiles.

Limitation 2 – Sensitivity of the method to input HDEMG data

While the method predicts a list of simulated spike trains $sp_j^{sim}(t)$ and a $eND_N$ that more accurately

describes the MN pool behaviour than the experimental $sp_i^{exp}(t)$ and $eND_{N_r}$, as discussed previously,

the accuracy of these predictions (Figure 12 and Figure 13) remains sensitive to the distribution in the

entire MN pool of the $N_r$ MNs identified experimentally, reported in the third column of Table 2.

Because of the definition of the current input $I(t)$, the $eND_N$ onset is defined by the recruitment time

$tf_{N_1}^1$ of the lowest-threshold MN $N_1$ identified experimentally, as shown in Figure 13, while the

unknown firing behaviour for $t < ft_{N_1}^1$ of the non-identified MNs of rheobase lower than $I_{N_1}^{th}$ is not

captured. This is not an issue in samples of homogeneously distributed MNs like dataset $DTA_{35}$, where

the 9th smallest MN (first 2% of the pool of threshold-increasing MNs) is identified (Figure 5BC), Table

2) and the $eND$ obtained from the 15 first MNs is not predicted during the short time range $[2.1, 2.3]s$,

where the whole muscle builds only 1%MVC (top-left plot in Figure 13). However in heterogeneous or

incomplete samples like dataset $HTA_{50}$, the lowest-threshold (232nd, Table 2) identified MN identified

is recruited at $t = tf_{N_1}^1 = 6.1s$ and the approach wrongly returns a zero muscle neural drive $eND = 0$

for the time period $[1.6; 6.1]s$ where the muscle actually builds 20%MVC during the ramp of

contraction (bottom-left plot in Figure 13). To tackle this issue, the normalized force trace, which is

non-zero for $t < ft_{N_1}^1$ and representative of the CSI in this time region, could be scaled and used in

lieu of the current definition of $I(t)$. However, this approach, which may not be suitable for non-

isometric conditions, is not applicable in forward predictions of unknown whole muscle force from

neural inputs. Experimental samples of homogeneously distributed MNs are also required to derive

realistic $S(j)$ and $IP(j)$ distributions with the four-step approach. As observed in Figure 12, ignoring in

the derivation of $I(t), IP(j)$ and $S(j)$ the spike trains $sp_i^{exp}(t)$ of the MNs that are representative of

a large subset of the MN pool affects the accuracy of the predicted recruitment and firing behaviour

of the MNs falling in that subset (Figure 12). More specifically, the non-physiological distribution

$IP(j)[s] = 5.6 \cdot 10^{-4} \cdot j^{0.80}$ was fitted to the experimental data of the 14 high-threshold MNs

identified in dataset $HTA_{50}$, where the neural information of the 60% smallest MNs of the MN pool

(Table 2) is lacking. Such $IP(j)$ distribution overestimates the LIF-predicted maximum firing frequency

of low-threshold MNs, which explains the overestimation of the $eND_N$ in Figure 13 (bottom-left plot,

orange curve). Non-physiological predictions can be avoided by adding artificial data points consistent

with other experiments or with the literature in the rheobase regions of the MN pool where no MNs

were experimentally identified. For example, using an $IP(j)$ relationship consistent with a dataset of

homogeneously distributed MNs ($DTA_{35}$) constrained the predicted maximum firing rates to

physiological values and returned more accurate predictions of the $eND_N$ (bottom-left plot, purple

curve).

Limitation 3 – LIF MN modelling

The LIF MN model described from (EQ. 7) to (EQ. 13) was shown, with credible sets of inter-related

parameters $S_{MN}, R, C$ and $\tau$ after [42], to accurately mimic (Figure 9) and blindly predict (Figure 12)

most of the key phenomenological features of the $N_r$ firing MNs, including their FIDFs and time of first

discharge, as well as nonlinear behaviours such as firing rate saturation (Figure 8) and hysteresis (Figure

4B and C). However, this MN model is mostly phenomenological and does not provide a realistic

description of the actual mechanisms underlying the dynamics of action potential firing for several

reasons. First, this single-compartment approach neglects the activity of the MN dendrites, which

account for more than 95% of the total MN surface area and gather most of the post-synaptic receptors

and MN PIC-generating channels responsible for the MN nonlinear input-output functions [40]. The

inherent difference in membrane voltage dynamics between the dendrites and the soma, also partially

mediated by hundred-fold differences in the value of passive electrophysiological parameters such as

$R$ which increases with somatofugal distance [126] is therefore neglected in this point model approach.

Then, the nonlinear dendritic activity being overlooked, the Common Synaptic Input CSI(t) in (EQ. 5),

which is the common net excitatory synaptic influx, is non-physiologically assimilated with a constant

gain to the post-synaptic total dendritic membrane current $I(t)$ that depolarizes the MN soma and is

responsible for spike generation. With this linear $CSI - I$ scaling in (EQ. 6), the approach neglects the

realistic description of the voltage-driven dynamics of the PIC-generating channels that are responsible

for some important nonlinear mechanisms, such as firing rate saturation [41] and hysteresis in the

MN's current-voltage relation [94,95], which dictate the nonlinear $CSI - I$ transformation [40]. In this

study, the firing rate saturation, which in real MNs is due to a decrease in driving force for synaptic

current flow as the dendrites become more depolarized [41], is captured by the phenomenological $IP$

parameter. The observed hysteresis between the recruitment and decruitment thresholds (Figure 4B)

and in the current-firing rate relation between recruitment and derecruitment phases (Figure 4C),

explained by an hysteresis in the MN's current-voltage relation that arises from the activity of long-

lasting PICs [20], are successfully but non-physiologically addressed by a tuning during derecruitment

of the $R$ and $C_m$ parameters respectively, although the values of the passive property $R$ and of $C_m$,

biological constant for the MN population [127], should be independent from the MN activity [44]. For

these reasons, even if the distributions of the $S_{MN}, R, C$ and $\tau$ parameters at recruitment take values

consistent with the literature, as discussed, the MN LIF model described in (EQ. 7) to (EQ. 12) is not

suitable for investigating the neurophysiology and the neuromechanics of individual MNs. The

phenomenological tuning of the $IP, R$ and $C_m$ parameters, although symptomatic of underlying

nonlinear mechanisms, does not provide any insight into the true MN neurophysiology. If a more

realistic MN model was considered in lieu of the LIF model, such as multiple-compartments Hodgkin-

Huxley-based approaches, the four-step workflow would be a tool for testing neurophysiological

hypotheses and investigating some mechanisms and properties of the complete MN pool that cannot

be directly observed experimentally in conditions of human voluntary contractions. A possible trade-

off would be considering a single-compartment LIF model with active conductances [44], in which case

(EQ. 7) is re-written as (EQ. 18), where $g_R = \frac{1}{R}$ is the constant MN passive conductance, $E_+ > \Delta V_{th}$ is

the constant reversal potential for excitatory channels, $g(t, CSI)$ the time-varying total active

excitatory synaptic conductance, and $I_{ps}(t, CSI, V_m) = g(t, CSI) \cdot (E_+ - V_m)$ the post-synaptic total

dendritic membrane current induced by the synaptic drive $CSI$, the driving force of which is decreased

by the term $(E_+ - V_m)$ when the dendritic membrane depolarizes, as discussed.

$$\frac{dV_m}{dt} = \frac{1}{C} \cdot [g(t, CSI) \cdot (E_+ - V_m) - g_R \cdot V_m] \qquad \text{(EQ. 18)}$$

In the literature, $g(t, CSI) = g_{max} \cdot T(t, CSI)$ where $g_{max}$ is the maximum active conductance of the

synapse and $T(t, CSI)$ can be interpreted as the fraction of bound synaptic receptors [30] or of opened

ionic channels in the range $[0; 1]$ or as a train of synaptic pulses [44]. Because setting $T(t, CSI) =$

$CSI(t)$, with $CSI(t)$ as defined in Figure 6B, does not meet the requirements in firing rate saturation

for large CSI input, the saturation in dendritic ionic channel activation for large CSI [41] must be

accounted for and $T(t, CSI)$ should be a nonlinear saturating function of CSI. In this approach, the MN-

specific hysteresis and firing rate behaviour could be obtained with a tuning during recruitment and

derecruitment and a distribution across the MN pool of the $g_{max}$ parameter and of the shape

parameter of the $S$ function based on the experimental data. Although this approach more realistically

describes the neurophysiological mechanisms underlying the nonlinear MN behaviour during

discharge events, it is more challenging to scale using the experimental information. Considering that

this study focuses on the phenomenological behaviour of individual MNs and on the overall activity of

reconstructed MN populations, the LIF model defined in (EQ. 7) was judged an adequate trade-off

between accuracy and modelling complexity, considering the overall accurate predictions of the MN

firing behaviours (Figure 12) and its low computational cost, and no other modelling approaches such

as (EQ. 18) were pursued.

It must be noted that, whatever the MN modelling approach, the experimental drive to the MN

currently considered in the four-step method is the CSI, which disregards the MN-specific synaptic

noise $SN(t)$, which is responsible for most of the inter-spike variability (ISV). Considering that the MN

pool and the MU neuromechanical mechanisms are expected to filter the MN-specific $SN(t)$, this

simplification is adequate for accurate predictions of the $eND_N$ and of the whole muscle force (Farina,

2015). Sometimes modelled as a random Gaussian-like event (Fuglevand 1993), the ISV can be

obtained as the response to a $SN(t)$ white noise added to the current definition of the synaptic input

as $CSI(t) + SN(t)$. The relative proportion of $CSI$ and $SN$ in % of the variance of the total synaptic

current can be approximated from Figure 2A in [47]. Accounting for $SN(t)$ might improve the accuracy

of the predicted firing behaviour of the largest MNs, for which the largest $\Delta ft^1$ and $nRMSE$ and lowest

$r^2$ values were obtained in Figure 12 for all four datasets, and the firing behaviour and recruitment

dynamics of which are dominantly dictated by random fluctuations of $SN(t)$.

1046

Limitation 4 – Limited experimental data in the literature

The four-step approach is constrained by the limited knowledge in the literature of the characteristics

of the human MU pool. Therefore, the LIF parameters and the MN rheobase distribution in the MN

pool (Figure 7A) are tuned with typical cat data from various hindlimb muscles [42] while the

experimental MN spike trains were obtained from the human TA muscle. While the normalized

mathematical relationships relating the MN electrophysiological parameters of the LIF model

$(R, C, \tau, \Delta V_{th})$ can be extrapolated between mammals [42] and thus to humans, no experimental data

is yet available to scale these relationships to typical human data.

Also, the typical $F^{th}(j)$ and $I_{th}(j)$ threshold distributions, derived for mapping the $N_r$ identified MNs

to the complete MN pool (Figure 5) and for scaling the $CSI$ to physiological values of $I(t)$ (Figure 7),

were obtained from studies which relied on experimental samples of MN populations of small size.

These source studies therefore did not ensure that the largest and lowest values were identified and

reported, and that the identification process was not biased towards a specific subset of the MN pool,

such as larger MNs. In such cases, the true threshold distributions would be more skewed and spread

over a larger range of values, as discussed in [42], than the distributions reported in Figure 5 and Figure

7. The $F^{th}(j)$ distribution is besides muscle-specific [20] with large hindlimb muscles being for example

recruited over a larger range of MVC than hand muscles. However, enough data is reported in the

literature to build the $F^{th}(j)$ distributions for the TA and first dorsal interossei human muscles only.

For these limitations, the two first steps of this approach could be made subject- species- and muscle-

specific by calibrating the $F^{th}(j)$ and $I^{th}(j)$ described as the 3-parameter power functions defined in

this study.

1068

Conclusion

This study presents a four-step workflow (Figure 1 and Figure 11) which predicts the spiking activity of

the discharging MNs that were not identified by decomposed HDEMG signals. The method is driven by

the common synaptic input, which is derived from the experimental data, and reconstructs, after a

calibration step, the distribution across the MN population of some MN properties involved into the

MN-specific recruitment and spiking behaviour of the discharging MNs (Figure 8 and Figure 10A). The

method blindly predicts the discharge behaviour of the $N_r$ experimentally identified MNs (Figure 12)

and accurately predicts the muscle neural drive (Figure 13) after the complete discharging MN

population is reconstructed (Figure 10C and D). With direct application in neuromuscular modelling

(Figure 14), this method addresses the bias of HDEMG identification towards high-threshold large units

and is relevant for neuroscientists, modelers and experimenters to investigate the MN pool dynamics

during force generation.

1081

REFERENCES

[1] Felici F, Del Vecchio A. Surface Electromyography: What Limits Its Use in Exercise and Sport
Physiology? *Frontiers in neurology.* 2020; 11 578504.

[2] Watanabe K, Vieira TM, Gallina A, Kouzaki M, Moritani T. Novel Insights Into Biarticular Muscle
Actions Gained From High-Density Electromyogram. *Exercise and sport sciences reviews.* 2021; 49 (3):
179-187.

[3] Maeda N, Komiya M, Nishikawa Y, Morikawa M, Tsutsumi S, Tashiro T, et al. Effect of Acute Static
Stretching on the Activation Patterns Using High-Density Surface Electromyography of the
Gastrocnemius Muscle during Ramp-Up Task. *Sensors (Basel, Switzerland).* 2021; 21 (14): 4841.

[4] Jordanić M, Rojas-Martínez M, Mañanas MA, Alonso JF. Prediction of isometric motor tasks and
effort levels based on high-density EMG in patients with incomplete spinal cord injury. *Journal of*
*Neural Engineering.* 2016; 13 (4): 046002.

[5] Fang C, He B, Wang Y, Cao J, Gao S. EMG-Centered Multisensory Based Technologies for Pattern
Recognition in Rehabilitation: State of the Art and Challenges. *Biosensors.* 2020; 10 (8): 85.

[6] Pilkar R, Momeni K, Ramanujam A, Ravi M, Garbarini E, Forrest GF. Use of Surface EMG in Clinical
Rehabilitation of Individuals With SCI: Barriers and Future Considerations. *Frontiers in Neurology.*
2020; 11 1680.

[7] Kisiel-Sajewicz K, Marusiak J, Rojas-Martínez M, Janecki D, Chomiak S, Kamiński Ł, et al. High-
density surface electromyography maps after computer-aided training in individual with congenital
transverse deficiency: a case study. *BMC musculoskeletal disorders.* 2020; 21 (1): 682.

[8] Nishikawa Y, Watanabe K, Holobar A, Maeda N, Maruyama H, Tanaka S. Identification of the
laterality of motor unit behavior in female patients with parkinson's disease using high-density
surface electromyography. *European Journal of Neuroscience.* 2021; 53 (6): 1938-1949.

[9] Callahan DM, Umberger BR, Kent-Braun JA. A computational model of torque generation: neural,
contractile, metabolic and musculoskeletal components. *PloS one.* 2013; 8 (2): e56013.

[10] Carriou V, Boudaoud S, Laforet J, Mendes A, Canon F, Guiraud D. Multiscale Hill-type modeling of
the mechanical muscle behavior driven by the neural drive in isometric conditions. *Computers in*
*biology and medicine.* 2019; 115 103480.

[11] Kim H, Kim M. PyMUS: python-based simulation software for virtual experiments on motor unit
system. *Frontiers in neuroinformatics.* 2018; 12 15.

[12] Potvin JR, Fuglevand AJ. A motor unit-based model of muscle fatigue. *PLoS computational*
*biology.* 2017; 13 (6): e1005581.

[13] Waasdorp R, Mugge W, Vos HJ, De Groot JH, Verweij MD, De Jong N, et al. Combining ultrafast
ultrasound and high-density EMG to assess local electromechanical muscle dynamics: a feasibility
study. *IEEE access.* 2021; 9 1.

[14] Martinez-Valdes E, Negro F, Botter A, Pincheira PA, Cerone GL, Falla D, et al. *Modulations in motor unit discharge are related to changes in fascicle length during isometric contractions.* Cold Spring Harbor Laboratory; 2021.

[15] Chen C, Yu Y, Ma S, Sheng X, Lin C, Farina D, et al. Hand gesture recognition based on motor unit spike trains decoded from high-density electromyography. *Biomedical signal processing and control.* 2020; 55 101637.

[16] Farina D, Vujaklija I, Brånemark R, Bull AMJ, Dietl H, Graimann B, et al. Toward higher-performance bionic limbs for wider clinical use. *Nature biomedical engineering.* 2021; 1-13.

[17] Farina D, Vujaklija I, Sartori M, Kapelner T, Negro F, Jiang N, et al. Man/machine interface based on the discharge timings of spinal motor neurons after targeted muscle reinnervation. *Nature biomedical engineering.* 2017; 1 (2): 1-12.

[18] Freund HJ, Budingen HJ, Dietz V. Activity of single motor units from human forearm muscles during voluntary isometric contractions. *J Neurophysiol.* 1975; 38 (4): 933-946.

[19] Dengler R, Stein RB, Thomas CK. Axonal conduction velocity and force of single human motor units. *Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine.* 1988; 11 (2): 136-145.

[20] Heckman CJ, Enoka RM. Motor Unit. *Compr. Physiol.* 2012; 2 (4): 2629-2682.

[21] Manuel M, Chardon M, Tysseling V, Heckman CJ. Scaling of Motor Output, From Mouse to Humans. *Physiology (Bethesda, Md.).* 2019; 34 (1): 5-13.

[22] Highlander MM, Allen JM, Elbasiouny SM. *Meta-analysis of biological variables' impact on spinal motoneuron electrophysiology data.* American Physiological Society; 2020.

[23] Holobar A, Farina D. Blind source identification from the multichannel surface electromyogram. *Physiological measurement.* 2014; 35 (7): R143-R165.

[24] Negro F, Muceli S, Castronovo AM, Holobar A, Farina D. Multi-channel intramuscular and surface EMG decomposition by convolutive blind source separation. *Journal of neural engineering.* 2016; 13 (2): 026027.

[25] Oliveira AS, Negro F. NEURAL CONTROL OF MATCHED MOTOR UNITS DURING MUSCLE SHORTENING AND LENGTHENING AT INCREASING VELOCITIES. *Journal of applied physiology (1985).* 2021; 130 (6): 1798-1813.

[26] Del Vecchio A, Casolo A, Negro F, Scorcelletti M, Bazzucchi I, Enoka R, et al. The increase in muscle force after 4 weeks of strength training is mediated by adaptations in motor unit recruitment and rate coding. *J Physiol.* 2019; 597 (7): 1873-1887.

[27] Del Vecchio A, Holobar A, Falla D, Felici F, Enoka RM, Farina D. Tutorial: analysis of motor unit discharge characteristics from high-density surface EMG signals. *Journal of Electromyography and Kinesiology.* 2020; 53 102426.

[28] Dideriksen JL, Farina D, Baekgaard M, Enoka RM. An integrative model of motor unit activity during sustained submaximal contractions. *Journal of Applied Physiology.* 2010; 108 (6): 1550-1562.

[29] Fuglevand AJ, Winter DA, Patla AE. Models of recruitment and rate coding organization in
motor-unit pools. *Journal of neurophysiology.* 1993; 70 (6): 2470-2488.

[30] Cisi RRL, Kohn AF. Simulation system of spinal cord motor nuclei and associated nerves and
muscles, in a Web-based architecture. *Journal of computational neuroscience.* 2008; 25 (3): 520-542.

[31] Farina D, Negro F, Dideriksen JL. The effective neural drive to muscles is the common synaptic
input to motor neurons. *The Journal of physiology.* 2014; 592 (16): 3427-3441.

[32] Negro F, Farina D. Decorrelation of cortical inputs and motoneuron output. *Journal of
Neurophysiology.* 2011; 106 (5): 2688-2697.

[33] Watanabe RN, Magalhães FH, Elias LA, Chaud VM, Mello EM, Kohn AF. Influences of
premotoneuronal command statistics on the scaling of motor output variability during isometric
plantar flexion. *Journal of Neurophysiology.* 2013; 110 (11): 2592-2606.

[34] Dideriksen JL, Enoka RM, Farina D. Neuromuscular adjustments that constrain submaximal EMG
amplitude at task failure of sustained isometric contractions. *Journal of Applied Physiology.* 2011; 111
(2): 485-494.

[35] Röhrle O, Yavuz U, Klotz T, Negro F, Heidlauf T. Multiscale modeling of the neuromuscular
system: Coupling neurophysiology and skeletal muscle mechanics. *Wiley interdisciplinary reviews.
Systems biology and medicine.* 2019; 11 (6): e1457.

[36] Fohlmeister JF. A nerve model of greatly increased energy-efficiency and encoding flexibility
over the Hodgkin–Huxley model. *Brain research.* 2009; 1296 225-233.

[37] Bondarenko VE, Szigeti GP, Bett GC, Kim S, Rasmusson RL. Computer model of action potential of
mouse ventricular myocytes. *American Journal of Physiology-Heart and Circulatory Physiology.* 2004;
287 (3): H1378-H1403.

[38] Smit JE, Hanekom T, Hanekom JJ. Predicting action potential characteristics of human auditory
nerve fibres through modification of the Hodgkin-Huxley equations. *South African Journal of Science.*
2008; 104 (7): 284-292.

[39] Raikova R, Celichowski J, Angelova S, Krutki P. A model of the rat medial gastrocnemius muscle
based on inputs to motoneurons and on an algorithm for prediction of the motor unit force. *Journal
of neurophysiology.* 2018; 120 (4): 1973-1987.

[40] Binder MD, Powers RK, Heckman CJ. Nonlinear input-output functions of motoneurons.
*Physiology.* 2020; 35 (1): 31-39.

[41] Powers RK, Heckman CJ. Synaptic control of the shape of the motoneuron pool input-output
function. *Journal of neurophysiology.* 2017; 117 (3): 1171-1184.

[42] Caillet AH, Phillips AT, Farina D, Modenese L. Mathematical relationships between spinal
motoneuron properties. *eLife.* 2022; 11 e76489.

[43] Teeter C, Iyer R, Menon V, Gouwens N, Feng D, Berg J, et al. Generalized leaky integrate-and-fire
models classify multiple neuron types. *Nat. Commun.* 2018; 9 (1): 1-15.

[44] Burkitt AN. A review of the integrate-and-fire neuron model: I. Homogeneous synaptic input.
*Biological cybernetics.* 2006; 95 (1): 1-19.

[45] Izhikevich EM. Which model to use for cortical spiking neurons? *IEEE Transactions on Neural*
*Networks.* 2004; 15 (5): 1063-1070.

[46] Dong Y, Mihalas S, Russell A, Etienne-Cummings R, Niebur E. Estimating parameters of
generalized integrate-and-fire neurons from the maximum likelihood of spike trains. *Neural*
*computation.* 2011; 23 (11): 2833-2867.

[47] Negro F, Yavuz U, Farina D. The human motor neuron pools receive a dominant slow-varying
common synaptic input. *J. Physiol.* 2016; 594 (19): 5491-5505.

[48] Henneman E. Relation between Size of Neurons and Their Susceptibility to Discharge. *Science.*
1957; 126 (3287): 1345-1347.

[49] Henneman E, Somjen G, Carpenter DO. Excitability and inhibitibility of motoneurons of different
sizes. *J Neurophysiol.* 1965; 28 (3): 599-620.

[50] Henneman E, Somjen G, Carpenter DO. Functional significance of cell size in spinal motoneurons.
*J Neurophysiol.* 1965; 28 (3): 560-580.

[51] Henneman E, Clamann HP, Gillies JD, Skinner RD. Rank order of motoneurons within a pool: law
of combination. *J Neurophysiol.* 1974; 37 (6): 1338-1349.

[52] Henneman E. Recruitment of motoneurons : The size principle. *Motor Unit Types, Recruitment*
*and Plasticity in Health and Disease.* 1981; 26-60.

[53] Henneman E. The size-principle: a deterministic output emerges from a set of probabilistic
connections. *J. Exp. Biol.* 1985; 115 (1): 105-112.

[54] Hug F, Avrillon S, Del Vecchio A, Casolo A, Ibanez J, Nuccio S, et al. Analysis of motor unit spike
trains estimated from high-density surface electromyography is highly reliable across operators.
*Journal of Electromyography and Kinesiology.* 2021; 58 102548.

[55] Hug F, Del Vecchio A, Avrillon S, Farina D, Tucker K. Muscles from the same muscle group do not
necessarily share common drive: evidence from the human triceps surae. *Journal of applied*
*physiology.* 2021; 130 (2): 342-354.

[56] De Luca CJ, Lefever RS, Mccue MP, Xenakis AP. Behaviour of human motor units in different
muscles during linearly varying contractions. *J. Physiol.* 1982; 329 (1): 113-128.

[57] Feinstein B, Lindegård B, Nyman E, Wohlfart G. Morphologic studies of motor units in normal
human muscles. *Cells Tissues Organs.* 1955; 23 (2): 127-142.

[58] Trojaborg W, Kaufmann P, Gooch CL. Motor unit estimate number in the anterior tibial muscle:
normative data versus findings in critically ill patients in intensive care units. *Journal of clinical*
*neuromuscular disease.* 2002; 3 (4): 139-142.

[59] Boe SG, Dalton BH, Harwood B, Doherty TJ, Rice CL. Inter-rater reliability of motor unit number
estimates and quantitative motor unit analysis in the tibialis anterior muscle. *Clinical*
*neurophysiology.* 2009; 120 (5): 947-952.

[60] Power GA, Dalton BH, Behm DG, Vandervoort AA, Doherty TJ, Rice CL. Motor unit number
estimates in masters runners: use it or lose it. *Med Sci Sports Exerc.* 2010; 42 (9): 1644-1650.

[61] Piasecki M, Ireland A, Coulson J, Stashuk DW, Hamilton-Wright A, Swiecicka A, et al. Motor unit
number estimates and neuromuscular transmission in the tibialis anterior of master athletes:
evidence that athletic older people are not spared from age-related motor unit remodeling.
*Physiological reports.* 2016; 4 (19): e12987.

[62] Hourigan ML, McKinnon NB, Johnson M, Rice CL, Stashuk DW, Doherty TJ. Increased motor unit
potential shape variability across consecutive motor unit discharges in the tibialis anterior and vastus
medialis muscles of healthy older subjects. *Clinical Neurophysiology.* 2015; 126 (12): 2381-2389.

[63] McNeil CJ, Doherty TJ, Stashuk DW, Rice CL. Motor unit number estimates in the tibialis anterior
muscle of young, old, and very old men. *Muscle & Nerve: Official Journal of the American Association*
*of Electrodiagnostic Medicine.* 2005; 31 (4): 461-467.

[64] Van Cutsem M, Feiereisen P, Duchateau J, Hainaut K. Mechanical properties and behaviour of
motor units in the tibialis anterior during voluntary contractions. *Can. J. Appl. Physiol.* 1997; 22 (6):
585-597.

[65] Xiong GX, Zhang JW, Hong Y, Guan Y, Guan H. Motor unit number estimation of the tibialis
anterior muscle in spinal cord injury. *Spinal Cord.* 2008; 46 (10): 696-702.

[66] Gooch CL, Doherty TJ, Chan KM, Bromberg MB, Lewis RA, Stashuk DW, et al. Motor unit number
estimation: a technology and literature review. *Muscle & nerve.* 2014; 50 (6): 884-893.

[67] Henriksson-Larsén KB, Lexell J, Sjöström M. Distribution of different fibre types in human
skeletal muscles. I. Method for the preparation and analysis of cross-sections of whole tibialis
anterior. *The Histochemical journal.* 1983; 15 (2): 167-178.

[68] Henriksson-Larsén K. Distribution, number and size of different types of fibres in whole cross-
sections of female m tibialis anterior. An enzyme histochemical study. *Acta Physiologica*
*Scandinavica.* 1985; 123 (3): 229-235.

[69] Christensen E. American Journal of Physical Medicine & Rehabilitation. *American Journal of*
*Physical Medicine & Rehabilitation.* 1959; 38 (2): 65-78.

[70] Handsfield GG, Meyer CH, Hart JM, Abel MF, Blemker SS. Relationships of 35 lower limb muscles
to height and body mass quantified using MRI. *Journal of Biomechanics.* 2014; 47 (3): 631-638.

[71] Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour contribute to the
increase in contraction speed after dynamic training in humans. *The Journal of physiology.* 1998; 513
(1): 295-305.

[72] Feiereisen P, Duchateau J, Hainaut K. Motor unit recruitment order during voluntary and
electrically induced contractions in the tibialis anterior. *Experimental brain research.* 1997; (114(1)):
117.123.

[73] Desmedt JE, Godaux E. Ballistic contractions in man: characteristic recruitment pattern of single
motor units of the tibialis anterior muscle. *The Journal of physiology.* 1977; 264 (3): 673-693.

[74] Andreassen S, Arendt-Nielsen L. Muscle fibre conduction velocity in motor units of the human
anterior tibial muscle: a new size principle parameter. *J. Physiol.* 1987; 391 (1): 561-571.

[75] Vander Linden DW, Kukulka CG, Soderberg GL. The effect of muscle length on motor unit
discharge characteristics in human tibialis anterior muscle. *Experimental brain research.* 1991; 84 (1):
210-218.

[76] Connelly DM, Rice CL, Roos MR, Vandervoort AA. Motor unit firing rates and contractile
properties in tibialis anterior of young and old men. *Journal of applied physiology.* 1999; 87 (2): 843-
852.

[77] Ankit R. *WebPlotDigitizer.* (4.4) Pacifica, California, USA: 2020.

[78] Garnett RAF, O'Donovan MJ, Stephens JA, Taylor A. Motor unit organization of human medial
gastrocnemius. *J. Physiol.* 1979; 287 (1): 33-43.

[79] Vieira TM, Loram ID, Muceli S, Merletti R, Farina D. Recruitment of motor units in the medial
gastrocnemius muscle during human quiet standing: is recruitment intermittent? What triggers
recruitment? *Journal of neurophysiology.* 2012; 107 (2): 666-676.

[80] Kallio J, Søgaard K, Avela J, Komi PV, Selänne H, Linnamo V. Motor unit firing behaviour of soleus
muscle in isometric and dynamic contractions. *PloS one.* 2013; 8 (2): e53425.

[81] Ballantyne BT, Kukulka CG, Soderberg GL. Motor unit recruitment in human medial
gastrocnemius muscle during combined knee flexion and plantarflexion isometric contractions.
*Experimental brain research.* 1993; 93 (3): 492-498.

[82] Héroux ME, Dakin CJ, Luu BL, Inglis JT, Blouin J. Absence of lateral gastrocnemius activity and
differential motor unit behavior in soleus and medial gastrocnemius during standing balance. *Journal
of applied physiology.* 2014; 116 (2): 140-148.

[83] Farina D, Negro F. Common synaptic input to motor neurons, motor unit synchronization, and
force control. *Exercise and sport sciences reviews.* 2015; 43 (1): 23-33.

[84] Krawitz S, Fedirchuk B, Dai Y, Jordan LM, McCrea DA. State-dependent hyperpolarization of
voltage threshold enhances motoneurone excitability during fictive locomotion in the cat. *The
Journal of physiology.* 2001; 532 (1): 271-281.

[85] Gustafsson, Pinter MJ. An investigation of threshold properties among cat spinal alpha-
motoneurones. *J. Physiol.* 1984; 357 (1): 453-483.

[86] Kernell D. Input Resistance, Electrical Excitability, and Size of Ventral Horn Cells in Cat Spinal
Cord. *Science.* 1966; 152 (3729): 1637-1640.

[87] Zengel JE, Reid SA, Sypert GW, Munson JB. Membrane electrical properties and prediction of
motor-unit type of medial gastrocnemius motoneurons in the cat. *J Neurophysiol.* 1985; 53 (5): 1323-
1344.

[88] Fleshman JW, Munson JB, Sypert GW, Friedman WA. Rheobase, input resistance, and motor-unit
type in medial gastrocnemius motoneurons in the cat. *J Neurophysiol.* 1981; 46 (6): 1326-1338.

[89] Bakels R, Kernell D. Matching between motoneurone and muscle unit properties in rat medial
gastrocnemius. *J. Physiol.* 1993; 463 (1): 307-324.

[90] Kernell D, Monster AW. Threshold current for repetitive impulse firing in motoneurones
innervating muscle fibres of different fatigue sensitivity in the cat. *Brain Res.* 1981; 229 (1): 193-196.

[91] Foehring RC, Sypert GW, Munson JB. Motor-unit properties following cross-reinnervation of cat
lateral gastrocnemius and soleus muscles with medial gastrocnemius nerve. II. Influence of muscle on
motoneurons. *J Neurophysiol.* 1987; 57 (4): 1227-1245.

[92] Gardiner PF. Physiological properties of motoneurons innervating different muscle unit types in
rat gastrocnemius. *J Neurophysiol.* 1993; 69 (4): 1160-1170.

[93] Caillet A, Phillips A, Farina D, Modenese L. Mathematical Relationships between Spinal
Motoneuron Properties. *bioRxiv.* 2021; .

[94] Lee RH, Heckman CJ. Bistability in Spinal Motoneurons In Vivo: Systematic Variations in
Persistent Inward Currents. *Journal of neurophysiology.* 1998; 80 (2): 583-593.

[95] Lee RH, Heckman CJ. Bistability in spinal motoneurons in vivo: systematic variations in rhythmic
firing patterns. *Journal of neurophysiology.* 1998; 80 (2): 572-582.

[96] Lloyd DG, Besier TF. An EMG-driven musculoskeletal model to estimate muscle forces and knee
joint moments in vivo. *Journal of Biomechanics.* 2003; 36 (6): 765-776.

[97] Seth A, Hicks JL, Uchida TK, Habib A, Dembia CL, Dunne JJ, et al. OpenSim: Simulating
musculoskeletal dynamics and neuromuscular control to study human and animal movement. *PLoS*
*computational biology.* 2018; 14 (7): e1006223.

[98] Rajagopal A, Dembia CL, DeMers MS, Delp DD, Hicks JL, Delp SL. Full-Body Musculoskeletal
Model for Muscle-Driven Simulation of Human Gait. *IEEE Transactions on Biomedical Engineering.*
2016; 63 (10): 2068-2079.

[99] Hatze H. *Myocybernetic control models of skeletal muscle - characteristics and applications.* b
ed.: University of South Africa Press, Pretoria; 1980.

[100] Foehring RC, Sypert GW, Munson JB. Properties of self-reinnervated motor units of medial
gastrocnemius of cat. II. Axotomized motoneurons and time course of recovery. *J Neurophysiol.*
1986; 55 (5): 947-965.

[101] Munson JB, Foehring RC, Lofton SA, Zengel JE, Sypert GW. Plasticity of medial gastrocnemius
motor units following cordotomy in the cat. *Journal of neurophysiology.* 1986; 55 (6): 1454.

[102] Kernell D, Zwaagstra B. Dendrites of cat's spinal motoneurones: relationship between stem
diameter and predicted input conductance. *J. Physiol.* 1989; 413 (1): 255-269.

[103] Burke RE, Dum RP, Fleshman JW, Glenn LL, Lev-Tov A, O'Donovan MJ, et al. An HRP study of the
relation between cell size and motor unit type in cat ankle extensor motoneurons. *J. Comp. Neurol.*
1982; 209 (1): 17-28.

[104] Ulfhake, Kellerth JO. A quantitative light microscopic study of the dendrites of cat spinal α-
motoneurons after intracellular staining with horseradish peroxidase. *J. Comp. Neurol.* 1981; 202 (4):
571-583.

[105] Ulfhake B, Kellerth JO. Electrophysiological and morphological measurements in cat
gastrocnemius and soleus α-motoneurones. *Brain Res.* 1984; 307 (1): 167-179.

[106] Ulfhake B, Cullheim S. Postnatal development of cat hind limb motoneurons. III: Changes in size
of motoneurons supplying the triceps surae muscle. *Journal of comparative neurology (1911).* 1988;
278 (1): 103-120.

[107] Barrett JN, Crill WE. Specific membrane properties of cat motoneurones. *J. Physiol.* 1974; 239
(2): 301-324.

[108] Gustafsson, Pinter MJ. Relations among passive electrical properties of lumbar alpha-
motoneurones of the cat. *J. Physiol.* 1984; 356 (1): 401-431.

[109] De Luca CJ, Hostage EC. Relationship between firing rate and recruitment threshold of
motoneurons in voluntary isometric contractions. *Journal of neurophysiology.* 2010; 104 (2): 1034-
1046.

[110] Kapelner T, Sartori M, Negro F, Farina D. Neuro-musculoskeletal mapping for man-machine
interfacing. *Scientific reports.* 2020; 10 (1): 1-10.

[111] Thompson CK, Negro F, Johnson MD, Holmes MR, McPherson LM, Powers RK, et al. Robust and
accurate decoding of motoneuron behaviour and prediction of the resulting force output. *The
Journal of physiology.* 2018; 596 (14): 2643-2659.

[112] Huxley AF, Niedergerke R. Structural changes in muscle during contraction: interference
microscopy of living muscle fibres. *Nature.* 1954; 173 (4412): 971-973.

[113] Huxley AF. Muscle structure and theories of contraction. *Prog.Biophys.Biophys.Chem.* 1957; 7
255-318.

[114] Hof AL, Van den Berg J. EMG to force processing I: An electrical analogue of the hill muscle
model. *Journal of biomechanics.* 1981; 14 (11): 747,755-753,758.

[115] Zajac Felix E. Muscle and Tendon: Properties, Models, Scaling, And Application To Biomechanics
And Motor Control. *Critical reviews in biomedical engineering.* 1989; 17 (4): 359-411.

[116] Sartori M, Yavuz U, Farina D. In Vivo Neuromechanics: Decoding Causal Motor Neuron Behavior
with Resulting Musculoskeletal Function. *Scientific reports.* 2017; 7 (1): 13465-14.

[117] Dick TJM, Biewener AA, Wakeling JM. Comparison of human gastrocnemius forces predicted by
Hill-type muscle models and estimated from ultrasound images. *Journal of experimental biology.*
2017; 220 (Pt 9): 1643-1653.

[118] Hatze H. A myocybernetic control model of skeletal muscle. *Biological cybernetics.* 1977; 25 (2):
103-119.

[119] Hussein M, Shebl S, Elnemr R, Elkaranshawy H. A New Muscle Activation Dynamics Model, That
Simulates the Calcium Kinetics and Incorporates the Role of Store-Operated Calcium Entry Channels,
to Enhance the Electromyography-Driven Hill-Type Models. *Journal of Biomechanical Engineering.*
2021; 144 (1): 011002.

[120] Dorgan SJ, O'Malley MJ. A nonlinear mathematical model of electrically stimulated skeletal
muscle. *IEEE transactions on rehabilitation engineering.* 1997; 5 (2): 179-194.

[121] Song D, Raphael G, Lan N, Loeb GE. Computationally efficient models of neuromuscular
recruitment and mechanics. *Journal of neural engineering.* 2008; 5 (2): 175-184.

[122] Hamouda A, Kenney L, Howard D. Dealing with time-varying recruitment and length in Hill-type
muscle models. *Journal of biomechanics.* 2016; 49 (14): 3375-3380.

[123] Legreneur P, Morlon B, Van Hoecke J. Simulation of in situ soleus isometric force output as a
function of neural excitation. *Journal of biomechanics.* 1996; 29 (11): 1455-1462.

[124] Sartori M, Reggiani M, Farina D, Lloyd DG. EMG-Driven Forward-Dynamic Estimation of Muscle
Force and Joint Moment about Multiple Degrees of Freedom in the Human Lower Extremity. *PloS*
*one.* 2012; 7 (12): e52618.

[125] Pizzolato C, Lloyd DG, Sartori M, Ceseracciu E, Besier TF, Fregly BJ, et al. CEINMS: A toolbox to
investigate the influence of different neural control solutions on the prediction of muscle excitation
and joint moments during dynamic motor tasks. *Journal of Biomechanics.* 2015; 48 (14): 3929-3936.

[126] Fleshman JW, Segev I, Cullheim S, Burke RE. Matching electrophysiological with morphological
measurements in cat α-motoneurons. *Soc. Neurosci. Abstr,* ; 1983. pp. 431.

[127] Gentet L, Stuart GJ, Clements JD. Direct Measurement of Specific Membrane Capacitance in
Neurons. *Biophys. J.* 2000; 79 (1): 314-320.

1392