1 **Antibacterial effects of propolis and brood comb extracts on the causative agent of**
2 **European Foulbrood (*Melissococcus plutonius*) in honey bees (*Apis mellifera*)**
3

4 Stephanie K. Murray[1][*], Colin M. Kurkul[1], Andrew J. Mularo[2], Vanessa L. Hale[3], Rachelle M. M.
5 Adams[2], Reed M. Johnson[1]
6
7 [1]Department of Entomology
8 The Ohio State University
9 Wooster, Ohio 44691
10
11 [2]Department of Evolution, Ecology, and Organismal Biology
12 The Ohio State University
13 Columbus, Ohio 43210
14
15 [3]Department of Veterinary Preventive Medicine
16 The Ohio State University
17 Columbus, Ohio 43210
18
19 *murray.968@osu.edu

20
21 Word count: 5643

Abstract

Among a long list of parasites and pathogens that threaten the European honey bee (*Apis*

mellifera), European Foulbrood (EFB) has become an urgent apiary disease, as epidemic

outbreaks are becoming increasingly common. EFB is a bacterial disease of larval honey bees,

caused by a gram-positive, anaerobic bacterium, *Melissococcus plutonius*. The most effective

current treatment for EFB, oxytetracycline hydrochloride, can disrupt the bee microbiome, cause

bee mortality and residues may persist in honey harvested for human consumption. In this study,

we explore the efficacy of more sustainable bee-derived solutions, including propolis, honey

comb and brood comb ethanol extracts. Using a series of dilutions of these extracts, we

determined the minimum inhibitory concertation (MIC) of each bee-derived product on *M.*

plutonius, as well as two model bacterial species, *Staphylococcus saprophyticus* (gram-positive)

and *Escherichia coli* (gram-negative). Overall, we found that propolis extract was most effective

at inhibiting growth of gram-positive bacteria, and that *M. plutonious* was also susceptible to

honey comb (MIC = 16.00 mg/mL) and brood comb (MIC = 45.33 mg/mL) extracts, but at much

higher concentrations than that of propolis (MIC = 1.14 mg/mL).

Introduction

The social life history of the European honey bee (*Apis mellifera*) is highly conducive to the spread of pathogens, such as foulbrood (i.e., bacterial disease of the larvae). Both American Foulbrood and European Foulbrood diseases, named for the regions in which they were initially discovered (Milbrath, 2021), are present in honey bee colonies worldwide. American Foulbrood (AFB) is typically more devastating, as the causative agent *Paenibacillus larvae* is spore-forming, and spores are known to remain viable on beekeeping equipment for at least 35 years (Haseman, 1961). While antibiotics such as oxytetracycline hydrochloride (OTC) can be effective against *P. larvae* in the vegetative stage, they are ineffective against spores (Lodesani & Costa, 2005). Additionally, tetracycline antibiotics decrease core microbiota of the honey bee gut, resulting in increased susceptibility to opportunistic bacteria and elevated mortality (Raymann et al., 2017; Soares et al., 2021). These changes have the potential to cause long-term effects on colony fitness (Bulson et al., 2021). Because antibiotics may simply mask AFB infection in a hive (Lodesani & Costa, 2005), colonies diagnosed with AFB are often killed and hive equipment destroyed by fire (Hansen & Brødsgaard, 1999), resulting in a substantial economic loss for beekeepers (estimated at 5 million USD/year; Eischen et al., 2005). In contrast, European Foulbrood (EFB) is caused by *Melissococcus plutonius*, a non-spore-forming bacterium (Bailey, 1957), and is treatable with OTC (Forsgren, 2010). Still, treatment with OTC can be economically damaging as antibiotic residues can prevent beekeepers from selling honey (Gilliam & Argauer, 1981; Matsuka & Nakamura, 1990; Milbrath, 2021; Sporns et al., 1986; Wilson, 1974). Additionally, EFB can easily return after OTC treatment, and outbreaks have become increasingly prominent worldwide (Grossar et al., 2020; Simone-Finstrom & Spivak, 2010).

EFB infection causes discoloration and loss of internal body pressure in affected larvae, leaving

them yellow-brown and flaccid (Forsgren, 2010). *Melissococcus plutonius* is a gram-positive,

microaerophilic to anaerobic bacterium that thrives in the larval honey bee midgut (Bailey, 1983;

Bailey & Collins, 1982; Forsgren, 2010). Infected larvae pass on the bacterium by either

defecating or dying within their wax cells, and bacteria are then spread to healthy larvae by adult

nurse bees through larval care activities (Bailey, 1983), often resulting in the spread of EFB

within a colony and throughout an apiary.

Propolis is a substance bees make by mixing resinous plant exudates with saliva and beeswax

(Simone-Finstrom & Spivak, 2010). Bees use propolis to seal cracks in the hive body, reduce

the hive entrance, and line the rims of wax cells (Evans & Spivak, 2010; Seeley & Morse, 1976;

Strehle et al., 2003). In a natural setting, feral colonies create a "propolis envelope" around the

entire surface of their rough nesting cavity (e.g., a tree cavity; Seeley & Morse, 1976), but this

behavior is exhibited much less inside hive boxes made from finely milled lumber. Instead,

propolis is frequently deposited between boxes, hive covers, and man-made frames that hold

beeswax combs. Due to this meticulous behavior, propolis can be easily harvested for human use

in folk medicine (Marcucci, 1995; Przybyłek & Karpiński, 2019). Propolis contains

antimicrobial compounds, such as flavonoids, phenolics, terpenoids and aromatic acids, that are

present in the source plant exudates (Marcucci, 1995; Przybyłek & Karpiński, 2019). These

compounds vary greatly depending on hive location and nearby plants from which the bees

collect material (Huang et al., 2014; Sforcin, 2016). For example, propolis in temperature regions

originates mostly from *Populus* species (Anđelković et al., 2017; Sforcin, 2016), and thus

propolis samples collected in temperate regions (e.g., Ohio) largely contain flavonoids and

phenolics (Johnson et al., 1994).

Propolis has been shown to combat various pathogenic threats to bees, including fungal parasites

like *Nosema* spp. and *Ascosphaera apis*, the causative agents of Nosemosis and larval

chalkbrood disease, respectively (Borba et al., 2015; Simone-Finstrom & Spivak, 2012). It has

also demonstrated effects against bee viral infections, including deformed wing virus and black

queen cell virus (Borba et al., 2015; Drescher et al., 2017). Additionally, bees from hives that

were supplemented with additional propolis were found to exhibit lower expression of immune

genes (Simone et al., 2009).

With a recent increase in EFB outbreaks and the need for more sustainable treatments, we aimed

to test the antibacterial properties of extracts from propolis and other bee-derived products (i.e.,

wax brood combs and honey combs) against *M. plutonius*. We are interested in the effects of wax

combs previously used for brood rearing on honey bee health, due to the deposition of both

antimicrobial substances (i.e., propolis and honey; Pasupuleti et al., 2017) and bee frass over

time. Bees use propolis to increase the structural integrity of all wax combs (Strehle et al., 2003),

but brood comb darkens in color and gradually becomes a composite material, as the wax

accumulates larval silk, castings, and frass with repeated cycles of brood rearing (Hepburn &

Kurstjens, 1988). Due to this build-up of waste materials, as well as the potential for build-up of

pesticide residues, beekeepers are often encouraged to replace their used brood comb after

several years (Jaycox, 1979) However, the presence and potential benefits of propolis in wax

cells is a relatively unexplored research topic. Surveys of beekeepers indicate that older brood

comb may be associated with improved winter survival

(https://research.beeinformed.org/survey/), possibly due to the propolis constituents or other

components accumulated by wax through repeated brood-rearing cycles.

In the present study, we tested the antibacterial properties of ethanol extracts of (1) propolis, (2)

brood comb (wax comb in which immature bees had previously been reared), and (3) honey

comb (wax comb used exclusively for honey storage) against *M. plutonious,* the bacterial agent

causing EFB. Additionally, these extracts were tested against two model bacteria, gram-negative

Escherichia coli and gram-positive *Staphylococcus saprophyticus*. The latter species are not

associated with honey bee disease; however, they are pathogenic bacteria in mammals and are

frequently used in antimicrobial testing. Historically, propolis extracts have not demonstrated

substantial inhibition of gram-negative bacteria (Przybyłek & Karpiński, 2019), therefore, *E. coli*

was included to serve as a negative control. We measured the optical density of all bacterial

cultures after exposure to propolis, brood comb and honey comb extracts to determine the lowest

concentration, or minimum inhibitory concentration (MIC), that inhibits bacterial growth.

Materials and Methods

Colony setup

A total of six experimental colonies were established in standard eight-frame Langstroth-style

hive equipment at three apiary locations, with two colonies at each site, in spring of 2018.

Apiaries were located at (1) the Waterman Agriculture and Natural Resources Laboratory in

Columbus, Ohio (40.0104° N, 83.0400° W), (2) the Ohio Agricultural Research and

Development Center in Wooster, Ohio (40.7818° N, 81.9305° W), and (3) the Muck Crops

Agricultural Research Station in Willard, Ohio (41.010200° N, 82.731350° W). New waxed

black plastic foundation (9FDBW, Pierco) in new wooden frames were fitted in either newly

constructed or thoroughly cleaned deep hive boxes (KD-603, Mann Lake). A new or thoroughly

cleaned metal queen excluder (HD-121, Mann Lake) was used to restrict the queen and rearing

of immature bees, or brood, to the bottom box so that the upper box could only be used for honey

storage (Fig. 1).

Wax and propolis collection and extraction

Using a freshly cleaned hive tool, empty combs (free of honey or brood) and propolis samples

were collected from each site in late-winter of 2019 (approximately ten months later) and stored

in darkness at -20°C until extraction. Before comb samples or propolis were extracted, they were

first ground to a fine powder using liquid nitrogen and a food processor (FP2500B, Black &

Decker). Separate food processors were used for each sample type to prevent cross

contamination. Two grams of each powdered sample were placed in 20 mL of 95% ethanol (1:10

w/v) and sonicated (HB-S-49DHT, Kenda) for 60 minutes at up to 60°C. The resulting extract

was gravity filtered using grade one filter paper to remove particulate matter, evaporated under a

stream of nitrogen at 20°C, and resuspended in 10 mL of 70% ethanol. Ethanol was chosen as it

is a commonly used solvent in propolis research and many of the antimicrobial compounds

present are soluble in it. In total, nine crude ethanol extracts were made, three each of honey

comb, brood comb and propolis corresponding to the three apiary locations. Concentrations were

measured after evaporating crude extracts. The average concentration of dry material in crude

extracts was 16.0 mg/mL for honey comb, 45.33 mg/mL for brood comb, and 72.83 mg/mL for

propolis. A series of eight two-fold dilutions were then made for testing against bacteria. Diluted

extract concentrations are listed in Fig. 2. All extracts were stored in darkness at 4°C.

Figure 1. Experimental colony design. Two eight-frame deep boxes separated by a queen

excluder, to restrict the queen, brood and brood comb to the lower box and allow construction of

comb for honey storage in the upper box. Propolis was collected from hive crevices.

Bacterial Cultures

Melissococcus plutonius was isolated from EFB-infected larvae collected at the Ohio State

University Wooster Campus in spring 2018. This microaerophilic-anaerobic bacterium was

isolated and cultured by inoculating modified basal medium (Forsgren et al., 2013) with fresh

infected larvae that were ground with a sterile pestle in 1x PBS solution. Growth medium (Table

S1) contained yeast extract, glucose, sucrose, L-cysteine, 1M KH_2PO_4 and 2.5M KOH

(AC400405000, D16-500 and P285-500, Fisher Scientific; P5958 and W326305, Sigma

Aldrich). To prevent the growth of secondary bacteria, nalidixic acid was dissolved in 0.1M

NaOH (AAB2509606 and S318-100, Fisher Scientific), filter sterilized, and added to the

medium at a concentration of 3 µg/ml. *Melissococcus plutonius* was cultured in a vinyl anaerobic

chamber (Coy Laboratory Products, Grass Lake, MI) at 37°C (Table S1). Sequencing using

primers from Haynes et al. (2013) confirmed the identity of bacteria in culture (GenBank

accession: MN886237). Cultures were stored in 50% glycerol for long-term storage at -80°C.

Strains of two aerobic bacterial species, *E. coli* (CSH36) and *S. saprophyticus* (ATCC15305),

were obtained from collections of the Department of Microbiology at The Ohio State University

and cultured using standard protocols in Mueller-Hinton media (OXCM0405B, Fisher Scientific)

at 37°C.

Bacterial Growth Inhibition Assays

We determined the MIC of each treatment using a broth microdilution method modeled after the

Clinical and Laboratory Standards Institute (Cockerill et al., 2012) as in Sozanski et al. (2020).

Before testing the inhibitory properties of each extract dilution series, bacterial isolates were first

inoculated in broth media appropriate for each bacterium (aerobic Meuller-Hinton media or

anaerobic *M. plutonius* media) and grown overnight at 37ºC under shaking aerobic conditions (*E.*

coli and *S. saprophyticus*) or still anaerobic conditions (*M. plutonius*). To inoculate microplate

test wells (96-well flat-bottomed microplates, Thermo Scientific™ 243656) at 5 x 10^5 CFU/mL,

we first diluted overnight cultures to a 0.5 McFarland standard (1.0×10^8 CFU/mL) and further

diluted cultures by a factor of five. We then added these diluted cultures to plate wells

containing 5 µL of either antimicrobial treatments or controls and 190 µL of appropriate broth

media, resulting in a final concentration of 5 x 10^5 CFU/mL. Each column of wells on the 96-

well plate contained the series of two-fold diluted propolis, brood comb or honey comb extracts,

with column one containing the most concentrated treatment and column eight containing the

most dilute treatment. Columns nine and 10 contained a positive oxytetracycline (OTC) control

(1 mg/mL) and a negative solvent control (70% ethanol), respectively. The final two columns on

the plate served as a standard growth control and a contamination control. Standard growth

control wells contained 195 µL of appropriate broth with 5 µL of diluted culture. Finally,

contamination control wells contained 200 µL of appropriate broth. Each 96-well plate was

considered one technical replicate. For every extract and bacterium tested, there were three

biological replicates, corresponding to the three different source apiaries, and three technical

replicates (n = 9).

To measure bacterial growth, plates were placed in an ELx808i™ Absorbance Microplate

Reader (BioTek Instruments) with incubation at 37°C. Plates were shaken and an absorbance

reading was made at 600 nm (OD_{600}) every five minutes for 12 hours (Quigley, 2008). Standard

growth controls for each plate were used to determine the mid-log phase of bacterial growth,

allowing MIC determination at that time point for each treatment.

Statistical Analysis and MIC Determination

We first analyzed standard growth wells to determine the mid-log phase of each plate using the R

package Growthcurver v3.6.0 (Sprouffske & Wagner, 2016). We recorded OD_{600} readings for

bacterial growth in all wells at their corresponding mid-log phase and tested each plate for

growth differences among treatments using a Kruskal-Wallis test. Dunn's Multiple Comparison

post-hoc test with Bonferroni correction was used to determine the MIC of each treatment

against all three bacteria. We defined the MIC as the lowest treatment concentration that was

significantly different from the solvent control. Propolis dilutions one through five were

ultimately excluded from analysis, as these wells were oversaturated, causing noise in OD

readings. All statistics were performed in R v4.0.3 (R Core Team, 2019).

Results

Growth of *M. plutonius* was inhibited by OTC ($p < 0.001$) and all three hive product extracts

(Fig. 2A). The MIC of honey comb extract ($p = 0.007$) and brood comb extract ($p < 0.001$) was

16.00 mg/mL and 45.33 mg/mL, respectively. The MIC of propolis extract against *M. plutonius*

was the second-lowest concentration tested, 1.14 mg/mL ($p < 0.001$). Growth of *S. saprophyticus*

was inhibited only by OTC ($p < 0.001$) and propolis extract (Fig. 2B; $p = 0.001$). The MIC of

propolis extract against *S. saprophyticus* was 2.28 mg/mL. Growth of *E. coli* was not

significantly inhibited by any extract, but was inhibited by OTC (Fig. 2C; $p = 0.001$). Statistical

analyses also revealed a significant effect by plate for each bacterium ($p < 0.001$), as well as a

significant effect of extract source location for *S. saprophyticus* ($p < 0.001$) and *M. plutonius* (p

$= 0.032$).

Figure 2. OD$_{600}$ of (A) *M. plutonius*, (B) *S. saprophyticus* and (C) *E. coli* at mid-log phase when

treated with a two-fold dilution series of 70% ethanol extracts of honey comb, brood comb and

propolis. Comb and propolis were collected from three locations, with three technical replicates

each (n=9). Ethanol (70%) was used as a solvent control (EtOH) and Oxytetracycline

hydrochloride (1mg/mL) was used as a positive control (OTC). Untreated wells were used as a

standard growth control (Std.). MICs and controls significantly different (p<0.05, Dunn's

Multiple Comparison test with Bonferonni correction) from solvent control are indicated with

asterisks (*).

Discussion

In this study, we set out to test the antibacterial effects of ethanol extracts of three hive

products— propolis, brood comb, and honey comb—against *M. plutonius*, the causative agent of

EFB in honey bees. Overall, *M. plutonius* growth was most susceptible to inhibition by extracts

of all three hive products. To our knowledge, this is the first study to test the antibacterial

properties of propolis and wax combs against *M. plutonius*. The highest concentrations of both

honey comb (16.00 mg/mL) and brood comb (45.33 mg/mL) extracts successfully inhibited the

growth of *M. plutonius*, but MICs for these extracts were an order of magnitude higher than that

of propolis extract (1.14 mg/mL; Fig 2A). We demonstrated strong antibacterial properties of

propolis against *M. plutonius*, at roughly the same concentration as our positive control OTC (1.0

mg/mL). The gram-positive model bacterium, *S. saprophyticus*, was more susceptible than *E.*

coli to inhibition by propolis extract (MIC = 2.28 mg/mL; Fig. 2B), which is consistent with

previous work showing that propolis extracts have more inhibitory activity against gram-positive

bacteria (Rahman et al., 2010; Ristivojević et al., 2016; Sforcin et al., 2000; Tosi et al., 1996;

Velikova et al., 2000). This tolerance is likely due to the structure of gram-negative bacterial

cells, which are characterized by a thin layer of peptidoglycans and an extra plasma membrane

(Harrop et al., 1989); however, the composition of these outer membranes—and thus the

tolerance to propolis activity—may be species-dependent (Mirzoeva et al., 1997). In line with

this previous work, propolis extracts here did not inhibit *E. coli* growth.

We observed significant differences between experimental plates, as well as between materials

collected from different apiary locations. These differences are likely due to a small number of

replicates. Though chemical composition of propolis is dependent upon the hive location and

surrounding flora, this was unlikely to have affected our experiment, as our hives were located

within 200 miles of one another, with similar surrounding flora. Additionally, propolis sourced

from North America, Europe and Asia is characterized by similar antibiotic compounds,

including flavonoids, aromatic acids, and esters (Sforcin, 2016).

The antibiotic properties of propolis are well-established (Marcucci, 1995; Przybyłek &

Karpiński, 2019), but this is not the case for properties of honey comb and brood comb. Our

results suggest that antibacterial compounds are much more abundant in propolis but are also

present in wax combs and can serve to inhibit the growth of *M. plutonius.* Understanding the

origins of this activity could inform beekeeping practices, such as brood comb replacement.

Beeswax itself is made up of hydrocarbon chains, free fatty acids, free fatty alcohols and wax

esters, but also contains plant-derived materials (e.g., nectar, pollen, resin) that are incorporated

as honey bees build and fill combs (Fratini et al., 2016). Due to its lipophilic composition,

beeswax captures many compounds, including pheromones and acaracides (Svečnjak et al.,

2019). Therefore, we assume that antimicrobial effects of wax comb extracts are likely due to

other compounds incorporated during and after comb construction. To disentangle the effects of

beeswax itself from those of other compounds taken up by or incorporated into the wax, future

research should focus on collecting wax scales directly from bees (as in Svečnjak et al., 2019).

Brood comb frames within a healthy colony will typically hold cells of pollen and honey along

the outer edges (Schneider, 2015), but using a queen excluder (Fig. 1; a common practice for

hobbyists and large-scale beekeepers) encourages honey storage to occur outside of the brood

boxes. Like propolis, honey is another bee-derived substance that has long been used in human

medicine (Bogdanov et al., 2008). The antimicrobial properties of honey derive from its low pH,

flavonoids present in pollen and nectar, as well as enzymes originating from both plants and

honey bee hypopharyngeal gland secretions (Aurongzeb & Azim, 2011). Though we collected

honey comb free of honey to make our ethanol extracts, there were likely traces of honey

present, and previous honey storage may have incorporated compounds, such as flavonoids, into

the wax cells.

The propolis envelope that surrounds the entire nest cavity of a feral colony has been

hypothesized to have several homeostatic functions, including antimicrobial protection, water-

repellency and even communication (Seeley & Morse, 1976). This extensive envelope is not

present in managed colonies, but propolis is incorporated into the rims of wax cells, possibly to

serve a structural function (Strehle et al., 2003) or confer an antimicrobial benefit (Evans &

Spivak, 2010), but these hypotheses remain relatively unexplored. The composite structure of

brood combs—containing silk, frass, propolis and exogenous plant material (Hepburn &

Kurstjens, 1988)—results in a darker, thicker comb than those that are used strictly for honey

storage, but studies on the quantity of propolis in brood or honey combs, and whether those

quantities differ between comb type, do not exist. Though there are potential benefits from

propolis build-up in well-used comb, there is still concern over build-up of pesticide residues and

pathogen persistence in old brood combs, which currently informs brood comb replacement

practices.

Propolis as a potential preventative or treatment for EFB

With well-established antibacterial properties, and our demonstrated activity against *M.*

plutonius, propolis and propolis extracts should be further developed for control of EFB in honey

bee colonies. There is now a growing body of literature supporting the benefits of propolis in

honey bee immunity, and bee stocks artificially selected for higher propolis collection

demonstrate increased survivorship and longevity (Nicodemo et al., 2014). Aside from breeding

bees for increased propolis collection, two of the most practical ways to supplement colonies

with propolis include applying propolis extract to colony walls or stimulating propolis collection

through hive design (i.e., propolis traps with many cervices; Borba et al., 2015; Drescher et al.,

2017). Colonies supplemented with propolis exhibit lower viral infection rates and immune gene

expression (Borba et al., 2015; Drescher et al., 2017; Simone et al., 2009). As a bee-derived

substance that is (1) already present in the hive, (2) continually changing in its chemical

constituents (based on available flora), and (3) safe for humans when present in harvested honey

(unlike OTC; Gilliam and Argauer, 1981; Matsuka and Nakamura, 1990; Sporns et al., 1986;

Wilson, 1974), propolis shows promise as a sustainable preventative or curative against EFB

outbreaks. Field experiments using propolis traps or extracts, as well as *in-vitro* experiments (as

in Grossar et al., 2020) in which EFB-infected larvae are treated with topical or ingestible

propolis extracts, are the next step to evaluating the effectiveness of propolis to manage EFB.

Acknowledgements

References

Anđelković, B., Vujisić, L., Vučković, I., Tešević, V., Vajs, V., & Gođevac, D. (2017).

Metabolomics study of *Populus* type propolis. *Journal of Pharmaceutical and Biomedical*

Analysis, *135*, 217–226. https://doi.org/10.1016/j.jpba.2016.12.003

Aurongzeb, M., & Azim, M. K. (2011). Antimicrobial properties of natural honey: a review of

literature. *Pakistan Journal of Biochemistry and Molecular Biology*, *44*(3), 118–124.

Bailey, L. (1957). The cause of European Foulbrood. *Bee World*, *38*(4), 85–89.

Bailey, L. (1983). *Melissococcus pluton*, the cause of European foulbrood of honey bees (*Apis*

spp.). *Journal of Applied Bacteriology*, *55*, 65–69.

Bailey, L., & Collins, M. D. (1982). Reclassification of *'Streptococcus pluton*' (White) in a new

genus *Melissococcus pluton* nom. rev.; comb. nov. *Journal of Applied Bacteriology*, *53*,

215–217.

Bogdanov, S., Jurendic, T., Sieber, R., & Gallmann, P. (2008). Honey for nutrition and health: A

review. *Journal of the American College of Nutrition*, *27*(6), 677–689.

https://doi.org/10.1080/07315724.2008.10719745

Borba, R. S., Klyczek, K. K., Mogen, K. L., & Spivak, M. (2015). Seasonal benefits of a natural

propolis envelope to honey bee immunity and colony health. *Journal of Experimental*

Biology, *218*(22), 3689–3699. https://doi.org/10.1242/jeb.127324

Bulson, L., Becher, M. A., McKinley, T. J., & Wilfert, L. (2021). Long-term effects of antibiotic

treatments on honeybee colony fitness: A modelling approach. *Journal of Applied Ecology*,

58(1), 70–79. https://doi.org/10.1111/1365-2664.13786

Cockerill, F. R., Wikler, M. A., Alder, J., Dudley, M. N., Eliopoulos, G. M., Ferraro, M. J.,

Hardy, D. J., Hecht, D. W., Hindler, J. A., Patel, J. B., Powell, M., Swenson, J. M.,

Thomson, R. B., Traczewski, M. M., Turnidge, J. D., Weinstein, M. P., & Zimmer, B. L.

(2012). *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow*

Aerobically (9th ed., Vol. 32). National Committee for Clinical Laboratory Standards.

www.clsi.org.

Drescher, N., Klein, A. M., Neumann, P., Yañez, O., & Leonhardt, S. D. (2017). Inside honeybee

hives: Impact of natural propolis on the ectoparasitic mite *Varroa destructor* and viruses.

Insects, *8*(15), 8010015. https://doi.org/10.3390/insects8010015

Eischen, F. A., Graham, R. H., & Cox, R. (2005). Regional distribution of *Paenibacillus larvae*

subspecies *larvae*, the causative organism of American Foulbrood, in honey bee colonies of

the western United States. *Journal of Economic Entomology*, *98*(4), 1087–1093.

https://academic.oup.com/jee/article/98/4/1087/883400

Evans, J. D., & Spivak, M. (2010). Socialized medicine: Individual and communal disease

barriers in honey bees. *Journal of Invertebrate Pathology*, *103*, S62–S72.

https://doi.org/10.1016/j.jip.2009.06.019

Forsgren, E. (2010). European foulbrood in honey bees. *Journal of Invertebrate Pathology*, *103*,

S5–S9. https://doi.org/10.1016/j.jip.2009.06.016

Forsgren, E., Budge, G. E., Charrière, J. D., & Hornitzky, M. A. Z. (2013). Standard methods for

European foulbrood research. *Journal of Apicultural Research*, *52*(1), 1–14.

https://doi.org/10.3896/IBRA.1.52.1.12

Fratini, F., Cilia, G., Turchi, B., & Felicioli, A. (2016). Beeswax: A minireview of its

antimicrobial activity and its application in medicine. *Asian Pacific Journal of Tropical*

Medicine, *9*(9), 839–843. https://doi.org/10.1016/j.apjtm.2016.07.003

Gilliam, M., & Argauer, R. J. (1981). Oxytetracycline residues in surplus honey, brood nest

honey, and larvae after medication of colonies of honey bees, *Apis mellifera*, with antibiotic

extender patties, sugar dusts, and syrup sprays. *Environmental Entomology*, *10*(4), 479–482.

https://academic.oup.com/ee/article/10/4/479/2392625

Grossar, D., Kilchenmann, V., Forsgren, E., Charrière, J. D., Gauthier, L., Chapuisat, M., &

Dietemann, V. (2020). Putative determinants of virulence in *Melissococcus plutonius*, the

bacterial agent causing European foulbrood in honey bees. *Virulence*, *11*(1), 554–567.

https://doi.org/10.1080/21505594.2020.1768338

Hansen, H., & Brødsgaard, C. J. (1999). American foulbrood: A review of its biology, diagnosis

and control. *Bee World*, *80*(1), 5–23. https://doi.org/10.1080/0005772X.1999.11099415

Harrop, A. J., Hocknull, M. D., & Lilly, M. D. (1989). Biotransformation in organic solvents: A

difference between gram-positive and gram-negative bacteria. *Biotechnology Letters*,

11(11), 807–810.

Haseman, L. (1961). How long can spores of American foulbrood live? *American Bee Journal*,

101, 298–299.

Haynes, E., Helgason, T., Young, J. P. W., Thwaites, R., & Budge, G. E. (2013). A typing

scheme for the honeybee pathogen *Melissococcus plutonius* allows detection of disease

transmission events and a study of the distribution of variants. *Environmental Microbiology*

Reports, *5*(4), 525–529. https://doi.org/10.1111/1758-2229.12057

Hepburn, H. R., & Kurstjens, S. P. (1988). The combs of honeybees as composite materials.

Apidologie, *19*(1), 25–36.

Huang, S., Zhang, C. P., Wang, K., Li, G. Q., & Hu, F. L. (2014). Recent advances in the

chemical composition of propolis. *Molecules*, *19*(12), 19610–19632.

https://doi.org/10.3390/molecules191219610

Jaycox, E. R. (1979). Comb foundation: Are we using enough? *American Bee Journal*, *119*(7),

515–516.

Johnson, K. S., Eischen, F. A., & Giannasi, D. E. (1994). Chemical composition of North

America bee propolis and biological activity towards larvae of greater wax moth

(*Lepidoptera: Pyralidae*). *Journal of Chemical Ecology*, *20*(7), 1783–1791.

Lodesani, M., & Costa, C. (2005). Limits of chemotherapy in beekeeping: Development of

resistance and the problem of residues. *Bee World*, *86*(4), 102–109.

https://doi.org/10.1080/0005772X.2005.11417324

Marcucci, M. C. (1995). Propolis: Chemical composition, biological properties and therapeutic

activity. *Apidologie*, *26*, 83–99.

Matsuka, M., & Nakamura, J. (1990). Oxytetracycline residues in honey and royal jelly. *Journal*

of Apicultural Research, *29*(2), 112–117. https://doi.org/10.1080/00218839.1990.11101206

Milbrath, M. (2021). Honey bee bacterial diseases. In T. R. Kane & Faux Cynthia M. (Eds.),

Honey Bee Medicine for the Veterinary Practitioner (First, pp. 277–293). John Wiley &

Sons.

Mirzoeva, O. K., Grishanin, R. N., & Calder, P. C. (1997). Antimicrobial action of propolis and

some of its components: The effects on growth, membrane potential and motility of

bacteria. *Microbiological Research*, *152*(3), 239–246. https://doi.org/10.1016/S0944-

5013(97)80034-1

Nicodemo, D., Malheiros, E. B., De Jong, D., & Couto, R. H. N. (2014). Increased brood

viability and longer lifespan of honeybees selected for propolis production. *Apidologie*,

45(2), 269–275. https://doi.org/10.1007/s13592-013-0249-y

Pasupuleti, V. R., Sammugam, L., Ramesh, N., & Gan, S. H. (2017). Honey, propolis, and royal

jelly: A comprehensive review of their biological actions and health benefits. *Oxidative*

Medicine and Cellular Longevity, *2017*, 1259510. https://doi.org/10.1155/2017/1259510

Przybyłek, I., & Karpiński, T. M. (2019). Antibacterial properties of propolis. *Molecules*, *24*(11),

2047. https://doi.org/10.3390/molecules24112047

Quigley, T. (2008). *Monitoring the growth of* E. coli *with light scattering using the Synergy^{TM} 4*

Multi-Mode Microplate Reader with Hybrid Technology^{TM}. www.biotek.com

R Core Team. (2019). *R: A language and environment for statistical computing*. https://www.R-

project.org/

Rahman, M. M., Richardson, A., & Sofian-Azirun, M. (2010). Antibacterial activity of propolis

and honey against *Staphylococcus aureus* and *Escherichia coli*. *African Journal of*

Microbiology Research, *4*(16), 1872–1878.

https://www.researchgate.net/publication/228662460

Raymann, K., Shaffer, Z., & Moran, N. A. (2017). Antibiotic exposure perturbs the gut

microbiota and elevates mortality in honeybees. *PLoS Biology*, *15*(3), e2001861.

https://doi.org/10.1371/journal.pbio.2001861

Ristivojević, P., Dimkić, I., Trifković, J., Berić, T., Vovk, I., Milojkovič-Opsenica, D., &

Stanković, S. (2016). Antimicrobial activity of Serbian propolis evaluated by means of

MIC, HPTLC, bioautography and chemometrics. *PLoS ONE*, *11*(6), e0157097.

https://doi.org/10.1371/journal.pone.0157097

Schneider, S. S. (2015). The honey bee colony: Life history. In J. M. Graham (Ed.), *The Hive*

and the Honey Bee (pp. 73–109). Dadant & Sons.

Seeley, T. D., & Morse, R. A. (1976). The nest of the honey bee (*Apis mellifera* L.). *Insectes*

Sociaux, *23*(4), 495–512. https://doi.org/10.1007/BF02223477

Sforcin, J. M. (2016). Biological properties and therapeutic applications of propolis.

Phytotherapy Research, *30*(6), 894–905. https://doi.org/10.1002/ptr.5605

Sforcin, J. M., Fernandes, A., Lopes, C. A. M., Bankova, V., & Funari, S. R. C. (2000). Seasonal

effect on Brazilian propolis antibacterial activity. *Journal of Ethnopharmacology*, *73*, 243–

249. www.elsevier.com/locate/jethpharm

Simone, M., Evans, J. D., & Spivak, M. (2009). Resin collection and social immunity in honey

bees. *Evolution*, *63*(11), 3016–3022. https://doi.org/10.1111/j.1558-5646.2009.00772.x

Simone-Finstrom, M., & Spivak, M. (2012). Increased resin collection after parasite challenge: A

case of self-medication in honey bees? *PLoS ONE*, *7*(3), e34601.

https://doi.org/10.1371/journal.pone.0034601

Simone-Finstrom, M., & Spivak, M. (2010). Propolis and bee health: The natural history and

significance of resin use by honey bees. *Apidologie*, *41*(3), 295–311.

https://doi.org/10.1051/apido/2010016

Soares, K. O., Oliveira, C. J. B. de, Rodrigues, A. E., Vasconcelos, P. C., Silva, N. M. V. da,

Cunha Filho, O. G. da, Madden, C., & Hale, V. L. (2021). Tetracycline exposure alters key

gut microbiota in Africanized honey bees (*Apis mellifera scutellata* x spp.). *Frontiers in*

Ecology and Evolution, *9*, 716660. https://doi.org/10.3389/fevo.2021.716660

Sozanski, K., Prado, L. P. do, Mularo, A. J., Sadowski, V. A., Jones, T. H., & Adams, R. M. M.

(2020). Venom function of a new species of *Megalomyrmex* Forel, 1885 (Hymenoptera:

Formicidae). *Toxins*, *12*(11), 679. https://doi.org/10.3390/toxins12110679

Sporns, P., Kwan, S., & Roth, L. A. (1986). HPLC analysis of oxytetracycline residues in honey.

Journal of Food Protection, *49*(5), 383–388. http://meridian.allenpress.com/jfp/article-

pdf/49/5/383/1657041/0362-028x-49_5_383.pdf

Sprouffske, K., & Wagner, A. (2016). Growthcurver: An R package for obtaining interpretable

metrics from microbial growth curves. *BMC Bioinformatics*, *17*, 172.

https://doi.org/10.1186/s12859-016-1016-7

Strehle, M. A., Jenke, F., Fröhlich, B., Tautz, J., Riederer, M., Kiefer, W., & Popp, J. (2003).

Raman spectroscopic study of spatial distribution of propolis in comb of *Apis mellifera*

carnica (Pollm.). *Biopolymers - Biospectroscopy Section*, *72*(4), 217–224.

https://doi.org/10.1002/bip.10379

Svečnjak, L., Chesson, L. A., Gallina, A., Maia, M., Martinello, M., Mutinelli, F., Muz, M. N.,

Nunes, F. M., Saucy, F., Tipple, B. J., Wallner, K., Waś, E., & Waters, T. A. (2019).

Standard methods for *Apis mellifera* beeswax research. *Journal of Apicultural Research*,

58(2), 1–108. https://doi.org/10.1080/00218839.2019.1571556

Tosi, B., Donini, A., Romagnoli, C., & Bruni, A. (1996). Antimicrobial activity of some

commercial extracts of propolis prepared with different solvents. *Phytotherapy Research*,

10, 335–336.

Velikova, M., Bankova, V., Marcucci, M. C., Tsvetkova, I., & Kujumgiev, A. (2000). Chemical

composition and biological activity of propolis from Brazilian Meliponinae. *Verlag Der*

Zeitschrift Für Naturforschung, *55*, 785–789. www.znaturforsch.com

Wilson, W. T. (1974). Residues of oxytetracycline in honey stored by *Apis mellifera*.

Environmental Entomology, *3*(4), 674–676.

https://academic.oup.com/ee/article/3/4/674/2395300

Table S1. Protocol for *Melissococcus plutonius* cultivation. Growth media adapted from Bailey

(1957) and Forsgren et al (2013). All steps were performed using sterile equipment and in a

sterile environment, including a biosafety cabinet and an anaerobic chamber.

Growth Media (makes 1 L)	Cultivation in Growth Media

1. Mix the following into 800 mL DI H$_2$0:
  a. 10 g yeast extract
  b. 10 g glucose
  c. 10 g starch (solid medium) or sucrose (liquid medium)
  d. 0.25 g L-cysteine
  e. 20 g agar (solid medium only)
2. Add 100 mL 1M KH$_2$PO$_4$
3. Adjust pH to 6.6 using 2.5M KOH
4. Adjust final volume to 1000 mL using distilled water
5. Autoclave for 15 min at 115°C
6. Dissolve nalidixic acid into 0.1M NaOH (30 mg/mL) and filter sterilize (store at -20°C)
7. After cooling media, add filter sterilized nalidixic acid at a final concentration of 3 µg/mL
8. Transfer to anaerobic chamber
	480
	1. Use sterile pestle to prepare suspension of 2-3 diseased brood in 1x PBS and transfer suspension to anaerobic chamber
	2. Inoculate growth medium:
	  a. Add ~20 µL suspension to 20 mL liquid growth medium OR
	  b. Pipette ~20 µL suspension onto agar plate and spread with sterile plate spreader
	3. Incubate in anaerobic conditions at 37°C and check daily