A high-throughput yeast display approach to profile pathogen proteomes for MHC-II binding

Brooke D. Huisman[1,2], Zheng Dai[3,4], David K. Gifford[2,3,4], Michael E. Birnbaum[1,2,5,]*

[1] Koch Institute for Integrative Cancer Research, Cambridge, MA, USA
[2] Department of Biological Engineering, MIT, Cambridge, MA, USA
[3] Computer Science and Artificial Intelligence Laboratory, MIT, Cambridge, MA, USA
[4] Department of Electrical Engineering and Computer Science, MIT, Cambridge, MA, USA
[5] Ragon Institute of MIT, MGH, and Harvard, Cambridge, MA, USA

* Corresponding author: mbirnb@mit.edu

Author contributions

Competing Interest Statement

D.K.G. is a founder of ThinkTx. M.E.B. is an equity holder in 3T Biosciences, and is a co-founder of Viralogic Therapeutics and Abata Therapeutics. The other authors declare no competing interests.

Keywords: yeast surface display, antigen prediction, peptide-MHC

Abstract

T cells play a critical role in the adaptive immune response, recognizing peptide antigens presented on the cell surface by Major Histocompatibility Complex (MHC) proteins. While assessing peptides for MHC binding is an important component of probing these interactions, traditional assays for testing peptides of interest for MHC binding are limited in throughput. Here we present a yeast display-based platform for assessing the binding of tens of thousands of user-defined peptides in a high throughput manner. We apply this approach to assess a tiled library covering the SARS-CoV-2 proteome and four dengue virus serotypes for binding to human class II MHCs, including HLA-DR401, -DR402, and -DR404. This approach identifies binders missed by computational prediction and serves as a framework for diverse objectives, including examining relationships between viral conservation and MHC binding.

Introduction

Major histocompatibility complex (MHC) proteins play a critical role in adaptive immunity by presenting peptide fragments on the surface of cells. Peptide-MHC (pMHC) complexes are then surveilled by T cells via their T cell receptors, enabling immune cells to sense intracellular dysfunction, such as the presence of pathogen-derived peptides. Class I MHCs (MHC-I) present peptides with relatively constrained lengths, typically from 8-11 amino acids, within a closed binding groove to CD8$^+$ cytotoxic T cells. In contrast, class II MHCs (MHC-II) present peptides to CD4$^+$ T cells, and they have an open binding groove, allowing for the display of longer peptides consisting of a 9 amino acid 'core' flanked by a variable number of additional residues on each side.

Generating reliable and rapid data on peptide-MHC binding could be useful for identifying important clinical targets, including for optimized T cell epitopes in vaccine design (Dai et al., 2021; Keskin et al., 2019; Liu et al., 2020, 2021a; Moise et al., 2015; Ott et al., 2017; Patronov and Doytchinova, 2013; Rosati et al., 2021). In fact, therapeutics to generate antigen-specific T cell responses have shown great promise in cancer (Keskin et al., 2019; Ott et al., 2017) and infectious disease (Gambino et al., 2021). Since understanding peptide-MHC binding is critical for identifying and engineering T cell epitopes, there have been sustained efforts to produce high-quality experimental data and predictive algorithms.

Initial experimental methods for determining peptide binding to MHC relied upon the analysis of synthesized candidate peptides via MHC stability or functional assays. While these methods typically produce high-confidence data, they can be difficult to scale beyond a relatively small number of candidate peptides (Altmann and Boyton, 2020; Justesen et al., 2009; Mateus et al., 2020; Sidney et al., 2010; Yin and Stern, 2014). More recently, mass spectrometry-based approaches have been demonstrated for determining the MHC-presented peptide repertoire of cells. These approaches include monoallelic mass spectrometry, which allows for the unambiguous assignment of presented peptides to a given MHC allele. However, mass spectrometry-based approaches are not necessarily quantitative measures of presented peptide affinity or abundance, although there have been advances in quantitation using internal standards (Stopfer et al., 2020, 2021). Additionally, peptides naturally expressed by a cell can potentially crowd out exogenously examined peptides of interest (Abelin et al., 2017, 2019; Parker et al., 2021; Purcell et al., 2019).

A wave of higher throughput approaches have been recently developed for studying peptide-MHC interactions, including yeast display (Jiang and Boder, 2010; Liu et al., 2021b; Rappazzo et al., 2020) and mammalian display-based methods (Obermair et al., 2021). Several of these approaches circumvent the bottlenecks of synthesizing or identifying peptides by utilizing DNA-based inputs and outputs (Jiang and Boder, 2010; Obermair et al., 2021; Rappazzo et al., 2020). These assays rely upon libraries that are often generated via DNA oligonucleotide synthesis, and use peptide stabilization and surface expression (Jiang and Boder, 2010; Liu et al., 2021b; Obermair et al., 2021) or peptide dissociation (Rappazzo et al., 2020) to assess peptide-MHC binding.

In addition to experimental advances, computational approaches for peptide-MHC binding prediction have advanced markedly over the past decade. These developments are due to algorithmic advances (O'Donnell et al., 2020; Racle et al., 2019; Reynisson et al., 2020; Zeng and Gifford, 2019) and the availability of large, high-quality training data (Abelin et al., 2017,

2019; Rappazzo et al., 2020; Reynisson et al., 2020). However, despite the improvements in
predicting peptide binding to MHC in a broad sense, the predictive power for individual
peptides often remain imperfect relative to experimental measurements (Rappazzo et al., 2020;
Zhao and Sher, 2018).
Here we present a yeast display approach to directly assess peptide-MHC binding for
large collections of defined peptide antigens, by adapting a previously described MHC-II
platform (Rappazzo et al., 2020) to screen whole viral proteomes for MHC-II binding in high-
throughput. We utilize this approach to screen the full proteome of SARS-CoV-2, a present,
global threat to public health. We additionally apply this approach to screen proteomes from
serotypes 1-4 of dengue viruses, in which antibody dependent enhancement results in more
severe disease upon second infection with a different serotype of the virus (Guzman et al.,
2016), and thus represents a potential important application area for T cell-directed
therapeutics.
95
Results
Generation of yeast display libraries for profiling the SARS-CoV-2 proteome
Previous studies have reported the use of yeast-displayed MHC-II for characterizing
peptide-MHC and pMHC-TCR interactions (Birnbaum et al., 2014, 2017; Rappazzo et al., 2020).
We adapted MHC-II yeast display constructs (Rappazzo et al., 2020) to generate a defined
library of peptides that cover the SARS-CoV-2 proteome to assess them for MHC binding. To
compare SARS-CoV-2 with a related coronavirus, we also included peptides from the spike and
nucleocapsid proteins from SARS-CoV.
Each protein was windowed into peptides of 15 amino acids in length, with a step size of
1 **(Figure 1a)**, and each peptide was encoded in DNA and cloned in a pooled format into yeast
vectors containing MHC-II proteins. The generated library was linked to three MHC-II alleles:
HLA-DR401 (HLA-DRA1*01:01, HLA-DRB1*04:01), HLA-DR402 (HLA-DRA1*01:01, HLA-
DRB1*04:02), and HLA-DR404 (HLA-DRA1*01:01, HLA-DRB1*04:04). Yeast were formatted as
previously described (Rappazzo et al., 2020), where a flexible linker connecting the peptide and
MHC contains a 3C protease site and a Myc epitope tag, which can be used for selections
(Figure 1a). The final library contained 11,040 unique peptides, with 99% of the designed
peptides present in each cloned yeast library, as assessed by next-generation sequencing.
113
Strategies for selecting defined libraries
To enrich for peptide binders, iterative selections were performed as previously
described (Rappazzo et al., 2020) (**Figure 1a**): the library is first incubated with competitor
peptide and 3C protease, which cleaves the covalent linkage between peptide and MHC,
followed by the addition of HLA-DM at lower pH. These conditions allow for the encoded
peptide to be displaced from the peptide-binding groove. The Myc epitope tag is proximal to
the peptide, which can be identified via incubation with an anti-epitope tag antibody followed
by enrichment via magnetic bead selection if the yeast-expressed peptide remains bound to the
MHC after the peptide exchange reaction.
Three rounds of selection were iteratively performed. Representative enrichment of
yeast expressing Myc-tagged peptides can be seen in **Figure 1c** ("undoped library"), for the
library displayed by HLA-DR401. Here the pre-selection Myc-positive population starts at 29.3%

and quickly converges, with 65.0% positive in the pre-selection Round 2 population and 74.1%
in the pre-selection Round 3 population.

128   Given the rapid convergence of the library, we performed a second set of selections in
which we doped the defined library into a randomized, null library to enable a greater degree of
enrichment as compared to non-binding peptides. The null library was generated by fully
randomizing ten amino acids in the peptide region of the peptide-MHC-II construct while fixing
three amino acids to encode stop codons. This library provides a baseline population of yeast
which should not express pMHC, and therefore not enrich in our selections. We doped our
defined peptide library into a 500-fold excess of null library, such that each peptide member
was represented at approximately the same frequency (**Figure 1b**). The null library provides
baseline competition, which true binders must enrich beyond, and increases the stringency of
the enrichment task.

138   We performed four rounds of selection on the doped library. Because of the excess of
null yeast, the initial pre-selection stain is low (1.6%) compared to the initial undoped library
(**Figure 1c**). This staining enriched over the first three rounds of selection, reflective of the
stringency of the task and clarity of enrichment. This is in contrast to the initial undoped library,
which began with a much higher pre-selection stain, with a lower fold-change in staining over
rounds of selection. The low frequency of each member in the starting doped library, however,
increases the likelihood of stochastic dropout for any given member.

Analysis of selection data
147   After selections, peptide identities were determined through deep sequencing of
enriched yeast populations, providing us with a dataset comprised of positive enrichment over
four rounds of selection from the doped library and both positive and negative enrichment for
three rounds of selection from the undoped library (**Supplemental Data**). **Supplemental Figure**
1 shows the correlation between defined library members on HLA-DR401. As expected, the
unselected library correlated poorly with post-selection rounds. Consistent with the observed
staining (**Figure 1c**), the doped library essentially converged after Round 3, making it likely that
changes between Rounds 3 and 4 were due to stochastic variation rather than improved
binding. Similarly, the undoped library appears converged following Round 2.

156   Next, we established metrics for enrichment for each mode of selection. Given the high
starting frequency of members in the undoped library, we classify enrichment based on fold
change between Round 1 and Round 2, and we define criteria for enriched yeast in the
undoped library as making up a higher fraction of reads following Round 2 compared to Round
1. In contrast, in the doped library, members start at low frequencies, and we define
enrichment based on presence above a threshold in Round 3 of selection, specifically as having
greater than or equal to 10 reads following Round 3. **Figure 2b** illustrates the correspondence
between enrichment metrics in the doped and undoped library for the library on HLA-DR401.
Of the 11,040 peptides in the library, 2,467 enriched in both the doped and undoped libraries
displayed by HLA-DR401 (**Figure 2a**). An additional 1,252 enriched in the doped library only and
797 enriched in the undoped library only.

167   In further data analysis, we identified a potentially confounding factor caused by the
construct design itself: the N-terminus of the peptide includes an extra alanine to ensure
consistent cleavage between the construct and its signal peptide, while the C-terminus of the

peptide is connected to the MHC construct via a Gly-Ser linker (**Figure 1a**). Since each 15mer
peptide is flanked by invariant sequences that are not present in the native sequences, it is
possible that individual peptide-construct fusions may produce high-affinity peptide binding
registers that do not occur natively. The small residues comprising the Gly-Ser linker are
favorable anchor residues at position 9 for each MHC allele in our study (Abelin et al., 2019;
Rappazzo et al., 2020; Reynisson et al., 2020). As a result, there are three non-native registers in
which a peptide could utilize a linker residue as a position 9 anchor without using a linker
residue at the position 6 anchor, which does not favor Gly or Ser (Abelin et al., 2019; Rappazzo
et al., 2020; Reynisson et al., 2020). In setting a threshold for high confidence enrichment, we
therefore sought to exclude peptides which rely on non-native linker residues in the groove for
enrichment.
Because the library is designed with a step size of one, we utilized redundancy between
adjacent peptides to determine high-confidence binders, including adjusting for peptides which
enrich with linker residues in the groove. To do this, we implement a smoothing method,
examining overlapping peptides for shared enrichment behavior. Classically, the strongest
determinant of peptide affinity for an MHC is the nine amino acid stretch sitting within the
peptide-binding groove (Jones et al., 2006; Stern, 1994), although proximal peptide flanking
residues can also affect binding (Lovitch et al., 2006; O'Brien et al., 2008; Zavala-Ruiz et al.,
2004). In our libraries, a given 9mer is present in seven overlapping 15mer peptides, and we
calculate how many of these seven 15mers have enriched. This calculation is shown
schematically in **Supplemental Figure 2a** with toy sequences and applied to enrichment data
for SARS-CoV-2 nucleocapsid on HLA-DR401 in **Supplemental Figure 2b**. Sequences with good
9mer cores should enrich along with neighboring sequences with the same 9mer sequence. In
contrast, sequences which enrich spuriously or due to linker sequence in the peptide groove or
other stochastic factors should have few neighbor sequences also enriching. Thus, we define a
cutoff for high confidence 9mer enrichment of five out of seven 9mer-containing sequences
enriching. This cutoff tolerates some stochastic dropout, while being above the probable sets of
three peptides that could enrich only with the Gly-Ser in the Position 9 pocket. Of the 2,467
peptides which enriched in both the doped and undoped libraries for HLA-DR401, 1,791 also
contain a 9mer sequence which enriched in five or more peptides of the seven neighboring
sequences containing it (**Figure 2a**), with 676 peptides enriching in both doped and undoped
libraries but not containing a 9mer core enriched in five or more peptides, and 788 15mers
containing a 9mer which enriched in five or more peptides but enriched in zero or one of the
doped and undoped libraries. These full relationships are captured in Venn diagrams in
Supplemental Figure 3 for all three MHC alleles studied here.
205
Sequence motifs of enriched peptides are consistent with known binders
To examine the 9mer core motifs of enriched peptides, we utilized a position weight
matrix method to infer the peptide register and generated visualizations of the 9mer cores
using Seq2Logo (Thomsen and Nielsen, 2012). **Figure 2c** shows a sequence logo of the aligned
9mer cores from the 2,467 15mer peptides which enriched on HLA-DR401 in both doped and
undoped libraries. The peptide motif is consistent with previously reported motifs for HLA-
DR401 (Abelin et al., 2019; Rappazzo et al., 2020): hydrophobic amino acids are preferred at P1,
acidic residues at P4, polar residues at P6, and small residues at P9. We also observe some

preference for glycine at P8 in the sequence logo, which is potentially an artifact of non-native
registers with linker at P8 and P9.
The other alleles used in the study, HLA-DR402 and HLA-DR404, have polymorphisms in
their peptide binding groove sequences as compared to HLA-DR401, which affect binding
preferences. HLA-DR401 differs from HLA-DR402 at four amino acids and from HLA-DR404 at
two amino acids, with all polymorphisms located in the beta chain. HLA-DR402 and HLA-DR404
share an amino acid distinct from HLA-DR401 affecting the P1 pocket (Gly86Val), resulting in a
preference for smaller hydrophobic residues (**Figure 3a**). Three polymorphisms in HLA-DR402
affect P4, P5, and P7 compared to HLA-DR401 (Leu67Ile, Gln70Asp, and Lys71Glu), while HLA-
DR404 has only one (Lys71Arg). Sequence logos for HLA-DR402 and HLA-DR404 are consistent
with previously reported motifs and MHC polymorphisms (**Supplemental Figure 4**). We observe
less P4 preference compared to the motif of HLA-DR402 binders enriched from a randomized
yeast display peptide library (Rappazzo et al., 2020), albeit consistent with mass spectrometry-
generated motifs which also showed minimal P4 preference for HLA-DR402 (Abelin et al.,
2019).
To probe the diminished P4 peptide preference observed for HLA-DR402 and HLA-
DR404, we examined differences between the compositions of randomized and defined
libraries. We hypothesized that skewed amino acid abundances in nature, which are reflected
in the defined library, could result in an apparent diminished amino acid preference. Indeed,
three of the most preferred P4 residues for binding HLA-DR402, Trp, His, and Met (Rappazzo et
al., 2020), are all low abundance in the SARS-CoV-2 proteome (Trp 1.1%, His 1.9%, Met 2.2%). In
comparison, a randomized peptide library for HLA-DR402 (Rappazzo et al., 2020) had a higher
representation of these amino acids (Trp 3.8%, His 2.9%, Met 3.8%). Additionally, the
randomized library had approximately nine thousand-fold more members than the defined
library, providing more instances of all amino acids. The low abundance and
underrepresentation of these amino acids likely underlies the apparent lack of amino acid
consensus at P4 in enriched peptides. Interestingly, Arg and Lys, which have also been reported
as preferred HLA-DR402 P4 residues, are more abundant than Trp, His, and Met in the SARS-
CoV-2 proteome (Arg 3.4% and Lys 5.9%; compare to Arg 9.7%, Lys 4.0% in the random library),
but still show less representation at P4 in the defined library enriched peptides compared to
the random library-enriched peptides. These differences in motifs between randomized and
defined libraries highlight the utility of randomized libraries for downstream applications such
as training prediction algorithms. Approaches influenced by amino acid abundance in nature,
such as defined libraries and mass spectrometry approaches, could inadvertently bias against
possible binders because of absence of amino acids in their null distribution, rather than true
binding preference.
Next, we wanted to examine the distribution of peptides among the possible 9mer
registers along each 15 amino acid sequence. Based on our register inference, of the 2,467
enriched peptides from the HLA-DR401 library, 1,610 peptides bound native 9mer cores
without using any linker sequence residues in the 9mer core, which is consistent with
theoretical ratios of possible native and non-native cores for a given 9mer (**Supplemental
Data**). The peptides with predicted native 9mer cores were approximately equally distributed
between possible registers, with the exception of the N-terminal register, which had one-third
fewer peptides. This register had only a single N-terminal flanking residue (the fixed Ala), which

is likely disfavored. We also identified a register with the linker filling positions 6 through 9 in
the groove, though this is the least frequent register, with 118 peptides.
Because the library was designed with step size of one, many of the 9mer cores will be
repeated among neighboring peptides. Of the 1,610 HLA-DR401 peptides which enriched using
a native 9mer core, there are 563 unique 9mer cores identified through register-inference.
Table 1 summarizes enrichment for each protein included in the library, highlighting the
number of 15mers which enriched in both the doped and undoped libraries, the number of
unique native 9mer cores, and the number of 15mers containing a 9mer enriched in at least
five of seven overlapping peptides.
267
MHC-specific relationships can be observed in binding to library peptides: MHC binding to SARS-
CoV and SARS-CoV-2 spike proteins
To further explore relationships between the MHCs studied here and their virally-
derived peptide repertoires, we compared the binding of SARS-CoV-2 and SARS-CoV spike
proteins to all three MHC alleles. Sequence alignment of these three MHC alleles is shown in
Figure 3a, with polymorphic regions highlighted on an HLA-DR401 structure (adapted from PDB
1J8H). Interplay between viral conservation and binding are illustrated in **Figure 3b**, highlighting
conserved regions of the proteome in black and binders to each allele in grey, red, and blue.
Regions are highlighted where sequences enrich in overlapping peptides; that is, for each 9
amino acid stretch along the proteome, we calculated how many of the seven 15mer peptides
enrich in the yeast display assay, and if a 9mer enriched five or more times, it is marked as a hit.
Specific examples of these relationships are probed in **Figure 3c, d, and e**, where individually
enriched 15mer sequences are represented as horizontal lines above 15mer stretches in the
proteome. Bolded 9mers are identified through register inference as consensus binding cores
for these peptides. Only 15mers which contain the bolded 9mer are included in this
representation. Non-conserved amino acids within this 9mer are highlighted in yellow.
Figure 3c illustrates a region that is not conserved between SARS-CoV-2 and SARS-CoV,
where the SARS-CoV-2 peptides containing the core IYQAGSTPC are enriched for binding to all
three MHCs, but mutations, including at both P1 and P4 to Proline, discourage binding of the
aligned SARS-CoV peptide. **Figure 3e** illustrates a core that is conserved between SARS-CoV and
SARS-CoV-2, which can bind only to HLA-DR401, but not to HLA-DR402 or HLA-DR404, likely due
to the size of the P1 hydrophobic residue and, for HLA-DR402, the acidic P4 residue. **Figure 3d**
illustrates relationships between both viral conservation and MHC preference. In **Figure 3d**, the
SARS-CoV peptides containing the core IKNQCVNFN can bind to all three alleles. However, the
aligned SARS-CoV-2 peptides containing the core VKNKCVNFN do not bind to HLA-DR401, likely
because of the less preferable P1 Valine and basic P4 Lysine, but can bind to HLA-DR402, which
prefers these residues. These peptides can bind to HLA-DR404, although only four of the
adjacent peptides containing this core enrich, which is below the cutoff of five or more, and
since no other adjacent peptides enriched, this would not have been classified as a binder
(reflected in **Figure 3b**). This marginal, but below-threshold binding is logical, given that the P4
pocket for HLA-DR404 is similar to HLA-DR401, which does not prefer P4 Lysine, but HLA-DR404
has the same P1 binding pocket as HLA-DR402, which both prefer the P1 Valine in the SARS-
CoV-2 peptide.
301

Comparing enriched sequences and algorithm predictions
Next, we compared our direct experimental assessments with results from
computational MHC binding predictions. Prediction algorithms allow for rapid computational
screening of potential peptide binders (Abelin et al., 2019; Reynisson et al., 2020), although
they can contain systemic biases (Rappazzo et al., 2020). To test the outputs of our direct
assessment approach and computational prediction algorithms, we assessed binding of several
peptides using a fluorescence polarization competition assay to determine IC_{50} values, as
described previously (Rappazzo et al., 2020; Yin and Stern, 2014). Yeast-formatted peptides
(Ala+15mer+Gly+Gly+Ser) from SARS-CoV-2 spike protein were run through NetMHCIIpan4.0
for binding to HLA-DR401, with binders defined as having ≤ 10% Rank (Eluted Ligand mode).
Yeast display binders to HLA-DR401 were defined via the stringent criteria of 1) enriching in
both in doped and undoped selections, and 2) containing a 9mer that enriched in five or more
of the overlapping seven 15mers. 15mers were selected such that they could contain a
maximum overlap of 8 amino acids with other selected peptides, to avoid selecting peptides
with redundant 9mer cores. An length-matched version of the commonly studied Influenza A
$HA_{306-318}$ peptide (APKYVKQNTLKLATG) known to bind HLA-DR401 (Hennecke and Wiley, 2002;
Rappazzo et al., 2020) was included as a positive control, along with sequences that yeast
display and NetMHCIIpan4.0 both classified as either binders or non-binders. **Supplemental**
Figure 5 shows a comparison of yeast-enriched and NetMHCpan4.0 predicted binders, with
boxed sequences selected for testing by fluorescence polarization.
The resulting fluorescence polarization IC_{50} data from the native 15mer peptides are
shown in **Table 2** and **Supplemental Figure 6**. Peptides which both enriched in yeast display and
were predicted by NetMHCIIpan4.0 to bind ('Agreed Binders') all showed IC_{50} values consistent
with binding, each with IC_{50} < 2.2 μM. Similarly, peptides which were agreed non-binders
showed no affinity for HLA-DR401, with IC_{50} > 50 μM.
All 8 'Yeast-Enriched Binders', which enriched in the yeast display assay but were not
predicted to bind via NetMHCIIpan4.0, showed some degree of binding, with IC_{50} values
distributed from 14 nM (higher affinity than the HA control peptide) to 18 μM (weak, but
measurable, binding). Retrospectively, the weakest two binders appear to be enriching in the
yeast assay using the peptide linker or have a binding core offset from center. Interestingly,
NetMHCIIpan4.0 predictions on the peptides identified via yeast display proved highly sensitive
to the length or content of the flanking sequences: if we repeat predictions on only the antigen-
derived 15mer sequences without the flanking sequences, NetMHCIIpan4.0 recovers four of its
former false negative peptides (**Table 3**; peptides listed at the top in each section of the table).
We will refer to these four peptides as 'flank-sensitive centered peptides', as they each have
the consensus 9mer core centered in the peptide.
To further investigate the relationship with flanking residues, we selected five additional
peptides ('offset peptides') matching three criteria; these offset peptides were 1) enriched in
the yeast display assay, 2) share an overlapping core with the four flank-sensitive centered
peptides, but are 3) not predicted by NetMHCIIpan4.0 to be binders (either with or without
invariant flanking sequence added). All five offset peptides have their predicted cores offset by
1-2 amino acids from center, leaving at minimum 1 amino acid on both ends of the 9mer core
for each peptide. All five offset peptides exhibit some binding, with IC_{50} values below 13 μM,
although each peptide is lower affinity than its overlapping centered counterpart, illustrating

effects of flanking residues on peptide binding, although some over-estimation of these effects
in NetMHCIIpan4.0 predictions are present.
We tested three 'NetMHC-Predicted Binders', which were predicted to bind by
NetMHCIIpan4.0, but were not enriched (nor did any neighboring sequences within an offset of
4 amino acids) in the yeast display assay (**Table 2**). Of these, one bound to HLA-DR401 (IC$_{50}$ 475
nM), while two showed minimal binding with IC50 > 35 µM, which is above the maximum 20
µM concentration tested. All three were predicted by NetMHCIIpan4.0 to bind with or without
the invariant flanking sequences (Eluted ligand mode % Rank: 5.7, 4.1, 8.7 (with flanking
residues) and 2.3, 0.6, 7.0 (without flanking residues), for ELDKYFKNHTSPDVD,
LQSYGFQPTNGVGYQ, and KTQSLLIVNNATNVV, respectively).
Of the eight 'Yeast-Enriched Binders' in **Table 2**, six contain cysteine residues, which
have been shown to be systematically absent from other datasets, including those from mono-
allelic mass spectrometry (Abelin et al., 2019; Barra et al., 2018), yet present in yeast display-
derived datasets (Rappazzo et al., 2020). To test for non-specific binding due to cysteine, two
cysteine-containing 'Agreed Non-Binders' were also tested and showed no affinity for HLA-
DR401, suggesting that cysteine itself is not causing non-specific binding. In the fluorescence
polarization dataset, the highest affinity binder (14 nM) contained cysteine and was missed by
NetMHCIIpan4.0 predictions (Eluted ligand mode % Rank: 71 (with flanking residues) and 28
(without flanking residues)).
The relationship between measured IC$_{50}$ values and NetMHCIIpan4.0 predicted values
for all 15mer SARS-CoV-2 spike peptides tested is shown in **Figure 4** and **Supplemental Figure 7**.
367
Yeast display approach can be used to compare dengue serotypes for MHC binding
Defined yeast display libraries can generate data for diverse objectives. Dengue viruses
typically cause most severe disease after a second infection with a serotype different from the
first infection, due to antibody dependent enhancement (Guzman et al., 2016), which makes T
cell-directed therapeutics a potentially attractive means of combatting disease. To profile and
compare MHC binding across serotypes, we generated libraries containing 12,672 dengue-
derived peptides, covering the entire proteomes of dengue serotypes 1-4. These libraries were
on HLA-DR401 and HLA-DR402 and had coverage of 98% and 96% of the dengue library
members after construction, respectively.
Peptides from homologous regions of the four dengue serotypes have different MHC
binding ability, as illustrated in **Figure 5a** for binding to HLA-DR401. The proteins encoded in the
dengue genome are indicated along the horizontal axis (C: capsid; M: membrane; E: envelope;
NS: nonstructural proteins). Peptides that enriched in the yeast display assay are marked by a
line (serotype 1 in blue, serotype 2 in purple, serotype 3 in red, and serotype 4 in grey). The
proteome is smoothed to 9 amino acid stretches (as in **Figure 3b**), with a given 9 amino acid
region marked as a hit if five or more of the seven adjacent peptides enrich. For each 9mer, the
maximum number of serotypes with a conserved identical 9mer at that position is indicated at
the top in black.
These data can reveal relationships between conservation and binding ability. **Figure 5b-**
d shows enrichment data for individual 15mer peptides, with consensus inferred 9mer cores in
bold and non-conserved amino acids in these cores highlighted in yellow, as in **Figure 3c-e**.
Conserved cores which show binding ability (**Figure 5c**) may be ideal T cell targets. However,

the permissiveness of the binding groove allows for peptides to bind that have mutations at the
anchors, such as in NS5 (**Figure 5d**), where P4 Asn and P4 Met both allow binding. Interestingly,
the serotype 3 core (LASNAICSA) only enriched in four peptides, which is below our described
cutoff for high-confidence peptide cores. However, three adjacent peptides enriched and
register-inference for these peptides identifies the non-native, linker-containing version of the
LASNAICSA core as binding in the MHC-binding groove. This results in an adjacent 9mer being
highlighted as a binder in this region (**Figure 5a**) because overlapping 15mers enrich in five or
more of the seven adjacent peptides. With this in mind, care must be taken for core
identification in enriched regions and can be aided by coupling enrichment with register-
inference of enriched peptides. Further, we can also see relationships between conservation
and binding in non-conserved regions, such as in the envelope protein (**Figure 5b**) with the
mutations in serotype 3 enabling binding.
402
Discussion
CD4[+] T cells responses play important roles in infection, autoimmunity, and cancer. By
extension, understanding peptide-MHC binding is critical for identifying and engineering T cell
epitopes. Here we present an approach to directly assess defined libraries of peptides covering
whole pathogen proteomes for binding to MHC-II proteins. We examine alternative modes of
selection and utilize overlapping peptides to determine high-confidence binders. We
demonstrate the utility of this approach by identifying binders that are missed by prediction
algorithms, highlighting a prediction algorithm bias against cysteine-containing peptides and
sensitivity to peptide flanking residues (**Table 2** and **Table 3**). Finally, this approach can be
utilized for different objectives, including comparing binding to multiple MHC alleles (**Figure 3**)
or comparing peptides from related pathogen sequences for MHC-II binding (**Figure 5**).
This approach for direct assessment shows benefit compared to prediction algorithms
for identifying binders, particularly for finding weak peptide binders. The overlapping peptides
in our library were useful for identifying enriched cores, especially when combined with register
inference to identify consensus cores shared between these overlapping peptides.
NetMHCIIpan4.0 exhibits a sensitivity to length and register, which may cause users to miss
binders, albeit potentially of lower affinity. Of the overlapping peptides we tested to study this
phenomenon, NetMHCIIpan4.0 correctly ranked the affinities of the overlapping peptides
(**Table 3**), but missed binders. **Supplemental Figure 5** also highlights the sensitivity of
NetMHCIIpan4.0 to flanking sequences, where neighboring peptides with shared cores often
are not predicted to bind, resulting in fewer clusters of peptides in **Supplemental Figure 5**.
Design of defined libraries with sources of redundancy, such as overlapping peptides,
was critical for determining binders with higher degrees of confidence and allowed us to apply
stringent cutoffs for individual peptides. Overlapping peptides allowed us to account for
construct-specific confounding effects, such as the peptides binding using non-native residues
in the linker. Future iterations can change the sequence of the linker, such as defining favorable
P-1 and P10 anchors to fix the register (Rappazzo et al., 2020), although these adaptations
would likely require MHC-specific knowledge in advance and may need to be altered for
different MHCs. Additionally, the engineered redundancy and multiple modes of selection
result in hyperparameters that can be tuned to meet users' stringency requirements, such as

defining different thresholds for calling individual 15mer binders or alternative integration of
overlapping binders.

Further, this approach can be used to study MHC binding between similar viruses, as
done with the dengue proteomes and the spike proteins from SARS-CoV-2 and SARS-CoV,
highlighting regions where mutations disrupt binding as well as regions where binding is
unperturbed. This method can also be rapidly adapted to study future sequences if pathogens
evolve over time.

As experimental approaches and computational approaches continue to co-develop,
they present complementary benefits. Though this platform allows for rapid assessment of
peptide-MHC binding, the speed of computational prediction surpasses experimental
approaches. NetMHCIIpan4.0 prediction and yeast display selections identified sets of non-
overlapping misses, highlighting a utility for both. Additionally, all agreed binders and non-
binders matched fluorescence polarization results, suggesting a consensus of yeast display
enrichment and algorithmic prediction provide high-confidence results. Approaches such as
yeast display assessment can be used to complement computational approaches, such as for
identifying cysteine-containing peptides which are still under-predicted by algorithms. Similarly,
prediction algorithms can be trained using large, quality datasets to account for biases. In
another application, our platform to assess peptide-MHC binding can be used to design high-
throughput assays to test peptide immunogenicity in clinical samples (Klinger et al., 2015;
Snyder et al., 2020).

Defined yeast display peptide libraries can also be readily applied to identification of T
cell ligands and present an opportunity for identifying unknown ligands from orphan TCRs
known to respond to a proteome of interest (Birnbaum et al., 2014; Gee et al., 2018). Indeed, as
DNA synthesis and sequencing continue to advance, defined peptide libraries expanding
beyond viral proteomes to covering whole bacterial or human proteomes will be possible, and
could present opportunities for investigating autoimmune diseases, which frequently have
strong MHC-II associations (Karnes et al., 2017). Such tools would be rich resources for
identifying both peptide-MHC binders and TCR ligands.

Methods
Library design and creation
Yeast display libraries were designed to cover all 15mer sequences within a given
proteome, with step size one. Reference proteomes used in creating defined libraries were
accessed from Uniprot, with the following Proteome IDs. SARS-CoV-2: UP000464024, SARS-CoV:
UP000000354, dengue serotype 1: UP000002500, dengue serotype 2: UP000180751, dengue
serotype 3: UP000007200, dengue serotype 4: UP000000275. The dengue proteome is
expressed as a single polypeptide, and peptides were generated from that contiguous stretch.
Each library peptide is encoded in DNA space, with specific codons selected randomly
from possible codons, with probabilities matching yeast codon usage (GenScript Codon Usage
Frequency Table). The DNA-encoded peptide sequences were flanked by invariant sequences
from the yeast construct for handles in amplification and cloning, and the DNA oligonucleotide
sequences were ordered from Twist Bioscience (South San Francisco, CA), with maximum
length of 120 nucleotides. The DNA oligo pool was amplified in low cycle PCR, followed by
amplification with construct DNA using overlap extension PCR. This extended product was
assembled in yeast with linearized pYal vector at a 5:1 insert:vector via electroporation with
electrocompetent RJY100 yeast.
HLA-DR401 and HLA-DR402 libraries were generated using previously described vectors
(Rappazzo et al., 2020) which contain mutations from wild type Metα36Leu, Valα132Met,
Hisβ33Asn, and Aspβ43Glu to enable proper folding without disrupting TCR or peptide contact
residues (Birnbaum et al., 2017). HLA-DR404 was generated using the same stabilizing
mutations.
The previously described null library (Dai et al., 2021) was generated with a peptide
encoded as "NNNTAANNNNNNNNNNTAGNNNNNNNNNNNNNNTGANNNNNN", where "N" indicates
any nucleotide and encodes ten random amino acids and three stop codons. This library was
similarly generated in yeast using electrocompetent RJY100 yeast.
487
Peptide visualizations and predictions
Data visualizations of viral conservation and enrichment were generated using custom
scripts. For each 9mer stretch in a protein of interest, there are seven 15mer sequences that
overlap and contain that 9mer. We calculate how many of these seven 15mers enriched in both
the doped and undoped libraries. If five or more of the seven 15mers enriched, that stretch is
marked as a 'hit'. To examine conservation between viruses, viral proteins are aligned using
ClustalOmega (Madeira et al., 2019). Aligned 9mer stretches are compared between viruses
and identical stretches are considered conserved. Hits are determined individually for each
virus before merging, such that gaps in sequence alignments do not affect calculations of
enrichment for a given virus.
Representations of 15mer hits (as in **Figure 3**, **Figure 5** and **Supplemental Figure 5**) were
generated using in-house scripts, such that a 15mer that enriched in both the doped and
undoped library was marked as a horizontal line above the relevant 15mer sequence. Only
15mers containing the bolded 9mer in **Figure 3** and **Figure 5** were included.
NetMHCIIpan4.0 webserver was used for computational predictions (Reynisson et al.,
2020), where a binder is defined as having a predicted percent rank ≤ 10%, as defined in the
webserver instructions.

505
506   *Yeast library selections*
507   Library selections were consistent with previous peptide-MHC-II yeast display
dissociation studies (Dai et al., 2021; Rappazzo et al., 2020). Yeast were washed into pH 7.2 PBS
to a concentration with 1 µM 3C protease and incubated at room temperature for 45 minutes.
Yeast were then washed into 4 °C acid saline (150mM NaCl, 20mM citric acid, pH5) with 1 µM
HLA-DM and incubated at 4 °C overnight. Each step takes place in the presence of competitor
peptide (HLA-DR401: $HA_{306-318}$ PKYVKQNTLKLAT, 1 µM; HLA-DR402: $CD48_{36-51}$
FDQKIVEWDSRKSKYF, 5 µM; HLA-DR404: NKVKSLRILNTRRKL, 5 µM (Vita et al., 2019)). Non-
specific binders are removed by incubating yeast with anti-AlexaFluor647 magnetic beads and
flowed over a magnetic Milltenyi column at 4 °C. A positive selection follows, comprised of
incubation with anti-Myc-AlexaFluor647 antibody (1:100 volume:volume) and anti-
AlexaFluor647 magnetic beads (1:10 volume:volume) and flowed over a Milltenyi column on a
magnet at 4 °C, such that yeast with bound peptide are retained on the column. These yeast
are eluted, grown to confluence in at 30 °C in SDCAA media (pH 5), and sub-cultured in at 20 °C
SGCAA media (pH 5) at OD600=1 for two days. The first round of selections of doped libraries
were conducted on 180 million yeast (SARS-CoV-2 library) or 400 million yeast (dengue library)
to ensure at least 20-fold coverage or peptides. Subsequent rounds of doped library selection,
and all rounds of undoped library selections, were performed on 20-25 million yeast.
524
525   *Library sequencing and analysis*
526   Libraries were deep sequenced to determine their composition after each round of
selection. Plasmid DNA was extracted from ten million yeast from each round of selection using
the Zymoprep Yeast Miniprep Kit (Zymo Research), following manufacturer instructions.
Amplicons were generated through PCR, covering the peptide sequence through the 3C cut site.
A second PCR round was performed to add i5 and i7 sequencing handles and in-line index
barcodes unique to each round of selection. Amplicons were sequenced on an Illumina MiSeq
using paired-end MiSeq v2 300bp kits at the MIT BioMicroCenter.
533   Paired-end reads were assembled using PandaSeq (Masella et al., 2012). Peptide
534   sequences were extracted by identifying correctly encoded flanking regions, and were filtered
535   to ensure they matched designed members of the library or the randomized null construct
536   encoding, providing a stringent threshold for contamination and PCR and read errors.
537
538   *Register inference and sequence logos*
539   The 9mer core of enriched sequences was inferred using an in-house alignment
algorithm. In this approach, we utilize a 9mer position weight matrix (PWM), which we assess
at different offsets along the peptide. We one-hot encode sequences and pad with zeros on the
C-terminus of the peptide; to assess seven native registers and four non-native registers, we
pad the peptides with four zeros. Three of the non-native registers utilize the linker at the P9
anchor but not the P6 anchor, and the addition of a fourth register captures a minority set of
peptides which utilize Gly-Gly-Ser-Gly of the linker at P6 through P9 in the groove. Register-
setting is performed with zero-padded 15mers, rather than 15mers flanked by invariant
flanking residues, because the PWM would otherwise align all sequences to the invariant
region.

At the start, we randomly assign peptides to registers and generate a 9mer PWM. Over
subsequent iterations, peptides are assigned to new registers and the PWM was updated.
Assignments are random but biased, such that clusters corresponding to registers that match
the PWM are favored. Specifically, at each assignment we first take out the sequence under
consideration from the PWM. The PWM then defines an energy value for each register shift of a
given peptide, which is then used to generate a Boltzmann distribution from which we sample
the updated register shift. The stochasticity is decreased over time by raising the inverse
temperature linearly from 0.05 to 1 over 60 iterations, simulating 'cooling' (Andreatta et al.,
2017). A final deterministic iteration was carried out, where the distribution concentrates
entirely on the optimal register shift.
After register inference, sequence logo visualizations of the 9mer cores were generated
using Seq2Logo-2.0 with default settings, except using background frequencies from the SARS-
CoV-2 proteome and SARS-CoV spike and nucleocapsid proteins (Thomsen and Nielsen, 2012).
For registers with the C-terminus utilizing the C-terminal linker, the relevant linker sequence
was added to achieve a full 9mer sequence for visualizing the full 9mer core. For HLA-DR401,
distribution among registers, starting from N-terminally to C-terminally aligned in the peptide,
is: 161, 237, 227, 238, 231, 279, 237, 266, 271, 202, 118.
566
Recombinant protein expression
HLA-DM and HLA-DR401 were expressed recombinantly in High Five insect cells (Thermo
Fisher) using a baculovirus expression system, as previously described (Birnbaum et al., 2014;
Rappazzo et al., 2020). Ectodomain sequences of each chain were formatted with a C-terminal
poly-histidine purification tag and cloned into pAcGP67a vectors. Each vector was individually
transfected into SF9 insect cells (Thermo Fisher) with BestBac 2.0 linearized baculovirus DNA
(Expression Systems; Davis, CA) and Cellfectin II Reagent (Thermo Fisher), and propagated to
high titer. Viruses were co-titrated for optimal expression to maximize balanced MHC
heterodimer formation, co-transduced into Hi5 cells, and grown for 48-72 hours at 27 °C. The
secreted protein was purified from pre-conditioned media supernatant with Ni-NTA resin and
purified via size exclusion chromatography with a S200 increase column on an AKTA PURE FPLC
(GE Healthcare). To improve protein yields, the HLA-DRB1*04:01 chain was expressed with a
$CLIP_{87-101}$ peptide (PVSKMRMATPLLMQA) connected to the N-terminus of the MHC chain via a
flexible, 3C protease-cleavable linker.
581
Fluorescence polarization experiments for peptide IC_{50} determination
Peptide IC_{50} values were determined following a protocol modified from Yin & Stern (Yin
and Stern, 2014), as in Rappazzo et al (Rappazzo et al., 2020). In the assay, recombinantly
expressed HLA-DR401 is incubated with fluorescently labelled modified $HA_{306-318}$ (APRFV{Lys(5,6
FAM)}QNTLRLATG) peptide and a titration series for each unlabeled competitor peptide is
added (1.28 nM – 20 uM). A change in polarization value resulting from displacement of
fluorescent peptide from the binding groove is used to determine IC_{50} values.
Relative binding at each concentration is calculated as $(FP_{sample} - FP_{free})/(FP_{no\_comp} -$
$FP_{free})$. Here, FP_{free} is the polarization value for the fluorescent peptide alone with no added
MHC, FP_{no_comp} is polarization value for MHC with no competitor peptide added, and FP_{sample} is
the polarization value with both MHC and competitor peptide added. Relative binding curves

were then generated and fit in Prism 9.3 to the equation $y = 1/(1+[pep]/IC_{50})$, where [pep] is the
concentration of un-labelled competitor peptide, in order to determine the concentration of
half-maximal inhibition, the IC_{50} value.

Each assay was performed at 200 uL, with 100 nM recombinant MHC, 25 nM fluorescent
peptide, and competitor peptide (GenScript). This mixture co-incubates in pH 5 binding buffer
at 37 °C for 72 hours in black flat bottom 96-well plates. Competitor peptide concentrations
ranged from 1.28 nM to 20 μM, as a five-fold dilution series. Three replicates are performed for
each peptide concentration. Fluorescent peptide-only, no competitor peptide, and binding
buffer controls were also included. Our MHC was expressed with a linked CLIP peptide, so prior
to co-incubation, the peptide linker is cleaved by addition of 3C protease at 1:10 molar ratio at
room temperature for one hour; the residual cleaved 100 nM CLIP peptide is not expected to
alter peptide binding measurements.

Measurements were taken on a Molecular Devices SpectraMax M5 instrument. G-value
was 1.1 for each plate, as calculated per manufacturer instructions for each plate based on
fluorescent peptide-only wells minus buffer blank wells, with 35 mP reference for 5,6FAM
(Fluorescein setting). Measurements were made with 470 nm excitation and 520 nm emission,
10 flashes per read, and default PMT gain high.

Data Availability
All deep sequencing data are deposited on the Sequence Read Archive (SRA), with accession
codes PRJNA806475 [https://www.ncbi.nlm.nih.gov/bioproject/PRJNA806475] and
PRJNA708266 [https://www.ncbi.nlm.nih.gov/bioproject/PRJNA708266]

Code Availability
Scripts used for data processing and visualization are publicly available at
https://github.com/birnbaumlab/Huisman-et-al-2022.

Acknowledgements
We would like to thank Patrick Holec for feedback on this manuscript, and the MIT BioMicro
Center for library sequencing. This work was supported in part by the Koch Institute Support
(core) Grant P30-CA14051 from the National Cancer Institute. This work was supported by
National Institute of Health (U19-AI110495), the Packard Foundation to M.E.B., a Schmidt
Futures grant to D.K.G. and M.E.B., and a National Science Foundation Graduate Research
Fellowship to B.D.H.

Figures

628
Figure 1. Overview of library and selections. a) The defined library contains pathogen
proteome peptides (length 15, sliding window 1). Poor binding peptides are displaced with
addition of protease, competitor peptide, and HLA-DM. **b)** Schematic of doped and undoped
libraries: in the doped selection strategy, the library is added to a library of null, non-expressing
constructs. **c)** Representative flow plots showing enrichment of MHC-expressing yeast over
rounds of selection for the library containing SARS-CoV-2 and SARS-CoV peptides on HLA-
DR401.

636
Figure 2. Output of selections and analysis of selection data. a) Overview of filtering peptides
and correspondence between selection strategies for SARS-CoV and SARS-CoV-2 library on HLA-
DR401. Peptides are filtered for enrichment in both doped and undoped libraries. Further, the
relationship between these peptides and peptides which contain a 9mer that is enriched in five
or more of the seven peptides containing it is shown. **b)** Relationships between enrichment in
doped and undoped libraries. Absolute counts following Round 3 of selection of the doped
library are plotted against the log2 fold change between read fraction for peptides in Round 2
and Round 1. Data are shown for the library on HLA-DR401. **c)** Sequence logo of 2,467 peptides
that enriched in both doped and undoped selected libraries for HLA-DR401. Registers are
inferred with a position weight matrix-based alignment method. Logos were generated with
Seq2Logo-2.0.

648
Figure 3. Comparing HLA-DR401, HLA-DR402, and HLA-DR404 for binding to related Spike
proteins from SARS-CoV-2 and SARS-CoV. a) Sequence alignment showing sequence
differences in HLA-DR402 and HLA-DR404 compared to HLA-DR401 and highlighted on HLA-
DR401 structure (PDB 1J8H). Colors are: red for amino acids shared between HLA-DR401 and
HLA-DR404, green for amino acids shared between HLA-DR402 and HLA-DR404, and yellow for
amino acids different in all 3 alleles. Affected peptide positions (P1, P4, P5, P7) are colored in
blue and labeled on the structure. **b)** Conservation and enrichment of 9mer peptides from
SARS-CoV-2 and SARS-CoV Spike proteins. Conserved 9mers are indicated in black. If a 9mer
along the proteome enriched in 5 or more of the adjacent peptides containing it, its enrichment
is indicated with a vertical line with color for allele (HLA-DR401: blue; HLA-DR402: red; HLA-
DR404: grey) and opacity for virus (SARS-CoV-2: dark; SARS-CoV: light). **b-e)** Zoomed regions
show enrichment of individual 15mer peptides. Only peptides containing the bolded 9mer
sequence are shown. Amino acids in the bolded 9mer that are not conserved between SARS-
CoV-2 and SARS-CoV are highlighted in yellow.

663
664 **Figure 4. Comparing measured IC$_{50}$ values and computational prediction.** Relationship
665 between measured IC$_{50}$ values and NetMHCIIpan4.0 predicted ranks in Eluted Ligand mode (EL)
666 on invariant-flanked sequences. Data points are colored by label, and IC$_{50}$ values ≥50 μM are set
667 to 50 μM.

668
Figure 5. Conservation and enrichment of dengue virus serotypes 1-4. a) Conservation and
enrichment of 9mer peptides along four aligned dengue serotypes. All stretches of 9 amino
acids are compared across the four serotypes and conservation is indicated with a black vertical
line (i.e. 2, 3, or 4 of 4 serotypes conserved). 9mers which enriched on HLA-DR401 are also
indicated, colored by virus serotype. **b-d)** Zoomed regions, showing enrichment for individual
15mer peptides to HLA-DR401. Only peptides which contain the bolded 9mer sequence are
shown. Amino acids in the bolded 9mer that are not conserved between serotypes are
highlighted in yellow. Insets show regions which are differently conserved and enriched: **b)** non-
conserved sequences with peptides from one serotype enriched; **c)** conserved sequences
enriched across all serotypes; **d)** non-conserved sequences which are enriched.

Virus	Protein	Protein length (# of amino acids)	MHC Allele	# of 15mers	# of 9mer cores	# of smoothed 15mers
SARS-CoV	Spike	1255	HLA-DR401	324	74	221
			HLA-DR402	217	65	110
			HLA-DR404	289	61	193
SARS-CoV	Nucleocapsid	422	HLA-DR401	40	8	34
			HLA-DR402	34	13	12
			HLA-DR404	31	6	20
SARS-CoV-2	Spike	1273	HLA-DR401	305	67	221
			HLA-DR402	230	62	130
			HLA-DR404	290	64	217
SARS-CoV-2	Nucleocapsid	419	HLA-DR401	34	8	24
			HLA-DR402	33	10	15
			HLA-DR404	30	8	18
SARS-CoV-2	Replicase polyprotein 1ab	7096	HLA-DR401	1652	388	1204
			HLA-DR402	1104	325	678
			HLA-DR404	1368	350	890
SARS-CoV-2	Non-structural protein 8	121	HLA-DR401	41	10	32
			HLA-DR402	21	7	17
			HLA-DR404	32	8	19
SARS-CoV-2	Protein 7a	121	HLA-DR401	27	8	18
			HLA-DR402	7	3	0
			HLA-DR404	13	2	6
SARS-CoV-2	Non-structural protein 6	61	HLA-DR401	0	0	0
			HLA-DR402	1	1	0
			HLA-DR404	0	0	0
SARS-CoV-2	Membrane protein	222	HLA-DR401	40	7	29
			HLA-DR402	26	6	19
			HLA-DR404	23	7	21
SARS-CoV-2	Envelope small membrane protein	75	HLA-DR401	6	1	0
			HLA-DR402	7	3	0
			HLA-DR404	6	1	0
SARS-CoV-2	Protein 3a	275	HLA-DR401	22	4	11
			HLA-DR402	13	4	10
			HLA-DR404	10	2	0
SARS-CoV-2	Replicase polyprotein 1a	4405	HLA-DR401	948	228	658
			HLA-DR402	657	196	409
			HLA-DR404	865	222	582
SARS-CoV-2	ORF10 protein	38	HLA-DR401	6	1	6
			HLA-DR402	2	0	0
			HLA-DR404	5	1	5
SARS-CoV-2	Protein non-structural 7b	43	HLA-DR401	0	0	0
			HLA-DR402	0	0	0
			HLA-DR404	0	0	0
SARS-CoV-2	Uncharacterized protein 14	73	HLA-DR401	8	4	6
			HLA-DR402	20	5	16
			HLA-DR404	22	4	21
SARS-CoV-2	Protein 9b	97	HLA-DR401	29	7	27
			HLA-DR402	35	6	31
			HLA-DR404	37	9	34

680 **Table 1.** Summary of enriched peptides for each source protein, including: the number of
681 unique 15mers which each enriched in both of the doped and undoped libraries; the number of
682 unique 9mers cores identified by register-inference in these enriched 15mers (native cores
683 only, so linker-containing inferred cores excluded); and the number of unique enriched 15mers
684 that contain 9mer sequences enriched in five or more of overlapping neighbors.

	Spike Position	Peptide+flank (A+15mer+GGS)	NetMHCIIpan4.0 Predicted Core (A+15mer+GGS)	NetMHCIIpan4.0 %Rank (A+15mer+GGS)	15mer Affinity from FP (IC$_{50}$, nM)
Agreed Binders	34-48	ARGVYYPDKVFRSSVLGGS	YYPDKVFRS	1.49	15.8
	87-101	ANDGVYFASTEKSNIIGGS	VYFASTEKS	4.28	2117
	303-317	ALKSFTVEKGIYQTSNGGS	FTVEKGIYQ	8.41	396.9
	362-376	AVADYSVLYNSASFSTGGS	YSVLYNSAS	8.36	113.7
	1015-1029	AAAEIRASANLAATKMGGS	IRASANLAA	3.13	105.4
	1112-1126	APQIITTDNTFVSGNCGGS	ITTDNTFVS	7.32	527.0
Yeast-Enriched Binders	165-179	ANCTFEYVSQPFLMDLGGS	YVSQPFLMD	64.83	14,652
	172-186	ASQPFLMDLEGKQGNFGGS	FLMDLEGKQ	20.34	123.2
	286-300	ATDAVDCALDPLSETKGGS	VDCALDPLS	32.68	521.6
	373-387	ASFSTFKCYGVSPTKLGGS	YGVSPTKLG	16.59	18,452
	469-483	ASTEIYQAGSTPCNGVGGS	IYQAGSTPC	18.22	67.7
	580-594	AQTLEILDITPCSFGGGGS	LEILDITPC	62	119.9
	739-753	ATMYICGDSTECSNLLGGS	YICGDSTEC	70.91	14.4
	920-934	AQKLIANQFNSAIGKIGGS	FNSAIGKIG	20.47	1121
NetMHC-Predicted Binders	1151-1165	AELDKYFKNHTSPDVDGGS	YFKNHTSPD	5.74	35,510
	492-506	ALQSYGFQPTNGVGYQGGS	YGFQPTNGV	4.11	454.7
	113-127	AKTQSLLIVNNATNVVGGS	IVNNATNVV	8.74	>50,000
Agreed Non-Binders	534-548	AVKNKCVNFNFNGLTGGGS	FNFNGLTGG	57.13	>50,000
	1079-1093	APAICHDGKAHFPREGGGS	ICHDGKAHF	80.47	>50,000

685
686 **Table 2.** Peptides selected for fluorescence polarization (FP) experiments for binding to HLA-
687 DR401. NetMHCIIpan4.0 predictions for HLA-DR401 binding are performed on 15mers plus
688 invariant flanking residues (N-terminal Ala, C-terminal Gly-Gly-Ser) and percent rank values
689 generated using Eluted Ligand mode. Fluorescence polarization is performed on native 15mer
690 peptides without invariant flanking residues.

Spike Position	Sequence	NetMHCIIpan4.0 Predicted Core (A+15mer+GGS)	NetMHCIIpan4.0 %Rank (A+15mer+GGS)	NetMHCIIpan4.0 Predicted Core (15mer)	NetMHCIIpan4.0 %Rank (15mer)	15mer Affinity from FP (IC50, nM)
172-186	SQP**FLMDLEGKQ**GNF	FLMDLEGKQ	20.34	FLMDLEGKQ	4.1	123.2
173-187	QP**FLMDLEGKQ**GNFK	FLMDLEGKQ	27.73	FLMDLEGKQ	12.21	8613
286-300	TDA**VDCALDPLS**ETK	VDCALDPLS	32.68	VDCALDPLS	9.8	1154
287-301	DA**VDCALDPLS**ETKC	VDCALDPLS	42.42	VDCALDPLS	22.57	4393
469-483	STE**IYQAGSTPC**NGV	IYQAGSTPC	18.22	IYQAGSTPC	5.41	67.7
467-481	DISTE**IYQAGSTPC**N	IYQAGSTPC	11.47	IYQAGSTPC	12.61	4875
471-485	E**IYQAGSTPC**NGVEG	YQAGSTPCN	39.17	YQAGSTPCN	21.81	12519
920-934	QKL**IANQFNSAI**GKI	FNSAIGKIG	20.47	IANQFNSAI	7.89	1495
921-935	KL**IANQFNSAI**GKIQ	FNSAIGKIQ	18.3	IANQFNSAI	19.79	11937

Table 3. Effects of peptide flanking sequences on NetMHCIIpan4.0 predictions for HLA-DR401 binding and measured fluorescence polarization (FP) values for overlapping peptides. Yeast display-enriched peptides that are predicted to bind by NetMHCIIpan4.0 when without flanking residues, plus offset variants of these peptides, which are not predicted to bind, with or without flanking sequence. Yeast display register-inferred consensus cores are highlighted in green. NetMHCIIpan4.0 percent rank values are generated using Eluted Ligand mode.

References

Abelin, J.G., Keskin, D.B., Sarkizova, S., Hartigan, C.R., Zhang, W., Sidney, J., Stevens, J., Lane, W.,
Zhang, G.L., Eisenhaure, T.M., et al. (2017). Mass spectrometry profiling of HLA-associated
peptidomes in mono-allelic cells enables more accurate Epitope prediction. Immunity *46*, 315–
326.

Abelin, J.G., Harjanto, D., Malloy, M., Suri, P., Colson, T., Goulding, S.P., Creech, A.L., Serrano,
L.R., Nasir, G., Nasrullah, Y., et al. (2019). Defining HLA-II ligand processing and binding rules
with mass spectrometry enhances cancer Epitope prediction. Immunity *51*, 766-779.e17.

Altmann, D.M., and Boyton, R.J. (2020). SARS-CoV-2 T cell immunity: Specificity, function,
durability, and role in protection. Sci. Immunol. *5*, eabd6160.

Andreatta, M., Alvarez, B., and Nielsen, M. (2017). GibbsCluster: unsupervised clustering and
alignment of peptide sequences. Nucleic Acids Res. *45*, W458–W463.

Barra, C., Alvarez, B., Paul, S., Sette, A., Peters, B., Andreatta, M., Buus, S., and Nielsen, M.
(2018). Footprints of antigen processing boost MHC class II natural ligand predictions. Genome
Med. *10*, 84.

Birnbaum, M.E., Mendoza, J.L., Sethi, D.K., Dong, S., Glanville, J., Dobbins, J., Ozkan, E., Davis,
M.M., Wucherpfennig, K.W., and Garcia, K.C. (2014). Deconstructing the peptide-MHC
specificity of T cell recognition. Cell *157*, 1073–1087.

Birnbaum, M.E., Mendoza, J., Bethune, M., Baltimore, D., and Garcia, K.C. (2017). Ligand
discovery for t cell receptors. US20170192011A1.

Dai, Z., Huisman, B.D., Zeng, H., Carter, B., Jain, S., Birnbaum, M.E., and Gifford, D.K. (2021).
Machine learning optimization of peptides for presentation by class II MHCs. Bioinformatics.

Gambino, F., Jr, Tai, W., Voronin, D., Zhang, Y., Zhang, X., Shi, J., Wang, X., Wang, N., Du, L., and
Qiao, L. (2021). A vaccine inducing solely cytotoxic T lymphocytes fully prevents Zika virus
infection and fetal damage. Cell Rep. *35*, 109107.

Gee, M.H., Han, A., Lofgren, S.M., Beausang, J.F., Mendoza, J.L., Birnbaum, M.E., Bethune, M.T.,
Fischer, S., Yang, X., Gomez-Eerland, R., et al. (2018). Antigen identification for orphan T cell
receptors expressed on tumor-infiltrating lymphocytes. Cell *172*, 549-563.e16.

Guzman, M.G., Gubler, D.J., Izquierdo, A., Martinez, E., and Halstead, S.B. (2016). Dengue
infection. Nat. Rev. Dis. Primers *2*, 16055.

Hennecke, J., and Wiley, D.C. (2002). Structure of a complex of the human alpha/beta T cell
receptor (TCR) HA1.7, influenza hemagglutinin peptide, and major histocompatibility complex
class II molecule, HLA-DR4 (DRA*0101 and DRB1*0401): insight into TCR cross-restriction and
alloreactivity. J. Exp. Med. *195*, 571–581.

Jiang, W., and Boder, E.T. (2010). High-throughput engineering and analysis of peptide binding
to class II MHC. Proc. Natl. Acad. Sci. U. S. A. *107*, 13258–13263.

Jones, E.Y., Fugger, L., Strominger, J.L., and Siebold, C. (2006). MHC class II proteins and disease:
a structural perspective. Nat. Rev. Immunol. *6*, 271–282.

Justesen, S., Harndahl, M., Lamberth, K., Nielsen, L.-L.B., and Buus, S. (2009). Functional
recombinant MHC class II molecules and high-throughput peptide-binding assays. Immunome
Res. *5*, 2.

Karnes, J.H., Bastarache, L., Shaffer, C.M., Gaudieri, S., Xu, Y., Glazer, A.M., Mosley, J.D., Zhao,
S., Raychaudhuri, S., Mallal, S., et al. (2017). Phenome-wide scanning identifies multiple
diseases and disease severity phenotypes associated with HLA variants. Sci. Transl. Med. *9*.

Keskin, D.B., Anandappa, A.J., Sun, J., Tirosh, I., Mathewson, N.D., Li, S., Oliveira, G., Giobbie-
Hurder, A., Felt, K., Gjini, E., et al. (2019). Neoantigen vaccine generates intratumoral T cell
responses in phase Ib glioblastoma trial. Nature *565*, 234–239.

Klinger, M., Pepin, F., Wilkins, J., Asbury, T., Wittkop, T., Zheng, J., Moorhead, M., and Faham,
M. (2015). Multiplex Identification of Antigen-Specific T Cell Receptors Using a Combination of
Immune Assays and Immune Receptor Sequencing. PLoS One *10*, e0141561.

Liu, G., Carter, B., Bricken, T., Jain, S., Viard, M., Carrington, M., and Gifford, D.K. (2020).
Computationally optimized SARS-CoV-2 MHC class I and II vaccine formulations predicted to
target human haplotype distributions. Cell Syst. *11*, 131-144.e6.

Liu, G., Carter, B., and Gifford, D.K. (2021a). Predicted cellular immunity population coverage
gaps for SARS-CoV-2 subunit vaccines and their augmentation by compact peptide sets. Cell
Syst. *12*, 102-107.e4.

Liu, R., Jiang, W., and Mellins, E.D. (2021b). Yeast display of MHC-II enables rapid identification
of peptide ligands from protein antigens (RIPPA). Cell. Mol. Immunol. *18*, 1847–1860.

Lovitch, S.B., Pu, Z., and Unanue, E.R. (2006). Amino-terminal flanking residues determine the
conformation of a peptide-class II MHC complex. J. Immunol. *176*, 2958–2968.

Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey,
A.R.N., Potter, S.C., Finn, R.D., et al. (2019). The EMBL-EBI search and sequence analysis tools
APIs in 2019. Nucleic Acids Res. *47*, W636–W641.

Masella, A.P., Bartram, A.K., Truszkowski, J.M., Brown, D.G., and Neufeld, J.D. (2012).
PANDAseq: paired-end assembler for illumina sequences. BMC Bioinformatics *13*, 31.

Mateus, J., Grifoni, A., Tarke, A., Sidney, J., Ramirez, S.I., Dan, J.M., Burger, Z.C., Rawlings, S.A.,
Smith, D.M., Phillips, E., et al. (2020). Selective and cross-reactive SARS-CoV-2 T cell epitopes in
unexposed humans. Science *370*, 89–94.

Moise, L., Gutierrez, A., Kibria, F., Martin, R., Tassone, R., Liu, R., Terry, F., Martin, B., and De
Groot, A.S. (2015). iVAX: An integrated toolkit for the selection and optimization of antigens
and the design of epitope-driven vaccines. Hum. Vaccin. Immunother. *11*, 2312–2321.

Obermair, F.J., Renoux, F., Heer, S., Lee, C., Cereghetti, N., Maestri, G., Haldner, Y., Wuigk, R.,
Iosefson, O., Patel, P., et al. (2021). High resolution profiling of MHC-II peptide presentation
capacity, by Mammalian Epitope Display, reveals SARS-CoV-2 targets for CD4 T cells and
mechanisms of immune-escape (bioRxiv).

O'Brien, C., Flower, D.R., and Feighery, C. (2008). Peptide length significantly influences in vitro
affinity for MHC class II molecules. Immunome Res. *4*, 6.

O'Donnell, T.J., Rubinsteyn, A., and Laserson, U. (2020). MHCflurry 2.0: Improved pan-allele
prediction of MHC class I-presented peptides by incorporating antigen processing. Cell Syst. *11*,
42-48.e7.

Ott, P.A., Hu, Z., Keskin, D.B., Shukla, S.A., Sun, J., Bozym, D.J., Zhang, W., Luoma, A., Giobbie-
Hurder, A., Peter, L., et al. (2017). An immunogenic personal neoantigen vaccine for patients
with melanoma. Nature *547*, 217–221.

Parker, R., Partridge, T., Wormald, C., Kawahara, R., Stalls, V., Aggelakopoulou, M., Parker, J.,
Powell Doherty, R., Ariosa Morejon, Y., Lee, E., et al. (2021). Mapping the SARS-CoV-2 spike
glycoprotein-derived peptidome presented by HLA class II on dendritic cells. Cell Rep. *35*,
109179.

Patronov, A., and Doytchinova, I. (2013). T-cell epitope vaccine design by immunoinformatics.
Open Biol. *3*, 120139.

Purcell, A.W., Ramarathinam, S.H., and Ternette, N. (2019). Mass spectrometry-based
identification of MHC-bound peptides for immunopeptidomics. Nat. Protoc. *14*, 1687–1707.

Racle, J., Michaux, J., Rockinger, G.A., Arnaud, M., Bobisse, S., Chong, C., Guillaume, P., Coukos,
G., Harari, A., Jandus, C., et al. (2019). Robust prediction of HLA class II epitopes by deep motif
deconvolution of immunopeptidomes. Nat. Biotechnol. *37*, 1283–1286.

Rappazzo, C.G., Huisman, B.D., and Birnbaum, M.E. (2020). Repertoire-scale determination of
class II MHC peptide binding via yeast display improves antigen prediction. Nat. Commun. *11*,
4414.

Reynisson, B., Alvarez, B., Paul, S., Peters, B., and Nielsen, M. (2020). NetMHCpan-4.1 and
NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif
deconvolution and integration of MS MHC eluted ligand data. Nucleic Acids Res. *48*, W449–
W454.

Rosati, E., Pogorelyy, M.V., Minervina, A.A., Scheffold, A., Franke, A., Bacher, P., and Thomas,
P.G. (2021). Characterization of SARS-CoV-2 public CD4+ αβ T cell clonotypes through reverse
epitope discovery. BioRxivorg.

Sidney, J., Steen, A., Moore, C., Ngo, S., Chung, J., Peters, B., and Sette, A. (2010). Divergent
motifs but overlapping binding repertoires of six HLA-DQ molecules frequently expressed in the
worldwide human population. J. Immunol. *185*, 4189–4198.

Snyder, T.M., Gittelman, R.M., Klinger, M., May, D.H., Osborne, E.J., Taniguchi, R., Zahid, H.J.,
Kaplan, I.M., Dines, J.N., Noakes, M.T., et al. (2020). Magnitude and dynamics of the T-cell
response to SARS-CoV-2 infection at both individual and population levels. MedRxiv.

Stern, L.J. (1994). Crystal structure of the human class II MHC protein HLA- DR1 complexed with
an influenza virus peptide. Nature *368*, 215–221.

Stopfer, L.E., Mesfin, J.M., Joughin, B.A., Lauffenburger, D.A., and White, F.M. (2020).
Multiplexed relative and absolute quantitative immunopeptidomics reveals MHC I repertoire
alterations induced by CDK4/6 inhibition. Nat. Commun. *11*, 2760.

Stopfer, L.E., Gajadhar, A.S., Patel, B., Gallien, S., Frederick, D.T., Boland, G.M., Sullivan, R.J., and
White, F.M. (2021). Absolute quantification of tumor antigens using embedded MHC-I
isotopologue calibrants. Proc. Natl. Acad. Sci. U. S. A. *118*, e2111173118.

Thomsen, M.C.F., and Nielsen, M. (2012). Seq2Logo: a method for construction and
visualization of amino acid binding motifs and sequence profiles including sequence weighting,
pseudo counts and two-sided representation of amino acid enrichment and depletion. Nucleic
Acids Res. *40*, W281-7.

Vita, R., Mahajan, S., Overton, J.A., Dhanda, S.K., Martini, S., Cantrell, J.R., Wheeler, D.K., Sette,
A., and Peters, B. (2019). The Immune Epitope Database (IEDB): 2018 update. Nucleic Acids Res.
47, D339–D343.

Yin, L., and Stern, L.J. (2014). Measurement of peptide binding to MHC class II molecules by
fluorescence polarization. Curr. Protoc. Immunol. *106*, 5.10.1-5.10.12.

Zavala-Ruiz, Z., Strug, I., Anderson, M.W., Gorski, J., and Stern, L.J. (2004). A polymorphic pocket
at the P10 position contributes to peptide binding specificity in class II MHC proteins. Chem.
Biol. *11*, 1395–1402.

Zeng, H., and Gifford, D.K. (2019). Quantification of uncertainty in peptide-MHC binding
prediction improves high-affinity peptide selection for therapeutic design. Cell Syst. *9*, 159-
166.e3.

Zhao, W., and Sher, X. (2018). Systematically benchmarking peptide-MHC binding predictors:
From synthetic to naturally processed epitopes. PLoS Comput. Biol. *14*, e1006457.