1
2
3
4
5 **Preferred synonymous codons are translated more accurately: Proteomic evidence, among-**
6 **species variation, and mechanistic basis**
7

8 Mengyi Sun and Jianzhi Zhang
9
10 Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan
11 48109, USA
12
13
14
15
*Correspondence to
Jianzhi Zhang
Department of Ecology and Evolutionary Biology
University of Michigan
4018 Biological Sciences Building
1105 North University Avenue
Ann Arbor, MI 48109, USA
Phone: 734-763-0527
Email: jianzhi@umich.edu

25
26
27
28
29 **Running title:** Preferred codons are decoded more accurately
30

31 **Keywords:** *Escherichia coli*, codon usage, tRNA, translational accuracy, translational efficiency,
32 evolution

33
34

ABSTRACT

A commonly stated cause of the widespread phenomenon of unequal uses of synonymous

codons is their differential translational accuracies. However, this long-standing translational

accuracy hypothesis (TAH) of codon usage bias has had no direct evidence beyond anecdotes.

Analyzing proteomic data from *Escherichia coli*, we observe higher translational accuracies of

more frequently used synonymous codons, offering direct, global evidence for the TAH. The

experimentally measured codon-specific translational accuracies validate a sequence-based

proxy; this proxy provides support for the TAH from the vast majority of over 1000 taxa

surveyed in all domains of life. We find that the relative translational accuracies of synonymous

codons vary substantially among taxa and are strongly correlated with the amounts of cognate

tRNAs relative to those of near-cognate tRNAs. These and other observations suggest a model

in which selections for translational efficiency and accuracy drive codon usage bias and its

coevolution with the tRNA pool.

INTRODUCTION

Eighteen of the 20 amino acids are each encoded by more than one codon, but the synonymous codons are usually unequally used in a genome (*1, 2*). Among the synonymous codons of an amino acid, those used more often than the average are referred to as preferred codons while the rest unpreferred. This phenomenon of codon usage bias (CUB), initially discovered over four decades ago from the first few determined gene sequences (*3-6*), is a result of the joint forces of mutation, genetic drift, and natural selection, but the specific selective agents have not been fully deciphered (*1, 2*). One long-standing hypothesis known as the translational accuracy hypothesis (TAH) asserts that different synonymous codons are translated with different accuracies and that CUB results at least in part from natural selection for translational accuracy (*7*). Indeed, the importance of accurate protein translation cannot be overstated, because mistranslation may lead to the loss of normal protein functions and gain of cellular toxicity (*8*) and cause severe diseases including cancer and neurodegenerative diseases (*9*). In fact, several cellular mechanisms are known to ensure the overall fidelity of protein synthesis. For example, conformational changes of the ribosome decoding center can be more efficiently induced by cognate codon-anticodon interactions than near-cognate codon-anticodon interaction (*10*), allowing discrimination against incorrect decoding. Additionally, the accuracy of many steps in translation, such as tRNA aminoacylation (*10*) and codon-anticodon matching, is enhanced by the energy-consuming kinetic proofreading (*11*). Notwithstanding, even if synonymous codons differ in translational accuracy, relatively accurate synonymous codons may not be preferentially used. This is because synonymous codons differ in other properties such as the translational elongation speed (*12, 13*); selection related to these other features(*14*) could triumph selection for translational accuracy.

Several groups have attempted to test the TAH of CUB. In particular, Akashi (*7*) developed an indirect test based on the idea that the benefit of using relatively accurate codons should be greater at evolutionarily conserved amino acid sites than unconserved sites of the same protein; consequently, a higher usage of preferred codons at conserved sites than at unconserved sites supports the TAH. While Akashi's test is positive for several model organisms investigated (*7, 15, 16*), this test does not directly compare translational accuracies of synonymous codons so cannot completely exclude alternative explanations (*7, 17*). In an early study, Precup and Parker used site-directed mutagenesis followed by peptide sequencing to show that AAU, an

unpreferred codon of Asn, is misread as Lys 4-9 times more often than is AAC, a preferred codon

of Asn, at a particular position of the coat protein gene of the bacteriophage MS2 under Asn

starvation (*18*). Similarly, Kramer and Farabaugh observed that AAU has a significantly higher

rate of mistranslation to Lys than AAC at a particular position of a reporter gene in *Escherichia*

coli (*19*). Nonetheless, Kramer and Farabaugh also observed that the unpreferred Arg codons of

CGA and CGG and the preferred Arg codons of CGU and CGC exhibited similar rates of

mistranslation to Lys (*19*). While the above experiments directly tested the TAH, they were each

based on the investigation of one amino acid site of one protein, so its genome-wide generality is

unknown. As such, a direct and global test of the TAH is needed.

Capitalizing on a proteome-wide probe of mistranslation in *E. coli* (*20*), we here provide

direct evidence that preferred codons are generally translated more accurately than unpreferred

codons. We then use the *E. coli* data to validate a sequence-based proxy for relative translational

accuracies of synonymous codons. Using this proxy, we show that the TAH of CUB is supported

in the vast majority of over 1000 diverse taxa surveyed, but that the relative translational

accuracies of synonymous codons vary substantially among taxa. We find that the relative

translational accuracy of a synonymous codon is strongly correlated with its cognate tRNA

abundance relative to near-cognate tRNA abundance. These and other results suggest a model in

which selections for translational efficiency and accuracy drive the CUB and its coevolution with

the tRNA pool.

RESULTS

Preferred codons are more accurately decoded

A direct test of the TAH of CUB requires comparing the mistranslation rate among

synonymous codons. Using mass spectrometry, Mordret *et al.* quantified mistranslations at

individual sites of the *E. coli* proteome (*20*). After removing sites and codons where

mistranslation rates cannot be quantified due to technical reasons (see Materials and Methods),

we grouped mistranslation events according to the identities of their original codons. We then

computed the absolute mistranslation rate of a codon as the ratio of the total intensity of

mistranslated peptides to that of all peptides mapped to the codon. Finally, we computed the

relative mistranslation rate (*RMR*) of a codon by dividing its absolute mistranslation rate by the

mean absolute mistranslation rate of all codons coding for the same amino acid. *RMR* >1 means

that the codon has a higher mistranslation rate than the average among all codons for the same

amino acid, whereas *RMR* <1 means the opposite. Codon usage was assessed by the relative

synonymous codon usage (*RSCU*). The *RSCU* of a codon equals its frequency in the genome

relative to the average frequency of all codons for the same amino acid (*21*). A codon with

RSCU >1 is preferred while a codon with *RSCU* <1 is unprefered.

We were able to estimate the *RMR* for 27 codons of nine amino acids (**Fig. 1a**). Except

for Gly, the most preferred synonymous codon of an amino acid shows *RMR* <1, providing a

significant support for the TAH ($P = 0.020$, one-tailed binomial test). Similarly, except for Gly

and Val, the least prevalent synonymous codon of an amino acid shows *RMR* >1 ($P = 0.090$, one-

tailed binomial test). Because both *RSCU* and *RMR* of a codon are relative to the mean of all

codons for the same amino acid, they can be compared among codons of different amino acids.

Indeed, a strong negative correlation was observed between *RSCU* and *RMR* among the 27

codons (Pearson's $r = -0.56$, $P < 0.001$, permutation test; Spearman's $\rho = -0.49$, $P = 0.005$,

permutation test; **Fig. 1b**). Together, these findings from the proteomic data of *E. coli*

demonstrate that preferred codons tend to have lower mistranslation rates, supporting the TAH of

CUB.

Relative translational accuracies of synonymous codons vary across taxa

How do certain synonymous codons achieve higher translational accuracies than others?

There are two general scenarios. In the first scenario, referred hereinafter as the constant

accuracy hypothesis, the translational accuracy is intrinsically higher for a synonymous codon

than another because of their different chemical nature (*22*). Consequently, the relative

translational accuracies of synonymous codons should be more or less the same in different

species. For instance, given that AAA (Lys) is more accurate than AAG (Lys) in *E. coli* (**Fig.**

1a), we expect the same trend in the vast majority if not all species. Alternatively, relative

translational accuracies of synonymous codons may be greatly influenced by species-specific

factors such as the tRNA pool. Under this scenario, referred to as the variable accuracy

hypothesis hereinafter, the relative accuracies of synonymous codons vary among species. That

is, AAA is more accurate than AAG in many species but the opposite is true in many other

species.

Measuring the relative translational accuracies of synonymous codons in a large number

of species will allow differentiating between the above two hypotheses, which will in turn help

understand the mechanism underlying the translational accuracy differences among synonymous

codons. Because codon-specific, proteome-based translational accuracies have not been

measured beyond *E. coli*, we resort to a sequence-based proxy referred to as the odds ratio (*OR*)

that originated from Akashi's test (*7*). Specifically, the *OR* of synonymous codon X that encodes

amino acid Y in a gene is the number of times that X is used at invariant Y sites relative to the

number of times that X is not used at invariant Y sites, divided by the number of times that X is

used at variant Y sites relative to the number of times that X is not used at variant Y sites (**Fig.**

2a). Here, invariant and variant Y sites refer to Y sites in the focal species whose counterparts in

the ortholog from a related species have Y and non-Y, respectively. The *OR* values computed

from individual genes can be combined to yield a single *OR* using the Mantel-Haenszel

procedure (see Materials and Methods). While *OR* was originally developed for preferred

codons, it can be computed for any codon of the 18 amino acids that have multiple synonymous

codons (*14*). Based on Akashi's test, *OR* has been used as a proxy for the relative translational

accuracy of a codon (*14*). To verify the relationship between *OR* and relative translational

accuracy, we computed *OR* values by aligning *E. coli* genes with their *Salmonella enterica*

orthologs. Indeed, for the 27 codons with *RMR* estimates, *OR* and *RMR* are strongly negatively

correlated ($r = -0.63$, $P = 0.001$; $\rho = -0.43$, $P = 0.01$; **Fig. 2b**), confirming that the *OR* of a codon

is a valid proxy for its relative translational accuracy.

To examine whether the relative translational accuracies of synonymous codons vary

among species, we took advantage of a recently built phylogenetic tree of 10,575 microbial taxa

(*23*). Because most taxa (9,906) in the tree are from the domain Bacteria, we first focused our

analysis on Bacteria. We picked all 1,197 pairs of sister bacterial taxa from the tree and aligned

their orthologous genes (see Materials and Methods). We randomly assigned one taxon in each

pair as the focal taxon and computed *OR* for each codon as described above. We found a

positive correlation between *RSCU* and *OR* across codons in 95% of the taxa examined (**Fig. 2c**),

demonstrating an overwhelming support for the TAH of CUB in Bacteria.

We computed ln(*OR*) to make its distribution relatively symmetric to aid visualization,

and examined as an example ln(*OR*) for codon CAT (His) in each of the focal taxa arranged

according to the bacterial tree (one taxon per order is presented in **Fig. 2d**). We found ln(*OR*) to

vary greatly from negative values to positive values, with a high density near 0 (**Fig. 2e**).

Furthermore, the extreme values of ln(OR) (bright red and bright green in **Fig. 2d**) are scattered

across the tree rather than concentrated in a few clades, suggesting that the relative translational

accuracy of CAT has changed substantially and frequently in evolution. The across-taxon

variation of OR indicates that CAT is the relatively inaccurate one of the two synonymous

codons of His in many taxa (red in **Fig. 2d**) but the relatively accurate one in many other taxa

(green), supporting the variable accuracy hypothesis. From **Fig. 2e**, which shows the 18 amino

acids each with multiple codons, it is clear that the pattern observed for CAT applies to all

codons. Furthermore, every codon has OR >1 in at least 8.9% of the taxa examined (**Fig. S1a**).

These results thus support the variable accuracy hypothesis for all synonymous codons. The

above observations of OR variation among taxa are not primarily caused by sampling error,

because a similar pattern was detected when we analyzed a subset of taxa for each amino acid

where the number of occurrences of each synonymous codon considered in OR estimation is at

least 1000 per taxon (**Fig. S1b**). They are not mainly caused by genetic drift either, because a

similar pattern was found when we analyzed a subset of taxa with strong signals of selection for

translational accuracy (correlation between $RSCU$ and OR exceeding 0.5) (**Fig. S1c**). It is worth

pointing out that, despite the general support for the variable accuracy hypothesis, for a minority

of codons such as ATA (Ile), AGA (Arg), and AGG (Arg), the distribution of ln(OR) is strongly

skewed toward negative values (**Fig. 2e**), suggesting that their relative translational accuracies

are somewhat constrained although not invariable in evolution.

To investigate if the above observations from Bacteria are generalizable to the other two

domains of life, we first expanded our analysis to Archaea represented in the large phylogeny

mentioned (*23*). We found that the correlation between $RSCU$ and OR is positive in 90% of taxa

examined and that ln(OR) varies greatly across taxa for each codon (**Fig. S2**), further supporting

the TAH and the variable accuracy hypothesis. For Eukaryota, we analyzed five commonly used

model organisms: human, mouse, worm, fly, and budding yeast (see Materials and Methods). In

each of these species, the correlation between $RSCU$ and OR is significantly positive (**Table S1**),

supporting the TAH. Except for the two mammals, which are closely related, the ORs estimated

from one species are not well correlated with those estimated from another species (**Fig. S3**).

Furthermore, the correlation in OR generally declines with the divergence time between the two

species (**Fig. S3**), consistent with the variable accuracy hypothesis. Taken together, our results

show that the TAH is generally supported in all domains of life but the relative translational

accuracies of synonymous codons vary across taxa.

Mechanistic basis of among-codon and across-taxon variations of translational accuracies

The empirical support for the variable accuracy hypothesis strongly suggests that the

determinants of the *RMR*s of synonymous codons vary among species. In the aforementioned

Kramer-Farabaugh study (*19*), the authors found that artificially increasing the expression level

of the cognate tRNA for Arg codons AGA and AGG reduces their mistranslations to Lys, so

proposed that the competition between cognate and near-cognate tRNAs determines the

mistranslation rate of a codon. Here, the cognate tRNA is the tRNA whose anticodon pairs with

the codon correctly (allowing wobble pairing), whereas the near-cognate tRNA corresponds to a

different amino acid and has an anticodon that mismatches the codon at one position. Consistent

with the above proposal, Mordret *et al.* (*20*) inferred that most of the mistranslation events in *E.*

coli arose from mispairing between codons and near-cognate tRNAs. They further noted that,

for certain types of mistranslation, there is a negative correlation across growth phases between

the mistranslation rate and the ratio ($R_{c/nc}$) in abundance between cognate and near-cognate

tRNAs, although the correlation was rarely statistically significant (*20*). Based on these past

observations, we hypothesize that the relative translational accuracy of a synonymous codon

increases with its relative $R_{c/nc}$, or $RR_{c/nc}$, which is $R_{c/nc}$ divided by the mean $R_{c/nc}$ of all codons

coding for the same amino acid (see Materials and Methods). We further hypothesize that,

because the tRNA pool varies substantially among species (*24*), the among-species variation of

relative translational accuracies arises from the among-species variation in $RR_{c/nc}$.

To test the above hypotheses, we computed $RR_{c/nc}$ for each codon using published tRNA

expression levels in *E. coli* (*20*). Indeed, we observed a significant negative correlation between

$RR_{c/nc}$ and *RMR* ($r = -0.47$, $P = 0.009$; $\rho = -0.53$, $P = 0.005$; **Fig. 3a**) and a significant positive

correlation between $RR_{c/nc}$ and *OR* ($r = 0.49$, $P = 0.07$; $\rho = 0.74$, $P = 0.00001$; **Fig. 3b**) across

codons, supporting the hypothesis that the relative ratio of cognate to near-cognate tRNA

abundances is a major determinant of a codon's relative translational accuracy in *E. coli*. Note

that the relative cognate tRNA abundance alone is not significantly correlated with *RMR* ($r = -$

0.22, $P = 0.2$; $\rho = -0.04$, $P = 0.4$; **Fig. S4a**), supporting the role of competition between cognate

and near-cognate tRNAs in determining *RMR*. As previously reported (*14*), the relative cognate

tRNA level is highly correlated with $RSCU$ ($r = 0.61$, $P = 0.03$; $\rho = 0.48$, $P = 0.02$; **Fig. S4b**),

which is likely a result of selection for high translational efficiency (i.e., more codons translated

per unit time per cell) because balanced codon usage relative to cognate tRNA concentrations

maximizes translational efficiency (*14*).

We next investigated whether the above finding in *E. coli* applies to other bacterial taxa

surveyed in Fig. 2. Because tRNA expression levels are unknown for the vast majority of these

taxa, we used the gene copy number of each tRNA species as a proxy for the total expression

level of the tRNA species (*25*). Indeed, *E. coli* $RR_{c/nc}$ computed from tRNA gene copy numbers

is highly correlated with that computed from tRNA expression levels ($r = 0.77$, $P = 1.64 \times 10^{-6}$; ρ

$= 0.90$, $P = 1.36 \times 10^{-10}$). Furthermore, *E. coli* $RR_{c/nc}$ computed from tRNA gene copy numbers is

significantly correlated with RMR (**Fig. 3c**), confirming the validity of using this proxy. We

obtained the tRNA gene annotations for 1094 of the 1197 focal bacterial taxa examined in Fig. 2.

However, in many of these taxa, there is little tRNA gene redundancy or variation in cognate

tRNA gene copy number among synonymous codons despite considerable CUB; in these taxa,

the tRNA gene copy number is unlikely a good proxy for tRNA abunadnce (*26*). Because the

tRNA gene copy number is a good proxy for tRNA abundance in *E. coli*, which has 85 tRNA

genes, we decided to filter out taxa with fewer than 81 tRNA genes to strike a balance between

the noise level and number of taxa in our analysis. This filtering left us with 59 taxa, in each of

which we correlated the OR of a codon with its $RR_{c/nc}$ computed from tRNA gene copy numbers.

The vast majority (92%) of the taxa show a positive correlation (**Fig. 3d**), supporting the

generality of our hypothesis on the role of $RR_{c/nc}$ in determining the relative translational

accuracy of a codon in Bacteria.

To investigate whether the above finding is generalizable to other domains of life, we

analyzed tRNA genes in Archaea taxa and Eukaryotic model organisms. Unfortunately, no

Archaea taxa examined have more than 80 tRNA genes. For each of the five eukaryotes (human,

mouse, fly, worm, and yeast), the correlation between OR and $RR_{c/nc}$ computed from tRNA gene

copy numbers is significantly positive for linear or rank correlation (**Table S2**). Together, our

findings strongly support that, in the diverse taxa surveyed, the ratio of cognate tRNA abundance

to near-cognate tRNA abundance is generally a major determinant of the relative translational

accuracy of a codon. Hence, the variation of the tRNA pool among species can explain the

across-species variation of the relative translational accuracies of synonymous codons.

DISCUSSION

Analyzing published proteomic data from *E. coli*, we provided direct, global evidence that preferred synonymous codons are generally decoded more accurately than unpreferred codons. We found that relative translational accuracies of synonymous codons vary substantially among species, supporting the variable accuracy hypothesis. We obtained strong evidence that the ratio of cognate tRNA abundance to near-cognate tRNA abundance is a major determinant of a codon's relative translational accuracy. Hence, the variable accuracies observed are mechanistically explained by the variation of the tRNA pool across species. These findings, together with the previous report on the selection for translational efficiency (*14*), suggest a model in which the tRNA pool and codon usage coevolve to improve both translational efficiency and accuracy (**Fig. 4a**). Specifically, mutation and drift can alter both codon frequencies and tRNA concentrations. The cellular translational efficiency is maximized when (transcriptomic) codon frequencies equal relative cognate tRNA concentrations (*14*), whereas the translational accuracy of a codon is maximized when the ratio of its cognate tRNA concentration to near-cognate tRNA concentration is maximized. Under this model, selections for translational efficiency and translational accuracy are related but not perfectly aligned, which could introduce tradeoffs between translational efficiency and accuracy (*27*). Indeed, our simulation of a simple genetic system with two amino acids, each encoded by two synonymous codons (**Fig. 4b**), found that imposing a selection for translational accuracy can lower translational efficiency (**Fig. 4c**).

Interestingly, our results imply that, even in the absence of selection for translational accuracy, the positive correlation between synonymous codon frequency and cognate tRNA concentration resulting from selection for translational efficiency (*14*) will likely render the cognate tRNA concentration relative to near-cognate tRNA concentration higher for more frequently used synonymous codons. Consequently, the positive correlation between the relative codon frequency and relative translational accuracy may arise in the absence of selection for translational accuracy. In fact, in *E. coli*, for 16 of the 18 amino acids with multiple synonymous codons, the codon with the highest cognate tRNA concentration has the highest $RR_{c/nc}$. Upon shuffling the expression levels among tRNA species, we found that, for over one half of the 18 amino acids, the codon with the highest cognate tRNA concentration has the highest $RR_{c/nc}$. This was true in each of 1000 shufflings. Nevertheless, in only 6 of these 1000 shufflings did all 18

amino acids exhibit the above feature. Thus, a high but non-perfect concordance between
translational efficiency and accuracy is expected. Therefore, strictly speaking, the correlation in
Fig. 1b by itself does not prove selection for translational accuracy. However, this correlation, in
conjunction with the correlation between *RSCU* and *OR*, demonstrates that evolutionarily
conserved sites tend to use preferred synonymous codons, which tend to be relatively accurately
translated, hence proving the role of selection for translational accuracy in causing CUB, or the
TAH. How codon usage and the tRNA pool evolve under the joint forces of selections for
translational efficiency and accuracy in addition to mutation and drift is quite complex. For
instance, because any tRNA is simultaneously a cognate tRNA for one or more codons and a
near-cognate tRNA for some other codons, increasing the translational accuracy of a particular
codon might be at the expense of the translational accuracy of another codon. Indeed, a previous
study showed that artificially increasing the cognate tRNA expression levels for arginine codons
can result in proteotoxic stress (*28*). This subtle tradeoff could cause non-independent uses of
codons of different amino acids. This was indeed observed in the simulation aforementioned
(**Fig. 4d**). Future modeling work with realistic parameters might shed more light on this issue.
In addition to impacting translational efficiency and accuracy, synonymous mutations also affect
mRNA folding (*29*), mRNA stability (*30*), mRNA concentration (*30-32*), pre-mRNA splicing
(*33*), and co-translational protein folding (*34, 35*), so additional selections may shape CUB and
its evolution.
Our study has several caveats. First, in our calculation of a codon's mistranslation rate,
we lumped all mistranslations of the codon regardless of the erroneous amino acid it is translated
to. Because different mistranslations of the same codon likely have differential fitness costs and
because selection for translational accuracy likely minimizes the total fitness reduction caused by
mistranslation instead of the mistranslation rate *per se*, properly weighting different
mistranslations in *RMR* calculation will likely strengthen its correlation with *RSCU*. Second,
when calculating the ratio of cognate tRNA concentration to near-cognate tRNA concentration,
we did not consider the difference in interaction strength between different codons and
anticodons (*36*). Future research that takes into account this interaction under physiological
conditions may significantly improve the signal in the correlation analysis of Fig. 3. Third, our
analysis in Fig. 3d was limited to taxa with >80 tRNA genes. Future research using tRNA
expression levels (*26*) when they become available can confirm if the same pattern holds for taxa

with fewer tRNA genes. Finally, due to data limitation, we did not consider tRNA expression

variations across environments, cell cycle stages, or tissues (*37*). In the future, it would be

interesting to study how such variations simultaneously impact translational efficiency and

accuracy.

Our results might help design organisms with expanded code tables (*38*). Expanding the

code table is realized by introducing unnatural tRNAs that are charged with non-canonical amino

acids. The introduction of these tRNAs often leads to fitness defects due to mistranslation of

normal codons (*39*). Our research suggests that one way to alleviate the proteotoxic stress is to

identify potential near-cognate codons that could be mistranslated by the unnatural tRNA and

adjust the natural tRNA pool to minimize the impact.

MATERIALS AND METHODS

Estimating relative mistranslation rates of synonymous codons from *E. coli* proteomic data

The proteomic data analyzed came from Table S1 in Modret *et al.* (*20*). The authors

separately measured mistranslation events from high-solubility and low-solubility proteins using

mass spectrometry, and both groups of events were considered in our analysis. We focused on

the data from the wild-type strain BW25113 in the MOPS complete medium because (i) this

dataset is the largest among datasets from all strain-medium combinations and (ii) no artificial

perturbation such as mutation, drug, or amino acid depletion was applied (*20*). We first removed

sites that cannot be traced to a unique original codon. We also filtered out sites showing an

intensity of "NaN" for the unmodified (aka base) peptide or mistranslated (aka dependent)

peptide. Because different synonymous codons tend to generate different mistranslations by

mispairing with different near-cognate tRNAs, if these different mistranslations have different

detection probabilities, the comparison between synonymous codons would be unfair.

Unfortunately, some mistranslations produce mass shifts indistinguishable from post-

translational modifications so cannot be reliably identified through mass spectrometry (*26*),

which would produce exactly this situation in some cases. Therefore, we removed amino acids

with undetectable mistranslations except for Leu and Ile. We kept these two amino acids because

the only undetectable mistranslations for them are Leu to Ile and Ile to Leu, both can be

considered benign due to the high physicochemical similarity between Leu and Ile (*40*).

Considering the structure of the genetic code table, we found that the underestimation of the

mistranslation rate due to the negligence of mistranslations between Leu and Ile is severer for

unpreferred than preferred codons, suggesting that the actual strength of evidence for higher

mistranslation rates of unpreferred than preferred synonymous codons is stronger than what is

shown in Fig. 1. We then computed each codon's absolute mistranslation rate by dividing the

total intensity of mistranslated (i.e., dependent) peptides by that of all (i.e., dependent + base)

peptides mapped to the codon. We divided each codon's absolute mistranslation rate by the mean

absolute mistranslation rate of all codons coding for the same amino acid to obtain the codon's

relative mistranslation rate (RMR). We removed amino acids without data for all of its

synonymous codons because calculating RMR requires having data for all synonymous codons of

an amino acid. In total, we computed RMR for 27 codons of 9 amino acids.

Relative synonymous codon usage ($RSCU$), odds ratio (OR), and relative ratio of cognate

tRNA concentration to near-cognate tRNA concentration ($RR_{c/nc}$) for *E. coli*

Peptide and cDNA sequences of *E. coli* (genome assembly ASM584v2) and *S. enterica*

(genome assembly ASM78381v1) were downloaded from Ensembl Bacteria (*41*). We computed

$RSCU$ of codon j of amino acid i from all coding sequences of *E. coli* by $RSCU_{i,j} = \frac{n_i x_{i,j}}{\sum_{j=1}^{n_i} x_{i,j}}$,

where n_i is the number of synonymous codons of amino acid i and $x_{i,j}$ is the number of codon j

of amino acid i in all coding sequences (*21*). Conventionally, $RSCU$ is computed from highly

expressed genes (*21*). However, due to the lack of gene expression information from most of the

species analyzed, we computed $RSCU$ from all genes. This should not qualitatively affect our

analysis, because $RSCU$ computed from highly expressed genes (e.g., the top 20% of genes) is

nearly perfectly correlated with that computed from all genes (e.g., in *E. coli*, $r = 0.96$, $P <$

2.2×10^{-16}).

To calculate the OR of each codon, we first identified one-to-one orthologous proteins

between *E. coli* and *S. enterica* using OrthoFinder (*42*). Next, we aligned these one-to-one

orthologs using MUSCLE (*43*), separating all amino acid sites into conserved and non-conserved

sites. For a focal codon in gene i, we tabulated a_i, number of times the focal codon is observed

at conserved amino acid sites; b_i, number of times the focal codon is observed at unconserved

sites; c_i, total number of times the focal codon's synonymous codons are observed at conserved

sites; and d_i, total number of times the focal codon's synonymous codons are observed at

unconserved sites. Here, the focal codon's synonymous codons do not include itself. *OR* for

gene *i* equals $(a_i d_i)/(b_i c_i)$. Using the Mantel-Haenszel procedure, we combined the odds ratios of

the focal codon from individual genes into one odds ratio (*7*) by $OR = \dfrac{\Sigma_i \frac{a_i d_i}{(a_i+b_i+c_i+d_i)}}{\Sigma_i \frac{b_i c_i}{(a_i+b_i+c_i+d_i)}}$.

To compute $RR_{c/nc}$ of a codon, we tabulated the cognate tRNAs and near-cognate tRNAs

of the codon. Cognate tRNAs are all tRNAs that can pair with the focal codon allowing wobble

pairing at the 3rd codon position, while near-cognate tRNAs are tRNAs coded for a different

amino acid but can pair with the focal codon with one base-pair mismatch (allowing wobble

pairing at the 3rd codon position). We then weighted each tRNA by their average relative

expression levels across three growth stages in the MOPS complete media (GEO number:

GSE128812). Finally, we normalized the ratio for each codon by the average ratio of all codons

coding for the same amino acid.

***RSCU*, *OR*, and *RR*_{c/nc} for other species**

$RSCU$, OR, and $RR_{c/nc}$ for other species

RSCU, *OR*, and $RR_{c/nc}$ were calculated for non-*E. coli* taxa as for *E. coli*, with the

differences noted below. For the non-*E. coli* prokaryotic taxa, we downloaded the phylogenetic

tree of 10,575 taxa from the Web of Life (*23*) (https://biocore.github.io/wol/) and identified sister

taxa from the tree. Briefly, each pair of sister taxa are two taxa that are the single closest relative

to each other in the tree. For each pair of sister taxa, we downloaded from the same web site

their protein-coding DNA sequences, protein sequences, and tRNA gene copy number data. For

eukaryotic model organisms, we downloaded protein-coding DNA sequences and protein

sequences of human (*Homo sapiens*), mouse (*Mus musculus*), fly (*Drosophila melanogaster*),

worm (*Caenorhabditis elegans*), and budding yeast (*Saccharomyces cerevisiae*) from the NCBI

refseq database (*44*). We further downloaded the protein sequences of *Macaca mulatta* (as a

relative of *H. sapiens*), *Rattus norvegicus* (as a relative of *M. musculus*), *Drosophila erecta* (as a

relative of *D. melanogaster*), *Caenorhabditis briggsae* (as a relative of *C. elegans*), and

Saccharomyces paradoxus (as a relative of *S. cerevisiae*) from the NCBI refseq database. The

tRNA gene annotations in the five model organisms were downloaded from GtRNAdb (*24*).

$RR_{c/nc}$ was computed using tRNA gene copy numbers instead of tRNA expression levels.

Statistical analysis

Many of the quantities estimated in our work, such as RMR, $RR_{c/nc}$, $RSCU$, and OR, are

not independent among synonymous codons. To deal with this non-independence in statistical

tests, we applied permutation tests. Specifically, in **Fig. 1b**, we generated 1000 permuted

samples by shuffling the absolute mistranslation rates among all codons and then re-estimated

RMR values. We then computed the correlation between RMR and $RSCU$ in each permuted

sample while holding the $RSCU$ value of each codon unchanged. P equals the fraction of

permuted samples with the correlation coefficient more negative than that observed in the

original sample. Similarly, when testing the correlation between RMR and OR (**Fig. 2b**), we

shuffled the absolute mistranslation rate among all codons and recomputed RMR while holding

the OR for each codon unchanged. When testing the correlation between RMR (or OR) and

$RR_{c/nc}$ (**Fig. 3**), we shuffled the absolute mistranslation rates among codons and the expression

levels (or gene copy numbers) among tRNAs. Finally, when testing the correlation between

RMR (or $RSCU$) and relative cognate tRNA concentration (**Fig. S4**), we shuffled the absolute

mistranslation rate among codons and the expression level among tRNAs.

To estimate the standard error (SE) of the RMR of each codon, we constructed 1000

bootstrap samples by resampling the sites in the original data with replacement. Similarly, we

estimated the SE of the OR of each codon by constructing 1000 bootstrapped *E. coli* genomes via

resampling its genes that have one-to-one orthologs in *S. enterica*.

Simulation

To assess the impact of selections for translational accuracy and efficiency on codon

usage, we built a toy model with two amino acids: aa_0 and aa_1. Amino acid aa_0 is encoded by

synonymous codons 00 and 01 while aa_1 is encoded by synonymous codons 10 and 11 (**Fig. 4b**).

Codon-anticodon pairing follows the rule that 0 pairs with 1 and vice versa. The cognate tRNA

of a codon has an anticodon that pairs perfectly with the codon, while the near-cognate tRNA has

an anticodon that pairs with the codon with exactly one mismatch and carries the other amino

acid.

We considered a unicellular organism with one gene consisting of n codons. We assumed

that the mRNA level of the gene does not change in the evolution simulated and that ribosomes

are in shortage. We defined the organismal fitness as follows.

461            $Absolute\ fitness = Function - Cost$, where

$$Function = TE \times \sum_{i=1}^{n} f_i \text{ and } Cost = \sum_{i=1}^{n} c_i.$$

Here, f_i and c_i are the function and cost of codon i, respectively. We set $f_i = F_i$ if codon i encodes

the pre-specified optimal amino acid at the codon; otherwise, $f_i = 0$. For each i, F_i is a random

variable sampled from an exponential distribution with the mean equal to 1 (*45*). Following a

previous study (*14*), we set the expected codon selection time per amino acid aa₀ as $t_0 =$

$p_1^2/q_1 + p_2^2/q_2$, where p_1 and $p_2 = 1 - p_1$ are the fractions of amino acid aa₀ encoded by codon 00

and 01, respectively, and q_1 and $q_2 = 1 - q_1$ are the fractions of corresponding cognate tRNAs

among all tRNAs of aa₀, respectively. We similarly set the expected codon selection time per

amino acid aa₁ and computed the total codon selection time of all codons. Translational

efficiency *TE*, which is the number of codons translated per unit time, is the inverse of the total

codon selection time. We set $c_i = TE \times C_i$ if codon i does not encode the pre-specified optimal

amino acid at the codon; otherwise, $c_i = C_i \times TE \times \frac{1}{RR_{c/nc}}$. When there is no selection for

translational accuracy, $C_i = 0$; otherwise, C_i for codon i is a random variable sampled from an

exponential distribution with mean equal to 1. Note that C_i and F_i are independent from each

other. $RR_{c/nc}$ is computed as described in Results; the inverse of $RR_{c/nc}$ measures the

mistranslation rate.

We started the simulation with a coding sequence of 200 nucleotides, coding for 100

amino acids. Each site had a 50% chance to be 0 or 1. For simplicity, we assumed that the initial

amino acid sequence is optimal such that the evolution in our simulation is largely about codon

usage. For each of the four different tRNAs (with anticodons of 00, 01, 10, and 11,

respectively), we sampled the initial copy number from 1, 2, and 3 with equal probabilities.

Next, we simulated the coevolution between the tRNA pool and codon usage following a

strong selection, weak mutation regime. We first generate a mutation. With a probability of

0.02, it alters the copy number of a tRNA. In this case, we randomly pick a tRNA species to

change its copy number by +1 or -1 with equal probabilities unless the copy number is 1, in

which case it is +1. With a probability of 0.98, the mutation is a random point mutation at a

randomly picked site of the coding sequence. The fitness of the mutant is then computed

following the above fitness definition. The mutation is fixed with a probability of $\frac{1-r^{-1}}{1-r^{-N}}$, where r

is the ratio of the absolute fitness of the mutant to that of the wild-type and N is the population

size (*46*). The above mutation-selection process was repeated 100,000 rounds in each simulation

to reach an equilibrium. For each *N*, we simulated 200 times with and 200 times without

selection for translational accuracy.

ACKNOWLEDGEMENTS

We thank Dr. Yitzhak Pilpel for consultation on the *E. coli* proteomic data in Mordret *et*

al. (2019) and Wenfeng Qian, Jian-Rong Yang, and members of the Zhang laboratory for

valuable comments. This work was supported by the U.S. National Institutes of Health research

grant R35GM139484 to J.Z.

REFERENCES

1. R. Hershberg, D. A. Petrov, Selection on codon bias. *Annu Rev Genet* **42**, 287-299
(2008).
2. J. B. Plotkin, G. Kudla, Synonymous but not the same: the causes and consequences of
codon bias. *Nat Rev Genet* **12**, 32-42 (2011).
3. T. Ikemura, Correlation between the abundance of Escherichia coli transfer RNAs and the
occurrence of the respective codons in its protein genes: a proposal for a synonymous
codon choice that is optimal for the E. coli translational system. *J Mol Biol* **151**, 389-409
(1981).
4. W. Fiers *et al.*, A-protein gene of bacteriophage MS2. *Nature* **256**, 273-278 (1975).
5. G. M. Air *et al.*, Gene F of bacteriophage phiX174. Correlation of nucleotide sequences
from the DNA and amino acid sequences from the gene product. *J Mol Biol* **107**, 445-458
(1976).
6. A. Efstratiadis, F. C. Kafatos, T. Maniatis, The primary structure of rabbit beta-globin
mRNA as determined from cloned DNA. *Cell* **10**, 571-585 (1977).
7. H. Akashi, Synonymous codon usage in Drosophila melanogaster: natural selection and
translational accuracy. *Genetics* **136**, 927-935 (1994).
8. D. A. Drummond, C. O. Wilke, The evolutionary consequences of erroneous protein
synthesis. *Nat Rev Genet* **10**, 715-724 (2009).
9. B. Chen, M. Retzlaff, T. Roos, J. Frydman, Cellular strategies of protein quality control.
Cold Spring Harb. Perspect. Biol. **3**, a004374 (2011).
10. M. Ibba, D. Söll, Quality control mechanisms during translation. *Science* **286**, 1893-1897
(1999).
11. J. J. Hopfield, Kinetic proofreading: a new mechanism for reducing errors in biosynthetic
processes requiring high specificity. *Proceedings of the National Academy of Sciences* **71**,
4135-4139 (1974).
12. D. E. Weinberg *et al.*, Improved Ribosome-Footprint and mRNA Measurements Provide
Insights into Dynamics and Regulation of Yeast Translation. *Cell Rep* **14**, 1787-1799
(2016).
13. J. A. Hussmann, S. Patchett, A. Johnson, S. Sawyer, W. H. Press, Understanding Biases in
Ribosome Profiling Experiments Reveals Signatures of Translation Dynamics in Yeast.
PLoS Genet **11**, e1005732 (2015).

14. W. Qian, J.-R. Yang, N. M. Pearson, C. Maclean, J. Zhang, Balanced codon usage
optimizes eukaryotic translational efficiency. *PLoS Genet* **8**, e1002603 (2012).
15. D. A. Drummond, C. O. Wilke, Mistranslation-induced protein misfolding as a dominant
constraint on coding-sequence evolution. *Cell* **134**, 341-352 (2008).
16. N. Stoletzki, A. Eyre-Walker, Synonymous codon usage in Escherichia coli: selection for
translational accuracy. *Mol Biol Evol* **24**, 374-381 (2007).
17. P. Shah, M. A. Gilchrist, Effect of correlated tRNA abundances on translation errors and
evolution of codon usage bias. *PLoS Genet* **6**, e1001128 (2010).
18. J. Precup, J. Parker, Missense misreading of asparagine codons as a function of codon
identity and context. *J Biol Chem* **262**, 11351-11355 (1987).
19. E. B. Kramer, P. J. Farabaugh, The frequency of translational misreading errors in E. coli
is largely determined by tRNA competition. *RNA* **13**, 87-96 (2007).
20. E. Mordret *et al.*, Systematic detection of amino acid substitutions in proteomes reveals
mechanistic basis of ribosome errors and selection for translation fidelity. *Mol. Cell* **75**,
427-441. e425 (2019).
21. P. M. Sharp, T. M. Tuohy, K. R. Mosurski, Codon usage in yeast: cluster analysis clearly
differentiates highly and lowly expressed genes. *Nucleic Acids Res.* **14**, 5125-5143
(1986).
22. R. Hershberg, D. A. Petrov, General rules for optimal codon choice. *PLoS Genet* **5**,
e1000556 (2009).
23. Q. Zhu *et al.*, Phylogenomics of 10,575 genomes reveals evolutionary proximity between
domains Bacteria and Archaea. *Nature communications* **10**, 1-14 (2019).
24. P. P. Chan, T. M. Lowe, GtRNAdb: a database of transfer RNA genes detected in genomic
sequence. *Nucleic Acids Res.* **37**, D93-D97 (2009).
25. T. Tuller *et al.*, An evolutionarily conserved mechanism for controlling the efficiency of
protein translation. *Cell* **141**, 344-354 (2010).
26. Y. Wei, J. R. Silke, X. Xia, An improved estimation of tRNA expression to better
elucidate the coevolution between tRNA abundance and codon usage in bacteria. *Sci.*
Rep. **9**, 1-11 (2019).
27. J. R. Yang, X. Chen, J. Zhang, Codon-by-codon modulation of translational speed and
accuracy via mRNA folding. *PLoS Biol* **12**, e1001910 (2014).
28. A. H. Yona *et al.*, tRNA genes rapidly change in evolution to meet novel translational
demands. *Elife* **2**, e01339 (2013).
29. C. Park, X. Chen, J. R. Yang, J. Zhang, Differential requirements for mRNA folding
partially explain why highly expressed proteins evolve slowly. *Proc Natl Acad Sci U S A*
110, E678-686 (2013).
30. V. Presnyak *et al.*, Codon optimality is a major determinant of mRNA stability. *Cell* **160**,
1111-1124 (2015).
31. S. Chen *et al.*, Codon-Resolution Analysis Reveals a Direct and Context-Dependent
Impact of Individual Synonymous Mutations on mRNA Level. *Mol Biol Evol* **34**, 2944-
2958 (2017).
32. Z. Zhou *et al.*, Codon usage is an important determinant of gene expression levels largely
through its effects on transcription. *Proc Natl Acad Sci U S A* **113**, E6117-E6125 (2016).
33. J. V. Chamary, J. L. Parmley, L. D. Hurst, Hearing silence: non-neutral evolution at
synonymous sites in mammals. *Nat Rev Genet* **7**, 98-108 (2006).
34. F. Buhr *et al.*, Synonymous Codons Direct Cotranslational Folding toward Different

Protein Conformations. *Mol Cell* **61**, 341-351 (2016).

35.   I. M. Walsh, M. A. Bowman, I. F. Soto Santarriaga, A. Rodriguez, P. L. Clark,
Synonymous codon substitutions perturb cotranslational protein folding in vivo and
impair cell fitness. *Proc Natl Acad Sci U S A* **117**, 3528-3534 (2020).

36.   M. d. Reis, R. Savva, L. Wernisch, Solving the riddle of codon usage preferences: a test
for translational selection. *Nucleic Acids Res.* **32**, 5036-5044 (2004).

37.   H. Gingold *et al.*, A dual program for translation regulation in cellular proliferation and
differentiation. *Cell* **158**, 1281-1292 (2014).

38.   E. Ros, A. G. Torres, L. R. de Pouplana, Learning from Nature to Expand the Genetic
Code. *Trends Biotechnol.*, (2020).

39.   J. W. Chin, Expanding and reprogramming the genetic code. *Nature* **550**, 53-60 (2017).

40.   S. Henikoff, J. G. Henikoff, Amino acid substitution matrices from protein blocks. *Proc*
Natl Acad Sci U S A **89**, 10915-10919 (1992).

41.   K. L. Howe *et al.*, Ensembl 2021. *Nucleic Acids Res.* **49**, D884-D891 (2021).

42.   D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 1-14 (2019).

43.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792-1797 (2004).

44.   N. A. O'Leary *et al.*, Reference sequence (RefSeq) database at NCBI: current status,
taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733-D745
(2016).

45.   A. Eyre-Walker, P. D. Keightley, The distribution of fitness effects of new mutations. *Nat.*
Rev. Genet **8**, 610-618 (2007).

46.   P. A. P. Moran, in *Mathematical proceedings of the cambridge philosophical society*.
(Cambridge University Press, 1958), vol. 54, pp. 60-71.

608
609
Figure 1. More frequently used synonymous codons tend to be decoded more accurately in
E. coli. **a**, Comparison of relative synonymous codon usage (*RSCU*, bars) and relative
mistranslation rate (*RMR*, dots) among synonymous codons for nine amino acids with proteome-
based *RMR* estimates. **b**, A significant negative correlation between *RSCU* and *RMR* across the
27 codons in panel a. The red line is the linear regression. In both panels, error bars represent one
standard error estimated by the bootstrap method. The standard error of *RSCU* estimated by the
bootstrap method is negligible due to the large number of each codon in the genome, so is not
shown. *P*-values are based on permutation tests.
618
619

Figure 2. Variation of relative translational accuracies of synonymous codons across taxa.
a, Diagram explaining the calculation of odds ratio (*OR*) of the codon CAT that serves as a proxy for its relative translational accuracy. Showing here is a hypothetical alignment of orthologous proteins (and the underlying coding sequences) between the focal species and a related species. **b**, *OR* is negatively correlated with *RMR* across codons in *E. coli*. *P*-values are based on permutation tests. The red line is the linear regression. **c**, Frequency distribution of Pearson's correlation between *RSCU* and *OR* in 1197 bacterial taxa. Ninety-five percent of these taxa show positive correlations. **d**, ln(*OR*) of codon CAT for each of 118 bacterial taxa, one per order, arranged according to their phylogeny shown in the middle. **e**, Violin plots showing frequency distributions of ln(*OR*) of individual codons of the 18 amino acids that have multiple synonymous codons across 1197 bacterial taxa.

Figure 3. The relative ratio of cognate tRNA concentration to near-cognate tRNA concentration ($RR_{c/nc}$) is a major determinant of a codon's relative translational accuracy. **a**, *RMR* is negatively correlated with $RR_{c/nc}$ across codons in *E. coli*. **b**, *OR* is positively correlated with $RR_{c/nc}$ across codons in *E. coli*. **c**, *RMR* is negatively correlated with $RR_{c/nc}$ computed using tRNA gene copy numbers instead of tRNA concentrations in *E. coli*. **d**, Frequency distribution of Pearson's correlation between *OR* and $RR_{c/nc}$ computed using tRNA gene copy numbers in bacterial taxa with >80 tRNA genes. All *P*-values are based on permutation tests. In a-c, the red line is the linear regression.

644
Figure 4. Selections for translational efficiency and accuracy shape the tRNA pool and
codon usage. a, A model for the coevolution of the tRNA pool and codon usage driven by
selections for translational efficiency and accuracy. **b.** A toy model with two amino acids, each
encoded by two synonymous codons. A dotted line connects a codon with its near-cognate
tRNA. **c,** Translational efficiency is significantly lower in the presence of selection for
translational accuracy than in the absence of this selection. **d,** The absolute difference between
the $RSCU$ of 00 ($RSCU_{00}$) and that of 11 ($RSCU_{11}$) is smaller under the selection for translational
accuracy than without this selection. With the selection, codon usage for aa_0 and that for aa_1
become coupled, because selection disfavors the cognate tRNA of the common codon of aa_0 to
become the near-cognate tRNA of the common codon of aa_1, and vice versa. In c and d, each box
plot shows the distribution from 200 replicates. The lower and higher edges of a box represent
the first (qu_1) and third (qu_3) quartiles, respectively; the horizontal line inside the box indicates
the median (md); the whiskers extend to the most extreme values inside inner fences, md $\pm$
1.5(qu_3-qu_1); and the dots are outliers. *, $0.01 \leq P < 0.05$, Wilcoxon rank-sum tests; **, $0.001 \leq P$
< 0.01; ***, $P < 0.001$.

Supplementary materials of

"Preferred synonymous codons are translated more accurately: Proteomic evidence, among-species variation, and mechanistic basis" by Sun & Zhang

The supplementary materials include:

Table S1. Pearson (*r*) and Spearman (*ρ*) correlations Between *OR* and *RSCU* in eukaryotic model organisms.

Species	*r*	*P*	*ρ*	*P*
Fly	0.84	<0.001	0.83	<0.001
Human	0.80	<0.001	0.74	<0.001
Mouse	0.74	<0.001	0.68	<0.001
Worm	0.42	<0.001	0.39	0.002
Yeast	0.35	0.01	0.39	0.003

Table S2. Pearson (*r*) and Spearman (*ρ*) correlations between *OR* and $RR_{c/nc}$ in eukaryotic model organisms.

Species	r	P	ρ	P
Fly	0.21	0.109	0.25	0.046
Human	0.39	0.007	0.29	0.055
Mouse	0.29	0.081	0.41	0.018
Worm	0.54	<0.001	0.57	<0.001
Yeast	0.42	0.045	0.37	0.053

Fig. S1. Observed patterns of variation of relative translational accuracies of synonymous codons across bacterial taxa are robust. a, Fraction of 1197 taxa with $OR > 1$ for each codon. **b**, Violin plots showing frequency distributions of ln(OR) of individual codons across a subset of taxa in which the number of occurrences of each synonymous codon considered in OR estimation is at least 1000. **c**, Violin plots showing frequency distributions of ln(OR) of individual codons across a subset of taxa with strong signals of selection for translational accuracy (i.e., Pearson's correlation between $RSCU$ and OR exceeds 0.5).

Fig. S2. Variation of relative translational accuracies of synonymous codons among Archaea taxa. a, Frequency distribution of Pearson's correlation between *RSCU* and *OR* in 63 taxa. Ninety percent of taxa show positive correlations. **b**, ln(*OR*) of codon CAT for each of the taxa arranged according to their phylogeny shown in the middle. **c**, Violin plots showing frequency distributions of ln(*OR*) of individual codons across taxa. ln(*OR*) appears less variable here than in Bacteria because of the much fewer Archaea taxa examined.

Fig. S3. Correlation in odds ratio (*OR*) across codons between eukaryotic model organisms. Each dot is a codon.

Fig. S4. Relationship between the cognate tRNA concentration and *RMR* or *RSCU* in *E. coli*. **a**, *RMR* of a codon is not significantly correlated with its relative cognate tRNA concentration, which is its cognate tRNA concentration divided by the mean cognate tRNA concentration of all codons coding for the same amino acid. **b**, *RSCU* of a codon is significantly positively correlated with its relative cognate tRNA concentration. *P*-values are based on permutation tests. Only the 27 codons with *RMR* estimates are analyzed in each panel to allow a direct comparison.