Trends in *Arabidopsis* Research post genome sequencing- A Scientometric study

Sandeep Kumar[1,2$], Amar Kant Kushwaha[1,3], Thribhuvan R[1,4], Balakrishnan M[1], Krishnan P[1$]

[1]ICAR-National Academy of Agricultural Research and Management, Hyderabad, India

[2]Quality and Productivity Improvement Division, ICAR-Indian Institute of Natural Resins and Gums,

Ranchi, India

[3]Division of Crop Improvement and Biotechnology, ICAR-Central Institute for Subtropical

Horticulture, Rehmankhera, P.O. Kakori, (Lucknow), India, 226101

[4]Division of Crop Improvement, ICAR-Central Research Institute for Jute and Allied fibres,

Barrackpore, India

$ Corresponding authors

Sandeep Kumar & Krishnan P

Mail: Sandeep.kumar8@icar.gov.in; krishnanars@naarm.org.in

ICAR-National Academy of Agricultural Research and Management,

Hyderabad, Telangana, India-500030

1. Abstract

Arabidopsis thaliana, a model plant, is intensively researched because of the intrinsic advantages associated with its life cycle, genetics, and other characteristics. In the present study, we analysed the publication data of research work done on *A. thaliana* during 2001-2020, a period when the whole genome sequence of the plant was available to the researchers. The current meta-analysis showed that 31965 research papers were published globally on *Arabidopsis* during the study period. The USA topped the countries with the maximum share of publications (28.97%), followed by China, Japan, and West European countries. The analysis showed a temporal shift in the number of publications from different countries and their research focus. After 2013, the research output from China was higher than that from the USA, and there was a shift in focus from developmental biology to stress-related topics. Plant Physiology journal carried the most research work (2490 publications) on Arabidopsis, which was followed by Plant Cell (2376), the Plant Journal (2260) and the Journal of Experimental Botany (1121). However, there was a progressive decline in the publication in these journals, and a shift was evident in favour of open access journals like Frontiers in Plant Science. The publication and citation numbers also show ongoing Arabidopsis research's relevance to plant sciences, particularly curiosity-driven and discovery-based science. This article delves into the patterns in the prominence of research areas, ideas and foci for the future Arabidopsis research roadmap.

2. Introduction

Arabidopsis thaliana is a plant native to the Eurasian and African region, where it is generally considered a weed that often grows along the roadsides. Researchers across the globe have used this plant to study basic plant biology. It was first described by a German physician named Johannes Thal, on whose name Carl Linnaeus named it *Arabis thaliana*. The name *Arabidopsis thaliana* was given by Gustav Heynhold, who created a new genus by the name of *Arabidopsis* (Meyerowitz, 2001). It is self-pollinated, annual and grows up to 20-25cm. The key qualities, including small plant size, a relatively shorter life cycle period, seed setting in large quantities and smaller genome size with fewer chromosome number, qualifies it as the best candidate as a model plant and amenable for genetic and physiological studies (Rédei, 1970; Somerville and Koornneef, 2002; Koornneef and Meinke, 2010). Primary emphasis was given to the screening and identification of auxotrophic mutants (Rédei, 1975), as a result of which the first Arabidopsis gene sequences came out in 1986 (Chang and Meyerowitz, 1986).

After that, T-DNA-mediated transformation and vacuum infiltration made it more amenable to studies (Koncz et al., 1989). Eventually, Arabidopsis became a plant amenable to generating gene mutations and implementing DNA transformation protocols. The culmination of these projects was the National Science Foundation (NSF) project on Arabidopsis gene identification and whole-genome sequencing (Arabidopsis Genome Initiative, 2000). This project was one of many initiatives of genome sequencing. Various technological advancements happened after the completion of Arabidopsis genome sequencing, which shaped the biological research in many organisms. Genome sequencing brought a tremendous pace and shift in Arabidopsi*s* research (Bevan and Walsh, 2005). Multinational Arabidopsis Steering Committee (MASC) has been actively involved in Arabidopsis research and constantly prepares the roadmaps for Arabidopsis research (http://arabidopsisresearch.org/index.php/en/about). The research interest in *Arabidopsis* as a model plant in the emerging research scenarios has been dwindling following the emergence of high throughput sequencing and other genomics tools (Fig. 1). In addition, there has been an increased application of these newer tools to non-model plants. The current research was designed to analyse

65 the trends in research on the *Arabidopsis thaliana*, in the post-genome sequencing period, with the

66 specific objectives of

67   (i)   Mapping the variation in the theme areas of Arabidopsis research and

68   (ii)   Profile the trends in research networks at various levels, *viz.,* authors, organisations and

69   countries.

3. Materials and methods

3.1 Data collection and analysis

This study uses the bibliometric methodology to quantify scientific collaborations in Arabidopsis

research worldwide. The Web of Science (WoS) database was used for the literature search on

Arabidopsis. Web of Science is a website published by Clarivate Analytics that offers subscription-

based access to numerous databases that provide detailed citation data for many academic subjects

(Li et al., 2018). The search string and filters used to select the publications were as follows (T.I. =

(Arabidopsis)) AND DOCUMENT TYPES: (Article) Timespan: 2001-2020. Indexes: SCI-

EXPANDED, CPCI-S, ESCI. The year 2001 was chosen as the starting point for the survey since

Arabidopsis genome sequencing had been finished by 2000 (Arabidopsis Genome Initiative, 2000).

Conference proceedings and books/book chapters were excluded from the study. The search phrase

was chosen based on our investigation of existing literature and discussions among authors and

subject matter experts. The *thaliana* was not added to the string as our preliminary scanning showed

that many publications did not use the full binomial name in the titles and used only '*Arabidopsis*'.

The data retrieved from the WoS database was exported in text format, with each record containing

the bibliometric data: title name, journal names, author names with Web of Science affiliation,

number of citations, document kinds, year of publication for each article, authors, Web of Science

keywords, and abstracts. The WoS database was also utilised for publication trends over the years, for

which five year moving average of publications was used. The Scientometric/bibliometric parameters

like the total number of publications and citations, average citations per article (ACPP), h-index,

research areas, collaborating countries and funding agencies were extracted from the WoS platform

for 20 years (2000-2020).

3.2 Network analysis:

The co-authorship and keyword occurrence maps were generated using the VOS viewer version

1.6.16 using metadata generated from the web of science (van Eck and Waltman, 2011)

(http://www.vosviewer.com/). For the network analysis, the top 100 results were selected and used in

case of both institutional collaboration and year wise keyword occurrence. Bibliometrix, an R-based

tool, was used for data filtering, research theme evolution and identifying journals based on

Bradford's law, wherein the metadata was fed in BibTeX files (http://www.bibliometrix.org)

(Aria and Cuccurullo, 2017).

4. Results & Discussion

4.1 Global output of research on *Arabidopsis*

The publication of the *Arabidopsis* genome led to a significant rise in *Arabidopsis* research as it

expanded the area of research from only isozyme and protein study to functional genomics

(Mysore et al., 2001; Nath Radhamony et al., 2005; Bevan and Walsh, 2005). The study showed that

31965 research articles were published on the subject during the 20 years from 2001 to 2020 (Fig. 2).

One intriguing element of the total number of articles was a five-year consecutive decline in the total

number of publications following 2013, except for 2019/2020. This decreasing trend in recent years

can be attributed to increased emphasis on the sequencing of commercially significant species such as

rice,   maise,   pigeon   pea,   rapeseed,   tomato   and   many   others

(International Rice Genome Sequencing Project, 2005; Schnable et al., 2009; Varshney et al., 2011;

Wang et al., 2011). This paradigm shift was made possible due to the remarkable progress of

sequencing techniques appropriate for diverse purposes, from high cost and poor throughput to low

cost and high throughput (Türktaş et al., 2015).

In the keyword co-occurrence, we found a clear shift from developmental pathways related to the

peripheral pathways like stress responses and the emergence of some new keywords like

transcriptome and proteomics in the last decade (Fig. 3). It also demonstrates the widespread usage of

various next-generation genomic tools and techniques in the previous decade (2011-2020). The

Arabidopsis gene functions have also been elucidated in the major signalling pathways

(Alonso and Stepanova, 2004; Atkinson et al., 2013). Bolker (2012) emphasised the need to go

beyond model organisms to get a complete picture of various biochemical and signalling pathways in

taxonomic groups. Moreover, the elucidation of biological function through *Arabidopsis* has trickled

into the agriculturally important crop sp. for their validation and utilisation

(Priyanka et al., 2010; Manmathan et al., 2013; Mishra et al., 2014).

4.2 Highly cited studies on Arabidopsis

After whole-genome sequencing of Arabidopsis, the research papers on insertional mutagenesis in

Arabidopsis thaliana received more citations, thus indicating the research trends (Table 1). Before

sequencing, it was tough to create a loss of function mutation. Till the sequencing of the first plant,

only unicellular organisms were subject to the creation of loss of mutation for understanding the

function of the gene(s) (Giaever et al., 2002) and gene silencing and RNAi were utilised for

understanding the developmental role of genes in *C. elegans* with their limitations of unstable

heritability, residual expression of genes and low throughput (Ashrafi et al., 2003). After the

publication of the paper titled "Genome-wide Insertional mutagenesis of *Arabidopsis thaliana"* by

Alonso et al (2003) depicting the technique overcoming the limitation of earlier methods, work started

in Arabidopsis mutant generation and their use in research. The article influenced a large stream of

research, as indicated by the maximum citations (3790) that any Arabidopsis paper received in the last

two decades and laid the foundations of functional biology in Arabidopsis.

Similarly, with the advancement in sequencing techniques, epigenomes were targeted to decipher the

developmental pathway. To detect such epigenetic modification, Cokus et al (2008) came up with a

new approach, BS-seq (Bisulphite sequencing), combining sequence information and bisulphate

treatment for mapping the methylation sites in the whole genome. Similarly, in the same area,

Lister et al (2008) mapped the methylation sites by sequencing smRNA, and methylcytosine to

decipher the role in the developmental pathway. The significance of these two articles is indicated by

the citations received by them, viz., 1516 and 1363, respectively. Some other papers featured on the

top 10 list by the number of citations were related to transcript normalisation, transient gene

expression and microarray database. In short, the protocol and database development paper featured

prominently on the most cited list. It is in line with the evolution of *Arabidopsis* as a model plant,

evidenced by the fact that novel tools and techniques to decipher plant biological functions has been

developed in Arabidopsis and later extended to other crop plants (Koornneef and Meinke, 2010).

4.3 Top countries working on the Arabidopsis research

Based on the number of publications, America contributed 28.97% of the total publications, followed

by China, Germany, and Japan with 21.23%, 14.11% and 12.06% of the publications, respectively.

Table 2 shows the top 10 countries and their per cent contribution, along with other publication

metrics *viz.,* h-index, T.C. (Total Citations) and ACPP (Average Citation Per Publication). The h-

index, which indirectly represents the research influence, was higher for publications from the USA

(133), which was followed by Germany (223), England (204) and Japan (200). Though China has the

second-highest per cent share in total research papers, its h-index is relatively lower, i.e., 166 and the

lowest ACPP score of 29.96 compared to other top 10 countries. The country collaboration map

shown in Fig. 5 provides a clear picture of the collaborations. The United States of America and the

countries in Western Europe like the UK, France and Germany showed a very high collaboration and

scientific output. However, some African and Baltic countries were missing from the Arabidopsis

research map entirely. The countries leading the research in terms of the number of publications were

part of the genome sequencing consortium or those collaborating with them

(https://www.nsf.gov/bio/pubs/reports/arab_prog78.htm). Though China was not the part as they were

not much into *Arabidopsis* research before 1995 (Chen et al., 2006), its research output has been high

in recent years (Fig 4). A plausible explanation of this is the acquisition of natural and mutant stocks

of Arabidopsis seeds post-genome sequencing (Provart et al., 2016).

Based on subject areas, most publications featured the plant sciences and developmental biology

across the countries (Table 3). Cell biology and developmental biology were laid less stress in China,

possibly due to more research being done later. However, the research areas of publication journals

are indicative, and there might exist many cross-cutting and multidisciplinary studies which do not

follow this classification.

4.4 Leading institutions in the Arabidopsis research

Collaborations between different institutes revealed the close links between American and European

institutes (Fig. 5). The top three institutions with the most publications were the Chinese Academy of

Science, the University of California, and the University of Tokyo. However, the Asian institutes had

relatively weak partnerships with other countries. The North American and European institutes were

active in the early research periods, as visible from the corresponding colour. The Chinese Academy

of Science and other Chinese Universities have published more papers than those from Japan and

Western countries.

The highest papers feature the Chinese Academy of Sciences (Fig. 6). However, the articles from

western Europe had relatively high-quality parameters, e.g. Chinese Academy of science has 1856

total publications, but the average citation per paper is lower (41.69) than the U.K. Research

Innovation (UKRI). UKRI has only 779 total papers but 90.03 average citations per paper.

4.5 Top Journals for Arabidopsis Research

Plant Physiology (2376), Plant cell (2490) and the Plant Journal (2260) were the top three journals in

terms of the number of relevant publications, h-index, and total citations (Table 4). Bradford law is an

essential consideration that decides the importance of different journals based on diminishing citation

of relevant articles over the year, following Pareto distribution (Nash-Stewart et al., 2012). Here it can

be easily seen that Plant Physiology, Plant Cell, Plant Journal, Journal of Experimental Botany, Plant

and Cell Physiology, Frontiers in Plant Science and PNAS were the core sources that carried most

publications on Arabidopsis and thus represent the choice journals for the researchers (Fig. 6). The

annual pattern of publication on Arabidopsis in these journals showed a shift in the preferred journals

over the years (Fig. 7), i.e. a decline in the publication in the Plant cell, plant physiology and the plant

journal; however, open publications models of frontiers in plant sciences have gained in popularity.

As a result, none of the top three journals based on the total number of publications was present

among the top three for the recent year, i.e. 2020. This trend is probably due to the citation advantage

of an article in open access (O.A.) due to an increase in the number of readers citing open access

articles (Cullen and Chawner, 2011). The radical initiative to O.A. rather than a subscription with

'Plans S' (science, speed, solution, shock) by 11 funding agencies in Europe indicates such implication

(Else, 2018).

5. Conclusion

We set out to determine the trajectory of Arabidopsis research in various areas. With the sequencing

of the Arabidopsis genome, there was a rise in interest in the research, but it decreased after the

advent and popularisation of new sequencing tools. Publication in Arabidopsis increased steadily in

the first two quarters after the genome sequencing, reducing in the third quarter. Primary research

interests spanned from mutant identification to protein investigation, as seen by co-word analysis,

whereas gene expression experiments were constant throughout 20 years. Some of the highly cited

articles were related to developing and validating novel techniques in plant science research.

Furthermore, Asian countries that have shown a strong interest in Arabidopsis research must

collaborate closely with their western counterparts to improve the quality and reach. Quality is a

significant consideration, as evidenced by the low number of citations, particularly from China, which

has shown a rising trend in the total number of publications. The funds worldwide are depicted more

toward the number of publications than ACPP. The target journal should be criteria based on impact

factors for grants by the funding agency, depicting quality (IF) rather than quantity (T.P.). This can

result in a good ACPP score, although ACCP is affected by several other factors also, viz., journal

rank, primary/reporting author's h-index, SCImago journal rank (SJR) and source normalised impact

per paper (SNIP) (Yaminfirooz and Ardali, 2018). This can be represented from the author's

collaboration point of view as it also has a significant correlation with citation, and it has been

witnessed that there is a lower collaboration among Asian countries compared to among European

countries and USA. A substantial part of Arabidopsis papers has gone to three leading journals: Plant

Physiology, Plant Cell, and Plant Journal. The growing trend toward open access publication is

unmistakably the way forward for journals seeking to remain relevant.

6. Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

7. References

Alonso, J. M., Stepanova, A. N., Leisse, T. J., Kim, C. J., Chen, H., Shinn, P., Stevenson, D. K.,

Zimmerman, J., Barajas, P., Cheuk, R., et al. (2003). Genome-wide insertional mutagenesis of

Arabidopsis thaliana. *Science* 301, 653–657. doi:10.1126/science.1086391.

Alonso, J. M., and Stepanova, A. N. (2004). The ethylene signaling pathway. *Science* 306, 1513–

1515. doi:10.1126/science.1104812.

Arabidopsis Genome Initiative (2000). Analysis of the genome sequence of the flowering plant

Arabidopsis thaliana. *Nature* 408, 796–815. doi:10.1038/35048692.

Aria, M., and Cuccurullo, C. (2017). bibliometrix□: An R-tool for comprehensive science mapping

analysis. *Journal of Informetrics* 11, 959–975. doi:10.1016/j.joi.2017.08.007.

Ashrafi, K., Chang, F. Y., Watts, J. L., Fraser, A. G., Kamath, R. S., Ahringer, J., and Ruvkun, G.

(2003). Genome-wide RNAi analysis of Caenorhabditis elegans fat regulatory genes. *Nature* 421,

268–272. doi:10.1038/nature01279.

Atkinson, N. J., Lilley, C. J., and Urwin, P. E. (2013). Identification of genes involved in the response

of Arabidopsis to simultaneous biotic and abiotic stresses. *Plant Physiol.* 162, 2028–2041.

doi:10.1104/pp.113.222372.

Bevan, M., and Walsh, S. (2005). The Arabidopsis genome: a foundation for plant research. *Genome*

Res. 15, 1632–1642. doi:10.1101/gr.3723405.

Bolker, J. (2012). Model organisms: There's more to life than rats and flies. *Nature* 491, 31–33.

doi:10.1038/491031a.

Chang, C., and Meyerowitz, E. M. (1986). Molecular cloning and DNA sequence of the Arabidopsis

thaliana alcohol dehydrogenase gene. *Proc Natl Acad Sci USA* 83, 1408–1412.

doi:10.1073/pnas.83.5.1408.

Chen, H., Karplus, V. J., Ma, H., and Deng, X. W. (2006). Plant biology research comes of age in

China. *Plant Cell* 18, 2855–2864. doi:10.1105/tpc.106.045393.

Cokus, S. J., Feng, S., Zhang, X., Chen, Z., Merriman, B., Haudenschild, C. D., Pradhan, S., Nelson,

S. F., Pellegrini, M., and Jacobsen, S. E. (2008). Shotgun bisulphite sequencing of the Arabidopsis

genome reveals DNA methylation patterning. *Nature* 452, 215–219. doi:10.1038/nature06745.

Cullen, R., and Chawner, B. (2011). Institutional Repositories, Open Access, and Scholarly

Communication: A Study of Conflicting Paradigms. *The Journal of Academic Librarianship* 37,

460–470. doi:10.1016/j.acalib.2011.07.002.

Else, H. (2018). Radical open-access plan could spell end to journal subscriptions. *Nature* 561, 17–18.

doi:10.1038/d41586-018-06178-7.

Giaever, G., Chu, A. M., Ni, L., Connelly, C., Riles, L., Véronneau, S., Dow, S., Lucau-Danila, A.,

Anderson, K., André, B., et al. (2002). Functional profiling of the Saccharomyces cerevisiae

genome. *Nature* 418, 387–391. doi:10.1038/nature00935.

International Rice Genome Sequencing Project (2005). The map-based sequence of the rice genome.

Nature 436, 793–800. doi:10.1038/nature03895.

Koncz, C., Martini, N., Mayerhofer, R., Koncz-Kalman, Z., Körber, H., Redei, G. P., and Schell, J.

(1989). High-frequency T-DNA-mediated gene tagging in plants. *Proc Natl Acad Sci USA* 86,

8467–8471. doi:10.1073/pnas.86.21.8467.

Koornneef, M., and Meinke, D. (2010). The development of Arabidopsis as a model plant. *Plant J.*

61, 909–921. doi:10.1111/j.1365-313X.2009.04086.x.

Li, K., Rollins, J., and Yan, E. (2018). Web of Science use in published research and review papers

1997-2017: a selective, dynamic, cross-domain, content-based analysis. *Scientometrics* 115, 1–20.

doi:10.1007/s11192-017-2622-5.

Lister, R., O'Malley, R. C., Tonti-Filippini, J., Gregory, B. D., Berry, C. C., Millar, A. H., and Ecker,

J. R. (2008). Highly integrated single-base resolution maps of the epigenome in Arabidopsis. *Cell*

133, 523–536. doi:10.1016/j.cell.2008.03.029.

Manmathan, H., Shaner, D., Snelling, J., Tisserat, N., and Lapitan, N. (2013). Virus-induced gene

silencing of Arabidopsis thaliana gene homologues in wheat identifies genes conferring improved

drought tolerance. *J. Exp. Bot.* 64, 1381–1392. doi:10.1093/jxb/ert003.

Meyerowitz, E. M. (2001). Prehistory and history of Arabidopsis research. *Plant Physiol.* 125, 15–19.

doi:10.1104/pp.125.1.15.

Mishra, S., Shukla, A., Upadhyay, S., Sanchita, Sharma, P., Singh, S., Phukan, U. J., Meena, A.,

Khan, F., Tripathi, V., et al. (2014). Identification, occurrence, and validation of DRE and ABRE

Cis-regulatory motifs in the promoter regions of genes of Arabidopsis thaliana. *J. Integr. Plant*

Biol. 56, 388–399. doi:10.1111/jipb.12149.

Mysore, K. S., Tuori, R. P., and Martin, G. B. (2001). Arabidopsis genome sequence as a tool for

functional genomics in tomato. *Genome Biol.* 2, REVIEWS1003. doi:10.1186/gb-2001-2-1-

reviews1003.

Nash-Stewart, C. E., Kruesi, L. M., and Del Mar, C. B. (2012). Does Bradford's Law of Scattering

predict the size of the literature in Cochrane Reviews? *Journal of the Medical Library Association:*

JMLA 100, 135.

Nath Radhamony, R., Mohan Prasad, A., and Srinivasan, R. (2005). T-DNA insertional mutagenesis

in Arabidopsis: a tool for functional genomics. *Electron. J. Biotechnol.* 8. doi:10.2225/vol8-issue1-

fulltext-4.

Priyanka, B., Sekhar, K., Sunita, T., Reddy, V. D., and Rao, K. V. (2010). Characterization of

expressed sequence tags (ESTs) of pigeonpea (Cajanus cajan L.) and functional validation of

selected genes for abiotic stress tolerance in Arabidopsis thaliana. *Molecular Genetics and*

Genomics 283, 273–287.

Provart, N. J., Alonso, J., Assmann, S. M., Bergmann, D., Brady, S. M., Brkljacic, J., Browse, J.,

Chapple, C., Colot, V., Cutler, S., et al. (2016). 50 years of Arabidopsis research: highlights and

future directions. *New Phytol.* 209, 921–944. doi:10.1111/nph.13687.

Rédei, G. P. (1970). Arabidopsis thaliana (L.) Heynh. A review of the genetics and biology.

Bibliographia Genetica. 20.

Rédei, G. P. (1975). "Induction of auxotrophic mutations in plants," in *Genetic manipulations with*

plant material (Springer), 329–350.

Schnable, P. S., Ware, D., Fulton, R. S., Stein, J. C., Wei, F., Pasternak, S., Liang, C., Zhang, J.,

Fulton, L., Graves, T. A., et al. (2009). The B73 maize genome: complexity, diversity, and

dynamics. *Science* 326, 1112–1115. doi:10.1126/science.1178534.

Somerville, C., and Koornneef, M. (2002). A fortunate choice: the history of Arabidopsis as a model

plant. *Nat. Rev. Genet.* 3, 883–889. doi:10.1038/nrg927.

Türktaş, M., Yücebilgili Kurtoğlu, K., Dorado, G., Zhang, B., Hernandez, P., and Ünver, T. (2015).

Sequencing of plant genomes – a review. *Turk. J. Agric. For.* 39, 361–376. doi:10.3906/tar-1409-

93.

Varshney, R. K., Chen, W., Li, Y., Bharti, A. K., Saxena, R. K., Schlueter, J. A., Donoghue, M. T. A.,

Azam, S., Fan, G., Whaley, A. M., et al. (2011). Draft genome sequence of pigeonpea (Cajanus

cajan), an orphan legume crop of resource-poor farmers. *Nat. Biotechnol.* 30, 83–89.

doi:10.1038/nbt.2022.

Wang, X., Wang, H., Wang, J., Sun, R., Wu, J., Liu, S., Bai, Y., Mun, J.-H., Bancroft, I., Cheng, F., et

al. (2011). The genome of the mesopolyploid crop species Brassica rapa. *Nat. Genet.* 43, 1035–

1039. doi:10.1038/ng.919.

Yaminfirooz, M., and Ardali, F. R. (2018). Identifying the Factors Affecting Papers' Citability in the

Field of Medicine: an Evidence-based Approach Using 200 Highly and Lowly-cited Papers. *Acta*

Inform. Med. 26, 10–14. doi:10.5455/aim.2018.26.10-14.

8. Figures

Fig. 1: Timeline of critical technological advancements in Genome studies after *A. thaliana* genome

Sequencing.   (Adapted   from   *Nature   Milestones: Genomic   Sequencing*

https://www.nature.com/collections/ genomic-sequencing-milestones (2021))

Figure 2: Research output in *Arabidopsis* over the years

Figure 3: Thematic evolution based on the major keywords co-occurrence in two decades (2001-10)

& (2011-20)

Countries Wise Publication in Two Decades

Figure 4: Country-wise publication over the different period

Figure 5: Country-wise Scientific production based on the number of publications and Interaction

among countries publishing articles on *A. thaliana*. The blue colour's intensity corresponds to the total

number of publications from the corresponding country, and the red lines highlight the interactions

among countries (co-authorship).

Figure 6: Collaboration among different institutions working on *Arabidopsis* in the first decade

(2001-2010) vs second decade (2011-2020)

Figure 7: Quantitative and qualitative parameters in the research backed by different funding

organisations (2001-2020)

Figure 8: The Bradford law displaying key Arabidopsis journals

Figure 9: Performance of core journals (as defined by Bradford's law) over 2001-2020 period based

on total number of annual articles on 'Arabidopsis'

9. Tables

No	Title of the paper	Author and year	Affiliation of corresponding author	Journal	Citations (WoS)
1.	Genome wide insertional mutagenesis of *Arabidopsis thaliana*	Alonso J.M. et al., 2003	Salk Inst Biol Studies, Genome Anal Lab, 10010 N Torry Pines Rd, La Jolla, CA 92037 USA.	Science	3790
2.	*Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis	Yoo S.D., Cho Y.H. and Sheen J. (2007)	Massachusetts Gen Hosp, Dept Mol Biol, Boston, MA 02114 USA.	Nature Protocols	2344
3.	Genome-wide identification and testing of superior reference genes for transcript normalisation in *Arabidopsis*	Czechowski, T. et al., 2005	Max Planck Inst Mol Plant Physiol, D-14476 Postdam, Germany.	Plant Physiology	2018
4.	GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox	Zimmermann P. et al., 2004	ETH, Inst Plant Sci, CH-8092 Zurich, Switzerland.	Plant Physiology	1888
5.	A gene expression map of *Arabidopsis thaliana* development	Schmid, M. et al., 2005	Max Planck Inst Dev Biol, Spemannstr 37-39, D-72076 Tibingen, Germany	Nature Genetics	1787
6.	MAP kinase signalling cascade in *Arabidopsis* innate immunity	Asai, T. et al., 2002	Harvard Univ, Sch Med, Dept Genet, Boston, MA 02114 USA.	Nature	1684

359   **Table 1:** Highly cited research publications on *Arabidopsis* since its full genome was sequenced

363 **Table 2:** Top 10 countries which published the most research papers on *Arabidopsis* in the world

364 during 2001-2020

No.	Country	TP	% Share to N	h-index	ACPP	TC	WSC	% Share of self-citation
1	USA	9262	28.97%	313	75.92	703214	640485	8.9
2	PR China	6787	21.23%	166	29.96	203311	179953	11.5
3	Germany	4510	14.11%	223	66.71	300856	279745	7.0
4	Japan	3857	12.06%	200	56.75	218899	204002	6.8
5	France	2613	8.17%	188	64.89	169568	160796	5.2
6	England	2447	7.65%	204	75.74	185338	177949	4.0
7	S. Korea	1685	5.27%	122	39.78	67025	63347	5.5
8	Canada	1422	4.44%	127	51.15	72733	69609	4.3
9	Spain	1333	4.17%	129	56.49	75304	72118	4.2
10	Australia	979	3.06%	137	64.65	63293	61007	3.6

365 TP=Total publications, TC= Total citations, WSC=Without-self citation, ACPP= average citation per paper

367 **Table 3:** Top 10 Country vs Subject field wise (No. of publications)

Country/Region	Subject							Total*
	Plant Sciences	Bio-chemistry and Molecular biology	Cell Biology	Muti-disciplinary Sciences	Genetic heredity	Bio-technology applied micro-biology	Developmental biology	
USA	5549	3014	1760	1053	795	419	243	9262
China	4351	1976	753	616	410	361	55	6787
Germany	2885	1479	857	407	313	131	106	4510
Japan	2559	1206	1031	285	194	366	75	3857
France	1583	885	507	223	193	86	59	2613
England	1495	770	477	257	158	88	66	2447
S. Korea	1080	595	320	96	65	110	17	1685
Canada	1011	367	191	89	114	94	30	1422
Spain	880	392	194	133	82	57	40	1333
Australia	689	251	149	90	45	28	18	979

368 *The total number of articles in sub-subjects does not add up as only the five major subjects are mentioned here,

369 and there exist many journals which fall under more than one category

371   **Table 4:** Top journals that published most *Arabidopsis* research articles from 2001 to 2020

No.	Title of the journal	TP	% share	TC	WCS	*H*-Index	ACPP	IF (2020)
1	*Plant Physiology*	2490	7.79	226960	220788	204	91.15	6.902
2	*Plant Cell*	2376	7.43	287284	277416	248	120.21	9.618
3	*Plant Journal*	2260	7.07	179334	174911	190	79.35	6.141
4	*Journal of Experimental Botany*	1121	3.51	44776	44089	99	39.94	5.908
5	*Plant and Cell Physiology*	999	3.12	46314	45320	105	46.36	4.062
6	*Frontiers in Plant Science*	950	2.97	13016	12696	44	13.7	4.407
7	*PNAS*	878	2.75	102939	101908	180	117.24	9.412
8	*PLoS ONE*	829	2.59	23548	23269	67	28.41	2.740
9.	*Plant Signalling Behaviour*	804	2.52	8810	8728	36	10.96	1.671
10.	*Plant Molecular Biology*	677	.12	33566	33234	90	49.58	3.302