Fate-resolved gene regulatory signatures of individual B lymphocytes in the early stages of Epstein-Barr Virus infection

Elliott D. SoRelle[1,2*], Joanne Dai[1,3], Nicolás M. Reinoso-Vizcaino[1], Ashley P. Barry[1], Cliburn Chan[2], & Micah A. Luftig[1*]

[1] Department of Molecular Genetics and Microbiology, Duke Center for Virology, Duke University School of Medicine, Durham, NC 27710
[2] Department of Biostatistics and Bioinformatics, Duke University School of Medicine, Durham, NC 27710
[3] Current address: Amgen Inc., 1120 Veterans Blvd, South San Francisco, CA 94080

*Corresponding Authors: Elliott D. SoRelle & Micah A. Luftig

Abstract

Epstein-Barr Virus (EBV) infection of B lymphocytes elicits diverse host responses via complex, well-adapted transcriptional control dynamics. Consequently, this host-pathogen interaction provides a powerful system to explore fundamental cellular processes that contribute to consensus fate decisions including cell cycle arrest, apoptosis, proliferation, and differentiation. Here we capture these responses and fates with matched single-cell transcriptomics and chromatin accessibility, from which we construct a genome-wide multistate model of early infection dynamics. Notably, our model captures a previously uncharacterized EBV$^+$ analog of a multipotent activated precursor state that can yield early memory B cells. We also find that a marked global reduction in host chromatin accessibility occurs during the first stages of infection in subpopulations of EBV$^+$ cells that display senescent and pre-apoptotic hallmarks induced by innate antiviral sensing and proliferation-linked DNA damage. However, cells in proliferative infection trajectories exhibit greater accessibility at select host sites linked to B cell activation and survival genes as well as key regions within the viral genome. To further investigate such loci, we implement a bioinformatic workflow (crisp-ATAC) to identify phenotype-resolved regulatory signatures. This customizable method applies user-specified logical criteria to produce genome-wide single-cell ATAC- and ChIP-seq range intersections that are used as inputs for *cis*-linkage prediction and ontology tools. The resulting tri-modal data yield exquisitely detailed hierarchical perspectives of the transforming regulatory landscape during critical stages of an oncogenic viral infection that simulates antigen-induced B cell activation and differentiation. We anticipate these resources will guide investigations of gene regulatory modules controlling EBV-host dynamics, B cell effector fates, and lymphomagenesis.

Introduction

Epstein-Barr Virus (EBV) is an oncogenic gammaherpesvirus present in >90% of adults (Rickinson and Kieff, 2007) and associated with up to 2% of human cancers (Cohen et al., 2011). Recent reports have also provided epidemiological and mechanistic evidence supporting an etiological role for EBV in Multiple Sclerosis (MS) (Bjornevik et al., 2022; Lanz et al., 2022). In its initial stages, EBV infection within primary B lymphocytes manifests an array of host and viral programs. Upon entry into the host cell, the linear dsDNA viral genome rapidly circularizes to form an episome that is retained within the nucleus (Lindahl et al., 1976; Nonoyama and Pagano, 1972). Within hours to days, host innate immune responses are generated to restrict viral progression (Lünemann et al., 2015; Martin et al., 2007; Smith et al., 2013; Tsai et al., 2011). Simultaneously, viral genes are expressed to counteract host defenses (Ressing et al., 2015), co-opt B cell-intrinsic activation and proliferation (Calender et al., 1987; Thorley-Lawson, 2001; Thorley-Lawson and Mann, 1985), and attenuate DNA damage and stress responses instigated by virus-induced growth (McFadden et al., 2016; Nikitin et al., 2010). A consequence of these intimately adapted host-pathogen dynamics is that EBV infection can precipitate diverse responses and outcomes for host B cells. These include unsuccessful infection routes resulting from effective antiviral restriction and DNA damage-induced growth arrest as well as successful infection leading to immortalization *in vitro* (Bird, 1981; Henle et al., 1967; Pope et al., 1968; Zhao et al., 2011) or lifelong latency *in vivo* within memory B cells (Babcock, 1998; Longnecker et al., 2013; Miyashita et al., 1997) that retain oncogenic potential (Raab-Traub, 2007; Thorley-Lawson and Gross, 2004).

Since its discovery in 1964 as the first human tumor virus (Epstein et al., 1964; Young and Rickinson, 2004), extensive research has revealed the molecular means by which EBV establishes infection and underlies various malignancies. The entire EBV genome is ~172 kilobases and contains at least 80 protein-coding sequences including six EBV nuclear antigens (EBNAs); several latent membrane proteins (LMPs); and loci that encode replicative and transcriptional machinery as well as structural proteins. The EBV genome also contains functional non-coding RNAs: the BHRF and BART microRNAs and the EBV-encoding regions (EBERs) (Rickinson and Kieff, 2007; Young et al., 2007).

EBNAs are especially important in establishing distinct forms of latency depending on their combinatorial expression (Price and Luftig, 2015). EBNA1 is a transcription factor (TF) that is essential for viral genome maintenance and B cell transformation and ubiquitously binds and epigenetically regulates host chromatin (Altmann et al., 2006; Canaan et al., 2009; Dheekollu et

al., 2021; Humme et al., 2003; Lamontagne et al., 2021; Lu et al., 2010; Lupton and Levine, 1985;

Wood et al., 2007; Yates et al., 1985). EBNALP is another essential factor (Mannick et al., 1991;

Szymula et al., 2018) that initiates host cell proliferation alongside its co-activated target, EBNA2

(Alfieri et al., 1991; Harada and Kieff, 1997; Sinclair et al., 1994), and interacts with several host

proteins including TFs (Han et al., 2001; Ling et al., 2005; Matsuda et al., 2003). EBNA2 is likewise

required for B cell immortalization (Cohen et al., 1989), notably through coordination with host

TFs and their binding sites (Lu et al., 2016; Zhao et al., 2011) and with EBNALP to drive early cell

proliferation and viral LMP1 expression (Peng et al., 2005). The EBNA3 proteins (EBNA3A,

EBNA3B, and EBNA3C) mediate a delicate balance of anti- and pro-oncogenic processes (Allday

et al., 2015; Banerjee et al., 2014; Parker et al., 1996; Tomkinson et al., 1993; White et al., 2010).

These include epigenetic repression of host tumor suppressor genes (*BIM, p14, p16*) and viral

promoters (Maruo et al., 2011; Paschos et al., 2012; Saha et al., 2015; Skalska et al., 2010; Styles

et al., 2017), competitive binding of the EBNA2-interacting host factor RBPJ (Robertson et al.,

1995; Wang et al., 2015), and inhibition of apoptosis (Price et al., 2017). Collectively, the EBNAs

reshape the nuclear regulatory and transcriptional landscape of EBV$^+$ B cells, effectively hijacking

B cell-intrinsic activation, expansion, and differentiation programs. Thus, EBV co-opts antigen-

responsive host immune mechanisms for the ulterior purposes of viral replication and

propagation.

While the EBNAs engage cell proliferation machinery at the epigenetic and transcriptional level

in the nucleus, the LMPs (LMP1, LMP2A, and LMP2B) do so at the cell membrane by simulating

antigen-induced signal transduction pathways. The essential LMP1 promotes B cell activation

through mimicry of a constitutively active CD40 receptor (Kilger et al., 1998; Uchida et al., 1999)

and interacts with Tumor Necrosis Factor (TNF) receptor-associated factors (TRAFs) to activate

NF-κB pathway signaling via IKK (Devergne et al., 1996; Eliopoulos et al., 2003; Greenfeld et al.,

2015; Luftig et al., 2003). These interactions induce anti-apoptotic pathways, MHC-mediated

immune recognition, pro-inflammatory responses, and cell migration. Downstream consequences

include oncogenic proliferation and survival but also induction of pro-apoptotic responses

(Devergne et al., 1998; Fries et al., 1999; Greenfeld et al., 2015; Henderson et al., 1991; Shair et

al., 2008; Wang et al., 2017). Thus, as in antigen-induced B cell activation (and subsequent

differentiation), adept regulatory control of NF-κB signaling (Hoffmann et al., 2002; Mitchell et al.,

2018; O'Dea et al., 2007; Roy et al., 2019) is dispositive for the fate of a given EBV$^+$ B cell.

Although it is not essential for transformation, LMP2A promotes cell survival through mimicry of a

stimulated B cell receptor (BCR), which activates signaling cascades complementary to those

induced by LMP1 (Anderson and Longnecker, 2008; Fish et al., 2020; Guasparri et al., 2008;

Portis and Longnecker, 2004). LMP2A expression further predisposes EBV+ B cells to survival by

lowering antigen affinity selection thresholds *in vivo* (Minamitani et al., 2015). Thus, EBV latent

membrane proteins play integral roles in B cell proliferation in the absence of antigen licensing

and in avoiding replicative dead ends effected by antiviral sensing.

Clearly, key EBV gene products manipulate diverse host programs at early stages to achieve

sustained latency (Mrozek-Gorska et al., 2019; Pich et al., 2019). Many such perturbations involve

extensive rewiring of epigenetic and transcriptional regulatory modules. EBV researchers have

used methods such as RNA-, ATAC- (Assay for Transposase-Accessible Chromatin), and ChIP-

seq (Chromatin Immunoprecipitation) to study these changes at various levels in the gene

regulatory hierarchy within early infected cells and transformed lymphoblastoid cell lines (LCLs)

(Arvey et al., 2012; Jiang et al., 2017; McClellan et al., 2013; Mrozek-Gorska et al., 2019; Wang

et al., 2019; Zhou et al., 2015). Recently, the epigenetic and transcriptional roles of EBNA1 were

interrogated through time-resolved multi-omics (Lamontagne et al., 2021). While these and other

studies provide indispensable insights regarding virus-induced genome-wide expression and

regulation, they have relied on bulk ensemble sequencing. Such assays yield population-

averaged measurements that obscure variation arising from intrinsic stochasticity (Raj et al.,

2006; Raj et al., 2010; Raj and Van Oudenaarden, 2008), asynchronous behaviors, and

heterogeneous cell subsets. Specifically, ensemble averaging fails to capture cell-matched

measurements across genes, which precludes identification of coordinated expression programs

or epigenomic regulatory patterns in specific phenotypes. By contrast, single-cell sequencing

provides refined genome-wide views of expression and regulation that preserve the ability for the

identification of heterogenous cell states with low bias (Buenrostro et al., 2015; Junker and van

Oudenaarden, 2014; Shalek et al., 2013; Shapiro et al., 2013; Wills et al., 2013). Given the

complexity of host-virus relationships, single-cell -omics approaches are essential to dissect the

early stages of EBV infection and the distinct fate trajectories it comprises. We previously used

single-cell RNA sequencing (scRNA-seq) to identify EBV-driven heterogeneity in LCLs (SoRelle

et al., 2021). Recent advances in single-cell multimodal -omics methods have made it possible to

integrate scRNA-seq with several levels of hierarchical regulation (Efremova and Teichmann,

2020), which can provide greater insight into the mechanistic origins underlying gene expression.

These include techniques for obtaining cell-matched measurements of mRNA transcripts,

chromatin accessibility, and DNA methylation status (Cao et al., 2018; Chen et al., 2019; Clark et

al., 2018; Zhu et al., 2019), as well as other molecular levels. In this work, we leverage single-cell

multiomics (scRNA-seq + scATAC-seq) to capture and explore the distinct gene expression and

regulatory signatures that determine the course of EBV infection in primary human B lymphocytes.

Results

EBV asynchronously induces primary B cells into distinct phenotypic states early after infection

To interrogate chromatin accessibility and gene expression changes that occur upon EBV infection, we isolated primary human B cells from the peripheral blood of two donors and infected them with the B95-8 strain of EBV. Infections were performed at a multiplicity of infection (MOI) of 5 to ensure latent gene expression in every cell (Nikitin et al., 2010). We cryopreserved samples of infected cells at 2-, 5-, and 8-days post-infection in addition to uninfected cells (Day 0) from each donor sample following B cell enrichment. Cell samples from each donor and timepoint were simultaneously thawed, prepared to >90% viability, and processed into single-cell multiome libraries. Single-cell matched transcript and accessibility data were obtained through standard NGS, alignment, counting, and quality control (QC) methods (**Table S1**).

EBV infection induced broad transcriptomic changes in B lymphocytes at high efficiency, as evidenced by the near-complete loss of resting phenotypes (Day 0) within two days of infection. New states emerged between Day 2 and Day 5, while subtle shifts in state proportions defined the period between Day 5 and Day 8 (**Figure 1A**). Total and unique transcripts per cell increased, particularly between Day 0 and Day 2, while mitochondrial gene expression increased gradually (**Figure 1B**). Total transcript and mitochondrial distributions at Day 2 exhibited two modes, which was consistent with the presence of both non-proliferative and mitotic cells identified by S-phase and G2M-phase marker scoring (**Figure 1C**).

Unsupervised methods revealed subpopulations (clusters) in cell cycle-regressed aggregated scRNA-seq time courses. Two clusters corresponded to uninfected B cells (c3, c8); seven were post-infection B cell phenotypes (c0, c1, c2, c4, c5, c6, c7); and two were T cells (c9) and CD14$^+$ monocytes (c10) carried over from PBMCs despite extensive B-cell enrichment (**Figure 1D**). Genome-wide expression correlation was higher among post-infection states relative to uninfected cells, and certain phenotypes were more strongly correlated (e.g., c0 with c1; c4 with c7, **Figure 1E**). Sorting cluster membership by day yielded coarse-grained dynamics of cell state transitions (**Figure 1F**). We determined top differential genes in each cluster (one-vs-all-others) to inform state identity annotations (**Figure 1G**). Identified clusters included many genes known to be modulated in EBV infection and were broadly consistent across both donors with respect to top marker genes, cell population frequencies, and temporal emergence (**Figures S1-S4**).

<u>Infected cell state heterogeneity is linked to antiviral and B cell-intrinsic responses</u>

Cluster analysis deconvolved heterogeneous biological states within each sample and

revealed phenotypes retained across multiple timepoints (**Figure 2**). At this resolution, we

estimated time- and state-level trends in viral gene expression, variation in metabolic activity, and

transcript diversity (**Figure 2A**). Overall, c0, c1, c2, c5, and c6 exhibited the highest levels of EBV

transcripts and more unique transcripts than c3, c4, c7, and c8. Mitochondrial gene fraction and

unique feature content were highest in c6 and lowest in c4 and c7, although c7 had a long-tailed

distribution of mitochondrial expression (20-80%) prior to QC, indicative of (pre-) apoptotic cells.

All clusters except uninfected B cells (c3, c8) displayed broad innate antiviral and interferon-

stimulated gene (*ISG* and *IFI* member) expression. Antiviral gene expression was generally higher

and exhibited greater variance in c7 than c4 and persisted at roughly uniform levels in c0, c1, c2,

c5, and c6 (**Figure 2B**). Through joint consideration of cluster-resolved expression trends for viral,

mitochondrial, and interferon-stimulated genes, we distinguished uninfected cells (c3, c8) and two

classes of cells with the hallmarks of antiviral response: those with low proliferation and negligible

viral expression (c4, c7) and those with viral and metabolic indicators of progressive EBV infection

(c0, c1, c2, c5, c6).

Next, we extensively analyzed differentially expressed genes (DEGs) among clusters and

groups, including pairwise comparisons of all post-infection phenotypes (**Figures 2C, S5-S11**).

The two resting cell phenotypes differed in their expression of *IGHD*, *IGHM*, *CD27*, and other

markers that distinguish naïve (c8) from memory (c3) B cells. In addition to interferon response

signatures, non-proliferating infected cells exhibited an overall reduction in gene expression and

upregulated stress response markers. These included the highest overall expression of actin

sequestration genes (*TMSB10*, *TMSB4X*) and, particularly within c7, numerous ribosomal subunit

genes (e.g., *RPS27A*). Cells in c4 were distinguished by elevated expression of *MARCH1*, which

encodes an E3 ubiquitin ligase that regulates the type I interferon response ([Wu et al., 2020](#)).

Unlike c4, c7 cells also contained high transcript levels for genes involved in oxidative stress

(*TXN*, *FTL*, *FTH1*), cytochrome oxidase subunits (e.g., *COX7C*), ubiquitin genes (*UBA52*, *UBL5*)

and highly variable mitochondrial fractions. Among EBV$^+$ cells with hallmarks of elevated

respiration, those in c6 were most clearly consistent with proliferating cells based on upregulated

cell cycle markers. Cells in c0 were distinguished by upregulation of *FCRL5* and *LY86-AS1*, an

antisense RNA to a lymphocyte antigen (*LY86*) that mediates innate immune responses. Cells in

this cluster also displayed markers consistent with the early stages of pre- germinal center

activated B cells (e.g., *CCR6, CD69, POU2AF1, TNFRSF13B, PIK3AP1*). Notably, cells in c1 and

c2 contained the highest levels of the EBV gene *BHRF1*. Between these two phenotypes, c2 was

enriched for genes involved in NF-κB signaling and known markers of EBV-mediated B cell
activation (*NFKBIA*, *TNFAIP3*, *EBI3*) while c1 appeared to be derived from naïve cells (based on
IGHD and other carryover genes) and exhibited near-unique expression of *SH3RF3/POSH2* and
FIRRE, a MYC-regulated long non-coding RNA (lncRNA). Finally, c5 displayed upregulation of
immunoglobulin heavy and light chains (*IGHA1, IGHG1, IGHM, IGKC, IGLC1-3*) as well as genes
involved in B cell differentiation (*MZB1, PRDM1/BLIMP1, XBP1*). Gene ontology (GO) networks
were also generated for top DEGs from one-versus-all-other comparisons to facilitate phenotype
annotations (**Figures S12-S16**).

<u>A map of B cell phenotypes and fate trajectories in early EBV infection</u>
Graph-based pseudotime ([Qiu et al., 2017](#)) approximated EBV-induced state transitions when
anchored from resting cells (**Figure 3A**). Pseudotime scoring was used to track state dynamics
of the top 25 marker genes for each phenotype and four example expression trajectories are
highlighted (**Figure 3B**). Collectively, flow cytometry for the B cell marker CD19 and CD23
(upregulated in EBV infection) at each timepoint (**Figure S17**), cluster-specific DEGs, network
ontologies, and pseudotime led us to propose a multi-phenotype model for heterogeneous cell
fate trajectories (**Figures 3C, S18-S19**) that manifest in early EBV infection *in vitro*. In this model,
naïve (c8) and memory (c3) B cells infected with EBV either undergo antiviral response-mediated
arrest (c4) or EBV-driven hyperproliferation (c6) within several days of infection.
Hyperproliferating cells can subsequently enter one of several activated states (c0, c1, c2) or
undergo growth arrest (c7). Further, differentiated B cells (c5) can develop following activation in
a manner analogous to effector cell exit from the germinal center reaction.
Among activated phenotypes, c2 matched classical EBV-mediated activation of NF-κB
pathway genes, apoptotic regulators, and other known biomarkers ([Cahir-McFarland et al., 2004](#);
[Messinger et al., 2019](#)). Cells in c1 were consistent with a related activation intermediate that
originated from EBV$^+$ naïve cells. Despite the relatively short timecourse, c2 and c5 began to
reflect the continuum of activation and differentiation phenotypes we previously characterized in
LCLs ([SoRelle et al., 2021](#)), which are considered to be immortalized at 21-28 days post-infection
([Nilsson et al., 1971](#)). We confirmed these similarities by merging Day 8 and the LCL GM12878,
for which scRNA-seq data was previously reported and analyzed ([Osorio et al., 2019](#); [Osorio et
al., 2020](#); [SoRelle et al., 2021](#)) (**Figure S20**). Conceivably, EBV$^+$ cells could also transition to a
plasmablast phenotype (c5) from memory cells (c3) through hyperproliferation (c6) via division-
linked differentiation ([Hodgkin et al., 1996](#)), effectively bypassing intermediate states.

Conversely, cells in c7 highlighted diverse origins of EBV[+] cell growth arrest, apoptosis, and

senescence, which each provide host defenses against oncogenic malignancies (Bartkova et al.,

2006; Nikitin et al., 2010). In addition to highly variable mitochondrial expression and the lowest

transcript levels of any state, this phenotype was defined by broad upregulation of genes involved

in ribosome biogenesis-mediated senescence (*RPS14*, *RPL29*, *RPS11*, *RPL5*) (Lessard et al.,

2018; Nishimura et al., 2015) and stress-associated sequestration of actin monomers that favor

G-actin formation (*TMSB4X*, *TMSB10*, *PFN1*) (Kwak et al., 2004). A subset of cells within c7 also

contained elevated levels of cell cycle markers (*MKI67*, *TOP2A*, *CCNB1*, *CENPF*) carried over

from pre-arrest hyperproliferation (**Figure S21**).

<u>Evidence for EBV induction of an activated precursor to early memory B cells (AP-eMBC)</u>

We next sought to compare early infected phenotypes from our multistate model with cells

isolated from secondary lymphoid organs. We acquired single-cell RNA-seq data from human

tonsil tissue and identified germinal center (GC) cell subsets (**Figure 4A**), which we analyzed

alongside early infection phenotypes of interest. EBV[+] NF-κB activated cells (c2) clearly mimicked

GC light zone (LZ) B cells; *MKI67*[hi] cells (c6) matched actively cycling cells (including GC dark

zone (DZ) B cells); and EBV[+] differentiated cells (c5) matched plasmablasts and plasma cells (PB

/ PC). Cells in c0 were most like pre-GC naïve and memory B cell (MBC) subsets (**Figure 4B-D**).

Further, numerous c0 markers were consistent with both pre-GC activated B cells (*SELL, BANK1,*

CD69, GPR183 (EBI2)) and memory B cell phenotypes (*SELL, BANK1, GPR183, PLAC8*)

recently identified from scRNA-seq of tonsils in response to antigen challenge (King et al., 2021)

(**Figure 4C-D**). Cells in c0 further exhibited upregulation of genes with essential roles in B cell

activation (*TNFRSF13B/TACI*) (Wu et al., 2000) and germinal center formation (*POU2AF1/OCA-*

B) (Kim et al., 1996; Luo and Roeder, 1995; Schubart et al., 1996) (**Figure S18**). Moreover, c0

displayed elevated *CCR6*, a marker of an activated precursor (AP) state that can generate early

memory B cells (eMBCs) (Glaros et al., 2021; Suan et al., 2017) (**Figure 5A**).

We subsequently validated the generation of CCR6[+] AP-eMBC B cells in response to EBV

infection through time-resolved FACS (**Figures 5B-D, S22-S24**). Resting B cells were CCR6[lo]

and remained so until between 2 and 5 days after infection. Further, we observed that the most

proliferative cell fraction at day 8 was CCR6[lo] and a moderately proliferative cell population was

CCR6[hi]. While the most proliferative cells were CCR6[lo]/CD23[lo], the proliferative CCR6[hi] cells

displayed variable CD23 levels (**Figure 5B**). Consistent with our scRNA time course,

CCR6[hi]/CD23[hi] and CCR6[hi]/CD23[lo] populations respectively corresponded to c1/c2 and c0 and

emerged within 5 days (**Figure 5C**). Based on CD27 and IgD status, these populations

predominantly originated from naïve or non-switched memory versus switched memory cells,
respectively; notably, cells from these different resting phenotypes were present in each
population gated by CCR6 and CD23 status (**Figure 5D, S24C**). Rapidly proliferative
CCR6^{lo}/CD23^{lo}/CD27^{hi}/IgD^{lo} cells were consistent with infected memory B cells transitioning to
plasmablasts (*c3→c6→c5* model trajectory; ~72% of CCR6^{lo}/CD23^{lo} cells). Marginally less
proliferative CCR6^{hi}/CD23^{lo}/CD27^{hi}/IgD^{lo} cells were consistent with stimulated AP-eMBCs
(*c3→c6→c0* model trajectory; ~74% of CCR6^{hi}/CD23^{lo} cells). We also observed an IgD^{hi} naïve
population that matched the pre-GC AP-eMBC phenotype (*c8→c6→c0*; (~25% of CCR6^{hi}/CD23^{lo}
cells). Finally, an even less proliferative CCR6^{hi}/CD23^{hi}/IgD^{hi} population matched activated naïve
(or non-switched memory) cells destined for GC BC (*c8→c6/c0→c1/c2*; ~80% of CCR6^{hi}/CD23^{hi}
cells) and the minor subset (~17%) of CCR6^{hi}/CD23^{hi} cells that was IgD^{lo} was consistent with
MBCs induced by EBV to undergo a pseudo-GC reaction (*c3→c6/c0→c2*) (**Figure 5C-E**).
Intriguingly, a subset of CCR6^{+} cells displaying the AP phenotype apparently persists long after
the early stages of infection based on scRNA-seq data from LCLs (**Figure S25**). Thus, c0 in our
model matches a virus-induced common progenitor state from which PBs, GC BCs, and early
MBCs have been shown to originate in response to antigen stimulation ([Taylor et al., 2015](#)). Our
results further indicate that both naïve and memory B cells can achieve this multipotent state at
different frequencies upon *in vitro* infection and that the AP-eMBC phenotype is perpetuated in
EBV-immortalized B cells.
294
<u>Linked expression and accessibility illuminate regulatory mechanisms in phenotype trajectories</u>
We next investigated potential regulatory mechanisms underlying DEGs observed across
phenotypes. Expression data were jointly analyzed with cell-matched measurements from single-
cell Assay for Transposase-Accessible Chromatin sequencing (scATAC-seq) and annotated by
state (**Figures 6A-B**, **S26A-B**). While total and unique transcripts per cell increased through early
infection, global chromatin accessibility decreased substantially upon infection. Resting (c3, c8)
and hyperproliferative (c6) cells had the highest overall accessibility. There were significantly
more peaks in the NF-κB activation state (c2) relative to other activation intermediates (c0 and
c1; two-tailed t-test, $p < 2.2 \times 10^{-16}$ and 1.4×10^{-14}, respectively) and differentiated cells (c5; $p <$
2.2×10^{-16}). Similar accessibility reduction occurred in both donors in the first five days, with
increased accessibility recovered between Day 5 and Day 8 (including to higher than resting
levels in one donor) (**Figures 6B, S27**). This indicated that EBV-induced heterochromatinization
is likely transient in successfully infected cells (i.e., those that evade innate- and damage-
mediated arrest).

We found 954 linked feature genes derived from the top 100 marker genes for each cluster

using the multimodal integration capabilities of the Signac package in R ([Stuart et al., 2021](#)). Of

these 954 genes, 177 were significant DEGs with linkages to 476 differentially accessible peaks

(DAPs). This translated to 18.6% of tested genes with potential DAP-linked regulation (**Figures**

6C, S26C-D). We identified genes linked (in *cis*) to DAPs to explore phenotype-associated gene

regulatory relationships (**Figure S28A**). Joint analysis of DAP-linked DEGs yielded four regulatory

patterns: higher accessibility with higher expression (+/+); lower accessibility with higher

expression (-/+); lower accessibility with lower expression (-/-); and higher accessibility with lower

expression (+/-). (**Figure S28B**). The +/+ and -/- patterns were characteristic of positive regulatory

sites. The less frequently observed -/+ and +/- patterns were consistent with closure or opening

of ATAC sites with negative regulatory functions, respectively. Analysis of genes of interest

including *CCR7*, *CXCR4*, *RUNX3*, *BACH2*, *JCHAIN*, and *PRDM1* provided examples of each

regulatory pattern and their variation among states (**Figures S28C, S29-S30**).

We developed joint scRNA + scATAC profiles for major infection fate trajectories in our model

(**Figure 6C-F**). The path from resting cells to EBV$^+$ arrested/senescent cells (*c38→c47*) was

characterized by global reductions in accessibility and expression. 34.3% of all DEGs between

resting cells and these non-proliferative EBV$^+$ fates were linked to DAPs that become inaccessible

after infection (*c38!c47* peaks). Top DAP-linked DEGs in innate arrested cells (c4) corresponded

to upregulation of interferon-responsive genes and downregulation of mitochondrial genes, while

stress- and damage-induced senescent cells (c7) were distinguished by their high expression of

interferon-stimulated genes (e.g., *ISG15*) and ribosomal transcripts (e.g., *RPS14*). Cells in each

of these clusters also displayed reduced expression of the proto-oncogenic tyrosine kinase gene

LYN linked to closure of multiple regulatory sites following infection (**Figure 6D**).

The reduction in ATAC peaks within c7 was consistent with the formation of senescence-

associated heterochromatin foci (SAHF) ([Courtois-Cox et al., 2008](#); [Di Micco et al., 2011](#); [Lenain](#)

[et al., 2017](#)) (**Figure S31**). Because senescence can arise from diverse mechanisms such as

innate immune sensing or growth-induced DNA damage, we used higher resolution clustering to

reveal c7 subsets (7a and 7b). These subsets displayed DEGs involved in the cell cycle and

antiviral sensing (**Figure S32**). Different *HMGB2* levels between 7a and 7b were notable, as this

gene's product mediates diverse roles in sensing ([Yanai et al., 2009](#)), double-stranded break

repair ([Krynetskaia et al., 2009](#)), and p53 downregulation ([Stros et al., 2002](#)). Relative to resting

B cells, *HMGB2* expression was strongly elevated in 7b (as in the hyperproliferative state, c6) but

only mildly so in 7a (similar to c4, which could precede senescence ([Glück et al., 2017](#))). Similarly,

cell cycle markers were lower in 7a than 7b. Thus, 7a was consistent with EBV$^+$ cells that arrest

almost immediately via innate sensing and become senescent, whereas 7b matched a trajectory

in which EBV⁺ hyperproliferative cells become senescent following replicative stress response

induction. Notably, both 7a and 7b exhibited elevated levels of ribosomal subunit mRNAs (**Figure**

S32C-D).

NF-κB activated EBV⁺ cells (c2) exhibited loss of accessibility at 1,142 sites present within

resting cells in both donors (8.7% of all resting cell peaks). This reduction paralleled upregulated

expression of the polycomb group repressor *EZH2* and a polycomb-interacting methyltransferase,

DNMT1 (**Figure S33**). However, EBV-activated cells possessed 668 peaks absent in resting cells

(*c2!c38*) that were linked to 595 unique genes. 154 of these (25.9%) were DEGs between resting

and activated cells (**Figure 6E**). These 154 *c2!c38* DAP-linked DEGs included 109 upregulated

and 45 downregulated genes from *c38→c2*. Upregulated genes included regulators of apoptosis

and tumor suppression (*BCL2A1, TNFRSF8, PDCD1LG2, ST7, IQGAP2, TOPBP1, CD86*);

proliferation (*CDCA7, MKI67*); B cell signaling (*NFKBIA, MAPK6, TNIP1, TRAF3*); inflammation

(*SLC7A11, RXRA, ZC3H12C*); oxidative stress (*SLC15A4, TXN*); and epigenetic remodeling

(*AHRR, NCOR2*). The 45 downregulated *c2!c38* DAP-linked DEGs included *CCR7,*

acetyltransferases (*EPC1, KAT6B*), apoptotic and stress response regulators (*STK39,*

STK17A/DRAK1, VOPP1, ZDHHC14), negative regulators of B cell signaling (*CBLB*), and the

tumor suppressor *ARRDC3* (**Figure S34**).

In a third example, we explored DAP-linked DEGs between EBV-induced activated (c2) and

differentiated (c5) phenotypes. Activated cells exhibited 999 called peaks that were absent in

differentiated cells (*c2!c5* DAPs) while only 13 new peaks emerged in differentiated cells (*c5!c2*

DAPs) in both donors. This corresponded to a 15% net reduction in accessible peaks in the

c2→c5 transition. Notably, *c2!c5* DAPs found in both donors were linked to 13.4% of all DEGs

identified between these states in the scRNA assay. Key *c2→c5* DAP-linked DEG dynamics

included downregulation of NF-κB family genes and upregulation of plasmablast-specific

transcriptional regulators, translation factors, and protein export machinery (i.e., facilitators of Ig

synthesis, secretion, and protein folding chaperones) (**Figures 6F, S10**).

By mapping multiome reads to a concatenated reference (human + EBV), we were able to

detect increased accessibility within the EBV genome over time after infection. We detected 21

unique viral ATAC peaks (20 of 21 common to both donors) including at TSSs for essential viral

genes such as the *EBNAs* and *LMP1* (**Figure S35**). Quantification of episome peak-containing

cells by phenotype revealed that EBV⁺ activated and hyperproliferative cells had the greatest

number of accessible loci relative to other post-infection phenotypes. These sites included the C

promoter (Cp) for *EBNA1*, *EBNA2*, and *EBNA3A-C*; the *LMP1* TSS; the TSS for *BMRF1*, a DNA

polymerase accessory protein; and the *BHLF1* locus, which was recently recognized as a
facilitator of latency and B cell immortalization (Yetming et al., 2020) (**Figure S36A-B**). Innate
arrested cells (c4) exhibited the lowest frequency of cells with accessible episomal loci, followed
by growth-arrested cells (c7) (**Figure S36C**).

<u>Post-infection cell fates exhibit differential enrichment of TF motifs</u>

To further investigate regulatory differences by phenotype, we assayed TF binding motif
enrichment by state (**Figure S37**). We identified variable motif enrichment linked to resting cell
phenotypes and among non-arrested post-infection states (**Figure S37A**). Variation in accessible
motifs broadly aligned with phenotypic gene expression with respect to antiviral response
induction, promotion of cell growth, and oncogenesis (**Figure S37B-C**). Activated B cell (c2)
ATAC peaks were enriched in binding sites for proto-oncogenic TFs including members of the
REL (cREL, RelA, RelB), AP-1 (FOS, FOSB, JUNB, JUND), and EGR (EGR1-4) families.
Enhanced accessibility at NF-κB family binding sites within activated cells was noteworthy, given
the observed concurrent upregulation of NF-κB pathway gene expression. Similar phenotypic
consistency was observed within differentiated cells (c5), which were enriched in accessible
motifs for IRF4, IRF8, and XBP1. Globally, both resting B cell phenotypes and the innate sensing
arrest state shared the greatest motif correlation with each other (R>0.75) and the lowest
correlation with EBV-activated and hyperproliferating cells (0.55<R<0.7) (**Figure S38**).

<u>An informatics pipeline to infer phenotype-resolved TF signatures and gene regulatory elements</u>

The prevalence of DAPs linked to DEGs known to be modulated in *trans* by EBV gene products
led us to interrogate phenotype-specific TF signatures genome-wide. To do so, we employed a
bioinformatic workflow to obtain <u>C</u>hIP-seq <u>r</u>eferenced <u>i</u>nferences of <u>s</u>ingle-cell <u>p</u>henotypes from
sc<u>ATAC</u> data, which we termed "crisp-ATAC" (**Figures S39, S40**). We expected that ensemble-
averaged ChIP data from an appropriate reference cell type would contain TF binding (and
epigenetic) data from a superposition of cell phenotypes at high coverage, thus maximizing
chances to identify overlaps with comparatively sparse scATAC cluster data. We further reasoned
that phenotypic variation in TF binding site accessibility would have biological consequences.
We sought to predict viral EBNA and LMP1-mediated NF-κB accessible sites at promoters,
enhancers, and actively transcribed genes in each state of our model. To do so, we applied crisp-
ATAC recipes to intersect peaks from each scATAC phenotype with ChIP-seq peaks for viral
EBNAs, NF-κB/Rel TFs, H3K4me1, H3K4me3, H3K27ac, H3K36me3, and RNA Pol II (Jiang and
Mortazavi, 2018) (**Figure S39C**). Hyperproliferative cells (c6), EBV-activated cells (c2), and

resting memory B cells (c3) exhibited up to 3-fold more enhancers and promoters at known

EBNA2 binding sites relative to naïve B cells (c8), other activation intermediates (c0, c1),

plasmablasts (c5), arrested states (c4, c7), and non-B cells (c9, c10). Similar patterns were found

for EBNA3C and EBNALP sites (**Figure S39C**, left column) as well as Rel family TF binding sites

(cRel, RelA, and RelB). Enhancers, promoters, and actively transcribed genes were consistently

enriched in c2, c3, and c6 and depleted in c4 and c7, with intermediate levels present in c0, c1,

c5, and c8. By accounting for peaks conserved in both biological replicates, we demonstrated

low-noise measurements of DAPs for use with crisp-ATAC and characterized DAP frequencies

and interval length distributions across all pairwise phenotype comparisons (**Figure S40**).

<u>crisp-ATAC finds TF-linked expression signatures that vary across distinct EBV$^+$ cell fates</u>

We applied crisp-ATAC to capture regulatory variation among infection phenotypes with

respect to key viral transcriptional co-activators. We first compared the innate sensing arrest (c4)

and NF-κB (c2) states (**Figure 7A**), as these represent starkly different post-infection fates. Peak

data were extracted and gated to obtain *c2!c4* DAPs present in both donors (n=1,873), which

yielded linked gene predictions (n=1,514) (**Figure 7B-C**). The *c2!c4* linked gene ontology network

was enriched for innate defense (inflammation, antimicrobial processes) and EBV-induced

responses (lymphocyte activation, regulation of apoptosis) (**Figure 7D**). Taking a macroscopic

view, we found that predicted *c2!c4* linked genes included 42.5% (71 of 167) of known EBV super-

enhancer (EBVSE) site-linked genes (Zhou et al., 2015). Consequently, 41-55% of EBNA-

associated *c2!c4* DAPs also overlapped a peak for the super-enhancer-associated host TF RelA.

Of the 71 EBVSE-linked genes identified for *c2!c4*, 19 (27%) were linked to EBNALP ChIP-seq

peaks; 22 (31%) to EBNA2 ChIP-seq peaks; and 15 (21%) to EBNA3C ChIP-seq peaks. EBVSE-

linked genes were enriched in EBNA-associated DAP-linked DEGs relative to size-matched

random samples of genes in the captured transcriptome (**Figure 7E**).

We analyzed specific genes of interest based on 1) EBVSE membership, 2) GO process

involvement, and/or 3) empirically demonstrated importance to EBV infection. The NF-κB

activated gene and signal transducer *TRAF1*, whose gene product interacts with viral LMP1, was

identified through all three of these routes (Devergne et al., 1996; Eliopoulos et al., 2003; Fries et

al., 1999; Greenfeld et al., 2015; Sandberg et al., 1997). We found *c2!c4* DAPs associated with

one or more EBNA at -3kb, +2kb, and +37kb relative to the *TRAF1* TSS, each with significant

positive correlation to *TRAF1* expression (p<0.05 for correlation z-score). Notably, these EBNA-

associated regulatory loci exhibited reduced accessibility in stress arrest (c7), activation

intermediate (c1) and differentiated (c5) states compared with c2 (**Figure 7F**).

We used a grouped crisp-ATAC comparison (*c256!c38*) to study changes associated with the
trajectory for successful EBV-induced B cell immortalization (**Figure S41**). We analyzed viral co-
activator-associated DAPs between proliferative (c6) and LCL-like phenotypes (c2, c5) versus
resting B cells (c3, c8). Despite the net reduction in accessibility after infection, we identified 245
unique genes linked to 1,824 peaks present in all tested EBV$^+$ states (c256) but absent in both
resting phenotypes (c38) (**Figure S41A-C**). 166 of the 245 genes were linked to a binding site for
at least one EBNA, and 18 of these genes overlapped with EBVSE targets (7.3% of predicted
genes, 10.8% of known EBVSE genes). Only 31 GO process terms were shared across the top
100 terms for each tested EBNA, accounting for 15% of unique terms (**Figure S41D**). We selected
the *c256!c38*∩EBNA targets *TNFRSF8* (CD30), *CD274* (PD-L1), and *PDCDL1G2* (PD-L2) based
on their therapeutic relevance to EBV-associated lymphomas. For each gene, we confirmed the
presence of EBNA-associated DAP-linked DEGs by phenotype. These included three
EBNA2$_{c256!c38}$ sites near the *TNFRSF8* TSS (-17kb, -12kb, and +16kb) and a shared multi-EBNA
site -17kb from the *PDCDL1G2* TSS and +43kb from the end of the *CD274* gene (**Figure S41E**).
These loci were enriched for Rel sites and activating histone marks in LCL reference data.
In a final example, we evaluated activated versus differentiated EBV$^+$ phenotypes (*c2!c5*) sites
with known viral transcriptional co-activator binding in both donors to explore regulatory
relationships that distinguished the phenotypes present in LCLs ([SoRelle et al., 2021](#)) (**Figure**
S42A). 519 of 999 identified *c2!c5* peaks intersected with at least one EBNA binding site, from
which 247 unique genes were predicted. 29 of 110 *c2!c5*∩EBNA2 site-linked genes (26.3%) were
c2!c5 DEGs in the scRNA assay, as were 34 of 115 *c2!c5*∩EBNALP site-linked genes (29.6%)
and 20 of 125 *c2!c5*∩EBNA3C site-linked genes (16.0%). 20 site-linked genes were identified
from all three viral co-activator recipes (**Figure S42B**), including the EBVSE-linked G protein
coupled receptor *GPR137B*, a lysosmal transmembrane receptor that regulates mTORC1 activity
and autophagy ([Gan et al., 2019](#); [Gao et al., 2012](#)). *GPR137B* was also identified as a *c2!c38*
DAP-linked DEG, indicating inaccessible regulatory loci within resting cells as well. We identified
two regulatory DAPs with significant positive correlation to gene expression at +14kb and +18kb
relative to the *GPR137B* TSS. One of these sites (+14kb) coincided with EBNALP and EBNA3C
binding sites but did not exhibit a *c2!c5* DAP. However, the second site (+18kb) overlapped with
all three intersected EBNAs and was a *c2!c5* DAP. Both sites also intersected with Rel family TFs
(cRel, RelA, and RelB) (**Figure S42C**). Other genes involved in lysosome-mediated processes
including autophagy and antigen presentation regulation were also identified from the *c2!c5*
comparison (*TFEB*, *LAMP3*), albeit with modestly elevated but significant differential expression
([Martina et al., 2012](#); [Nagelkerke et al., 2014](#); [Settembre et al., 2011](#)). We also found numerous

other genes involved in immune activation signaling, apoptosis, and transcriptional regulation that
displayed EBNA-associated *c2!c5* DAP-linked DEGs (**Figure S42D-E**). Collectively, these
vignettes of post-infection cell trajectories highlight the genome-wide dynamics of diverse EBV-
induced B cell responses captured within the single-cell multiomics dataset.

Discussion

Our data reveal heterogeneity in coordinated gene expression and chromatin accessibility dynamics within individual cells during the critical early stages of a viral infection. By capturing the initial phases of EBV infection with high resolution -omics, we discern the gene regulatory environments associated with diverse infected cell fates and their respective developmental trajectories. These include genome-wide expression and chromatin signatures associated with effective host antiviral response, virus-triggered oncogene-induced senescence, and the path to sustained EBV latency and host cell immortalization, which is accessed via simulated B cell activation. Moreover, we develop a bioinformatic workflow to characterize post-infection outcomes through gene- and peak-level investigations at loci matching specific epigenetic patterns as well as host and viral transcription factor binding profiles. The combined high resolution multiomics data and integrative analytical framework reported herein yield a vividly detailed representation of the genome-wide interplay of host and virus.

We captured diverse post-infection B cell fate trajectories that, due to their asynchronous parallel emergence, cannot be resolved by ensemble sequencing methods. Remarkably, we find that large-scale euchromatin-to-heterochromatin transitions (20-40% reductions in genome-wide accessibility) can occur in post-infection trajectories and fates, including several that display increases in transcript levels. The scATAC data implicate Simpson's paradox with respect to heterogeneous chromatin accessibility dynamics following infection, since the total number of unique peaks increases in early infected populations while the peaks per cell can, in fact, decrease (Simpson, 1951; Trapnell, 2015) (**Table S1**).

Global epigenetic silencing via SAHF formation (Di Micco et al., 2011) is most prominent both in cells that rapidly undergo innate sensing-mediated arrest and in cells that evade this response but arrest due to DNA damage response activation during virus-induced hyperproliferation. Our data also indicate a possible role for ribosome biogenesis in the transition from virus-induced arrest to senescence, likely through a p53-MDM2 axis (Deisenroth and Zhang, 2010), which critically regulates EBV transformation (Forte and Luftig, 2009). Intriguingly, the trajectory of successful infection is distinguished by increased accessibility at key sites against the larger backdrop of heterochromatin formation. A substantial number of these sites (many associated with viral TFs) have predicted *cis* linkages to genes enriched for regulation of apoptosis, tumor suppression, inflammation, and chromatin remodeling, all of which are determinants of successful EBV infection. Moreover, viral episome ATAC profile heterogeneity across infected phenotypes

indicates that latency establishment depends on retained accessibility to viral genes within the
repressive host milieu.

The identification of an EBV-infected analog of an AP-eMBC phenotype is consistent with
results from *in vitro* and *in vivo* antigen stimulation experiments (Suan et al., 2017; Taylor et al.,
2015), as well as previous work from our lab that identified CCR6 as an EBV Latency IIb program
biomarker that becomes downregulated in the transition to Latency III in LCLs (Messinger et al.,
2019).The development of this state *in vitro* implies that EBV may gain access to the memory
pool *in vivo* via progenitors that have limited involvement in the GC reaction. In the context of
normal antigen stimulation, this subset of eMBCs undergoes early exit from the cell cycle and GC
reaction as a consequence of restricted access to or engagement with cognate antigen (Glaros
et al., 2021). It is thus conceivable that EBV-infected B cells may differentially develop into GC
BCs, PBs, or eMBCs from the AP state in a manner dependent on the extent to which the LMPs,
EBNAs and other viral gene products induce mimicry of BCR activation and signaling. This model
accommodates a surprising possibility – that EBV may access long-term persistence and survival
not only within post-GC high-affinity MBCs but also via GC-independent eMBCs that avoid
extensive proliferation.

The activated B cell and plasmablast phenotypes that developed within 5 days are generally
consistent with our findings in LCLs (SoRelle et al., 2021) and resemble *in vivo* tonsil cell subset
transcriptomes. Furthermore, differentiated plasmablasts exhibited fewer accessible sites in both
host chromatin and viral episomes relative to activated cells. Notably, prior studies found that only
50% of EBV-infected cells that secrete immunoglobulin go on to become immortalized (Tosato et
al., 1985). Additional work within LCLs demonstrated that EBV$^+$ cells with upregulated Ig
production exhibited reduced DNA synthesis and EBNA downregulation (Wendel-Hansen et al.,
1987). Collectively, these findings support a model for continuous EBV-driven B cell differentiation
in vitro, wherein plasmablasts are continually generated through activation-induced maturation
yet selected against by their reduced proliferation. While this disadvantage limits viral replication
via cell division, the reduced MYC levels, increased XBP1, and endoplasmic reticulum stress in
these cells may support EBV lytic reactivation (Guo et al., 2020; Laichalk and Thorley-Lawson,
2005; Sun and Thorley-Lawson, 2007; Taylor et al., 2011).

Depending on the phenotype comparison, roughly 10-35% of all DEGs were correlated with
DAPs. This range was similar to the frequencies of both differential accessibility of expressed TF
binding motifs (23%) and DEGs associated with differentially accessible promoters (25%)
identified in dexamethasone-treated A549 cells profiled with sci-CAR (Cao et al., 2018). The
observation of similar DAP-linked DEG frequencies in response to diverse stimuli (e.g., chemically

induced glucocorticoid receptor activation, viral infection) raises intriguing questions regarding the
fundamental frequency of genes regulated, at least in part, by accessibility changes. The
observed proportion implies that most DEGs may be regulated by higher order chromatin
interactions and differential recruitment of transcriptional activators and/or RNA Pol II.

The crisp-ATAC method provides a simple, flexible informatic approach to map ChIP-seq
profiles to cell phenotypes discovered from scATAC-seq. Thus, it provides a potential route for
effectively bootstrapping scATAC resolution to datasets from other -omics modalities. When data
from suitable reference cell types and/or states are available, crisp-ATAC can be used to predict
phenotype-resolved gene regulation by evaluating simple or complex combinations of factor
binding sites, histone modifications, and/or nuclear chromatin compartmentalization across all
differentially accessible sites. In our case, cell-matched empirical scRNA-seq data can be cross-
referenced with crisp-ATAC outputs for methodological validation and identification of genes of
interest for future studies. This method is adaptable to comparisons of phenotype-resolved ATAC
profiles in contexts such as other infections, development, and cell responses to stimuli or
therapies. We expect this approach to be particularly powerful for exploring TF-associated
regulatory changes in time-resolved studies of cellular behaviors.

<u>Limitations of the study</u>

The reported single-cell multiomics data have several limitations. They do no capture aspects
of host-virus dynamics acting at other molecular levels. Examples include epigenetic
modifications (e.g., DNA methylation status), three-dimensional chromatin architecture changes,
modulation of translation and protein abundance, post-translational modifications, protein-protein
interactions, and signaling cascades (e.g., phosphorylation status). While we present a method
for inferring DAP-linked TF binding and epigenetic modifications based on empirical scATAC data,
we do not have direct ChIP-seq measurements at the single-cell level.

The reported bioinformatic methodology (crisp-ATAC) also has notable constraints. This
approach is limited by the availability of ChIP-seq data from an appropriate cell type and target
reference state. Moreover, regulatory site predictions must be empirically tested to validate
potential functions in gene expression control. Finally, distance limits imposed for identifying *cis-*
regulatory linkages preclude identification of distal gene regulatory elements formed via 3D
nuclear conformation.

Acknowledgments

The authors wish to thank members of the Duke University Molecular Genomics Core (MGC) and the Duke Center for Genomic and Computational Biology (GCB), especially Emily Hocke, Karen Abramson, Dr. Simon Gregory, and Dr. Nicolas Devos. Special thanks are also due to the anonymous donors, without whose blood donations this work would not have been possible. E.D.S. wishes to acknowledge funding support from the Department of Molecular Genetics and Microbiology Viral Oncology Training Grant (NIH T32 # 5T32CA009111-42). This work was supported by funding from the National Institute of Dental and Craniofacial Research (NIDCR award #R01DE025994-06).

Experimental Methods

PBMC isolation and B lymphocyte enrichment

Whole blood (50 mL each from two anonymous donors; TX1241/Donor 1 & TX1242/Donor 2) was obtained from the Gulf Coast Regional Blood Center (Houston, TX). Upon receipt, peripheral blood mononuclear cells (PBMCs) from each donor sample were isolated via Ficoll gradient separation (Histopaque®-1077, Sigma #H8889), resuspended at $10x10^6$ cells/mL in RPMI 1640 + 15% heat-inactivated fetal bovine serum (FBS, v/v, Corning) (R15 media), and incubated overnight at 37°C and 5% CO_2. The next day, $CD19^+$ B cells were enriched from donor PBMCs via negative isolation (BD iMag Negative Isolation Kit, BD Biosciences #558007) and resuspended at $2x10^6$ cells/mL in R15 supplemented with 2 mM L-glutamine, 100 units/mL penicillin, 100 µg/mL streptomycin, and 0.5 µg/mL cyclosporine A ($R15^+$ media). Roughly $45x10^6$ B cells were recovered per donor post-enrichment. Following $CD19^+$ validation (see below: Flow cytometry), enriched B cell aliquots (1-2 mL at $3x10^6$ cells/mL) were viably frozen in 90% FBS + 10% DMSO and stored in liquid N_2.

EBV infection and cell culture

The type 1 EBV strain B95-8 was obtained in-house as viral supernatant from the inducible B95-9 Z-HT cell line as reported previously (Johannsen et al., 2004). Immediately after withholding and cryopreserving uninfected enriched B cells for each donor (day 0 samples), the remaining cells in culture were infected with B95-8 via resuspension in viral supernatant (100 µL per $1x10^6$ cells) for 1 h at 37°C and 5% CO_2. Infected B cells from each donor were rinsed with 1x PBS, resuspended in $R15^+$ media, and incubated at the conditions described above throughout the course of infection. Aliquots were taken from each infected donor culture at 2-, 5-, and 8-days post-infection and viably frozen as described for uninfected day 0 samples.

Flow cytometry

The extent of B cell enrichment was quantified for each donor using flow cytometry. Following negative isolation, cell samples ($2x10^5$ per donor) were rinsed with FACS buffer (1x PBS + 2% FBS), stained with phycoerythrin (PE)-conjugated mouse anti-human CD19 (BioLegend, clone HIB19; catalog #302208; lot #B273508) in the dark for 30 min at room temperature, then rinsed again prior to analysis. Cell samples at each timepoint were prepared as described and co-stained with PE-anti-CD19 and allophycocyanin-conjugated mouse anti-human CD23 (APC-anti-CD23, BioLegend, clone EBVCS-5; catalog #338513; lot #B273489) to validate successful EBV

infection. To validate the AP-eMBC phenotype (c0), enriched resting B cells from two additional

donors (TX1253 and TX1254) were labeled with CellTrace Violet (ThermoFisher / Invitrogen, Cat

#34571) and stained with one of the following combinations at days 0, 2, 5, and 8: CCR6/Memory

panel (FITC-anti-CD27, PE-anti-CCR6, and APC-anti-CD23); Naïve/Memory panel (FITC-anti-

IgD, PE-anti-CD19, and APC-anti-CD27); or CCR6/Naïve panel (FITC-anti-IgD, PE-anti-CCR6,

and APC-anti-CD23). FITC-anti-CD27, FITC-anti-IgD, and APC-anti-CD27 were purchased from

BioLegend (Cat #302806, #348206, and #356410, respectively) and PE-anti-CCR6 was

purchased from Invitrogen (Cat #12-1969-42). Compensation matrices were calculated from

single-stain controls for FITC and PE and applied to all samples for analysis. All cytometry

measurements were acquired with a BD FACSCanto II (BD Biosciences) and analyzed using

FlowJo version 10 (Ashland / BD Biosciences).

<u>Human tonsil sample preparation</u>

Tonsillar B cells were isolated from discarded, anonymized tonsillectomies from the Duke

Biospecimen Repository and Processing Core (BRPC; Durham, NC). Tonsil tissue samples were

manually disaggregated, filtered through a cell strainer, and isolated by layering over a cushion

made from Histopaque-1077 (H8889; Sigma-Aldrich). Harvested lymphocytes were washed three

times with FACS buffer (5% FBS in PBS) prior to scRNA library preparation.

<u>Preparation of scRNA and scATAC libraries</u>

Cryopreserved samples from each early infection timepoint of interest were simultaneously

thawed by donor and purified to > 90% viable cells by Ficoll gradient separation. Viable cells from

each timepoint and donor were then prepared as single-cell matched gene expression (scRNA)

and chromatin accessibility (scATAC) libraries by the Duke Molecular Genomics Core staff with

the 10x Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Kit (10x Genomics,

Pleasanton, CA) (Satpathy et al., 2019; Zheng et al., 2017). Briefly, nuclei were isolated from

each sample and subjected to transposition at accessible chromatin sites. Next, transposed

nuclei, barcoding master mix, and gel beads containing unique barcode sequences were

prepared into single-cell GEMs (Gel bead emulsions) using the Chromium Controller and Chip J.

Within each GEM, poly-adenylated (poly-A) mRNA transcripts from individual nuclei are captured

by barcoded, indexed poly-T primers and reverse transcribed into cDNA. Simultaneously, a

separate barcoded sequence containing a spacer and Illumina P5 adaptor sequence is added to

transposed regions within the captured nucleus. Barcoded multiomes were then purified, pooled,

and pre-amplified by PCR prior to library construction. The scATAC library for each sample is

generated by PCR amplification and incorporation of sample index and Illumina P7 adaptor
sequences. Separately, pre-amplified gene expression cDNA is further PCR amplified,
fragmented, and size selected. The scRNA library for each sample is then constructed using PCR
to incorporate the P5 and P7 sequencing adaptors. Two biological replicates of tonsillar
lymphocytes were prepared as scRNA libraries using the 10x Genomics Next GEM 3' Gene
Expression kit with v3 chemistry (10x Genomics, Pleasanton, CA), and sequenced, processed,
aligned, and analyzed as described above for early infection scRNA samples.
667
Sequencing, alignment, and count matrix generation
The 8 paired-end scATAC libraries (4 timepoints per 2 donors) were pooled onto two lanes of
an Illumina S2 flow cell and sequenced at a target depth of 25,000 reads per cell on an Illumina
NovaSeq (Illumina, San Diego, CA). The 8 paired-end scRNA libraries were similarly pooled and
sequenced at a target depth of 50,000 reads per cell. Tonsil scRNA libraries were likewise pooled
and sequenced at 50,000 reads per cell. All sequencing runs were performed by staff at the Duke
Center for Genomic and Computational Biology. Raw base calls for each assay were prepared
as sample-demultiplexed FASTQ files using *cellranger-arc mkfastq* (Cellranger, 10x Genomics),
a wrapper of the Illumina *bcl2fastq* function. Next, sample-matched scRNA and scATAC reads
were aligned against genomic references to produce multiome count matrices using *cellranger-*
arc count. One set of count matrices was generated by mapping reads to a concatenated genomic
reference constructed from the human genome (GRCh38) with the ~172 kB type 1 EBV genome
(NC_007605) included as an extra chromosome. These outputs were used for downstream RNA-
only analyses to capture host and viral gene expression. A second set of count matrices
generated by mapping to GRCh38 only was used for downstream joint RNA and ATAC analyses.
Compatible reference packages were assembled from the relevant genome (.fa) and annotation
(.gtf) files using *cellranger-arc mkref*.
685
Data QC and scMultiome analysis
All direct analysis of scRNA and scATAC data was conducted in R using Signac (Stuart et al.,
2021), an extension of Seurat (Macosko et al., 2015; Satija et al., 2015; Stuart et al., 2019).
Following read mapping and counting, linked scRNA and scATAC data were obtained from
between 8,934-20,000 cells per sample. After QC filtering by mitochondrial expression (n < 20%),
total transcripts (n < 25,000), unique transcripts (n > 1,000), and minimum cells expressing a
given feature (n > 3), data from between 8,376-19,310 cells per sample were analyzed (see **Table**
S1). The mitochondrial gene expression threshold was selected based on the high metabolic

activity of early-infected B cells and the high cell viability observed in each sample (> 90%)
immediately prior to library preparation to preserve biologically relevant phenotypes (Osorio and
Cai, 2021). After QC filtering, a total of 52,271 and 44,920 cells were analyzed across the infection
timecourse for donors TX1241 and TX1242, respectively. For RNA-only analysis, gene
expression data (host and viral) across all timepoints for a given donor were merged into a single
object, log normalized, scored for cell cycle markers, and scaled with cell cycle scoring regressed
out. The top 2,000 differentially expressed features over the early infection timecourse data were
identified and used for principal component analysis (PCA). The top 30 principal components
were further dimensionally reduced via uniform manifold approximation projection (UMAP,
(McInnes et al., 2018)), and clustering was performed to identify biologically distinct cell
subpopulations. Merged scRNA dataset pseudotime trajectories were calculated using Monocle3
(Qiu et al., 2017), and were mapped along with cluster identities to 3D UMAP coordinates for
visualization (Qadir, 2019; Qadir et al., 2020). For joint ATAC and RNA analysis, host gene
expression and chromatin accessibility were analyzed for each separate timepoint and for a
merged object containing Day 0 and Day 8 multiome data. Nucleosome signal and transcription
start site (TSS) enrichment were calculated and used for QC filtering (Nucleosome.signal < 2,
TSS.enrichment > 1). ATAC peaks were called using macs2 (Liu, 2014) with hg38 annotations.
Gene expression data in the joint analysis was processed as described for the RNA-only analysis
with the exception of using Signac's SCTransform function instead of log normalization for
expression counts. Top differential features in each assay ('peaks' and 'SCT') were determined,
and multimodal neighboring and UMAP were performed for integrated data visualization. Cluster
identities defined in the RNA-only assay were mapped to this merged joint dataset, which
contained cells representing all identified subpopulations. Peaks with significant (anti-) correlation
($p < 0.05$ for z-score of correlation coefficients) to differentially expressed genes were identified
using the LinkPeaks function in Signac, which was informed by SHARE-seq (Ma et al., 2020).

crispATAC workflow and reference data curation

ChIP-referenced inference from single-cell phenotype ATAC (crispATAC) was developed to
predict subpopulation-resolved gene regulatory features. In a typical workflow, cluster-level
chromatin accessibility tracks are cross-referenced against ChIP-Seq (Chromatin
Immunoprecipitation Sequencing) profiles for epigenetic marks and TFs of interest measured from
a reference cell phenotype (in this study, lymphoblastoid cell lines such as GM12878). In this
study, cluster-specific called peaks from the joint scATAC + scRNA dataset were extracted and
prepared as simplified genomic range files (3-column .bed file format). Next, the desired ChIP-

Seq datasets for the reference phenotype were downloaded and, where applicable, converted to

.bedgraph format to be used as input for peak calling with the *macs2* function *bdgpeakcall*. The

ChIP (and Hi-C) datasets used for crispATAC in this study (**Table S2**) are all publicly available

from the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO).

Once all ATAC-seq and ChIP-seq peak files were generated, all were used as inputs to a single

call of the bedtools (Quinlan and Hall, 2010) function *multiinter*, which output a matrix of all

genomic range intersection intervals where at least one input file exhibited a peak. This

intersection matrix was imported to R as a data frame and analyzed to identify common and/or

differential intervals (matrix rows) among scATAC cluster phenotypes, epigenetic marks, and TFs

using Boolean logic gating by dataset (matrix columns). For a given crispATAC recipe (e.g., peaks

in scATAC cluster 1 not in scATAC cluster 2 intersected with EBNA2 ChIP peaks = [c1 ∩ !c2] ∩

EBNA2), the genome intervals matching the gating criteria were returned and converted to .bed

files. Lists of differentially accessible, transcription-factor associated sites generated in this way

were subsequently analyzed with the Genomic Regions Enrichment of Annotations Tool (GREAT)

(McLean et al., 2010) to identify potential *cis*-regulated genes within 1 megabase of each query

site. As a final step, output lists of potential linked genes were intersected with the top marker

genes identified from the corresponding cluster-wise comparison in the scRNA assay, thus

integrating direct single-cell RNA and ATAC measurements with subpopulation-resolved

regulatory inferences from ensemble ChIP profiles. In a similar but separate approach, scATAC

and ChIP peaks were intersected with topologically associated domain (TAD) boundaries

(prepared using hicExplorer, (Ramírez et al., 2018; Wolff et al., 2018; Wolff et al., 2020)) and

nuclear subcompartments from GM12878 Hi-C data to study differentially accessible TF-

associated sites in the context of 3D nuclear architecture.

<u>Visualization of crispATAC outputs, gene ontologies, and networks</u>

Data for genes of interest identified from crispATAC recipes were explored using dimensionally

reduced (UMAP) expression maps and cluster-level accessibility tracks (Signac, (Stuart et al.,

2021)), called peaks aligned with TFs and epigenetic marks (IGV, (Robinson et al., 2011)), and

local neighborhoods in Hi-C contact maps (Juicebox, (Durand et al., 2016)). Cluster-resolved

gene ontologies were generated and quantified by GREAT (McLean et al., 2010). Top scRNA

assay cluster markers and GREAT output gene lists were also visualized as annotated networks

using Cytoscape (Shannon et al., 2003).

Main Figures

Figure 1. Time-resolved single-cell gene expression during early EBV infection of B lymphocytes

(A) Single-cell gene expression timecourse data from early EBV infection.

(B) General expression trends during early infection. Total mRNA refers to all transcripts captured, while Feature mRNA refers to the number of unique transcripts (per cell).

(C) Cell phase scoring of expression data after cell cycle marker regression.

(D) Unsupervised clustering of early infected cell expression in merged timepoint data.

(E) Pairwise correlation of identified clusters.

(F) Cluster membership by timepoint. Fit lines show coarse changes in phenotype frequency over time.

(G) Single-cell expression of the top 15 gene markers by cluster.

See also Figures S1-S4

Figure 2. High-resolution dissection of infected B cell phenotypes

(A) Overview of global gene expression trends by phenotype.

(B) Induction of interferon response genes in all EBV+ clusters.

(C) Phenotype-resolved transcriptomic signatures in resting and EBV+ B cells. Select cluster-resolved comparisons of gene expression were evaluated via the Kolmogorov-Smirnov D statistic (KS D) and associated p value (* p < 1e-5; ** p < 1e-10; *** p < 1e-15) from 500 randomly sampled cells per cluster.

See also Figures S5-S11

789
790
Figure 3. A model of B cell fate trajectories in early EBV infection
(A) Monocle3 pseudotime scoring of merged timecourse expression data relative to resting B cells
(day 0). Unlike the 2D UMAP, 3D UMAPs depict closer proximity of c6 (first observed at day 2,
blue dashed circle) to resting cells, consistent with the temporal emergence of the c6 phenotype
prior to the c0, c1, c2, and c5 phenotypes.
(B) Pseudotime-resolved expression dynamics of top differentially expressed genes (DEGs)
across phenotypes. Genes are hierarchically clustered by pseudo-temporal expression pattern
similarity. Spline interpolant fits are shown for expression of select genes in pseudotime (insets i-
iv). After sorting cells by pseudotime score in ascending order, the average pseudotime score of
every 25-cell interval was computed for efficient visualization (i.e., pseudotime for 52,271 cells at
25 cell resolution).
(C) Annotated state model of EBV+ B cell phenotypes and fate trajectories. Empirically observed
and putative directed state transitions are depicted in solid and dashed edges, respectively.
Edges drawn to groups of phenotypes (dotted ovals) indicate transitions to/from each cluster
within the group.
See also Figures S12-S19
807

808
809
Figure 4. A subset of early infected cells exhibits hallmarks of a multipotent activated
precursor to early memory B cells (eMBCs)
(A) Top phenotype markers of healthy human tonsil subsets identified from scRNA-seq.
(B) UMAP merging of tonsil and key early infection cluster scRNA-seq assays. Tonsil clusters are
colored to match the closest corresponding cells from early infection.
(C) UMAP Correspondence of key gene expression across tonsillar subsets and early infection
phenotypes. Select markers of multipotent progenitors and eMBCs were informed by data from
(Suan et al., 2017) and (Glaros et al., 2021).
(D) Dot plot visualization of key genes across early infection (ei) c0, c2, c5, and c6 and their
analogs within tonsils (t).
820

821
822
Figure 5. FACS validation of CCR6$^+$ AP-eMBCs derived from naïve and memory B cells.
(A) Relative expression of *CCR6* and *FCER2/CD23* by model phenotype determined by scRNA-seq.
(B) CCR6 surface expression on uninfected and EBV-infected B cells by number of cell divisions (CellTrace Violet). CCR6^{hi} cells (blue gate) exhibits reduced proliferation relative to CCR6^{lo} (magenta gate) cells.
(C) CCR6 and CD23 (FCER2) surface expression over the early infection timecourse.
(D) Cell divisions and IgD status of cells at Day 8 gated by CCR6 and CD23 expression. Gated fractions are colored by approximate correspondence to scRNA-seq model phenotypes.
(E) A fate model for EBV-induced AP-eMBC-like cells. $P_{0\rightarrow...n}$ signifies the probability of the eventual transition of a cell from cluster 0 (AP-eMBC analog) to a given cluster n. Relative probability relationships for naïve and memory cells are proposed based on empirical findings from FACS.
See also Figures S22-25
837

838
839
Figure 6. Cell-matched expression and chromatin accessibility cell fate trajectories
(A) UMAP visualization of scRNA + scATAC data generated using weighted nearest neighbors
(WNN) multimodal integration (Hao et al., 2021; Stuart et al., 2020). Merged multimodal data from
the first and last timepoints (day 0 & day 8) contain cells representative of all identified
phenotypes.
(B) Distribution of called ATAC peaks per cell by timepoint and phenotype in both donors (***p <
1e-15, one-sided Kolmogorov-Smirnov test).
(C) Overview of global differentially accessible peak (DAP) -linked differentially expressed genes
(DEGs) and cluster comparisons for major trajectories of interest. DAPs are identified by their
presence in one or more peaks but not in (!) one or more other peaks. DAP-linked DEGs were
explored in resting versus arrested cells (c38 vs c47), EBV[+] activated versus resting cells (c2 vs
c38), and EBV[+] activated vs EBV+ differentiated cells (c2 vs c5).
(D) Virus-induced arrest responses. State-resolved DAPs (*c38!c47*) and joint assay trend
summaries are presented in addition to example DEGs and DAP linkages resolved by phenotype.
(E) Successful infection trajectory. State-resolved DAPs (*c2!c38*), joint assay summaries, and
trajectory-specific examples are presented as in (D).
(F) EBV-induced B cell activation/differentiation continuum. State-resolved DAPs (*c2!c5*), joint
assay summaries, and trajectory-specific examples are presented as in (D and E).
See also Figures S26-S38
859

860
861
Figure 7. crisp-ATAC analysis of DAP-linked DEGs in activated versus innate arrested
EBV⁺ B cells
(A) Schematic of NF-κB activation (c2) and innate arrest (c4) model phenotypes.
(B) Multimodal assay gating to extract *c2!c4* DAPs.
(C and D) Prediction of *cis*-regulatory linkage. All *c2!c4* peak intervals (n=1,873) are used as
inputs to the GREAT (McLean et al., 2010) to predict *c2!c4* DAP-linked genes (n=1,514).
(E) Occurrence of EBVSE-linked genes identified as *c2!c4* DAP-linked DEGs associated with
select EBNA binding sites relative to expected frequency due to random overlap (n=100
simulation trials, error bars depict mean +/- standard deviation). Random samples were size-
matched relative to each EBNA-associated gene list.
(F) Example gene of interest analysis for *TRAF1*, an EBVSE-linked gene identified as a *c2!c4*
DAP-linked DEG with phenotype-variable accessibility at multiple EBNA binding sites.
See also Figures S39-S46

References

References

Alfieri, C., Birkenbach, M., and Kieff, E. (1991). Early events in Epstein-Barr virus infection of human B
lymphocytes. Virology *181*, 595-608.

Allday, M.J., Bazot, Q., and White, R.E. (2015). The EBNA3 family: two oncoproteins and a tumour
suppressor that are central to the biology of EBV in B cells. Epstein Barr Virus Volume 2, 61-117.

Altmann, M., Pich, D., Ruiss, R., Wang, J., Sugden, B., and Hammerschmidt, W. (2006). Transcriptional
activation by EBV nuclear antigen 1 is essential for the expression of EBV's transforming genes.
Proceedings of the National Academy of Sciences *103*, 14188-14193.

Anderson, L.J., and Longnecker, R. (2008). EBV LMP2A provides a surrogate pre-B cell receptor signal
through constitutive activation of the ERK/MAPK pathway. The Journal of general virology *89*, 1563.

Arvey, A., Tempera, I., Tsai, K., Chen, H.-S., Tikhmyanova, N., Klichinsky, M., Leslie, C., and Lieberman,
P.M. (2012). An atlas of the Epstein-Barr virus transcriptome and epigenome reveals host-virus regulatory
interactions. Cell host & microbe *12*, 233-245.

Babcock, G.J.D., L.L.; Volk, M.; Thorley-Lawson, D.A. (1998). EBV Persistence in Memory B Cells In
Vivo. Immunity *9*, 395-404.

Banerjee, S., Lu, J., Cai, Q., Sun, Z., Jha, H.C., and Robertson, E.S. (2014). EBNA3C augments Pim-1
mediated phosphorylation and degradation of p21 to promote B-cell proliferation. PLoS Pathog *10*,
e1004304.

Bartkova, J., Rezaei, N., Liontos, M., Karakaidos, P., Kletsas, D., Issaeva, N., Vassiliou, L.-V.F., Kolettas,
E., Niforou, K., and Zoumpourlis, V.C. (2006). Oncogene-induced senescence is part of the
tumorigenesis barrier imposed by DNA damage checkpoints. Nature *444*, 633-637.

Bird, A.G.M., S.M.; Britton, S. (1981). Cyclosporin A promotes spontaneous outgrowth in vitro of Epstein-
Barr virus-induced B-cell lines. Nature *289*, 300-301.

Bjornevik, K., Cortese, M., Healy, B.C., Kuhle, J., Mina, M.J., Leng, Y., Elledge, S.J., Niebuhr, D.W.,
Scher, A.I., and Munger, K.L. (2022). Longitudinal analysis reveals high prevalence of Epstein-Barr virus
associated with multiple sclerosis. Science.

Buenrostro, J.D., Wu, B., Litzenburger, U.M., Ruff, D., Gonzales, M.L., Snyder, M.P., Chang, H.Y., and
Greenleaf, W.J. (2015). Single-cell chromatin accessibility reveals principles of regulatory variation.
Nature *523*, 486-490.

Cahir-McFarland, E.D., Carter, K., Rosenwald, A., Giltnane, J.M., Henrickson, S.E., Staudt, L.M., and
Kieff, E. (2004). Role of NF-κB in cell survival and transcription of latent membrane protein 1-expressing
or Epstein-Barr virus latency III-infected cells. Journal of virology *78*, 4108-4119.

Calender, A., Billaud, M., Aubry, J.-P., Banchereau, J., Vuillaume, M., and Lenoir, G.M. (1987). Epstein-
Barr virus (EBV) induces expression of B-cell activation markers on in vitro infection of EBV-negative B-
lymphoma cells. Proceedings of the National Academy of Sciences *84*, 8060-8064.

Canaan, A., Haviv, I., Urban, A.E., Schulz, V.P., Hartman, S., Zhang, Z., Palejev, D., Deisseroth, A.B.,
Lacy, J., and Snyder, M. (2009). EBNA1 regulates cellular gene expression by binding cellular promoters.
Proceedings of the National Academy of Sciences *106*, 22421-22426.

Cao, J., Cusanovich, D.A., Ramani, V., Aghamirzaie, D., Pliner, H.A., Hill, A.J., Daza, R.M., McFaline-
Figueroa, J.L., Packer, J.S., and Christiansen, L. (2018). Joint profiling of chromatin accessibility and
gene expression in thousands of single cells. Science *361*, 1380-1385.

Chen, S., Lake, B.B., and Zhang, K. (2019). High-throughput sequencing of the transcriptome and
chromatin accessibility in the same cell. Nature biotechnology *37*, 1452-1457.

Clark, S.J., Argelaguet, R., Kapourani, C.-A., Stubbs, T.M., Lee, H.J., Alda-Catalinas, C., Krueger, F.,
Sanguinetti, G., Kelsey, G., and Marioni, J.C. (2018). scNMT-seq enables joint profiling of chromatin
accessibility DNA methylation and transcription in single cells. Nature communications *9*, 1-9.

Cohen, J.I., Fauci, A.S., Varmus, H., and Nabel, G.J. (2011). Epstein-Barr virus: an important vaccine
target for cancer prevention. Science translational medicine *3*, 107fs107-107fs107.

Cohen, J.I., Wang, F., Mannick, J., and Kieff, E. (1989). Epstein-Barr virus nuclear protein 2 is a key
determinant of lymphocyte transformation. Proceedings of the National Academy of Sciences *86*, 9558-
9562.

Courtois-Cox, S., Jones, S.L., and Cichowski, K. (2008). Many roads lead to oncogene-induced
senescence. Oncogene *27*, 2801-2809.

Deisenroth, C., and Zhang, Y. (2010). Ribosome biogenesis surveillance: probing the ribosomal protein-
Mdm2-p53 pathway. Oncogene *29*, 4253-4260.

Devergne, O., Hatzivassiliou, E., Izumi, K.M., Kaye, K.M., Kleijnen, M.F., Kieff, E., and Mosialos, G.
(1996). Association of TRAF1, TRAF2, and TRAF3 with an Epstein-Barr virus LMP1 domain important for
B-lymphocyte transformation: role in NF-kappaB activation. Molecular and cellular biology *16*, 7098-7108.

Devergne, O., McFarland, E.C., Mosialos, G., Izumi, K.M., Ware, C.F., and Kieff, E. (1998). Role of the
TRAF binding site and NF-κB activation in Epstein-Barr virus latent membrane protein 1-induced cell
gene expression. Journal of virology *72*, 7900-7908.

Dheekollu, J., Wiedmer, A., Ayyanathan, K., Deakyne, J.S., Messick, T.E., and Lieberman, P.M. (2021).
Cell-cycle-dependent EBNA1-DNA crosslinking promotes replication termination at oriP and viral episome
maintenance. Cell *184*, 643-654. e613.

Di Micco, R., Sulli, G., Dobreva, M., Liontos, M., Botrugno, O.A., Gargiulo, G., Dal Zuffo, R., Matti, V.,
d'Ario, G., and Montani, E. (2011). Interplay between oncogene-induced DNA damage response and
heterochromatin in senescence and cancer. Nature cell biology *13*, 292-302.

Durand, N.C., Robinson, J.T., Shamim, M.S., Machol, I., Mesirov, J.P., Lander, E.S., and Aiden, E.L.
(2016). Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell systems
3, 99-101.

Efremova, M., and Teichmann, S.A. (2020). Computational methods for single-cell omics across
modalities. Nature methods *17*, 14-17.

Eliopoulos, A.G., Waites, E.R., Blake, S.M., Davies, C., Murray, P., and Young, L.S. (2003). TRAF1 is a
critical regulator of JNK signaling by the TRAF-binding domain of the Epstein-Barr virus-encoded latent
infection membrane protein 1 but not CD40. Journal of virology *77*, 1316-1328.

Epstein, M., Achong, B., and Barr, Y. (1964). Virus particles in cultured lymphoblasts from Burkitt's
lymphoma. Lancet, 702-703.

Fish, K., Comoglio, F., Shaffer, A.L., Ji, Y., Pan, K.-T., Scheich, S., Oellerich, A., Doebele, C., Ikeda, M.,
and Schaller, S.J. (2020). Rewiring of B cell receptor signaling by Epstein–Barr virus LMP2A.
Proceedings of the National Academy of Sciences *117*, 26318-26327.

Forte, E., and Luftig, M.A. (2009). MDM2-dependent inhibition of p53 is required for Epstein-Barr virus B-
cell growth transformation and infected-cell survival. Journal of virology *83*, 2491-2499.

Fries, K.L., Miller, W.E., and Raab-Traub, N. (1999). The A20 protein interacts with the Epstein–Barr virus
latent membrane protein 1 (LMP1) and alters the LMP1/TRAF1/TRADD complex. Virology *264*, 159-166.

Gan, L., Seki, A., Shen, K., Iyer, H., Han, K., Hayer, A., Wollman, R., Ge, X., Lin, J.R., and Dey, G.
(2019). The lysosomal GPCR-like protein GPR137B regulates Rag and mTORC1 localization and activity.
Nature cell biology *21*, 614-626.

Gao, J., Xia, L., Lu, M., Zhang, B., Chen, Y., Xu, R., and Wang, L. (2012). TM7SF1 (GPR137B): a novel
lysosome integral membrane protein. Molecular biology reports *39*, 8883-8889.

Glaros, V., Rauschmeier, R., Artemov, A.V., Reinhardt, A., Ols, S., Emmanouilidi, A., Gustafsson, C.,
You, Y., Mirabello, C., and Björklund, Å.K. (2021). Limited access to antigen drives generation of early B
cell memory while restraining the plasmablast response. Immunity *54*, 2005-2023. e2010.

Glück, S., Guey, B., Gulen, M.F., Wolter, K., Kang, T.-W., Schmacke, N.A., Bridgeman, A., Rehwinkel, J.,
Zender, L., and Ablasser, A. (2017). Innate immune sensing of cytosolic chromatin fragments through
cGAS promotes senescence. Nature cell biology *19*, 1061-1070.

Greenfeld, H., Takasaki, K., Walsh, M.J., Ersing, I., Bernhardt, K., Ma, Y., Fu, B., Ashbaugh, C.W., Cabo,
J., and Mollo, S.B. (2015). TRAF1 coordinates polyubiquitin signaling to enhance Epstein-Barr virus
LMP1-mediated growth and survival pathway activation. PLoS pathogens *11*, e1004890.

Guasparri, I., Bubman, D., and Cesarman, E. (2008). EBV LMP2A affects LMP1-mediated NF-κB
signaling and survival of lymphoma cells by regulating TRAF2 expression. Blood, The Journal of the
American Society of Hematology *111*, 3813-3820.

Guo, R., Jiang, C., Zhang, Y., Govande, A., Trudeau, S.J., Chen, F., Fry, C.J., Puri, R., Wolinsky, E., and
Schineller, M. (2020). MYC controls the Epstein-Barr virus lytic switch. Molecular cell *78*, 653-669. e658.

Han, I., Harada, S., Weaver, D., Xue, Y., Lane, W., Orstavik, S., Skalhegg, B., and Kieff, E. (2001).
EBNA-LP associates with cellular proteins including DNA-PK and HA95. Journal of virology *75*, 2475-
2481.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck III, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., and Zager, M. (2021). Integrated analysis of multimodal single-cell data. Cell.

Harada, S., and Kieff, E. (1997). Epstein-Barr virus nuclear protein LP stimulates EBNA-2 acidic domain-
mediated transcriptional activation. Journal of virology *71*, 6611-6618.

Henderson, S., Rowe, M., Gregory, C., Croom-Carter, D., Wang, F., Longnecker, R., Kieff, E., and
Rickinson, A. (1991). Induction of bcl-2 expression by Epstein-Barr virus latent membrane protein 1
protects infected B cells from programmed cell death. Cell *65*, 1107-1115.

Henle, W., Diehl, V., Kohn, G., Zur Hausen, H., and Henle, G. (1967). Herpes-type virus and
chromosome marker in normal leukocytes after growth with irradiated Burkitt cells. Science *157*, 1064-
1065.

Hodgkin, P.D., Lee, J.-H., and Lyons, A.B. (1996). B cell differentiation and isotype switching is related to
division cycle number. Journal of Experimental Medicine *184*, 277-281.

Hoffmann, A., Levchenko, A., Scott, M.L., and Baltimore, D. (2002). The IκB-NF-κB signaling module:
temporal control and selective gene activation. science *298*, 1241-1245.

Humme, S., Reisbach, G., Feederle, R., Delecluse, H.-J., Bousset, K., Hammerschmidt, W., and
Schepers, A. (2003). The EBV nuclear antigen 1 (EBNA1) enhances B cell immortalization several
thousandfold. Proceedings of the National Academy of Sciences *100*, 10989-10994.

Jiang, S., and Mortazavi, A. (2018). Integrating ChIP-seq with other functional genomics data. Briefings in
functional genomics *17*, 104-115.

Jiang, S., Zhou, H., Liang, J., Gerdt, C., Wang, C., Ke, L., Schmidt, S.C., Narita, Y., Ma, Y., and Wang, S.
(2017). The Epstein-Barr virus regulome in lymphoblastoid cells. Cell host & microbe *22*, 561-573. e564.

Johannsen, E., Luftig, M., Chase, M.R., Weicksel, S., Cahir-McFarland, E., Illanes, D., Sarracino, D., and
Kieff, E. (2004). Proteins of purified Epstein-Barr virus. Proceedings of the National Academy of Sciences
101, 16286-16291.

Junker, J.P., and van Oudenaarden, A. (2014). Every cell is special: genome-wide studies add a new
dimension to single-cell biology. Cell *157*, 8-11.

Kilger, E., Kieser, A., Baumann, M., and Hammerschmidt, W. (1998). Epstein–Barr virus-mediated B-cell
proliferation is dependent upon latent membrane protein 1, which simulates an activated CD40 receptor.
The EMBO journal *17*, 1700-1709.

Kim, U., Qin, X.-F., Gong, S., Stevens, S., Luo, Y., Nussenzweig, M., and Roeder, R.G. (1996). The B-
cell-specific transcription coactivator OCA-B/OBF-1/Bob-1 is essential for normal production of
immunoglobulin isotypes. Nature *383*, 542-547.

King, H.W., Orban, N., Riches, J.C., Clear, A.J., Warnes, G., Teichmann, S.A., and James, L.K. (2021).
Single-cell analysis of human B cell maturation predicts how antibody class switching shapes selection
dynamics. Science Immunology *6*, eabe6291.

Krynetskaia, N.F., Phadke, M.S., Jadhav, S.H., and Krynetskiy, E.Y. (2009). Chromatin-associated
proteins HMGB1/2 and PDIA3 trigger cellular response to chemotherapy-induced DNA damage.
Molecular cancer therapeutics *8*, 864-872.

Kwak, I.H., Kim, H.S., Choi, O.R., Ryu, M.S., and Lim, I.K. (2004). Nuclear accumulation of globular actin
as a cellular senescence marker. Cancer research *64*, 572-580.

Laichalk, L.L., and Thorley-Lawson, D.A. (2005). Terminal differentiation into plasma cells initiates the
replicative cycle of Epstein-Barr virus in vivo. J Virol *79*, 1296-1307.

Lamontagne, R.J., Soldan, S.S., Su, C., Wiedmer, A., Won, K.J., Lu, F., Goldman, A.R., Wickramasinghe,
J., Tang, H.-Y., Speicher, D.W.*, et al.* (2021). A multi-omics approach to Epstein-Barr virus
immortalization of B-cells reveals EBNA1 chromatin pioneering activities targeting nucleotide metabolism.
PLOS Pathogens *17*, e1009208.

Lanz, T.V., Brewer, R.C., Ho, P.P., Moon, J.-S., Jude, K.M., Fernandez, D., Fernandes, R.A., Gomez,
A.M., Nadj, G.-S., Bartley, C.M.*, et al.* (2022). Clonally Expanded B Cells in Multiple Sclerosis Bind EBV
EBNA1 and GlialCAM. Nature.

Lenain, C., de Graaf, C.A., Pagie, L., Visser, N.L., de Haas, M., de Vries, S.S., Peric-Hupkes, D., van
Steensel, B., and Peeper, D.S. (2017). Massive reshaping of genome–nuclear lamina interactions during
oncogene-induced senescence. Genome research *27*, 1634-1644.

Lessard, F., Igelmann, S., Trahan, C., Huot, G., Saint-Germain, E., Mignacca, L., Del Toro, N., Lopes-
Paciencia, S., Le Calvé, B., and Montero, M. (2018). Senescence-associated ribosome biogenesis
defects contributes to cell cycle arrest through the Rb pathway. Nature cell biology *20*, 789-799.

Lindahl, T., Adams, A., Bjursell, G., Bornkamm, G.W., Kaschka-Dierich, C., and Jehn, U. (1976).
Covalently closed circular duplex DNA of Epstein-Barr virus in a human lymphoid cell line. Journal of
molecular biology *102*, 511-530.

Ling, P.D., Peng, R.S., Nakajima, A., Yu, J.H., Tan, J., Moses, S.M., Yang, W.H., Zhao, B., Kieff, E., and
Bloch, K.D. (2005). Mediation of Epstein–Barr virus EBNA-LP transcriptional coactivation by Sp100. The
EMBO journal *24*, 3565-3575.

Liu, T. (2014). Use model-based analysis of ChIP-Seq (MACS) to analyze short reads generated by
sequencing protein–DNA interactions in embryonic stem cells. In Stem Cell Transcriptional Networks
(Springer), pp. 81-95.

Longnecker, R.M., Kieff, E., and Cohen, J.I. (2013). Epstein-barr virus. In Fields Virology: Sixth Edition
(Wolters Kluwer Health Adis (ESP)).

Lu, F., Chen, H.-S., Kossenkov, A.V., DeWispeleare, K., Won, K.-J., and Lieberman, P.M. (2016). EBNA2
drives formation of new chromosome binding sites and target genes for B-cell master regulatory
transcription factors RBP-jκ and EBF1. PLoS pathogens *12*, e1005339.

Lu, F., Wikramasinghe, P., Norseen, J., Tsai, K., Wang, P., Showe, L., Davuluri, R.V., and Lieberman,
P.M. (2010). Genome-wide analysis of host-chromosome binding sites for Epstein-Barr Virus Nuclear
Antigen 1 (EBNA1). Virology journal *7*, 1-17.

Luftig, M., Prinarakis, E., Yasui, T., Tsichritzis, T., Cahir-McFarland, E., Inoue, J.-I., Nakano, H., Mak,
T.W., Yeh, W.-C., and Li, X. (2003). Epstein–Barr virus latent membrane protein 1 activation of NF-κB
through IRAK1 and TRAF6. Proceedings of the National Academy of Sciences *100*, 15595-15600.

Lünemann, A., Rowe, M., and Nadal, D. (2015). Innate immune recognition of EBV. Epstein Barr Virus
Volume 2, 265-287.

Luo, Y., and Roeder, R.G. (1995). Cloning, functional characterization, and mechanism of action of the B-
cell-specific transcriptional coactivator OCA-B. Molecular and cellular biology *15*, 4115-4124.

Lupton, S., and Levine, A.J. (1985). Mapping genetic elements of Epstein-Barr virus that facilitate
extrachromosomal persistence of Epstein-Barr virus-derived plasmids in human cells. Molecular and
Cellular Biology *5*, 2533-2542.

Ma, S., Zhang, B., LaFave, L.M., Earl, A.S., Chiang, Z., Hu, Y., Ding, J., Brack, A., Kartha, V.K., and Tay,
T. (2020). Chromatin potential identified by shared single-cell profiling of RNA and chromatin. Cell *183*,
1103-1116. e1120.

Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A.R.,
Kamitaki, N., and Martersteck, E.M. (2015). Highly parallel genome-wide expression profiling of individual
cells using nanoliter droplets. Cell *161*, 1202-1214.

Mannick, J., Cohen, J., Birkenbach, M., Marchini, A., and Kieff, E. (1991). The Epstein-Barr virus nuclear
protein encoded by the leader of the EBNA RNAs is important in B-lymphocyte transformation. Journal of
virology *65*, 6826-6837.

Martin, H.J., Lee, J.M., Walls, D., and Hayward, S.D. (2007). Manipulation of the toll-like receptor 7
signaling pathway by Epstein-Barr virus. Journal of virology *81*, 9748-9758.

Martina, J.A., Chen, Y., Gucek, M., and Puertollano, R. (2012). MTORC1 functions as a transcriptional
regulator of autophagy by preventing nuclear transport of TFEB. Autophagy *8*, 903-914.

Maruo, S., Zhao, B., Johannsen, E., Kieff, E., Zou, J., and Takada, K. (2011). Epstein-Barr virus nuclear
antigens 3C and 3A maintain lymphoblastoid cell growth by repressing p16INK4A and p14ARF
expression. Proceedings of the National Academy of Sciences *108*, 1919-1924.

Matsuda, G., Nakajima, K., Kawaguchi, Y., Yamanashi, Y., and Hirai, K. (2003). Epstein-Barr virus (EBV)
nuclear antigen leader protein (EBNA-LP) forms complexes with a cellular anti-apoptosis protein Bcl-2 or
its EBV counterpart BHRF1 through HS1-associated protein X-1. Microbiology and immunology *47*, 91-
99.

McClellan, M.J., Wood, C.D., Ojeniyi, O., Cooper, T.J., Kanhere, A., Arvey, A., Webb, H.M., Palermo,
R.D., Harth-Hertle, M.L., Kempkes, B.*, et al.* (2013). Modulation of enhancer looping and differential gene
targeting by Epstein-Barr virus transcription factors directs cellular reprogramming. PLoS Pathog *9*,
e1003636.

McFadden, K., Hafez, A.Y., Kishton, R., Messinger, J.E., Nikitin, P.A., Rathmell, J.C., and Luftig, M.A.
(2016). Metabolic stress is a barrier to Epstein–Barr virus-mediated B-cell immortalization. Proceedings of
the National Academy of Sciences *113*, E782-E790.

McInnes, L., Healy, J., and Melville, J. (2018). Umap: Uniform manifold approximation and projection for
dimension reduction. arXiv preprint arXiv:180203426.

McLean, C.Y., Bristor, D., Hiller, M., Clarke, S.L., Schaar, B.T., Lowe, C.B., Wenger, A.M., and Bejerano,
G. (2010). GREAT improves functional interpretation of cis-regulatory regions. Nature biotechnology *28*,
495-501.

Messinger, J.E., Dai, J., Stanland, L.J., Price, A.M., and Luftig, M.A. (2019). Identification of Host
Biomarkers of Epstein-Barr Virus Latency IIb and Latency III. mBio *10*.

Minamitani, T., Yasui, T., Ma, Y., Zhou, H., Okuzaki, D., Tsai, C.-Y., Sakakibara, S., Gewurz, B.E., Kieff,
E., and Kikutani, H. (2015). Evasion of affinity-based selection in germinal centers by Epstein–Barr virus
LMP2A. Proceedings of the National Academy of Sciences *112*, 11612-11617.

Mitchell, S., Roy, K., Zangle, T.A., and Hoffmann, A. (2018). Nongenetic origins of cell-to-cell variability in
B lymphocyte proliferation. Proc Natl Acad Sci U S A *115*, E2888-E2897.

Miyashita, E.M., Yang, B., Babcock, G.J., and Thorley-Lawson, D.A. (1997). Identification of the site of
Epstein-Barr virus persistence in vivo as a resting B cell. Journal of virology *71*, 4882-4891.

Mrozek-Gorska, P., Buschle, A., Pich, D., Schwarzmayr, T., Fechtner, R., Scialdone, A., and
Hammerschmidt, W. (2019). Epstein-Barr virus reprograms human B lymphocytes immediately in the
prelatent phase of infection. Proc Natl Acad Sci U S A *116*, 16046-16055.

Nagelkerke, A., Sieuwerts, A.M., Bussink, J., Sweep, F., Look, M.P., Foekens, J.A., Martens, J., and
Span, P.N. (2014). LAMP3 is involved in tamoxifen resistance in breast cancer cells through the
modulation of autophagy. Endocr Relat Cancer *21*, 101-112.

Nikitin, P.A., Yan, C.M., Forte, E., Bocedi, A., Tourigny, J.P., White, R.E., Allday, M.J., Patel, A., Dave,
S.S., and Kim, W. (2010). An ATM/Chk2-mediated DNA damage-responsive signaling pathway
suppresses Epstein-Barr virus transformation of primary human B cells. Cell host & microbe *8*, 510-522.

Nilsson, K., Klein, G., Henle, W., and Henle, G. (1971). The establishment of lymphoblastoid lines from
adult and fetal human lymphoid tissue and its dependence on EBV. International Journal of Cancer *8*,
443-450.

Nishimura, K., Kumazawa, T., Kuroda, T., Katagiri, N., Tsuchiya, M., Goto, N., Furumai, R., Murayama,
A., Yanagisawa, J., and Kimura, K. (2015). Perturbation of ribosome biogenesis drives cells into
senescence through 5S RNP-mediated p53 activation. Cell reports *10*, 1310-1323.

Nonoyama, M., and Pagano, J.S. (1972). Separation of Epstein-Barr virus DNA from large chromosomal
DNA in non-virus-producing cells. Nature New Biology *238*, 169-171.

O'Dea, E.L., Barken, D., Peralta, R.Q., Tran, K.T., Werner, S.L., Kearns, J.D., Levchenko, A., and
Hoffmann, A. (2007). A homeostatic model of IkappaB metabolism to control constitutive NF-kappaB
activity. Mol Syst Biol *3*, 111.

Osorio, D., and Cai, J.J. (2021). Systematic determination of the mitochondrial proportion in human and
mice tissues for single-cell RNA-sequencing data quality control. Bioinformatics *37*, 963-967.

Osorio, D., Yu, X., Yu, P., Serpedin, E., and Cai, J.J. (2019). Single-cell RNA sequencing of a European
and an African lymphoblastoid cell line. Sci Data *6*, 112.

Osorio, D., Yu, X., Zhong, Y., Li, G., Serpedin, E., Huang, J.Z., and Cai, J.J. (2020). Single-cell
expression variability implies cell function. Cells *9*, 14.

Parker, G.A., Crook, T., Bain, M., Sara, E.A., Farrell, P.J., and Allday, M.J. (1996). Epstein-Barr virus
nuclear antigen (EBNA) 3C is an immortalizing oncoprotein with similar properties to adenovirus E1A and
papillomavirus E7. Oncogene *13*, 2541-2549.

Paschos, K., Parker, G.A., Watanatanasup, E., White, R.E., and Allday, M.J. (2012). BIM promoter
directly targeted by EBNA3C in polycomb-mediated repression by EBV. Nucleic acids research *40*, 7233-
7246.

Peng, R., Moses, S.C., Tan, J., Kremmer, E., and Ling, P.D. (2005). The Epstein-Barr virus EBNA-LP
protein preferentially coactivates EBNA2-mediated stimulation of latent membrane proteins expressed
from the viral divergent promoter. Journal of virology *79*, 4492-4505.

Pich, D., Mrozek-Gorska, P., Bouvet, M., Sugimoto, A., Akidil, E., Grundhoff, A., Hamperl, S., Ling, P.D.,
and Hammerschmidt, W. (2019). First days in the life of naive human B lymphocytes infected with
Epstein-Barr virus. MBio *10*, e01723-01719.

Pope, J., Horne, M., and Scott, W. (1968). Transformation of foetal human leukocytes in vitro by filtrates
of a human leukaemic cell line containing herpes-like virus. International journal of cancer *3*, 857-866.

Portis, T., and Longnecker, R. (2004). Epstein–Barr virus (EBV) LMP2A mediates B-lymphocyte survival
through constitutive activation of the Ras/PI3K/Akt pathway. Oncogene *23*, 8619-8628.

Price, A.M., Dai, J., Bazot, Q., Patel, L., Nikitin, P.A., Djavadian, R., Winter, P.S., Salinas, C.A., Barry,
A.P., Wood, K.C.*, et al.* (2017). Epstein-Barr virus ensures B cell survival by uniquely modulating
apoptosis at early and late times after infection. Elife *6*.

Price, A.M., and Luftig, M.A. (2015). To be or not IIb: a multi-step process for Epstein-Barr virus latency
establishment and consequences for B cell tumorigenesis. PLoS Pathog *11*, e1004656.

Qadir, F.S., Saad; Dominguez-Bendala, Juan (2019). 3D Plotting of scRNAseq data using Seurat objects
(1.3) (Zenodo).

Qadir, M.M.F., Álvarez-Cubela, S., Klein, D., van Dijk, J., Muñiz-Anquela, R., Moreno-Hernández, Y.B., Lanzoni, G., Sadiq, S., Navarro-Rubio, B., and García, M.T. (2020). Single-cell resolution analysis of the human pancreatic ductal progenitor cell niche. Proceedings of the National Academy of Sciences *117*, 10876-10887.

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017). Reversed graph embedding resolves complex single-cell trajectories. Nature methods *14*, 979.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics *26*, 841-842.

Raab-Traub, N. (2007). EBV-induced oncogenesis. Human herpesviruses: biology, therapy, and immunoprophylaxis.

Raj, A., Peskin, C.S., Tranchina, D., Vargas, D.Y., and Tyagi, S. (2006). Stochastic mRNA synthesis in mammalian cells. PLoS Biol *4*, e309.

Raj, A., Rifkin, S.A., Andersen, E., and Van Oudenaarden, A. (2010). Variability in gene expression underlies incomplete penetrance. Nature *463*, 913-918.

Raj, A., and Van Oudenaarden, A. (2008). Nature, nurture, or chance: stochastic gene expression and its consequences. Cell *135*, 216-226.

Ramírez, F., Bhardwaj, V., Arrigoni, L., Lam, K.C., Grüning, B.A., Villaveces, J., Habermann, B., Akhtar, A., and Manke, T. (2018). High-resolution TADs reveal DNA sequences underlying genome organization in flies. Nature communications *9*, 1-15.

Ressing, M.E., van Gent, M., Gram, A.M., Hooykaas, M.J., Piersma, S.J., and Wiertz, E.J. (2015). Immune evasion by Epstein-Barr virus. Epstein Barr Virus Volume 2, 355-381.

Rickinson, A., and Kieff, E. (2007). Epstein-Barr virus, p 2655–2700. Fields virology *2*.

Robertson, E.S., Grossman, S., Johannsen, E., Miller, C., Lin, J., Tomkinson, B., and Kieff, E. (1995). Epstein-Barr virus nuclear protein 3C modulates transcription through interaction with the sequence-specific DNA-binding protein J kappa. Journal of virology *69*, 3108-3116.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nature biotechnology *29*, 24-26.

Roy, K., Mitchell, S., Liu, Y., Ohta, S., Lin, Y.-s., Metzig, M.O., Nutt, S.L., and Hoffmann, A. (2019). A regulatory circuit controlling the dynamics of NFκB cRel transitions B cells from proliferation to plasma cell differentiation. Immunity *50*, 616-628. e616.

Saha, A., Jha, H.C., Upadhyay, S.K., and Robertson, E.S. (2015). Epigenetic silencing of tumor suppressor genes during in vitro Epstein–Barr virus infection. Proceedings of the National Academy of Sciences *112*, E5199-E5207.

Sandberg, M., Hammerschmidt, W., and Sugden, B. (1997). Characterization of LMP-1's association with TRAF1, TRAF2, and TRAF3. Journal of Virology *71*, 4649-4656.

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. Nature biotechnology *33*, 495-502.

Satpathy, A.T., Granja, J.M., Yost, K.E., Qi, Y., Meschi, F., McDermott, G.P., Olsen, B.N., Mumbach, M.R., Pierce, S.E., and Corces, M.R. (2019). Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion. Nature biotechnology *37*, 925-936.

Schubart, D.B., Rolink, A., Kosco-Vilbois, M.H., Botteri, F., and Matthias, P. (1996). B-cell-specif ic coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre formation. Nature *383*, 538-542.

Settembre, C., Di Malta, C., Polito, V.A., Arencibia, M.G., Vetrini, F., Erdin, S., Erdin, S.U., Huynh, T., Medina, D., and Colella, P. (2011). TFEB links autophagy to lysosomal biogenesis. science *332*, 1429-1433.

Shair, K.H., Schnegg, C.I., and Raab-Traub, N. (2008). EBV latent membrane protein 1 effects on
plakoglobin, cell growth, and migration. Cancer research *68*, 6997-7005.

Shalek, A.K., Satija, R., Adiconis, X., Gertner, R.S., Gaublomme, J.T., Raychowdhury, R., Schwartz, S.,
Yosef, N., Malboeuf, C., and Lu, D. (2013). Single-cell transcriptomics reveals bimodality in expression
and splicing in immune cells. Nature *498*, 236-240.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and
Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction
networks. Genome research *13*, 2498-2504.

Shapiro, E., Biezuner, T., and Linnarsson, S. (2013). Single-cell sequencing-based technologies will
revolutionize whole-organism science. Nature Reviews Genetics *14*, 618-630.

Simpson, E.H. (1951). The interpretation of interaction in contingency tables. Journal of the Royal
Statistical Society: Series B (Methodological) *13*, 238-241.

Sinclair, A.J., Palmero, I., Peters, G., and Farrell, P. (1994). EBNA-2 and EBNA-LP cooperate to cause
G0 to G1 transition during immortalization of resting human B lymphocytes by Epstein-Barr virus. The
EMBO journal *13*, 3321-3328.

Skalska, L., White, R.E., Franz, M., Ruhmann, M., and Allday, M.J. (2010). Epigenetic repression of
p16INK4A by latent Epstein-Barr virus requires the interaction of EBNA3A and EBNA3C with CtBP. PLoS
pathogens *6*, e1000951.

Smith, N., Tierney, R., Wei, W., Vockerodt, M., Murray, P.G., Woodman, C.B., and Rowe, M. (2013).
Induction of interferon-stimulated genes on the IL-4 response axis by Epstein-Barr virus infected human b
cells; relevance to cellular transformation. PLoS One *8*, e64868.

SoRelle, E.D., Dai, J., Bonglack, E.N., Heckenberg, E.M., Zhou, J.Y., Giamberardino, S.N., Bailey, J.A.,
Gregory, S.G., Chan, C., and Luftig, M.A. (2021). Single-cell RNA-seq reveals transcriptomic
heterogeneity mediated by host–pathogen dynamics in lymphoblastoid cell lines. Elife *10*, e62586.

Stros, M., Ozaki, T., Bacıková, A., Kageyama, H., and Nakagawara, A. (2002). HMGB1 and HMGB2 cell-
specifically down-regulate the p53-and p73-dependent sequence-specific transactivation from the human
Bax gene promoter. Journal of Biological Chemistry *277*, 7157-7164.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck III, W.M., Hao, Y., Stoeckius, M.,
Smibert, P., and Satija, R. (2019). Comprehensive integration of single-cell data. Cell *177*, 1888-1902.
e1821.

Stuart, T., Srivastava, A., Lareau, C., and Satija, R. (2020). Multimodal single-cell chromatin analysis with
Signac. BioRxiv.

Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2021). Single-cell chromatin state
analysis with Signac. Nature Methods, 1-9.

Styles, C.T., Bazot, Q., Parker, G.A., White, R.E., Paschos, K., and Allday, M.J. (2017). EBV
epigenetically suppresses the B cell-to-plasma cell differentiation pathway while establishing long-term
latency. PLoS Biol *15*, e2001992.

Suan, D., Kräutler, N.J., Maag, J.L., Butt, D., Bourne, K., Hermes, J.R., Avery, D.T., Young, C., Statham,
A., and Elliott, M. (2017). CCR6 defines memory B cell precursors in mouse and human germinal centers,
revealing light-zone location and predominant low antigen affinity. Immunity *47*, 1142-1153. e1144.

Sun, C.C., and Thorley-Lawson, D.A. (2007). Plasma cell-specific transcription factor XBP-1s binds to
and transactivates the Epstein-Barr virus BZLF1 promoter. J Virol *81*, 13566-13577.

Szymula, A., Palermo, R.D., Bayoumy, A., Groves, I.J., Ba Abdullah, M., Holder, B., and White, R.E.
(2018). Epstein-Barr virus nuclear antigen EBNA-LP is essential for transforming naive B cells, and
facilitates recruitment of transcription factors to the viral genome. PLoS pathogens *14*, e1006890.

Taylor, G.M., Raghuwanshi, S.K., Rowe, D.T., Wadowsky, R.M., and Rosendorff, A. (2011). Endoplasmic
reticulum stress causes EBV lytic replication. Blood, The Journal of the American Society of Hematology
118, 5528-5539.

Taylor, J.J., Pape, K.A., Steach, H.R., and Jenkins, M.K. (2015). Apoptosis and antigen affinity limit
effector cell differentiation of a single naïve B cell. Science *347*, 784-787.

Thorley-Lawson, D.A. (2001). Epstein-Barr virus: exploiting the immune system. Nature Reviews
Immunology *1*, 75-82.

Thorley-Lawson, D.A., and Gross, A. (2004). Persistence of the Epstein–Barr virus and the origins of
associated lymphomas. New England Journal of Medicine *350*, 1328-1337.

Thorley-Lawson, D.A., and Mann, K.P. (1985). Early events in Epstein-Barr virus infection provide a
model for B cell activation. The Journal of experimental medicine *162*, 45-59.

Tomkinson, B., Robertson, E., and Kieff, E. (1993). Epstein-Barr virus nuclear proteins EBNA-3A and
EBNA-3C are essential for B-lymphocyte growth transformation. Journal of virology *67*, 2014-2025.

Tosato, G., Blaese, R., and Yarchoan, R. (1985). Relationship between immunoglobulin production and
immortalization by Epstein Barr virus. The Journal of Immunology *135*, 959-964.

Trapnell, C. (2015). Defining cell types and states with single-cell genomics. Genome research *25*, 1491-
1498.

Tsai, K., Thikmyanova, N., Wojcechowskyj, J.A., Delecluse, H.-J., and Lieberman, P.M. (2011). EBV
tegument protein BNRF1 disrupts DAXX-ATRX to activate viral early gene transcription. PLoS pathogens
7, e1002376.

Uchida, J., Yasui, T., Takaoka-Shichijo, Y., Muraoka, M., Kulwichit, W., Raab-Traub, N., and Kikutani, H.
(1999). Mimicry of CD40 signals by Epstein-Barr virus LMP1 in B lymphocyte responses. Science *286*,
300-303.

Wang, A., Welch, R., Zhao, B., Ta, T., Keleş, S., and Johannsen, E. (2015). Epstein-Barr virus nuclear
antigen 3 (EBNA3) proteins regulate EBNA2 binding to distinct RBPJ genomic sites. Journal of virology
90, 2906-2919.

Wang, C., Li, D., Zhang, L., Jiang, S., Liang, J., Narita, Y., Hou, I., Zhong, Q., Zheng, Z., and Xiao, H.
(2019). RNA sequencing analyses of gene expression during Epstein-Barr virus infection of primary B
lymphocytes. Journal of virology *93*, e00226-00219.

Wang, L.W., Jiang, S., and Gewurz, B.E. (2017). Epstein-Barr virus LMP1-mediated oncogenicity. Journal
of virology *91*, e01718-01716.

Wendel-Hansen, V., Rosén, A., and Klein, G. (1987). EBV-transformed lymphoblastoid cell lines down-
regulate ebna in parallel with secretory differentiation. International journal of cancer *39*, 404-408.

White, R.E., Groves, I.J., Turro, E., Yee, J., Kremmer, E., and Allday, M.J. (2010). Extensive co-operation
between the Epstein-Barr virus EBNA3 proteins in the manipulation of host gene expression and
epigenetic chromatin modification. PloS one *5*, e13979.

Wills, Q.F., Livak, K.J., Tipping, A.J., Enver, T., Goldson, A.J., Sexton, D.W., and Holmes, C. (2013).
Single-cell gene expression analysis reveals genetic associations masked in whole-tissue experiments.
Nature biotechnology *31*, 748-752.

Wolff, J., Bhardwaj, V., Nothjunge, S., Richard, G., Renschler, G., Gilsbach, R., Manke, T., Backofen, R.,
Ramírez, F., and Grüning, B.A. (2018). Galaxy HiCExplorer: a web server for reproducible Hi-C data
analysis, quality control and visualization. Nucleic acids research *46*, W11-W16.

Wolff, J., Rabbani, L., Gilsbach, R., Richard, G., Manke, T., Backofen, R., and Grüning, B.A. (2020).
Galaxy HiCExplorer 3: a web server for reproducible Hi-C, capture Hi-C and single-cell Hi-C data
analysis, quality control and visualization. Nucleic acids research *48*, W177-W184.

Wood, V., O'neil, J., Wei, W., Stewart, S., Dawson, C., and Young, L. (2007). Epstein–Barr virus-encoded
EBNA1 regulates cellular gene transcription and modulates the STAT1 and TGF β signaling pathways.
Oncogene *26*, 4135-4147.

Wu, J., Xia, L., Yao, X., Yu, X., Tumas, K.C., Sun, W., Cheng, Y., He, X., Peng, Y.-c., and Singh, B.K.
(2020). The E3 ubiquitin ligase MARCH1 regulates antimalaria immunity through interferon signaling and
T cell activation. Proceedings of the National Academy of Sciences *117*, 16567-16578.

Wu, Y., Bressette, D., Carrell, J.A., Kaufman, T., Feng, P., Taylor, K., Gan, Y., Cho, Y.H., Garcia, A.D.,
and Gollatz, E. (2000). Tumor necrosis factor (TNF) receptor superfamily member TACI is a high affinity
receptor for TNF family members APRIL and BLyS. Journal of Biological Chemistry *275*, 35478-35485.

Yanai, H., Ban, T., Wang, Z., Choi, M.K., Kawamura, T., Negishi, H., Nakasato, M., Lu, Y., Hangai, S.,
and Koshiba, R. (2009). HMGB proteins function as universal sentinels for nucleic-acid-mediated innate
immune responses. Nature *462*, 99-103.

Yates, J.L., Warren, N., and Sugden, B. (1985). Stable replication of plasmids derived from Epstein–Barr
virus in various mammalian cells. Nature *313*, 812-815.

Yetming, K.D., Lupey-Green, L.N., Biryukov, S., Hughes, D.J., Marendy, E.M., Miranda, J.L., and Sample,
J.T. (2020). The BHLF1 locus of Epstein-Barr virus contributes to viral latency and B-cell immortalization.
Journal of virology *94*, e01215-01220.

Young, L.S., Arrand, J.R., and Murray, P.G. (2007). EBV gene expression and regulation. Human
herpesviruses: biology, therapy, and immunoprophylaxis.

Young, L.S., and Rickinson, A.B. (2004). Epstein-Barr virus: 40 years on. Nat Rev Cancer *4*, 757-768.

Zhao, B., Zou, J., Wang, H., Johannsen, E., Peng, C.-w., Quackenbush, J., Mar, J.C., Morton, C.C.,
Freedman, M.L., and Blacklow, S.C. (2011). Epstein-Barr virus exploits intrinsic B-lymphocyte
transcription programs to achieve immortal cell growth. Proceedings of the National Academy of Sciences
108, 14902-14907.

Zheng, G.X., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B., Wheeler, T.D.,
McDermott, G.P., and Zhu, J. (2017). Massively parallel digital transcriptional profiling of single cells.
Nature communications *8*, 1-12.

Zhou, H., Schmidt, S.C., Jiang, S., Willox, B., Bernhardt, K., Liang, J., Johannsen, E.C., Kharchenko, P.,
Gewurz, B.E., and Kieff, E. (2015). Epstein-Barr virus oncoprotein super-enhancers control B cell growth.
Cell host & microbe *17*, 205-216.

Zhu, C., Yu, M., Huang, H., Juric, I., Abnousi, A., Hu, R., Lucero, J., Behrens, M.M., Hu, M., and Ren, B.
(2019). An ultra high-throughput method for single-cell joint analysis of open chromatin and
transcriptome. Nature structural & molecular biology *26*, 1063-1070.

1323