Title:

Chloride ions evoke taste sensations by binding to the extracellular ligand-binding

domain of sweet/umami taste receptors

Authors:

Nanako Atsumi[1†], Keiko Yasumatsu[2†], Yuriko Takashina[3], Chiaki Ito[1], Norihisa Yasui[1], Robert

F. Margolskee[4], Atsuko Yamashita[1]

[†]These authors contributed equally to this work.

Affiliation:

[1]Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University,

Okayama, Japan, [2]Oral Health Science Center, Tokyo Dental College, Tokyo, [3]School of

Pharmaceutical Sciences, Okayama University, Okayama, Japan, [4]Monell Chemical Senses

Center, Philadelphia, PA, USA.

Corresponding author:

Atsuko Yamashita, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

Okayama University, 1-1-1, Tsushima-naka, Kita-ku, Okayama 600-8530, Japan. Fax: +81-86-

251-7974. E-mail: a_yama@okayama-u.ac.jp.

Abstract

Salt taste sensation is multifaceted: NaCl at low or high concentrations is preferably or

aversively perceived through distinct pathways. Cl^- is thought to participate in taste sensation

through an unknown mechanism. Here we describe Cl^- ion binding and the response of taste

receptor type 1 (T1r), a receptor family composing sweet/umami receptors. The T1r2a/T1r3

heterodimer from the medaka fish, currently the sole T1r amenable to structural analyses,

exhibited a specific Cl^- binding in the vicinity of the amino-acid-binding site in the ligand-

binding domain (LBD) of T1r3, which is likely conserved across species, including human T1r3.

The Cl^- binding induced a conformational change in T1r2a/T1r3LBD at sub- to low-mM

concentrations similar to canonical taste substances. Furthermore, oral Cl^- application to mice

increased impulse frequencies of taste nerves connected to T1r-expressing taste cells and

promoted their behavioral preferences attenuated by a T1r-specific blocker or T1r3 knock-out.

These results suggest that the Cl^- evokes taste sensations by binding to T1r, thereby serving as

another preferred salt taste pathway at a low concentration.

Introduction

The taste sensation is initiated by specific interactions between chemicals in food and taste receptors in taste buds in the oral cavity. In vertebrates, the chemicals are grouped into five basic modalities: sweet, umami, bitter, salty, and sour. This sensation occurs through taste receptor recognition specific to a group of chemicals representing each taste modality (Taruno *et al*, 2021). Regarding the salty taste, the preferable taste, ~100 mM concentration of table salt, is evoked by specific interaction between the epithelial sodium channel (ENaC) and sodium ion (Chandrashekar *et al*, 2010) (Fig 1). Notably, salt sensation exhibits multifaced properties (Roper, 2015), thereby suggesting the existence of an adequate concentration range for salt intake to maintain the homeostasis of body fluid concentration. For example, high concentrations of salt over levels perceived as a preferred taste, such as ~500 mM, stimulate bitter and sour taste cells and are perceived as an aversive taste (Oka *et al*, 2013). Conversely, low salt concentrations under the "preferable" concentration, such as several mM to a hundred mM concentrations, are perceived as sweet by human panels (Bartoshuk *et al*, 1964; Bartoshuk *et al*, 1978; Cardello, 1979). However, its mechanism has never been extensively pursued.

Figure 1. Salt taste sensation. Approximate concentration ranges of salt taste perceptions in humans (Bartoshuk *et al.*, 1978) and qualities of taste sensation with known cells and receptors responsible for their sensing are summarized.

The various impacts of the salt taste sensation indicate multiple salt detection pathways in taste

buds (Roper, 2015). Moreover, another component of table salt, the chloride ion, participates

in taste sensation because of the existence of the "anion effect": the salty taste is most strongly

perceived when the counter anion is a chloride ion (Ye et al, 1991). Several reports suggested

a certain cellular/molecular machinery underlying anion-sensitive Na^+-detection or Cl^--

detection, which is independent of ENaC (Lewandowski et al, 2016; Roebber et al, 2019).

Indeed, a recent study indicated that transmembrane channel-like 4 (TMC4) expressed in taste

buds involves high-concentration Cl^--sensation (Kasahara et al, 2021). Nevertheless, no

candidate molecule capable of sensing low or preferable concentrations of Cl^- has been

elucidated. Therefore, the complete understanding of salt taste sensation, including the

mechanism of chloride ion detection, remains unclear.

Unlike salt taste sensation, those for nutrients as sugars, amino acids, and nucleotides

are understood as sweet and umami sensations through specific receptor proteins (Li et al, 2002;

Nelson et al, 2002; Nelson et al, 2001). Sweet and umami receptors are composed of taste

receptor type 1 (T1r) proteins in the class C G protein-coupled receptor family. In humans, the

T1r1/T1r3 heterodimer serves as the umami taste receptor and responds to amino acids as L-

glutamate and aspartate, and nucleotides. In contrast, the T1r2/T1r3 heterodimer is the sweet

taste receptor and responds to sugars. We previously elucidated the crystallographic structure

of the medaka fish T1r2a/T1r3 extracellular ligand-binding domain (LBD) (Nuemket et al,

2017), which is currently the sole reported structure of T1rs. In the structure, the amino acid-

binding was observed in the middle of the LBD of T1r2a and T1r3 subunits, which was

consistent with the fact that T1r2a/T1r3 is an amino-acid receptor (Oike et al, 2007).

Furthermore, chloride ion binding was found in the vicinity of the amino-acid binding site in

T1r3 (Fig 2A). So far, the physiological significance of Cl^--binding for T1rs functions remains

unexplored. Nevertheless, chloride ions regulate other receptors in class C G protein-coupled

receptors (GPCRs), such as metabotropic glutamate receptors (mGluRs) and calcium-sensing

receptors (CaSRs), and act as positive modulators for agonist binding (Eriksen & Thomsen,

1995; Kuang & Hampson, 2006; Liu *et al*, 2020; Tora *et al*, 2018; Tora *et al*, 2015). The

potential effect of Cl⁻-binding on T1r receptor function is of significant interest under these

conditions.

Here, we investigated the Cl⁻ actions on T1rs using structural, biophysical, and

physiological analyses. The Cl⁻ binding to the LBD was investigated using the medaka fish

T1r2a/T1r3LBD, which is amenable to structural and biophysical analyses. Since the Cl⁻-

binding site in T1r3 was conserved across various species, taste nerve recordings from mice

were used to investigate the physiological significance of Cl⁻. The results suggest that Cl⁻

induces the moderate response via T1rs, thereby implying that T1rs are involved in Cl⁻

sensation in taste buds.

Results

Cl⁻-binding site in T1r3

In the previously reported structure of T1r2a/T1r3LBD crystallized in the presence of NaCl,

bound Cl⁻ was identified based on electron density and binding distances (Nuemket *et al.*, 2017).

To verify the Cl⁻-binding, Cl⁻ in the T1r2a/T1r3LBD crystal was substituted with Br⁻, a halogen

ion amenable for specific detection by anomalous scattering using a synchrotron light source

(Table S1). The diffraction data from the crystal resulted in an anomalous difference Fourier

peak at 14.1 σ at the site in the vicinity of the amino-acid binding site in T1r3, where Cl⁻ was

originally bound (Fig 2B, S1A). For further confirmation, the anomalous data of the original

crystal containing Cl⁻ was collected at 2.7 Å, where the anomalous peak for Cl and several

other elements such as Ca or S can be detected (Table S1). The resultant anomalous difference

Fourier map showed a peak at the bound Cl⁻ position, while all the other peaks were observed

at the S atoms in the protein (Fig 2C, S1B). These results verify that the site is able to bind

halogen ions, likely accommodating Cl⁻ under physiological conditions.

Figure 2. Cl⁻-binding sites in the medaka fish taste receptor T1r2a/T1r3LBD. (A)

Schematic drawing of the overall architecture of T1r2a/T1r3. The crystal structure (PDB ID:

5X2M) (Nuemket *et al.*, 2017) is shown at the LBD region, and helices B and C in T1r3 are

labeled. (B) Anomalous difference Fourier map (4.5 σ, red) of the Br⁻-substituted

T1r2a/T1r3LBD crystal. (C) Anomalous difference Fourier map (4.5 σ, red) of the Cl⁻-bound

T1r2a/T1r3LBD crystal derived from the diffraction data collected at the wavelength of 2.7 Å.

In panels B and C, the site originally identified the Cl⁻-binding was framed. (D) A close-up

view of the Cl⁻-binding site in T1r3LBD in the Cl⁻-bound T1r2a/T1r3LBD (PDB ID: 5X2M).

(E) Amino-acid sequence alignment of T1r proteins and the related receptors at the Cl⁻-binding

site. The "h," "m," and "mf" prefixes to T1rs indicate human, mouse, and medaka fish,

respectively. The position corresponding to Thr105 in T1r3 from medaka fish is highlighted.

(F) The structures of ANPR (PDB ID: 1T34, left) (Ogawa *et al*, 2004) and mGluR2 (PDB ID:

5CNI, right) (Monn *et al*, 2015a) bound with Cl⁻. (G) Superposition of the Cl⁻-binding site in

T1r3, ANPR, and mGluR2.

Cl⁻ was coordinated at the binding site by the side-chain hydroxyl group of Thr105

and the main-chain amide groups of Gln148 and Ser149 (Fig 2D). These main-chain

coordinating residues are followed by Ser150, a critical residue for binding amino-acid ligands

(Nuemket *et al.*, 2017). Furthermore, the loop regions where Thr105 and the Gln148-Ser150

locate are followed by helices B and C, respectively. These helices are essential structural units

at the heterodimer interface (Nuemket *et al.*, 2017) (Fig 2A). They are known to reorient upon

agonist binding, resulting in conformation rearrangement of the subunits in the dimer, likely

inducing receptor activation in class C GPCRs (Koehl *et al*, 2019; Kunishima *et al*, 2000). In

addition, the side-chain hydroxyl group of Ser149, which serves as a cap for the positive helix

dipole of helix C, simultaneously functions as a distal ligand for Cl⁻ coordination. Therefore,

the Cl⁻ binding at this site is important for organizing the structure of the amino-acid binding

site and the heterodimer interface.

The Cl⁻-binding site observed in the crystal structure is most likely conserved among

T1r3s in various organisms, such as humans (Fig 2E). Thr105, the residue that provides the

side-chain-coordinating ligand for Cl$^-$-binding, is strictly conserved as either serine or threonine

among T1r3s. Additionally, the amino-acid sequence motifs surrounding the main-chain-

coordinating ligands, FP^{45}L and VIGP$\zeta^{148}\zeta^{149}$, where "ζ" is a hydrophilic amino acid, are well

conserved to present the main-chain amide groups to coordinate Cl$^-$ with an appropriate

geometry. Notably, the site structurally corresponds to the Cl$^-$-binding site in the hormone-

binding domain of the atrial natriuretic peptide receptor (ANPR) (Fig 2F, G), in which Cl$^-$

positively regulates the peptide hormone binding (Misono, 2000). Although ANPR is not a

member of class C GPCR, the hormone-binding domain in ANPR shares a similar structural

fold with LBD of T1rs and other class C GPCRs, and bacterial periplasmic-binding proteins

(Kunishima *et al.*, 2000; van den Akker *et al*, 2000). Accordingly, the conservation of the

structure and the sequence motif at the Cl$^-$-binding site at ANPR is also observed on mGluRs

(Ogawa *et al*, 2010). Indeed, Cl$^-$ binding at the site corresponding to that in T1r3 was observed

in several mGluR and CaSR structures (Monn *et al*, 2015b; Zhang *et al*, 2016) (Fig 2F, G) and

was identified as a potential site responsible for regulating agonist binding (Liu *et al.*, 2020;

Tora *et al.*, 2015). These results strongly imply the possibility that Cl$^-$ has some actions on T1r

receptor functions.

In contrast, conservation at the Thr105 position was not observed among T1r1 and

T1r2 (Fig 2E). Evidently, no significant anomalous peak derived from Br$^-$- or Cl$^-$-binding was

observed in the crystal structure at the corresponding site in T1r2a (Fig 2B, C). His100 in T1r2a,

which corresponds to Thr105 in T1r3, adopted a significantly different side-chain conformation

from that of Thr105 in T1r3 (Fig S1C). Therefore, T1r1 and T1r2's ability to bind Cl$^-$ is unlikely.

In addition to the Cl$^-$-binding site discussed above, the Br$^-$-substituted crystal

exhibited an anomalous peak at 8.5 σ in T1r2a, at a position close to the Lys265 side-chain ε-

amino group (Fig 2B, S1D). Nevertheless, Cl$^-$-binding was not observed in the original Cl$^-$-

contained crystal. This is further confirmed by the absence of an anomalous peak at this position

in the data collected at 2.7 Å (Fig 2C, S1E). Therefore, the site might have the ability to bind

anions such as Br^- or larger; but be not specific to Cl^-. In human T1r1, the residue corresponding

to Lys265 (Arg277) was suggested as a critical residue for activities of inosine monophosphate,

an umami enhancer (Zhang *et al*, 2008).

Cl^--binding properties in T1r2a/T1r3LBD

To investigate the Cl^- actions on T1r functions, we first examined the properties of the Cl^--

binding to medaka T1r2a/T1r3LBD using various biophysical techniques. For this purpose, the

purified T1r2a/T1r3LBD was subjected to differential scanning fluorimetry (DSF), which we

previously used for the amino acid-binding analysis (Yoshida *et al*, 2019). In order to prepare

a Cl^--free condition, Cl^- in the sample was substituted with gluconate, as it is unlikely

accommodated in the Cl^--site due to its much larger size. We confirmed that gluconate does not

serve as a ligand for T1r2a/T1r3LBD (Fig S2A).

The addition of Cl^- to the Cl^--free T1r2a/T1r3LBD sample resulted in thermal

stabilization of the protein (Fig 3A), which is indicative of Cl^- binding to the protein. The

apparent K_d value for Cl^- estimated by the melting temperatures (T_m) at various Cl^-

concentrations was ~110 µM (Fig 3B, Table 1). The Cl^--dependent thermal stabilization was

confirmed by the fluorescence-detection size-exclusion chromatography-based thermostability

(FSEC-TS) assay (Hattori *et al*, 2012) (Fig 3C, Table 1). However, the Cl^--dependent

stabilization was not observed on T1r2a/T1r3 with the Cl^--site mutation, T105A in T1r3. In the

case of this mutant, the T_m values for both in the presence and absence of Cl^- was similar to the

values obtained for the wild-type protein in the absence of Cl^- (Fig 3C, S2B, Table 1). These

results indicate that the Cl^- effect attributed to the identical site where the Cl^--binding was

observed in the crystal structure of T1r3.

Figure 3. The Cl⁻-binding properties of T1r2a/T1r3LBD. (A) Representative thermal melt

curves of T1r2a/T1r3LBD in the presence of 0.003 –10 mM concentrations of Cl⁻ measured

using DSF. (B) Dose-dependent T_m changes of T1r2a/T1r3LBD by addition of Cl⁻. ($n = 4$) (C)

Thermal melting curves of WT and the T1r3-T105A mutant of T1r2a/T1r3LBD in the presence

and absence of Cl⁻, analyzed by FSEC-TS. ($n = 1$) (D) Dose-dependent FRET signal changes

of the T1r2aLBD-Cerulean and T1r3LBD-Venus heterodimer by addition of Cl⁻. ($n = 3$) (E)

FRET index increases by adding 10-mM Cl⁻ or 1-mM L-glutamine to the WT or T1r3-T105A

mutant T1r2aLBD-Cerulean/T1r3LBD-Venus heterodimer relative to that in the absence of any

ligand in the absence of Cl⁻. ($n = 3$) (F) Dose-dependent FRET signal changes of the

T1r2aLBD-Cerulean and T1r3LBD-Venus heterodimer induced by the addition of L-glutamine

in the presence and absence of Cl⁻. ($n = 3$) The experiments were performed two (panels A, B,

D, and E), three (C and WT), four (F and +Cl⁻ condition), or one (C and mutant; F and –Cl⁻

condition) time(s), and the results from one representative experiment are shown with numbers

204   of technical replicates. Data points represent mean and s.e.m.

206   **Table 1.** Properties of the Cl⁻-binding to T1r2a/T1r3LBD.

Cl⁻-binding, DSF ($n = 4$)		
$K_{d\text{-app}}$ (mM)	0.111 ± 0.046	
Protein thermal stability, DSF ($n = 6$)		
condition	+Cl⁻	-Cl⁻
Melting temperature (˚C)	55.2 ± 0.03	46.6 ± 0.06
Protein thermal stability, FSEC-TS		
wild type T1r2a/T1r3LBD ($n = 3$)		
condition	+Cl⁻	-Cl⁻
Melting temperature (˚C)	56.4 ± 5.1	46.0 ± 0.3
mutant T1r2a/T1r3-T105ALBD		
condition	+Cl⁻	-Cl⁻
Melting temperature (˚C)	42.7 ± 0.1	46.7 ± 0.7
Cl⁻- binding, FRET ($n = 3$)		
FRET index minimum	1.00 ± 0.005	
FRET index change	0.119 ± 0.014	
EC_{50} (mM)*	1.23 ± 0.53	
L-glutamine binding, FRET ($n = 3$)		
condition	+Cl⁻	-Cl⁻
FRET index minimum	1.08 ± 0.01	0.86 ± 0.01
FRET index change	0.11	0.17
EC_{50} (µM)	3.78 ± 1.29	3.87 ± 1.76
Hill coefficient	1.20 ± 0.88	0.94 ± 0.29
L-glutamine binding, ITC		
condition	+Cl⁻	-Cl⁻
N (sites)	0.389 ± 0.028	0.303 ± 0.023
K_a (M⁻¹)	(2.85 ± 0.65) × 10⁵	(2.12 ± 0.46) × 10⁵
[converted to K_d (µM)]	[3.51]	[4.72]
ΔH (kcal/mol)	⁻12.3 ± 1.2	−12.9 ± 1.3
ΔS (cal/mol/deg)	−16.3	⁻18.8

*Hill coefficient was fixed to 1 for fitting.

Next, we examined the consequence of the Cl^--binding to T1r2a/T1r3LBD via

Förster resonance energy transfer (FRET) using the fluorescent protein-fused sample. Class C

GPCRs commonly exhibit agonist-induced conformational changes in LBD, such as the dimer

rearrangement, which is essential for receptor activation and signaling (Ellaithy *et al*, 2020;

Koehl *et al.*, 2019; Kunishima *et al.*, 2000; Lin *et al*, 2021). Consistent with this, we previously

reported that T1r2a/T1r3LBD shows conformational change concomitant with the binding of

amino acids, which can be detected as increased FRET intensity (Nango *et al*, 2016). Notably,

adding Cl^- to the fluorescent protein-fused T1r2a/T1r3LBD also increased FRET intensities,

similar to amino acids (Fig 3D). The EC_{50} for Cl^--induced FRET signal change was determined

as ~1 mM (Table 1). Note that both DSF and FRET estimations have some degree of error: the

former produced slightly lower values than the latter, particularly in the case of weak affinities

in the mM concentration range (Yoshida *et al.*, 2019). As such, although the EC_{50} value

determined by FRET was slightly higher than the apparent K_d value of Cl^- determined by DSF,

the two are most likely relevant. Considering that the Cl^--binding site is located adjacent to the

dimer interface, which exhibits reorientation upon agonist binding (Fig 2D), the results suggest

that the Cl^-binding to T1r2a/T1r3LBD induces a conformational rearrangement of

T1r2a/T1r3LBD similar to its agonist amino acid. Nevertheless, the extent of the FRET change

induced by Cl^- was smaller than the changes induced by amino acids, such as ~1/2 of the latter

(Fig 3E). Therefore, the results suggest that the extent of the conformational change induced by

Cl^- is smaller than the change induced by amino acids. The extent of Cl^--dependent FRET index

change was reduced on T1r2a/T1r3 with the Cl^--site mutation, T105A in T1r3 (Fig 3E).

Considering that the amino-acid-dependent change in the mutant was also significantly reduced

(Fig 3E), the T105A mutation on T1r3 might result in losing the ability of the conformational

change induced by Cl^-- and amino-acid bindings, although the possibility of deactivation of the

protein during preparation due to its low stability cannot be excluded.

233   In addition to the Cl⁻-binding effect, the Cl⁻ effect on amino-acid binding to

234 T1r2a/T1r3LBD was investigated by FRET and isothermal calorimetry (ITC). The K_d values

235 for L-glutamine binding determined by ITC, as well as the EC_{50} values and the other parameters

236 for L-glutamine-induced conformational change determined by FRET, did not differ in the

237 presence and absence of Cl⁻ (Fig 3F, S2C, 2D, and Table 1). These results indicated that the Cl⁻

238 binding had no significant effect on the binding of L-glutamine, a representative taste substance,

239 at least for T1r2a/T1r3LBD from medaka fish.

241 **Taste response to Cl⁻ through T1rs in mouse**

242 Biophysical studies on T1r2a/T1r3LBD from medaka fish suggested that Cl⁻binding to

243 T1r3LBD induces a conformational change similar to that of an agonist without affecting

244 agonist binding. As described above, the Cl⁻-binding site is likely conserved among T1r3 in

245 various species, such as those in mammals. Therefore, we analyzed single fiber responses from

246 mouse chorda tympani nerve to investigate the physiological effect of Cl⁻ on taste sensation.

247 While conventional cell-based receptor assay systems are affected by changes in extracellular

248 ionic components, the application of various solutions to the taste pore side of the taste buds

249 projected to taste nerve systems, are transduced exclusively as "taste" signals, without inducing

250 the other cellular responses derived from the ionic component changes in the surrounding

251 environment.

252   We first identified nerve fibers that connect to T1r-expressing taste cells in wild-type

253 (WT) mice, i.e., that receive taste information from cells likely possessing sweet (T1r2/T1r3)

254 and umami (T1r1/T1r3) receptors (Yasumatsu *et al*, 2012). The identification was evidenced by

255 responses to T1r agonists, such as sugars and amino acids, which were inhibited by gurmarin

256 (Gur), a T1r-specific blocker (Daly *et al*, 2013; Margolskee *et al*, 2007; Ninomiya & Imoto,

257 1995; Ninomiya *et al*, 1999). Then, we examined the responses to Cl⁻ in these fibers.

Remarkably, the fibers also exhibited responses induced by Cl^-, which was applied as a form

of NMDG-Cl devoid of the known salty taste stimulant, sodium ion (Fig 4A). Cl^--induced

impulse frequencies from the nerves increased in a concentration-dependent manner (Fig 4B).

The responses to NMDG-Cl, NaCl, and KCl at the same concentrations did not differ

significantly (repeated measures analysis of variance [ANOVA], $P > 0.05$), whereas responses

to NMDG-gluconate were significantly smaller than those to NMDG-Cl (repeated measures

ANOVA: $F_{(1, 43)} = 31.33$, $P < 0.001$) and did not induce explicit responses up to 10 mM. These

results confirmed that the observed responses were attributed to Cl^-. All responses to Cl^-,

regardless of the type of counter cations, were significantly decreased by lingual treatment with

Gur (Fig 4B, repeated measures ANOVA: $F_{(1, 42)} = 56.65$, $P < 0.001$ for NMDG-Cl; $F_{(1, 50)} =$

24.78, $P < 0.001$ for NaCl; and $F_{(1, 45)} = 35.72$, $P < 0.001$ for KCl). Furthermore, responses to

Cl^- in T1r3-KO mice were significantly lower than those in WT mice (repeated measures

ANOVA: $F_{(1, 43)} = 25.36$, $P < 0.001$; Fig 4C). The results indicate that the observed Cl^--

dependent responses were mediated by T1r. Notably, the Cl^--concentration range that induced

nerve responses was lower ($\leq \sim 10$ mM) than that for Na^+-detection by ENaC when applied as

the NaCl form ($\geq \sim 30$ mM) (Chandrashekar *et al.*, 2010) but is consistent with those for Cl^--

binding and Cl^--induced conformational change observed on T1r2a/T1r3LBD (Table 1). These

results suggest that a low Cl^- concentration is sensorily detected via T1r in taste buds. Although

the responses induced by known taste substances for T1rs, such as sugars and amino acids,

range from tens to hundreds of impulse frequencies per 10 s (Yasumatsu *et al.*, 2012); the

maximum response level induced by Cl^- was low, ~ 10 per 10 s (Fig 4B). According to our

observations, Cl^- likely produces a "light" taste sensation compared with other known taste

substances.

Figure 4. Electrophysiological and behavioral analyses of the T1r-mediated Cl⁻ responses

in mouse. (A–D) Results of single fiber recordings from the mouse chorda tympani nerve. (A)

Representative recordings of single fibers that connect to T1r-expressing taste cells. The stimuli

were 10-mM NMDG-Cl, 100-mM sucrose, 100-mM sucrose + 10-mM NMDG-Cl, 20-mM L-

glutamine, or 20-mM L-glutamine + 10-mM NMDG-Cl. Lines indicate the application of

stimuli to the tongue. All the responses were suppressed by lingual treatment with a T1r blocker,

Gur (right). (B) Impulse frequencies in response to the concentration series of NMDG-Cl, NaCl,

or KCl before and after Gur treatment in WT mice. Responses to NMDG-gluconate are also

shown. The mean number of net impulses per 10 s (mean response) ± s.e.m. in Gur-sensitive

fibers (n = 5–6 from six mice). (C) Impulse frequencies in response to the concentration series

of NMDG-Cl before and after Gur treatment were measured in T1r3-KO mice (n = 4–5 from

three mice). Responses to NMDG-gluconate are also shown. (D) Impulse frequencies to 20-

mM L-glutamine or 100-mM sucrose in the absence or presence of 10-mM NMDG-Cl before

and after Gur treatment. Responses to 20-mM L-glutamine or 100-mM sucrose by T1r3-KO

mouse are also shown. Values are mean ± s.e.m. (n = 3–5 from three mice each). *, **: paired

t-test; $P < 0.05$ (*) and < 0.01 (**). (E) Amount of fluid intake for water and 10-mM NMDG-

Cl in the two-bottle preference tests. Values are mean ± s.e.m. (n = nine mice) (F) NMDG-Cl

intake shown in (E) normalized to water intake (preference score) in the two-bottle preference

tests. A score > 50% indicates that the taste solution was preferred over water.

Next, to examine the physiological interaction between a canonical taste substance

for mouse T1r and Cl⁻, we recorded responses to 20-mM L-glutamine or 100-mM sucrose from

T1r1/T1r3- and T1r2/T1r3-expressing cells, respectively, with or without NMDG-Cl from the

same T1r-connecting single fibers (Fig 4A). The concentrations for the taste substances were

set to induce responses greater than the baseline but less than maximum. As shown in Figure

4D, the response to L-glutamine or sucrose increased significantly by adding 10-mM NMDG-

Cl (paired t-test, $t2 = 7.56$, $p = 0.017$ for L-glutamine and $t2 = 5.05$, $p = 0.037$ for sucrose). We

confirmed that these responses had been suppressed by a lingual treatment of Gur in the

presence ($t2 = 6.73$, $p = 0.021$ for L-glutamine and $t2 = 8.80$, $p = 0.013$ for sucrose) or absence

($t2 = 8.32$, $p = 0.014$ for L-glutamine and $t2 = 11.72$, $p = 0.007$ for sucrose) of NMDG-Cl to a

similar extent. Moreover, the responses to the mixtures did not differ significantly from the

summation of the responses to each solution ($t2 = 2.34$, $p = 0.145$ for L-glutamine and $t2 = 2.31$,

$p = 0.147$ for sucrose). The results suggest that the simultaneous binding of Cl⁻ and a canonical

taste substance, such as amino acids and sugars, to T1r do not cause synergistic responses.

Finally, we addressed whether T1r-mediated Cl⁻ responses observed in the taste

nerves involve taste perception. Thus far, most reported behavioral assays for examining the

gustatory detection of NaCl were performed with concentrations above ~30 mM, which can

induce ENaC-mediated responses. Nevertheless, a few reports have shown that NaCl solution

induced higher consumption relative to water even below 10-mM concentration (Dyr et al,

2014; Stewart *et al*, 1994), which is below the range inducing ENaC-mediated responses

(Chandrashekar *et al.*, 2010) but within what we observed in the Cl⁻-induced taste nerve

responses via T1r. For verification, we performed a mouse two-bottle choice test using NMDG-

Cl solution and analyzed the preference relative to water. The mouse preferred water containing

10-mM NMDG-Cl, which was abolished by the application of Gur (Fig 4E, F). These results

suggest that Cl⁻ is preferably perceived through taste signal transduction mediated by T1r.

Discussion

In this study, Cl⁻ is found to specifically interact with the LBD in T1r3, a common component

of sweet and umami taste receptors, and induces a conformational change in the receptor's LBD

region. The Cl⁻ binding to T1r in taste cells is likely further transmitted to the sweet taste

nervous system, resulting in a light yet preferable taste sensation. Since T1rs are conserved

across vertebrates and the Cl⁻-binding site is likely conserved among T1r3 in various organisms,

T1r-mediated Cl⁻ responses might be common in many animals, such as humans. Evidently,

the concentration range for the Cl⁻-induced conformational change of medaka T1r2a/T1r3LBD

and increase in murine sweet nerve impulses observed in this study (i.e., ≤ ~10 mM) agrees

with the NaCl concentration perceived as "sweet" by humans (Bartoshuk *et al.*, 1964;

Bartoshuk *et al.*, 1978; Cardello, 1979) (Fig 1). Additionally, the sweet sensation induced by

NaCl was reportedly suppressed by topical application of *Gymnema sylvestre* (Bartoshuk *et al.*,

1978), containing Gymnemic acids, which are specific inhibitors of human sweet taste receptor

T1r2/T1r3 (Sanematsu *et al*, 2014). Overall, these results agree with the involvement of T1rs

in the Cl⁻-taste detection. These findings agree with an earlier hypothesis by Bartoshuk and

colleagues that dilute NaCl contains a sweet stimulus that interacts with the same receptor

molecules as sucrose (Bartoshuk *et al.*, 1978). The reported "sweet" sensation by low NaCl

concentration faded out at the concentrations detected as "salty" (Bartoshuk *et al.*, 1964;

Bartoshuk *et al.*, 1978; Cardello, 1979), thereby agreeing with the fact that we might be unaware

that table salt is sweet at such concentrations. The phenomenon could be explained by "mixture

suppression" (Bartoshuk, 1975; Keast & Breslin, 2002; Stevens, 1996), such that a light sweet

sensation is masked by an intense salty sensation in a higher concentration range of NaCl.

The Cl^- perception at low salt concentrations found in this study is achieved via the

T1rs-mediating taste system, which transduces information as a preferred taste by nature. Since

Cl^- is also a component of table salt, this system might serve as another pathway for preferred

salt perception promoting intake along with the pathway for Na^+ perception as a preferred taste

via taste cells that express the Na^+ receptor ENaC and a purinergic neurotransmission channel

CALHM1/3 (Nomura *et al*, 2020). In contrast, the T1r-mediated Cl^--sensing observed in this

study shows several different properties from those of cells or the molecules reportedly

exhibiting anion-sensitive Na^+ detection or Cl^- detection in taste buds thus far. Specifically, the

anion-sensitive Na^+-responding taste cells reported by Lewandowski *et al.* are type 3 cells,

which include sour- but not sweet-responding cells (Lewandowski *et al.*, 2016). The Cl^--

detection pathway reported by Roebber *et al.* was observed in type 2 cells, which include both

sweet- and bitter-responding cells, but was inhibited by a blocker of phospholipase C, which

mediates sweet- and bitter-signaling downstream of T1r and bitter receptors (Roebber *et al.*,

2019). The Cl^--responses mediated by TMC4 were observed in the glossopharyngeal nerve but

not in the chorda tympani nerve wherein T1r-mediated responses were observed (Kasahara *et*

al., 2021). Notably, responses by these reported pathways were tested by much higher

concentrations of NaCl, typically in the several hundred mM range (Kasahara *et al.*, 2021;

Lewandowski *et al.*, 2016; Roebber *et al.*, 2019), than those used in this study. These results

imply that there may be multiple distinct concentration-dependent pathways for Cl^- detection

in taste buds. Given that high NaCl concentration is transduced as an aversive taste through

bitter and sour taste cells (Oka *et al.*, 2013), the relevance between the reported high Cl^--

responsive pathways and high NaCl responses by bitter and sour cells is of interest, as pointed

out by the previous studies, and will require further investigation.

Salt taste sensation and natriuresis are critical physiological processes that regulate

sodium intake and excretion to maintain body fluid homeostasis. Intriguingly, both processes

were found to use the counter anion Cl^- to regulate the molecular functions of the receptors,

T1rs and ANPRs, which share a similar extracellular protein architecture with a conserved Cl^-

-binding site. In the case of ANPR, mGluRs, and CaSR, positive allosteric modulations by Cl^-

for agonist binding have been observed, with some variations in the extent of the enhancement

(Eriksen & Thomsen, 1995; Kuang & Hampson, 2006; Liu *et al.*, 2020; Tora *et al.*, 2018; Tora

et al., 2015). Whether the Cl^- actions on T1rs of other subtypes or from other organisms have

some variance is yet to be examined.

Materials and methods

Crystallography

All protein samples used for structural and functional analyses were prepared using Drosophila

S2 cells (Invitrogen) or high-expression clones for each protein sample established from S2

cells in previous studies (Nuemket *et al.*, 2017; Yamashita *et al*, 2017). No authentication or

test for mycoplasma contamination was performed.

The L-glutamine-bound T1r2a/T1r3LBD crystals, in complex with a crystallization

chaperone Fab16A, were prepared in the presence of NaCl as described (Nuemket *et al.*, 2017).

For the preparation of the Br^--substituted crystals, the obtained crystals were soaked in a mother

liquor consisting of 100 mM MES-Tris, pH 6.0, 50 mM NaBr, 17% PEG1500, 5% PEG400, 5

mM L-glutamine, 2 mM $CaCl_2$, cryoprotected by gradually increasing the concentration of

glycerol to 10%, incubated for 2 hours, and flash-frozen.

The X-ray diffraction data were collected at the SPring-8 beamline BL41XU using a

PILATUS6M detector (DECTRIS) at wavelength 0.9194 Å or at the Photon Factory beamline

BL-1A using an EIGER X4M detector (DECTRIS) at wavelength 2.7 Å. The data were

processed with XDS (Kabsch, 2010) (Table S1). The phases for anomalous difference Fourier

map calculation were obtained by molecular replacement methods with the program PHASER

(McCoy *et al*, 2007), using the structures of a single unit of the mfT1r2a-3LBD-Fab16A

complex (PDB ID: 5X2M; ligands and water models were removed) (Nuemket *et al.*, 2017) as

the search model (the R/R_{free} of the model for the Br^--data collected at 0.9194 Å: 0.248/0.350;

the Cl^--data collected at 2.7 Å: 0.233/0.339).

Differential scanning fluorimetry (DSF)

Differential scanning fluorimetry was performed as previously described (Yoshida *et al.*, 2019).

The purified mfT1R2a/3LBD heterodimer protein was prepared (Nango *et al.*, 2016) and

dialyzed with buffer A (20 mM HEPES-NaOH, 300 mM sodium gluconate, pH 7.5) to remove

Cl^-. 1 μg of the dialyzed protein sample was mixed with Protein Thermal Shift Dye (Applied

Biosystems) and 0.003 – 10 mM NaCl in 20 μL of buffer A. The mixture solutions were then

loaded to a MicroAmpR Fast Optical 48-Well Reaction Plate (Applied Biosystems).

Fluorescent intensities were measured by the StepOne Real-Time PCR System (Applied

Biosystems) while the temperature raised from 25 °C to 99 °C with a velocity of 0.022 °C/sec.

For detection, the reporter and quencher were set as "ROX" and "none," respectively. The

apparent melting transition temperature (T_m) was determined using the maximum of the

derivatives of the melt curve (dFluorescence/dT) by Protein Thermal Shift Software version 1.3

(Applied Biosystems). The apparent dissociation constant ($K_{d\text{-app}}$) for Cl^- derived from the T_m

values at different NaCl concentrations was estimated using a thermodynamic model proposed

by Schellman (Schellman, 1975) as described (Yoshida *et al.*, 2019). The sample sizes for the

analyses by DSF, as well as Förster resonance energy transfer and isothermal titration

calorimetry described below, were set to obtain reliable values based on the experiences in the

previous studies (Nango *et al.*, 2016; Nuemket *et al.*, 2017; Yoshida *et al.*, 2019).

Förster resonance energy transfer (FRET)

FRET analysis was performed as described previously (Nango *et al.*, 2016; Nuemket *et al.*,

2017). The WT T1r2aLBD-Cerulean and T1r3LBD-Venus fusion heterodimer proteins were

prepared as described (Nuemket *et al.*, 2017). For the mutant protein preparation, a T1r3-T105A

mutation was introduced into the vector pAc-mfT1r3aL-Ve (Nango *et al.*, 2016) using

polymerase chain reaction. The mutant expression vector was co-introduced with pAc-

mfT1r2aL-Ce (Nango *et al.*, 2016) to Drosophila S2 cells using Polyethyleneimine (PEI)

"MAX" (Polysciences) as previously described (Bleckmann *et al*, 2019) with a ratio of 0.5-µg

pAc-mfT1r2aL-Ce, 0.5-µg pAc- mfT1r3aL-Ve-T105A, and 10-µg PEI to ~1 × 10^6 cells.

Protein expression and purification were conducted similarly as for the WT protein.

436   For the Cl⁻-titration, the purified protein samples were dialyzed against buffer A in

the presence of 1-mM L-alanine. Afterward, the samples were diluted with buffer A to reduce

the remaining L-alanine concentration below 1 µM (< ~1/100 of EC_{50} (Nango *et al.*, 2016;

Nuemket *et al.*, 2017)), and then incubated in the presence of 0.001–10-mM NaCl or 1-mM L-

glutamine in buffer A at 4°C overnight. For L-glutamine titration in the presence or absence of

Cl⁻, the protein solution was dialyzed with buffer B (20-mM HEPES-Tris, 300-mM NaCl, pH

7.5) or buffer C (20-mM HEPES-Tris, 300-mM sodium gluconate, pH 7.5) in the presence of

1-mM L-alanine to prepare the conditions with or without Cl⁻, respectively. Then, the samples

were diluted with buffer B or C to reduce the remaining L-alanine concentration below 1-µM

and then incubated in the presence of 0.01–1000-µM L-glutamine at 4°C overnight.

Fluorescence spectra were recorded at 298 K with a FluoroMax4 spectrofluorometer (Horiba).

The sample was excited at 433 nm, and FRET was detected via the emission at 526 nm. The

emission at 475 nm was also recorded for the FRET index calculation. The FRET index

(intensity at 526 nm/intensity at 475 nm) was plotted against the Cl⁻ or L-glutamine

concentration, and the titration curves were fitted to the Hill equation using KaleidaGraph

(Synergy Software) or ORIGIN (OriginLab).

Isothermal titration calorimetry

In order to prepare the conditions with or without Cl⁻, the purified mfT1R2a/3LBD heterodimer

protein was dialyzed with buffers B or C, respectively. The dialyzed protein solution (~50 µM)

was then loaded into the sample cell in iTC200 (GE Healthcare) after the removal of insoluble

materials by centrifugation (10,000×g, 15 min, 277 K). The titration was performed by injecting

2 µL of 400 µM L-glutamine at intervals of 120 s at 298 K. The thermograms and the binding

isotherms were analyzed with Origin software, assuming one set of binding sites for fitting.

Fluorescence-detection size-exclusion chromatography-based thermostability assay

(FSEC-TS)

A T1r3-T105A mutation was introduced in the vector pAc_mfT1r3L (Yamashita *et al.*, 2017)

by PCR. The mutant expression vector was co-introduced with pAc_mft1r2aL (Yamashita *et*

al., 2017) to Drosophila S2 cells to establish a stable high-expression clone cell as previously

described (Yamashita *et al.*, 2017). The wild-type T1r2a/T1r3LBD and mutant T1r2a/T1r3-

105A-LBD proteins were expressed and purified as previously described (Nango *et al.*, 2016)

with several modifications as listed below. After the protein binding to ANTI-FLAG M2 affinity

gel, the resin was washed with either buffer D (20 mM HEPES-NaOH, 0.3 M NaCl, 2 mM

CaCl₂, 5 mM L-Gln, pH 7.5) or buffer E (20 mM HEPES-NaOH, 0.3 M Na gluconate, 2 mM

Ca gluconate, 5 mM L-Gln, pH 7.5). Then, the protein was eluted with 100 µg/mL FLAG

peptide in buffer D or E.

The protein solutions (50 µg/mL) in buffer D or E were incubated at 4 °C, 37 °C,

50 °C, 70 °C, or 90 °C at 2 hours. Subsequently, the samples were loaded on an SEC-5 column,

500 Å, 4.6 × 300 mm (Agilent) connected to a Prominence HPLC system (Shimadzu), using

buffer D or E as a running buffer at a flow rate of 0.3 ml min^{-1}. The elution profiles were

detected with an RF-20A fluorometer (Shimadzu), using excitation and emission wavelengths

of 280 and 340 nm for the detection of intrinsic tryptophan fluorescence.

The residual ratio after incubation at each temperature was estimated using the

fluorescence intensity at the elution peak, which corresponded to the T1rLBD dimer, *i.e.,* the

peak height at ~11.6 min. The values were normalized to the intensity of the sample incubated

at 4 °C as 1. In order to estimate the apparent melting temperature ($T_{\text{m-app}}$) of the sample, the

values of residual ratio at each temperature were fitted to the Gibbs-Helmholtz equation

transformed as shown below (assuming that the sample protein is under equilibrium between a

folding and unfolding state under each condition):

486
$$Residual\ ratio\ =\ 1 - \frac{1}{1 + exp\left[-\dfrac{\Delta H\left(1 + \frac{T}{T_{m-app}}\right) - \Delta C_P\left\{(T_{m-app} - \mathrm{T}) + T ln\left(\frac{T}{T_{m-app}}\right)\right\}}{RT}\right]}$$

where ΔH and ΔC_{p} are the enthalpy and heat capacity change of unfolding, respectively; T is

the temperature of the sample incubation; R is the gas constant. The fittings were performed

with KaleidaGraph (Synergy Software), with ΔH, ΔC_{p}, and $T_{\text{m-app}}$ are set as valuables.

Single fiber recording from mouse chorda tympani (CT) nerve

All animal experiments were conducted following the National Institutes of Health Guide for

the Care and Use of Laboratory Animals and approved by the committee for Laboratory Animal

Care and Use and the local ethics committee at Tokyo Dental College and Okayama University

Japan. The subjects were six adult male C57BL/6JCrj mice (Charles River Japan, Tokyo, Japan)

and four T1r3GFP-KO mice, which were obtained by mating T1r3-GFP (Damak *et al*, 2008)

and T1r3-KO (Damak *et al*, 2003) mice. Mice were maintained on a 12/12-h light/dark cycle

and fed standard rodent chow and 8–20 weeks of age ranging in weight from 20 g to 30 g.

The mice were anesthetized with an injection of sodium pentobarbital (40–50 mg/kg

ip) and maintained at a surgical level of anaesthesia, with additional injections (8–10 mg/kg ip

every hour). Under anaesthesia, each mouse was fixed in the supine position with a head holder,

and the trachea cannulated. The right CT nerve was dissected, free from surrounding tissues,

after the removal of the pterygoid muscle and cut at the point of its entry to the tympanic bulla.

A single or a few nerve fibers were teased apart with a pair of needles and lifted onto an Ag-

AgCl electrode, and an indifferent electrode was placed in nearby tissue. Their neural activities

were amplified (K-1; Iyodenshikagaku, Nagoya, Japan) and recorded on a computer using a

PowerLab system (PowerLab/sp4; AD Instruments, Bella Vista, NSW, Australia). For taste

stimulation of fungiform papillae, the anterior half of the tongue was enclosed in a flow

chamber. Taste solutions or rinses (distilled water) (~24 °C) were delivered to the tongue by

gravity flow at the same flow rate (~0.1 mL s^{-1}). For data analysis, we used the net average

frequency for 10 sec after the stimulus onset, which was obtained by subtracting the

spontaneous frequency for the 10 sec duration before stimulation from after stimulation. In the

initial survey to identify a nerve fiber connecting to T1r-expressing cells, test stimuli such as

100 mM NaCl, 10 mM HCl, 500 mM sucrose, 100 mM monopotassium glutamate, 20 mM

quinine HCl were separately applied. If the fiber responded to sucrose, we applied 10 μM–100

mM NMDG-Cl, NaCl, KCl, or either of 20 mM L-glutamine or 100 mM sucrose with or without

10 mM NMDG-Cl to the tongue. The criteria for the occurrence of response were the following:

the number of spikes was larger than the mean + two standard deviations of the spontaneous

discharge for three 10-s periods before stimulation, and at least three spikes were evoked by

taste stimulation (Yasumatsu *et al.*, 2012). In the case of T1r3GFP-KO mice, as the mice

showed a significant response to 0.5-M sucrose, we could identify sweet-responsive fibers

(impulse frequency of 13.4 ± 1.31) (Yasumatsu *et al.*, 2012). The reagents used were purchased

from Wako Pure Chemical Industries (Osaka, Japan; others). To block responses via T1r (Daly

et al., 2013; Margolskee *et al.*, 2007; Ninomiya & Imoto, 1995; Ninomiya *et al.*, 1999), each

tongue was treated with 30-µg ml^{-1} (~7 µM) gurmarin (Gur) dissolved in 5-mM phosphate

buffer (pH 6.8) for 10 min, similarly as described by Ninomiya & Imoto (Ninomiya & Imoto,

1995). To assure the detection of responses from T1r-expressing cells, recordings from Gur-

insensitive sweet-responsive fibers were defined as those retaining impulse frequencies to 0.5-

M sucrose more than 60% after the Gur treatment (Ninomiya *et al.*, 1999) were excluded from

the data. The number of Gur-sensitive and Gur-insensitive fibers were six and three,

respectively, among 0.5-M sucrose responding fibers. The sucrose application was repeated 3–

6 times during the recordings. Additionally, the recovery of the suppressed responses was

confirmed using 15-mM β-cyclodextrin, which could remove the effect of Gur from the tongue

(Ninomiya *et al.*, 1999). All Gur-sensitive fibers recovered up to 60%–150% of responses

before Gur. At the end of the experiment, animals were killed by administering an overdose of

the anaesthetic. Repeated measures ANOVA and Student's paired *t*-test were used to statistically

evaluate the effects of chemicals or gene deletion. The sample size was calculated according to

power analysis, thereby resulting in three per group due to the effect size (d) of 6–9.3 to detect

the effects of blocking or deleting T1r3.

Two-bottle preference tests

All training and testing sessions occurred during the light phase of the light/dark cycle. On the

first day of training, the WT mice (adult male C57BL/6JCrj) were water deprived for 23 h and

then placed in a test box with two bottles: one filled with water and the other empty. The amount

of fluid intake was measured after a 5-min presentation. After 4 days of training, mice were

used for test sessions if they drank water evenly on either side (nine mice). In the test sessions,

they were provided with two bottles, one containing 10-mM NMDG-Cl and the other

containing water, with or without Gur, for 5 min. The amount of liquid consumed was measured

by weighing the bottles, and a preference score of NMDG-Cl was calculated using the following

equation:

$$Preference\ score(\%) = \frac{V_{Cl}}{V_{Cl} + V_w} \times 100$$

where V_{cl} and V_w are the amount of NMDG-Cl intake and water intake, respectively.

Acknowledgments

We thank Drs. Kazuya Hasegawa, Nobuhiro Mizuno, Naohiro Matsugaki for help with X-ray

data collection; Junya Nitta and Hikaru Ishida for help with protein preparation; Ryusuke

Yoshida for help with the single fiber recording; Yuko Kusakabe for attempt at cell-based

receptor assay in the early stage of the study; Haruo Ogawa for sharing knowledge about ANPR;

Yuzo Ninomiya for valuable discussions. We also thank the reviewers of BioPhysics Colab for

their helpful comments and Enago (www.enago.jp) for the English language review. The

synchrotron radiation experiments at the BL41XU, SPring-8 were performed with approvals of

the Japan Synchrotron Radiation Research Institute (JASRI) (Proposal No. 2016B2534). The

synchrotron radiation experiment at the BL-1A, Photon Factory was supported by the Platform

for Drug Discovery, Informatics, and Structural Life Science (Proposal No. 1264). This work

was financially supported by JSPS KAKENHI Grant Numbers JP17H03644, JP18H04621,

JP20H03195, JP20H04778, JP21H05524 (to A.Y.) and JP20H03855 (to K.Y.), Mishima Kaiun

Memorial Foundation, and the Salt Science Research Foundation (Proposal No. 2039) (to A.Y.).

Competing interests

The authors declare that they have no conflict of interest.

References

Bartoshuk LM (1975) Taste mixtures: is mixture suppression related to compression? *Physiol Behav* 14: 643-649

Bartoshuk LM, McBurney DH, Pfaffmann C (1964) Taste of Sodium Chloride Solutions after Adaptation to Sodium Chloride: Implications for the "Water Taste". *Science* 143: 967-968

Bartoshuk LM, Murphy C, Cleveland CT (1978) Sweet taste of dilute NaCl: psychophysical evidence for a sweet stimulus. *Physiol Behav* 21: 609-613

Bleckmann M, Schurig M, Endres M, Samuels A, Gebauer D, Konisch N, van den Heuvel J (2019) Identifying parameters to improve the reproducibility of transient gene expression in High Five cells. *PLoS One* 14: e0217878

Cardello AV (1979) Taste quality changes as a function of salt concentration in single human taste papillae. *Chem Senses* 4: 1-13

Chandrashekar J, Kuhn C, Oka Y, Yarmolinsky DA, Hummler E, Ryba NJ, Zuker CS (2010) The cells and peripheral representation of sodium taste in mice. *Nature* 464: 297-301

Daly K, Al-Rammahi M, Moran A, Marcello M, Ninomiya Y, Shirazi-Beechey SP (2013) Sensing of amino acids by the gut-expressed taste receptor T1R1-T1R3 stimulates CCK secretion. *Am J Physiol Gastrointest Liver Physiol* 304: G271-282

Damak S, Mosinger B, Margolskee RF (2008) Transsynaptic transport of wheat germ agglutinin expressed in a subset of type II taste cells of transgenic mice. *BMC Neurosci* 9: 96

Damak S, Rong M, Yasumatsu K, Kokrashvili Z, Varadarajan V, Zou S, Jiang P, Ninomiya Y, Margolskee RF (2003) Detection of sweet and umami taste in the absence of taste receptor T1r3. *Science* 301: 850-853

Dyr W, Wyszogrodzka E, Mierzejewski P, Bienkowski P (2014) Drinking of flavored solutions by high preferring (WHP) and low preferring (WLP) alcohol-drinking rats. *Pharmacol Rep* 66: 28-33

Ellaithy A, Gonzalez-Maeso J, Logothetis DA, Levitz J (2020) Structural and Biophysical Mechanisms of Class C G Protein-Coupled Receptor Function. *Trends Biochem Sci* 45: 1049-1064

Eriksen L, Thomsen C (1995) [3H]-L-2-amino-4-phosphonobutyrate labels a metabotropic glutamate receptor, mGluR4a. *Br J Pharmacol* 116: 3279-3287

Hattori M, Hibbs RE, Gouaux E (2012) A fluorescence-detection size-exclusion chromatography-based thermostability assay for membrane protein precrystallization screening. *Structure* 20: 1293-1299

Kabsch W (2010) Xds. *Acta Crystallogr D Biol Crystallogr* 66: 125-132

Kasahara Y, Narukawa M, Ishimaru Y, Kanda S, Umatani C, Takayama Y, Tominaga M, Oka Y,

Kondo K, Kondo T *et al* (2021) TMC4 is a novel chloride channel involved in high-
concentration salt taste sensation. *J Physiol Sci* 71: 23

Keast RS, Breslin PA (2002) An overview of binary taste–taste interactions. *Food Quality and*
Preference 14: 111-124

Koehl A, Hu H, Feng D, Sun B, Zhang Y, Robertson MJ, Chu M, Kobilka TS, Laeremans T,
Steyaert J *et al* (2019) Structural insights into the activation of metabotropic glutamate receptors.
Nature 566: 79-84

Kuang D, Hampson DR (2006) Ion dependence of ligand binding to metabotropic glutamate
receptors. *Biochem Biophys Res Commun* 345: 1-6

Kunishima N, Shimada Y, Tsuji Y, Sato T, Yamamoto M, Kumasaka T, Nakanishi S, Jingami H,
Morikawa K (2000) Structural basis of glutamate recognition by a dimeric metabotropic
glutamate receptor. *Nature* 407: 971-977

Lewandowski BC, Sukumaran SK, Margolskee RF, Bachmanov AA (2016) Amiloride-
Insensitive Salt Taste Is Mediated by Two Populations of Type III Taste Cells with Distinct
Transduction Mechanisms. *J Neurosci* 36: 1942-1953

Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E (2002) Human receptors for sweet and
umami taste. *Proc Natl Acad Sci U S A* 99: 4692-4696

Lin S, Han S, Cai X, Tan Q, Zhou K, Wang D, Wang X, Du J, Yi C, Chu X *et al* (2021) Structures
of Gi-bound metabotropic glutamate receptors mGlu2 and mGlu4. *Nature* 594: 583-588

Liu H, Yi P, Zhao W, Wu Y, Acher F, Pin JP, Liu J, Rondard P (2020) Illuminating the allosteric
modulation of the calcium-sensing receptor. *Proc Natl Acad Sci U S A* 117: 21711-21722

Margolskee RF, Dyer J, Kokrashvili Z, Salmon KS, Ilegems E, Daly K, Maillet EL, Ninomiya
Y, Mosinger B, Shirazi-Beechey SP (2007) T1R3 and gustducin in gut sense sugars to regulate
expression of Na+-glucose cotransporter 1. *Proc Natl Acad Sci U S A* 104: 15075-15080

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ (2007) Phaser
crystallographic software. *J Appl Crystallogr* 40: 658-674

Misono KS (2000) Atrial natriuretic factor binding to its receptor is dependent on chloride
concentration: A possible feedback-control mechanism in renal salt regulation. *Circ Res* 86:
1135-1139

Monn JA, Prieto L, Taboada L, Hao J, Reinhard MR, Henry SS, Beadle CD, Walton L, Man T,
Rudyk H *et al* (2015a) Synthesis and Pharmacological Characterization of C4-(Thiotriazolyl)-
substituted-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylates. Identification of
(1R,2S,4R,5R,6R)-2-Amino-4-(1H-1,2,4-triazol-3-ylsulfanyl)bicyclo[3.1.0]hexane-2, 6-
dicarboxylic Acid (LY2812223), a Highly Potent, Functionally Selective mGlu2 Receptor
Agonist. *J Med Chem* 58: 7526-7548

Monn JA, Prieto L, Taboada L, Pedregal C, Hao J, Reinhard MR, Henry SS, Goldsmith PJ,
Beadle CD, Walton L *et al* (2015b) Synthesis and pharmacological characterization of C4-
disubstituted analogs of 1S,2S,5R,6S-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylate:
identification of a potent, selective metabotropic glutamate receptor agonist and determination
of agonist-bound human mGlu2 and mGlu3 amino terminal domain structures. *J Med Chem*
58: 1776-1794

Nango E, Akiyama S, Maki-Yonekura S, Ashikawa Y, Kusakabe Y, Krayukhina E, Maruno T,
Uchiyama S, Nuemket N, Yonekura K *et al* (2016) Taste substance binding elicits
conformational change of taste receptor T1r heterodimer extracellular domains. *Sci Rep* 6:
25745

Nelson G, Chandrashekar J, Hoon MA, Feng L, Zhao G, Ryba NJ, Zuker CS (2002) An amino-
acid taste receptor. *Nature* 416: 199-202

Nelson G, Hoon MA, Chandrashekar J, Zhang Y, Ryba NJ, Zuker CS (2001) Mammalian sweet
taste receptors. *Cell* 106: 381-390

Ninomiya Y, Imoto T (1995) Gurmarin inhibition of sweet taste responses in mice. *Am J Physiol*
268: R1019-1025

Ninomiya Y, Imoto T, Sugimura T (1999) Sweet taste responses of mouse chorda tympani
neurons: existence of gurmarin-sensitive and -insensitive receptor components. *J Neurophysiol*
81: 3087-3091

Nomura K, Nakanishi M, Ishidate F, Iwata K, Taruno A (2020) All-Electrical Ca(2+)-
Independent Signal Transduction Mediates Attractive Sodium Taste in Taste Buds. *Neuron* 106:
816-829 e816

Nuemket N, Yasui N, Kusakabe Y, Nomura Y, Atsumi N, Akiyama S, Nango E, Kato Y, Kaneko
MK, Takagi J *et al* (2017) Structural basis for perception of diverse chemical substances by T1r
taste receptors. *Nat Commun* 8: 15530

Ogawa H, Qiu Y, Ogata CM, Misono KS (2004) Crystal structure of hormone-bound atrial
natriuretic peptide receptor extracellular domain: rotation mechanism for transmembrane signal
transduction. *J Biol Chem* 279: 28625-28631

Ogawa H, Qiu Y, Philo JS, Arakawa T, Ogata CM, Misono KS (2010) Reversibly bound
chloride in the atrial natriuretic peptide receptor hormone-binding domain: possible allosteric
regulation and a conserved structural motif for the chloride-binding site. *Protein Sci* 19: 544-
557

Oike H, Nagai T, Furuyama A, Okada S, Aihara Y, Ishimaru Y, Marui T, Matsumoto I, Misaka
T, Abe K (2007) Characterization of ligands for fish taste receptors. *J Neurosci* 27: 5584-5592

Oka Y, Butnaru M, von Buchholtz L, Ryba NJ, Zuker CS (2013) High salt recruits aversive

taste pathways. *Nature* 494: 472-475

Roebber JK, Roper SD, Chaudhari N (2019) The Role of the Anion in Salt (NaCl) Detection by

Mouse Taste Buds. *J Neurosci* 39: 6224-6232

Roper SD (2015) The taste of table salt. *Pflugers Arch* 467: 457-463

Sanematsu K, Kusakabe Y, Shigemura N, Hirokawa T, Nakamura S, Imoto T, Ninomiya Y

(2014) Molecular mechanisms for sweet-suppressing effect of gymnemic acids. *J Biol Chem*

289: 25711-25720

Schellman JA (1975) Macromolecular Binding. *Biopolymers* 14: 999-1018

Stevens JC (1996) Detection of tastes in mixture with other tastes: issues of masking and aging.

Chem Senses 21: 211-221

Stewart RB, Russell RN, Lumeng L, Li TK, Murphy JM (1994) Consumption of sweet, salty,

sour, and bitter solutions by selectively bred alcohol-preferring and alcohol-nonpreferring lines

of rats. *Alcohol Clin Exp Res* 18: 375-381

Taruno A, Nomura K, Kusakizako T, Ma Z, Nureki O, Foskett JK (2021) Taste transduction and

channel synapses in taste buds. *Pflugers Arch* 473: 3-13

Tora AS, Rovira X, Cao AM, Cabaye A, Olofsson L, Malhaire F, Scholler P, Baik H, Van

Eeckhaut A, Smolders I *et al* (2018) Chloride ions stabilize the glutamate-induced active state

of the metabotropic glutamate receptor 3. *Neuropharmacology* 140: 275-286

Tora AS, Rovira X, Dione I, Bertrand HO, Brabet I, De Koninck Y, Doyon N, Pin JP, Acher F,

Goudet C (2015) Allosteric modulation of metabotropic glutamate receptors by chloride ions.

FASEB J 29: 4174-4188

van den Akker F, Zhang X, Miyagi M, Huo X, Misono KS, Yee VC (2000) Structure of the

dimerized hormone-binding domain of a guanylyl-cyclase-coupled receptor. *Nature* 406: 101-

104

Yamashita A, Nango E, Ashikawa Y (2017) A large-scale expression strategy for multimeric

extracellular protein complexes using Drosophila S2 cells and its application to the recombinant

expression of heterodimeric ligand-binding domains of taste receptor. *Protein Sci* 26: 2291-

2301

Yasumatsu K, Ogiwara Y, Takai S, Yoshida R, Iwatsuki K, Torii K, Margolskee RF, Ninomiya

Y (2012) Umami taste in mice uses multiple receptors and transduction pathways. *J Physiol*

590: 1155-1170

Ye Q, Heck GL, DeSimone JA (1991) The anion paradox in sodium taste reception: resolution

by voltage-clamp studies. *Science* 254: 724-726

Yoshida T, Yasui N, Kusakabe Y, Ito C, Akamatsu M, Yamashita A (2019) Differential scanning

fluorimetric analysis of the amino-acid binding to taste receptor using a model receptor protein,

the ligand-binding domain of fish T1r2a/T1r3. *PLoS One* 14: e0218909

Zhang C, Zhang T, Zou J, Miller CL, Gorkhali R, Yang JY, Schilmiller A, Wang S, Huang K,

Brown EM *et al* (2016) Structural basis for regulation of human calcium-sensing receptor by

magnesium ions and an unexpected tryptophan derivative co-agonist. *Sci Adv* 2: e1600241

Zhang F, Klebansky B, Fine RM, Xu H, Pronin A, Liu H, Tachdjian C, Li X (2008) Molecular

mechanism for the umami taste synergism. *Proc Natl Acad Sci U S A* 105: 20930-20934