Genetic deletion of Kiaa1109 in mice leads to developmental defects at the neuromuscular junction

Yun Liu and Weichun Lin

Department of Neuroscience
UT Southwestern Medical Center
6000 Harry Hines Blvd
Dallas, TX 75390-9111
USA

Abstract

KIAA1109 (4932438A13Rik) is a novel gene linked to Alkuraya-Kucinska Syndrome, an autosomal recessive disorder with severe brain malformations and arthrogryposis in humans. The role of KIAA1109 in mammalian development and function is unknown. Here, we characterize mutant mice deficient in *Kiaa1109* (*Kiaa1109$^{-/-}$*). We report that *Kiaa1109$^{-/-}$* mice died during perinatal stages. These *Kiaa1109$^{-/-}$* embryos exhibited impaired intramuscular nerve growth and reduced sizes of the neuromuscular junction (NMJ) compared with their littermate controls. Electrophysiological analysis further revealed defects in neuromuscular synaptic transmission in *Kiaa1109$^{-/-}$* embryos. Notably, the frequency of spontaneous neurotransmitter release was markedly increased, whereas evoked neurotransmitter release and quantal content were reduced. Furthermore, neuromuscular synapses in *Kiaa1109$^{-/-}$* embryos failed to respond to a repetitive, low frequency stimulation (10Hz). These results demonstrate that KIAA1109 is required for survival in mice and for proper development and function of the NMJ.

Key words: KIAA1109; neuromuscular junction; synaptic transmission; mice

Significance Statement

This is the first report characterizing the phenotype of mutant mice deficient in KIAA1109 (4932438A13Rik), a novel gene in mammals. We show that KIAA1109 is required for survival in mice and that KIAA1109 plays important roles in normal development and function of the NMJ in mice.

Introduction

KIAA1109 is a novel, uncharacterized gene that has been linked to fetal defects and multiple diseases in humans. For example, loss-of-function mutations in *KIAA1109* result in miscarriages and perinatal deaths (Filatova et al., 2019). Homozygous or compound heterozygous mutations in *KIAA1109* are associated with Alkuraya-Kucinskas Syndrome – an autosomal recessive severe neurodevelopmental disorder characterized by developmental delay, severe brain malformations, arthrogryposis and clubfoot (Filatova et al., 2019; Gueneau et al., 2018; Kumar et al., 2020; Kvarnung et al., 2018; Meszarosova et al., 2020). In addition, the linkage of *KIAA1109* with genes *IL2* and *IL21* within chromosome 4q27 region has been implicated in increased susceptibility to autoimmune disorders such as type I diabetes, Crohn's disease, rheumatoid arthritis and celiac disease (Balasopoulou et al., 2016; Bouzid et al., 2014; Hollis-Moffatt et al., 2010; Louahchi et al., 2016; Teixeira et al., 2009; Zhernakova et al., 2007).

KIAA1109 was initially cloned from a size-fractionated adult human brain cDNA library (Kikuno et al., 1999) and was later mapped to the chromosome 4q27 region (van Heel et al., 2007). Based on its coding sequence, *KIAA1109* encodes for a large protein of 5005 amino acids. Despite its direct linkage to human diseases, its function is uncertain, and it has not yet been studied in the context of an intact mammal. In Zebrafish, *kiaa1109* gene knockdown results in hydrocephaly and abnormally curved notochords in fish embryos reminiscent of clinical features in Alkuraya-Kucinskas Syndrome in human patients (Gueneau et al., 2018). In *Drosophila*, loss of *Tweek* (an ortholog of KIAA1109 in flies) leads to lethality at larval stage –*Tweek* homozygous flies rarely survive to the adulthood; for those that do survive, the mutant flies exhibit seizure and severe neurological defects (Verstreken et al., 2009). In *Drosophila*, Tweek is involved in regulating membrane endocytosis and synaptic vesicle recycling (Verstreken et al., 2009) and in regulating the size of the NMJs via a phosphatidylinositol 4,5-biophosphate (PI(4,5)P2)- and WSP-dependent pathway (Khuong et al., 2010). In human patient-derived dermal fibroblasts carrying

compound heterozygous *KIAA1109* variants, endocytosis and endosome trafficking are also notably

compromised (Kane et al., 2019), suggesting the function of KIAA1109 in humans may overlap with the

function of its fly ortholog *Tweek*.

Here, we present the first study of the roles of KIAA1109 in development of the neuromuscular junction

by investigating mutant mice deficient in KIAA1109 (*Kiaa1109$^{-/-}$*). We found that ablating KIAA1109 in

mice led to perinatal lethality; no *Kiaa1109$^{-/-}$* mice survived to postnatal day 0 (P0). Furthermore,

Kiaa1109$^{-/-}$ embryos exhibited marked defects in motor innervation and neurotransmission at the NMJ.

The results suggest KIAA1109 plays key roles in the neurodevelopment and synaptic neurotransmission

at the mouse NMJ. This study provides novel evidence in terms of the function of KIAA1109 in

mammals.

Results

KIAA1109 deficiency leads to perinatal lethality in mice

The heterozygous mice (*Kiaa1109$^{+/-}$*) were viable and fertile. These *Kiaa1109$^{+/-}$* mice were interbred to

generate *Kiaa1109$^{-/-}$* mice. A total of 9 litters (48 mice) were genotyped at postnatal day 21 (P21) – all

were identified as either wild type (+/+) or heterozygote (+/-), no homozygote *Kiaa1109$^{-/-}$* mice were

identified at P21. We therefore turned our attention to newborn pups and we noticed that several pups (5

pups in 6 litters) died shortly after birth. These dead pups were typically smaller than their littermates and

were later identified as *Kiaa1109$^{-/-}$* by genotyping. Since respiratory failure is a common cause for

neonatal death, we collected lung tissues from pups at P0. We performed a lung floatation assay using a

small piece of lung tissue and noticed that the lung tissue from the mutant pups sank to the bottom in

PBS, whereas the lung tissue from the control pups (*Kiaa1109$^{+/+}$*) remained afloat (Fig. 1 B), suggesting

that the mutant lungs were not inflated by the air. This was further confirmed by histological analysis of the lung sections – the alveoli in the WT were fully expanded whereas the alveoli in the mutant remained unexpanded (Fig.1 C). These observations suggest that $Kiaa1109^{-/-}$ mice likely died from respiratory failure.

This perinatal lethality precluded further analyses of $Kiaa1109^{-/-}$ mice at postnatal stages. We therefore focused our subsequent studies on embryonic stages. We collected 107 embryos (at E18.5) from 16 litters (Table 1). Among these embryos, 16 embryos (5 dead, 11 alive) were identified as $Kiaa1109^{-/-}$ at E18.5. Again, these $Kiaa1109^{-/-}$ embryos were noticeably smaller than their littermates (Fig. 1 A). In addition, $Kiaa1109^{-/-}$ embryos typically appeared hunchback with wrist dropping (Fig. 1 A); they exhibited little spontaneous movements but were able to respond to a mild pinch on their skin or tails. Table 1 summarizes total number of embryos at E14.5, E16.5 and E18.5. The numbers in parentheses are the numbers of embryos predicted based on the Mendelian ratio (1:2:1). The observed numbers of $Kiaa1109^{-/-}$ embryos at E14.5 and at E16.5 were fewer than those predicted, although the differences did not reach statistical significance at these stages (E14.5-E16.5). However, at E18.5, the number of $Kiaa1109^{-/-}$ mice was significantly reduced ($\chi2 = 7.1245$, P = 0.0284). At P0, no live $Kiaa1109^{-/-}$ mice were observed. These data indicate that the majority of $Kiaa1109^{-/-}$ embryos died perinatally and that KIAA1109 is required for survival in mice.

Abnormal motor innervation in $Kiaa1109^{-/-}$ mice

The perinatal lethality in $Kiaa1109^{-/-}$ embryos prompted us to examine the formation and function of the NMJ, as the NMJ is crucial for survival. First, we analyzed the innervation pattern of embryonic diaphragm muscles by wholemount immunofluorescence staining using anti-syntaxin 1 antibodies. In

both of WT and *Kiaa1109*[-/-] mice, the diaphragm muscles were innervated (Fig. 2). However, compared to the complex nerve branching in the WT diaphragm, the intramuscular nerve branches in the *Kiaa1109*[-/-] diaphragm exhibited much less complexity. Using a method described previously (Liu et al., 2019), we detected a significant reduction in the number of nerve branches and innervation area in *Kiaa1109*[-/-] diaphragms at E14.5 (Fig. 2 A and C). A similar phenotype was also observed at E18.5 (Fig. 2 B and D).

To determine if acetylcholine receptors (AchR) clustered normally in the diaphragm muscles, we used Texas-Red conjugated α-bungoratoxin to label postsynaptic AChR clusters in embryonic diaphragm muscles. At E14.5, AChRs clusters were detected at the center region of the diaphragm muscles; the average width of the endplate band was 159.27 ± 6.43 μm in *Kiaa1109*[-/-] muscles (dorsal quadrant), comparable to the band width of 181.38 ± 16.99 μm in control muscles (Fig. 3 C). At E18.5, the average width of the endplate band was significantly reduced in *Kiaa1109*[-/-] muscles (250.75 ± 10.75 μm), compared with that of the controls (382.46 ± 8.4 μm) (Fig. 3 D).

This narrow distribution of the endplate band in *Kiaa1109*[-/-] muscles appeared consistent with their branching pattern of intramuscular nerves, which extend numerous fine branches (pre-terminal axons) to muscle fibers. Compared to controls, *Kiaa1109*[-/-] muscles exhibited fewer and shorter pre-terminal axons (Fig. 3 E). We measured the distance between nerve terminals and intramuscular nerve trunks from which the nerve terminals sprouted and generated scatter plots (Fig. 3 F). The average distance was significantly reduced in mutants (12.02 ± 0.19 μm), compared to that of the control (28.9 ± 2.89 μm) (Fig. 3 G), suggesting an insufficient nerve sprouting from nerve trunks prior to muscle fibers in *Kiaa1109*[-/-] mice.

To further analyze the development of the NMJ, we labeled diaphragm muscles with antibodies against synaptotagmin 2 (Syt2), choline transporter (CHT), or acetylcholinesterase (AChE) (Fig. 4). In both control and mutant muscles, nerve terminals were intensely labeled with anti-Syt2 or anti-CHT antibodies, and were juxtaposed to postsynaptic AChRs labeled with Texas-Red conjugated α-bungoratoxin (arrows in Fig. 4 A and B). Similarly, anti-AChE labeling was superimposed with α-bungoratoxin labeled endplates in both control and mutant muscles (arrows in Fig. 4 C). However, the sizes of endplates in mutants were significantly smaller compare with the control (Fig. 4 D).

Since defects in muscle development may impair the formation of the neuromuscular junction and retrogradely affect presynaptic nerve growth (Rodriguez Cruz et al., 2020; Sanes and Lichtman, 1999; Tintignac et al., 2015), we examined the skeletal muscle in *Kiaa1109*$^{-/-}$ mice. Cross-sections of steromastoid muscles at E18.5 revealed that muscle fiber sizes in mutants were comparable to the WT. Muscle nuclei were peripherally localized in both mutant and WT muscles (Fig. 1 D). Overall, no signs of developmental retardation, degeneration or atrophy were detected in the mutant muscles. Taken together, these results show that that KIAA1109 deficiency leads to insufficient nerve branching, narrower endplate bands and smaller endplates.

Increased spontaneous neurotransmitter release in *Kiaa1109*$^{-/-}$ mice

We next analyzed neuromuscular synaptic transmission in *Kiaa1109*$^{-/-}$ mice (E18.5). First, we monitored spontaneous neurotransmitter release by analyzing miniature endplate potentials (mEPPs) at resting membrane potentials. In control muscles, the average frequency of mEPPs was 1.64 ± 0.11 events/min (N = 6, n = 70) (Fig. 5 A and C), consistent with the mEPP frequencies observed in our previous studies (Liu et al., 2009; Liu et al., 2012). However, in *Kiaa1109*$^{-/-}$ muscles, a significant increase in mEPP frequency

was detected (2.39 ± 0.32 events/min, N = 3 mice, n = 33 cells) (Fig. 5 A and C). The average amplitude

of mEPPs in *Kiaa1109*$^{-/-}$ mice (2.48 ± 0.07 mV) was similar with that of the control (2.24 ± 0.22 mV)

(Fig. 5 B and D), suggesting the quantal size was not affected in *Kiaa1109*$^{-/-}$ mice. Furthermore, the

parameters of mEPP kinetics, the rise time and half-width were comparable to those of the control mice

(Fig. 5 E and F). These results indicate that KIAA1109 deficiency resulted in an increase in spontaneous

transmitter release at the NMJ and suggest that the changes in spontaneous release in mutant NMJs is

likely due to a presynaptic alteration.

Defects in evoked synaptic transmission in *Kiaa1109*$^{-/-}$ mice

To further examine activity-dependent neurotransmission in *Kiaa1109*$^{-/-}$ mice, we stimulated the phrenic

nerve with supra threshold stimuli (2 - 5V, 0.1ms) and recorded evoked responses from the diaphragm

muscle. Muscle action potentials (Fig. 6 C) were readily elicited by nerve stimulation and muscle

contractions were observed in both control and *Kiaa1109*$^{-/-}$ mice. However, muscle movements in

Kiaa1109$^{-/-}$ mice appeared uncoordinated and much slower and weaker, compared to those in control

mice. In order to obtain stable recordings of evoked responses, we applied 50 μM 3-(N-

butylethanimidoyl)-4-hydroxy-2H-chromen-2-one (BHC), a specific myosin inhibitor (Heredia et al.,

2016), in Ringer's solution to suppress muscle movements. In both control and mutant muscles, muscle

contraction was markedly suppressed after incubation in BHC for 30 minute – only slight muscle

contractions occasionally remained. Upon suppression of muscle contraction, we were able to record

stable EPPs, which appeared as either single-peak EPPs (Fig. 6 A) or complex EPPs (Fig. 6 B). To

quantitate the parameters of EPP amplitude and kinetics, we focused our analyses on the traces with

single-peak EPPs (Fig. 6 A). The average amplitude of EPPs in controls was 13.41 ± 0.45 mV (N = 6, n =

50); whereas the average amplitude of EPPs in *Kiaa1109*$^{-/-}$ mice was reduced by 35% to 8.76 ± 0.77 mV

(N = 3, n = 38) (Fig. 6 D). The quantal content of $Kiaa1109^{-/-}$ mice was 4.37 ± 0.46, significantly less than that of the control mice (6.62 ± 0.44) (Fig. 6 E). The other parameters of EPP kinetics such as the rise time and half-width were not affected in $Kiaa1109^{-/-}$ mice (Fig. 6 F and G). These results indicate that the evoked neurotransmitter release was compromised in $Kiaa1109^{-/-}$ mice.

To further analyze the neuromuscular transmission, we applied a train of 10 stimuli (1 second each, for 5 sweeps) to the phrenic nerves, and measured the corresponding EPPs in the diaphragm muscle fibers. In control mice, we recorded 85 muscle cells from 8 mice. 97.9% of the cells responded faithfully to each stimulus with an evoked response (EPP) and we observed only 2.1% of the cells exhibiting 2 failures during 5 sweeps. In $Kiaa1109^{-/-}$ mice, we recorded 38 cells from 3 mice. Only 54.7% of the cells faithfully responded to each stimulus, whereas 45.3% of the cells exhibited at least 1 or more (up to 18) neurotransmission failures during 5 sweeps (Figure 7). This result indicates that neuromuscular synapses in $Kiaa1109^{-/-}$ embryos failed to respond to a repetitive, low frequency stimulation (10Hz). Together, our electrophysiological analyses demonstrate that KIAA1109 is involved in regulating synaptic activity at mouse NMJs.

Discussion

In this study, we examined mutant mice deficient in KIAA1109. We observed that the loss of KIAA1109 leads to perinatal lethality. Some mutant embryos die in uterus but the majority survive throughout embryonic stages and die at birth. This is consistent with a previously report showing flies carrying homozygous mutations in *Tweek*, the fly ortholog of *KIAA1109*, die at larval stages (Verstreken et al., 2009). In humans, mutations in *KIAA1109* result in miscarriages and perinatal deaths (Gueneau et al., 2018; Kumar et al., 2020). We provide evidence that the evolutionarily conserved gene *KIAA1109* is

essential for embryonic development and survival. In addition, KIAA1109 has been reported to be ubiquitously expressed in human adult and fetal tissues, including skeletal muscles and neural tissues such as brain and spinal cord (Https://www.gtexportal.org/home/gene/KIAA1109#geneExpression). The ubiquitous expression of KIAA1109 is suggestive of its crucial roles in fundamental biological processes, which may explains the global developmental delays in human patients suffering with Alkuraya-Kucinskas syndrome (Gueneau et al., 2018).

The defects in nerve growth resulting from the loss of KIAA1109 is consistent with brain malformations associated with Alkuraya-Kucinskas Syndrome (Filatova et al., 2019; Gueneau et al., 2018). The reduced nerve branching and narrower innervation territory in *Kiaa1109* KO mice suggest that KIAA1109 is involved in nerve outgrowth during development. Mutant mice exhibit less nerve branching throughout embryonic development. Membrane trafficking and addition to the tip of growth cones are key events that lead to axon outgrowth and elongation and synapse formation (Bloom and Morgan, 2011; Futerman and Banker, 1996; Pfenninger, 2009; Rodemer et al., 2020). Studies on fibroblasts obtained from patients with compound heterozygous *KIAA1109* variants show abnormal endocytosis and endosomal recycling in patient fibroblasts, suggesting that KIAA1109 plays a role in membrane trafficking (Kane et al., 2019). Thus, it is plausible that KIAA1109 in neurons may play a role in axon growth and nerve branching by regulating membrane trafficking and recycling.

Electrophysiological studies reveal marked defects in neuromuscular synaptic transmission in *KIAA1109* mutant. These defects appear primarily as pre-synaptic in nature as demonstrated by the significant increase in spontaneous transmitter release (mEPP frequency) and decreased evoked transmitter release (EPP amplitude and quantal content). These phenotypes share a common feature with mutant mice

deficient in synaptic vesicle proteins such as synaptobrevin (Liu et al., 2011; Liu et al., 2019), SNAP25 (Washbourne et al., 2002), synaptotagmin 2 (Pang et al., 2006), and syntaxin 1B (Wu et al., 2015). For example, evoked synaptic transmission was significantly reduced in synaptobrevin 2 Knockout NMJs and completely abolished in mutant NMJs deficient in both synaptobrevin 1 (Syb1) and synaptobrevin 2 (Syb2). In contrast, spontaneous synaptic transmission was increased in the Syb2 mutant or Syb1/Syb2 double mutant NMJs (Liu et al., 2019). Since both spontaneous and evoked neurotransmission require the fusion of synaptic vesicles with the pre-synaptic membrane which leads to exocytosis and neurotransmitter release, the defects in neuromuscular synaptic transmission in *KIAA1109* mutant mice suggest that KIAA1109 may be involved, albeit indirectly, in vesicle exocytosis.

More importantly, the intermittent transmission failures in mutant NMJs in response to repetitive stimulation suggest that, in the absence of KIAA1109, neuromuscular synaptic transmission cannot be maintained at normal levels. This finding is consistent with the phenotype observed in *Tweek* mutants in Drosophila (Verstreken et al., 2009). The failure to maintain normal levels of neurotransmission could result from multiple factors including a smaller synaptic vesicle pool, compromised vesicle mobilization and defective membrane trafficking and recycling. Indeed, synaptic vesicle numbers at the NMJ and neuromuscular endocytosis is reduced in *Tweek* mutant flies (Verstreken et al., 2009). In addition, defective cargo sorting and abnormal endosomal trafficking have been reported in fibroblasts derived from patients with *KIAA1109* mutations (Kane et al., 2019). Since endosome trafficking is crucial for presynaptic membrane retrieval and vesicle recycling (Buckley et al., 2000; Chanaday et al., 2019; Morgan et al., 2013; Sudhof, 2000), it is plausible that KIAA1109 functions in mammals in a manner similar to Tweek in *Drosophila.* Nevertheless, the mechanisms whereby KIAA1109 regulates synaptic transmission require further investigation.

Methods and materials

Mice

Kiaa1109 mutant mice [B6N(Cg)-4932438A13Rik[tm1b(EUCOMM)Hmgu]/J, Stock No.026878] were

generated by the Knockout Mouse Project (KOMP) at The Jackson Laboratory (Bar Harbor, Maine,

USA) using embryonic stem cells provided by the International Knockout Mouse Consortium.

Heterozygotes (*Kiaa1109*$^{+/-}$) were viable and fertile. *Kiaa1109*$^{-/-}$ mice were obtained from interbreeding

of heterozygotes (time-mated). The day when a vaginal plug first appeared was designated as embryonic

(E) day 0.5. Embryos were collected by Cesarean section of anesthetized pregnant female mice.

Kiaa1109 mice were genotyped using the following PCR primers: wild-type allele forward GGG ATA

TGG CAG AGA AGC TG, reverse AAA ACA ATT GGC TTA GAG ACT TCA; mutant allele forward

CGG TCG CTA CCA TTA CCA GT, reverse GAC CAC ACA AAT CCC TTG GT.

All experimental protocols followed National Institutes of Health (NIH) guidelines and were approved by

the University of Texas Southwestern Institutional Animal Care and Use Committee.

Histology staining

Embryonic (E18.5) and postnatal (P0) tissues were fixed with 2% paraformaldehyde (PFA) at 4°C

overnight. Samples were washed with PBS and equilibrated with 30% sucrose. Samples were then

transversely sectioned at a thickness of 12 μm and sections were stained with 0.5% Cresyl Violet. Images

were acquired on an Olympus BX51 upright microscope with Nomarski optics.

Whole mount immunofluorescence staining

Immunofluorescence staining was carried out on whole mount muscles as previously described (Liu et al., 2008). Diaphragm muscles were dissected from mouse embryos (E14.5 - E18.5). Muscle samples were fixed in 2% paraformaldehyde (PFA) at 4°C overnight. Then samples were extensively washed with PBS and incubated with Texas-Red conjugated α-bungarotoxin (α-bgt) (2 nM, Molecular Probes) at room temperature for 30 minutes. Samples were then incubated with primary antibodies at 4°C overnight. Primary antibodies used were: syntaxin-1 (I379), synaptotagmin-2 (I735) (1:1000, gifts from Dr. Thomas Südhof, Stanford University School of Medicine, Palo Alto, CA) (Pang et al., 2006), choline transporter (1:1000, gift from Dr. Randy Blakely, Vanderbilt University School of Medicine, Nashville, TN) (Ferguson et al., 2004) and acetylcholinesterase (1:1000, gifts from Dr. Palmer Taylor, Skaggs School of Pharmacy & Pharmaceutical Sciences, UC San Diego, CA, USA). After three washes with PBS samples were then incubated with fluorescein isothiocyanate-conjugated (FITC) secondary antibodies (1:600) at 4°C overnight. Samples were washed in PBS and mounted in Vectashield mounting medium (H-1000, Vector Laboratories, Inc., Burlingame, CA, USA). Fluorescence images were acquired with a Zeiss LSM 880 confocal microscope.

Morphometric analysis

Nerve branching was quantified using the methods describe previously (Kaplan et al., 2018; Liu et al., 2019). Confocal images were captured on the right hemi-diaphragms of E14.5 and E18.5 embryos. A vertical straight line perpendicular to the long axis of the muscle was drawn at the center of the endplate band. Then, three straight lines in parallel with the first line were respectively drawn to the left and right sides of the first line at 100 μm interval. The numbers of intersections between nerve branches and the lines were manually counted. The width of the endplate band was measured from the right dorsal quadrant of E14.5 and E18.5 diaphragms using NIH Image J. First, the endplate band labeled with Texas-

Red conjugated α-bgt was manually contoured. Then, 50 straight lines in parallel with the long axis of muscle fibred were drawn evenly within the contoured area. The length of each straight line was measured and the average length of the 50 lines was calculated as the width of the endplate band. The sizes of individual endplates were measured from high power confocal images (63x, oil) by using NIH Image J. The distance of individual synapses to the intramuscular nerve trunk was measure based on high power confocal images (63x, oil) obtained from the right ventral region of the diaphragm muscles (E18.5). A straight line in parallel with the long axis of the nerve trunk was drawn at the central of the nerve and designated as the midline. Then a vertical straight line from the center of the endplate to the midline was drawn and the length of the line was measured as the distance of the synapse to the nerve trunk.

Electrophysiology

Intracellular recordings were carried out on diaphragm muscles from $Kiaa1109^{-/-}$ and littermate control (both $Kiaa1109^{+/+}$ and $Kiaa1109^{+/-}$) mice at E18.5 as previously described (Liu et al., 2008). Acutely isolated phrenic nerve-diaphragm muscles were mounted on a Sylgard coated dish and bathed in oxygenated (95% O_2, 5% CO_2) Ringer's solution (136.8 mM NaCl, 5 mM KCl, 12 mM $NaHCO_3$, 1 mM NaH_2PO_4, 1 mM $MgCl_2$, 2 mM $CaCl_2$, and 11 mM d-glucose, pH 7.3). The muscle samples were then incubated in Ringer's solution containing 50 μM 3-(N-butylethanimidoyl)-4-hydroxy-2H-chromen-2-one (BHC), a specific myosin inhibitor (Heredia et al., 2016), for 30 minutes prior to the recording in the same solution. To evoke synaptic transmission, the phrenic nerve was stimulated with supra threshold stimuli (2-5V, 0.1ms) via a glass suction electrode connected to an extracellular stimulator (SD9, Grass-Telefactor, West Warwick, RI). Endplate regions were identified using a water-immersion objective (Olympus BX51WI). Muscle membranes were penetrated with sharp microelectrodes (resistance 20–40

MΩ) filled with 2 M potassium citrate and 10 mM potassium chloride. Muscle membrane potentials were acquired with an intracellular amplifier (AxoClamp-2B) and digitized with Digidata 1332A (Molecular Devices, Sunnyvale, CA, USA). MEPPs were analyzed with Mini Analysis Program (Synaptosoft, Inc., Decatur, GA) and EPPs were analyzed with pClamp 10.7 (Molecular Devices, CA). The rise time of mEPPs or EPPs was calculated as the time taken by the membrane potential to rise from 10% to 90% of the peak values of mEPPs or EPPs. Quantal content, defined as the number of acetylcholine quanta released in response to a single nerve impulse, was estimated by using a direct method: dividing the mean amplitude of EPPs by the mean amplitude of mEPPs of the same cell (Boyd and Martin, 1956; Hubbard et al., 1969; Wood and Slater, 2001).

Data analysis

Data were presented as mean ± standard error of the mean (SEM). Statistical analyses were carried out using SigmaPlot or Microsoft Excel software. Chi-square test was used to analyze the survival of mice. For quantitative morphometric analyses, student paired *t*-test was used to compared nerve branching between mutant and control groups. Student *t*-test was used to determine the difference of endplate band width and endplate size between mutant and control groups. For electrophysiological analyses, the mEPP frequency, amplitude, mEPP rise time (10%~90%), half width and EPP amplitude, quantal content, EPP rise time (10%~90%) and half width were compared between mutant and control groups by using student *t*-test. The difference between mutant and control groups is considered statistically significant if the P-value is less than 0.05.

Acknowledgments

We would like to thank Ms Qiaohong Ye for her excellent technical assistant, and Drs. Beverly Rothermel, Jane Johnson and Joseph McArdle for their comments on earlier drafts of the manuscript. This work was supported by grants from NIH/NINDS (R01 NS055028), the Edward Mallinckrodt, Jr. Foundation, and the Cain Foundation in Medical Research.

Declaration of Interests

The authors declare no competing interests.

Table 1. A summary of *Kiaa1109* mouse survival numbers during embryonic and perinatal stages

Age	Litter	Live Mouse	+/+	+/–	–/–	χ^2	P
E14.5	6	46	13 (11.5)	26 (23)	7 (11.5)	1.5289	0.4656
E16.5	2	13	2 (3.25)	10 (6.5)	1 (3.25)	1.7294	0.4212
E18.5	16	102	32 (25.5)	59 (51)	11 ((25.5)	7.1245	0.0284
P0	6	23	7 (5.75)	16	0 (5.75)	6.6301	0.0363

The expected numbers indicated in the parentheses are based on the Mendelian ratio (1:2:1).

Figure legends

Figure 1. KIAA1109 deficiency leads to perinatal lethality in mice. A: External phenotype of *Kiaa1109*$^{-/-}$ and *Kiaa1109*$^{+/+}$ embryos at E18.5. Note the mutant embryo (–/–) is smaller and curved (hunchback), compared with its WT littermate (+/+). B: Lung floatation assay: two small pieces of lung tissue (one from WT, and the other from *Kiaa1109*$^{-/-}$ at P0) were placed in a beaker filled with PBS. The *Kiaa1109*$^{-/-}$ lung tissue sank to the bottom in PBS, whereas the WT lung tissue remained afloat. C: Histological analysis of the lung section (Cresyl Violet staining) shows that the alveoli (a) in WT mouse were fully expanded whereas the alveoli in *Kiaa1109*$^{-/-}$ mouse were compressed, indicating breathing failure in the mutant. D: Cross sections of steromastoid muscles show that the muscle fiber sizes were similar between WT and *Kiaa1109*$^{-/-}$ mice (E18.5). Arrowheads point to muscle nuclei peripherally localized in both mutant and WT muscle fibers. a: alveoli; b: bronchiole. Scale bars: C: 100 μm; D: 20 μm.

Figure 2. Aberrant innervation pattern in *Kiaa1109*$^{-/-}$ muscles. Hemi-diaphragm muscle (right side) at E14.5 (A) and E18.5 (B) were labeled by anti-syntaxin-1 antibodies for the visualization of the phrenic nerves. The innervation territories appear narrower in *Kiaa1109*$^{-/-}$ muscles, compared with the WT. C-D: Quantification shows that the numbers of nerve branches in *Kiaa1109*$^{-/-}$ embryos are significantly reduced at both E14.5 (P = 0.0012, student paired *t*-test) and E18.5 (P = 0.0014). Scale bars: A, 200 μm; B: 800 μm.

Figure 3. Reduced endplate band in *Kiaa1109*$^{-/-}$ muscles. Confocal images of the dorsal right quadrant of the diaphragm muscles labeled by Texas Red conjugated α-bungoratoxin at E14.5 (A) and E18.5 (B). The width of the endplate band is reduced in *Kiaa1109*$^{-/-}$ muscles compared to the WT. C-D: Quantification of the width of the endplate band at E14.5 (*Kiaa1109*$^{-/-}$: 159.27 ± 6.43 μm; control: 181.38 ± 16.99 μm) and E18.5 (*Kiaa1109*$^{-/-}$: 250.75 ± 10.75 μm, control: 382.46 ± 8.4 μm). E: High-power views of nerves and individual AChR clusters. F: Scatter plots display the distance of the nerve terminals to the midline of the intramuscular nerve trunk. G: Comparison of average distance of the nerve terminals to the nerve trunk in *Kiaa1109*$^{-/-}$ (12.02 ± 0.19 μm, N = 3 mice, n = 185 synapses) and the control (28.9 ± 2.89 μm, N = 3 mice, n = 157 synapses). Scale bars: A and C, 200 μm; E: 20 μm.

Figure 4. Reduced neuromuscular synapse sizes in *Kiaa1109*$^{-/-}$ muscles. A-C: High power images of the neuromuscular synapses doubly labeled with Texas Red conjugated α-bungoratoxin (red) and anti-synaptotagmin 2 (syt 2) (A), anti- choline transporter (CHT) (B), or anti- acetylcholinesterase (AChE) (C). Arrows point to the NMJs. D: Quantification of the size of the endplate. The endplate size was significantly reduced in mutant mice (52.53±5.52 μm^2, N = 3 mice, n = 481 endplates), compared to that of the control (66.52±2.24 μm^2, N = 3 mice, n = 363 endplates).

Figure 5. Spontaneous synaptic activity is increased at the NMJs in *Kiaa1109*$^{-/-}$ muscles. A: Sample mEPP traces of 2-minute continuous recordings in the diaphragm muscles from the control and *Kiaa1109*$^{-/-}$ embryos (E18.5). B: Scale-up examples of mEPPs. C-F: Average mEPP frequency (C), amplitude (D), rise time (E) and half-width (F) are shown in the bar graphs. Notably, the mEPP frequency was increased in *Kiaa1109*$^{-/-}$ mice (2.39 ± 0.32 events/min, N = 3 mice, n = 34 cells), compared with controls (1.647 ± 0.11 events/min, N = 6 mice, n = 70 cells). MEPP amplitudes were comparable between *Kiaa1109*$^{-/-}$ (2.24

± 0.22 mV) and control mice (2.48 ± 0.07 mV). The rise time (E) and half-width (F) are also similar

between *Kiaa1109*$^{-/-}$ and control (rise time: control, 5.28 ± 0.36 ms, *Kiaa1109*$^{-/-}$, 6.24 ± 0.61 ms; half-

width: control, 15.92 ± 0.75 ms; *Kiaa1109*$^{-/-}$, 15.99 ± 1.26 ms).

Figure 6. Evoked neurotransmitter release is reduced at the NMJs in *Kiaa1109*$^{-/-}$ muscles. A-B:

Representative traces of endplate potentials (EPPs) recorded from E18.5 diaphragm muscles. Arrows in B

point to EPPs with multiple peaks (complex EPPs). C: Examples of muscle action potentials. D: EPP

amplitude is significantly reduced in *Kiaa1109*$^{-/-}$ (8.76 ± 0.77 mV, N = 3 mice, n = 38 cells), compared

with controls (13.08 ± 0.49 mV, N = 6 mice, n = 50 cells). E: Quantal content is also significantly reduced

in *Kiaa1109*$^{-/-}$ mice (4.37 ± 0.46) compared with control (6.29 ± 0.37). F-G: Quantification of rise time

(F) and half-width (G) between *Kiaa1109*$^{-/-}$ and controls (rise time: control, 5.25 ± 0.34 ms; *Kiaa1109*$^{-/-}$,

4.74 ± 0.44 ms; half-width: control, 29.67 ± 2.16 ms; *Kiaa1109*$^{-/-}$, 29.41 ± 3 ms).

Figure 7. Intermittent synaptic transmission failure in *Kiaa1109*$^{-/-}$ muscles. A-B: Sample traces of

EPPs in diaphragm muscle (E18.5) responding to 1-second train of supra-threshold stimuli (10 Hz)

applied to the phrenic nerve. Such a stain of stimuli evoked a train of 10 EPPs, corresponding to the

frequency of the stimulation (10Hz) in control embryos (A). In contrast, the NMJs in *Kiaa1109*$^{-/-}$ embryos

failed to follow – the absence of EPPs occurred following a train of stimulation (arrowheads point to the

stimulus artifacts). C: Quantification of the percentages of cells with success or failures in responding to

the repetitive stimuli. In controls, 97.9% of the cells responded to every stimulus (N = 8 mice, n = 85

cells). In contrast, only 54.7% of the mutant cells responded to every stimuli, and 45.3% exhibited

intermittent synaptic transmission failures (N = 3 mice, n = 38 cells).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Key reagents

Reagent or resource	Source	Identifier
Antibodies		
Rabbit polyclonal anti-syntaxin	Dr. Thomas Südhof, Stanford University School of Medicine, Palo Alto, CA	I379
Rabbit polyclonal anti-synaptotagmin-2	Dr. Thomas Südhof, Stanford University School of Medicine, Palo Alto, CA	I735
Rabbit polyclonal anti-acetylcholinesterase	Dr. Palmer Taylor Skaggs School of Pharmacy & Pharmaceutical Sciences, UC San Diego, CA	
Rabbit polyclonal anti-choline transporter	Dr. Randy Blakely, Vanderbilt University School of Medicine, Nashville, TN	
Fluorescein(FITC)-Conjugated AffiniPure Goat Anti-Rabbit IgG (H+L)	Jackson ImmunoResearch, West Grove, PA	Code# 111-095-144
Texas-Red conjugated α-bungarotoxin (α-bgt)	Molecular Probes (Thermo Fisher Scientific)	B-7489
Software		
pClamp 10.7	Molecular Devices, San Jose, CA	
Mini Analysis	Synaptosoft, Inc., Decatur, GA	
Image J	NIH	
SigmaPlot 11.0	SPSS Inc., Chicago, IL	

References

Balasopoulou, A., Stankovic, B., Panagiotara, A., Nikcevic, G., Peters, B.A., John, A., Mendrinou, E., Stratopoulos, A., Legaki, A.I., Stathakopoulou, V., Tsolia, A., Govaris, N., Govari, S., Zagoriti, Z., Poulas, K., Kanariou, M., Constantinidou, N., Krini, M., Spanou, K., Radlovic, N., Ali, B.R., Borg, J., Drmanac, R., Chrousos, G., Pavlovic, S., Roma, E., Zukic, B., Patrinos, G.P., Katsila, T., 2016. Novel genetic risk variants for pediatric celiac disease. Hum Genomics 10, 34.

Bloom, O.E., Morgan, J.R., 2011. Membrane trafficking events underlying axon repair, growth, and regeneration. Mol Cell Neurosci 48, 339-348.

Bouzid, D., Fourati, H., Amouri, A., Marques, I., Abida, O., Tahri, N., Penha-Goncalves, C., Masmoudi, H., 2014. Autoimmune diseases association study with the KIAA1109-IL2-IL21 region in a Tunisian population. Mol Biol Rep 41, 7133-7139.

Boyd, I.A., Martin, A.R., 1956. The end-plate potential in mammalian muscle. J Physiol 132, 74-91.

Buckley, K.M., Melikian, H.E., Provoda, C.J., Waring, M.T., 2000. Regulation of neuronal function by protein trafficking: a role for the endosomal pathway. J Physiol 525 Pt 1, 11-19.

Chanaday, N.L., Cousin, M.A., Milosevic, I., Watanabe, S., Morgan, J.R., 2019. The Synaptic Vesicle Cycle Revisited: New Insights into the Modes and Mechanisms. J Neurosci 39, 8209-8216.

Ferguson, S.M., Bazalakova, M., Savchenko, V., Tapia, J.C., Wright, J., Blakely, R.D., 2004. Lethal impairment of cholinergic neurotransmission in hemicholinium-3-sensitive choline transporter knockout mice. Proc Natl Acad Sci U S A 101, 8762-8767.

Filatova, A., Freire, V., Lozier, E., Konovalov, F., Bessonova, L., Iudina, E., Gnetetskaya, V., Kanivets, I., Korostelev, S., Skoblov, M., 2019. Novel KIAA1109 variants affecting splicing in a Russian family with ALKURAYA-KUCINSKAS syndrome. Clin Genet 95, 440-441.

Futerman, A.H., Banker, G.A., 1996. The economics of neurite outgrowth--the addition of new membrane to growing axons. Trends Neurosci 19, 144-149.

Gueneau, L., Fish, R.J., Shamseldin, H.E., Voisin, N., Tran Mau-Them, F., Preiksaitiene, E., Monroe, G.R., Lai, A., Putoux, A., Allias, F., Ambusaidi, Q., Ambrozaityte, L., Cimbalistiene, L., Delafontaine, J., Guex, N., Hashem, M., Kurdi, W., Jamuar, S.S., Ying, L.J., Bonnard, C., Pippucci, T., Pradervand, S., Roechert, B., van Hasselt, P.M., Wiederkehr, M., Wright, C.F., Study, D.D.D., Xenarios, I., van Haaften, G., Shaw-Smith, C., Schindewolf, E.M., Neerman-Arbez, M., Sanlaville, D., Lesca, G., Guibaud, L., Reversade, B., Chelly, J., Kucinskas, V., Alkuraya, F.S., Reymond, A., 2018. KIAA1109 Variants Are Associated with a Severe Disorder of Brain Development and Arthrogryposis. Am J Hum Genet 102, 116-132.

Heredia, D.J., Schubert, D., Maligireddy, S., Hennig, G.W., Gould, T.W., 2016. A Novel Striated Muscle-Specific Myosin-Blocking Drug for the Study of Neuromuscular Physiology. Front Cell Neurosci 10, 276.

Hollis-Moffatt, J.E., Gearry, R.B., Barclay, M.L., Merriman, T.R., Roberts, R.L., 2010. Consolidation of evidence for association of the KIAA1109-TENR-IL2-IL21 rs6822844 variant with Crohn's disease. Am J Gastroenterol 105, 1204-1205; author reply 1206-1207.

Hubbard, J.I., Llinas, R., Quastel, D.M.J., 1969. Investigation of Presynaptic Function. The Williams & Wilkins Company, Baltimore.

Kane, M.S., Diamonstein, C.J., Hauser, N., Deeken, J.F., Niederhuber, J.E., Vilboux, T., 2019. Endosomal trafficking defects in patient cells with KIAA1109 biallelic variants. Genes Dis 6, 56-67.

Kaplan, M.M., Sultana, N., Benedetti, A., Obermair, G.J., Linde, N.F., Papadopoulos, S., Dayal, A., Grabner, M., Flucher, B.E., 2018. Calcium Influx and Release Cooperatively Regulate AChR Patterning and Motor Axon Outgrowth during Neuromuscular Junction Formation. Cell Rep 23, 3891-3904.

Khuong, T.M., Habets, R.L., Slabbaert, J.R., Verstreken, P., 2010. WASP is activated by phosphatidylinositol-4,5-bisphosphate to restrict synapse growth in a pathway parallel to bone morphogenetic protein signaling. Proc Natl Acad Sci U S A 107, 17379-17384.

Kikuno, R., Nagase, T., Ishikawa, K., Hirosawa, M., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N., Ohara, O., 1999. Prediction of the coding sequences of unidentified human genes. XIV. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. DNA Res 6, 197-205.

Kumar, K., Bellad, A., Prasad, P., Girimaji, S.C., Muthusamy, B., 2020. KIAA1109 gene mutation in surviving patients with Alkuraya-Kucinskas syndrome: a review of literature. BMC Med Genet 21, 136.

Kvarnung, M., Taylan, F., Nilsson, D., Anderlid, B.M., Malmgren, H., Lagerstedt-Robinson, K., Holmberg, E., Burstedt, M., Nordenskjold, M., Nordgren, A., Lundberg, E.S., 2018. Genomic screening in rare disorders: New mutations and phenotypes, highlighting ALG14 as a novel cause of severe intellectual disability. Clin Genet 94, 528-537.

Liu, Y., Oppenheim, R.W., Sugiura, Y., Lin, W., 2009. Abnormal development of the neuromuscular junction in Nedd4-deficient mice. Dev Biol 330, 153-166.

Liu, Y., Padgett, D., Takahashi, M., Li, H., Sayeed, A., Teichert, R.W., Olivera, B.M., McArdle, J.J., Green, W.N., Lin, W., 2008. Essential roles of the acetylcholine receptor gamma-subunit in neuromuscular synaptic patterning. Development 135, 1957-1967.

Liu, Y., Sugiura, Y., Lin, W., 2011. The role of Synaptobrevin1/VAMP1 in Ca2+-triggered neurotransmitter release at the mouse neuromuscular junction. J Physiol 589, 1603-1618.

Liu, Y., Sugiura, Y., Sudhof, T.C., Lin, W., 2019. Ablation of All Synaptobrevin vSNAREs Blocks Evoked But Not Spontaneous Neurotransmitter Release at Neuromuscular Synapses. J Neurosci 39, 6049-6066.

Liu, Y., Sugiura, Y., Wu, F., Mi, W., Taketo, M.M., Cannon, S., Carroll, T., Lin, W., 2012. beta-Catenin stabilization in skeletal muscles, but not in motor neurons, leads to aberrant motor innervation of the muscle during neuromuscular development in mice. Dev Biol 366, 255-267.

Louahchi, S., Allam, I., Raaf, N., Berkani, L., Boucharef, A., Abdessemed, A., Khaldoun, N., Bahaz, N., Ladjouze-Rezig, A., Nebbab, A., Ghaffor, M., Djidjik, R., 2016. Association of rs6822844 within the KIAA1109/TENR/IL2/IL21 locus with rheumatoid arthritis in the Algerian population. HLA 87, 160-164.

Meszarosova, A.U., Lastuvkova, J., Rennerova, L., Hitka, P., Cihlar, F., Seeman, P., Safka Brozkova, D., 2020. Two novel pathogenic variants in KIAA1109 causing Alkuraya-Kucinskas syndrome in two Czech Roma brothers. Clin Dysmorphol 29, 197-201.

Morgan, J.R., Comstra, H.S., Cohen, M., Faundez, V., 2013. Presynaptic membrane retrieval and endosome biology: defining molecularly heterogeneous synaptic vesicles. Cold Spring Harb Perspect Biol 5, a016915.

Pang, Z.P., Melicoff, E., Padgett, D., Liu, Y., Teich, A.F., Dickey, B.F., Lin, W., Adachi, R., Sudhof, T.C., 2006. Synaptotagmin-2 is essential for survival and contributes to Ca2+ triggering of neurotransmitter release in central and neuromuscular synapses. J Neurosci 26, 13493-13504.

Pfenninger, K.H., 2009. Plasma membrane expansion: a neuron's Herculean task. Nat Rev Neurosci 10, 251-261.

Rodemer, W., Gallo, G., Selzer, M.E., 2020. Mechanisms of Axon Elongation Following CNS Injury: What Is Happening at the Axon Tip? Front Cell Neurosci 14, 177.

Rodriguez Cruz, P.M., Cossins, J., Beeson, D., Vincent, A., 2020. The Neuromuscular Junction in Health and Disease: Molecular Mechanisms Governing Synaptic Formation and Homeostasis. Front Mol Neurosci 13, 610964.

Sanes, J.R., Lichtman, J.W., 1999. Development of the vertebrate neuromuscular junction. Annu Rev Neurosci 22, 389-442.

Sudhof, T.C., 2000. The synaptic vesicle cycle revisited. Neuron 28, 317-320.

Teixeira, V.H., Pierlot, C., Migliorini, P., Balsa, A., Westhovens, R., Barrera, P., Alves, H., Vaz, C., Fernandes, M., Pascual-Salcedo, D., Bombardieri, S., Dequeker, J., Radstake, T.R., Van Riel, P., van de Putte, L., Lopes-Vaz, A., Bardin, T., Prum, B., Cornelis, F., Petit-Teixeira, E., European Consortium on

Rheumatoid Arthritis, F., 2009. Testing for the association of the KIAA1109/Tenr/IL2/IL21 gene region with rheumatoid arthritis in a European family-based study. Arthritis Res Ther 11, R45.

Tintignac, L.A., Brenner, H.R., Ruegg, M.A., 2015. Mechanisms Regulating Neuromuscular Junction Development and Function and Causes of Muscle Wasting. Physiol Rev 95, 809-852.

van Heel, D.A., Franke, L., Hunt, K.A., Gwilliam, R., Zhernakova, A., Inouye, M., Wapenaar, M.C., Barnardo, M.C., Bethel, G., Holmes, G.K., Feighery, C., Jewell, D., Kelleher, D., Kumar, P., Travis, S., Walters, J.R., Sanders, D.S., Howdle, P., Swift, J., Playford, R.J., McLaren, W.M., Mearin, M.L., Mulder, C.J., McManus, R., McGinnis, R., Cardon, L.R., Deloukas, P., Wijmenga, C., 2007. A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. Nat Genet 39, 827-829.

Verstreken, P., Ohyama, T., Haueter, C., Habets, R.L., Lin, Y.Q., Swan, L.E., Ly, C.V., Venken, K.J., De Camilli, P., Bellen, H.J., 2009. Tweek, an evolutionarily conserved protein, is required for synaptic vesicle recycling. Neuron 63, 203-215.

Washbourne, P., Thompson, P.M., Carta, M., Costa, E.T., Mathews, J.R., Lopez-Bendito, G., Molnar, Z., Becher, M.W., Valenzuela, C.F., Partridge, L.D., Wilson, M.C., 2002. Genetic ablation of the t-SNARE SNAP-25 distinguishes mechanisms of neuroexocytosis. Nat Neurosci 5, 19-26.

Wood, S.J., Slater, C.R., 2001. Safety factor at the neuromuscular junction. Prog Neurobiol 64, 393-429.

Wu, Y.J., Tejero, R., Arancillo, M., Vardar, G., Korotkova, T., Kintscher, M., Schmitz, D., Ponomarenko, A., Tabares, L., Rosenmund, C., 2015. Syntaxin 1B is important for mouse postnatal survival and proper synaptic function at the mouse neuromuscular junctions. J Neurophysiol 114, 2404-2417.

Zhernakova, A., Alizadeh, B.Z., Bevova, M., van Leeuwen, M.A., Coenen, M.J., Franke, B., Franke, L., Posthumus, M.D., van Heel, D.A., van der Steege, G., Radstake, T.R., Barrera, P., Roep, B.O., Koeleman, B.P., Wijmenga, C., 2007. Novel association in chromosome 4q27 region with rheumatoid arthritis and confirmation of type 1 diabetes point to a general risk locus for autoimmune diseases. Am J Hum Genet 81, 1284-1288.