Complex effects of chytrid parasites on the growth of the cyanobacterium *Planktothrix rubescens* across interacting temperature and light gradients

Joren Wierenga[1], Mridul K. Thomas[1], Ravi Ranjan[2,3], Bas W. Ibelings[1]

[1] Department F.A. Forel for Environmental and Aquatic Sciences and Institute for Environmental Sciences, University of Geneva, Geneva, Switzerland

[2] Helmholtz Institute of Functional Marine Biodiversity at the University of Oldenburg (HIFMB), Ammerländer Heerstraße 231, D-26129 Oldenburg, Germany

[3] Alfred-Wegener-Institute, Helmholtz-Centre for Polar and Marine Research (AWI), Am Handelshafen 12, 27570 Bremerhaven, Germany

Corresponding author:

Bas W. Ibelings

Dept. F.A. Forel for Environmental and Aquatic Sciences and Institute for Environmental Sciences, University of Geneva

Uni Carl Vogt (room C211)

66 Blvd Carl Vogt

Geneva 1205, Switzerland

Tel +41223790313

Email: Bastiaan.Ibelings@unige.ch

Competing Interests:

The authors declare no competing financial interests

Abstract

Chytrids are important drivers of aquatic ecosystems as phytoplankton parasites. The interaction between these parasites and their hosts are shaped by abiotic factors such as temperature and light. Here, we performed a full-factorial experiment to study how temperature and light interact to affect the dynamics of the bloom-forming toxic cyanobacterium *Planktothrix rubescens* and its chytrid parasite. We used a dynamic host-parasite model to explore how temperature and light affect long term dynamics. At low temperatures, chytrids do not survive. Higher light and temperature levels stimulated both phytoplankton and chytrid growth, with complex effects on their dynamics. Model exploration indicates that increasing temperature and light shifts equilibrium outcomes from *P. rubescens* persisting alone to stable coexistence and then to limit cycles. This provides an alternative biological explanation for why *P. rubescens* is mainly found in the relatively cold and dark lake metalimnion – it may enable avoidance of its parasite. Our study emphasizes the importance of investigating how abiotic factors interact with biotic interactions to drive complex outcomes.

Introduction

Cyanobacteria are important primary producers in aquatic systems [1] that can be harmful when they form dense blooms, which can sometimes contain high concentrations of toxins [2-4]. *Planktothrix rubescens* is a common cyanobacterium that often produces toxic blooms, with high concentrations of hepatotoxins like microcystins or neurotoxins like anatoxin-a [5, 6]. It is typically found in the metalimnion of deep stratifying lakes, and performs better at lower light and temperature conditions than other cyanobacteria [7 - 11]. It has an efficient light-harvesting complex composed of phycocyanin and phycoerythrin in addition to chlorophyll, which allows it to photosynthesize at these low light levels [12, 13]. But even though *P. rubescens* can grow at low light and temperature, its growth rate increases with temperature and light, and is highest at values well outside the range they typically experience in their natural habitats. Why *P. rubescens* is rarely seen in high-light, high-temperature conditions that would appear to favour its growth remains an open question. One possible explanations that we investigate here is that temperature and light shape biotic interactions such as parasitism.

Phytoplankton growth rates increase with light and temperature up to optimal values [14], and decrease thereafter – slowly for light, very rapidly for temperature i.e. they are oppositely skewed unimodal functions [15-17]. The optimal temperature and light conditions for

P. rubescens are poorly constrained based on existing experiments, but Oberhaus et al. (2007)

measured growth rates at 15 & 25 °C and showed that it was faster at 25 °C. This implies an

optimum temperature close to or above 25 °C because growth rate decreases rapidly above the

optimum temperature [17]. Similarly, Bright & Walsby (2000) and Oberhaus et al. (2007) found no

meaningful decrease in growth till the maximum irradiance levels they measured in white light (200

and 300 μE m^{-2} s^{-1}) [13, 18]. The optimum may be lower than this at 10-15 °C, but the measured

difference in growth rate was well within the range of experimental error and an optimum above

200 μE m^{-2} s^{-1} cannot be ruled out at these temperatures either. But in lakes, the species appears

to be most abundant at temperatures of 10-15 °C, although there is based on limited data;

abundance peaks from 6.5 to 20 °C have been seen [7, 19-22]. And it is often found at the depth

where irradiance is 0.1% – 1% of that at the surface (~ 1 – 20 μE m^{-2} s^{-1}). They maintain

themselves at this depth through light-mediated buoyancy regulation, with lift provided by

gas vesicles offset by carbohydrate ballast [23].

The dynamics of cyanobacteria are driven not just by abiotic factors such as temperature and

light, but also by biotic interactions with predators and parasites [24-26]. Parasites are

important but neglected biotic drivers of ecological dynamics, commonly affecting

cyanobacteria as well as other species of phytoplankton [27-30]. Parasites have been shown to

impact host populations of primary producers, with knock-on-effects at higher trophic levels

and community structure [31-35]. One of the most ubiquitous groups of aquatic parasites is the

Chytridiomycota, a large and diverse group of fungi that are best known for the role they are

playing in amphibian extinctions worldwide [36-38]. Chytrids have complex life cycles

(described in [39]) and also infect many species of algae and cyanobacteria, including

P. rubescens. They are often involved in the termination of blooms of cyanobacteria and other

phytoplankton [40-44]. Chytrids can act as a shunt in aquatic ecosystems, transferring

nutrients and carbon to higher trophic levels through infection of phytoplankton that are

otherwise resistant to grazing [45-47].

The chytrid parasite is also affected by temperature and light, not just through the direct

effects of temperature on its physiological processes, but also through temperature and light

effects on the host [9-11, 48, 49]. For example, light affects the release of dissolved organic

carbon (DOC) by phytoplankton [50, 51], which subsequently impacts chemotaxis of chytrid

zoospores, and thus infection dynamics [9, 11, 52, 53]. However, we know little about how light

and temperature interact to affect chytrid infections generally, and even less in the case of

P. rubescens. Much of the work on chytrid – host interactions has been done on the diatom

Asterionella formosa and its associated chytrids, *Rhizophydium planktonicum* and

Zygorhizidium planktonicum. In these host-parasite pairs, both low temperature and low light

can (independently) provide refuge for the host [9, 52]; 'refuge' in this case indicates a set of

environmental conditions that prevents chytrid infection and not a physical location.

Asterionella appears to have both a low-temperature and a high-temperature refuge,

indicating that the physiological tolerance range of chytrids is narrow relative to at least some

phytoplankton [48]. For *P. rubescens,* only low temperatures have been shown to offer a refuge

from chytrid infections with high temperature not investigated [10]. The closely-related

species *Planktothrix agardhii* does appear to have a high-temperature refuge; McKindles et al.

(2021a) found that chytrid infection rates (measured as % increase in filaments infected per

day) declined above approximately 22 °C to nearly zero by 29 °C [49]. This environmental

sensitivity of the chytrids has important consequences for *P. rubescens*, which is believed to

experience substantial variation in temperature and light due to its typical habitat in the

metalimnion. Understanding how the environment shapes the dynamics and location of *P.*

rubescens requires us to examine how these abiotic factors interact to shape its growth and

biotic interactions. The size and complexity of such experiments means that they are rarely

performed, weakening our ability to understand how environmental change affects natural

populations and communities.

Here we investigated how light and temperature interact to shape *Planktothrix*-chytrid

interactions using a combination of experiments and theory. We performed a full-factorial

experiment that features four temperatures, four light levels and two infection statuses

(infected and uninfected *P. rubescens* cultures). We extended a dynamic host-parasite model

[54-57] to explore the consequences of temperature-light interactions for *P. rubescens*-chytrid

dynamics. To examine this specific system, we made model parameters dependent on

temperature and light and estimated them from our experiment where possible. We

investigated whether the thermal refugium for *P. rubescens* depends on light intensity,

indicating that abiotic interactions shape these ecologically important biotic interactions. We

demonstrate that increasing light and temperature stimulated both phytoplankton and

chytrid growth, with complex effects on host-parasite interactions.

Materials and Methods

Experimental design

We used a full factorial experimental design with 4 temperatures (6, 11, 16 and 21 °C), 4 light

levels (2, 7, 14 and 21 µE m^{-2} s^{-1}) and 2 infection status levels (infected/uninfected *P. rubescens*

cultures), for a total of 32 treatment combinations. We used 4 replicates at each combination,

for a total of 128 experimental units. Fig. S1 shows a schematic illustrating the experimental

design.

Culture details and experimental conditions

Planktothrix rubescens strain NIVA-CYA98 and the chytrid parasite Chy-Kol2008

(Rhizophydiales) that were used in this study are monoclonal but non-axenic; bacterial

biomass was kept low by semi continuous growth, confirmed by microscopy. WC medium was

used to grow *P. rubescens* [58], modified to exclude silicon as a macronutrient.

Before the start of the experiment the *P. rubescen*s cultures were acclimatised to the different

experimental conditions. Temperature was regulated by using heated water baths in a

refrigerated room at 4 °C, logged with calibrated temperature loggers (HOBO Onset, UX120-

006M). Light was provided by fluorescent tubes (Osram neutral-white, 4000K), and light

intensities were manipulated using neutral density filters shading the culture flasks

(manufactured by Lee Filters, Hampshire UK, filters numbers 211, 210 and 209). Light

intensities were measured inside the shaded flasks (Licor LI-250A light meter, US-SQS/WB

quantum sensor). Photoperiod was set to a 16:8 hour cycle of light and dark. The experiment

was performed in 250 mL culture flasks (Greiner Bio One, Item No.: 658195). Cultures were

shaken daily during acclimatisation and during the experiment.

At the start of the experiment, the culture flasks were filled with 100 mL of *P. rubescens*

suspension, diluted to a biovolume of 4 nL mL^{-1}, so all treatments had the same density at the

start of the experiment. Subsequently, half of these flasks inoculated with a dense zoospore

suspension, constituting the infected treatment. The zoospore suspension was made by

filtering a heavily infected culture (16 °C, 11 µE m^{-2} s^{-1}) of *P. rubescens* through a glass

serological pipette filled with loosely packed glass fibers that had been autoclaved. The glass

fiber filter that is created this way lets through most of the zoospores but is highly efficient in

removing the filaments of *P. rubescens*. One mL of this dense zoospore suspension was added

to the infected treatment, resulting in 70 zoospores mL^{-1} at the start of the experiment. The

experiment lasted for 20 days.

Sampling and imaging

2 mL samples were taken every other day during the course of the experiment and fixed in

0.5% final concentration glutaraldehyde. The samples were stored in 2 mL sample tubes at 4

°C.

To analyze the samples from 128 experimental units, a high throughput biovolume

measurement method was developed (Wierenga et al., in prep). The method relies on high

resolution composite images of the entire surface of a clear bottom well, imaged by an

automated microscope. 300 μL of each sample was pipetted in duplicate into a clear bottom

96-well plate (Greiner Bio one Item No: 655096) for image analysis. The plates were put in the

fridge for 24 hours, to allow enough time for settling of the filaments. Using a Biotek cytation-

3, image composites were taken of the entire surface of each well bottom with a 4x

magnification objective. Images were taken in the chlorophyll and phycoerythrin

autofluorescence channel (586 nm excitation and 647 nm emission). This yields high quality

images with bright filaments against a dark background, ideal for image analysis. For infection

measurements 300 μL fixed sample was pipetted in duplicate into clear-bottom 96-well plates.

The plates were put in the fridge for 24 hours, to allow enough time for settling of the

filaments. To visualize the chytrid sporangia, 10 μL calcofluor-white (CFW) was added to each

well to get a 5 μg/mL final solution. To increase fluorescence output of the CFW stain, 10 μL of

sodium hydroxide (pH > 13.5) was added to increase the pH in the well. After incubation for 10

minutes, the plates were imaged with the Biotek cytation-3, in three different channels:

brightfield, chlorophyll and phycoerythrin autofluorescence (586 nm excitation, 647 nm

emission), and CFW (377 nm excitation, 447 nm emission). These composite images were

combined to generate images which clearly show the outline of the filaments and stained

sporangia. Furthermore, the viability of filaments can be assessed from the autofluorescence

signal in the images.

Image analysis

To measure the biovolume, the autofluorescence image composites were analyzed with the

open-source image analysis tool *ImageJ*, using the plugin *skeletonize*. In short, this method

uses a "Skeletonization algorithm" to compute the total length of all filaments in a picture.

These are subsequently converted to biovolume using the equation for calculating the volume

of a cylinder ($V = \pi r^2 l$), where r is the mean radius of filaments and l is the total length of all

filaments. The method is robust and can handle relatively high concentrations in samples

because overlapping filaments is not a big problem. It requires only limited human input,

reducing bias. Furthermore, with this method a relatively big sample of 300 μL is measured in

its entirety, compared to manual counting of only a fraction of that volume, thereby

improving precision.

To measure infections during the experiment, in the composite infection images a minimum

of 50 filaments were measured and marked as infected or uninfected. Instead of using the

frequency of infected and uninfected filaments to calculate the prevalence of infection, we

used the length of infected and uninfected filaments. This results in a better representation of

the prevalence of infection because there is a large variation in length, and moreover, infected

filaments are typically shorter than uninfected filaments (as also seen in [59]).

Data analysis

Data analysis and graphing was done using R version 4.0.4 and RStudio [60, 61]. ggplot2

(version 3.3.5) was used to make the graphs and the data processing was done with the help of

the tidyverse packages (version 1.3.1) [60-63]. Comparisons of means were done using the

robust yuen t-test from the package PairedData version 1.1.1 [64].

Dynamical model

We used a dynamical host – parasite model to assess the dependence of chytrid –

phytoplankton dynamics on temperature and light. We modified a model from Frenken et al.

(2020) [55] by incorporating temperature- and light-dependence of the parameters. The model

consists of three equations, which track the dynamics of uninfected hosts (H_u), infected hosts

(H_i) and free-swimming zoospores (Z).

The uninfected host H_u follows logistic growth with a carrying capacity K (nL ml^{-1}) and a

growth rate r (day^{-1}). Losses of uninfected host are defined by a density-independent mortality

rate: m_u (day^{-1}), and by infection of uninfected host by a zoospore, which converts uninfected

host into infected host. The infectivity constant I (mL cell^{-1}) is a measure for the infection

efficiency of zoospores (mL cell^{-1}).

$$\frac{dH_u}{dt} = rH_u\left(1 - \frac{H_u}{K}\right) - m_uH_u - IH_uZ \tag{1}$$

213   Infected hosts H_i increase with every uninfected host that gets infected by the attachment of a

214   zoospore. This is the same element as the loss due to infection for the uninfected host (IH_uZ).

215   Infected host decreases with a density-independent mortality rate m_i (day $^{-1}$) and with

216   maturation of sporangia given by the development time τ (days). The maturation of sporangia

217   assumes infected host H_i is converted into zoospores.

$$\frac{dH_i}{dt} = IH_uZ - \left(\frac{1}{\tau}\right)H_i - m_iH_i \tag{2}$$

218   Chytrid zoospores Z increase when sporangia develop with development time τ and release

219   new zoospores. The zoospore production parameter ρ (cells nL^{-1}) defines the number of spores

220   produced per biovolume of infected host. Spores are lost when they infect an uninfected host,
221   and additionally at a density-independent mortality rate m_z (day $^{-1}$).

$$\frac{dZ}{dt} = \left(\frac{p}{\tau}\right) H_i - IH_u Z - m_z Z \qquad (3)$$

223   *Model parameter values*: Some model parameters (Table 1) were calculated directly from our
224   experimental data: the infectivity parameter I and the growth rate r, making them functions
225   of light and temperature. The infectivity parameter I is calculated by assuming exponential
226   decay of H_u with Z, and is directly calculated from experimental data according to the
227   following equation:

$$I = -\frac{ln\left[\frac{H_u(t)}{H_u(t_0)}\right]}{Z(t_0)\, t} \qquad (4)$$

where t_0 is the beginning of the experiment and t is the length of the chosen period over which
I was calculated (2 days; see supplementary information for details). The phytoplankton
growth rate r was determined by fitting an exponential model to *P. rubescens* density over
time in the uninfected treatments. To run the dynamical model at a higher resolution for
temperature and light than the intervals used for the experiment, a generalized additive model
(GAM) was fitted to calculated values of r and I across the temperature and light levels used
in the experiment. This GAM-model was subsequently used in the dynamical host – parasite
model to generate values for r and I at interpolated temperature and light. The parameters K
and ρ are kept constant. The carrying capacity is set at 600 nL mL^{-1}, based on a logistic fit
through experimental data at high light and temperature. The production of zoospores is ρ
calculated from supplementary data from Frenken et al. (2020) [55], which has zoospore
production data on a very closely related strain of *P. rubescens* and chytrids. It is calculated by
dividing newly produced zoospores by the biovolume of the host 2 days prior, and is found to
be 20 cells nL^{-1}, which is set for each temperature and light combination. Infective lifetime of
zoospores was found to be 2.71 days at 17 °C. This translates into a mortality rate m_z of 0.37
(day^{-1}) which we set to exponentially increase with temperature. Development time for
sporangia is found to be between 2 and 3 days based on routine observations of infected
cultures and is set to exponentially decrease with temperature. The dynamical model was
analyzed in R using the R Package: deSolve, version 1.28 [65].

There is no straightforward way to calculate the mortality rates m_u and m_i from available data;

we therefore assumed m_u to be equal to 1% of the maximum growth rate at each temperature

(i.e. growth rate at the highest light level, 21 µE m⁻¹ s⁻¹). The mortality rate of infected

P. rubescens is based on the same fit, with a 50 percent higher dependency on temperature

and a higher initial value to account for the increased mortality due to infection. For more

details about this temperature and light dependence, please refer to the supplementary

information.

Table 1: Overview of all parameters and variables used in the dynamical model. Where
possible parameters were based on data from our experiment or from previous experiments
with closely related organisms. The dependence of the parameters on temperature, light or
both is indicated.

Symbol	Shape/Dependency	Meaning	Value/equation	Units
Variables				
H_u		Uninfected host		nL mL⁻¹
H_i		Infected host		nL mL⁻¹
Z		Free-swimming zoospores		cells mL⁻¹
Parameters				
K	Constant	Carrying capacity	600	nL mL⁻¹
I	GAM with temperature and light	Infectivity	GAM model	mL cell⁻¹ day⁻¹
r	GAM with temperature and light	Growth rate of the uninfected host	GAM model	day⁻¹
m_u	Exponential increase with temperature	Death rate of uninfected host	$m_u = 4.62 \cdot 10^{-4} \cdot e^{0.0783 \cdot T}$	day⁻¹
m_I	Exponential increase with temperature	Death rate of infected host	$m_i = 0.0305 \cdot e^{0.117 \cdot T}$	day⁻¹
p	Constant	Zoospore production per biovolume of infected host	20	cells nL⁻¹
m_Z	Exponential increase with temperature	Death rate zoospores	$m_z = 0.035 \cdot e^{0.128 \cdot T}$	day⁻¹
τ	Exponential decrease with temperature	Development time of zoospores in sporangia	$\tau = 16.42 \cdot e^{-0.099 \cdot T}$	Time in days

Results

Temperature and light effects on uninfected and infected Planktothrix growth rates

Temperature and light interacted to shape the growth of *P. rubescens* and its susceptibility to chytrid infections (Fig. 1, Table 1). The expected cold refuge [10] occurred in our experiments at temperatures of 11 °C and lower; no infections occurred at 11 °C and below. The growth rate of *P. rubescens* generally increased with both temperature and light at the levels tested in this experiment, but the shape of the interaction was complex. At low temperature, increasing light had little effect on growth. At low light, increasing temperature above 11 °C decreased growth rate (Fig. 1). Furthermore, the positive effect of temperature on the growth rate increased with light, whereas the positive effect of light on the growth rate, increased only till 14 µE m^{-2} s^{-1} (Fig. 1D, lines on top of each other).

Figure 1. *Planktothrix* growth rate varies strongly with temperature, light and infection status (**A**) Growth rate of uninfected and infected *P. rubescens* as a function of light at each of the 4 temperatures tested. The effect of the interaction between light and temperature is clearly visible both in infected and uninfected treatments. Infection reduces growth rate at 16 and 21 °C but not at 6 and 11 °C. Within each temperature, changing light has little effect on the difference in growth rate between infected and uninfected treatments. (**B**) The same data, viewed with temperature on X-axis and with light levels indicated by color. Within each light level, changing temperature has a large effect on the difference in growth rate between infected and uninfected treatments. In all panels dots represent mean growth rates and error bars represent the standard error of the mean.

Effects of chytrid infection on Planktothrix growth

Chytrids did not strongly affect the growth rate of *P. rubescens* at temperatures of 11 °C and lower. The growth rate remained essentially unchanged between the infected and uninfected treatment, confirming the refuge for *P. rubescens* at low temperatures (Fig. 1C). At temperatures of 16 °C and higher, the chytrid infection led to a substantially lower growth rate at all light levels tested. The reduction is similar across light levels (Fig. 1C). The biggest effect of the chytrid infection occurs at the highest temperature, with a big change in growth rate. This results in negative growth rates of *P. rubescens* at lower light levels (Fig. 1B, 1E). Growth of *P. rubescens* is very low at high temperature and low light (21 °C, 2 μE m^{-2} s^{-1}); infection led to further deterioration, resulting in negative growth rates for *P. rubescens* in the infected treatment (21 °C, at 2 and 7 μE m^{-2} s^{-1} and 16 °C at 2 μE m^{-2} s^{-1}).

Temperature and light effects on chytrid growth and infection prevalence

Temperature and light interact to shape chytrid growth, as indicated by the varying prevalence of infection on the last day of the experiment (Fig. 2A), and the growth and development of the infected and uninfected fractions of the *Planktothrix* biovolume (Fig. 1, 2B). Temperature is a major factor for chytrid growth showing increased prevalence of infection, especially at high light (Fig. 2A), and increased growth of infected BV with higher temperature (Fig. 2B). The effect of temperature is stronger at light. At the lowest light level (2 μE m^{-2} s^{-1}), the effect of temperature is absent, and the prevalence of infection is similar between 16 °C and 21 °C. Chytrid growth, as indicated by the growing fraction of infected *P. rubescens* biovolume, was only higher at 21 °C than at 16 °C with light levels at or above 14 μE m^{-2} s^{-1}. Furthermore, growth of uninfected biovolume at 16 °C was higher than at 21 °C even though growth of *P. rubescens* in the uninfected treatment is higher at 21 °C than at 16 °C (Fig. 2B). This indicates that at 21 °C *P. rubescens* is more vulnerable to chytrid infection than at 16 °C, especially at higher light levels. At 21 °C increased light also leads to a greater increase of the prevalence of infection compared to 16 °C. We did not find a low light refuge for *P. rubescens*. On the contrary, our experimental results show a large reduction in the growth rate of *P. rubescens* at low light with chytrid infection, especially at high temperatures (Fig. 1).

Figure 2. *P. rubescens* is affected more strongly by the chytrid parasite at 21 °C, especially at high light. (A) Prevalence of infection as a function of light level on the last day of the experiment. The prevalence of infection increases with increasing light at 21 °C, but not at 16 °C. At 6 and 11 °C no infected filaments were observed, and lines are not shown. (B) *P. rubescens* biovolume in the infected treatment, divided into uninfected and infected fractions; uninfected biovolume is shown in the left- and infected in the right-hand panels respectively. Asterisks indicate a statistically significant difference between means: * $p<0.05$, ** $p<0.01$

Dynamical model

We investigated how temperature and light would interact to drive *P. rubescens* - chytrid dynamics using a host – parasite model. We see a clear refugium defined largely by conditions of low temperature (Fig. 3A), identified by the absence of zoospores (Fig. 3B) and the distinct outcomes of the model (the 'only host' region in Fig. 3C). The low temperature refuge is explained by the absence of chytrid infections at low temperature. This refuge is also dependent on light level; between approximately 13 and 15 °C, a refuge exists at very low light levels and at high (for this experiment) light levels, while intermediate light levels allow for persistence of the parasite. The highest densities of *P. rubescens* are found in refuge conditions and decrease with increasing temperature and light (Fig. 3A). The chytrid abundance is at its highest at an intermediate temperature between 14 and 16 °C, and light levels above 10 µE m^{-2} s^{-1}.

Figure 3. Outcomes of modelled host-parasite dynamics in constant temperature and light
conditions from our theoretical model. Fig. 4 shows example time series from individual
points on the temperature-light surface. (**A**) Host biovolume is greatest at temperatures
below 13 °C and decreases above it. At high temperatures, host biovolume decreases with
increasing light. (**B**) Chytrid zoospore abundance is highest at intermediate temperature and
medium-to-high light levels. (**C**) The model shows 3 possible outcomes. The host persists
alone at low temperatures, it exhibits limit cycles at high temperature and high light, and it
coexists stably at intermediate conditions. Abundances shown in the limit cycle region are
averages over the course of one cycle. Note that the model is intended to explore qualitative
outcomes across parameter space and not to make precise quantitative predictions, for
which additional factors (such as nutrients) would need to be considered.

Three distinct outcomes are observed at the different combinations of temperature and light

we explored (Fig. 3C, Fig. 4). At low temperatures (< 13 °C), only the uninfected host can

survive irrespective of light levels (Fig. 3A). The host (uninfected and infected) and the chytrid

coexist stably at intermediate to high temperatures (Fig. 3C). However, the temperature range

for stable coexistence is drastically reduced at high light levels (Fig. 3C). Finally, at high

temperature and light levels, we find a limit cycle involving the uninfected host, infected host

and the chytrid.

Figure 4. Example dynamics from the *P. rubescens* – chytrid model (summarized in Fig. 3), at the parameter combinations we measured experimentally. The solid and dashed black lines indicate uninfected and infected host densities respectively, while the red line indicates the chytrid density. At low temperatures, *P. rubescens* has a thermal refuge and its density reaches the carrying capacity. With increasing light and temperature there is a shift from stable coexistence (16 °C with 7, 14, 21 µE m^{-2} s^{-1}, and 21 °C with 7 µE m^{-2} s^{-1}) to limit cycles (21 °C with 14, 21 µE m^{-2} s^{-1}).

Discussion

P. rubescens – chytrid growth and infection dynamics

We show that dynamics of *P. rubescens* and their chytrid parasites are heavily shaped by the interaction of temperature and light (Figs. 1-3). It has been previously shown that temperature and light are important factors [10, 11, 48, 49, 66, 67], but the effect of the interaction between temperature and light has not been well studied. To our knowledge, the only work on the interaction of temperature and light on chytrid-host dynamics is by Bruning [9], showing that in the diatom *Asterionella* and its chytrid parasite, that the two factors interact to affect infection dynamics. *P. rubescens* growth is most affected by chytrid infections at higher light and temperature levels. Even though we find relatively low infectivity of chytrids under low light, intrinsic slow growth due to low light means that chytrids can arrest *P. rubescens*

373 growth. We thus do not find a physiological refuge for *P. rubescens* at low light as we

374 hypothesized and was found for *Asterionella* [9].

Indirect light resource utilization by chytrids

The chytrids in this experiment are obligate parasites [68], feeding on the host as a resource,

and thus light is not a direct resource for the chytrids. Increased light does increase the release

of dissolved organic carbon (DOC) by phytoplankton, affecting the affinity of zoospores to

their host through chemotaxis and thus the interaction between the host and the parasite [9,

53, 69, 70]. However, here we find that at the lowest light level, increased temperature does

not lead to increased prevalence of infection, and that increased light leads to a very small

increase in the prevalence of infection at 16 °C. At 16 °C, temperature may be the limiting

factor for chytrid growth, and more light does not lead to increased chytrid growth. At 21 °C

however, the chytrid growth increases with light and thus might be resource-limited, with

increased photosynthetic resources available at higher light intensity indicating indirect light

utilization by acquiring energy resources from photosynthesis of the host while it is infected.

Optimizing resource utilization by feeding while the host is actively photosynthesizing makes

sense from an evolutionary perspective and is further suggested by the absence of a

temperature effect at the lowest light level. This would not be expected if the chytrid uses only

existing host resources at the time of infection for its reproduction and growth. Instead, it may

feed off the host while alive, as occurs in most parasites [71]. A recent study has shown that

increased light intensity and quality leads to bigger sporangia and increased chytrid

transmission [11], indicating that chytrids indirectly utilize light through host photosynthesis.

Another explanation could be that hosts grown under higher light stores more carbohydrates,

and as such are a better energy source for the chytrid parasite.

Dynamic model outcome and implications

The model shows three distinct outcomes over the range of conditions we explored: 1) only the

host survives. 2) stable coexistence between the parasite and the host. 3) limit cycles of host

and parasite. (Fig. 3C, Fig 4). Limit cycles occur at high temperatures and high light intensities

(Fig. 3C, Fig. 4). Under these conditions, *P. rubescens* can grow rapidly but is vulnerable to

infection, and infection under these conditions leads to a rapid decline of the host to very low

densities. In a natural setting, chytrids are found both at low densities with only minimal

impact on the host [72-74], as well as at high densities and high infection incidence that can

rapidly terminate host blooms [34, 44, 66, 75]. In natural conditions, it is possible that periods

of very low density could lead to local extinction of both host and parasite and so may not be

congenial to the persistence of this host-parasite system despite the model outcome of limit
cycles.

This suggests an interesting biotic alternative to abiotic factors explaining the niche of
P. rubescens, which is almost always found around the 0.1 – 1% light level in stratified lakes
[23]. The prevailing hypothesis is that they have adapted to a specific niche in the lake with
relatively high nutrient levels, lower temperatures, and low light, where they outcompete
other phytoplankton because they possess the accessory pigments phycoerythrin and
phycocyanin [13]. However, this is inconsistent with the experimental data showing that
growth rates of *P. rubescens* increase when temperature and light are increased beyond the
limiting conditions found at the bottom of the euphotic zone; light levels up to 300 μE m^{-2} s^{-1},
and temperatures up to at least 21 °C still promote growth [12, 13, 18, this study]. But at these
high temperature and light levels, *P. rubescens* may be very sensitive to chytrid infections, and
therefore unable to sustain growth, especially in competition with other phytoplankton
species. Chytrid infection might thus limit *P. rubescens* to regions of the water column that are
suboptimal for its growth, but safe from infection. If so, temperature, light and chytrids may
together determine the vertical distribution of *P. rubescens* in stratified lakes. The model
furthermore shows that the thermal refuge extends to higher temperatures at higher light
levels, although we do not have experimental data to confirm this. We also cannot say
whether a high-temperature refuge also exists as in the case of *P. agardhii* and *A. formosa* [48,
49]. Our explanation here for the *P. rubescens* vertical distribution would only hold if there
was no such high-temperature refuge.

Although we believe this model offers us useful insights into the *P. rubescens* vertical
distribution, its purpose is not to make precise quantitative predictions but instead to explore
how temperature-light interactions may affect host-parasite dynamics. To make quantitative
predictions, the model would at least need to account for several additional factors, including
nutrient dynamics, temperature & light variation, and the effects of sinking, none of which we
account for. Also, for some model parameter values, we used estimates based on the literature
instead of measurements from our experiments (Table 1). These estimates are based on
empirical data and physiological understanding, but there is unavoidable uncertainty about
their true values. The values we chose may have led to lower prevalence of infection in the
model (Fig. 3) than that seen in our experiments (Fig. 2), suggesting that some of these
parameters could be improved upon if a predictive model is developed in future.

Conclusion

We have shown that the host – parasite interaction of *P. rubescens* and chytrids is strongly affected by the interaction of temperature and light. Our results confirm the existence of a thermal refugium below at least 11 °C, and that this may vary with light intensity although there appears to be no specific low light refuge. In general, vulnerability of *P. rubescens* to chytrid infections increases with temperature and light. This may explain why *P. rubescens* is hardly ever found under warmer or high-light conditions, even though lab experiments show that these conditions lead to higher growth rates in the absence of chytrids. Our model does neglect some important complexities, such as the effects of nutrients on the biotic interaction, and the effects of light on host stoichiometry and consequently chytrid growth. Our results highlight how abiotic and biotic environmental factors can interact in complex ways to affect ecological dynamics.

Acknowledgements

The authors gratefully acknowledge Thomas Rohrlack for early discussions which greatly helped putting us in the research direction described here, and Thijs Frenken & Takeshi Miki for discussions of their dynamic model and calculation of model parameters.

Competing Interests

The authors declare no competing financial interests

Data availability statement

The datasets generated during and/or analysed during the current study will be made available in the Zenodo repository on acceptance of this manuscript.

References

1. Díez, B., & Ininbergs, K. (2013). Ecological importance of cyanobacteria. In *Cyanobacteria* (pp. 41–63). John Wiley & Sons, Ltd. https://doi.org/10.1002/9781118402238.ch3

2. Fristachi, A., Sinclair, J. L., Hall, S., Berkman, J. A. H., Boyer, G., Burkholder, J., Burns, J., Carmichael, W., Dufour, A., Frazier, W., Morton, S. L., O'Brien, E., & Walker, S. (2008). Occurrence of cyanobacterial harmful algal blooms workgroup report. *Advances in Experimental Medicine and Biology*, *619*(January), 45–103. https://doi.org/10.1007/978-0-387-75865-7_3

3. Huisman, J., Codd, G. A., Paerl, H. W., Ibelings, B. W., Verspagen, J. M. H., & Visser, P. M. (2018). Cyanobacterial blooms. *Nature Reviews Microbiology 2018 16:8*, *16*(8), 471–483. https://doi.org/10.1038/s41579-018-0040-1

4. Plaas, H. E., & Paerl, H. W. (2021). Toxic Cyanobacteria: A Growing Threat to Water and Air Quality. In *Environmental Science and Technology* (Vol. 55, Issue 1, pp. 44–64). American Chemical Society. https://doi.org/10.1021/acs.est.0c06653

5. Kurmayer, R., Deng, L., & Entfellner, E. (2016). Role of toxic and bioactive secondary metabolites in colonization and bloom formation by filamentous cyanobacteria Planktothrix. *Harmful Algae*, *54*, 69–86. https://doi.org/10.1016/j.hal.2016.01.004

6. Rohrlack, T., Christiansen, G., & Kurmayer, R. (2013). Putative antiparasite defensive system involving ribosomal and nonribosomal oligopeptides in cyanobacteria of the genus planktothrix. *Applied and Environmental Microbiology*, *79*(8), 2642–2647. https://doi.org/10.1128/AEM.03499-12

7. Legnani, E., Copetti, D., Oggioni, A., Tartari, G., Palumbo, M. T., & Morabito, G. (2005). Planktothrix rubescens' seasonal dynamics and vertical distribution in. *Journal of Limnology*, *64*(1), 61–73.

8. Walsby, A., Ng, G., Dunn, C., & Davis, P. A. (2004). Comparison of the depth where Planktothrix rubescens stratifies and the depth where the daily insolation supports its neutral buoyancy. *New Phytologist*, *162*(1), 133–145. https://doi.org/10.1111/j.1469-8137.2004.01020.x

9. Bruning, K. (1991a). Effects of temperature and light on the population dynamics of the Asterionella-Rhizophydium association. *Journal of Plankton Research*, *13*(4), 707–719. https://doi.org/10.1093/plankt/13.4.707

10. Rohrlack, T., Haande, S., Molversmyr, Å., & Kyle, M. (2015). *Environmental Conditions Determine the Course and Outcome of Phytoplankton Chytridiomycosis*. 1–17. https://doi.org/10.1371/journal.pone.0145559

11. Tao, Y., Wolinska, J., Hölker, F., & Agha, R. (2020). Light intensity and spectral distribution affect chytrid infection of cyanobacteria via modulation of host fitness. *Parasitology*, *147*(11), 1206–1215. https://doi.org/10.1017/S0031182020000931

12. Davis, P. A., & Walsby, A. (2002). Comparison of measured growth rates with those calculated from rates of photosynthesis in Planktothrix spp. isolated from Blelham Tarn, English Lake District. *New Phytologist*, *156*(2), 225–239. https://doi.org/10.1046/j.1469-8137.2002.00495.x

13. Oberhaus, L., Briand, J. F., Leboulanger, C., Jacquet, S., & Humbert, J. F. (2007). Comparative effects of the quality and quantity of light and temperature on the growth of Planktothrix agardhii and P. rubescens 1. *Journal of Phycology*, *43*(6), 1191–1199. https://doi.org/10.1111/j.1529-8817.2007.00414.x

14. Reynolds, C. S. (2009). Growth and replication of phytoplankton. In *The Ecology of Phytoplankton* (pp. 145–177). Cambridge University Press. https://doi.org/10.1017/CBO9780511542145.005

15. Litchman, E., & Klausmeier, C. A. (2008). Trait-based community ecology of
phytoplankton. *Annual Review of Ecology, Evolution, and Systematics*, 615-639.
16. Edwards, K. F., Thomas, M. K., Klausmeier, C. A., & Litchman, E. (2016). Phytoplankton growth
and the interaction of light and temperature: A synthesis at the species and community
level. *Limnology and Oceanography*, *61*(4), 1232-1244.
17. Thomas, M. K., Kremer, C. T., & Litchman, E. (2016). Environment and evolutionary history
determine the global biogeography of phytoplankton temperature traits. *Global Ecology and
Biogeography*, 25(1), 75–86. https://doi.org/10.1111/geb.12387
18. Bright, D. I., & Walsby, A. (2000). The daily integral of growth by Planktothrix rubescens
calculated from growth rate in culture and irradiance in Lake Zürich. *New Phytologist*, *146*(2),
301–316. https://doi.org/10.1046/j.1469-8137.2000.00640.x
19. Jann-Para, G., Schwob, I., & Feuillade, M. (2004). Occurrence of toxic Planktothrix rubescens
blooms in lake Nantua, France. *Toxicon*, 43(3), 279-285.
20. Jacquet, S., Briand, J. F., Leboulanger, C., Avois-Jacquet, C., Oberhaus, L., Tassin, B., Vinçon-Leite,
B., Paolini, G., Druart, J.-C., Anneville, O. & Humbert, J. -F. (2005). The proliferation of the toxic
cyanobacterium *Planktothrix rubescens* following restoration of the largest natural French lake
(Lac du Bourget). *Harmful Algae*, 4(4), 651-672.
21. Lenard, T. (2009). Metalimnetic bloom of *Planktothrix rubescens* in relation to environmental
conditions. *Oceanological and Hydrobiological Studies*, 38(2), 45-53.
22. Van den Wyngaert, S., Salcher, M. M., Pernthaler, J., Zeder, M., & Posch, T. (2011). Quantitative
dominance of seasonally persistent filamentous cyanobacteria (*Planktothrix rubescens*) in the
microbial assemblages of a temperate lake. *Limnology and Oceanography*, 56(1), 97-109.
23. Walsby, A. (2005). Stratification by cyanobacteria in lakes: A dynamic buoyancy model indicates
size limitations met by Planktothrix rubescens filaments. *New Phytologist*, *168*(2), 365–376.
https://doi.org/10.1111/j.1469-8137.2005.01508.x
24. Conroy, J. D., Kane, D. D., Quinlan, E. L., Edwards, W. J., & Culver, D. A. (2017). Abiotic and biotic
controls of phytoplankton biomass dynamics in a freshwater tributary, estuary, and large lake
ecosystem: Sandusky bay (lake erie) chemostat. *Inland Waters*, *7*(4), 473–492.
https://doi.org/10.1080/20442041.2017.1395142
25. Sommer, U., Maciej Gliwics, Z., Lampert, W., & Duncan, A. (1986). The PEG-model of seasonal
succession of planktonic events in fresh waters. *Archiv Für Hydrobiologie*, *106*(4), 433–471.
26. Sommer, U., Adrian, R., De Senerpont Domis, L., Elser, J. J., Gaedke, U., Ibelings, B., Jeppesen, E.,
Carlos Molinero, J., Mooij, W. M., van Donk, E., Winder, M., Lürling, M., Molinero, J. C., Mooij,
W. M., van Donk, E., Winder, M., Carlos Molinero, J., Mooij, W. M., van Donk, E., & Winder, M.
(2012). Beyond the plankton ecology group (PEG) model: Mechanisms driving plankton
succession. *Annual Review of Ecology, Evolution, and Systematics*, *43*(December), 429–448.
https://doi.org/10.1146/annurev-ecolsys-110411-160251
27. Hatcher, M. J., & Dunn, A. M. (2011). *Parasites in ecological communities: from interactions to
ecosystems*. Cambridge University Press.
28. Marcogliese, D. J. (2004). Parasites: Small Players with Crucial Roles in the Ecological Theater.
EcoHealth, *1*(2), 151–164. https://doi.org/10.1007/s10393-004-0028-3
29. Sime-Ngando, T., Lafferty, K. D., & Biron, D. G. (2007). *Roles and Mechanisms of Parasitism in
Aquatic Microbial Communities*. https://doi.org/10.3389/978-2-88919-588-6
30. Frenken, T., Alacid, E., Berger, S. A., Bourne, E. C., Gerphagnon, M., Grossart, H. P., Gsell, A. S.,
Ibelings, B. W., Kagami, M., Küpper, F. C., Letcher, P. M., Loyau, A., Miki, T., Nejstgaard, J. C.,

Rasconi, S., Reñé, A., Rohrlack, T., Rojas-Jimenez, K., Schmeller, D. S., … Rica, C. (2017). Integrating chytrid fungal parasites into plankton ecology: research gaps and needs. *Environmental Microbiology*, *19*(10), 3802–3822. https://doi.org/10.1111/1462-2920.13827

31. Brussaard, C. P. D., Kuipers, B., & Veldhuis, M. J. W. (2005). A mesocosm study of Phaeocystis globosa population dynamics: I. Regulatory role of viruses in bloom control. *Harmful Algae*, *4*(5), 859–874. https://doi.org/10.1016/j.hal.2004.12.015

32. Gerphagnon, M., Macarthur, D. J., Gachon, C., Ogtrop, F. Van, Latour, D., Lilje, O., Gleason, F. H., & Sime-ngando, T. (2009). *The biological factors affecting the dynamics of cyanobacterial blooms*.

33. Gleason, F. H., Jephcott, T. G., Upper, F. C. K., Karpov, S. A., Guillou, L., Ogtrop, F. F. V. A. N., Pascal, B., Küpper, F. C., Gerphagnon, M., Sime-Ngando, T., Karpov, S. A., Guillou, L., & van Ogtrop, F. F. (2015). Potential roles for recently discovered chytrid parasites in the dynamics of harmful algal blooms. *Fungal Biology Reviews*, *29*(1), 20–33. https://doi.org/10.1016/j.fbr.2015.03.002

34. Ibelings, B. W., Gsell, A. S., Mooij, W. M., Van Donk, E., Van Den Wyngaert, S., & De Senerpont Domis, L. N. (2011). Chytrid infections and diatom spring blooms: Paradoxical effects of climate warming on fungal epidemics in lakes. *Freshwater Biology*, *56*(4), 754–766. https://doi.org/10.1111/j.1365-2427.2010.02565.x

35. Kagami, M., De Bruin, A., Ibelings, B. W., & Van Donk, E. (2007). Parasitic chytrids: Their effects on phytoplankton communities and food-web dynamics. *Hydrobiologia*, *578*(1), 113–129. https://doi.org/10.1007/s10750-006-0438-z

36. Lips, K. R., Brem, F., Brenes, R., Reeve, J. D., Alford, R. A., Voyles, J., Carey, C., Livo, L., Pessier, A. P., & Collins, J. P. (2005). *Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community*.

37. McKenzie, V. J., & Peterson, A. C. (2012). Pathogen pollution and the emergence of a deadly amphibian pathogen. *Molecular Ecology*, *21*(21), 5151–5154. https://doi.org/10.1111/mec.12013

38. Skerratt, L. F., Berger, L., Speare, R., Cashins, S., Mcdonald, K. R., Phillott, A. D., Hines, H. B., & Kenyon, N. (2007). Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth*, *4*(2), 125–134. https://doi.org/10.1007/s10393-007-0093-5

39. Ibelings, B. W., De Bruin, A., Kagami, M., Rijkeboer, M., Brehm, M., & Donk, E. V. (2004). Host parasite interactions between freshwater phytoplankton and chytrid fungi (Chytridiomycota). *Journal of Phycology*, 40(3), 437-453.

40. Bosch, J., Martínez-Solano, I., García-París (2001). Evidence of a chytrid fungus infection involved in the decline of the common midwife toad (Alytes obstetricans) in protected areas of central Spain. *Biological Conservation*, *97*(3), 331–337. https://doi.org/10.1016/S0006-3207(00)00132-4

41. Bruning, K., Lingeman, R., & Ringelberg, J. (1992). Estimating the impact of fungal parasites on phytoplankton populations. *Limnology and Oceanography*, *37*(2), 252–260. https://doi.org/10.4319/lo.1992.37.2.0252

42. Paterson, R. A. (1960). Infestation of Chytridiaceous Fungi on Phytoplankton in Relation to Certain Environmental Factors. *Ecology*, *41*(3), 416–424. https://doi.org/10.2307/1933316

43. Şen, B. (1988). Fungal parasitism of planktonic algae in Shearwater. IV: Parasitic occurrence of a new chytrid species on the blue-green alga *Microcystis aeruginosa* Kuetz. emend. Elenkin.

44. van Donk, E., & Ringelberg, J. (1983). The effect of fungal parasitism on the succession of
diatoms in Lake Maarsseveen I (The Netherlands). *Freshwater Biology*, *13*(3), 241–251.
https://doi.org/10.1111/j.1365-2427.1983.tb00674.x
45. Agha, R., Saebelfeld, M., Manthey, C., Rohrlack, T., & Wolinska, J. (2016). Chytrid parasitism
facilitates trophic transfer between bloom-forming cyanobacteria and zooplankton (Daphnia).
Scientific Reports, *6*. https://doi.org/10.1038/srep35039
46. Frenken, T., Wierenga, J., van Donk, E., Declerck, S. A. J., de Senerpont Domis, L. N., Rohrlack, T.,
& Van de Waal, D. B. (2018). Fungal parasites of a toxic inedible cyanobacterium provide food to
zooplankton. *Limnology and Oceanography*, *63*(6), 2384–2393.
https://doi.org/10.1002/lno.10945
47. Kagami, M., von Elert, E., Ibelings, B. W., de Bruin, A., & van Donk, E. (2007). The parasitic
chytrid, *Zygorhizidium*, facilitates the growth of the cladoceran zooplankter, *Daphnia*, in cultures
of the inedible alga, *Asterionella*. *Proceedings. Biological Sciences / The Royal Society*, *274*(April),
1561–1566. https://doi.org/10.1098/rspb.2007.0425
48. Gsell, A. S., de Senerpont Domis, L. N., van Donk, E., & Ibelings, B. W. (2013). Temperature alters
host genotype-specific susceptibility to chytrid infection. *PLoS One*, 8(8), e71737.
https://doi.org/10.1371/journal.pone.0071737
49. McKindles, K. M., Manes, M. A., McKay, R. M., Davis, T. W., & Bullerjahn, G. S. (2021a).
Environmental factors affecting chytrid (Chytridiomycota) infection rates on *Planktothrix*
agardhii. Journal of Plankton Research, 43(5), 658-672.
50. Fallowfield, H. J., & Daft, M. J. (1988). The extracellular release of dissolved organic carbon by
freshwater cyanobacteria and algae and the interaction with Lysobacter CP-1. *Br Phycol J*,
1617(23), 317–326. https://doi.org/10.1080/00071618800650351
51. Mueller, B., den Haan, J., Visser, P. M., Vermeij, M. J. A., & van Duyl, F. C. (2016). Effect of light
and nutrient availability on the release of dissolved organic carbon (DOC) by Caribbean turf
algae. *Scientific Reports 2016 6:1*, *6*(1), 1–9. https://doi.org/10.1038/srep23248
52. Bruning, K. (1991c). Infection of the diatom Asterionella by a chytrid. II. Effects of light on
survival and epidemic development of the parasite. *Journal of Plankton Research*, *13*(1), 119–
129. https://doi.org/10.1093/plankt/13.1.119
53. Van den Wyngaert, S., Gsell, A. S., Spaak, P., & Ibelings, B. W. (2013). Herbicides in the
environment alter infection dynamics in a microbial host-parasite system. *Environmental*
Microbiology, *15*(3), 837–847. https://doi.org/10.1111/j.1462-2920.2012.02874.x
54. Almocera, A. E. S., Hsu, S. B., & Sy, P. W. (2019). Extinction and uniform persistence in a
microbial food web with mycoloop: Limiting behavior of a population model with parasitic fungi.
Mathematical Biosciences and Engineering, *16*(1), 516–537.
55. Frenken, T., Miki, T., Kagami, M., Van de Waal, D. B., Van Donk, E., Rohrlack, T., & Gsell, A. S.
(2020). The potential of zooplankton in constraining chytrid epidemics in phytoplankton hosts.
Ecology, *101*(1). https://doi.org/10.1002/ecy.2900
56. Gerla, D. J., Gsell, A. S., Kooi, B. W., Ibelings, B. W., Van Donk, E., & Mooij, W. M. (2013).
Alternative states and population crashes in a resource-susceptible-infected model for
planktonic parasites and hosts. *FMeier, M. H. et al. (2015) 'Neuropsychological Decline in*
Schizophrenia from the Premorbid to Post-Onset Period: Evidence from a Population-
Representative Longitudinal Study', American Journal of Psychiatry, 171(1), Pp. 91–101. Doi:
10.1176/Appi.Ajp.2013, *58*(3), 538–551. https://doi.org/10.1111/fwb.12010

57. Miki, T., Takimoto, G., & Kagami, M. (2011). Roles of parasitic fungi in aquatic food webs: A
theoretical approach. *Freshwater Biology*, *56*(6), 1173–1183. https://doi.org/10.1111/j.1365-
2427.2010.02562.x

58. Guillard, R. R. L., & J.Lorenzen, C. (1972). Yellow-green algae with chlorophyllid C. In *Phycology*
(Vol. 8, pp. 10–14).

59. McKindles, K. M., Jorge, A. N., McKay, R. M., Davis, T. W., & Bullerjahn, G. S. (2021b). Isolation
and characterization of Rhizophydiales (Chytridiomycota), obligate parasites of *Planktothrix*
agardhii in a Laurentian Great Lakes embayment. *Applied and Environmental Microbiology*,
87(4), e02308-20.

60. R Core Team. (2021). *R: A Language and Environment for Statistical Computing*.

61. RStudio Team. (2021). *RStudio: Integrated Development Environment for R* (1.4.1106).

62. Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

63. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K.,
Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the {tidyverse}.
Journal of Open Source Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

64. Champely, S. (2018). *PairedData* (1.1.1).

65. Soetaert, K., Petzoldt, T., & Setzer, R. W. (2010). Solving Differential Equations in {R}: Package
deSolve. *Journal of Statistical Software*, *33*(9), 1–25. https://doi.org/10.18637/jss.v033.i09

66. Frenken, T., Velthuis, M., de Senerpont Domis, L. N., Stephan, S., Aben, R., Kosten, S., van Donk,
E., & Van de Waal, D. B. (2016). Warming accelerates termination of a phytoplankton spring
bloom by fungal parasites. *Global Change Biology*, *22*(1), 299–309.
https://doi.org/10.1111/gcb.13095

67. Scholz, B., Vyverman, W., Küpper, F. C., Ólafsson, H. G., & Karsten, U. (2017). Effects of
environmental parameters on chytrid infection prevalence of four marine diatoms: A laboratory
case study. *Botanica Marina*, *60*(4), 419–431. https://doi.org/10.1515/bot-2016-0105

68. Sønstebø, J. H., & Rohrlack, T. (2011). Possible implications of Chytrid parasitism for population
subdivision in freshwater cyanobacteria of the genus Planktothrix. *Applied and Environmental*
Microbiology, *77*(4), 1344–1351. https://doi.org/10.1128/AEM.02153-10

69. Bruning, K. (1991b). Infection of the diatom Asterionella by a chytrid. I. Effects of light on
reproduction and infectivity of the parasite. *Journal of Plankton Research*, *13*(1), 103–117.
https://doi.org/10.1093/plankt/13.1.103

70. Muehlstein, L. K., Amon, J. P., & Leffler, D. L. (1988). Chemotaxis in the Marine Fungus
Rhizophydium littoreum. *Applied and Environmental Microbiology*, *54*(7), 1668–1672.
https://doi.org/10.1128/aem.54.7.1668-1672.1988

71. Esch, G. W., & Fernández, J. C. (1993). Introduction. In *A Functional Biology of Parasitism* (pp. 1–
25). Springer Netherlands. https://doi.org/10.1007/978-94-011-2352-5_1

72. Gerphagnon, M., Colombet, J., Latour, D., & Sime-Ngando, T. (2017). Spatial and temporal
changes of parasitic chytrids of cyanobacteria. *Scientific Reports*, *7*(1), 6056.
https://doi.org/10.1038/s41598-017-06273-1

73. Maier, M. A., & Peterson, T. D. (2017). Prevalence of chytrid parasitism among diatom
populations in the lower Columbia River (2009–2013). *Freshwater Biology*, *62*(2), 414–428.
https://doi.org/10.1111/fwb.12876

74. Sime-Ngando, T. (2012). Phytoplankton chytridiomycosis: Fungal parasites of phytoplankton and
their imprints on the food web dynamics. In *Frontiers in Microbiology* (Vol. 3, Issue OCT, p. 361).
Frontiers Research Foundation. https://doi.org/10.3389/fmicb.2012.00361
75. Kagami, M., & Urabe, J. (2002). Mortality of the planktonic desmid, Staurastrum
dorsidentiferum, due to interplay of fungal parasitism and low light conditions. *SIL Proceedings,*
1922-2010, *28*(2), 1001–1005. https://doi.org/10.1080/03680770.2001.11901868