Tropical hummingbird pollination networks are resistant to short-term experimental

removal of a common flowering plant

Kara G. Leimberger[1, *], Adam S. Hadley[1,2], Sarah J.K. Frey[1,3], and Matthew G. Betts[1]

[1]Forest Biodiversity Research Network, Department of Forest Ecosystems and Society, Oregon

State University, Corvallis, Oregon, USA

[2]Biodiversity Section, Department of Natural Resources and Energy Development, Fredericton,

New Brunswick, Canada

[3]Department of Animal and Rangeland Sciences, Oregon State University, Corvallis, Oregon,

USA

[*]Author for correspondence (E-mail: kleimberger@gmail.com).

ABSTRACT

Theory predicts that the structure of plant-pollination networks should withstand

disturbance, but experiments testing this prediction remain uncommon. In this study, we

simulated the local extinction of a hummingbird-pollinated understory plant, *Heliconia tortuosa,*

from tropical forest fragments using a replicated Before-After-Control-Impact design while

quantifying hummingbird space use (383 hummingbird captures and 72 radio-tagged

individuals), floral visitation rates (>19,000 observation hours), and pollination success (529

flowers). We expected that *H. tortuosa* removal would either result in (i) network collapse, in

which hummingbirds vacate fragments and compromise the reproductive success of other

flowering plants, or (ii) increased hummingbird reliance on alternative resources (rewiring),

leading to sustained fragment use. In our experiment, hummingbird behavior and pollination

were remarkably resistant to loss of *H. tortuosa*, a locally common plant species representing 30-

40% of the available nectar resources on average. The exact mechanisms enabling short-term

hummingbird persistence after resource removal remain unclear, as we did not discover evidence

of rewiring. Physiological adaptations (e.g., torpor and insectivory) may have allowed

hummingbird persistence, perhaps alongside high movement ability. With the important caution

that short-term experiments may not emulate natural extinction processes, our study provides

support for predictions that pollination networks may be robust to plant species loss.

Key words: mutualistic networks, ecosystem function, foraging ecology, functional

connectivity, fragmentation, Trochilidae

INTRODUCTION

Species loss can have cascading consequences for the broader community, including

coextinctions of dependent species and declines in ecosystem function [1,2]. Linked extinctions

may be especially likely for mutualistic interactions, such as those between plants and pollinators

[3,4]. For instance, plants that are not adequately pollinated produce fewer seeds, accruing

extinction debt that becomes realized when the existing generation cannot be replaced by new

recruits [5–7]. This repeated reproductive failure will eventually cause demographic collapse of

plant populations [8,9], which could then hasten the decline of animals that consume pollen or

nectar [10–13]. Indeed, plants and pollinators are experiencing declines worldwide [10,14,15],

which not only imperils the essential ecosystem services on which humans depend but also

threatens the maintenance of global biodiversity [16–18].

However, parallel declines of pollinators and plants are at odds with theoretical

predictions suggesting that species loss should generally not result in coextinction of interaction

partners [19,20]. Tolerance to subsequent extinctions (i.e., robustness) primarily arises from the

expectation that interaction networks exhibit strongly nested topologies, such that specialists

interact with subsets of partners linked to generalists [21]. Under this interaction pattern,

connections between specialists are uncommon, so the functional role of a specialist can be

absorbed by its more abundant generalist partners [21,22]. In addition to the redundancy afforded

by a nested structure, models suggest that flexibility in pollinator foraging behavior – i.e.,

rewiring to use alternative resources – may buffer networks against disturbance [23–26].

Despite these overall predictions and mechanisms of robustness, simulation studies have

revealed that removal of highly connected, generalist species thwarts a network's capacity to

buffer perturbations, leading to rapid partner extinction [19,23,27]. Thus, if external stressors

such as habitat fragmentation threaten highly connected species ['topological keystones' sensu

28], an extinction cascade could occur – particularly without rewiring by pollinators. Uncertainty

about the robustness of real-world plant-pollinator networks still exists, however, because

relatively few studies have quantified pollinator responses to deletions of other species

(Appendix S1: Table S1).

In this study, we tested whether declines of a hummingbird-pollinated plant *(Heliconia*

tortuosa) could have cascading consequences for communities of hummingbirds and flowering

plants within a fragmented tropical forest landscape – or whether rewiring might forestall

network collapse. In Neotropical forests, the genus *Heliconia* (Heliconiaceae) comprises

understory herbs that have been suggested to function as "keystone mutualists" [29]. Large,

brightly colored *Heliconia* inflorescences not only provide nectar to large-bodied hermit

hummingbirds (Trochilidae: Phaethornithinae) throughout the year, but also support a variety of

other hummingbird species during periods of resource scarcity [30,31]. Because most

hummingbird species visit multiple plant species [32], loss of *H. tortuosa* and its associated

hummingbird partners could jeopardize the reproduction of co-occurring plants, thus generating

an extinction cascade [29]. Although *H. tortuosa* is locally common in the mid-elevation,

premontane tropical forests of Costa Rica, Hadley *et al.* [33] found that *H. tortuosa* in smaller

forest fragments produced fewer seeds, possibly due to short-distance pollen transfer associated

with pollinator movement limitation.

To assess the potential consequences of *H. tortuosa* declines, we conducted one of most

highly replicated, large-scale removal experiments conducted within the context of plant-

pollinator interaction networks (Appendix S1: Table S1). We measured hummingbird space use,

floral visitation rates, and plant pollination success before and after temporary experimental

removal of *H. tortuosa* within forest fragments. We developed two alternative hypotheses about

how hummingbirds would respond to declines of *H. tortuosa* (hereafter *Heliconia*) and

potentially affect the pollination of co-occuring plant species (Table 1). Under the *robust*

network hypothesis (**H1**), hummingbirds do not permanently emigrate from forest fragments

following *Heliconia* decline, leading to uninterrupted visitation to remaining plants. We

hypothesized that hummingbirds compensate for *Heliconia* extinction by increasing their use of

non-*Heliconia* floral resources (rewiring). Alternatively, hummingbirds may be highly dependent

on *H. tortuosa* and inflexible in their foraging choices; therefore, local *Heliconia* declines should

reduce nectar availability such that hummingbirds cannot afford the energetic costs to remain in

(or travel to) focal forest fragments, leading to decreases in fragment use and pollination success

(**H2***: parallel extinction hypothesis*). Under both hypotheses, we expected hummingbirds to

change their behavior due to their exceptionally rapid metabolisms, high dependency on nectar,

and limited fat stores [34,35]. Lastly, we expected that changes in *Heliconia* density would affect

green hermits (*Phaethornis guy*) and violet sabrewings (*Campylopterus hemileucurus*) more

strongly than other hummingbird species (**H3**). These two species (hereafter '*Heliconia*

specialists') have bills that are morphologically specialized for long, curved flowers and are

Heliconia's primary pollinators [36–39]. Both species inhabit the forest understory and behave

as high-reward trapliners [sensu 40], visiting spatially dispersed, nectar-rich floral resources –

such as *Heliconia* – in a routine sequence [41].

METHODS

Study system

We conducted this study within a ~122 km^2 area surrounding the Las Cruces Biological

Station in Costa Rica (Coto Brus Canton, Puntarenas Province, 8°47´7´´ N, 82°57´32´´ W, Fig.

1: A-B). This study area comprises cattle pastures, coffee plantations, and fragments of tropical

premontane wet forest created during a period of extensive land clearing concentrated between

1960 and 1980 [42]. Remnant forest fragments (~30% of the study area) span a wide range of

sizes (<1 to >1000 ha), with otherwise isolated fragments often connected by narrow riparian

corridors and living fences [42,43]. Mean annual temperature at the Las Cruces Biological

Station is ~21 ℃, mean annual rainfall is 3500-4000 mm, and a dry season occurs from

December to March [41,42].

Based on visual examination of satellite imagery and previous work in this study area

[33,44,45], we selected 14 focal fragments ('sites', Fig. 1A). Sites were then included based on

presence of flowering *Heliconia***,** landowner permission, vehicle accessibility, and navigable

terrain. To ensure site accessibility, we conducted our study during the dry season and beginning

of the rainy season (February – May), because localized flooding can occur during periods of

heavy rainfall. Elevation of sites ranged from ~900-1500 m (Appendix S2: Table S1).

Experimental design

Site details and experimental timeline

This study used a replicated BACI (Before-After-Control-Impact) design. A BACI design

requires two paired sites: an Impact site (here, 'Treatment') that experiences a disturbance and a

Control site that does not. Additionally, each site is monitored concurrently throughout time

(here, 'Pre' and 'Post' disturbance) (Fig. 2). Therefore, the comparison of interest is the pre-to-

post change in the treatment site relative to the pre-to-post change in the control site. This design

separates treatment effects from natural processes occurring in the study system, either across

space (control *versus* treatment) or time (pre *versus* post).

Here, we paired control and treatment sites based on physical characteristics (elevation

and connectivity to additional forest: Appendix S2: Table S1) and ensured that paired sites were

at least 1000 m apart. This distance threshold corresponds to the estimated daily movement

distance of traplining hummingbirds in this study system [46], making it unlikely that

hummingbirds from the treatment site would relocate to the control site. Actual distances

between sites ranged from 1917-5637 m (mean ± SD: 4256 ± 1528 m, median: 5086 m), and we

never detected hummingbird movement between paired sites — both in terms of hummingbird

captures and radio telemetry (see *'Response variables'*, below). Using the 14 sites, we repeated

the *Heliconia* removal experiment 16 times across three years (2016-2018), yielding a maximum

sample size of 16 control replicates and 16 treatment replicates (Appendix S2: Table S2). Thus,

most sites were used multiple times across the three-year study period, but only once per year

(Appendix S2; Table S3). Sites used multiple times received the opposite treatment of the

previous year, except for 3 of 16 instances (when covering *Heliconia* in that site again would

have posed substantial logistical challenges). Initial treatment assignments were random, which

reduces the likelihood that any treatment differences would be due to factors associated with

individual sites.

Each experimental replicate lasted for 9 days. Starting around noon on the fourth day, we

removed *Heliconia* from the treatment site and designated the following 36 hours as a behavioral

adjustment period during which no data were collected. Although this adjustment period was

necessarily relatively short due to the limited battery lifespan of hummingbird radio-transmitters

(see *'Radio telemetry'*, below), we expected hummingbirds to respond rapidly due to their

exceptionally high metabolisms and limited fat stores [34,35].

Heliconia removal (hummingbird exclusion)

Across all study sites and years, we removed between 8 and 526 *Heliconia* inflorescences

per treatment replicate (mean ± SD: 172 ± 138, median: 138, Appendix S2: Table 4) by covering

Heliconia inflorescences (mature and immature) with dark-colored fabric bags (Fig. 2A). This

approach prevented hummingbird access and obscured flowers from view while avoiding long-

term changes to floral resources associated with destructive sampling methods (e.g., cutting).

The covered inflorescences represented, on average, 97% (± 5% SD) of the *Heliconia*

inflorescences encountered during removal (range: 80-99%); two focal *Heliconia* plants/site

remained uncovered for pre-to-post monitoring, and sometimes plant inaccessibility limited our

covering capacity. While covering *Heliconia*, we also collected data on the abundance and

location of resources available to hummingbirds (*Heliconia* and non-*Heliconia*). Resource data

were used to estimate the percentage of calories removed by our manipulation (see Appendix S3

for methods) and delineate 'focal areas' (see below). In 2017-2018, we also collected resource

data in the control sites, mimicked the trampling effect experienced by the treatment sites.

Focal areas

Within each site, we focused our observations within designated 'focal areas'. Focal areas

encompassed locations where we captured hummingbirds, installed trail cameras, and set up

plant arrays (Fig. 1C, see *'Response variables'*). We delineated focal areas as the 100%

minimum convex polygons of the plant GPS points. Treatment focal areas ranged from 0.27 to

7.49 ha (mean ± SD: 1.7 ± 1.9, median: 1.25, Appendix S2: Table S4), depending on the size of

the forest fragment and the navigability of the terrain. We acknowledge that our focal areas were

smaller, on average, than hummingbird home ranges in this study system [e.g., mean home range

size of 2.2 ha for green hermits: 47]. However, here we were interested in any behavioral

changes occurring *within* the focal area, following similar removal studies (e.g., 20 *x* 20-m plots

within the home range of a bumblebee: Brosi & Briggs 2013).

Response variables

Hummingbird captures

Within each focal area, we captured hummingbirds at the beginning and end of the experimental period (Days 1 and 9). During each capture session, positioned 5-20 mist nets (12-m and 6-m in length) near flowering plants visited by hummingbirds (e.g., the genera *Heliconia, Centropogon, Palicourea, Musa*). Nets were opened 30 minutes after sunrise and remained open for 2-5 hours per session. Capture effort (net-hours) and net locations were kept constant for both capture sessions, except in one instance (Appendix S2: Fig. S1). Following the North American Banding Council's hummingbird protocol [48], we tagged hummingbirds with an aluminum tarsus band, measured body mass to the nearest 0.01 g using an electronic balance, and measured wing length to the nearest 0.5 mm using a wing rule. Tarsus bands allowed detection of recaptured hummingbirds within and between capture sessions; individuals recaptured during the same capture session were only counted once. In 2017-2018, we also marked hummingbirds' heads with unique color combinations of nail polish following Hadley *et al.* [33]; this additional mark allowed us to identify individual hummingbirds seen on camera (see *'Flower visitation'*). All birds were handled in accordance with Oregon State University Animal Care and Use Protocols (ACUP #4655, #5015).

Radio telemetry

We attached miniaturized radio transmitters (0.15-0.25 g, Blackburn Transmitters, Nacogdoches, TX) to 72 hummingbirds according to the methods of Hadley & Betts [43]. We then followed hummingbirds on foot using portable receivers (Wildlife Materials Inc TRX 1000S) and handheld Yagi antennae, recorded locations continuously when the bird was found within receiver range (~250 m in the absence of topographical barriers). In total, we completed

450 hours of telemetry observation across Days 2-4 and Days 6-8 of the experiment, averaging

3.1 ± 0.9 hours per session (typically ~8:00-11:00) and 5.4 ± 0.9 sessions per site (mean ± SD).

From the recorded locations, we estimated the proportion of observation time that radio-tagged

birds spent in the focal area. Thirty-six hummingbirds spent time in the focal area during the

'pre' period and were included in analyses of the *Heliconia* removal experiment; for full methods

and a comprehensive summary of radio-tracking outcomes, see Appendix S2.

Flower visitation

To understand hummingbird resource use within each site and to monitor for the presence

of marked individuals, we positioned trail cameras (PlotWatcher Pro, Day 6 Outdoors) near

inflorescences of plant species known or suspected to be visited by hummingbirds. Effort within

each site was concentrated at two locations ('stations', Fig. 1C) located on roughly opposite ends

of the focal area (mean ± SD distance between stations: 60 ± 44 m). Stations were centered

around a focal *Heliconia* plant expected to flower throughout the experimental period, not in an

extremely steep area, and >10 m from the fragment edge. The two focal *Heliconia* plants

remained uncovered for the entire experimental period, which allowed us to observe whether

visitation rates changed after declines in surrounding *Heliconia* density. To partially standardize

plant species composition between control and treatment replicates, we added a floral array

comprising ~3-5 additional plant species at each station (see Appendix S2 and *'Pollination*

success').

Trail cameras took one photograph per second during daylight hours (5:30-17:30 on Days

2-4 and Days 6-8) and automatically combined these photographs into a time-lapse video. We

reviewed videos for hummingbird sightings using the motion detection program MotionMeerkat

[49] or by manually watching the sped-up video. When hummingbirds were detected on camera,

we identified the species, recorded any color marks, and noted whether they visited any flowers.

From video, we also recorded the number of open flowers available per day. Dates with no

available flowers were excluded from analysis, yielding 20,735 total hours of video. We used

subsets of these data for a variety of purposes, including estimating the total among of floral

resources available for hummingbirds under 'normal' (non-experimental) circumstances. The

analysis for the experiment used 19,870 video hours (Appendix S2: Table S6).

Pollination success

To investigate whether *Heliconia* declines influenced pollination success, we examined

styles for the presence of pollen tubes, which indicate pollen grain germination and subsequent

movement toward the flower ovule(s) [50]. We focused our pollen tube investigation on the two

focal *Heliconia* plants per replicate, as well as several species from the focal arrays (Appendix

S2: Table 7). Throughout the experimental period, we collected flowers from these focal plants

every 1-2 days, which corresponds to the maximum floral longevity for most of the selected

species [*Heliconia tortuosa*, *Hamelia patens*: 51; *Stachytarpheta frantzii*: 52]. Styles were fixed

in formalin-acetic acid-alcohol for at least 24 hours, stained with aniline blue, and examined for

pollen tubes using epifluorescence microscopy [39,50]. Due to low pollen tube presence and/or

challenges with pollen tube visualization, we were only able to analyze two plant species: *H.*

tortuosa ($N = 327$ styles) and *H. patens* ($N = 202$ styles). Additional details about style collection

and lab procedures are available in Appendix S2.

Additional response variables

To supplement the primary response variables described above, we examined several

additional metrics: (1) recapture probability (i.e., proportion of hummingbirds recaptured in the

'post' period), (2) floral visitation rates from individual hummingbirds marked with nail polish,

and (3) body mass of hummingbirds captured during both experimental periods (pre and post).

The first two response variables assessed the possibility that hummingbirds caught during the

initial capture sessions were rapidly replaced by new individuals, potentially obscuring treatment

effects. Body mass measurements allowed us to explore whether birds that remained in (or

returned to) treatment focal areas paid an energetic cost. To compare mass changes across

hummingbirds of different structural sizes, we calculated a relative body mass measure: observed

mass divided by the mass predicted by species-specific allometric equations (i.e., Ln(body mass)

~ Ln(wing length), Appendix S2: Fig. S4).

Statistical analysis

We used linear mixed models (LMMs) and generalized linear mixed models (GLMMs) to

analyze how experimental *Heliconia* extinction influenced (i) hummingbird abundance, (ii)

proportion of time that radio-tagged hummingbirds spent in focal area, (iii) visitation rates to

flowering plants (*Heliconia* and non-*Heliconia*), (iv) plant pollination success, and (v) the

presence and condition of individual hummingbirds (see '*Additional response variables*'). To

determine whether declines in *Heliconia* affected *Heliconia* specialists more strongly than other

hummingbird species (**H3**), we analyzed all hummingbird responses with (i) the full dataset

containing all species, and (ii) a dataset including green hermits and violet sabrewings. We

modeled each response variable using appropriate statistical distributions and controlled for

variable effort when applicable (Appendix S2: Tables S8-S9). All statistical analyses were

conducted in R version 4.2.0 [53] using 'glmmTMB' [54]. We checked statistical assumptions

by visually inspecting plots from the 'performance' package [55] for LMMs and 'DHARMa'

[56] for GLMMs.

In a BACI experimental design, an overall treatment effect is indicated by a statistically

significant interaction between the variables representing treatment and experimental period

[57,58]. Therefore, we tested the *parallel extinction hypothesis* (**H1**) and *robust network*

hypothesis (**H2**) using a two-way interaction between *Heliconia* removal and period (pre *versus*

post) for all response variables measured during both experimental periods (recapture probability

was measured in the post period only). All models included a random intercept for 'Site'. Except

when modeling recapture probability, we also included a nested random effect for 'Replicate'

(Site-Year), which paired pre-to-post observations within each replicate. Additional nested

random effects, for plant species and individual birds, were included where applicable (Appendix

S2: Tables S8-9). Because we were interested in pre-to-post changes in our response variables,

we only included covariates that varied between periods. Specifically, when analyzing flower

visitation, we accounted for the attractiveness of a given focal plant by including a covariate for

the number of open flowers per day, averaged across the experimental period. Since the

magnitude of our experimental *Heliconia* extinction varied widely (Appendix S2: Table S3), we

ran parallel models using a categorical treatment variable (Control/Treatment) and a quantitative

treatment variable, Ln(calories removed/hectare + 1). Because results were qualitatively similar,

only results from the categorical approach are presented in the main text.

RESULTS

Across all sites and years, we detected 19 hummingbird species in the forest fragments

surrounding the Las Cruces Biological Station (Appendix S4: Table S1). We estimate that, on

average, the covered *Heliconia* plants supplied 30% (± 25%, range: 0.4 – 87%) of the calories

available for this entire hummingbird community and 37% (± 28%, range: 0.4 – 94%) of the

calories available for green hermits and violet sabrewings (Appendix S4: Table S2). Although

green hermits and violet sabrewings visited >20 plant species, highest visitation rates were

observed for *Pitcairnia imbricata*, *Erythrina costaricensis*, *Musa x paradisiaca*, *Heliconia*

tortuosa, and *Costus barbatus* (Appendix S4: Fig. S1). *Heliconia* inflorescences were visited by

hummingbirds 6.4 ± 5.7 times per day (range: 0-25, median: 5), mostly by green hermits (75% of

1,055 visits, Appendix S4: Table S3).

Mist net captures & radio telemetry

We found little evidence that *Heliconia* removal reduced captured hummingbird

abundance or decreased the time that radio-tagged hummingbirds spent in the focal area. In

treatment replicates, we observed a pre-to-post decrease in hummingbird captures that was, on

average, 19% greater relative to control replicates but confidence intervals suggest a wide range

of possible effects (95% CI: -47% to +20%, Fig. 3, Appendix S4: Table S4). Radio-tagged

hummingbirds from treatment replicates showed a similar pre-to-post decrease in the percentage

of time spent in the focal area (-18% relative to control replicates, 95% CI: -53% to +40%, Fig.

3, Appendix S4: Table S4). Results were consistent when examining all species together and

Heliconia specialists separately (Fig. 3, Appendix S4: Fig. S2, Tables S5-S6).

Flower visitation & pollination success

We continuously monitored hummingbird visits to the two focal *Heliconia* inflorescences

that remained uncovered throughout the experimental period, as well as numerous other

flowering plant species (mean ± SD: 7.8 ± 2 species/replicate; range: 2-11). Under the *robust*

network hypothesis, we expected hummingbirds to rewire and increasingly use non-*Heliconia*

resources. However, pre-to-post changes in hummingbird visitation rate were non-

distinguishable between control and treatment replicates, both when examining the focal

Heliconia plants (95% CI: -18% to +90%) and non-*Heliconia* plant species (-29% to +10%) (Fig.

3). Responses by *Heliconia* specialists mirrored the responses observed for all species (Fig. 3,

Appendix S4: Tables S7-S8), although these species visited non-*Heliconia* species relatively

infrequently (<1 visit/day, on average: Appendix S4: Fig. S3). We also found no evidence that

experimental *Heliconia* removal influenced pollination success for the focal *Heliconia* plants ($z =$

0.15, $P = 0.88$) or another plant species from the floral arrays (*Hamelia patens: $z = 1.03$, $P =$

0.30) (Fig. 3, Appendix S3: Fig. S4, Table S9).

Additional response variables

When focusing further on the responses of marked individual hummingbirds, we also did

not observe an effect of *Heliconia* removal. On average, the estimated probability of recapturing

a hummingbird from the 'pre' capture session was 18% in control replicates (95% CI: 11% to

29%), compared to 14% in treatment replicates (95% CI: 8% to 23%). This difference was not

statistically significant, and recaptures of *Heliconia* specialists showed qualitatively similar

results (Fig. 4, Appendix S4: Table S10). Results were consistent when examining pre-to-post

changes in flower visitation for individual color-marked hummingbirds; hummingbirds from

treatment replicates only showed a 1% greater pre-to-post decline in visitation rate compared to

those from control replicates (95% CI: -42% to + 70%, Fig. 4, Appendix S4: Fig. S5, Tables

S11-12). Finally, we found no evidence that pre-to-post changes in hummingbird body condition

differed between control and treatment replicates (Fig. 4), either for all hummingbird species ($z =$

-0.49, $P = 0.62$) or *Heliconia* specialists ($z = -0.22$, $P = 0.83$) (Appendix S4: Table S13).

DISCUSSION

In this study, we experimentally induced the local extinction of a locally abundant

tropical plant, *Heliconia tortuosa*, to assess the robustness of plant-hummingbird interaction

networks in a fragmented tropical landscape. Based on suite of complementary response

variables we found that, even when high proportions of total available calories were

experimentally removed (~30-40% on average), hummingbird responses fell within the natural

variability of control replicates. As might be expected with such small changes in hummingbird

behavior, we did not observe any cascading effects of experimental manipulation on pollination

success of flowering plants, which supports the *robust network hypothesis* (**H1**). However, the

mechanism underlying the robust network hypothesis — hummingbirds expanding their diet to

include to alternative floral resources — did not receive strong support, suggesting other

explanations for hummingbird persistence.

Our finding that that experimental removal of *Heliconia* did not elicit parallel extinctions

in pollinators aligns with theoretical predictions [19,23] and several other experiments that

removed plant species from mutualistic networks [59–61]. If hummingbirds remained in forest

fragments despite *Heliconia* declines, we assumed that they would rewire to use alternative floral

resources; hummingbirds as a taxonomic group are highly opportunistic foragers [62,63] and the

most frequent *Heliconia* visitor, the green hermit, has a relatively wide niche breadth [39,41].

Unexpectedly, we found no evidence that hummingbirds increasingly visited non-*Heliconia*

resources when *Heliconia* became less available, suggesting that the mechanisms underlying

network robustness in this study may not necessarily align with the rewiring observed in

previous empirical tests [e.g., 26,61,64]. This apparent lack of foraging flexibility may emerge

from various constraints, such as morphological trait-matching, search costs, and/or competition

from other hummingbirds [see also 65]. For example, high levels of exploitative competition for

non-defendable resources may have discouraged hummingbirds from modifying their trapline

and risking overlap with another bird's foraging route [66–68]. Lastly, is possible that fine-scale

rewiring did occur but was not detected using our analytical approach, which aggregated

visitation rates to the level of hummingbird species; further analysis examining the diet of

individual hummingbirds may shed light on the possibility of rewiring at finer scales.

In the absence of rewiring, the resilience in hummingbird behavior, abundance, and

pollination services following experimental *Heliconia* removal is somewhat surprising. Brief

pauses in nectar intake can disrupt a hummingbird's daily energy balance, given their high

metabolic rate and limited energy reserves [69–71]. Moreover, previous nectar manipulations

with hummingbirds – albeit within defended territories – led to rapid changes in territory size,

foraging patterns, and/or defense behavior, generally within a day [e.g., 72–75]. One explanation

for our result is that hummingbirds permanently emigrated from fragments after *Heliconia*

removal but, because of high intruder pressure, were quickly replaced by new individuals [76].

However, we monitored individual birds using several different approaches (radio telemetry,

recaptures, flower visitation) and found no evidence that *Heliconia* removal caused emigration

and subsequent replacement. An alternative possibility is that hummingbirds relied on

physiological adaptations, such as a torpor and/or insectivory, to maintain body condition despite

large declines in nectar availability – resulting in the stable body mass measurements we

observed in recaptured hummingbirds. Torpor is an energy-saving adaptation characterized by

declines in body temperature and metabolic rate [77,78], and even tropical hummingbirds

inhabiting low-to-mid elevations appear to possess this capability, including the green hermit

[79–82]. Additionally, hummingbirds are highly insectivorous and, although they cannot

completely replace nectar with insects [71,83], they may supplement their diet with arthropods

when nectar becomes scarce [66,84–86]. Finding torpid hummingbirds at night is extremely

challenging [87], but future research could explore changes in arthropod consumption using

techniques such as DNA metabarcoding [88].

Another explanation for hummingbird persistence following experimental *Heliconia*

removal involves their high movement ability. *H. tortuosa* is a patchily distributed species that

necessitates long-distance foraging movements, as a single clump does not make territoriality

energetically profitable [30]. Densities can also vary substantially across short distances; for

instance, we observed a 150-fold difference in *Heliconia* availability across sites and years in the

present study (Appendix S4: Table S14). Similarly, Stiles [36] notes that some Costa Rican

Heliconia species can be common in one watershed, yet nearly absent in an adjacent watershed <

1 km away [see also 89]. In accordance with this natural history, the most frequent *Heliconia*

visitor, the green hermit, can travel long distances (>500 m) while foraging within home ranges

of >150 ha [although home range sizes of ~2 ha are more typical: ,46,47]. Thus, returning to our

focal areas to visit the few remaining *Heliconia* resources could be perceived as a relatively low-

cost option for long-distance trapliners, at least in the short term. Notably, although highly

mobile pollinators might buffer against network collapse [47], they may be vulnerable to habitat

loss and fragmentation once thresholds are crossed, leading to parallel pollinator declines

observed in nature [90,91]. This strategy is also unlikely to be profitable over longer time scales

than measured by our experiment, which only removed *Heliconia* for five days.

Although a short adjustment period (<1 day) is standard in similar species removal

experiments [e.g., 72,86,92], extending the experimental timeline may have more realistically

simulated a future extinction and provided more time for behavioral or demographic adjustment.

If hummingbirds were not permanent residents of our focal areas – as suggested by our relatively

low recapture rates (~15%) and proportion of time spent in the focal area (~20%) – then transient

individuals might have not immediately noticed changes in *Heliconia* density and changed their

foraging patterns accordingly. Likewise, hummingbirds might not have persisted had we

removed *Heliconia* over larger areas, because traveling to isolated plants would become even

more energetically costly [93]. Unfortunately, covering (and then uncovering) *Heliconia* plants

in steep terrain with dense understories limited the area we could manipulate.

We also acknowledge that our ability to detect subtle experimental effects may have been

limited by low statistical power, owing to high individual-level variation and limited sample size.

Nevertheless, we can rule out extreme declines in hummingbird space use, flower visitation, and

pollination success that would be indicative of complete network collapse; treatment replicates

generally showed no more than a 50% pre-to-post decline compared to control replicates (based

on the lower confidence interval of the relevant contrast: Fig. 3, Appendix S4: Table S4). We

also emphasize that the spatial scale of our experiment and amount of replication surpasses all

but two empirical studies simulating species extinctions within plant-pollinator networks

(Appendix S1: Table S1). Still, future removal experiments with highly mobile pollinators

should consider increasing the amount of replication, and, if possible, manipulate larger areas.

CONCLUSIONS

Hummingbird pollinators seemed able to persist during our experiments that removed

floral resources for a short period of time (several days), but we caution that this persistence may

not be sustainable or reflect the outcome of a natural extinction process. Longer-term

experiments will be key for predicting consequences of species loss and understanding

mechanisms of network robustness – at least for pollinator species that forage at large spatial

scales. Our study also suggests that behavioral flexibility to use alternative floral resources might

be limited, even in taxonomic groups thought to be highly opportunistic foragers. Thus, even

rewiring rules with strong biological underpinnings [e.g., 94] may not necessarily predict real-

world responses to species loss, perhaps due to additional constraints such as competition. To

further understand the mechanisms underlying network robustness or collapse, we encourage

future experiments that consider how temporal scale, species traits (e.g., movement ability),

competition, and landscape context interact to determine the outcome of species extinctions.

These experiments could be especially valuable at geographic range limits, where climate change

will likely lead to species replacements [95], and in the tropics where landscape change and

species loss are most rapid [96].

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation (NSF) Graduate

Research Fellowship Program under Grant No. 1840998 to KGL. Funding for the experiment

was provided primarily by NSF-DEB-1457837 to MGB and ASH, with additional grants to KGL

from the American Ornithological Society and Sigma Xi. We also thank the Las Cruces

Biological Station and the private landowners who allowed us to work on their property. Field

assistance was provided by Jorge Araya Paniagua, Michael Atencio Picado, Marion Donald,

Dustin Gannon, Jessica Greer, Matt Hadley, Urs Kormann, Daniel Moreno, Mauricio Paniagua

Castro, Ignacio Ruiz Ilama, Esteban Sandi Paniagua, Andrés Solorzano Vargas, Laura Sutcliffe,

and Felipe Torres Vanegas. Videos from trail cameras were reviewed with the help of Marion

Donald, Colleen Franklin, Jessica Greer, Zoe Griffith, Elena Hart, Billy Hilgert, Zachary

Kendall, Jeremy Lee, Ana Medina Roman, Briley Mullin, Amber Newell, Thanh Nguyen, Zoe

Sallada, Allison Simoni, Andrew Stokes, Claire Woods, and Mel Xiao. Nozomi Birkett and

Marion Donald helped collect nectar measurements, and Chandler Spruill assisted with pollen

tube processing. Mark Novak provided feedback on an earlier version of the manuscript, and

Ariel Muldoon provided valuable statistical advice. We also thank Nickolas Waser, Justin Bain,

and one anonymous reviewer for their suggestions.

DATA AVAILABILITY STATEMENT

Data will be made available through the Dryad Digital Repository. Code for all analyses will be

archived through Zenodo (code is currently available at

github.com/kleimberger/Heliconia_removal_Hummingbird_persistence).

LITERATURE CITED

1.   Colwell RK, Dunn RR, Harris NC. 2012 Coextinction and persistence of dependent species
in a changing world. *Annual Review of Ecology, Evolution, and Systematics* **43**, 183–203.
(doi:10.1146/annurev-ecolsys-110411-160304)

2.   Koh LP, Dunn RR, Sodhi NS, Colwell RK, Proctor HC, Smith VS. 2004 Species
Coextinctions and the Biodiversity Crisis. *Science* **305**, 1632–1634.
(doi:10.1126/science.1101101)

3.   Dunn RR, Harris NC, Colwell RK, Koh LP, Sodhi NS. 2009 The sixth mass coextinction:
are most endangered species parasites and mutualists? *Proceedings of the Royal Society B:*
Biological Sciences **276**, 3037–3045. (doi:10.1098/rspb.2009.0413)

4.   Aslan CE, Zavaleta ES, Tershy B, Croll D. 2013 Mutualism disruption threatens global plant
biodiversity: A systematic review. *PLOS ONE* **8**, e66993.
(doi:10.1371/journal.pone.0066993)

5.   Ashman TL *et al.* 2004 Pollen limitation of plant reproduction: ecological and evolutionary
causes and consequences. *Ecology* **85**, 2408–2421.

6.   Vellend M, Verheyen K, Jacquemyn H, Kolb A, Calster HV, Peterken G, Hermy M. 2006
Extinction Debt of Forest Plants Persists for More Than a Century Following Habitat
Fragmentation. *Ecology* **87**, 542–548. (doi:10.1890/05-1182)

7.   Kuussaari M *et al.* 2009 Extinction debt: a challenge for biodiversity conservation. *Trends in*
ecology & evolution **24**, 564–571.

8.   Anderson SH, Kelly D, Ladley JJ, Molloy S, Terry J. 2011 Cascading effects of bird
functional extinction reduce pollination and plant density. *Science* **331**, 1068–1071.
(doi:10.1126/science.1199092)

9.   Phillips RD *et al.* 2015 Pollinator rarity as a threat to a plant with a specialized pollination
system. *Botanical Journal of the Linnean Society* **179**, 511–525. (doi:10.1111/boj.12336)

10. Biesmeijer JC *et al.* 2006 Parallel declines in pollinators and insect-pollinated plants in
Britain and the Netherlands. *Science* **313**, 351–354. (doi:10.1126/science.1127863)

11. Pauw A, Hawkins JA. 2011 Reconstruction of historical pollination rates reveals linked
declines of pollinators and plants. *Oikos* **120**, 344–349. (doi:https://doi.org/10.1111/j.1600-
0706.2010.19039.x)

12. Weiner CN, Werner M, Linsenmair KE, Blüthgen N. 2014 Land-use impacts on plant–
pollinator networks: interaction strength and specialization predict pollinator declines.
Ecology **95**, 466–474. (doi:https://doi.org/10.1890/13-0436.1)

13. Scheper J, Reemer M, Kats R van, Ozinga WA, Linden GTJ van der, Schaminée JHJ, Siepel
H, Kleijn D. 2014 Museum specimens reveal loss of pollen host plants as key factor driving
wild bee decline in The Netherlands. *PNAS* **111**, 17552–17557.
(doi:10.1073/pnas.1412973111)

14. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global
pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* **25**, 345–353.
(doi:10.1016/j.tree.2010.01.007)

15. Regan EC, Santini L, Ingwall☐King L, Hoffmann M, Rondinini C, Symes A, Taylor J,
Butchart SHM. 2015 Global trends in the status of bird and mammal pollinators.
Conservation Letters **8**, 397–403. (doi:10.1111/conl.12162)

16. Allen-Wardell G *et al.* 1998 The potential consequences of pollinator declines on the
conservation of biodiversity and stability of food crop yields. *Conservation Biology* **12**, 8–
17.

17. Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C,
Tscharntke T. 2007 Importance of pollinators in changing landscapes for world crops.
Proceedings of the Royal Society B: Biological Sciences **274**, 303–313.
(doi:10.1098/rspb.2006.3721)

18. Ollerton J, Winfree R, Tarrant S. 2011 How many flowering plants are pollinated by
animals? *Oikos* **120**, 321–326. (doi:10.1111/j.1600-0706.2010.18644.x)

19. Memmott J, Waser NM, Price MV. 2004 Tolerance of pollination networks to species
extinctions. *Proceedings of the Royal Society B: Biological Sciences* **271**, 2605–2611.

20. Bascompte J, Jordano P, Olesen JM. 2006 Asymmetric coevolutionary networks facilitate
biodiversity maintenance. *Science* **312**, 431–433.

21. Bascompte J, Jordano P, Melián CJ, Olesen JM. 2003 The nested assembly of plant–animal
mutualistic networks. *Proceedings of the National Academy of Sciences* **100**, 9383–9387.

22. Bascompte J, Jordano P. 2007 Plant-animal mutualistic networks: the architecture of
biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**, 567–593.
(doi:10.1146/annurev.ecolsys.38.091206.095818)

23. Kaiser-Bunbury CN, Muff S, Memmott J, Müller CB, Caflisch A. 2010 The robustness of
pollination networks to the loss of species and interactions: a quantitative approach
incorporating pollinator behaviour. *Ecology Letters* **13**, 442–452.

24. Ramos-Jiliberto R, Valdovinos FS, Moisset de Espanés P, Flores JD. 2012 Topological
plasticity increases robustness of mutualistic networks. *Journal of Animal Ecology* **81**, 896–
904. (doi:10.1111/j.1365-2656.2012.01960.x)

25. Valdovinos FS, Moisset de Espanés P, Flores JD, Ramos-Jiliberto R. 2013 Adaptive foraging
allows the maintenance of biodiversity of pollination networks. *Oikos* **122**, 907–917.
(doi:10.1111/j.1600-0706.2012.20830.x)

26. Timóteo S, Ramos JA, Vaughan IP, Memmott J. 2016 High resilience of seed dispersal webs
highlighted by the experimental removal of the dominant disperser. *Current Biology* **26**,
910–915. (doi:10.1016/j.cub.2016.01.046)

27. Traveset A, Tur C, Eguíluz VM. 2017 Plant survival and keystone pollinator species in
stochastic coextinction models: role of intrinsic dependence on animal-pollination. *Sci Rep* **7**,
6915. (doi:10.1038/s41598-017-07037-7)

28. Jordán F. 2009 Keystone species and food webs. *Philosophical Transactions of the Royal
Society B: Biological Sciences* **364**, 1733–1741. (doi:10.1098/rstb.2008.0335)

29. Gilbert LE. 1980 Food web organization and conservation of neotropical diversity. In
Conservation Biology: An Evolutionary-Ecological Perspective (eds ME Soulé, BA Wilcox),
pp. 11–33. Sunderland: Sinauer.

30. Stiles FG. 1975 Ecology, flowering phenology, and hummingbird pollination of some Costa
Rican Heliconia species. *Ecology* **56**, 285–301. (doi:10.2307/1934961)

31. Stiles FG. 1985 Seasonal patterns and coevolution in the hummingbird-flower community of
a Costa Rican subtropical forest. *Ornithological Monographs* **36**, 757–787.
(doi:10.2307/40168315)

32. Rodríguez-Flores CI, Ornelas JF, Wethington S, Arizmendi MC. 2019 Are hummingbirds
generalists or specialists? Using network analysis to explore the mechanisms influencing
their interaction with nectar resources. *PLOS ONE* **14**, e0211855.
(doi:10.1371/journal.pone.0211855)

33. Hadley AS, Frey SJK, Robinson WD, Kress WJ, Betts MG. 2014 Tropical forest
fragmentation limits pollination of a keystone understory herb. *Ecology* **95**, 2202–2212.

34. Suarez RK. 1992 Hummingbird flight: sustaining the highest mass-specific metabolic rates
among vertebrates. *Experientia* **48**, 565–570. (doi:10.1007/BF01920240)

35. Powers DR, Brown AR, Van Hook JA. 2003 Influence of normal daytime fat deposition on
laboratory measurements of torpor use in territorial versus nonterritorial hummingbirds.
Physiological and Biochemical Zoology **76**, 389–397. (doi:10.1086/374286)

36. Stiles FG. 1979 Notes on the natural history of Heliconia (Musaceae) in Costa Rica. Notas
sobre la historia natural de Heliconia (Musaceae) en Costa Rica. *Brenesia* **15**, 151–180.

37. Wolf LL, Stiles FG. 1989 Adaptations for the 'fail-safe' pollination of specialized
ornithophilous flowers. *The American Midland Naturalist* **121**, 1–10. (doi:10.2307/2425651)

38. Taylor J, White SA. 2007 Observations of hummingbird feeding behavior at flowers of
Heliconia beckneri and H. tortuosa in southern Costa Rica. *Ornitologia Neotropical* **18**, 133–
138.

39. Betts MG, Hadley AS, Kress WJ. 2015 Pollinator recognition by a keystone tropical plant.
Proceedings of the National Academy of Sciences **112**, 3433–3438.

40. Feinsinger P, Colwell RK. 1978 Community organization among Neotropical nectar-feeding
birds. *American Zoologist* **18**, 779–795. (doi:10.1093/icb/18.4.779)

41. Borgella R, Snow AA, Gavin TA. 2001 Species richness and pollen loads of hummingbirds
using forest fragments in southern Costa Rica. *Biotropica* **33**, 90–109. (doi:10.1111/j.1744-
7429.2001.tb00160.x)

42. Zahawi RA, Duran G, Kormann U. 2015 Sixty-seven years of land-use change in southern
Costa Rica. *PLOS ONE* **10**, e0143554. (doi:10.1371/journal.pone.0143554)

43. Hadley AS, Betts MG. 2009 Tropical deforestation alters hummingbird movement patterns.
Biol Lett **5**, 207–210. (doi:10.1098/rsbl.2008.0691)

44. Hadley AS, Frey SJK, Robinson WD, Betts MG. 2018 Forest fragmentation and loss reduce
richness, availability, and specialization in tropical hummingbird communities. *Biotropica*
50, 74–83. (doi:10.1111/btp.12487)

45. Kormann U, Scherber C, Tscharntke T, Klein N, Larbig M, Valente JJ, Hadley AS, Betts
MG. 2016 Corridors restore animal-mediated pollination in fragmented tropical forest
landscapes. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152347.
(doi:10.1098/rspb.2015.2347)

46. Volpe NL, Hadley AS, Robinson WD, Betts MG. 2014 Functional connectivity experiments
reflect routine movement behavior of a tropical hummingbird species. *Ecological*
Applications **24**, 2122–2131.

47. Volpe NL, Robinson WD, Frey SJK, Hadley AS, Betts MG. 2016 Tropical forest
fragmentation limits movements, but not occurrence of a generalist pollinator species. *PLOS*
ONE **11**, e0167513.

48. Russell SM, Russell RO. 2001 *The North American Banders' Manual for Banding*
Hummingbirds. Point Reyes Station, CA: North American Banding Council.

49. Weinstein BG. 2015 MotionMeerkat: integrating motion video detection and ecological
monitoring. *Methods in Ecology and Evolution* **6**, 357–362.
(doi:https://doi.org/10.1111/2041-210X.12320)

50. Kearns CA, Inouye DW. 1993 *Techniques for Pollination Biologists*. Niwot: University
Press of Colorado.

51. Stratton DA. 1989 Longevity of individual flowers in a Costa Rican cloud forest: ecological
correlates and phylogenetic constraints. *Biotropica* **21**, 308–318. (doi:10.2307/2388281)

52. Thomas-Granger K. 2003 Light sum, nitrogen level and growth retardant effects on flower
and nectar production in Stachytarpheta frantzii. Master's thesis, University of Guelph.

53. R Core Team. 2022 *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing. See https://www.R-project.org/.

54. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, Skaug HJ,
Maechler M, Bolker BM. 2017 glmmTMB Balances Speed and Flexibility Among Packages
for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* **9**, 378–400.

55. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. 2021 performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *JOSS* **6**, 3139.
(doi:10.21105/joss.03139)

56. Hartig F. 2020 *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*
Regression Models. See https://CRAN.R-project.org/package=DHARMa.

57. Underwood AJ. 1994 On Beyond BACI: sampling designs that might reliably detect
environmental disturbances. *Ecological Applications* **4**, 3–15. (doi:10.2307/1942110)

58. McDonald TL, Erickson WP, McDonald LL. 2000 Analysis of count data from before-after
control-impact studies. *Journal of Agricultural, Biological, and Environmental Statistics* **5**,
262–279. (doi:10.2307/1400453)

59. Ferrero V, Castro S, Costa J, Acuña P, Navarro L, Loureiro J. 2013 Effect of invader
removal: pollinators stay but some native plants miss their new friend. *Biol Invasions* **15**,
2347–2358. (doi:10.1007/s10530-013-0457-4)

60. Goldstein J, Zych M. 2016 What if we lose a hub? Experimental testing of pollination
network resilience to removal of keystone floral resources. *Arthropod-Plant Interactions* **10**,
263–271. (doi:10.1007/s11829-016-9431-2)

61. Costa JM *et al.* 2018 Rewiring of experimentally disturbed seed dispersal networks might
lead to unexpected network configurations. *Basic and Applied Ecology* **30**, 11–22.
(doi:10.1016/j.baae.2018.05.011)

62. Arizmendi MC, Ornelas JF. 1990 Hummingbirds and their floral resources in a tropical dry
forest in Mexico. *Biotropica* **22**, 172–180. (doi:10.2307/2388410)

63. Maruyama PK, Oliveira GM, Ferreira C, Dalsgaard B, Oliveira PE. 2013 Pollination
syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna
hummingbirds. *Naturwissenschaften* **100**, 1061–1068. (doi:10.1007/s00114-013-1111-9)

64. Kaiser-Bunbury CN, Mougal J, Whittington AE, Valentin T, Gabriel R, Olesen JM,
Blüthgen N. 2017 Ecosystem restoration strengthens pollination network resilience and
function. *Nature* **542**, 223–227. (doi:10.1038/nature21071)

65. Weinstein BG, Graham CH. 2017 Persistent bill and corolla matching despite shifting
temporal resources in tropical hummingbird-plant interactions. *Ecol. Lett.* **20**, 326–335.
(doi:10.1111/ele.12730)

66. Young AM. 1971 Foraging for insects by a tropical hummingbird. *The Condor* **73**, 36–45.
(doi:10.2307/1366122)

67. Gill FB. 1988 Trapline foraging by hermit hummingbirds: competition for an undefended,
renewable resource. *Ecology* **69**, 1933–1942. (doi:10.2307/1941170)

68. Ohashi K, Thomson JD. 2009 Trapline foraging by pollinators: its ontogeny, economics and
possible consequences for plants. *Ann Bot* **103**, 1365–1378. (doi:10.1093/aob/mcp088)

69. Calder WA, Booser J. 1973 Hypothermia of broad-tailed hummingbirds during incubation in
nature with ecological correlations. *Science* **180**, 751–753.
(doi:10.1126/science.180.4087.751)

70. Tooze ZJ, Gass CL. 1985 Responses of rufous hummingbirds to midday fasts. *Can. J. Zool.*
63, 2249–2253. (doi:10.1139/z85-333)

71. Brice AT. 1992 The essentiality of nectar and arthropods in the diet of the Anna's
hummingbird (Calypte anna). *Comparative Biochemistry and Physiology Part A: Physiology*
101, 151–155.

72. Hixon MA, Carpenter FL, Paton DC. 1983 Territory area, flower density, and time budgeting
in hummingbirds: an experimental and theoretical analysis. *The American Naturalist* **122**,
366–391. (doi:10.1086/284141)

73. Gass CL, Sutherland GD. 1985 Specialization by territorial hummingbirds on experimentally
enriched patches of flowers: energetic profitability and learning. *Can. J. Zool.* **63**, 2125–
2133. (doi:10.1139/z85-313)

74. Eberhard JR, Ewald PW. 1994 Food availability, intrusion pressure and territory size: an
experimental study of Anna's hummingbirds (Calypte anna). *Behav Ecol Sociobiol* **34**, 11–
18. (doi:10.1007/BF00175453)

75. Temeles EJ, Muir AB, Slutsky EB, Vitousek MN. 2004 Effect of food reductions on
territorial behavior of purple-throated caribs. *Condor* **106**, 691–695.
(doi:10.1093/condor/106.3.691)

76. Stiles FG, Wolf LL. 1970 Hummingbird territoriality at a tropical flowering tree. *The Auk*
87, 467–491. (doi:10.2307/4083791)

77. Hainsworth FR, Collins BG, Wolf LL. 1977 The function of torpor in hummingbirds.
Physiological Zoology **50**, 215–222. (doi:10.1086/physzool.50.3.30155724)

78. Spence AR, Tingley MW. 2021 Body size and environment influence both intraspecific and
interspecific variation in daily torpor use across hummingbirds. *Functional Ecology* **35**, 870–
883. (doi:10.1111/1365-2435.13782)

79. Krüger K, Prinzinger R, Schuchmann K-L. 1982 Torpor and metabolism in hummingbirds.
Comparative Biochemistry and Physiology Part A: Physiology **73**, 679–689.
(doi:10.1016/0300-9629(82)90275-4)

80. Bech C, Abe AS, Steffensen JF, Berger M, Bicudo JEPW. 1997 Torpor in three species of
Brazilian hummingbirds under semi-natural conditions. *The Condor* **99**, 780–788.
(doi:10.2307/1370489)

81. Shankar A, Schroeder RJ, Wethington SM, Graham CH, Powers DR. 2020 Hummingbird
torpor in context: duration, more than temperature, is the key to nighttime energy savings.
Journal of Avian Biology **51**, e02305. (doi:10.1111/jav.02305)

82. Schuchmann K-L, Prinzinger R. 1988 Energy metabolism, nocturnal torpor, and respiration
frequency in a Green Hermit (Phaethornis guy). *J Ornithol* **129**, 469–472.
(doi:10.1007/BF01644492)

83. Stiles FG. 1995 Behavioral, ecological and morphological correlates of foraging for
arthropods by the hummingbirds of a tropical wet forest. *The Condor* **97**, 853–878.
(doi:10.2307/1369527)

84. Kuban JF, Neill RL. 1980 Feeding ecology of hummingbirds in the highlands of the Chisos
Mountains, Texas. *The Condor* **82**, 180–185. (doi:10.2307/1367475)

85. Montgomerie RD, Redsell CA. 1980 A nesting hummingbird feeding solely on arthropods.
The Condor **82**, 463. (doi:10.2307/1367577)

86. Hazlehurst JA, Karubian JO. 2018 Impacts of nectar robbing on the foraging ecology of a
territorial hummingbird. *Behavioural Processes* **149**, 27–34.
(doi:10.1016/j.beproc.2018.01.001)

87. Carpenter FL, Hixon MA. 1988 A new function for torpor: fat conservation in a wild migrant
hummingbird. *The Condor* **90**, 373–378.

88. Moran AJ, Prosser SWJ, Moran JA. 2019 DNA metabarcoding allows non-invasive
identification of arthropod prey provisioned to nestling Rufous hummingbirds (Selasphorus
rufus). *PeerJ* **7**, e6596. (doi:10.7717/peerj.6596)

89. Bruna EM, Kress WJ. 2002 Habitat fragmentation and the demographic structure of an
Amazonian understory herb (Heliconia acuminata). *Conservation Biology* **16**, 1256–1266.
(doi:https://doi.org/10.1046/j.1523-1739.2002.99494.x)

90. Andrén H. 1994 Effects of habitat fragmentation on birds and mammals in landscapes with
different proportions of suitable habitat: A review. *Oikos* **71**, 355–366.
(doi:10.2307/3545823)

91. Hadley AS, Betts MG. 2012 The effects of landscape fragmentation on pollination dynamics:
absence of evidence not evidence of absence. *Biological Reviews* **87**, 526–544.
(doi:10.1111/j.1469-185X.2011.00205.x)

92. Brosi BJ, Briggs HM. 2013 Single pollinator species losses reduce floral fidelity and plant
reproductive function. *Proceedings of the National Academy of Sciences* **110**, 13044–13048.

93. Heinrich B, Raven PH. 1972 Energetics and pollination ecology. *Science* **176**, 597–602.

94. Vizentin-Bugoni J, Debastiani VJ, Bastazini VAG, Maruyama PK, Sperry JH. 2020
Including rewiring in the estimation of the robustness of mutualistic networks. *Methods in*
Ecology and Evolution **11**, 106–116. (doi:10.1111/2041-210X.13306)

95. HilleRisLambers J, Harsch MA, Ettinger AK, Ford KR, Theobald EJ. 2013 How will biotic
interactions influence climate change–induced range shifts? *Annals of the New York*
Academy of Sciences **1297**, 112–125. (doi:10.1111/nyas.12182)

96. Hansen MC *et al.* 2013 High-resolution global maps of 21st-century forest cover change.
Science **342**, 850–853. (doi:10.1126/science.1244693)

97. Mendenhall CD, Wrona AM. 2018 Improving tree cover estimates for fine-scale landscape
ecology. *Landscape Ecol* **33**, 1691–1696. (doi:10.1007/s10980-018-0704-2)

98. Lenth R. 2020 *emmeans: Estimated Marginal Means, aka Least-Squares Means*. See
https://CRAN.R-project.org/package=emmeans.

TABLES

Table 1. Hypotheses testing whether declines of a locally common, tropical flowering plant (*Heliconia tortuosa*) affect hummingbird

space use, hummingbird floral visitation rates, and plant pollination success. Predicted effects are indicated in the column below each

hypothesis. We experimentally removed *H. tortuosa* using a Before-After-Control-Impact (BACI) experimental design (Fig. 2), so

predicted effects can be conceptualized as pre-to-post changes in treatment replicates compared to pre-to-post changes in control

replicates.

	Response variable	Methodological approach	How do hummingbirds respond to declines in *Heliconia,* and are there cascading effects on the remaining plants?		
			Robust network hypothesis (H1)	Parallel extinction hypothesis (H2)	Results from this study
Space use	Abundance	Mist net captures	No change	Decrease	**No change**
	Time spent in focal area	Radio telemetry			
Flower visitation rates	Visitation to focal *Heliconia* plants	Remote camera observations	No change or increase	Decrease	**No change**
	Visitation to non-*Heliconia* plants	Remote camera observations	Increase (rewiring)	Decrease	**No change**
Pollination services	Pollination success	Pollen tubes	No change or increase	Decrease	**No change**

  FIGURES

Figure 1. This study was conducted in the fragmented landscape surrounding the Las Cruces

Biological Station (A) in southern Costa Rica (B). Within the study system (rectangular extent),

we utilized fourteen sites. Forested areas are shown in grey, and non-forested areas (white)

primarily comprise pasture. The forest layer is a combination of areas digitized for this study

(generally within 1000 m of each site) and those digitized by Mendenhall & Wrona [97]. (C) An

example site, with red line representing a focal area [i.e., where we removed *Heliconia* (for

treatment replicates) and/or surveyed for floral resources (for treatment and control replicates)].

Within focal areas, we captured hummingbirds using mist nets (triangles) and monitored plant-

hummingbird interactions using trail cameras (camera icons).

Figure 2. We temporarily removed *Heliconia* by covering inflorescences with fabric bags (A) as part of our experiment Before-After-Control-Impact experimental design (B). Within a year (2016–2018), each site received either *Heliconia* removal (Treatment) or was assigned to be a Control. Treatments were reversed in alternate years for a total maximum sample size of 16 control replicates and 16 treatment replicates. Additionally, sites were sampled during two experimental periods: Before ('Pre') and After ('Post'). With this experimental design, the comparison of interest is the pre-to-post change in treatment replicates compared to the pre-to-post change in control replicates. *Photos provided by M. Donald.*

Figure 3. Effects of experimental *Heliconia* removal on hummingbird space use, flower visitation, and plant pollination success.

Following the Before-After-Control-Impact experimental design, the *y*-axis reflects the pre-to-post change in treatment replicates

relative to pre-to-post change in control replicates. Estimates and 95% confidence intervals were calculated using the 'contrasts'

function of 'emmeans' [98]. The dashed line at 1 indicates no experimental effect; the grey area indicates support for the *network*

collapse hypothesis. *Heliconia* specialists refers to the two hummingbird species expected to rely most strongly on *H. tortuosa*: green

hermits (*P. guy*) and violet sabrewings (*C. hemileucurus*).

Figure 4. Effects of experimental *Heliconia* removal on additional response variables, which investigate whether hummingbirds

present at the start of the experiment were replaced by new individuals (mist net recaptures and flower visitation of marked birds) or

suffered an energetic cost by remaining in the focal area (bird condition). Following the Before-After-Control-Impact experimental

design, the *y*-axis reflects treatment replicates relative to control replicates (mist net recaptures) or the pre-to-post change in treatment

replicates relative to pre-to-post change in control replicates (flower visitation of marked birds and bird condition). Estimates and 95%

confidence intervals were calculated using the 'contrasts' function of 'emmeans' [98]. The dashed lines at 1 and 0 indicate no

experimental effect; the grey area indicates support for the *network collapse hypothesis*. *Heliconia* specialists refers to the two

hummingbird species expected to rely most strongly on *H. tortuosa*: green hermits (*P. guy*) and violet sabrewings (*C. hemileucurus*).