Are tropical hummingbird pollination networks resistant to experimental extirpation of a

common flowering plant?

Kara G. Leimberger[1, *], Adam S. Hadley[1,2], Sarah J.K. Frey[1,3], and Matthew G. Betts[1]

[1]Forest Biodiversity Research Network, Department of Forest Ecosystems and Society, Oregon

State University, Corvallis, Oregon, USA

[2]Biodiversity Section, Department of Natural Resources and Energy Development, Fredericton,

New Brunswick, Canada

[3]Department of Animal and Rangeland Sciences, Oregon State University, Corvallis, Oregon,

USA

[*]Author for correspondence (E-mail: kara.leimberger@oregonstate.edu).

ABSTRACT

Plants and pollinators are experiencing parallel declines worldwide, despite theory predicting that pollination networks should withstand disturbance due to redundancy of pollinators, rarity of interactions between specialists, and flexible pollinator foraging behavior (rewiring). Experiments evaluating the stability of plant-pollinator networks may help resolve this incongruity but remain uncommon. In this study, we simulated the extirpation of a hummingbird-pollinated understory plant, *Heliconia tortuosa,* from tropical forest fragments using a replicated Before-After-Control-Impact experimental design while quantifying hummingbird space use (383 hummingbird captures and 72 radio-tagged individuals), floral visitation rates (6,759 visitations from 20,725 observation hours), and plant pollination success (529 flowers). To complement this experimental approach, we also examined these responses across a natural gradient in *H. tortuosa* density. We expected that declines of *H. tortuosa* would either result in (a) network collapse, in which hummingbirds vacate fragments and compromise the reproductive success of other flowering plants, or (b) increased hummingbird reliance on alternative resources, leading to sustained fragment use. We also hypothesized that landscape and local context (i.e., connectivity to additional forest area and alternative resource abundance) could mediate hummingbird responses to *H. tortuosa* declines; for example, connectivity could facilitate hummingbird visitation to areas of reduced food availability. In our removal experiment, hummingbird persistence and plant pollination success were remarkably resistant to loss of *H. tortuosa*, a locally common plant species representing >40% of the available nectar resources on average. However, naturally low *H. tortuosa* densities were associated with reduced floral visitation rates and decreased pollination. Although landscape context (connectivity) led to higher hummingbird abundance, we found little evidence that connectivity or resource availability mediated hummingbird responses to declines in *H. tortuosa* availability. The exact

mechanisms enabling short-term hummingbird persistence after resource removal remain

unclear, as we did not discover evidence of rewiring. Physiological adaptations (e.g., torpor and

insectivory), may have contributed to hummingbird resilience, perhaps alongside high movement

abilities that facilitated visitation to spatially dispersed floral resources. With the important

caution that short-term experiments may not realistically emulate natural extinction processes,

our study provides partial support for theoretical predictions that pollination networks may be

relatively robust to plant species loss.

Key words: mutualistic networks, ecosystem function, foraging ecology, functional

connectivity, fragmentation, Trochilidae

INTRODUCTION

Species loss can have cascading consequences for the broader community, including

coextinctions of dependent species and declines in ecosystem function (Colwell, Dunn, and

Harris 2012; Koh et al. 2004). Linked extinctions may be especially likely for mutualistic

interactions, such as those between plants and pollinators (Dunn et al. 2009; Aslan et al. 2013).

For instance, plants that are not adequately pollinated produce fewer seeds, accruing extinction

debt that becomes realized when the existing generation cannot be replaced by new recruits

(Ashman et al. 2004; Vellend et al. 2006; Kuussaari et al. 2009). This repeated reproductive

failure will eventually cause demographic collapse of plant populations (Anderson et al. 2011;

Phillips et al. 2015), which could then hasten the decline of animals that consume pollen or

nectar (Biesmeijer et al. 2006; Pauw and Hawkins 2011; Weiner et al. 2014; Scheper et al. 2014).

Indeed, plants and pollinators are experiencing declines worldwide (Biesmeijer et al. 2006; Potts

et al. 2010; Regan et al. 2015), which not only imperils the essential ecosystem services on

which humans depend but also threatens the maintenance of global biodiversity (Allen-Wardell

et al. 1998; Klein et al. 2007; Ollerton, Winfree, and Tarrant 2011).

Although the underlying reasons for plant and pollinator declines are not fully

understood, one prominent hypothesis invokes the landscape-level processes of habitat loss and

fragmentation (Kearns, Inouye, and Waser 1998; Aguilar et al. 2006; Potts et al. 2010; Hadley

and Betts 2012). Habitat destruction often creates small habitat patches unable to sustain large

populations, which can increase rates of local extinction (Hanski 1998), reduce functional

connectivity (sensu P. D. Taylor et al. 1993), and impede pollinator movement (Hadley and

Betts 2012). Likely due to these combined mechanisms, land-use change and habitat isolation

have been repeatedly linked to declines in pollinator diversity (Winfree, Bartomeus, and

Cariveau 2011; Montero-Castaño and Vilà 2012; Senapathi et al. 2015) and plant reproduction

(Steffan-Dewenter and Tscharntke 1999; Taki, Kevan, and Ascher 2007; Farwig et al. 2009;

Garibaldi et al. 2011).

However, parallel declines of pollinators and plants are at odds with theoretical

predictions suggesting that species loss should generally not result in coextinction of interaction

partners (Memmott, Waser, and Price 2004; Bascompte, Jordano, and Olesen 2006). Tolerance

to subsequent extinctions (i.e., robustness) primarily arises from the expectation that interaction

networks exhibit strongly nested topologies, such that specialists interact with subsets of partners

linked to generalists (Bascompte et al. 2003). Under this interaction pattern, connections between

specialists are uncommon, so the functional role of a specialist can be absorbed by its more

abundant generalist partners (Bascompte et al. 2003; Bascompte and Jordano 2007). In addition

to the redundancy afforded by a nested structure, models suggest that flexibility in pollinator

foraging behavior – i.e., rewiring to use alternative resources – may buffer networks against

disturbance (Kaiser-Bunbury et al. 2010; Ramos-Jiliberto et al. 2012; Valdovinos et al. 2013;

Timóteo et al. 2016).

Despite these overall predictions and mechanisms of robustness, pollination networks

may be vulnerable to habitat loss and fragmentation under certain conditions. Although

systematic loss of generalist pollinators may be less realistic than extinctions of rare, specialist

pollinators (Aizen, Sabatino, and Tylianakis 2012), simulation studies have revealed that

removal of highly connected, generalist species thwarts a network's capacity to buffer

perturbations, leading to rapid partner extinction (Memmott, Waser, and Price 2004; Kaiser-

Bunbury et al. 2010; Traveset, Tur, and Eguíluz 2017). Thus, if external stressors such as habitat

fragmentation threaten highly connected species ('topological keystones' sensu Jordán 2009), an

extinction cascade could occur – particularly without rewiring by pollinators. However,

uncertainty about the robustness of real-world plant-pollinator networks still exists because

relatively few studies have quantified pollinator responses to deletions of other species

(Appendix S1: Table S1).

In this study, we tested whether declines of a hummingbird-pollinated plant *(Heliconia*

tortuosa) could have cascading consequences for communities of hummingbirds and flowering

plants within a fragmented tropical forest landscape – or whether rewiring might forestall

network collapse. In Neotropical forests, the genus *Heliconia* (Heliconiaceae) comprises

understory herbs that have been suggested to function as "keystone mutualists" (Gilbert 1980).

Large, brightly colored *Heliconia* inflorescences not only provide nectar to large-bodied hermit

hummingbirds (Trochilidae: Phaethornithinae) throughout the year, but also support a variety of

other hummingbird species during periods of resource scarcity (Stiles 1975; 1985). Because most

hummingbird species visit multiple plant species (Rodríguez-Flores et al. 2019), loss of *H.*

tortuosa and its associated hummingbird partners could jeopardize the reproduction of co-

occurring plants, thus generating an extinction cascade (Gilbert 1980). Although *H. tortuosa* is

locally common in the mid-elevation, premontane tropical forests of Costa Rica, Hadley *et al.*

(2014) found that *H. tortuosa* in smaller forest fragments produced fewer seeds, possibly due to

short-distance pollen transfer associated with pollinator movement limitation. Although

numerous hummingbird species visit *H. tortuosa* opportunistically, its primary pollinators are the

green hermit (*Phaethornis guy*) and violet sabrewing (*Campylopterus hemileucurus*) (Stiles

1979; Wolf and Stiles 1989; J. Taylor and White 2007; Betts, Hadley, and Kress 2015). These

relatively large, forest-associated hummingbirds are high-reward trapliners (sensu Feinsinger and

Colwell 1978) that visit spatially dispersed, nectar-rich floral resources in a routine sequence

(Borgella, Snow, and Gavin 2001). However, their movements become restricted in fragmented

habitats (Hadley and Betts 2009; Volpe et al. 2016; Kormann et al. 2016).

To assess the potential consequences of *H. tortuosa* declines, we measured hummingbird

space use, floral visitation rates, and plant pollination success before and after temporary

experimental removal of *H. tortuosa* within forest fragments. Additionally, we assessed how

these responses varied across a natural gradient in *Heliconia* density. We developed two

alternative hypotheses about how hummingbirds would respond to declines of *H. tortuosa*

(hereafter *Heliconia*) and potentially affect the pollination of co-occuring plant species (Table 1).

Under the *robust network hypothesis* (**H1a**), hummingbirds do not emigrate from forest

fragments following *Heliconia* decline, leading to uninterrupted visitation to remaining plants.

Under this scenario, we hypothesized that hummingbirds compensate for *Heliconia* extinction by

increasing their use of non-*Heliconia* floral resources (rewiring). Alternatively, hummingbirds

may be highly dependent on *H. tortuosa* and inflexible in their foraging choices; therefore, local

Heliconia declines should reduce nectar availability such that hummingbirds cannot afford the

energetic costs to remain in (or travel to) focal forest fragments, leading to decreases in fragment

use and pollination success (**H1b***: parallel extinction hypothesis*). Under both hypotheses, we

expected hummingbirds to change their behavior due to their exceptionally rapid metabolisms,

high dependency on nectar, and limited fat stores (Suarez 1992; Powers, Brown, and Van Hook

2003). Additionally, we hypothesized that landscape and local context could mediate the

influence of *Heliconia* declines on hummingbird behavior (Table 1). Connectivity to additional

forest area (*connectivity hypothesis*, **H2**) may sustain hummingbird fragment use by facilitating

travel to fragments (*rescue hypothesis*, **H2a**) or promote steeper declines in fragment use by

facilitating hummingbird departure (*abandonment hypothesis*, **H2b**). Further, greater local

availability of non-*Heliconia* floral resources may promote hummingbird persistence through

increased opportunities for rewiring (*resource hypothesis,* **H3**)*.* Lastly, we expected that changes

in *Heliconia* density would affect green hermits and violet sabrewings more strongly than other

hummingbird species (**H4**). Green hermits and violet sabrewings (hereafter '*Heliconia*

specialists') inhabit the forest understory, have bills that are morphologically specialized to long,

curved *H. tortuosa* flowers, and are thought to visit *H. tortuosa* more frequently than other

hummingbirds (Borgella, Snow, and Gavin 2001; Betts, Hadley, and Kress 2015).

To our knowledge, this study not only constitutes one of most highly replicated, large-

scale removal experiments conducted within the context of plant-pollinator interaction networks

(Appendix S1: Table S1), but also represents the first test of how landscape and local context

influence hummingbird responses to declines in a common floral resource. Importantly,

understanding mechanisms of pollinator persistence in fragmented landscapes has implications

for conservation planning, since fragmentation effects may be ameliorated by habitat restoration,

preservation of riparian corridors, and/or implementation of living fences (Renjifo 2001;

Kormann et al. 2016).

METHODS

Study area

This study was conducted in southern Costa Rica within a ~122 km^2 area surrounding the Las

Cruces Biological Station (Coto Brus Canton, Puntarenas Province, 8°47´7´´ N, 82°57´32´´ W,

Fig. 1: A-B). This study area comprises cattle pastures, coffee plantations, and fragments of

tropical premontane wet forest created during a period of extensive land clearing concentrated

between 1960 and 1980 (Zahawi, Duran, and Kormann 2015). Remnant forest fragments (~30%

of the study area) span a wide range of sizes (<1 to >1000 ha), with otherwise isolated fragments

often connected by narrow riparian corridors and living fences (Hadley and Betts 2009; Zahawi,

Duran, and Kormann 2015). Mean annual temperature at the Las Cruces Biological Station is

~21 ℃, mean annual rainfall is 3500-4000 mm, and a dry season occurs from December to

March (Borgella, Snow, and Gavin 2001; Zahawi, Duran, and Kormann 2015).

Study sites

We selected 14 focal fragments embedded within focal landscapes of 1000 m radius

('sites', Fig. 1: A, C). We chose the spatial extent of focal landscapes *a priori* based on the

maximum estimated daily movement distance for large, forest-associated hummingbird species

in this study system (Volpe et al. 2014).We identified potential sites using a stratified design

with two categories: 'connected to forest' or 'unconnected to forest', based on visual

examination of satellite imagery and previous work in this study area (Hadley et al. 2014; 2018;

Kormann et al. 2016). This design was only for site establishment, as we later quantified

connectivity as a continuous variable (see *'Quantifying connectivity'* below). Sites were then

included based on presence of flowering *Heliconia*, landowner permission, vehicle accessibility,

and navigable terrain. To ensure site accessibility, we conducted our study during the dry season

and beginning of the rainy season (February – May), because localized flooding can occur during

periods of heavy rainfall. Elevation of sites ranged from ~900-1500 m (Appendix S2: Table S1).

Study design

Using the 14 focal-patch landscapes, we repeated the *Heliconia* removal experiment 16

times across three years (2016-2018). Each iteration of the experiment followed a BACI (Before-

After-Control-Impact) design and used two sites paired based on elevation and connectivity

(Appendix S2: Table S2). One of the paired sites ('treatment site') received the treatment of

Heliconia removal, and we designated the other site as a control ($N = 16$ control replicates, $N =$

16 treatment replicates). Most sites were used multiple times across the three-year study period,

but only once per year (Appendix S2: Tables S2-S3). Initial treatment assignments were random,

and sites used multiple times received the opposite treatment of the previous year (except in 3 of

16 instances); this design reduces the likelihood that treatment-control differences were due to

factors associated with individual sites. To prevent hummingbirds from the treatment site

relocating to the control site, paired sites were located at least 1000 m apart (mean $\pm$ SD: 2266 $\pm$

1511 m, median: 3086 m).

Each experimental iteration lasted nine days and began with a period of pre-treatment

data collection in each site (Fig. 2A). Starting around noon on the fourth day, we removed

Heliconia from the treatment site and designated the following 36 hours as a behavioral

adjustment period during which no data were collected. Although this adjustment period was

necessarily relatively short due to the limited battery lifespan of hummingbird radio-transmitters

(see *'Radio telemetry'*), we expected hummingbirds to respond rapidly due to their exceptionally

high metabolisms and limited fat stores (Suarez 1992; Powers, Brown, and Van Hook 2003). To

ensure that *Heliconia* inflorescences were not overlooked during removal, field technicians

positioned themselves ~2-3 meters apart and slowly walked through the forest while covering all

Heliconia inflorescences (mature and immature) with dark-colored fabric bags (Fig. 2B). These

bags prevented hummingbird access, obscured flowers from view, and allowed us to avoid long-

term changes to floral resources associated with destructive sampling methods (e.g., cutting).

The location of each *Heliconia* plant was recorded using a handheld GPS unit, and we retrieved

the covering bags at the end of the experimental period. In each site, we also recorded the

location and identity of all non-*Heliconia* floral resources available to hummingbirds (see

'Quantifying resource availability'). The areas where we removed *Heliconia* and/or surveyed for

floral resources are referred to as 'focal areas' (Fig. 1D), which we delineated using 100%

minimum convex polygons. Focal areas also encompass locations where we captured

hummingbirds, installed trail cameras, and set up plant arrays (Fig. 1D, see *'Response*

variables').

Response variables

Hummingbird captures

Within each focal area, we captured hummingbirds at the beginning and end of the

experimental period (Fig. 2A) by deploying 5 - 20 mist nets (12-m and 6-m in length) near

flowering plants visited by hummingbirds (e.g., the genera *Heliconia*, *Centropogon, Palicourea,*

Musa). Nets were opened 30 minutes after sunrise and remained open for 2 - 5 hours per session.

Capture effort (net-hours) and net locations were kept constant for both capture sessions, except

in one instance (Appendix S2: Figure S1). Following the North American Banding Council's

hummingbird protocol (Russell and Russell 2001), we tagged hummingbirds with an aluminum

tarsus band, measured body mass to the nearest 0.01 g using an electronic balance, and measured

wing length to the nearest 0.5 mm using a wing rule. Tarsus bands allowed detection of

recaptured hummingbirds within and between capture sessions; individuals recaptured during the

same capture session were only counted once. In 2017-2018, we also marked hummingbirds'

heads with unique color combinations of nail polish following Hadley *et al.* (2014); this

additional mark allowed us to identify individual hummingbirds seen on trail cameras (see

'Camera observations'). All birds were handled in accordance with Oregon State University

Animal Care and Use Protocols (ACUP #4655, #5015).

Radio telemetry

We attached miniaturized radio transmitters (0.15-0.25g, Blackburn Transmitters,

Nacogdoches, TX) to 72 hummingbirds according to the methods of Hadley & Betts (2009). We

then followed hummingbirds on foot using portable receivers (Wildlife Materials Inc TRX

1000S) and handheld Yagi antennae, recorded locations continuously when the bird was found

within receiver range (~250 m in the absence of topographical barriers). In total, we completed

447 hours of telemetry observation, averaging 3.1 ± 0.8 hours per session (typically ~8:00-11:00)

and 5.3 ± 0.9 sessions per site (mean $\pm$ SD). From the recorded locations, we estimated the

proportion of observation time that radio-tagged birds spent in the focal area (Appendix S2).

Thirty-six hummingbirds spent time in the focal area during the 'pre' period and were included

in analyses of the *Heliconia* removal experiment; for a comprehensive summary of radio-

tracking outcomes, see Appendix S2 (Table S4).

Floral visitation rates

To understand hummingbird resource use within each site and to monitor for the presence

of marked hummingbirds, we positioned trail cameras (PlotWatcher Pro, Day 6 Outdoors) near

inflorescences of plant species known or suspected to be visited by hummingbirds. Effort within

each site was concentrated at two locations ('stations', Fig. 1D) located on roughly opposite ends

of the focal area (mean ± SD distance between stations: 60 ± 44 m). Stations were centered

around a focal *Heliconia* plant that we expected to flower throughout the experimental period,

was not in an extremely steep area, and was >10 m from the fragment edge. The two focal

Heliconia plants remained uncovered throughout the entire experimental period to assess how

visitation rates changed after declines in surrounding *Heliconia* density. To partially standardize

plant species composition between control and treatment replicates, we added a floral array

comprising ~3-5 additional plant species at each station (see Appendix S2: Table S5 and

'Pollination success').

Trail cameras took one photograph per second during daylight hours (5:30-17:30) and

automatically combined these photographs into a time-lapse video. We reviewed videos for

hummingbird sightings using the motion detection program MotionMeerkat (Weinstein 2015) or

by manually watching the sped-up video. When hummingbirds were detected on camera, we

identified the species, recorded any color marks, and noted whether they visited any flowers.

From video, we also recorded the number of open flowers available per day. Dates with no

available flowers were excluded from analysis, yielding 20,735 total hours of video (Appendix

S2: Table S6).

Pollination success

To investigate whether *Heliconia* declines influenced pollination success, we examined

styles for the presence of pollen tubes, which indicate pollen grain germination and subsequent

growth toward the flower ovule(s) (Kearns and Inouye 1993). We focused our pollen tube

investigation on the two focal *Heliconia* plants per replicate, as well as several species from the

focal arrays (Appendix S2: Table S5). Throughout the experimental period, we collected flowers

from these focal plants every 1-2 days, which corresponds to the maximum floral longevity for

most of the selected species (*Heliconia tortuosa*, *Hamelia patens*: Stratton 1989; *Stachytarpheta*

frantzii: Thomas-Granger 2003). Styles were fixed in formalin-acetic acid-alcohol for at least 24

hours, stained with aniline blue, and examined for pollen tubes using epifluorescence microscopy

(Kearns and Inouye 1993; Betts, Hadley, and Kress 2015). Due to low pollen tube presence

and/or challenges with pollen tube visualization, we were only able to analyze two plant species:

H. tortuosa (N = 327 styles) and *H. patens* (N = 202 styles). Additional details about style

collection and lab procedures are available in Appendix S2.

Additional response variables (Heliconia removal experiment only)

In addition to the primary response variables described above, we examined several other

metrics for the *Heliconia* removal experiment: (1) recapture probability (i.e., proportion of birds

captured during the 'pre' period and recaptured during the 'post' period), (2) floral visitation

rates from individual birds marked with nail polish, and (3) hummingbird body mass. The first

two response variables assessed the possibility that hummingbirds caught during the initial

capture sessions were rapidly replaced by new individuals, potentially obscuring treatment

effects. To explore whether birds that remained in (or returned to) treatment focal areas were

paying an energetic cost, we also analyzed changes in the body mass of hummingbirds captured

during both experimental periods. To compare body mass change across hummingbirds of

different structural sizes, we calculated body mass relative to the mass predicted by species-

specific allometric equations (i.e., ln(body mass) ~ ln(wing length), Appendix S2: Fig. S4).

Quantifying connectivity

Within each focal landscape, we used Google Earth Pro to digitize closed-canopy forest

and regenerating forest (hereafter 'forest') from high resolution (1 m) Airbus/CNES satellite

imagery. These two forest types were combined because functional connectivity experiments

suggest that regenerating forest is nearly equivalent to closed-canopy forest in facilitating

hummingbird movement (Jackson 2017). We then delineated connected areas of forest

('patches') assuming a hummingbird gap-crossing ability of 50 m (Volpe et al. 2016; Kormann

et al. 2016). Lastly, we quantified functional connectivity as the amount of reachable forest

within each focal landscape, following a modified version of Hanski's (1994) incidence function

model for metapopulation connectivity. Substituting forested pixels for patches used in the

original model, we assigned exponentially lower importance to pixels located farther away from

the focal area (Fig. 1E, Appendix S2: Fig. S5). These weights were calculated as:

299                               $W_j = \exp(-\alpha\, d_{ij})$,   **(Eq. 1)**

where α is the decay rate parameter that scales the effect of distance and d is the shortest

cumulative distance between the focal area centroid i and each forested pixel j, staying within the

patch boundary. The decay rate parameter is inversely related to the average movement distance

of the focal species (Moilanen and Nieminen 2002) and was calculated using the mean home

range length of green hermit hummingbirds (282 m: Volpe et al. 2016). Then, to calculate

functional connectivity, we then summed the weighted pixel areas as follows:

306                               $S_i = \Sigma\, W_j\, A_{j,}$ **(Eq. 2)**

where W_j is the weight of each forested pixel and A is the area of each forested pixel. Geospatial

processing was conducted using Google Earth Engine through the JavaScript API (Gorelick et al.

2017).

Quantifying resource availability

From the floral resource surveys (see *'Study design'*), we estimated the energetic value of

the *Heliconia* flowers and the remaining non-*Heliconia* resources. Briefly, we converted

resource counts (i.e., number flowers and inflorescences) into calories, the most biologically

relevant unit for hummingbird energetics (e.g., Wolf, Hainsworth, and Stiles 1972; Stiles 1975).

To calculate calories, we measured nectar volume and concentration from >1400 flowers ($N = 38$

species) growing in the focal landscape and/or the Wilson Botanical Garden at the Las Cruces

Biological Station, supplementing these measurements with literature data when necessary. We

then converted average nectar volume and concentration per flower to calories per plant (see

Appendix S2 for full details). Because tropical hummingbirds generally interact with subsets of

the available floral resources (e.g., Stiles 1985; Maglianesi et al. 2014; Vizentin-Bugoni,

Maruyama, and Sazima 2014), we then generated estimates of caloric availability specific to

each hummingbird group examined. These estimates were developed prior to any analysis based

on relative visitation rates of each hummingbird species to 37 flowering plant species (Fig. 3).

We divided these calorie estimates by focal area size, then natural log-transformed the resulting

densities. Floral corolla length was measured from flowers collected in the field, supplemented

with literature values when necessary (Appendix S2: Table S16).

Statistical analysis

We used linear mixed models (LMMs) and generalized linear mixed models (GLMMs) to

analyze how experimental *Heliconia* extinction and natural variation in *Heliconia* density

influenced our primary response variables: (i) captured hummingbird abundance, (ii) proportion

of time spent in focal area, (iii) visitation rates to flowering plants (*Heliconia* and non-

Heliconia), and (iv) plant pollination success. Analyses of the *Heliconia* removal experiment

also included several additional, individual-level response variables (see '*Additional response*

variables'). To determine whether declines in *Heliconia* affected *Heliconia* specialists more

strongly than other hummingbird species (**H4**), we analyzed all hummingbird responses with (i)

the full dataset containing all species, and (ii) a dataset including green hermits and violet

sabrewings.

We modeled each response variable using appropriate statistical distributions and

controlled for variable effort when applicable (Appendix S2: Tables S17-S18). All statistical

analyses were conducted in R version 4.1.1 (R Core Team 2021) using 'glmmTMB' (Brooks et

al. 2017). For LMMs, we checked for normality, homoscedasticity, and linearity using the

'performance' package (Lüdecke et al. 2021). For GLMMs, we checked for overdispersion, zero-

inflation, and linearity by visually inspecting residual plots from 'DHARMa' (Hartig 2020).

Estimated marginal means were calculated using 'emmeans' (Lenth 2020) and visualized using

the 'ggemmeans' function within 'ggeffects' (Lüdecke 2018). Results were considered

statistically significant if $P < 0.05$.

Heliconia removal experiment

In a BACI experimental design, an overall treatment effect is indicated by a statistically

significant interaction between the variables representing treatment and experimental period

(Underwood 1994; McDonald, Erickson, and McDonald 2000). We adopted this standard

analytical approach rather than calculating change scores (e.g., pre-post differences) because

change scores are nearly always negatively correlated with baseline measurements; when

baseline measurements also covary with predictor variables of interest, misleading (i.e.,

spuriously negative) relationships can emerge (Allison 1990). Therefore, we tested the *parallel*

extinction hypothesis (**H1a**) and *robust network hypothesis* (**H1b**) using a two-way interaction

between *Heliconia* removal and period (pre *versus* post) for all response variables measured

during both experimental periods (recaptures were measured in the post period only). All models

included a random intercept for 'Site'. Except when modeling recapture probability, we also

included a nested random effect for 'Replicate ID' (Site-Year), which paired pre-post

observations within each replicate. Additional nested random effects, namely for plant species

and individual birds, were included where applicable (Appendix S2: Table S17). Because we

were interested in pre-post changes in our response variables, we only included covariates that

varied between pre and post periods. For example, when analyzing hummingbird visitation rates,

we accounted for the attractiveness of a given focal plant by including a covariate for the number

of open flowers per day, averaged across the experimental period. Since the magnitude of our

experimental *Heliconia* extinction varied widely (see '*Results*'), we ran parallel models using a

categorical treatment variable (Control/Treatment) and a quantitative treatment variable

(Calories removed/hectare). Because results were qualitatively similar, only results from the

categorical approach are presented in the main text.

Natural variation in Heliconia density

To complement the *Heliconia* removal experiment, we also examined hummingbird

responses over a natural gradient in *Heliconia* availability using data from the pre period of the

experimental sites. This 'space-for-time' approach elucidates how hummingbird space use and

foraging behavior respond to natural variation present in the study area – presumably emerging

over longer time scales than those of our removal experiment. We also used this approach to test

the connectivity hypothesis and resource hypothesis by modeling the interaction term between

Heliconia density and either connectivity (**H2**) or non-*Heliconia* resource availability (**H3**).

These hypotheses were not tested using the experimental framework due to limited sample size

(and thus statistical power) to enable estimation of a three-way interaction between treatment,

experimental period, and resources/connectivity. Because resource availability data were not

collected in 2016 control replicates, these replicates ($N = 6$) were excluded from this analysis.

In these analyses, we included several additional covariates not included in the analyses of the

Heliconia removal experiment. First, we used estimates of non-*Heliconia* resource availability

tailored to each hummingbird group being examined. Because resource availability was

moderately correlated with connectivity (Pearson's $r = 0.57$, $P = 0.002$, Appendix S2: Figure

S6), we included both covariates in all models to statistically control for their independent

effects. Analyses of radio telemetry data (i.e., proportion of time spent in focal area) included a

covariate for the focal area size. When analyzing visitation rates of *Heliconia* specialists to non-

Heliconia resources, we also included plant corolla length as a covariate, since these relatively

long-billed hummingbirds (mean bill lengths of 40 and 33 mm: Borgella, Snow, and Gavin 2001)

are expected to select plant species with long corolla tubes (e.g., Maglianesi et al. 2014;

Vizentin-Bugoni, Maruyama, and Sazima 2014; Dalsgaard et al. 2021). We included random

effects for Site in all models; additional random effects were included where applicable

(Appendix S2: Table S18).

RESULTS

Across all study sites and years, *Heliconia* density ranged from 3 to 1131 inflorescences

per ha (mean ± SD: 186 ± 228, median: 145, Appendix S3: Table S1). The magnitude of our

experiment therefore varied widely; we removed between 8 and 526 *Heliconia* inflorescences per

treatment replicate (mean ± SD: 172 ± 138, median: 138, Appendix S3: Table S2). These

inflorescences represented, on average, 96% (± 5.5%) of the *Heliconia* inflorescences

encountered during removal (two focal plants remained uncovered and sometimes plant

inaccessibility limited our capacity to cover). Sizes of focal treatment areas ranged from 0.27 to

7.49 ha (mean ± SD: 1.7 ± 1.9, median: 1.25, Appendix S3: Table S2), depending on the size of

the forest fragment and the navigability of the terrain. We estimate that, on average, the covered

Heliconia plants supplied 42.5% (± 30.9%, range: 0.9 – 90.4%) of the calories available for the

entire hummingbird community (N = 19 species, Appendix S3: Table S3) and at least 65.5% (±

31.3%, range: 2.4-98%) of calories for *Heliconia* specialists (Fig. 2C, Appendix S3: Table S4).

Focal *Heliconia* plants were visited by hummingbirds 6.4 ± 5.7 times per day (range: 0-25,

median: 5), with green hermits visiting most frequently (75% of 1,065 sightings).

Overall responses to changes in Heliconia density (H1)

Hummingbird behavior and pollination success were remarkably resistant to

experimentally induced *Heliconia* extinction. Despite the magnitude of *Heliconia* removal and

confirmed use of *Heliconia* by hummingbirds, we did not detect an overall effect of local

Heliconia removal hummingbird space use, floral visitation rates, or plant pollination success.

However, when examining these response variables across a natural gradient in *Heliconia*

availability (i.e., the space-for-time approach), we found that floral visitation rates and

pollination success declined as *Heliconia* became scarce. We provide detailed evidence for these

findings below.

Hummingbird captures & radio telemetry

Overall, we did not find that experimental *Heliconia* removal decreased hummingbird

abundance or the proportion of time that radio-tagged hummingbirds spent in the focal area.

Although we caught 23% fewer hummingbirds in the 'post' capture session than the 'pre'

capture session in treatment replicates (95% CI: -41% to 1%), this pre-post decline fell within the

variability of the control replicates (all species: $z = -1.11$, $P = 0.27$; *Heliconia* specialists: $z = -$

0.5, $P = 0.62$; Fig. 4: A-B; Appendix S3: Table S5). Similarly, although radio-tagged

hummingbirds spent 25-30% less time in focal areas during the 'post' period, we found no

evidence that pre-post changes in space use were linked to our experimental manipulation (95%

CI, treatment: -56% to 11%; 95% CI, control: -56% to 28%). Results were similar when

examining all species together (Fig. 4C) and *Heliconia* specialists separately (Fig. 4D). We also

found no evidence that natural variation in *Heliconia* density influenced the hummingbird space

use, quantified either with mist net captures or radio telemetry (Appendix S3: Tables S5-S6).

434   When examining individual-level hummingbird responses, we also did not detect changes

associated with *Heliconia* removal. On average, the estimated probability of recapturing a

hummingbird from the 'pre' capture session was 18% in control replicates (95% CI: 11% to 29

%), compared to 14% in treatment replicates (95% CI: 97% to 23%). This difference was not

statistically significant, and recaptures of *Heliconia* specialists showed qualitatively similar

results (Appendix S3: Fig. S1: A-B, Table S7). Among the individual hummingbirds caught

during both sessions, we also found no evidence that pre-post changes in the relative body mass

differed between control and treatment replicates (Appendix S3: Table S8).

Floral visitation rates & pollination success

443   Using trail cameras, we continuously monitored hummingbird visitation at the two focal

Heliconia inflorescences that remained uncovered throughout the experimental period, as well as

numerous other flowering plants (mean ± SD: 9.3 ± 2 species/replicate; range: 6-14). Under the

robust network hypothesis, we expected hummingbirds to rewire and increasingly use alternative

(non-*Heliconia*) resources. However, we did not find that pre-post changes in hummingbird

visitation rate differed between control and treatment replicates, either when examining the focal

Heliconia plants ($z = 0.83$, $P = 0.41$, Fig. 5: A-B, Appendix S3: Table S9) or non-*Heliconia* plant

species ($z = -0.69$, $P = 0.49$, Fig. 6: A-B, Appendix S3: Table S10). Responses by *Heliconia*

specialists mirrored the responses observed for all species, although these species visited non-

Heliconia species relatively infrequently (<1 visit/day, on average: Fig. 6B). Results were similar

when examining pre-post changes in visitation rates for individual color-marked hummingbirds

observed on camera (Appendix S3: Fig. S1: C-F, Tables S11-S12).

In contrast to the experimental results, we found that natural variation in *Heliconia*

availability strongly influenced visitation to focal *Heliconia* plants (Fig. 5: D-E). For every

standard deviation decrease in the caloric density of *Heliconia*, we found that overall *Heliconia*

visitation rates decreased by 49% (95% CI: 15% to 93%, $z = 3.03$, $P = 0.0024$, Appendix S3:

Table S9). Visitation to non-*Heliconia* plants also declined when natural *Heliconia* availability

was low, but only for *Heliconia* specialists ($z = 2.02$, $P = 0.044$, Fig. 6E, Appendix S3: Table

S10).

We found no evidence that experimental *Heliconia* removal influenced pollination

success for the focal *Heliconia* plants ($z = 0.23$, $P = 0.82$, Fig. 5C, Appendix S3: Table S13) or

another plant species from the floral arrays (*Hamelia patens:* $z = 0.19$, $P = 0.85$, Appendix S3:

Table S13). However, when *Heliconia* availability was naturally low, we observed declines in

the pollination success of the focal *Heliconia* plants; for every standard deviation decrease in the

caloric density of *Heliconia*, the odds of pollination declined by 2.4 times (95% CI: 1.1 times to

5.4 times, $z = 2.2$, $P = 0.028$, Fig. 5F).

Effects of connectivity (H2) and resource availability (H3)

Hummingbird captures & radio telemetry

Connectivity influenced hummingbird abundance; hummingbird captures increased by

6% for every one-unit increase in connectivity (95% CI: +1% to +12%, $z = 2.31$, $P = 0.021$,

Appendix S3: Table S5). Further, radio-tagged birds spent more time in focal areas when

alternative (non-*Heliconia*) resources were abundant; the odds of being in the focal area

increased by 2.3 times (95% CI: 1.3 to 4.1 times) for every standard deviation increase in caloric

density (all species only: $z = 2.78$, $P = 0.0055$, Appendix S3: Table S6). However, neither

connectivity or resource availability appeared to influence hummingbird responses to *Heliconia*

declines, either with experimental removal or across the natural density gradient ($P > 0.05$ for all

interactions of interest, Appendix S3: Tables S5-S6).

Floral visitation rates & pollination success

We did not observe any statistically significant main effects of connectivity on

hummingbird visitation rates, either to focal *Heliconia* plants or non-*Heliconia* species.

However, after *Heliconia* removal, visitation rates to focal *Heliconia* plants by *Heliconia*

specialists declined more sharply when connectivity was low ($z = -1.76$, $P = 0.08$, Appendix S3:

Fig. S2, Table S9). This potential connectivity effect did not emerge when examining *Heliconia*

pollination success ($z = -0.55$, $P = 0.58$). Although we expected that interaction rewiring might

be more likely to occur when alternative resources were abundant, we did not find that *Heliconia*

declines in otherwise resource-rich areas caused hummingbirds to rely increasingly on non-

Heliconia species (Appendix S3: Table S10).

DISCUSSION

In this study, we experimentally induced the local extinction of a locally abundant

tropical plant, *Heliconia tortuosa*, to assess the robustness of plant-hummingbird interaction

networks in a fragmented tropical landscape. Additionally, across a natural gradient in *Heliconia*

density, we assessed whether low *Heliconia* abundance would lead to parallel declines in

hummingbirds and their pollination services – and whether landscape context (connectivity) and

local context (alternative resource availability) could buffer against potential declines. Based on

suite of complementary response variables we found that, even when very high proportions of

total available calories were experimentally removed (~43-66% on average), hummingbird

responses fell within the natural variability of control replicates. As might be expected with such

small changes in hummingbird behavior, we did not observe any cascading effects of

experimental manipulation on pollination success of flowering plants, which supports the *robust*

network hypothesis (**H1b**). On the other hand, when examining hummingbird responses and

pollination success across a natural gradient in *Heliconia* density, we found evidence for the

parallel extinction hypothesis (**H1a**). In sites where *Heliconia* density was low, hummingbirds

visited individual *Heliconia* plants less frequently, and these plants experienced reduced

pollination success. Other flowering plants also exhibited lower visitation rates from *Heliconia*-

associated hummingbirds, suggesting that high *Heliconia* densities may attract these

hummingbirds (and/or incentivize repeat visitation) and therefore facilitate pollination for other

species.

Our finding that that experimental removal of *Heliconia* did not elicit parallel extinctions

in pollinators aligns with theoretical predictions (Memmott, Waser, and Price 2004; Kaiser-

Bunbury et al. 2010) and several other experiments that removed plant species from mutualistic

networks (Ferrero et al. 2013; Goldstein and Zych 2016; Costa et al. 2018). However, if

hummingbirds remained in forest fragments despite *Heliconia* declines, we assumed that they

would rewire to use alternative floral resources; hummingbirds as a taxonomic group are highly

opportunistic foragers (Arizmendi and Ornelas 1990; Maruyama et al. 2013) and the most

frequent *Heliconia* visitor, the green hermit, has a relatively wide niche breadth (Fig. 3, see also

Borgella, Snow, and Gavin 2001; Betts, Hadley, and Kress 2015). Unexpectedly, we found no

evidence that hummingbirds increasingly visited non-*Heliconia* resources when *Heliconia*

became less available, suggesting that the mechanisms underlying network robustness in this

study may not necessarily align with the rewiring observed in previous empirical tests (e.g.,

Timóteo et al. 2016; Kaiser-Bunbury et al. 2017; Costa et al. 2018). This apparent lack of

foraging flexibility may emerge from various constraints, such as morphological trait-matching,

search costs, and/or competition from other hummingbirds (see also Weinstein and Graham

2017). For example, high levels of exploitative competition for non-defendable resources may

have discouraged hummingbirds from modifying their trapline and risking overlap with another

bird's foraging route (Young 1971; Gill 1988; Ohashi and Thomson 2009). Lastly, is possible

that fine-scale rewiring did occur but was not detected using our analytical approach, which

aggregated visitation rates to the level of hummingbird species; further analysis examining the

diversity of pollen grains carried by individual hummingbirds may shed light on the possibility

of rewiring at finer scales.

In the absence of rewiring, the resilience in hummingbird behavior, abundance, and

pollination services following experimental *Heliconia* removal is surprising. Brief pauses in

nectar intake can disrupt a hummingbird's daily energy balance, given their high metabolic rate

and limited energy reserves (Calder and Booser 1973; Tooze and Gass 1985; Brice 1992).

Moreover, previous nectar manipulations with hummingbirds – albeit within defended territories

– led to rapid changes in territory size, foraging patterns, and/or defense behavior, generally

within a day (e.g., Hixon, Carpenter, and Paton 1983; Gass and Sutherland 1985; Eberhard and

Ewald 1994; Temeles et al. 2004). One explanation for our result is that hummingbirds

permanently emigrated from fragments after *Heliconia* removal but, because of high intruder

pressure, were quickly replaced by new individuals (Stiles and Wolf 1970). However, we

monitored individual birds using several different approaches (radio telemetry, recaptures,

visitation rates) and found no evidence that *Heliconia* removal caused emigration and subsequent

replacement. An alternative possibility is that hummingbirds relied on physiological adaptations,

such as a torpor and/or insectivory, to maintain body condition despite large declines in nectar

availability – resulting in the stable body mass measurements we observed in recaptured

hummingbirds. Torpor is an energy-saving adaptation characterized by declines in body

temperature and metabolic rate (Hainsworth, Collins, and Wolf 1977; Spence and Tingley 2021),

and even tropical hummingbirds inhabiting low-to-mid elevations appear to possess this

capability, including the green hermit (Krüger, Prinzinger, and Schuchmann 1982; Bech et al.

1997; Shankar et al. 2020; Schuchmann and Prinzinger 1988). Additionally, hummingbirds are

highly insectivorous and, although they cannot completely replace nectar with insects (Brice

1992; Stiles 1995), they may supplement their diet with arthropods when nectar becomes scarce

(Young 1971; Kuban and Neill 1980; Montgomerie and Redsell 1980; Hazlehurst and Karubian

2018). Finding torpid hummingbirds at night is extremely challenging (Carpenter and Hixon

1988), but future research could explore changes in arthropod consumption using techniques

such as DNA metabarcoding (Moran, Prosser, and Moran 2019).

Another explanation for hummingbird persistence following experimental *Heliconia*

removal involves their high movement ability. *H. tortuosa* is a patchily distributed species that

necessitates long-distance foraging movements, as a single clump does not make territoriality

energetically profitable (Stiles 1975). Densities can also vary substantially across short distances;

for example, Stiles (1979) notes that some Costa Rican *Heliconia* species can be common in one

watershed, yet nearly absent in an adjacent watershed < 1 km away (see also Bruna and Kress

2002). The most frequent *Heliconia* visitor, the green hermit, is therefore capable of traveling

long distances (>500m) while foraging within home ranges of >150 ha, although home range

sizes of ~2 ha are more typical (Volpe et al. 2014; 2016). Thus, returning to our focal areas

(typically ~1.5 ha) to visit the few remaining *Heliconia* resources could be perceived as a

relatively low-cost option for long-distance trapliners, at least in the short term. Notably,

although highly mobile pollinators might buffer against network collapse (Volpe et al. 2016),

they may be vulnerable to habitat loss and fragmentation once thresholds are crossed, leading to

parallel pollinator declines observed in nature (Andrén 1994; Hadley and Betts 2012). This

strategy is also unlikely to be profitable over longer time scales than measured by our

experiment, which only removed *Heliconia* for five days.

The relatively short length of our removal experiment might also explain why – when

examining hummingbird responses across a natural gradient in *Heliconia* availability – lower

Heliconia densities were indeed associated with reduced floral visitation rates. Although a short

adjustment period (<1 day) is standard in similar species removal experiments (e.g., Hixon,

Carpenter, and Paton 1983; Brosi and Briggs 2013; Hazlehurst and Karubian 2018), extending

the experimental timeline may have more realistically simulated a future extinction and provided

more time for behavioral or demographic adjustment. If hummingbirds were not permanent

residents of our focal areas – as suggested by our relatively low recapture rates (~15%) and

proportion of time spent in the focal area (~20%) – then transient individuals might have not

immediately noticed changes in *Heliconia* density and changed their foraging patterns

accordingly. Likewise, hummingbirds might not have persisted had we removed *Heliconia* over

larger areas, because traveling to isolated plants would become even more energetically costly

(Heinrich and Raven 1972). Unfortunately, covering (and then uncovering) *Heliconia* plants in

steep terrain with dense understories limited the area we could manipulate.

We also acknowledge that our ability to detect experimental effects may have been

limited by low statistical power, owing to high individual-level variation and limited sample size.

Low statistical power may have also weakened our tests of the connectivity hypothesis and

resource hypothesis, as those datasets only included a subset of the replicates included in the

experimental analyses (resource availability data were not available for all sites). Still, we

emphasize that the spatial scale of our experiment and amount of replication surpasses most

other empirical studies simulating species extinctions within plant-pollinator networks

(Appendix S1: Table S1). Nevertheless, future removal experiments with highly mobile

pollinators should consider increasing the amount of replication, and, if possible, manipulate

larger areas.

In the space-for-time component of our study, we found that visitation rates to individual

plants decreased with natural declines in *Heliconia* density, suggesting that *Heliconia* and its

pollinators might facilitate pollination of co-flowering plants in this study system. Indeed,

facilitative effects arising from the overall attractiveness of the floral neighborhood are expected

to operate most strongly in systems where plants are spatially dispersed and pollinator density is

low (Schemske 1981; Ghazoul 2005; Moeller 2004), as is the case in the tropical forest

understory. If *Heliconia* population densities decrease due to habitat fragmentation (Hadley et al.

2014), plants visited exclusively by *Heliconia*-associated hummingbirds might therefore

experience a higher risk of reproductive failure (e.g., *Pitcairnia imbricata*, *Calathea*

guzmanoides, *Aphelandra golfodulcensis*, Fig. 3). Whether visitation rates map onto pollination

success for these species remains an open question, since pollinator sharing may lead to

heterospecific pollen transfer and pollen loss (Ashman and Arceo-Gómez 2013; Morales and

Traveset 2008) – tipping the balance toward interspecific competition rather than facilitation

(Ghazoul 2005; Mitchell et al. 2009; Rathcke 1983). However, we did discover a positive

relationship between visitation rate and pollination success for *Heliconia*. This relationship has

been predicted for other *Heliconia* species but yielded mixed results (Stein et al. 2013; Bruna et

al. 2004), in accordance with accumulating evidence that facilitation effects can be highly

context-specific (Bergamo et al. 2020). Nevertheless, this finding suggests that any reproductive

declines of *Heliconia* in this study system might accelerate once extinction debt is realized and

conspecific density begins to decline.

CONCLUSIONS

Although pollinators seemed able to persist during our experiments that removed floral

resources for a short period of time (several days), this persistence may not be sustainable or

reflect the outcome of a natural extinction process. Indeed, we observed declines in

hummingbird visitation when *Heliconia* availability was naturally low; this reduced

hummingbird visitation had cascading consequences for pollination success. Thus, longer-term

experiments could be key for predicting consequences of species loss and understanding

mechanisms of network robustness – at least for pollinator species that forage at large spatial

scales. Our study also suggests that, over the short term, behavioral flexibility to use alternative

floral resources might be limited, even in taxonomic groups thought to be highly opportunistic

foragers. Thus, even rewiring rules with strong biological underpinnings (e.g., Vizentin-Bugoni

et al. 2020) may not necessarily predict real-world responses to species loss, perhaps due to

additional constraints such as competition. To further understand the mechanisms underlying

network robustness or collapse, we encourage future experiments that consider how temporal

scale, species traits (e.g., movement ability), competition, and landscape context interact to

determine the outcome of species extinctions. These experiments could be especially valuable at

geographic range limits, where climate change will likely lead to species replacements

(HilleRisLambers et al. 2013), and in the tropics where landscape change and species loss are

most rapid (Hansen et al. 2013).

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation (NSF) Graduate

Research Fellowship Program under Grant No. 1840998 to KGL. Funding for the experiment

was provided primarily by NSF-DEB-1457837 to MGB and ASH, with additional grants to KGL

from the American Ornithological Society and Sigma Xi. We also thank the Las Cruces

Biological Station and the private landowners who allowed us to work on their property. Field

assistance was provided by Jorge Araya Paniagua, Michael Atencio Picado, Marion Donald,

Dustin Gannon, Jessica Greer, Matt Hadley, Urs Kormann, Daniel Moreno, Mauricio Paniagua

Castro, Ignacio Ruiz Ilama, Esteban Sandi Paniagua, Andrés Solorzano Vargas, Laura Sutcliffe,

and Felipe Torres Vanegas. Videos from trail cameras were reviewed with the help of Marion

Donald, Colleen Franklin, Jessica Greer, Zoe Griffith, Elena Hart, Billy Hilgert, Zachary

Kendall, Jeremy Lee, Ana Medina Roman, Briley Mullin, Amber Newell, Thanh Nguyen, Zoe

Sallada, Allison Simoni, Claire Woods, and Mel Xiao. Nozomi Birkett and Marion Donald

helped collect nectar measurements, and Chandler Spruill assisted with pollen tube processing.

Ariel Muldoon provided valuable statistical advice, and Mark Novak provided feedback on an

earlier version of the manuscript. Response variable icons were created from the following Noun

Project images: Hummingbird by Agne Alesiute, net by Guillaume Kurkdjian, Stopwatch by

Thomas Marijinissen, leaves by Tatiana, Camera by Stanislav Levin, and Microscope by Gregor

Cresnar.

LITERATURE CITED

Aguilar, R., L. Ashworth, L. Galetto, and M.A. Aizen. 2006. "Plant Reproductive Susceptibility

to Habitat Fragmentation: Review and Synthesis through a Meta-Analysis." *Ecology*

Letters 9 (8): 968–80.

Aizen, M.A., M. Sabatino, and J.M. Tylianakis. 2012. "Specialization and Rarity Predict

Nonrandom Loss of Interactions from Mutualist Networks." *Science* 335 (6075): 1486–

89.

Allen-Wardell, G., P. Bernhardt, R. Bitner, A. Burquez, S. Buchmann, J. Cane, P.A. Cox, et al.

1998. "The Potential Consequences of Pollinator Declines on the Conservation of

Biodiversity and Stability of Food Crop Yields." *Conservation Biology* 12 (1): 8–17.

Allison, P.D. 1990. "Change Scores as Dependent Variables in Regression Analysis."

Sociological Methodology 20: 93–114.

Anderson, S.H., D. Kelly, J.J. Ladley, S. Molloy, and J. Terry. 2011. "Cascading Effects of Bird

Functional Extinction Reduce Pollination and Plant Density." *Science* 331 (6020): 1068–

71.

Andrén, H. 1994. "Effects of Habitat Fragmentation on Birds and Mammals in Landscapes with

Different Proportions of Suitable Habitat: A Review." *Oikos* 71 (3): 355–66.

Arizmendi, M.C., and J.F. Ornelas. 1990. "Hummingbirds and Their Floral Resources in a

Tropical Dry Forest in Mexico." *Biotropica* 22 (2): 172–80.

675 Ashman, T.L., and G. Arceo-Gómez. 2013. "Toward a Predictive Understanding of the Fitness

676    Costs of Heterospecific Pollen Receipt and Its Importance in Co-Flowering

677    Communities." *American Journal of Botany* 100 (6): 1061–70.

678 Ashman, T.L., T.M. Knight, J.A. Steets, P. Amarasekare, M. Burd, D.R. Campbell, M.R.

679    Dudash, M.O. Johnston, S.J. Mazer, and R.J. Mitchell. 2004. "Pollen Limitation of Plant

680    Reproduction: Ecological and Evolutionary Causes and Consequences." *Ecology* 85 (9):

681    2408–21.

682 Aslan, C.E., E.S. Zavaleta, B. Tershy, and D. Croll. 2013. "Mutualism Disruption Threatens

683    Global Plant Biodiversity: A Systematic Review." *PLOS ONE* 8 (6): e66993.

684 Bascompte, J., and P. Jordano. 2007. "Plant-Animal Mutualistic Networks: The Architecture of

685    Biodiversity." *Annual Review of Ecology, Evolution, and Systematics* 38 (1): 567–93.

686 Bascompte, J., P. Jordano, C.J. Melián, and J.M. Olesen. 2003. "The Nested Assembly of Plant–

687    Animal Mutualistic Networks." *Proceedings of the National Academy of Sciences* 100

688    (16): 9383–87.

689 Bascompte, J., P. Jordano, and J.M. Olesen. 2006. "Asymmetric Coevolutionary Networks

690    Facilitate Biodiversity Maintenance." *Science* 312 (5772): 431–33.

691 Bech, C., A.S. Abe, J.F. Steffensen, M. Berger, and J.E.P.W. Bicudo. 1997. "Torpor in Three

692    Species of Brazilian Hummingbirds under Semi-Natural Conditions." *The Condor* 99 (3):

693    780–88.

694 Bergamo, P.J., N.S. Streher, A. Traveset, M. Wolowski, and M. Sazima. 2020. "Pollination

695    Outcomes Reveal Negative Density-Dependence Coupled with Interspecific Facilitation

696    among Plants." *Ecology Letters* 23 (1): 129–39.

697   Betts, M.G., A.S. Hadley, and W.J. Kress. 2015. "Pollinator Recognition by a Keystone Tropical

698   Plant." *Proceedings of the National Academy of Sciences* 112 (11): 3433–38.

699   Biesmeijer, J.C., S.P.M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A.P.

700   Schaffers, et al. 2006. "Parallel Declines in Pollinators and Insect-Pollinated Plants in

701   Britain and the Netherlands." *Science* 313 (5785): 351–54.

702   Borgella, R., A.A. Snow, and T.A. Gavin. 2001. "Species Richness and Pollen Loads of

703   Hummingbirds Using Forest Fragments in Southern Costa Rica." *Biotropica* 33 (1): 90–

704   109.

705   Brice, A.T. 1992. "The Essentiality of Nectar and Arthropods in the Diet of the Anna's

706   Hummingbird (Calypte Anna)." *Comparative Biochemistry and Physiology Part A:

707   Physiology* 101 (1): 151–55.

708   Brooks, M.E., K. Kristensen, K.J. van Benthem, A. Magnusson, C.W. Berg, A. Nielsen, H.J.

709   Skaug, M. Maechler, and B.M. Bolker. 2017. "GlmmTMB Balances Speed and

710   Flexibility Among Packages for Zero-Inflated Generalized Linear Mixed Modeling." *The

711   R Journal* 9 (2): 378–400.

712   Brosi, B.J., and H.M. Briggs. 2013. "Single Pollinator Species Losses Reduce Floral Fidelity and

713   Plant Reproductive Function." *Proceedings of the National Academy of Sciences* 110

714   (32): 13044–48.

715   Bruna, E.M., and W.J. Kress. 2002. "Habitat Fragmentation and the Demographic Structure of

716   an Amazonian Understory Herb (Heliconia Acuminata)." *Conservation Biology* 16 (5):

717   1256–66.

Bruna, E.M., W.J. Kress, F. Marques, and O.F. da Silva. 2004. "Heliconia Acuminata

Reproductive Success Is Independent of Local Floral Density." *Acta Amazonica* 34 (3):

467–71.

Calder, W.A., and J. Booser. 1973. "Hypothermia of Broad-Tailed Hummingbirds during

Incubation in Nature with Ecological Correlations." *Science* 180 (4087): 751–53.

Carpenter, F.L., and M.A. Hixon. 1988. "A New Function for Torpor: Fat Conservation in a

Wild Migrant Hummingbird." *The Condor* 90 (2): 373–78.

Colwell, R.K., R.R. Dunn, and N.C. Harris. 2012. "Coextinction and Persistence of Dependent

Species in a Changing World." *Annual Review of Ecology, Evolution, and Systematics* 43

(1): 183–203.

Costa, J.M., J.A. Ramos, L.P. da Silva, S. Timóteo, P. Andrade, P.M. Araújo, C. Carneiro, et al.

2018. "Rewiring of Experimentally Disturbed Seed Dispersal Networks Might Lead to

Unexpected Network Configurations." *Basic and Applied Ecology* 30: 11–22.

Dalsgaard, B., P.K. Maruyama, J. Sonne, K. Hansen, T.B. Zanata, S. Abrahamczyk, R. Alarcón,

et al. 2021. "The Influence of Biogeographical and Evolutionary Histories on

Morphological Trait-Matching and Resource Specialization in Mutualistic

Hummingbird–Plant Networks." *Functional Ecology* 35 (5): 1120–33.

Dunn, R.R., N.C. Harris, R.K. Colwell, L.P. Koh, and N.S. Sodhi. 2009. "The Sixth Mass

Coextinction: Are Most Endangered Species Parasites and Mutualists?" *Proceedings of

the Royal Society B: Biological Sciences* 276 (1670): 3037–45.

Eberhard, J.R., and P.W. Ewald. 1994. "Food Availability, Intrusion Pressure and Territory Size:

An Experimental Study of Anna's Hummingbirds (Calypte Anna)." *Behavioral Ecology

and Sociobiology* 34 (1): 11–18.

Farwig, N., D. Bailey, E. Bochud, J.D. Herrmann, E. Kindler, N. Reusser, C. Schüepp, and M.H.

Schmidt-Entling. 2009. "Isolation from Forest Reduces Pollination, Seed Predation and

Insect Scavenging in Swiss Farmland." *Landscape Ecology* 24 (7): 919–27.

Feinsinger, P., and R.K. Colwell. 1978. "Community Organization among Neotropical Nectar-

Feeding Birds." *American Zoologist* 18 (4): 779–95.

Ferrero, V., S. Castro, J. Costa, P. Acuña, L. Navarro, and J. Loureiro. 2013. "Effect of Invader

Removal: Pollinators Stay but Some Native Plants Miss Their New Friend." *Biological*

Invasions 15 (10): 2347–58.

Garibaldi, L.A., I. Steffan-Dewenter, C. Kremen, J.M. Morales, R. Bommarco, S.A.

Cunningham, L.G. Carvalheiro, N.P. Chacoff, J.H. Dudenhöffer, and S.S. Greenleaf.

2011. "Stability of Pollination Services Decreases with Isolation from Natural Areas

despite Honey Bee Visits." *Ecology Letters* 14 (10): 1062–72.

Gass, C.L., and G.D. Sutherland. 1985. "Specialization by Territorial Hummingbirds on

Experimentally Enriched Patches of Flowers: Energetic Profitability and Learning."

Canadian Journal of Zoology 63 (9): 2125–33.

Ghazoul, J. 2005. "Pollen and Seed Dispersal among Dispersed Plants." *Biological Reviews* 80

(03): 413–43.

Gilbert, L.E. 1980. "Food Web Organization and Conservation of Neotropical Diversity." In

Conservation Biology: An Evolutionary-Ecological Perspective, edited by M.E. Soulé

and B.A. Wilcox, 11–33. Sunderland: Sinauer.

Gill, F.B. 1988. "Trapline Foraging by Hermit Hummingbirds: Competition for an Undefended,

Renewable Resource." *Ecology* 69 (6): 1933–42.

Goldstein, J., and M. Zych. 2016. "What If We Lose a Hub? Experimental Testing of Pollination

Network Resilience to Removal of Keystone Floral Resources." *Arthropod-Plant*

Interactions 10 (3): 263–71.

Gorelick, N., M. Hancher, M. Dixon, S. Ilyushchenko, D. Thau, and R. Moore. 2017. "Google

Earth Engine: Planetary-Scale Geospatial Analysis for Everyone." *Remote Sensing of*

Environment 202: 18–27.

Hadley, A.S., and M.G. Betts. 2009. "Tropical Deforestation Alters Hummingbird Movement

Patterns." *Biology Letters* 5 (2): 207–10.

———. 2012. "The Effects of Landscape Fragmentation on Pollination Dynamics: Absence of

Evidence Not Evidence of Absence." *Biological Reviews* 87 (3): 526–44.

Hadley, A.S., S.J.K. Frey, W.D. Robinson, and M.G. Betts. 2018. "Forest Fragmentation and

Loss Reduce Richness, Availability, and Specialization in Tropical Hummingbird

Communities." *Biotropica* 50 (1): 74–83.

Hadley, A.S., S.J.K. Frey, W.D. Robinson, W.J. Kress, and M.G. Betts. 2014. "Tropical Forest

Fragmentation Limits Pollination of a Keystone Understory Herb." *Ecology* 95 (8):

2202–12.

Hainsworth, F.R., B.G. Collins, and L.L. Wolf. 1977. "The Function of Torpor in

Hummingbirds." *Physiological Zoology* 50 (3): 215–22.

Hanski, I. 1994. "Patch-Occupancy Dynamics in Fragmented Landscapes." *Trends in Ecology &*

Evolution 9 (4): 131–35.

———. 1998. "Metapopulation Dynamics." *Nature* 396 (6706): 41–49.

Hartig, F. 2020. *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*

Regression Models. https://CRAN.R-project.org/package=DHARMa.

Hazlehurst, J.A., and J.O. Karubian. 2018. "Impacts of Nectar Robbing on the Foraging Ecology

787   of a Territorial Hummingbird." *Behavioural Processes* 149: 27–34.

788   Heinrich, B., and P.H. Raven. 1972. "Energetics and Pollination Ecology." *Science* 176 (4035):

789   597–602.

790   HilleRisLambers, J., M.A. Harsch, A.K. Ettinger, K.R. Ford, and E.J. Theobald. 2013. "How

791   Will Biotic Interactions Influence Climate Change–Induced Range Shifts?" *Annals of the*

792   *New York Academy of Sciences* 1297 (1): 112–25.

793   Hixon, M.A., F.L. Carpenter, and D.C. Paton. 1983. "Territory Area, Flower Density, and Time

794   Budgeting in Hummingbirds: An Experimental and Theoretical Analysis." *The American*

795   *Naturalist* 122 (3): 366–91.

796   Jackson, E.R. 2017. "Does the Matrix Matter? Using Radio Frequency Identification (RFID)

797   Technology to Examine Hummingbird Movements in Fragmented Tropical Landscapes."

798   Master's thesis, Oregon State University.

799   https://ir.library.oregonstate.edu/concern/graduate_thesis_or_dissertations/cj82kd06n.

800   Jordán, F. 2009. "Keystone Species and Food Webs." *Philosophical Transactions of the Royal*

801   *Society B: Biological Sciences* 364 (1524): 1733–41.

802   Kaiser-Bunbury, C.N., J. Mougal, A.E. Whittington, T. Valentin, R. Gabriel, J.M. Olesen, and N.

803   Blüthgen. 2017. "Ecosystem Restoration Strengthens Pollination Network Resilience and

804   Function." *Nature* 542 (7640): 223–27.

805   Kaiser-Bunbury, C.N., S. Muff, J. Memmott, C.B. Müller, and A. Caflisch. 2010. "The

806   Robustness of Pollination Networks to the Loss of Species and Interactions: A

807   Quantitative Approach Incorporating Pollinator Behaviour." *Ecology Letters* 13 (4): 442–

808   52.

Kearns, C.A., and D.W. Inouye. 1993. *Techniques for Pollination Biologists*. Niwot: University

Press of Colorado.

Kearns, C.A., D.W. Inouye, and N.M. Waser. 1998. "Endangered Mutualisms: The Conservation

of Plant-Pollinator Interactions." *Annual Review of Ecology and Systematics* 29 (1): 83–

112.

Klein, A.-M., B.E. Vaissière, J.H. Cane, I. Steffan-Dewenter, S.A. Cunningham, C. Kremen, and

T. Tscharntke. 2007. "Importance of Pollinators in Changing Landscapes for World

Crops." *Proceedings of the Royal Society B: Biological Sciences* 274 (1608): 303–13.

Koh, L.P., R.R. Dunn, N.S. Sodhi, R.K. Colwell, H.C. Proctor, and V.S. Smith. 2004. "Species

Coextinctions and the Biodiversity Crisis." *Science* 305 (5690): 1632–34.

Kormann, U., C. Scherber, T. Tscharntke, N. Klein, M. Larbig, J.J. Valente, A.S. Hadley, and

M.G. Betts. 2016. "Corridors Restore Animal-Mediated Pollination in Fragmented

Tropical Forest Landscapes." *Proceedings of the Royal Society B: Biological Sciences*

283 (1823): 20152347.

Krüger, K., R. Prinzinger, and K.-L. Schuchmann. 1982. "Torpor and Metabolism in

Hummingbirds." *Comparative Biochemistry and Physiology Part A: Physiology* 73 (4):

679–89.

Kuban, J.F., and R.L. Neill. 1980. "Feeding Ecology of Hummingbirds in the Highlands of the

Chisos Mountains, Texas." *The Condor* 82 (2): 180–85.

Kuussaari, M., R. Bommarco, R.K. Heikkinen, A. Helm, J. Krauss, R. Lindborg, E. Öckinger, et

al. 2009. "Extinction Debt: A Challenge for Biodiversity Conservation." *Trends in*

Ecology & Evolution 24 (10): 564–71.

Lenth, R. 2020. *Emmeans: Estimated Marginal Means, Aka Least-Squares Means*.

https://CRAN.R-project.org/package=emmeans.

Lüdecke, D. 2018. "Ggeffects: Tidy Data Frames of Marginal Effects from Regression Models."

Journal of Open Source Software 3 (26): 772.

Lüdecke, D., M. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. "Performance: An

R Package for Assessment, Comparison and Testing of Statistical Models." *Journal of*

Open Source Software 6 (60): 3139.

Maglianesi, M.A., N. Blüthgen, K. Böhning-Gaese, and M. Schleuning. 2014. "Morphological

Traits Determine Specialization and Resource Use in Plant–Hummingbird Networks in

the Neotropics." *Ecology* 95 (12): 3325–34.

Maruyama, P.K., G.M. Oliveira, C. Ferreira, B. Dalsgaard, and P.E. Oliveira. 2013. "Pollination

Syndromes Ignored: Importance of Non-Ornithophilous Flowers to Neotropical Savanna

Hummingbirds." *Naturwissenschaften* 100 (11): 1061–68.

McDonald, T.L., W.P. Erickson, and L.L. McDonald. 2000. "Analysis of Count Data from

Before-after Control-Impact Studies." *Journal of Agricultural, Biological, and*

Environmental Statistics 5 (3): 262–79.

Memmott, J., N.M. Waser, and M.V. Price. 2004. "Tolerance of Pollination Networks to Species

Extinctions." *Proceedings of the Royal Society of London B: Biological Sciences* 271

(1557): 2605–11.

Mendenhall, C.D., and A.M. Wrona. 2018. "Improving Tree Cover Estimates for Fine-Scale

Landscape Ecology." *Landscape Ecology* 33 (10): 1691–96.

Mitchell, R.J., R.J. Flanagan, B.J. Brown, N.M. Waser, and J.D. Karron. 2009. "New Frontiers in

Competition for Pollination." *Annals of Botany* 103 (9): 1403–13.

Moeller, D.A. 2004. "Facilitative Interactions among Plants via Shared Pollinators." *Ecology* 85

(12): 3289–3301.

Moilanen, A., and M. Nieminen. 2002. "Simple Connectivity Measures in Spatial Ecology."

Ecology 83 (4): 1131–45.

Montero-Castaño, A., and M. Vilà. 2012. "Impact of Landscape Alteration and Invasions on

Pollinators: A Meta-Analysis." *Journal of Ecology* 100 (4): 884–93.

Montgomerie, R.D., and C.A. Redsell. 1980. "A Nesting Hummingbird Feeding Solely on

Arthropods." *The Condor* 82 (4): 463.

Morales, C.L., and A. Traveset. 2008. "Interspecific Pollen Transfer: Magnitude, Prevalence and

Consequences for Plant Fitness." *Critical Reviews in Plant Sciences* 27 (4): 221–38.

Moran, A.J., S.W.J. Prosser, and J.A. Moran. 2019. "DNA Metabarcoding Allows Non-Invasive

Identification of Arthropod Prey Provisioned to Nestling Rufous Hummingbirds

(Selasphorus Rufus)." *PeerJ* 7: e6596.

Ohashi, K., and J.D. Thomson. 2009. "Trapline Foraging by Pollinators: Its Ontogeny,

Economics and Possible Consequences for Plants." *Annals of Botany* 103 (9): 1365–78.

Ollerton, J., R. Winfree, and S. Tarrant. 2011. "How Many Flowering Plants Are Pollinated by

Animals?" *Oikos* 120 (3): 321–26.

Pauw, A., and J.A. Hawkins. 2011. "Reconstruction of Historical Pollination Rates Reveals

Linked Declines of Pollinators and Plants." *Oikos* 120 (3): 344–49.

Phillips, R.D., R. Peakall, B.A. Retter, K. Montgomery, M.H.M. Menz, B.J. Davis, C. Hayes,

G.R. Brown, N.D. Swarts, and K.W. Dixon. 2015. "Pollinator Rarity as a Threat to a

Plant with a Specialized Pollination System." *Botanical Journal of the Linnean Society*

179 (3): 511–25.

Potts, S.G., J.C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W.E. Kunin. 2010.

"Global Pollinator Declines: Trends, Impacts and Drivers." *Trends in Ecology &*

Evolution 25 (6): 345–53.

Powers, D.R., A.R. Brown, and J.A. Van Hook. 2003. "Influence of Normal Daytime Fat

Deposition on Laboratory Measurements of Torpor Use in Territorial versus

Nonterritorial Hummingbirds." *Physiological and Biochemical Zoology* 76 (3): 389–97.

R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Ramos-Jiliberto, R., F.S. Valdovinos, P. Moisset de Espanés, and J.D. Flores. 2012.

"Topological Plasticity Increases Robustness of Mutualistic Networks." *Journal of*

Animal Ecology 81 (4): 896–904.

Rathcke, B. 1983. "Competition and Facilitation among Plants for Pollination." In *Pollination*

Biology, edited by L.A. Real, 305–29. New York: Academic Press.

Regan, E.C., L. Santini, L. Ingwall-King, M. Hoffmann, C. Rondinini, A. Symes, J. Taylor, and

S.H.M. Butchart. 2015. "Global Trends in the Status of Bird and Mammal Pollinators."

Conservation Letters 8 (6): 397–403.

Renjifo, L.M. 2001. "Effect of Natural and Anthropogenic Landscape Matrices on the

Abundance of Subandean Bird Species." *Ecological Applications* 11 (1): 14–31.

Rodríguez-Flores, C.I., J.F. Ornelas, S. Wethington, and M.C. Arizmendi. 2019. "Are

Hummingbirds Generalists or Specialists? Using Network Analysis to Explore the

Mechanisms Influencing Their Interaction with Nectar Resources." *PLOS ONE* 14 (2):

e0211855.

Russell, S.M., and R.O. Russell. 2001. *The North American Banders' Manual for Banding*

Hummingbirds. North American Banding Council.

Schemske, D.W. 1981. "Floral Convergence and Pollinator Sharing in Two Bee-Pollinated

Tropical Herbs." *Ecology* 62 (4): 946–54.

Scheper, J., M. Reemer, R. van Kats, W.A. Ozinga, G.T.J. van der Linden, J.H.J. Schaminée, H.

Siepel, and D. Kleijn. 2014. "Museum Specimens Reveal Loss of Pollen Host Plants as

Key Factor Driving Wild Bee Decline in The Netherlands." *Proceedings of the National*

Academy of Sciences 111 (49): 17552–57.

Schuchmann, K.-L., and R. Prinzinger. 1988. "Energy Metabolism, Nocturnal Torpor, and

Respiration Frequency in a Green Hermit (Phaethornis Guy)." *Journal Für Ornithologie*

129 (4): 469–72.

Senapathi, D., L.G. Carvalheiro, J.C. Biesmeijer, C.-A. Dodson, R.L. Evans, M. McKerchar,

R.D. Morton, et al. 2015. "The Impact of over 80 Years of Land Cover Changes on Bee

and Wasp Pollinator Communities in England." *Proceedings of the Royal Society B:*

Biological Sciences 282 (1806): 20150294.

Shankar, A., R.J. Schroeder, S.M. Wethington, C.H. Graham, and D.R. Powers. 2020.

"Hummingbird Torpor in Context: Duration, More than Temperature, Is the Key to

Nighttime Energy Savings." *Journal of Avian Biology* 51 (5): e02305.

Spence, A.R., and M.W. Tingley. 2021. "Body Size and Environment Influence Both

Intraspecific and Interspecific Variation in Daily Torpor Use across Hummingbirds."

Functional Ecology 35 (4): 870–83.

Steffan-Dewenter, I., and T. Tscharntke. 1999. "Effects of Habitat Isolation on Pollinator

Communities and Seed Set." *Oecologia* 121 (3): 432–40.

Stein, K., M. Templin, I. Hensen, M. Fischer, D. Matthies, and M. Schleuning. 2013. "Negative

Effects of Conspecific Floral Density on Fruit Set of Two Neotropical Understory

Plants." *Biotropica* 45 (3): 325–32.

Stiles, F.G. 1975. "Ecology, Flowering Phenology, and Hummingbird Pollination of Some Costa

Rican Heliconia Species." *Ecology* 56 (2): 285–301.

———. 1979. "Notes on the Natural History of Heliconia (Musaceae) in Costa Rica. Notas

Sobre La Historia Natural de Heliconia (Musaceae) En Costa Rica." *Brenesia* 15: 151–

80.

———. 1985. "Seasonal Patterns and Coevolution in the Hummingbird-Flower Community of a

Costa Rican Subtropical Forest." *Ornithological Monographs* 36: 757–87.

———. 1995. "Intraspecific and Interspecific Variation in Molt Patterns of Some Tropical

Hummingbirds." *The Auk* 112 (1): 118–32.

Stiles, F.G., and L.L. Wolf. 1970. "Hummingbird Territoriality at a Tropical Flowering Tree."

The Auk 87 (3): 467–91.

Stratton, D.A. 1989. "Longevity of Individual Flowers in a Costa Rican Cloud Forest: Ecological

Correlates and Phylogenetic Constraints." *Biotropica* 21 (4): 308–18.

Suarez, R.K. 1992. "Hummingbird Flight: Sustaining the Highest Mass-Specific Metabolic Rates

among Vertebrates." *Experientia* 48 (6): 565–70.

Taki, H., P.G. Kevan, and J.S. Ascher. 2007. "Landscape Effects of Forest Loss in a Pollination

System." *Landscape Ecology* 22 (10): 1575–87.

Taylor, J., and S.A. White. 2007. "Observations of Hummingbird Feeding Behavior at Flowers

of Heliconia Beckneri and H. Tortuosa in Southern Costa Rica." *Ornitologia Neotropical*

18: 133–38.

Taylor, P.D., L. Fahrig, K. Henein, and G. Merriam. 1993. "Connectivity Is a Vital Element of

946   Landscape Structure." *Oikos*, 571–73.

947   Temeles, E.J., A.B. Muir, E.B. Slutsky, and M.N. Vitousek. 2004. "Effect of Food Reductions on

948   Territorial Behavior of Purple-Throated Caribs." *The Condor* 106 (3): 691–95.

949   Thomas-Granger, K. 2003. "Light Sum, Nitrogen Level and Growth Retardant Effects on Flower

950   and Nectar Production in Stachytarpheta Frantzii." Master's thesis.

951   Timóteo, S., J.A. Ramos, I.P. Vaughan, and J. Memmott. 2016. "High Resilience of Seed

952   Dispersal Webs Highlighted by the Experimental Removal of the Dominant Disperser."

953   *Current Biology* 26 (7): 910–15.

954   Tooze, Z.J., and C.L. Gass. 1985. "Responses of Rufous Hummingbirds to Midday Fasts."

955   *Canadian Journal of Zoology* 63 (10): 2249–53.

956   Traveset, A., C. Tur, and V.M. Eguíluz. 2017. "Plant Survival and Keystone Pollinator Species

957   in Stochastic Coextinction Models: Role of Intrinsic Dependence on Animal-Pollination."

958   *Scientific Reports* 7: 6915.

959   Underwood, A.J. 1994. "On Beyond BACI: Sampling Designs That Might Reliably Detect

960   Environmental Disturbances." *Ecological Applications* 4 (1): 3–15.

961   Valdovinos, F.S., P. Moisset de Espanés, J.D. Flores, and R. Ramos-Jiliberto. 2013. "Adaptive

962   Foraging Allows the Maintenance of Biodiversity of Pollination Networks." *Oikos* 122

963   (6): 907–17.

964   Vellend, M., K. Verheyen, H. Jacquemyn, A. Kolb, H.V. Calster, G. Peterken, and M. Hermy.

965   2006. "Extinction Debt of Forest Plants Persists for More Than a Century Following

966   Habitat Fragmentation." *Ecology* 87 (3): 542–48.

Vizentin-Bugoni, J., V.J. Debastiani, V.A.G. Bastazini, P.K. Maruyama, and J.H. Sperry. 2020.

"Including Rewiring in the Estimation of the Robustness of Mutualistic Networks."

Methods in Ecology and Evolution 11 (1): 106–16.

Vizentin-Bugoni, J., P.K. Maruyama, and M. Sazima. 2014. "Processes Entangling Interactions

in Communities: Forbidden Links Are More Important than Abundance in a

Hummingbird–Plant Network." *Proceedings of the Royal Society B: Biological Sciences*

281 (1780): 20132397.

Volpe, N.L., A.S. Hadley, W.D. Robinson, and M.G. Betts. 2014. "Functional Connectivity

Experiments Reflect Routine Movement Behavior of a Tropical Hummingbird Species."

Ecological Applications 24 (8): 2122–31.

Volpe, N.L., W.D. Robinson, S.J.K. Frey, A.S. Hadley, and M.G. Betts. 2016. "Tropical Forest

Fragmentation Limits Movements, but Not Occurrence of a Generalist Pollinator

Species." *PLOS ONE* 11 (12): e0167513.

Weiner, C.N., M. Werner, K.E. Linsenmair, and N. Blüthgen. 2014. "Land-Use Impacts on

Plant–Pollinator Networks: Interaction Strength and Specialization Predict Pollinator

Declines." *Ecology* 95 (2): 466–74.

Weinstein, B.G. 2015. "MotionMeerkat: Integrating Motion Video Detection and Ecological

Monitoring." *Methods in Ecology and Evolution* 6 (3): 357–62.

Weinstein, B.G., and C.H. Graham. 2017. "Persistent Bill and Corolla Matching despite Shifting

Temporal Resources in Tropical Hummingbird-Plant Interactions." *Ecology Letters* 20

(3): 326–35.

Winfree, R., I. Bartomeus, and D.P. Cariveau. 2011. "Native Pollinators in Anthropogenic

Habitats." *Annual Review of Ecology, Evolution, and Systematics* 42 (1): 1–22.

Wolf, L.L., F.R. Hainsworth, and F.G. Stiles. 1972. "Energetics of Foraging: Rate and Efficiency

of Nectar Extraction by Hummingbirds." *Science* 176 (4041): 1351–52.

Wolf, L.L., and F.G. Stiles. 1989. "Adaptations for the 'fail-Safe' Pollination of Specialized

Ornithophilous Flowers." *The American Midland Naturalist* 121 (1): 1–10.

Young, A.M. 1971. "Foraging for Insects by a Tropical Hummingbird." *The Condor* 73 (1): 36–

45.

Zahawi, R.A., G. Duran, and U. Kormann. 2015. "Sixty-Seven Years of Land-Use Change in

Southern Costa Rica." *PLOS ONE* 10 (11): e0143554.

 TABLES

Table 1. Hypotheses testing whether declines of a locally common, tropical flowering plant (*Heliconia tortuosa*) affect hummingbird

space use, hummingbird floral visitation rates, and plant pollination success. Predicted effects are indicated in the column below each

hypothesis. To address Question 1, we (i) experimentally removed *H. tortuosa* using a BACI (Before-After-Control-Impact)

experimental design and (ii) studied a natural gradient in *H. tortuosa* density. Under the experimental framework, predicted effects can

be conceptualized as pre-post changes in treatment replicates, compared to to pre-post changes in control replicates. To address

Question 2, we examined changes in hummingbird and plant responses across a natural gradient in *H. tortuosa* density only.

			Question 1: How do hummingbirds respond to declines in *Heliconia,* and are there cascading effects on the remaining plants?		Question 2: What factors mediate hummingbird responses to *Heliconia* declines?		
					Connectivity hypothesis: Rescue (H-2a)	Connectivity hypothesis: Abandonment (H-2b)	Resource hypothesis (H-3)
	Response variable	Methodological approach	Robust network hypothesis (H-1a)	Parallel extinction hypothesis (H-1b)	*In highly connected forest fragments...*	*In highly connected forest fragments...*	*In forest fragments with high resource availability...*
Space use	Abundance	Mist net captures	No change	Decrease			
Space use	Time spent in focal area	Radio telemetry	No change	Decrease			
Floral visitation rates	Visitation to focal *Heliconia* plants	Remote camera observations	No change or increase	Decrease	Smaller decrease	Larger decrease	Smaller decrease
Floral visitation rates	Visitation to non-*Heliconia* plants	Remote camera observations	Increase (rewiring)	Decrease			
Pollination services	Pollination success	Pollen tubes	No change or increase	Decrease			

 FIGURES

Figure 1. This study was conducted within forest fragments in the landscape surrounding the Las Cruces Biological Station (A) in southern Costa Rica (B). Map in (A) illustrates forest (grey) and non-forest (white), with non-forest primarily comprising pasture. The dark grey forest layer was hand-digitized for this study; the light grey forest layer, available within the geographic boundary of Coto Brus Canton, was obtained from Mendenhall & Wrona (2018). We utilized fourteen focal landscapes of 1000 m radius ('sites', dashed circles), numbered here by increasing connectivity to other areas of forest. Most sites were used more than once across three years of study (2016-2018). Within a year, each site received either the *Heliconia* removal treatment or was assigned to be a control. Treatments were reversed in alternate years for a total sample size of 16 control replicates and 16 treatment replicates. An example site is depicted in (C). Red outlines in (C) and (D) represent a focal area in which we removed *Heliconia* (in treatment replicates) and surveyed floral resources (in all replicates). Within each focal area, we captured hummingbirds using mist nets (triangles in D) and surveyed plant-hummingbird interactions using trail cameras (icons in D). To quantify functional connectivity, we summed the total area of connected forest within the focal landscape, with connectedness based on 50-m gap crossing ability and pixels down-weighted based on distance to focal area (E; see Appendix S2 for details).

1026

Figure 2. (A) During the *Heliconia* removal experiment, we measured four main response

variables: captured hummingbird abundance, proportion of time radio-tagged hummingbirds

spent in the focal area, floral visitation rates, and plant pollination success. Each response

variable was measured during both experimental periods (pre and post); the sampling schedule

for each is denoted with an 'X'. Each experimental replicate used two sites (control and

treatment) sampled using this schedule, offset by one day. Starting around noon on the fourth

day, we covered *Heliconia* inflorescences in the treatment site. (B) *Heliconia* inflorescences

were covered with dark-colored bags that prevented hummingbirds from accessing any nectar

present. (C) Our manipulation removed, on average, 42.5% (± 30.9% SD) of the calories

available for the entire hummingbird community and 65.5% (± 31.3% SD) of the available

calories for *Heliconia* specialists (green hermits and violet sabrewings). Mean values are denoted

by red diamonds, and the darker black line represents the median. Lower and upper lines within

boxplots represent the first and third quartiles. Whiskers extend from the first and third quartiles

to the lowest and highest values within 1.5 times the interquartile range. Individual data points

1041   appear next to each boxplot. *Response variable icons in (A) were created using images from the*

1042   *Noun Project (see 'Acknowledgments'). Photos in (B) were provided by Marion Donald.*

1043

Figure 3. The overall plant-hummingbird visitation network for forest fragments in the

landscape surrounding the Las Cruces Biological Station (A), created from 15,196 hours of

camera observations across 37 plant species, 14 study sites, and three field seasons (2016-2018).

(B) The network in A, but subset to focus on visitation by green hermits (*P. guy*) and violet

sabrewings (*C. hemileucurus*), the two most frequent visitors to *Heliconia*. The height of each

rectangle represents the overall observation frequency for that species (larger rectangle = more

observations), while width of the connecting links reflects relative visitation rate to different

partners (wider link = higher visitation rate). Five plant species (*Calathea lutea*, *Ctenanthe*

dasycarpa, *Erythrina poeppigiana*, *Maripa nicaraguensis*, *Mucuna globulifera*) received no

hummingbird visitation during the observation period and are thus not represented in these

networks.

1055

Figure 4. Hummingbird space use before and after experimental local extinction of an

understory plant, *Heliconia tortuosa*, within forest fragments in southern Costa Rica. Despite

removing at least 43% (± 31 SD) of calories available to hummingbirds within focal areas of 1.7

ha (± 1.9 SD), we did not observe an effect of *Heliconia* removal on the number of

hummingbirds captured (A-B) or the proportion of time that radio-tagged hummingbirds spent in

the focal areas (C-D). Results are shown for all species analyzed (left) and green hermits and

violet sabrewings (right). Estimated marginal means from GLMMs are presented alongside 95%

confidence intervals. *P*-values refer to the interaction testing the effect of *Heliconia* removal,

relative to control replicates; conceptually, a treatment effect is indicated by non-parallel lines.

The estimated number of captured hummingbirds is calculated for 45.8 net-hours, the mean

capture effort across all replicates.

Figure 5. Hummingbird visitation rates to focal *Heliconia* plants and *Heliconia* pollination

success before and after experimental *Heliconia* removal (A-C) and across a natural gradient in

Heliconia density (D-F). For hummingbird visitation rates, results are shown for the entire

hummingbird community (A, D) and green hermits and violet sabrewings separately (B, E).

Estimated marginal means from GLMMs are presented alongside 95% confidence intervals. (A-

C) *P*-values refer to the interaction testing the effect of *Heliconia* removal, relative to control

replicates; conceptually, a treatment effect is indicated by non-parallel lines. (D-F) Values of

Heliconia availability (*x*-axis) are standard deviations (1 SD = 1.35), with zero representing the

mean log-transformed caloric density (mean = 8). Note that the *x*-axis is reversed, such that

declining visitation is associated with declining *Heliconia* availability. *P*-values refer to the

coefficient for *Heliconia* availability.

1079

Figure 6. Hummingbird visitation rates to non-*Heliconia* plant species before and after

experimental *Heliconia* removal (A-B) and across a natural gradient in *Heliconia* density (C-D).

Results are shown for the entire hummingbird community (A, C) and green hermits and violet

sabrewings separately (B, D). Estimated marginal means from GLMMs are presented alongside

95% confidence intervals. (A-B) *P*-values refer to the interaction testing the effect of *Heliconia*

removal, relative to control replicates; conceptually, a treatment effect is indicated by non-

parallel lines. (C-D) Values of *Heliconia* availability (*x*-axis) are standard deviations (1 SD =

1.35), with zero representing the mean log-transformed caloric density (mean = 8). Note that the

x-axis is reversed, such that declining visitation is associated with declining *Heliconia*

availability. *P*-values refer to the coefficient for *Heliconia* availability.