Are tropical hummingbird pollination networks resistant to experimental extirpation of a

common flowering plant?

Kara G. Leimberger[1, *], Adam S. Hadley[1,2], Sarah J.K. Frey[1,3], and Matthew G. Betts[1]

[1]Forest Biodiversity Research Network, Department of Forest Ecosystems and Society, Oregon

State University, Corvallis, Oregon, USA

[2]Biodiversity Section, Department of Natural Resources and Energy Development, Fredericton,

New Brunswick, Canada

[3]Department of Animal and Rangeland Sciences, Oregon State University, Corvallis, Oregon,

USA

[*]Author for correspondence (E-mail: kara.leimberger@oregonstate.edu).

ABSTRACT

Plants and pollinators are experiencing parallel declines worldwide, despite theory predicting that pollination networks should withstand disturbance due to redundancy of pollinators, rarity of interactions between specialists, and flexible pollinator foraging behavior (rewiring). Experiments evaluating the stability of plant-pollinator networks may help resolve this incongruity but remain uncommon. In this study, we simulated the extirpation of a hummingbird-pollinated understory plant, *Heliconia tortuosa,* from tropical forest fragments using a replicated Before-After-Control-Impact experimental design while quantifying hummingbird space use (383 hummingbird captures and 72 radio-tagged individuals), floral visitation rates (6474 visits from >19,000 observation hours), and plant pollination success (529 flowers). To complement this experimental approach, we also examined these responses across a natural gradient in *H. tortuosa* density. We expected that declines of *H. tortuosa* would either result in (i) network collapse, in which hummingbirds vacate fragments and compromise the reproductive success of other flowering plants, or (ii) increased hummingbird reliance on alternative resources, leading to sustained fragment use. We also hypothesized that landscape and local context (i.e., connectivity to additional forest area and alternative resource abundance) could mediate hummingbird responses to *H. tortuosa* declines; for example, connectivity could facilitate hummingbird visitation to areas of reduced food availability. In our removal experiment, hummingbird persistence and plant pollination success were remarkably resistant to loss of *H. tortuosa*, a locally common plant species representing >40% of the available nectar resources on average. However, naturally low *H. tortuosa* densities were associated with reduced floral visitation rates and decreased pollination. Although landscape context (connectivity) led to higher hummingbird abundance, we found little evidence that connectivity or resource availability mediated hummingbird responses to declines in *H. tortuosa* availability. The exact

mechanisms enabling short-term hummingbird persistence after resource removal remain

unclear, as we did not discover evidence of rewiring. Physiological adaptations (e.g., torpor and

insectivory), may have contributed to hummingbird resilience, perhaps alongside high movement

abilities that facilitated visitation to spatially dispersed floral resources. With the important

caution that short-term experiments may not realistically emulate natural extinction processes,

our study provides partial support for theoretical predictions that pollination networks may be

relatively robust to plant species loss.

Key words: mutualistic networks, ecosystem function, foraging ecology, functional

connectivity, fragmentation, Trochilidae

INTRODUCTION

Species loss can have cascading consequences for the broader community, including

coextinctions of dependent species and declines in ecosystem function (Koh et al. 2004, Colwell

et al. 2012). Linked extinctions may be especially likely for mutualistic interactions, such as

those between plants and pollinators (Dunn et al. 2009, Aslan et al. 2013). For instance, plants

that are not adequately pollinated produce fewer seeds, accruing extinction debt that becomes

realized when the existing generation cannot be replaced by new recruits (Ashman et al. 2004,

Vellend et al. 2006, Kuussaari et al. 2009). This repeated reproductive failure will eventually

cause demographic collapse of plant populations (Anderson et al. 2011, Phillips et al. 2015),

which could then hasten the decline of animals that consume pollen or nectar (Biesmeijer et al.

2006, Pauw and Hawkins 2011, Weiner et al. 2014, Scheper et al. 2014). Indeed, plants and

pollinators are experiencing declines worldwide (Biesmeijer et al. 2006, Potts et al. 2010, Regan

et al. 2015), which not only imperils the essential ecosystem services on which humans depend

but also threatens the maintenance of global biodiversity (Allen-Wardell et al. 1998, Klein et al.

2007, Ollerton et al. 2011).

Although the underlying reasons for plant and pollinator declines are not fully

understood, one prominent hypothesis invokes the landscape-level processes of habitat loss and

fragmentation (Kearns et al. 1998, Aguilar et al. 2006, Potts et al. 2010, Hadley and Betts 2012).

Habitat destruction often creates small habitat patches unable to sustain large populations, which

can increase rates of local extinction (Hanski 1998), reduce functional connectivity (sensu P. D.

Taylor et al. 1993), and impede pollinator movement (Hadley and Betts 2012). Likely due to

these combined mechanisms, land-use change and habitat isolation have been repeatedly linked

to declines in pollinator diversity (Winfree et al. 2011, Montero-Castaño and Vilà 2012,

Senapathi et al. 2015) and plant reproduction (Steffan-Dewenter and Tscharntke 1999, Taki et al.

2007, Farwig et al. 2009, Garibaldi et al. 2011).

However, parallel declines of pollinators and plants are at odds with theoretical

predictions suggesting that species loss should generally not result in coextinction of interaction

partners (Memmott et al. 2004, Bascompte et al. 2006). Tolerance to subsequent extinctions (i.e.,

robustness) primarily arises from the expectation that interaction networks exhibit strongly

nested topologies, such that specialists interact with subsets of partners linked to generalists

(Bascompte et al. 2003). Under this interaction pattern, connections between specialists are

uncommon, so the functional role of a specialist can be absorbed by its more abundant generalist

partners (Bascompte et al. 2003, Bascompte and Jordano 2007). In addition to the redundancy

afforded by a nested structure, models suggest that flexibility in pollinator foraging behavior –

i.e., rewiring to use alternative resources – may buffer networks against disturbance (Kaiser-

Bunbury et al. 2010, Ramos-Jiliberto et al. 2012, Valdovinos et al. 2013, Timóteo et al. 2016).

Despite these overall predictions and mechanisms of robustness, pollination networks

may be vulnerable to habitat loss and fragmentation under certain conditions. Although

systematic loss of generalist pollinators may be less realistic than extinctions of rare, specialist

pollinators (Aizen et al. 2012), simulation studies have revealed that removal of highly

connected, generalist species thwarts a network's capacity to buffer perturbations, leading to

rapid partner extinction (Memmott et al. 2004, Kaiser-Bunbury et al. 2010, Traveset et al. 2017).

Thus, if external stressors such as habitat fragmentation threaten highly connected species

('topological keystones' sensu Jordán 2009), an extinction cascade could occur – particularly

without rewiring by pollinators. However, uncertainty about the robustness of real-world plant-

pollinator networks still exists because relatively few studies have quantified pollinator

responses to deletions of other species (Appendix S1: Table S1).

In this study, we tested whether declines of a hummingbird-pollinated plant *(Heliconia*

tortuosa) could have cascading consequences for communities of hummingbirds and flowering

plants within a fragmented tropical forest landscape – or whether rewiring might forestall

network collapse. In Neotropical forests, the genus *Heliconia* (Heliconiaceae) comprises

understory herbs that have been suggested to function as "keystone mutualists" (Gilbert 1980).

Large, brightly colored *Heliconia* inflorescences not only provide nectar to large-bodied hermit

hummingbirds (Trochilidae: Phaethornithinae) throughout the year, but also support a variety of

other hummingbird species during periods of resource scarcity (Stiles 1975, 1985). Because most

hummingbird species visit multiple plant species (Rodríguez-Flores et al. 2019), loss of *H.*

tortuosa and its associated hummingbird partners could jeopardize the reproduction of co-

occurring plants, thus generating an extinction cascade (Gilbert 1980). Although *H. tortuosa* is

locally common in the mid-elevation, premontane tropical forests of Costa Rica, Hadley *et al.*

(2014) found that *H. tortuosa* in smaller forest fragments produced fewer seeds, possibly due to

short-distance pollen transfer associated with pollinator movement limitation. Although

numerous hummingbird species visit *H. tortuosa* opportunistically, its primary pollinators are the

green hermit (*Phaethornis guy*) and violet sabrewing (*Campylopterus hemileucurus*) (Stiles

1979, Wolf and Stiles 1989, Taylor and White 2007, Betts et al. 2015). These relatively large,

forest-associated hummingbirds are high-reward trapliners (sensu Feinsinger and Colwell 1978)

that visit spatially dispersed, nectar-rich floral resources in a routine sequence (Borgella et al.

2001). However, their movements become restricted in fragmented habitats (Hadley and Betts

2009, Volpe et al. 2016, Kormann et al. 2016).

To assess the potential consequences of *H. tortuosa* declines, we measured hummingbird

space use, floral visitation rates, and plant pollination success before and after temporary

experimental removal of *H. tortuosa* within forest fragments. Additionally, we assessed how

these responses varied across a natural gradient in *Heliconia* density. We developed two

alternative hypotheses about how hummingbirds would respond to declines of *H. tortuosa*

(hereafter *Heliconia*) and potentially affect the pollination of co-occuring plant species (Table 1).

Under the *robust network hypothesis* (**H1a**), hummingbirds do not emigrate from forest

fragments following *Heliconia* decline, leading to uninterrupted visitation to remaining plants.

Under this scenario, we hypothesized that hummingbirds compensate for *Heliconia* extinction by

increasing their use of non-*Heliconia* floral resources (rewiring). Alternatively, hummingbirds

may be highly dependent on *H. tortuosa* and inflexible in their foraging choices; therefore, local

Heliconia declines should reduce nectar availability such that hummingbirds cannot afford the

energetic costs to remain in (or travel to) focal forest fragments, leading to decreases in fragment

use and pollination success (**H1b***: parallel extinction hypothesis*). Under both hypotheses, we

expected hummingbirds to change their behavior due to their exceptionally rapid metabolisms,

high dependency on nectar, and limited fat stores (Suarez 1992, Powers et al. 2003).

Additionally, we hypothesized that landscape and local context could mediate the influence of

Heliconia declines on hummingbird behavior (Table 1). Connectivity to additional forest area

(*connectivity hypothesis*, **H2**) may sustain hummingbird fragment use by facilitating travel to

fragments (*rescue hypothesis*, **H2a**) or promote steeper declines in fragment use by facilitating

hummingbird departure (*abandonment hypothesis*, **H2b**). Further, greater local availability of

non-*Heliconia* floral resources may promote hummingbird persistence through increased

opportunities for rewiring (*resource hypothesis,* **H3**)*.* Lastly, we expected that changes in

Heliconia density would affect green hermits and violet sabrewings more strongly than other

hummingbird species (**H4**). Green hermits and violet sabrewings (hereafter '*Heliconia*

specialists') inhabit the forest understory, have bills that are morphologically specialized to long,

curved *H. tortuosa* flowers, and are thought to visit *H. tortuosa* more frequently than other

hummingbirds (Borgella et al. 2001, Betts et al. 2015).

To our knowledge, this study not only constitutes one of most highly replicated, large-

scale removal experiments conducted within the context of plant-pollinator interaction networks

(Appendix S1: Table S1), but also represents the first test of how landscape and local context

influence hummingbird responses to declines in a common floral resource. Importantly,

understanding mechanisms of pollinator persistence in fragmented landscapes has implications

for conservation planning, since fragmentation effects may be ameliorated by habitat restoration,

preservation of riparian corridors, and/or implementation of living fences (Renjifo 2001,

Kormann et al. 2016).

METHODS

Study area

This study was conducted in southern Costa Rica within a ∼122 km^2 area surrounding the Las

Cruces Biological Station (Coto Brus Canton, Puntarenas Province, 8°47´7´´ N, 82°57´32´´ W,

Fig. 1: A-B). This study area comprises cattle pastures, coffee plantations, and fragments of

tropical premontane wet forest created during a period of extensive land clearing concentrated

between 1960 and 1980 (Zahawi et al. 2015). Remnant forest fragments (∼30% of the study area)

span a wide range of sizes (<1 to >1000 ha), with otherwise isolated fragments often connected

by narrow riparian corridors and living fences (Hadley and Betts 2009, Zahawi et al. 2015).

Mean annual temperature at the Las Cruces Biological Station is ∼21 ºC, mean annual rainfall is

3500-4000 mm, and a dry season occurs from December to March (Borgella et al. 2001, Zahawi

et al. 2015).

Study sites

We selected 14 focal fragments embedded within focal landscapes of 1000 m radius

('sites', Fig. 1: A, C). We chose the spatial extent of focal landscapes *a priori* based on the

maximum estimated daily movement distance for large, forest-associated hummingbird species

in this study system (Volpe et al. 2014).We identified potential sites using a stratified design

with two categories: 'connected to forest' or 'unconnected to forest', based on visual

examination of satellite imagery and previous work in this study area (Hadley et al. 2014, 2018,

Kormann et al. 2016). This design was only for site establishment, as we later quantified

connectivity as a continuous variable (see *'Quantifying connectivity'* below). Sites were then

included based on presence of flowering *Heliconia***,** landowner permission, vehicle accessibility,

and navigable terrain. To ensure site accessibility, we conducted our study during the dry season

and beginning of the rainy season (February – May), because localized flooding can occur during

periods of heavy rainfall. Elevation of sites ranged from ~900-1500 m (Appendix S2: Table S1).

Study design

Using the 14 focal-patch landscapes, we repeated the *Heliconia* removal experiment 16

times across three years (2016-2018). Each iteration of the experiment followed a BACI (Before-

After-Control-Impact) design and used two sites paired based on elevation and connectivity

(Appendix S2: Tables S2-S3). One of the paired sites received the treatment of *Heliconia*

removal, and we designated the other site as a control (16 control replicates, 16 treatment

replicates). To prevent hummingbirds from the treatment site relocating to the control site, paired

sites were located at least 1000 m apart (mean $\pm$ SD: 2266 $\pm$ 1511 m, median: 3086 m). Most

sites were used multiple times across the three-year study period, but only once per year

(Appendix S2: Tables S2-S4). Sites used multiple times received the opposite treatment of the

previous year, except for 3 of 16 instances (e.g., when covering *Heliconia* in that site again

would have posed substantial logistical challenges). Initial treatment assignments were random.

This design reduced the likelihood that any treatment differences were due to factors associated

with individual sites (Appendix S2: Fig. S1).

Each experimental iteration lasted nine days and began with a period of pre-treatment

data collection in each site (Fig. 2A). Starting around noon on the fourth day, we removed

Heliconia from the treatment site and designated the following 36 hours as a behavioral

adjustment period during which no data were collected. Although this adjustment period was

necessarily relatively short due to the limited battery lifespan of hummingbird radio-transmitters

(see *'Radio telemetry'*), we expected hummingbirds to respond rapidly due to their exceptionally

high metabolisms and limited fat stores (Suarez 1992, Powers et al. 2003). To ensure that

Heliconia inflorescences were not overlooked during removal, field technicians positioned

themselves ~2-3 meters apart and slowly walked through the forest while covering all *Heliconia*

inflorescences (mature and immature) with dark-colored fabric bags (Fig. 2B). These bags

prevented hummingbird access, obscured flowers from view, and allowed us to avoid long-term

changes to floral resources associated with destructive sampling methods (e.g., cutting). The

location of each *Heliconia* plant was recorded using a handheld GPS unit, and we retrieved the

covering bags at the end of the experimental period. In each site, we also recorded the location

and identity of all non-*Heliconia* floral resources available to hummingbirds (see *'Quantifying*

resource availability'). The areas where we removed *Heliconia* and/or surveyed for floral

resources are referred to as 'focal areas' (Fig. 1D), which we delineated using 100% minimum

convex polygons. Focal areas also encompass locations where we captured hummingbirds,

installed trail cameras, and set up plant arrays (Fig. 1D, see *'Response variables'*).

Response variables

Hummingbird captures

Within each focal area, we captured hummingbirds at the beginning and end of the

experimental period (Fig. 2A) by deploying 5 - 20 mist nets (12-m and 6-m in length) near

flowering plants visited by hummingbirds (e.g., the genera *Heliconia*, *Centropogon, Palicourea,*

Musa). Nets were opened 30 minutes after sunrise and remained open for 2 - 5 hours per session.

Capture effort (net-hours) and net locations were kept constant for both capture sessions, except

in one instance (Appendix S2: Fig. S2). Following the North American Banding Council's

hummingbird protocol (Russell and Russell 2001), we tagged hummingbirds with an aluminum

tarsus band, measured body mass to the nearest 0.01 g using an electronic balance, and measured

wing length to the nearest 0.5 mm using a wing rule. Tarsus bands allowed detection of

recaptured hummingbirds within and between capture sessions; individuals recaptured during the

same capture session were only counted once. In 2017-2018, we also marked hummingbirds'

heads with unique color combinations of nail polish following Hadley *et al.* (2014); this

additional mark allowed us to identify individual hummingbirds seen on camera (see *'Camera*

observations'). All birds were handled in accordance with Oregon State University Animal Care

and Use Protocols (ACUP #4655, #5015).

Radio telemetry

We attached miniaturized radio transmitters (0.15-0.25g, Blackburn Transmitters,

Nacogdoches, TX) to 72 hummingbirds according to the methods of Hadley & Betts (2009). We

then followed hummingbirds on foot using portable receivers (Wildlife Materials Inc TRX

1000S) and handheld Yagi antennae, recorded locations continuously when the bird was found

within receiver range (~250 m in the absence of topographical barriers). In total, we completed

447 hours of telemetry observation, averaging 3.1 ± 0.8 hours per session (typically ~8:00-11:00)

and 5.3 ± 0.9 sessions per site (mean $\pm$ SD). From the recorded locations, we estimated the

proportion of observation time that radio-tagged birds spent in the focal area (Appendix S2: Fig.

S3-S4). Thirty-six hummingbirds spent time in the focal area during the 'pre' period and were

included in analyses of the *Heliconia* removal experiment; for a comprehensive summary of

radio-tracking outcomes, see Appendix S2 (Table S5).

Floral visitation rates

To understand hummingbird resource use within each site and to monitor for the presence

of marked individuals, we positioned trail cameras (PlotWatcher Pro, Day 6 Outdoors) near

inflorescences of plant species known or suspected to be visited by hummingbirds. Effort within

each site was concentrated at two locations ('stations', Fig. 1D) located on roughly opposite ends

of the focal area (mean ± SD distance between stations: 60 ± 44 m). Stations were centered

around a focal *Heliconia* plant expected to flower throughout the experimental period, not in an

extremely steep area, and >10 m from the fragment edge. The two focal *Heliconia* plants

remained uncovered for the entire experimental period, which allowed us to observe whether

visitation rates changed after declines in surrounding *Heliconia* density. To partially standardize

plant species composition between control and treatment replicates, we added a floral array

comprising ~3-5 additional plant species at each station (see Appendix S2: Table S6 and

'Pollination success').

Trail cameras took one photograph per second during daylight hours (5:30-17:30) and

automatically combined these photographs into a time-lapse video. We reviewed videos for

hummingbird sightings using the motion detection program MotionMeerkat (Weinstein 2015) or

by manually watching the sped-up video. When hummingbirds were detected on camera, we

identified the species, recorded any color marks, and noted whether they visited any flowers.

From video, we also recorded the number of open flowers available per day. Dates with no

available flowers were excluded from analysis, yielding 20,735 total hours of video (Appendix

S2: Table S7).

Pollination success

To investigate whether *Heliconia* declines influenced pollination success, we examined

styles for the presence of pollen tubes, which indicate pollen grain germination and subsequent

movement toward the flower ovule(s) (Kearns and Inouye 1993). We focused our pollen tube

investigation on the two focal *Heliconia* plants per replicate, as well as several species from the

focal arrays (Appendix S2: Table S6). Throughout the experimental period, we collected flowers

from these focal plants every 1-2 days, which corresponds to the maximum floral longevity for

most of the selected species (*Heliconia tortuosa*, *Hamelia patens*: Stratton 1989, *Stachytarpheta*

frantzii: Thomas-Granger 2003). Styles were fixed in formalin-acetic acid-alcohol for at least 24

hours, stained with aniline blue, and examined for pollen tubes using epifluorescence microscopy

(Kearns and Inouye 1993, Betts et al. 2015). Due to low pollen tube presence and/or challenges

with pollen tube visualization, we were only able to analyze two plant species: *H. tortuosa* (N =

327 styles) and *H. patens* (N = 202 styles). Additional details about style collection and lab

procedures are available in Appendix S2.

Additional response variables (Heliconia removal experiment only)

To supplement the primary response variables described above, we examined several

additional metrics for the *Heliconia* removal experiment: (1) recapture probability (i.e.,

proportion of hummingbirds recaptured in the 'post' period), (2) floral visitation rates from

individual hummingbirds marked with nail polish, and (3) body mass of hummingbirds captured

during both experimental periods (pre and post). The first two response variables assessed the

possibility that hummingbirds caught during the initial capture sessions were rapidly replaced by

new individuals, potentially obscuring treatment effects. Body mass measurements allowed us to

explore whether birds that remained in (or returned to) treatment focal areas paid an energetic

cost. To compare mass changes across hummingbirds of different structural sizes, we calculated

a relative body mass measure: observed mass divided by the mass predicted by species-specific

allometric equations (i.e., Ln(body mass) ~ Ln(wing length), Appendix S2: Fig. S5).

Quantifying connectivity

Within each focal landscape, we digitized forested habitat using high resolution (1 m)

Airbus/CNES satellite imagery in Google Earth Pro. When digitizing, we combined closed-

canopy forest and regenerating forest because functional connectivity experiments suggest that

both forest types facilitate hummingbird movement (Jackson 2017). We then delineated

connected areas of forest ('patches') assuming a hummingbird gap-crossing ability of 50 m

(Volpe et al. 2016, Kormann et al. 2016). Lastly, we quantified functional connectivity as the

amount of reachable forest within each focal landscape, following a modified version of

Hanski's (1994) incidence function model for metapopulation connectivity. Substituting forested

pixels for patches used in the original model, we assigned exponentially less importance to pixels

farther away from the focal area (Fig. 1E, Appendix S2: Fig. S6). These weights were calculated

as:

$$W_j = \exp(-\alpha\, d_{ij}), \hspace{4cm} \textbf{(Eq. 1)}$$

where α is the decay rate parameter that scales the effect of distance and d is the shortest

cumulative distance between the focal area centroid i and each forested pixel j, staying within the

patch boundary. The decay rate parameter is inversely related to the average movement distance

of the focal species (Moilanen and Nieminen 2002) and was calculated using the mean home

range length of green hermit hummingbirds (282 m: Volpe et al. 2016). Then, to calculate

functional connectivity, we then summed the weighted pixel areas as follows:

$$S_i = \Sigma\, W_j\, A_j, \hspace{4cm} \textbf{(Eq. 2)}$$

where Wj is the weight of each forested pixel and A is the area of each forested pixel. Geospatial

processing was conducted using Google Earth Engine (Gorelick et al. 2017).

Quantifying resource availability

From the floral resource surveys (see *'Study design'*), we estimated the combined

energetic value of *Heliconia* and non-*Heliconia* flowers. Briefly, we converted resource counts

(i.e., numbers of flowers and inflorescences) into calories, the most biologically relevant unit for

hummingbird energetics (e.g., Wolf et al. 1972, Stiles 1975). To calculate calories, we measured

nectar volume and concentration from >1400 flowers (N = 38 species) growing in the focal

landscape and/or the Wilson Botanical Garden at the Las Cruces Biological Station,

supplementing these measurements with literature data when necessary. We then converted

average nectar volume and concentration per flower to calories per plant. Full methods and

nectar data used in calorie calculations are available in Appendix S2 (Tables S8-S14).

Because tropical hummingbirds generally interact with subsets of the available floral

resources (e.g., Stiles 1985, Maglianesi et al. 2014, Vizentin-Bugoni et al. 2014), we generated

estimates of caloric availability specific to each hummingbird group examined (e.g., all species

versus *Heliconia* specialists). We developed these calorie estimates before conducting any

statistical analysis by summarizing visitation rates to 35 flowering plant species (15,119 video

hours; Appendix S2: Table S7). The most frequently visited plant species for each hummingbird

group was given a weight of one; other species were down-weighted accordingly (Fig. 3, Tables

S15-S16). We divided these calorie estimates by focal area size, then natural log-transformed the

resulting densities. Floral corolla length was measured from flowers collected in the field,

supplemented with literature values when necessary (Appendix S2: Table S17).

Statistical analysis

We used linear mixed models (LMMs) and generalized linear mixed models (GLMMs) to

analyze how experimental *Heliconia* extinction and natural variation in *Heliconia* density

influenced our primary response variables: (i) captured hummingbird abundance, (ii) proportion

of time spent in focal area, (iii) visitation rates to flowering plants (*Heliconia* and non-

Heliconia), and (iv) plant pollination success. Analyses of the *Heliconia* removal experiment

also included several additional, individual-level response variables (see '*Additional response*

variables'). To determine whether declines in *Heliconia* affected *Heliconia* specialists more

strongly than other hummingbird species (**H4**), we analyzed all hummingbird responses with (i)

the full dataset containing all species, and (ii) a dataset including green hermits and violet

sabrewings.

We modeled each response variable using appropriate statistical distributions and

controlled for variable effort when applicable (Appendix S2: Tables S18-S20). All statistical

analyses were conducted in R version 4.1.1 (R Core Team 2021) using 'glmmTMB' (Brooks et

al. 2017). For LMMs, we checked for normality, homoscedasticity, and linearity using the

'performance' package (Lüdecke et al. 2021). For GLMMs, we checked for overdispersion, zero-

inflation, and linearity by visually inspecting residual plots from 'DHARMa' (Hartig 2020).

Estimated marginal means were calculated using 'emmeans' (Lenth 2020) and visualized using

the 'ggemmeans' function within 'ggeffects' (Lüdecke 2018). Results were considered

statistically significant if $P < 0.05$.

Heliconia removal experiment

In a BACI experimental design, an overall treatment effect is indicated by a statistically

significant interaction between the variables representing treatment and experimental period

(Underwood 1994, McDonald et al. 2000). We adopted this standard analytical approach rather

than calculating change scores (e.g., pre-to-post differences) because change scores are nearly

always negatively correlated with baseline measurements; when baseline measurements also

covary with predictor variables of interest, misleading (i.e., spuriously negative) relationships

can emerge (Allison 1990). Therefore, we tested the *parallel extinction hypothesis* (**H1a**) and

robust network hypothesis (**H1b**) using a two-way interaction between *Heliconia* removal and

period (pre *versus* post) for all response variables measured during both experimental periods

(recaptures were measured in the post period only). All models included a random intercept for

'Site'. Except when modeling recapture probability, we also included a nested random effect for

'Replicate ID' (Site-Year), which paired pre-to-post observations within each replicate.

Additional nested random effects, namely for plant species and individual birds, were included

where applicable (Appendix S2: Tables S18-S19). Because we were interested in pre-to-post

changes in our response variables, we only included covariates that varied between pre and post

periods. For example, when analyzing hummingbird visitation rates, we accounted for the

attractiveness of a given focal plant by including a covariate for the number of open flowers per

day, averaged across the experimental period. Since the magnitude of our experimental

Heliconia extinction varied widely (see '*Results*'), we ran parallel models using a categorical

treatment variable (Control/Treatment) and a quantitative treatment variable (Calories

removed/hectare). Because results were qualitatively similar, only results from the categorical

approach are presented in the main text.

Natural variation in Heliconia density

To complement the *Heliconia* removal experiment, we also examined hummingbird

responses over a natural gradient in *Heliconia* availability using data from the 'pre' period of the

experimental sites. This 'space-for-time' approach elucidates how hummingbird space use and

foraging behavior respond to natural variation present in the study area – presumably emerging

over longer time scales than those of our removal experiment. We also used this approach to test

the connectivity hypothesis and resource hypothesis by modeling the interaction term between

Heliconia density and either connectivity (**H2**) or non-*Heliconia* resource availability (**H3**).

These hypotheses were not tested using the experimental framework due to limited sample size

(and thus statistical power) for estimating a three-way interaction between treatment,

experimental period, and resources/connectivity. Because we did not collect resource availability

data in the control sites from 2016, these replicates ($N = 6$) were excluded.

We included several additional covariates not included in the analyses of the *Heliconia*

removal experiment. First, we used estimates of non-*Heliconia* resource availability tailored to

each hummingbird group being examined. Because resource availability was moderately

correlated with connectivity (Pearson's $r = 0.57$, $P = 0.002$, Appendix S2: Fig. S7), we included

both covariates in all models to statistically control for their independent effects. Analyses of

radio telemetry data (i.e., proportion of time spent in focal area) included a covariate for the focal

area size. When analyzing visitation rates of *Heliconia* specialists to non-*Heliconia* resources, we

also included plant corolla length as a covariate, since these relatively long-billed hummingbirds

(mean bill lengths of 40 and 33 mm: Borgella et al. 2001) are expected to select plant species

with long corolla tubes (e.g., Maglianesi et al. 2014, Vizentin-Bugoni et al. 2014, Dalsgaard et

al. 2021). We included random effects for 'Site' in all models; additional random effects were

included where applicable (Appendix S2: Table S20).

RESULTS

Across all study sites and years, *Heliconia* density ranged from 3 to 1131 inflorescences

per ha (mean ± SD: 186 ± 228, median: 145, Appendix S3: Table S1). The magnitude of our

experiment therefore varied widely; we removed between 8 and 526 *Heliconia* inflorescences per

treatment replicate (mean ± SD: 172 ± 138, median: 138, Appendix S3: Table S2). These

inflorescences represented, on average, 96% (± 5.5%) of the *Heliconia* inflorescences

encountered during removal (two focal *Heliconia* plants/site remained uncovered, and sometimes

plant inaccessibility limited our covering capacity). Sizes of focal treatment areas ranged from

0.27 to 7.49 ha (mean ± SD: 1.7 ± 1.9, median: 1.25, Appendix S3: Table S2), depending on the

size of the forest fragment and the navigability of the terrain. We estimate that, on average, the

covered *Heliconia* plants supplied 42% (± 31%, range: 1 – 90%) of the calories available for the

entire hummingbird community ($N = 19$ species total, Appendix S3: Table S3), and at least 66%

(± 32%, range: 2 – 98%) of calories for *Heliconia* specialists (Fig. 2C, Appendix S3: Table S4).

Focal *Heliconia* plants were visited by hummingbirds 6.4 ± 5.7 times per day (range: 0-25,

median: 5), with green hermits visiting most frequently (75% of 1,055 visits, Appendix S3:

Table S5).

Overall responses to changes in Heliconia density (H1)

Hummingbird behavior and pollination success were remarkably resistant to

experimentally induced *Heliconia* extinction. Despite the magnitude of *Heliconia* removal and

confirmed hummingbird visitation to *Heliconia*, we found no overall effect of extirpation on

hummingbird space use, floral visitation rates, or plant pollination success. However, when

examining these response variables across a natural gradient in *Heliconia* availability (i.e., the

space-for-time approach), we found that floral visitation rates and pollination success declined as

Heliconia became scarce. We provide detailed evidence for these findings below.

Hummingbird captures & radio telemetry

We found little evidence that *Heliconia* removal reduced hummingbird abundance or

decreased time spent in the focal area. Although we caught 23% fewer hummingbirds in the

'post' capture session than the 'pre' capture session in treatment replicates (95% CI: -41% to

1%), this pre-to-post decline fell within the natural variability of the control replicates (all

species: $z = -1.11$, $P = 0.27$; *Heliconia* specialists: $z = -0.5$, $P = 0.62$; Fig. 4: A-B; Appendix S3:

Table S6). Similarly, although radio-tagged hummingbirds spent 25-30% less time in focal areas

during the 'post' period, we found no evidence that pre-to-post changes in space use were linked

to our experimental manipulation (95% CI, treatment: -56% to 11%; 95% CI, control: -56% to

28%). Results were similar when examining all species together (Fig. 4C) and *Heliconia*

specialists separately (Fig. 4D). We also found no evidence that natural variation in *Heliconia*

density influenced hummingbird space use, quantified either with mist net captures or radio

telemetry (Appendix S3: Tables S6-S7).

When examining individual-level hummingbird responses, we also did not observe an

effect of *Heliconia* removal. On average, the estimated probability of recapturing a hummingbird

from the 'pre' capture session was 18% in control replicates (95% CI: 11% to 29%), compared to

14% in treatment replicates (95% CI: 9% to 23%). This difference was not statistically

significant, and recaptures of *Heliconia* specialists showed qualitatively similar results

(Appendix S3: Fig. S1: A-B, Table S8). We also found no evidence that pre-to-post changes in

relative body mass differed between control and treatment replicates (Appendix S3: Table S9).

Floral visitation rates & pollination success

We continuously monitored hummingbird visits to the two focal *Heliconia* inflorescences

that remained uncovered throughout the experimental period, as well as numerous other

flowering plants (mean ± SD: 9.3 ± 2 species/replicate; range: 6-14). Under the *robust network*

hypothesis, we expected hummingbirds to rewire and increasingly use alternative (non-

Heliconia) resources. However, pre-to-post changes in hummingbird visitation rate were similar

between control and treatment replicates, both when examining the focal *Heliconia* plants (z =

0.83, P = 0.41, Fig. 5: A-B, Appendix S3: Table S10) and non-*Heliconia* plant species (z = -0.69,

P = 0.49, Fig. 6: A-B, Appendix S3: Table S11). Responses by *Heliconia* specialists mirrored the

responses observed for all species, although these species visited non-*Heliconia* species

relatively infrequently (<1 visit/day, on average: Fig. 6B). Results were similar when examining

pre-to-post changes in visitation rates for individual color-marked hummingbirds observed on

camera (Appendix S3: Fig. S1: C-F, Tables S12-S13).

In contrast to the experimental results, we found that natural variation in *Heliconia*

availability strongly influenced visitation to focal *Heliconia* plants (Fig. 5: D-E). For every

standard deviation decrease in the caloric density of *Heliconia*, we found that overall *Heliconia*

visitation rates decreased by 49% (95% CI: 15% to 93%, $z = 3.03$, $P = 0.0024$, Appendix S3:

Table S10). Visitation to non-*Heliconia* plants also declined when natural *Heliconia* availability

was low, but only for *Heliconia* specialists ($z = 2.02$, $P = 0.044$, Fig. 6E, Appendix S3: Table

S11).

We found no evidence that experimental *Heliconia* removal influenced pollination

success for the focal *Heliconia* plants ($z = 0.23$, $P = 0.82$, Fig. 5C, Appendix S3: Table S14) or

another plant species from the floral arrays (*Hamelia patens: $z = 0.19$, $P = 0.85$*, Appendix S3:

Table S14). However, focal *Heliconia* plants were pollinated less when surrounding *Heliconia*

availability was naturally low; for every standard deviation decrease in the caloric density of

Heliconia, the odds of pollination declined by 2.4 times (95% CI: 1.1 times to 5.4 times, $z = 2.2$,

$P = 0.028$, Fig. 5F).

Effects of connectivity (H2) and resource availability (H3)

Hummingbird captures & radio telemetry

Connectivity to forested habitat influenced hummingbird abundance, and alternative

(non-*Heliconia*) resource availability was associated with the amount of time radio-tagged birds

spent in focal areas. Specifically, hummingbird captures increased by 6% for every one-unit

increase in functional connectivity (95% CI: +1% to +12%, $z = 2.31$, $P = 0.021$, Appendix S3:

Table S6), and the odds of being in the focal area increased by 2.3 times (95% CI: 1.3 to 4.1

times) for every standard deviation increase in caloric density (all species only: $z = 2.78$, $P =$

0.0055, Appendix S3: Table S7). However, neither connectivity or resource availability appeared

to mediate hummingbird responses to *Heliconia* declines, either with experimental removal or

across the natural density gradient ($P > 0.05$ for all interactions of interest, Appendix S3: Tables

S6-S7).

Floral visitation rates & pollination success

Connectivity showed no statistically significant main effects on hummingbird visitation

rates, either to focal *Heliconia* plants or non-*Heliconia* species. However, after *Heliconia*

removal, visitation rates to focal *Heliconia* plants by *Heliconia* specialists declined more sharply

when connectivity was low ($z = -1.76$, $P = 0.08$, Appendix S3: Fig. S2, Table S10). This

potential interactive effect did not emerge when examining *Heliconia* pollination success ($z = -$

0.55, $P = 0.58$). Although we expected that rewiring might be more likely to occur when

alternative resources were abundant, we found no evidence for this hypothesis in the visitation

rates to non-*Heliconia* plant species (Appendix S3: Table S11).

DISCUSSION

In this study, we experimentally induced the local extinction of a locally abundant

tropical plant, *Heliconia tortuosa*, to assess the robustness of plant-hummingbird interaction

networks in a fragmented tropical landscape. Additionally, across a natural gradient in *Heliconia*

density, we assessed whether low *Heliconia* abundance would lead to parallel declines in

hummingbirds and their pollination services – and whether landscape context (connectivity) and

local context (alternative resource availability) could buffer against potential declines. Based on

suite of complementary response variables we found that, even when very high proportions of

total available calories were experimentally removed (~42-66% on average), hummingbird

responses fell within the natural variability of control replicates. As might be expected with such

small changes in hummingbird behavior, we did not observe any cascading effects of

experimental manipulation on pollination success of flowering plants, which supports the *robust*

network hypothesis (**Hlb**). On the other hand, when examining hummingbird responses and

pollination success across a natural gradient in *Heliconia* density, we found evidence for the

parallel extinction hypothesis (**H1a**). In sites where *Heliconia* density was low, hummingbirds

visited individual *Heliconia* plants less frequently, and these plants experienced reduced

pollination success. Other flowering plants also exhibited lower visitation rates from *Heliconia*-

associated hummingbirds, suggesting that high *Heliconia* densities may attract these

hummingbirds (and/or incentivize repeat visitation) and therefore facilitate pollination for other

species.

Our finding that that experimental removal of *Heliconia* did not elicit parallel extinctions

in pollinators aligns with theoretical predictions (Memmott et al. 2004, Kaiser-Bunbury et al.

2010) and several other experiments that removed plant species from mutualistic networks

(Ferrero et al. 2013, Goldstein and Zych 2016, Costa et al. 2018). However, if hummingbirds

remained in forest fragments despite *Heliconia* declines, we assumed that they would rewire to

use alternative floral resources; hummingbirds as a taxonomic group are highly opportunistic

foragers (Arizmendi and Ornelas 1990, Maruyama et al. 2013) and the most frequent *Heliconia*

visitor, the green hermit, has a relatively wide niche breadth (Borgella et al. 2001, Betts et al.

2015). Unexpectedly, we found no evidence that hummingbirds increasingly visited non-

Heliconia resources when *Heliconia* became less available, suggesting that the mechanisms

underlying network robustness in this study may not necessarily align with the rewiring observed

in previous empirical tests (e.g., Timóteo et al. 2016, Kaiser-Bunbury et al. 2017, Costa et al.

2018). This apparent lack of foraging flexibility may emerge from various constraints, such as

morphological trait-matching, search costs, and/or competition from other hummingbirds (see

also Weinstein and Graham 2017). For example, high levels of exploitative competition for non-

defendable resources may have discouraged hummingbirds from modifying their trapline and

risking overlap with another bird's foraging route (Young 1971, Gill 1988, Ohashi and Thomson

2009). Lastly, is possible that fine-scale rewiring did occur but was not detected using our

analytical approach, which aggregated visitation rates to the level of hummingbird species;

further analysis examining the diversity of pollen grains carried by individual hummingbirds

may shed light on the possibility of rewiring at finer scales.

In the absence of rewiring, the resilience in hummingbird behavior, abundance, and

pollination services following experimental *Heliconia* removal is surprising. Brief pauses in

nectar intake can disrupt a hummingbird's daily energy balance, given their high metabolic rate

and limited energy reserves (Calder and Booser 1973, Tooze and Gass 1985, Brice 1992).

Moreover, previous nectar manipulations with hummingbirds – albeit within defended territories

– led to rapid changes in territory size, foraging patterns, and/or defense behavior, generally

within a day (e.g., Hixon et al. 1983, Gass and Sutherland 1985, Eberhard and Ewald 1994,

Temeles et al. 2004). One explanation for our result is that hummingbirds permanently emigrated

from fragments after *Heliconia* removal but, because of high intruder pressure, were quickly

replaced by new individuals (Stiles and Wolf 1970). However, we monitored individual birds

using several different approaches (radio telemetry, recaptures, visitation rates) and found no

evidence that *Heliconia* removal caused emigration and subsequent replacement. An alternative

possibility is that hummingbirds relied on physiological adaptations, such as a torpor and/or

insectivory, to maintain body condition despite large declines in nectar availability – resulting in

the stable body mass measurements we observed in recaptured hummingbirds. Torpor is an

energy-saving adaptation characterized by declines in body temperature and metabolic rate

(Hainsworth et al. 1977, Spence and Tingley 2021), and even tropical hummingbirds inhabiting

low-to-mid elevations appear to possess this capability, including the green hermit (Krüger et al.

1982, Schuchmann and Prinzinger 1988, Bech et al. 1997, Shankar et al. 2020). Additionally,

hummingbirds are highly insectivorous and, although they cannot completely replace nectar with

insects (Brice 1992, Stiles 1995), they may supplement their diet with arthropods when nectar

becomes scarce (Young 1971, Kuban and Neill 1980, Montgomerie and Redsell 1980,

Hazlehurst and Karubian 2018). Finding torpid hummingbirds at night is extremely challenging

(Carpenter and Hixon 1988), but future research could explore changes in arthropod

consumption using techniques such as DNA metabarcoding (Moran et al. 2019).

Another explanation for hummingbird persistence following experimental *Heliconia*

removal involves their high movement ability. *H. tortuosa* is a patchily distributed species that

necessitates long-distance foraging movements, as a single clump does not make territoriality

energetically profitable (Stiles 1975). Densities can also vary substantially across short distances;

for example, Stiles (1979) notes that some Costa Rican *Heliconia* species can be common in one

watershed, yet nearly absent in an adjacent watershed < 1 km away (see also Bruna and Kress

2002). The most frequent *Heliconia* visitor, the green hermit, is therefore capable of traveling

long distances (>500m) while foraging within home ranges of >150 ha, although home range

sizes of ~2 ha are more typical (Volpe et al. 2014, 2016). Thus, returning to our focal areas

(typically ~1.5 ha) to visit the few remaining *Heliconia* resources could be perceived as a

relatively low-cost option for long-distance trapliners, at least in the short term. Notably,

although highly mobile pollinators might buffer against network collapse (Volpe et al. 2016),

they may be vulnerable to habitat loss and fragmentation once thresholds are crossed, leading to

parallel pollinator declines observed in nature (Andrén 1994, Hadley and Betts 2012). This

strategy is also unlikely to be profitable over longer time scales than measured by our

experiment, which only removed *Heliconia* for five days.

The relatively short length of our removal experiment might also explain why – when

examining hummingbird responses across a natural gradient in *Heliconia* availability – lower

Heliconia densities were indeed associated with reduced floral visitation rates. Although a short

adjustment period (<1 day) is standard in similar species removal experiments (e.g., Hixon et al.

1983, Brosi and Briggs 2013, Hazlehurst and Karubian 2018), extending the experimental

timeline may have more realistically simulated a future extinction and provided more time for

behavioral or demographic adjustment. If hummingbirds were not permanent residents of our

focal areas – as suggested by our relatively low recapture rates (~15%) and proportion of time

spent in the focal area (~20%) – then transient individuals might have not immediately noticed

changes in *Heliconia* density and changed their foraging patterns accordingly. Likewise,

hummingbirds might not have persisted had we removed *Heliconia* over larger areas, because

traveling to isolated plants would become even more energetically costly (Heinrich and Raven

1972). Unfortunately, covering (and then uncovering) *Heliconia* plants in steep terrain with

dense understories limited the area we could manipulate.

We also acknowledge that our ability to detect experimental effects may have been

limited by low statistical power, owing to high individual-level variation and limited sample size.

Low statistical power may have also weakened our tests of the connectivity hypothesis and

resource hypothesis, as those datasets only included a subset of the replicates included in the

experimental analyses (resource availability data were not available for all sites). Still, we

emphasize that the spatial scale of our experiment and amount of replication surpasses most

other empirical studies simulating species extinctions within plant-pollinator networks

(Appendix S1: Table S1). Nevertheless, future removal experiments with highly mobile

pollinators should consider increasing the amount of replication, and, if possible, manipulate

larger areas.

In the space-for-time component of our study, we found that visitation rates to individual

plants decreased with natural declines in *Heliconia* density, suggesting that *Heliconia* and its

pollinators might facilitate pollination of co-flowering plants in this study system. Indeed,

facilitative effects arising from the overall attractiveness of the floral neighborhood are expected

to operate most strongly in systems where plants are spatially dispersed and pollinator density is

low (Schemske 1981, Moeller 2004, Ghazoul 2005), as is the case in the tropical forest

understory. If *Heliconia* population densities decrease due to habitat fragmentation (Hadley et al.

2014), plants visited frequently by *Heliconia*-associated hummingbirds might therefore

experience a higher risk of reproductive failure (e.g., *Pitcairnia imbricata*: Fig. 3). Whether

visitation rates map onto pollination success for these species remains an open question, since

pollinator sharing may lead to heterospecific pollen transfer and pollen loss (Morales and

Traveset 2008, Ashman and Arceo-Gómez 2013) – tipping the balance toward interspecific

competition rather than facilitation (Rathcke 1983, Ghazoul 2005, Mitchell et al. 2009).

However, we did discover a positive relationship between visitation rate and pollination success

for *Heliconia*. This relationship has been predicted for other *Heliconia* species but yielded mixed

results (Bruna et al. 2004, Stein et al. 2013), in accordance with accumulating evidence that

facilitation effects can be highly context-specific (Bergamo et al. 2020). Nevertheless, this

finding suggests that any reproductive declines of *Heliconia* in this study system might

accelerate once extinction debt is realized and conspecific density begins to decline.

CONCLUSIONS

Although pollinators seemed able to persist during our experiments that removed floral resources for a short period of time (several days), this persistence may not be sustainable or reflect the outcome of a natural extinction process. In fact, we observed declines in hummingbird visitation when *Heliconia* availability was naturally low, with cascading consequences for pollination success. Thus, longer-term experiments will be key for predicting consequences of species loss and understanding mechanisms of network robustness – at least for pollinator species that forage at large spatial scales. Our study also suggests that behavioral flexibility to use alternative floral resources might be limited, even in taxonomic groups thought to be highly opportunistic foragers. Thus, even rewiring rules with strong biological underpinnings (e.g., Vizentin-Bugoni et al. 2020) may not necessarily predict real-world responses to species loss, perhaps due to additional constraints such as competition. To further understand the mechanisms underlying network robustness or collapse, we encourage future experiments that consider how temporal scale, species traits (e.g., movement ability), competition, and landscape context interact to determine the outcome of species extinctions. These experiments could be especially valuable at geographic range limits, where climate change will likely lead to species replacements (HilleRisLambers et al. 2013), and in the tropics where landscape change and species loss are most rapid (Hansen et al. 2013).

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation (NSF) Graduate Research Fellowship Program under Grant No. 1840998 to KGL. Funding for the experiment was provided primarily by NSF-DEB-1457837 to MGB and ASH, with additional grants to KGL from the American Ornithological Society and Sigma Xi. We also thank the Las Cruces Biological Station and the private landowners who allowed us to work on their property. Field

LITERATURE CITED

Aguilar, R., L. Ashworth, L. Galetto, and M. A. Aizen. 2006. Plant reproductive susceptibility to

habitat fragmentation: review and synthesis through a meta-analysis. Ecology letters

9:968–980.

Aizen, M. A., M. Sabatino, and J. M. Tylianakis. 2012. Specialization and rarity predict

nonrandom loss of interactions from mutualist networks. Science 335:1486–1489.

Allen-Wardell, G., P. Bernhardt, R. Bitner, A. Burquez, S. Buchmann, J. Cane, P. A. Cox, V.

Dalton, P. Feinsinger, M. Ingram, D. Inouye, C. E. Jones, K. Kennedy, P. Kevan, H.

Koopowitz, R. Medellin, S. Medellin-Morales, G. P. Nabhan, B. Pavlik, V. Tepedino, P.

Torchio, and S. Walker. 1998. The Potential Consequences of Pollinator Declines on the

Conservation of Biodiversity and Stability of Food Crop Yields. Conservation Biology

12:8–17.

Allison, P. D. 1990. Change scores as dependent variables in regression analysis. Sociological

Methodology 20:93–114.

Anderson, S. H., D. Kelly, J. J. Ladley, S. Molloy, and J. Terry. 2011. Cascading effects of bird

functional extinction reduce pollination and plant density. Science 331:1068–1071.

Andrén, H. 1994. Effects of habitat fragmentation on birds and mammals in landscapes with

different proportions of suitable habitat: A review. Oikos 71:355–366.

Arizmendi, M. C., and J. F. Ornelas. 1990. Hummingbirds and their floral resources in a tropical

dry forest in Mexico. Biotropica 22:172–180.

Ashman, T. L., and G. Arceo-Gómez. 2013. Toward a predictive understanding of the fitness

costs of heterospecific pollen receipt and its importance in co-flowering communities.

American Journal of Botany 100:1061–1070.

Ashman, T. L., T. M. Knight, J. A. Steets, P. Amarasekare, M. Burd, D. R. Campbell, M. R.

Dudash, M. O. Johnston, S. J. Mazer, and R. J. Mitchell. 2004. Pollen limitation of plant

reproduction: ecological and evolutionary causes and consequences. Ecology 85:2408–

2421.

Aslan, C. E., E. S. Zavaleta, B. Tershy, and D. Croll. 2013. Mutualism Disruption Threatens

Global Plant Biodiversity: A Systematic Review. PLOS ONE 8:e66993.

Bascompte, J., and P. Jordano. 2007. Plant-animal mutualistic networks: the architecture of

biodiversity. Annual Review of Ecology, Evolution, and Systematics 38:567–593.

680 Bascompte, J., P. Jordano, C. J. Melián, and J. M. Olesen. 2003. The nested assembly of plant–

681   animal mutualistic networks. Proceedings of the National Academy of Sciences

682   100:9383–9387.

683 Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks

684   facilitate biodiversity maintenance. Science 312:431–433.

685 Bech, C., A. S. Abe, J. F. Steffensen, M. Berger, and J. E. P. W. Bicudo. 1997. Torpor in three

686   species of Brazilian hummingbirds under semi-natural conditions. The Condor 99:780–

687   788.

688 Bergamo, P. J., N. S. Streher, A. Traveset, M. Wolowski, and M. Sazima. 2020. Pollination

689   outcomes reveal negative density-dependence coupled with interspecific facilitation

690   among plants. Ecology Letters 23:129–139.

691 Betts, M. G., A. S. Hadley, and W. J. Kress. 2015. Pollinator recognition by a keystone tropical

692   plant. Proceedings of the National Academy of Sciences 112:3433–3438.

693 Biesmeijer, J. C., S. P. M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A. P.

694   Schaffers, S. G. Potts, R. Kleukers, C. D. Thomas, J. Settele, and W. E. Kunin. 2006.

695   Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands.

696   Science 313:351–354.

697 Borgella, R., A. A. Snow, and T. A. Gavin. 2001. Species richness and pollen loads of

698   hummingbirds using forest fragments in southern Costa Rica. Biotropica 33:90–109.

699 Brice, A. T. 1992. The essentiality of nectar and arthropods in the diet of the Anna's

700   hummingbird (Calypte anna). Comparative Biochemistry and Physiology Part A:

701   Physiology 101:151–155.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.

Skaug, M. Maechler, and B. M. Bolker. 2017. glmmTMB Balances Speed and Flexibility

Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal

9:378–400.

Brosi, B. J., and H. M. Briggs. 2013. Single pollinator species losses reduce floral fidelity and

plant reproductive function. Proceedings of the National Academy of Sciences

110:13044–13048.

Bruna, E. M., and W. J. Kress. 2002. Habitat Fragmentation and the Demographic Structure of

an Amazonian Understory Herb (Heliconia acuminata). Conservation Biology 16:1256–

1266.

Bruna, E. M., W. J. Kress, F. Marques, and O. F. da Silva. 2004. Heliconia acuminata

reproductive success is independent of local floral density. Acta Amazonica 34:467–471.

Calder, W. A., and J. Booser. 1973. Hypothermia of broad-tailed hummingbirds during

incubation in nature with ecological correlations. Science 180:751–753.

Carpenter, F. L., and M. A. Hixon. 1988. A new function for torpor: fat conservation in a wild

migrant hummingbird. The Condor 90:373–378.

Colwell, R. K., R. R. Dunn, and N. C. Harris. 2012. Coextinction and persistence of dependent

species in a changing world. Annual Review of Ecology, Evolution, and Systematics

43:183–203.

Costa, J. M., J. A. Ramos, L. P. da Silva, S. Timóteo, P. Andrade, P. M. Araújo, C. Carneiro, E.

Correia, P. Cortez, M. Felgueiras, C. Godinho, R. J. Lopes, C. Matos, A. C. Norte, P. F.

Pereira, A. Rosa, and R. H. Heleno. 2018. Rewiring of experimentally disturbed seed

dispersal networks might lead to unexpected network configurations. Basic and Applied Ecology 30:11–22.

Dalsgaard, B., P. K. Maruyama, J. Sonne, K. Hansen, T. B. Zanata, S. Abrahamczyk, R. Alarcón, A. C. Araujo, F. P. Araújo, S. Buzato, E. Chávez-González, A. G. Coelho, P. A. Cotton, R. Díaz-Valenzuela, M. F. Dufke, P. L. Enríquez, M. M. D. Filho, E. Fischer, G. Kohler, C. Lara, F. M. G. Las-Casas, L. R. Lasprilla, A. O. Machado, C. G. Machado, M. A. Maglianesi, T. S. Malucelli, O. H. Marín-Gómez, V. Martínez-García, S. M. de Azevedo-Júnior, E. N. da S. Neto, P. E. Oliveira, J. F. Ornelas, R. Ortiz-Pulido, R. Partida-Lara, B. I. Patiño-González, S. N. de P. Queiroz, M. B. Ramírez-Burbano, A. R. Rech, M. A. Rocca, L. C. Rodrigues, A. M. Rui, I. Sazima, M. Sazima, B. I. Simmons, B. A. Tinoco, I. G. Varassin, M. F. Vasconcelos, J. Vizentin-Bugoni, S. Watts, J. D. Kennedy, C. Rahbek, M. Schleuning, and A. M. M. González. 2021. The influence of biogeographical and evolutionary histories on morphological trait-matching and resource specialization in mutualistic hummingbird–plant networks. Functional Ecology 35:1120–1133.

Dunn, R. R., N. C. Harris, R. K. Colwell, L. P. Koh, and N. S. Sodhi. 2009. The sixth mass coextinction: are most endangered species parasites and mutualists? Proceedings of the Royal Society B: Biological Sciences 276:3037–3045.

Eberhard, J. R., and P. W. Ewald. 1994. Food availability, intrusion pressure and territory size: an experimental study of Anna's hummingbirds (Calypte anna). Behavioral Ecology and Sociobiology 34:11–18.

Farwig, N., D. Bailey, E. Bochud, J. D. Herrmann, E. Kindler, N. Reusser, C. Schüepp, and M. H. Schmidt-Entling. 2009. Isolation from forest reduces pollination, seed predation and insect scavenging in Swiss farmland. Landscape Ecology 24:919–927.

Feinsinger, P., and R. K. Colwell. 1978. Community organization among Neotropical nectar-

feeding birds. American Zoologist 18:779–795.

Ferrero, V., S. Castro, J. Costa, P. Acuña, L. Navarro, and J. Loureiro. 2013. Effect of invader

removal: pollinators stay but some native plants miss their new friend. Biological

Invasions 15:2347–2358.

Garibaldi, L. A., I. Steffan-Dewenter, C. Kremen, J. M. Morales, R. Bommarco, S. A.

Cunningham, L. G. Carvalheiro, N. P. Chacoff, J. H. Dudenhöffer, and S. S. Greenleaf.

2011. Stability of pollination services decreases with isolation from natural areas despite

honey bee visits. Ecology letters 14:1062–1072.

Gass, C. L., and G. D. Sutherland. 1985. Specialization by territorial hummingbirds on

experimentally enriched patches of flowers: energetic profitability and learning. Canadian

Journal of Zoology 63:2125–2133.

Ghazoul, J. 2005. Pollen and seed dispersal among dispersed plants. Biological Reviews 80:413–

443.

Gilbert, L. E. 1980. Food web organization and conservation of neotropical diversity. Pages 11–

33 *in* M. E. Soulé and B. A. Wilcox, editors. Conservation Biology: An Evolutionary-

Ecological Perspective. Sinauer, Sunderland.

Gill, F. B. 1988. Trapline foraging by hermit hummingbirds: competition for an undefended,

renewable resource. Ecology 69:1933–1942.

Goldstein, J., and M. Zych. 2016. What if we lose a hub? Experimental testing of pollination

network resilience to removal of keystone floral resources. Arthropod-Plant Interactions

10:263–271.

Gorelick, N., M. Hancher, M. Dixon, S. Ilyushchenko, D. Thau, and R. Moore. 2017. Google

Earth Engine: Planetary-scale geospatial analysis for everyone. Remote Sensing of

Environment 202:18–27.

Hadley, A. S., and M. G. Betts. 2009. Tropical deforestation alters hummingbird movement

patterns. Biology Letters 5:207–210.

Hadley, A. S., and M. G. Betts. 2012. The effects of landscape fragmentation on pollination

dynamics: absence of evidence not evidence of absence. Biological Reviews 87:526–544.

Hadley, A. S., S. J. K. Frey, W. D. Robinson, and M. G. Betts. 2018. Forest fragmentation and

loss reduce richness, availability, and specialization in tropical hummingbird

communities. Biotropica 50:74–83.

Hadley, A. S., S. J. K. Frey, W. D. Robinson, W. J. Kress, and M. G. Betts. 2014. Tropical forest

fragmentation limits pollination of a keystone understory herb. Ecology 95:2202–2212.

Hainsworth, F. R., B. G. Collins, and L. L. Wolf. 1977. The function of torpor in hummingbirds.

Physiological Zoology 50:215–222.

Hanski, I. 1994. Patch-occupancy dynamics in fragmented landscapes. Trends in Ecology &

Evolution 9:131–135.

Hanski, I. 1998. Metapopulation dynamics. Nature 396:41–49.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models.

Hazlehurst, J. A., and J. O. Karubian. 2018. Impacts of nectar robbing on the foraging ecology of

a territorial hummingbird. Behavioural Processes 149:27–34.

Heinrich, B., and P. H. Raven. 1972. Energetics and pollination ecology. Science 176:597–602.

HilleRisLambers, J., M. A. Harsch, A. K. Ettinger, K. R. Ford, and E. J. Theobald. 2013. How

will biotic interactions influence climate change–induced range shifts? Annals of the

New York Academy of Sciences 1297:112–125.

Hixon, M. A., F. L. Carpenter, and D. C. Paton. 1983. Territory area, flower density, and time

budgeting in hummingbirds: an experimental and theoretical analysis. The American

Naturalist 122:366–391.

Jackson, E. R. 2017. Does the matrix matter? Using radio frequency identification (RFID)

technology to examine hummingbird movements in fragmented tropical landscapes.

Master's thesis, Oregon State University.

Jordán, F. 2009. Keystone species and food webs. Philosophical Transactions of the Royal

Society B: Biological Sciences 364:1733–1741.

Kaiser-Bunbury, C. N., J. Mougal, A. E. Whittington, T. Valentin, R. Gabriel, J. M. Olesen, and

N. Blüthgen. 2017. Ecosystem restoration strengthens pollination network resilience and

function. Nature 542:223–227.

Kaiser-Bunbury, C. N., S. Muff, J. Memmott, C. B. Müller, and A. Caflisch. 2010. The

robustness of pollination networks to the loss of species and interactions: a quantitative

approach incorporating pollinator behaviour. Ecology Letters 13:442–452.

Kearns, C. A., and D. W. Inouye. 1993. Techniques for pollination biologists. University Press

of Colorado, Niwot.

Kearns, C. A., D. W. Inouye, and N. M. Waser. 1998. Endangered mutualisms: the conservation

of plant-pollinator interactions. Annual Review of Ecology and Systematics 29:83–112.

Klein, A.-M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen,

and T. Tscharntke. 2007. Importance of pollinators in changing landscapes for world

crops. Proceedings of the Royal Society B: Biological Sciences 274:303–313.

Koh, L. P., R. R. Dunn, N. S. Sodhi, R. K. Colwell, H. C. Proctor, and V. S. Smith. 2004.

Species Coextinctions and the Biodiversity Crisis. Science 305:1632–1634.

Kormann, U., C. Scherber, T. Tscharntke, N. Klein, M. Larbig, J. J. Valente, A. S. Hadley, and

M. G. Betts. 2016. Corridors restore animal-mediated pollination in fragmented tropical

forest landscapes. Proceedings of the Royal Society B: Biological Sciences

283:20152347.

Krüger, K., R. Prinzinger, and K.-L. Schuchmann. 1982. Torpor and metabolism in

hummingbirds. Comparative Biochemistry and Physiology Part A: Physiology 73:679–

689.

Kuban, J. F., and R. L. Neill. 1980. Feeding ecology of hummingbirds in the highlands of the

Chisos Mountains, Texas. The Condor 82:180–185.

Kuussaari, M., R. Bommarco, R. K. Heikkinen, A. Helm, J. Krauss, R. Lindborg, E. Öckinger,

M. Pärtel, J. Pino, F. Roda, and others. 2009. Extinction debt: a challenge for biodiversity

conservation. Trends in ecology & evolution 24:564–571.

Lenth, R. 2020. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lüdecke, D. 2018. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

Journal of Open Source Software 3:772.

Lüdecke, D., M. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. performance: An

R Package for Assessment, Comparison and Testing of Statistical Models. Journal of

Open Source Software 6:3139.

Maglianesi, M. A., N. Blüthgen, K. Böhning-Gaese, and M. Schleuning. 2014. Morphological

traits determine specialization and resource use in plant–hummingbird networks in the

neotropics. Ecology 95:3325–3334.

Maruyama, P. K., G. M. Oliveira, C. Ferreira, B. Dalsgaard, and P. E. Oliveira. 2013. Pollination

syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna

hummingbirds. Naturwissenschaften 100:1061–1068.

McDonald, T. L., W. P. Erickson, and L. L. McDonald. 2000. Analysis of count data from

before-after control-impact studies. Journal of Agricultural, Biological, and

Environmental Statistics 5:262–279.

Memmott, J., N. M. Waser, and M. V. Price. 2004. Tolerance of pollination networks to species

extinctions. Proceedings of the Royal Society of London B: Biological Sciences

271:2605–2611.

Mendenhall, C. D., and A. M. Wrona. 2018. Improving tree cover estimates for fine-scale

landscape ecology. Landscape Ecology 33:1691–1696.

Mitchell, R. J., R. J. Flanagan, B. J. Brown, N. M. Waser, and J. D. Karron. 2009. New frontiers

in competition for pollination. Annals of Botany 103:1403–1413.

Moeller, D. A. 2004. Facilitative interactions among plants via shared pollinators. Ecology

85:3289–3301.

Moilanen, A., and M. Nieminen. 2002. Simple Connectivity Measures in Spatial Ecology.

Ecology 83:1131–1145.

Montero-Castaño, A., and M. Vilà. 2012. Impact of landscape alteration and invasions on

pollinators: a meta-analysis. Journal of Ecology 100:884–893.

Montgomerie, R. D., and C. A. Redsell. 1980. A nesting hummingbird feeding solely on

arthropods. The Condor 82:463.

Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: magnitude, prevalence and

consequences for plant fitness. Critical Reviews in Plant Sciences 27:221–238.

Moran, A. J., S. W. J. Prosser, and J. A. Moran. 2019. DNA metabarcoding allows non-invasive

identification of arthropod prey provisioned to nestling Rufous hummingbirds

(Selasphorus rufus). PeerJ 7:e6596.

Ohashi, K., and J. D. Thomson. 2009. Trapline foraging by pollinators: its ontogeny, economics

and possible consequences for plants. Annals of Botany 103:1365–1378.

Ollerton, J., R. Winfree, and S. Tarrant. 2011. How many flowering plants are pollinated by

animals? Oikos 120:321–326.

Pauw, A., and J. A. Hawkins. 2011. Reconstruction of historical pollination rates reveals linked

declines of pollinators and plants. Oikos 120:344–349.

Phillips, R. D., R. Peakall, B. A. Retter, K. Montgomery, M. H. M. Menz, B. J. Davis, C. Hayes,

G. R. Brown, N. D. Swarts, and K. W. Dixon. 2015. Pollinator rarity as a threat to a plant

with a specialized pollination system. Botanical Journal of the Linnean Society 179:511–

525.

Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin. 2010.

Global pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution

25:345–353.

Powers, D. R., A. R. Brown, and J. A. Van Hook. 2003. Influence of normal daytime fat

deposition on laboratory measurements of torpor use in territorial versus nonterritorial

hummingbirds. Physiological and Biochemical Zoology 76:389–397.

R Core Team. 2021. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Ramos-Jiliberto, R., F. S. Valdovinos, P. Moisset de Espanés, and J. D. Flores. 2012.

Topological plasticity increases robustness of mutualistic networks. Journal of Animal

Ecology 81:896–904.

Rathcke, B. 1983. Competition and facilitation among plants for pollination. Pages 305–329 *in*

L. A. Real, editor. Pollination biology. Academic Press, New York.

Regan, E. C., L. Santini, L. Ingwall-King, M. Hoffmann, C. Rondinini, A. Symes, J. Taylor, and

S. H. M. Butchart. 2015. Global trends in the status of bird and mammal pollinators.

Conservation Letters 8:397–403.

Renjifo, L. M. 2001. Effect of natural and anthropogenic landscape matrices on the abundance of

subandean bird species. Ecological Applications 11:14–31.

Rodríguez-Flores, C. I., J. F. Ornelas, S. Wethington, and M. C. Arizmendi. 2019. Are

hummingbirds generalists or specialists? Using network analysis to explore the

mechanisms influencing their interaction with nectar resources. PLOS ONE

14:e0211855.

Russell, S. M., and R. O. Russell. 2001. The North American banders' manual for banding

hummingbirds. North American Banding Council.

Schemske, D. W. 1981. Floral convergence and pollinator sharing in two bee-pollinated tropical

herbs. Ecology 62:946–954.

Scheper, J., M. Reemer, R. van Kats, W. A. Ozinga, G. T. J. van der Linden, J. H. J. Schaminée,

H. Siepel, and D. Kleijn. 2014. Museum specimens reveal loss of pollen host plants as

key factor driving wild bee decline in The Netherlands. Proceedings of the National

Academy of Sciences 111:17552–17557.

Schuchmann, K.-L., and R. Prinzinger. 1988. Energy metabolism, nocturnal torpor, and

respiration frequency in a Green Hermit (Phaethornis guy). Journal für Ornithologie

129:469–472.

Senapathi, D., L. G. Carvalheiro, J. C. Biesmeijer, C.-A. Dodson, R. L. Evans, M. McKerchar,

R. D. Morton, E. D. Moss, S. P. M. Roberts, W. E. Kunin, and S. G. Potts. 2015. The

impact of over 80 years of land cover changes on bee and wasp pollinator communities in

England. Proceedings of the Royal Society B: Biological Sciences 282:20150294.

Shankar, A., R. J. Schroeder, S. M. Wethington, C. H. Graham, and D. R. Powers. 2020.

Hummingbird torpor in context: duration, more than temperature, is the key to nighttime

energy savings. Journal of Avian Biology 51:e02305.

Spence, A. R., and M. W. Tingley. 2021. Body size and environment influence both intraspecific

and interspecific variation in daily torpor use across hummingbirds. Functional Ecology

35:870–883.

Steffan-Dewenter, I., and T. Tscharntke. 1999. Effects of habitat isolation on pollinator

communities and seed set. Oecologia 121:432–440.

Stein, K., M. Templin, I. Hensen, M. Fischer, D. Matthies, and M. Schleuning. 2013. Negative

Effects of Conspecific Floral Density on Fruit Set of Two Neotropical Understory Plants.

Biotropica 45:325–332.

Stiles, F. G. 1975. Ecology, flowering phenology, and hummingbird pollination of some Costa

Rican Heliconia species. Ecology 56:285–301.

Stiles, F. G. 1979. Notes on the natural history of Heliconia (Musaceae) in Costa Rica. Notas

sobre la historia natural de Heliconia (Musaceae) en Costa Rica. Brenesia 15:151–180.

Stiles, F. G. 1985. Seasonal patterns and coevolution in the hummingbird-flower community of a

Costa Rican subtropical forest. Ornithological Monographs 36:757–787.

Stiles, F. G. 1995. Intraspecific and interspecific variation in molt patterns of some tropical

hummingbirds. The Auk 112:118–132.

Stiles, F. G., and L. L. Wolf. 1970. Hummingbird territoriality at a tropical flowering tree. The

Auk 87:467–491.

Stratton, D. A. 1989. Longevity of Individual Flowers in a Costa Rican Cloud Forest: Ecological

Correlates and Phylogenetic Constraints. Biotropica 21:308–318.

Suarez, R. K. 1992. Hummingbird flight: sustaining the highest mass-specific metabolic rates

among vertebrates. Experientia 48:565–570.

Taki, H., P. G. Kevan, and J. S. Ascher. 2007. Landscape effects of forest loss in a pollination

system. Landscape Ecology 22:1575–1587.

Taylor, J., and S. A. White. 2007. Observations of hummingbird feeding behavior at flowers of

Heliconia beckneri and H. tortuosa in southern Costa Rica. Ornitologia Neotropical

18:133–138.

Taylor, P. D., L. Fahrig, K. Henein, and G. Merriam. 1993. Connectivity is a vital element of

landscape structure. Oikos:571–573.

Temeles, E. J., A. B. Muir, E. B. Slutsky, and M. N. Vitousek. 2004. Effect of food reductions on

territorial behavior of purple-throated caribs. The Condor 106:691–695.

Thomas-Granger, K. 2003. Light sum, nitrogen level and growth retardant effects on flower and

nectar production in Stachytarpheta frantzii. Master's thesis.

Timóteo, S., J. A. Ramos, I. P. Vaughan, and J. Memmott. 2016. High resilience of seed

dispersal webs highlighted by the experimental removal of the dominant disperser.

Current Biology 26:910–915.

Tooze, Z. J., and C. L. Gass. 1985. Responses of rufous hummingbirds to midday fasts. Canadian

Journal of Zoology 63:2249–2253.

Traveset, A., C. Tur, and V. M. Eguíluz. 2017. Plant survival and keystone pollinator species in

stochastic coextinction models: role of intrinsic dependence on animal-pollination.

Scientific Reports 7:6915.

Underwood, A. J. 1994. On Beyond BACI: sampling designs that might reliably detect

environmental disturbances. Ecological Applications 4:3–15.

Valdovinos, F. S., P. Moisset de Espanés, J. D. Flores, and R. Ramos-Jiliberto. 2013. Adaptive

foraging allows the maintenance of biodiversity of pollination networks. Oikos 122:907–

917.

Vellend, M., K. Verheyen, H. Jacquemyn, A. Kolb, H. V. Calster, G. Peterken, and M. Hermy.

2006. Extinction Debt of Forest Plants Persists for More Than a Century Following

Habitat Fragmentation. Ecology 87:542–548.

Vizentin-Bugoni, J., V. J. Debastiani, V. A. G. Bastazini, P. K. Maruyama, and J. H. Sperry.

2020. Including rewiring in the estimation of the robustness of mutualistic networks.

Methods in Ecology and Evolution 11:106–116.

Vizentin-Bugoni, J., P. K. Maruyama, and M. Sazima. 2014. Processes entangling interactions in

communities: forbidden links are more important than abundance in a hummingbird–

plant network. Proceedings of the Royal Society B: Biological Sciences 281:20132397.

Volpe, N. L., A. S. Hadley, W. D. Robinson, and M. G. Betts. 2014. Functional connectivity

experiments reflect routine movement behavior of a tropical hummingbird species.

Ecological Applications 24:2122–2131.

Volpe, N. L., W. D. Robinson, S. J. K. Frey, A. S. Hadley, and M. G. Betts. 2016. Tropical forest

fragmentation limits movements, but not occurrence of a generalist pollinator species.

PLOS ONE 11:e0167513.

Weiner, C. N., M. Werner, K. E. Linsenmair, and N. Blüthgen. 2014. Land-use impacts on

plant–pollinator networks: interaction strength and specialization predict pollinator

declines. Ecology 95:466–474.

Weinstein, B. G. 2015. MotionMeerkat: integrating motion video detection and ecological

monitoring. Methods in Ecology and Evolution 6:357–362.

Weinstein, B. G., and C. H. Graham. 2017. Persistent bill and corolla matching despite shifting

temporal resources in tropical hummingbird-plant interactions. Ecology Letters 20:326–

335.

Winfree, R., I. Bartomeus, and D. P. Cariveau. 2011. Native Pollinators in Anthropogenic

Habitats. Annual Review of Ecology, Evolution, and Systematics 42:1–22.

Wolf, L. L., F. R. Hainsworth, and F. G. Stiles. 1972. Energetics of Foraging: Rate and

Efficiency of Nectar Extraction by Hummingbirds. Science 176:1351–1352.

Wolf, L. L., and F. G. Stiles. 1989. Adaptations for the "fail-safe" pollination of specialized

ornithophilous flowers. The American Midland Naturalist 121:1–10.

Young, A. M. 1971. Foraging for insects by a tropical hummingbird. The Condor 73:36–45.

Zahawi, R. A., G. Duran, and U. Kormann. 2015. Sixty-seven years of land-use change in

southern Costa Rica. PLOS ONE 10:e0143554.

 TABLES

 Table 1. Hypotheses testing whether declines of a locally common, tropical flowering plant (*Heliconia tortuosa*) affect hummingbird

994   space use, hummingbird floral visitation rates, and plant pollination success. Predicted effects are indicated in the column below each

995   hypothesis. To address Question 1, we (i) experimentally removed *H. tortuosa* using a BACI (Before-After-Control-Impact)

996   experimental design and (ii) studied a natural gradient in *H. tortuosa* density. Under the experimental framework, predicted effects can

997   be conceptualized as pre-to-post changes in treatment replicates, compared to pre-to-post changes in control replicates. To address

998   Question 2, we examined changes in hummingbird and plant responses across a natural gradient in *H. tortuosa* density only.

| | Response variable | Methodological approach | **Question 1: How do hummingbirds respond to declines in *Heliconia*, and are there cascading effects on the remaining plants?** | | **Question 2: What factors mediate hummingbird responses to *Heliconia* declines?** | | |
			Robust network hypothesis (H-1a)	Parallel extinction hypothesis (H-1b)	Connectivity hypothesis: Rescue (H-2a) *In highly connected forest fragments…*	Connectivity hypothesis: Abandonment (H-2b) *In highly connected forest fragments…*	Resource hypothesis (H-3) *In forest fragments with high resource availability…*
Space use	Abundance	Mist net captures	No change	Decrease			
	Time spent in focal area	Radio telemetry					
Floral visitation rates	Visitation to focal *Heliconia* plants	Remote camera observations	No change or increase	Decrease	Smaller decrease	Larger decrease	Smaller decrease
	Visitation to non-*Heliconia* plants	Remote camera observations	Increase (rewiring)	Decrease			
Pollination services	Pollination success	Pollen tubes	No change or increase	Decrease			

1000

FIGURES

Figure 1. This study was conducted in the fragmented landscape surrounding the Las Cruces

Biological Station in southern Costa Rica **(A-B).** Forested areas are shown in grey. The dark grey

forest layer was hand-digitized for this study; the light grey forest layer, available within the

geographic boundary of Coto Brus Canton, was obtained from Mendenhall & Wrona *(2018)*.

Non-forested areas (white) primarily comprise pasture. **(A)** Within the study system (rectangular

extent), we utilized fourteen focal landscapes of 1000 m radius ('sites', dashed circles). Sites are

numbered by increasing connectivity to other areas of forest (1 = least connected, 14 = most

connected). Within a year (2016-2018), each site received either the *Heliconia* removal treatment

or was assigned to be a control. Treatments were reversed in alternate years for a total sample

size of 16 control replicates and 16 treatment replicates. **(C)** An example site, with red line

representing a focal area [i.e., where we removed *Heliconia* (for treatment replicates) and/or

surveyed for floral resources (for treatment and control replicates)]. **(D)** An example focal area,

showing where we captured hummingbirds using mist nets (triangles) and monitored plant-

hummingbird interactions using trail cameras (camera icons). **(E)** An illustration of how we

calculated functional connectivity. We first identified any forested pixels (1 x 1 m) that were

connected to the focal area, assuming a hummingbird gap-crossing ability of 50 m; in the

example shown, all pixels were connected to the focal area (center of map). We then down-

weighted pixel areas based on their distance to the focal area centroid (higher weights = closer to

focal area). Finally, we summed all weighted pixel areas within the 1000 m focal landscape (see

Appendix S2 for details).

Figure 2. (A) During the *Heliconia* removal experiment, we measured four main response

variables: captured hummingbird abundance, proportion of time radio-tagged hummingbirds

spent in the focal area, floral visitation rates, and plant pollination success. Each response

variable was measured during both experimental periods (pre and post); the sampling schedule

for each is denoted with an 'X'. Each experimental replicate used two sites (control and

treatment) sampled using this schedule, offset by one day. Starting around noon on the fourth

day, we covered *Heliconia* inflorescences in the treatment site. **(B)** *Heliconia* inflorescences

were covered with dark-colored bags that prevented hummingbirds from accessing any nectar

present. **(C)** Our manipulation removed, on average, 42% (± 31% SD) of the calories available

for the entire hummingbird community and 66% (± 32% SD) of the available calories for

Heliconia specialists (green hermits and violet sabrewings). Mean values are denoted by red

diamonds, and the darker black line represents the median. Lower and upper lines within

boxplots represent the first and third quartiles. Whiskers extend from the first and third quartiles

to the lowest and highest values within 1.5 times the interquartile range. Individual data points

1038   appear next to each boxplot. *Response variable icons in (A) were created using images from the*

1039   *Noun Project (see 'Acknowledgments'). Photos in (B) were provided by Marion Donald.*

Figure 3. Relative visitation rates (weights) used to tailor our resource availability estimates to different groups of hummingbird species. Within each hummingbird group, the most frequently visited plant species was assigned a weight of one. Visitation rates are summarized from the 15,119 hours of 'normal' (i.e., unmanipulated) camera observations collected in the landscape surrounding the Las Cruces Biological Station across 14 study sites, three field seasons (2016-2018), and 35 plant species. **(A)** Weights summarized across all hummingbird species detected on camera (*N* = 17 species). **(B)** Weights for green hermits (*P. guy*) and violet sabrewings (*C. hemileucurus*), two hummingbird species with morphologically specialized bill shapes that mirror the curvature of *Heliconia* flowers. These two species are also the most frequent *Heliconia* visitors (Appendix S3: Table S5).

1052

Figure 4. Hummingbird space use before and after experimental local extinction of an

understory plant, *Heliconia tortuosa*, within forest fragments in southern Costa Rica. Despite

removing at least 42% (± 31 SD) of calories available to hummingbirds within focal areas of 1.7

ha (± 1.9 SD), we did not observe an effect of *Heliconia* removal on the number of

hummingbirds captured (A-B) or the proportion of time that radio-tagged hummingbirds spent in

the focal areas (C-D). Results are shown for all species analyzed (left) and green hermits and

violet sabrewings (right). Estimated marginal means from GLMMs are presented alongside 95%

confidence intervals. *P*-values refer to the interaction testing the effect of *Heliconia* removal,

relative to control replicates; conceptually, a treatment effect is indicated by non-parallel lines.

The estimated number of captured hummingbirds is calculated for 45.8 net-hours, the mean

capture effort across all replicates.

1064

Figure 5. Hummingbird visitation rates to focal *Heliconia* plants and *Heliconia* pollination

success before and after experimental *Heliconia* removal (A-C) and across a natural gradient in

Heliconia density (D-F). For hummingbird visitation rates, results are shown for the entire

hummingbird community (A, D) and green hermits and violet sabrewings separately (B, E).

Estimated marginal means from GLMMs are presented alongside 95% confidence intervals. (A-

C) *P*-values refer to the interaction testing the effect of *Heliconia* removal, relative to control

replicates; conceptually, a treatment effect is indicated by non-parallel lines. (D-F) Values of

Heliconia availability (*x*-axis) are standard deviations (1 SD = 1.35), with zero representing the

mean log-transformed caloric density (mean = 8). Note that the *x*-axis is reversed, such that

declining visitation is associated with declining *Heliconia* availability. *P*-values refer to the

coefficient for *Heliconia* availability.

1076

Figure 6. Hummingbird visitation rates to non-*Heliconia* plant species before and after

experimental *Heliconia* removal (A-B) and across a natural gradient in *Heliconia* density (C-D).

Results are shown for the entire hummingbird community (A, C) and green hermits and violet

sabrewings separately (B, D). Estimated marginal means from GLMMs are presented alongside

95% confidence intervals. (A-B) *P*-values refer to the interaction testing the effect of *Heliconia*

removal, relative to control replicates; conceptually, a treatment effect is indicated by non-

parallel lines. (C-D) Values of *Heliconia* availability (*x*-axis) are standard deviations (1 SD =

1.35), with zero representing the mean log-transformed caloric density (mean = 8). Note that the

x-axis is reversed, such that declining visitation is associated with declining *Heliconia*

availability. *P*-values refer to the coefficient for *Heliconia* availability.