1 *Title*
2 **Copepod functional traits and groups show divergent biogeographies in the global ocean**
3
Running title
Global marine copepod traits biogeography

Authors
Fabio Benedetti[1*] (0000-0002-7554-3646)

Jonas Wydler[1]
Meike Vogt[1] (0000-0002-0608-1935)

11
[1]Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, 8092
Zürich, Switzerland.
*Corresponding author: fabio.benedetti@usys.ethz.ch

Acknowledgements
This project has received funding from the European Union's Horizon 2020 research and

innovation program under grant agreement No. 862923. This output reflects only the author's

view and the European Union cannot be held responsible for any use that may be made of the

information contained therein. We thank all contributors involved in the plankton species

field sampling and identification throughout the world and we acknowledge the efforts made

to deposit the data on publicly available online archives. We thank Luke Gregor for editing

the language of an early version of the manuscript. We thank Dr. Maria Grazia Mazzocchi

and an anonymous reviewer for their constructive comments. No permits were needed to

carry out the present study.

26
Conflict of interest
The authors declare no conflict of interests.

Abstract

Aim: The distribution of zooplankton functional traits is a key factor for regulating food web

dynamics and carbon cycling in the oceans. Yet, we lack a clear understanding of how many

functional groups (FGs) exist in the zooplankton and how their traits are distributed on a

global scale. Here, we model and map the environmental habitats of copepod (i.e. the main

component of marine zooplankton) FGs to identify regions sharing similar functional trait

expression, at the community level.

Taxon: Marine planktonic Neocopepoda.

Location: Global ocean.

Methods: Factor analysis on mixed data and hierarchical clustering were used to identify

copepod FGs based on five species-level functional traits. An ensemble of species distribution

models was used to estimate the environmental niches of the species modelled and the

community weighted mean values of the traits studied. Ocean regions were defined based on

their community-level mean trait expression using a principal component analysis and

hierarchical clustering.

Results: Eleven global copepod FGs were identified. They displayed contrasting latitudinal

patterns in mean annual habitat suitability that could be explained by differences in

environmental niche preferences: two FGs were associated with polar conditions, one

followed the global temperature gradient, five were associated with tropical oligotrophic

gyres, and the remaining three with boundary currents and counter currents. Four main

regions of varying community weighted mean trait values emerged: the Southern Ocean, the

northern and southern high latitudes, the tropical gyres, and the boundary currents and

upwelling systems.

Conclusions: The present FGs will improve the representation of copepods in global marine

ecosystem models. This study improves the understanding of the patterns and drivers of

copepods trait biogeography and will serve as a basis for studying links between zooplankton

biodiversity and ecosystem functioning in a context of climate change.

Keywords

Community weighted mean trait, Functional groups, Global Ocean, Marine copepods, Species

Distribution Models, Trait-based approach, Zooplankton

1. Introduction

Copepods are crustaceans that dominate the biomass of the mesozooplankton (0.2-2.0 mm) and rank amongst the most abundant animals in the oceans (Huys & Boxshall, 1991). They are morphologically and functionally diverse and adapted to all aquatic ecosystems, with particular success in marine systems (Huys & Boxshall, 1991; Kiørboe, 2011a). Copepods play a pivotal role in food webs, both as microplankton grazers and prey for higher trophic levels (Beaugrand, Edwards, & Legendre, 2010). Copepods also play a critical role in the marine biological carbon pump by performing extensive vertical migrations, grazing on phytoplankton in the sunlight layers and releasing relatively fast-sinking fecal pellets, exporting organic carbon with dead carcasses down to the deep layers (Turner, 2015; Steinberg & Landry, 2017).

The relative contribution of planktonic copepods to the above-mentioned processes is mediated by their diversity and trait expression (Barton et al., 2013). For instance, larger copepods perform stronger diel vertical migrations and produce larger fecal pellets that sink faster and thus increase the proportion of carbon exported to depth (Stamieszkin et al., 2015; Ohman & Romagnan, 2016). The dominance of certain copepod feeding modes impacts food-web dynamics as feeding modes affect grazing and mortality rates (Kenitz, Visser, Mariani, & Andersen, 2017; van Someren Gréve, Almeda, & Kiørboe, 2017). Yet, mechanistic ecosystem models usually represent zooplankton only through a few size classes (Le Quéré et al., 2005), which oversimplifies the contribution of species and functional diversity to ecosystem functioning and biogeochemical cycles (Flynn et al., 2015). Multiple studies show that the number and grazing characteristics of zooplankton functional groups (FGs) control the biomass and diversity of other functional groups, with consequences for global biogeochemical cycles (Prowe et al., 2012; Sailley et al., 2013; Vallina et al., 2014; Le Quéré et al., 2016). Ecosystem models rely strongly on parameterisations of traits governing biological processes and food-web interactions based on scarce empirical evidence (Barton et al., 2013). Consequently, investigating the potential links between copepod trait distribution and ecosystem functioning contributes to improving these models (Stocker, 2014).

To better represent the role of zooplankton in marine systems, ecologists are increasingly adopting trait-based approaches (Litchman, Ohman, & Kiørboe, 2013; Hébert, Beisner, & Maranger, 2017). Functional traits are species- or organism-level characteristics that affect their fitness and are linked to survival, feeding, growth, and reproduction (Violle et al., 2007; Litchman et al. 2013). Due to trade-offs between them, organisms cannot maximize all functions at the same time. For example, copepods that rely on active ambush feeding (i.e.,

copepods that only move to capture prey) show 8.5 times lower mortality rates than copepods

that rely on active feeding strategies (van Someren Gréve, Almeda, & Kiørboe, 2017). Yet,

adult male copepods need to move actively to find a mate for reproduction, which undermines

the benefits of their feeding mode (Kiørboe, 2011b; Litchman et al., 2013).

Copepod traits dynamics can be investigated by grouping species into FGs with similar trait

combinations (Benedetti, Gasparini, & Ayata, 2016; Becker et al., 2021). Categorizing

species according to their similarity in functional traits rather than their taxonomic

classification enables to summarize their diversity into distinct, parsimonious groups. At the

regional scale, Pomerleau et al. (2015) showed that 42 zooplankton species in the North East

Pacific Ocean could be clustered into five groups based on their body length, feeding and

reproduction mode, and trophic group. Similarly, Benedetti, Vogt et al. (2018) grouped 106

dominant Mediterranean copepod species into seven FGs with distinct environmental niche

characteristics. Such groupings can improve the representation of zooplankton in ecosystem

models since FGs increase the representation of ecological function without adding

taxonomic diversity and complexity.

The functional composition of zooplankton can be examined using functional diversity

indices which indicate how changes in species richness and composition affect the emergent

expression of traits (Villéger, Mason & Mouillot, 2008). These indices allow to investigate

biodiversity and ecosystem functioning relationships as they indicate changes in ecological

processes that drive community assembly. For instance, Becker et al. (2021) documented a

poleward decrease in copepod functional diversity in the South Atlantic, which was driven by

the decreasing contribution of cruising carnivores and small and medium-sized broadcasting

copepods. The authors attributed this poleward decrease in functional diversity to niche

partitioning as only a subset of functionally very dissimilar groups of copepods thrived in the

cold and productive waters of the Southern Ocean.

The functional composition of zooplankton can also be investigated through community

weighted mean (CWM) values of traits (Ricotta, 2005; Brun et al., 2016), i.e. the proportion

of species in a community that exhibits a particular trait value weighed by their relative

contribution to community composition. Thereby, CWM values of functional traits help to

identify the relationships between the emerging expression of functional traits on a

community-level, the environmental drivers of their biogeographical patterns of trait

expression, and the potential trade-offs underlying trait expression. The first truly global study

by Brun et al. (2016) found larger copepod body sizes, higher proportions of myelinated

species, and relatively smaller clutch size at latitudes >50°, associated with gradients in

temperature and phytoplankton seasonality. To our knowledge, other qualitative traits such as
trophic group or spawning mode have been investigated but not using a global CWM trait
approach (McGinty et al., 2018; Benedetti, Vogt et al., 2018). If patterns of CWM values are
clustered in space and time, they can be used to define ocean regions sharing common
ecological characteristics (Longhurst, 2010; Reygondeau et al., 2017; Hofmann Elizondo et
al., 2021). Although new data compilations make it possible, neither copepod FGs nor CMW-
based regionalization schemes have been investigated on a global scale, yet this could be
highly informative of large-scale marine food web and ecosystem dynamics.
Here, we use hierarchical clustering and a state-of-the-art species distribution models (SDMs)
framework to address the following questions: (i) What are the global copepod FGs emerging
from the clustering of the functional traits of commonly sampled copepod species? (ii) Do
copepod FGs show different environmental niche preferences (i.e. niche centers and widths)?
(iii) What are the habitat distribution patterns of these FGs and their community-level trait
expression, and which regions of the global ocean share similar trait expression?
144
145 **2. Materials and Methods**
146 Figure 1 summarizes the analytical framework we developed to identify emergent copepod
147 FGs and their habitats in the global ocean using an ensemble species distribution model
148 (SDMs) approach.
149

Figure 1: Flowchart summarizing the numerical analyses implemented in our study. SDMs =
Species distribution models, PCA = Principal component analysis.

2.1. Species occurrence data

We used the zooplankton occurrences dataset (geolocated and dated presences) compiled by

Benedetti et al. (2021). Occurrences corresponding to benthic and parasitic copepods,

occurrences with missing spatial coordinates, sampling dates, sampling depth, or not

identified at species-level were removed. Data from drilling holes, grid cells within 25 km of

the nearest shoreline, or with a maximum sampling depth >500 m were also discarded (see

Benedetti et al., 2021). To address sampling effort biases in the copepod occurrence data

(Appendix S1) that could inflate models performance metrics or over-represent portions of the

environmental space and thus hinder model predictability and interpretation (Veloz, 2009;

Hijmans, 2012), we thinned occurrences using a randomization algorithm for each month and

each species (30 randomizations per species dataset) so that monthly species occurrences

were at least 500 km apart (Aiello-Lammens, Boria, Radosavljevic, Vilela, & Anderson,

2015). Only those 385 species displaying at least 50 occurrences were retained to constitute
the list of species for which functional traits data were retrieved (Appendix S2).
168
2.2. Species functional trait data
Five functional traits were included based on data availability from the literature (Appendices
S2 and S3):
Body size (quantitative continuous): mean maximum adult female body size (i.e. length of the
cephalothorax) in mm. Body size is considered a master trait as it impacts all life functions,
scales with most physiological rates and influences predator-prey interactions (Hansen,
Bjornsen, & Hansen, 1994; Kiørboe & Hirst, 2014; Hébert, Beisner, & Maranger, 2017). The
mean maximum body size values used here were derived from Razouls et al. (2015) and range
from 0.46 to 11.00 mm.
Trophic group (categorical): most marine planktonic copepods are omnivorous but they can
be grouped according to their primary food source to describe their role in food-web
dynamics (Pomerleau et al., 2015; Benedetti et al., 2016). Here, the following five groups
were defined: Omnivore-Herbivore, Omnivore-Carnivore, Omnivore-Detritivore, strict
Carnivore, and Omnivore.
Feeding mode (categorical): copepods rely on various strategies to detect and capture their
prey. We follow the definitions of Kiørboe (2011a): ambush-feeding, current-feeding, cruise-
feeding, particle-feeding, current-cruise feeding, and current-ambush feeding (the last two
refer to mixed-feeding species with both strategies). Ambush-feeding copepods lurk in the
water column, detect the vibration generated by motile preys and capture them through quick
jumps. Current-feeding species use a scanning current to detect and capture immobile preys
like phytoplankton cells (Kiørboe, 2011a). Cruise-feeders are copepods that swim actively
through the water column in search of their prey. Lastly, particle-feeders are species known to
live on aggregates of sinking organic particles such as discarded appendicularians' houses.
Myelination (binary): myelinated copepod species have a lipid-rich myelin sheath around
their nerves that enables faster attack or evasive reactions (Lenz, 2012). Myelin sheaths play a
key role in modulating mortality and feeding rates and improve energy savings at low food
conditions.
Spawning mode (binary): eggs are either released into the open water after fertilization (free-
spawning) or are carried by females in egg sacs or in egg masses (sac-spawning). Sac
spawning copepods display lower fecundity rates and longer hatching times (Kiørboe &
Sabatini, 1994).

2.3. Definition of functional groups

Only those copepod species with a body size value and no missing data for three out of the

five traits were retained (n = 343). The species' trophic groups and feeding modes were re-

coded binarily to accurately represent species that belong to several feeding modes/ trophic

groups (Appendix S2). Thus, the analyses described below were performed on 343 species

based on ten functional trait dimensions.

A factor analysis on mixed data (FAMD; Pagès, 2004) that summarize the main modes of trait

combinations was used to estimate the functional distances between species. The use of a

FAMD is an improvement to the Multiple Correspondence Analysis used in Benedetti et al.

(2016) as it allowed to retain body size as a quantitative continuous variable. Four FAMD

components explaining 80.15% of the variance in species traits were retained (Audigier et al.,

2016). A Euclidean distance matrix was computed from the species' score along the four

FAMD components, and Ward's agglomeration link was used to generate a functional

dendrogram. The number FGs was determined by the height at which the dendrogram is cut.

We investigated the sensitivity of the resulting FGs to the main parameters of our clustering

approach (Appendix S4) and inspected each dendrogram to ensure the ecological relevance of

the final FGs (i.e., avoid few large groups that are functionally heterogenous or numerous

small groups that are functionally redundant). Ultimately, eleven FGs were identified (Table

S4.2).

2.4. Species distribution modelling

2.4.1. Environmental predictor selection

We considered 14 commonly used environmental drivers of the spatial ranges of marine

copepod species (see Appendix S5; Benedetti et al., 2021). The monthly climatologies of all

predictor variables were projected onto the 1°x1° cell grid of the World Ocean Atlas (WOA;

Boyer et al., 2013) and matched with the species occurrence data.

We used a two-stage procedure to select environmental predictors. First, we removed

collinear predictors (mean Spearman rank correlation coefficient > |0.7|) to avoid increasing

the uncertainty in regression model projections through coefficient inflation (Dormann et al.,

2013). This first step retained the following ten predictors: sea surface temperature (SST, °C),

photosynthetically active radiation (PAR, μmol m^{-2} s^{-1}), logged nitrate concentration (logNO$_3$,

μM), mixed layer depth (MLD, m), logged chlorophyll a concentration (logChla, mg m^{-3}),

logged eddy kinetic energy (logEKE, m^2 s^{-2}), surface wind speed (Wind, m s^{-1}), surface

carbon dioxide partial pressure (pCO$_2$, µatm), the excess of silicates to nitrates (Si* = [SiO$_2$] -

[NO$_3^-$], µM) and the excess of nitrate over phosphate (N* = [NO$_3$] - 16[PO$_4^{3-}$], µM).

Second, the number of environmental predictors included in the SDMs was restricted to five

to achieve a 10:1 ratio of occurrences to predictors (Guisan et al., 2017). For each species and

SDM, one of the ten predictors was randomly reshuffled prior to training a model (n = 30

repetitions). The correlation coefficients between the predictions of the original dataset and

those of the reshuffled datasets were used to estimate the relative ranking of each predictor

(Appendix S6). We used the top five predictors for each FG and SDM separately to train the

SDMs and project the species' habitat suitability (Appendix S6).

2.4.2. Background data, model configuration and habitat suitability projections

Pseudo-absence data were generated to use correlative SDMs following the target-group

approach of Phillips et al. (2009) which is appropriate to model plankton groups with sparse

data (Righetti et al., 2019; Benedetti et al., 2021). We chose to randomly draw pseudo-

absences from the total pool of sites (i.e., monthly 1° x 1° grid cells) with at least ten

occurrences across all species, except for FG2, FG4 and FG11 which showed sampling efforts

that were distinct from the other groups (Appendices S1 and S4). For species of FG2, FG4

and FG11, pseudo-absences were randomly drawn only from those sites where at least 10

different species of their FG were found. Ten times more background data than presences

were generated and presences were weighted ten times more than pseudo-absences (Barbet-

Massin et al., 2012).

We developed an ensemble modelling approach that combined three SDMs types which were

configured to avoid model overfitting (Merow et al., 2014; Thuiller et al., 2020): Generalized

Additive Models (GAMs), Generalized Linear Models (GLMs), and Artificial Neural

Networks (ANNs). The GLM were run using a quadratic formula and a logit link function.

The GAMs were set up to use a logit link function and a maximum of five smoothing terms.

ANNs were run using five cross validations with 200 iterations to optimize number of unities

in the hidden layer and the parameter for weight decay. The SDMs were trained on 80% of

the presence/pseudo-absence data chosen at random and tested on the remaining 20%. Ten

random cross evaluations were carried out and the true skill statistic (TSS) was calculated to

evaluate model performance. Only species displaying a mean TSS score >0.30 were used for

the final species and functional trait projections. The SDMs were then projected onto the

monthly environmental predictor conditions of the global ocean to obtain species-level maps

of habitat suitability indices (HSI). For each FG, annual mean HSI values were obtained by

averaging the monthly HSI maps across each species constituting the FG and SDMs.

2.5. Positioning FGs in environmental niche space

The mean univariate response curves emerging from GAMs were used to estimate the relative

optimal conditions (i.e., niche center) and tolerance (i.e., niche width) for each species and

predictor (Appendix S7; Benedetti, Vogt et al., 2018). The niche center was calculated as the

HSI-weighted median and the niche width as the difference between the weighted 10^{th} and

90^{th} percentiles. For this analysis, a set of eight predictors common to all species was defined

based on the overall ranking of predictors (i.e., unlike the top five FG-specific predictors used

for global SDMs projections; Appendix S6): SST, logChl, $logNO_3$, MLD, Si^*, PAR, N^* and

logEKE. We do this because a principal component analysis (PCA; Legendre & Legendre,

2012) was then performed on the niche centers and niche widths of all eight common

predictors to ordinate the species in an environmental "niche space". Non-parametric variance

analyses (Kruskal-Wallis tests and post-hoc Dunn's test) were carried out to test if FGs differ

in their position in niche space.

2.6. Community weighted median (CWM) proportions of traits

To explore the biogeography of all copepod traits, the monthly CWM trait values were

computed as the weighted sum of all HSI values belonging to species exhibiting that trait over

the sum of HSI values across all species in that community (Cormen et al., 2009). Like for

FGs, monthly CWM projections were performed for every SDM and mean annual CWM trait

values were derived based on the ensemble of projections (Appendices S8 and S9).

To explore the regional patterns of copepod traits expression, clustering was used to divide

the global ocean into regions with similar mean annual projections of: CWM body size,

trophic groups, feeding modes, myelination and spawning mode (Appendix S10). The spatial

patterns of the twelve CWM traits values were summarized through a PCA and the scores of

each grid cell along the first four components (97.9% of total variance, Appendix S10) were

used to calculate an Euclidean distance matrix. Various clustering approaches were explored

based on this distance matrix (Appendix S11). Ultimately, we chose to present the CWM trait

patterns obtained for k = 4 under Ward's linkage based on the profiles of cluster stability

metrics and because this method minimizes intra-cluster variance.

3. Results

3.1. Copepod FGs

The 343 copepod species retained were clustered based on their combinations of body size,

feeding mode, trophic group, myelination, and spawning mode. Eleven copepod FGs could be

derived from the functional dendrogram (Fig. 2; Appendix S4). The first dichotomy in the

dendrogram occurred between non carnivorous cruise feeders (FG1, 2, 3) and all other FGs.

FG1 was composed of 14 cruise-feeding herbivores of rather small sizes (median ± IQR =

1.10 ± 0.52 mm). Almost all species (13 out of 14) were sac-spawning, myelinated and

belonged to the *Clausocalanus* genera. FG2 was defined by small (0.80 ± 0.30 mm) cruise-

feeding detritivores. The 22 species of this group were all amyelinated, sac-spawning and

belonged to the Oncaeidae family. FG3 was composed of 46 detritivores spanning a broader

ranges of sizes (2.61 ± 1.70 mm) than the two previous groups. Most species in FG3 were

myelinated (74%) and free-spawning (80%). Cruise- and current-feeding were the two

predominant feeding modes, with *Spinocalanus*, *Metridia* and *Scaphocalanus* being the

dominant genera in this FG.

The second dichotomy in the dendrogram separated the small and large sac-spawning

carnivores (FG4 and 5) from the rest. FG4 consisted of 29 rather small (1.22 ± 0.70 mm)

carnivorous, sac-spawning ayelinated ambush-feeders part of the Corycaeidae family and thus

belonged to either the *Corycaeus*, *Farranula* or *Vettoria* genera. FG5 was made up of 29 of

larger (4.58 ± 2.75 mm) species of sac-spawning current- and cruise-feeding carnivores from

the Sapphirinidae and Euchaetidae families.

The third dichotomy of the dendrogram separated the large and medium-sized current feeders

(FG6 and 7) from the rest. FG6 was a group of 13 rather large (6.80 ± 2.15 mm) myelinated,

free-spawning species that were either current-feeding herbivores (85%) or fully omnivorous

(15%). The two most represented genera were *Calanus* and *Eucalanus*. FG7 was the largest

FG with 55 and current-feeding herbivores spanning a rather wide size range (1.80 ± 2.26

mm). Most were myelinated (95%) and free-spawning (89%). The main genera contributing

to the composition of FG7 were the smaller *Calanus* (e.g., *C. helgolandicus*, *C. finmarchicus*

or *C. pacificus*), and *Calocalanus* and *Paracalanus*.

The fourth dichotomy occurred between the small and medium-sized omnivores and the

medium free spawning carnivores of FG8. FG8 consisted of 49 predominantly amyelinated

(86%) and free-spawning (80%) medium-sized omnivorous-carnivorous species of various

size (3.40 ± 1.7 mm). This diverse group mixed current- (60%) and ambush-feeders (40%)

and gathered 12 genera with *Candacia*, *Haloptilus* and *Heterorhabdus* being the most

dominant. FG9 was composed of 40 free-spawning and mostly amyelinated (72%). current-

feeding omnivorous species showing a body sizes range of 2.76 ± 1.98 mm. *Pleuromamma*,

Gaetanus, and *Labidocera* were the main genera.

Finally, the fifth and last main dichotomy separated the mixed-feeding omnivores (FG11)

from the smaller ambush feeders (FG10). FG10 was rather homogeneous and contained 14

small (1.10 ± 0.52 mm) ambush-feeding, amyelinated and sac-spawning omnivores. All

species belonged to either the *Oithona* or the *Dioithona* genera from the Oithonidae family.

FG11 was a group of 32 amyelinated, free-spawning, current-ambush feeding species

predominantly belonging to the *Acartia* (omnivorous-herbivorous) and *Centropages*

(omnivorous) genera. All species in FG11 were rather small (1.50 ± 0.67 mm).

Figure 2: Functional dendrogram representing the inter-species traits dissimilarity for 343
copepod species. The hierarchical clustering was performed on a Euclidean distance matrix
issued from the species coordinates ensuing from a Factor Analysis of Mixed Data (FAMD).
Each leaf of the functional dendrogram represents a copepod species and the eleven
functional groups (FGs) are numbered and highlighted in color. The range (median ± IQR) of
the mean maximum body size of each FG is given in brackets.

The functional space defined by the PCs of the FAMD describes the reduced functional space

(Appendix S12). The first four PC of the FAMD explained 79.21% of the total variance in

functional traits. The largest functional distance along the first component was found between
FG4 and FG6.

3.2. FGs in environmental niche space

The univariate niche centers and widths of the copepod species modelled were used to
position FGs in niche space (Fig. 3; Appendix S13). The main niche characteristics that
contributed the most positively to PC1 (relative contribution to PC1 given in brackets) were:
logChl center (12.16%), logChl width (12.51%), EKE width (11.78%), MLD center (9.96%),
and MLD width (10.78%). The most important niche characteristics scoring PC2 were: Si*
center (30.88%), Si* width (27.90%) and SST center (11.17%). Species with positive PC1
scores were those affiliated with wider niches and conditions of higher concentrations of
nutrients and chlorophyll-a, stronger seasonal variations, and overall higher water column
turbulence and mixing.

Figure 3: Position of a) functional groups (FGs, larger circles) and species (smaller circles) in
environmental niche space according to the first two principal components (PCs) of a
principal component analysis (PCA) performed on b) the species-level niche characteristics
derived from GAM-based univariate response curves. The semi-transparent ellipses indicate
two times the value of the standard errors associated with the mean PC scores of the FGs.

Post-hoc variance analyses showed significant (Dunn's tests; $p < 0.05$) inter-FGs variations in
niche characteristics and PC scores (Appendix S14). The largest distance in niche space (PC1)
was found between FG4 and FG6 (Fig. 3a; $p = 6.5e\text{-}10$). Along PC1, FG4 also differed
significantly from FG1 ($p = 0.042$), FG3 ($p = 1.6e\text{-}06$), FG7 ($p = 1.3e\text{-}08$), FG8 ($p = 1.4e\text{-}4$),
FG9 ($p = 2.3e\text{-}06$), FG10 ($p = 0.011$) and FG11 ($p = 0.005$). Conversely, FG6 differed
significantly from FG2 ($p = 4.5e\text{-}4$), FG5 ($p = 0.001$) and FG8 ($p = 0.049$) along PC1. FG6

also differed significantly from FG2 (p = 0.001) and FG5 (p = 0.002). None of the FGs

showed significant variations along PC2. Only FG3 and FG7 showed significant variations

along PC3 (p = 4.2e-3; Appendices S13 and S14). To summarize, when FGs 4 and 6 were not

accounted for, none of the remaining nine FGs showed significant variations in PC1 and PC2

scores (all p > 0.05).

3.3. Mean annual habitat suitability indices (HSI) patterns

For all three SDMs and 11 FGs combined, we found SST to be the most important predictor,

followed by PAR, $logNO_3$, MLD, logChl, logEKE, Si^*, N^*, Wind, and pCO_2 (Appendix S6).

Thus, most FGs displayed mean annual HSI patterns that were driven by the latitudinal

temperature gradient at the first-order (Fig. 4).

Figure 4: Mean annual habitat suitability index (HSI) of the eleven copepod functional groups (FG)
for the global surface ocean (Mercator projection): a) FG1 (myelinated cruise-feeding omnivores-
herbivores), b) FG2 (amyelinated cruise-feeding detritivores), c) FG3 (myelinated mixed-feeding or
cuise-feeding detritivores), d) FG4 (amyelinated ambush-feeding carnivores), e) FG5 (current- or

cruise-feeding carnivores), f) FG6 (myelinated current-feeding omnivores-herbivores), g) FG7
(myelinated current-feeding omnivores-herbivores), h) FG8 (amyelinated ambush- or current-feeding
carnivores), i) FG9 (amyelinated current-feeding omnivores), j) FG10 (amyelinated ambush-feeding
omnivores), k) FG11 (amyelinated mixed-feeding omnivores) and for l) all species together. Mean
annual estimates were derived from the 12 monthly estimates of mean HSI obtained for three species
distribution models (SDMs).

The mean annual HSI of FG3 was found to decrease progressively from the poles towards the

equator. In contrast, the mean annual HSI of FG1, FG2, FG4, FG7, FG8, and FG10 were

found to be highest in the tropics and decreased towards higher latitudes. For those five FGs,

slighter decreases in HSI were modelled towards the tropical upwelling systems (e.g., Peru,

Benguela), the northern part of the Indian Ocean and the Pacific Equatorial counter current.

Therefore, these five FGs reached maximal HSI in the oligotrophic conditions of the tropical

gyres. Meanwhile, FG5, FG9, and FG11 show less marked latitudinal gradients in mean

annual HSI values, but they all displayed lower HSI in higher latitudes than in the tropics.

3.4. Functional trait biogeography from CMW trait values and regionalization

We identified four ocean regions that displayed significant contrasts in CWM trait proportions

and that overlap with major oceanographic features (Fig. 5; Appendix S15).

Figure 5: Partitioning of the global surface ocean into a) four regions according to the principal
components of a Principal Component Analysis (PCA) based on the CWM trait values (Mollweide
projection). Major oceanographic circulation features were overlayed to highlight their overlap
between the regions' boundaries. The distribution of CWM trait values between the four regions are
shown through the boxplots: b) median body size, CWM values of c) myelinated species, d) sac
spawning species, e) carnivores, f) herbivores, g) detritivores, h) omnivores, i) ambush-feeders, j)
current-feeders, k) cruise-feeders, l) current-ambush-feeders, and m) current-cruise-feeders. The
lower, middle, and upper boundaries of the boxplots correspond to the 25th, 50th, and 75th percentiles
respectively. The lower and upper whiskers extend no further than 1.5*IQR (interquartile range) from
the lower and upper hinges.

Region 1 fell within the equatorial band and comprised coastal upwelling regions and the

main oxygen minimum zones. The copepod communities of Region 1 displayed lower CWM

body size (median $\pm$ IQR = 2.043 mm $\pm$ 0.194), higher proportions of carnivores (0.342 $\pm$

0.040) and herbivores (0.335 ± 0.029) but lower proportions of omnivores (0.213 ±

and detritivores (0.098 ± 0.023). The CWM values of myelinated (0.446 ± 0.026), sac-

spawning (0.377 ± 0.026), and ambush-feeding (0.212 ± 0.036) were lower than in Region 2

and higher than in Regions 3 and 4 (Appendix S13).

Region 2 comprised the tropical gyres. We found that communities in Region 2 showed the

lowest CW median body size (2.002 ± 0.057 mm) but the highest CWM values of ambush-

feeding (0.259 ± 0.027) carnivorous (0.372 ± 0.028), and sac-spawning (0.440 ± 0.030)

copepods. Conversely, the CWM values of myelinated (0.413 ± 0.016) species in Region 2

were the lowest across all four regions.

Region 3 was located poleward to the previous two regions and covered the North Atlantic,

the North Pacific and the waters located between the Polar Front and the Antarctic

Circumpolar Current. The CWM body size (2.691 ± 0.342 mm), the CWM values of

myelinated (0.514 ± 0.028) and omnivorous (0.279 ± 0.021) species were higher there than in

Regions 1 and 2. In contrast, the CWM values of sac-spawners (0.368 ± 0.041), carnivores

(0.266 ± 0.026), and ambush feeders (0.132 ± 0.027) were lower.

Region 4 mainly corresponded to the Southern Ocean (i.e., grid cells south of the Antarctic

Polar Front, Fig. 5a). Region 4 displayed the highest CWM body size (3.035 ± 0.321 mm),

higher CWM values of myelinated (0.541 ± 0.018), omnivorous (0.278 ± 0.013), and

detritivorous (0.176 ± 0.030) species, and the lowest CWM values of sac-spawners (0.351 ±

0.035). The CWM values of current-feeding (0.522 ± 0.015) and cruise-feeding (0.218 ±

0.009) species were higher than those of ambush-feeding (0.120 ± 0.016).

4. Discussion

4.1. Towards meaningful global copepod FGs in marine ecology and ecosystem modelling

We identified eleven copepod FGs based on combinations of species-level functional traits.

Nine of these eleven FGs had also been found, or were nested within larger groups, in

previous studies based on regional species pools (Table 1), which suggests that most of the

functions performed by copepods on regional scales can also be found in the upper hundreds

of meters of the global ocean.

Table 1: Comparison of the copepod functional groups (FGs) defined in the present study and those
found in previous studies based on species and functional trait composition. The range (median ± IQR)
of the species body size is given in brackets for our FGs. List of acronyms describing the functional
traits: *Sma* = Small; *Lar* = Large; *Mye* = Myelinated; *Amye* = Amyelinated; *Sac* = Sac-spawning; *Cru*

This study	*Main clade*	Group composition	Pomerleau et al. (2015) - North Pacific Ocean	Benedetti et al. (2016) - Mediterranean Sea	Benedetti et al. (2018) - Mediterranean Sea	Becker et al. South Atlanti
FG1	*Clausocalanus*	Mye, Cru, Omni-Herb (1.10 ± 0.52)	Sma, Cur+Cru, Omni-Herb (Group 6)	Sma, Cru, Omni-Herb (subset of Group 6)	Sma, Cru, Omni-Herb (Group 7)	Sma+Lar, Cru+Cur, Om... (subset of Gr...
FG2	*Oncaeidae*	Amye, Cru, Detr (0.80 ± 0.30)	No equivalent	Sma, Cru, Omni-Detr (subset of Group 6)	Sma, Sac, Detr (subset of Group 5)	Sma, Amye, Cru, Carn o... (subset of Gr...
FG3	*Spinocalanus, Scaphocalanus, Metridia*	Mye, Mix or Cru, Detr (2.61 ± 1.70)	No equivalent	Sma, Cru, Omni-Detr (subset of Group 6)	Sma, Sac, Detr (subset of Group 5)	No equiv...
FG4	*Corycaeidae*	Amye, Amb, Carn (1.22 ± 0.70)	No equivalent	Sma, Amb, Carn (Group 2)	Sma, Amb, Carn (Group 2)	Sma, Amye, Cru, Carn o... (subset of Gr...
FG5	*Sapphirinidae, Euchaetidae*	Cur or Cru, Carn (4.58 ± 2.75)	No equivalent	Lar, Cru, Carn (Group 1)	Lar, Cru or Cur, Carn (Group 1)	Sma+Lar, My... Cur, Omni... (subset of Gr...
FG6	*Calanus, Eucalanus*	Mye, Cur, Omni-Herb (6.80 ± 2.15)	No equivalent	No equivalent	No equivalent	No equiv...
FG7	*Calanus, Paracalanus, Calocalanus*	Mye, Cur, Omni-Herb (1.80 ± 2.26)	Sma+Lar, Cur, Omni-Herb (Group 5c)	Sma+Lar, Cur, Omni-Herb (Group 4)	Sma+Lar, Cur, Omni-Herb (Groups 3 and 4)	Lar, Mye, Omni-Herb (s... Group ...
FG8	*Candacia, Haloptilus, Heterorhabdus*	Amy, Amb or Cur, Carn (3.40 ± 1.97)	No equivalent	Lar, Cru, Carn (Group 1)	Lar, Cru or Cur, Carn (Group 1)	No equiv...
FG9	*Pleuromamma, Gaetanus, Labidocera*	Amy, Cur, Omni (2.76 ± 1.98)	No equivalent	Sma+Lar, Cur, Omni-Herb (subset of Group 4)	Sma+Lar, Cur, Omni-Herb (subset of Group 4)	No equiv...
FG10	*Oithonidae*	Amy, Amb, Omni (1.10 ± 0.52)	No equivalent	Sma, Amb, Omni (Group 5)	Sma, Amb, Omni (Group 6)	Sma, Amye, Amb, Omni ... A)
FG11	*Acartia, Centropages*	Amy, Mix, Omni (1.50 ± 0.67)	Sma, Amb, Omni (subset of Group 4)	Sma, Mix, Omni (Group 3)	Sma, Mix, Omni (subset of Group 4)	Sma, Amye, Amb, Omni ... A)

Only the new FG3 and FG6 have no counterparts in previous studies. Their most suitable
habitats lie towards the poles (Fig. 4), whose copepod communities were not covered by the
regional studies mentioned. The copepods of FG3 (*Spinocalanus* spp., *Metridia* spp., or
Scaphocalanus spp.) are known to mainly inhabit deeper ocean layers where they feed on
sinking particulate organic matter and zooplankton carcasses (Yamaguchi et al., 2002; Sano et
al., 2013). Therefore, this group contributes to the remineralization of organic matter and

marine snow at higher latitudes and/or in colder conditions. FG6 comprises the largest (6.80 ±
2.15 mm) current-feeding omnivorous-herbivorous copepods from the Calanidae (e.g.,
Calanus hyperboreus, *C. glacialis*, *Eucalanus bungii* or *Rhincalanus gigas*). Many of these
large copepods are not present in the warm and oligotrophic conditions of the Mediterranean
Sea, hence their absence in Benedetti, Gasparini, & Ayata (2016) and Benedetti, Vogt et al.
(2018). FG6 is a key group for the biological carbon pump (Jónasdóttir et al., 2015; Visser et
al., 2017; Steinberg & Landry, 2017), as it represents large-bodied grazers that can actively
feed on microphytoplankton, perform relatively strong vertical migrations and generate large,
fast sinking pellets (Stamieszkin et al., 2015; Ohman & Romagnan 2016). This is confirmed
by the position of FG6 in niche space (Fig. 3) as it is affiliated with turbulent, seasonally
varying conditions with higher nutrient and chlorophyll-a concentrations. The species of FG7
fill similar functions but display smaller body sizes (1.80 ± 2.26mm; e.g., *C. helgolandicus*,
C. pacificus and the Paracalanidae), hence we expect their contribution to size-mediated
functions to be lower. The ecological roles and functions of the other nine FGs have already
been discussed in previous studies, and our results support the description of their ecological
roles. Here, we found a larger range of FGs compared to previous studies due to three main
reasons: (i) we investigated a global pool of species (hundreds of species instead of tens), (ii)
we accounted for myelination as an additional trait contrary to Benedetti, Gasparini, & Ayata
(2016) and Benedetti, Vogt et al. (2018), and (iii) we retained body size as a continuous trait
instead of using size classes.
Overall, the largest differences in niche space (Appendix S14) occurred between two main
sets of FGs: (i) myelinated free-spawning current-feeding herbivores of various sizes (FG3, 6,
7 and 9), and (ii) amyelinated sac-spawning small-bodied, or large-bodied, cruise- and
ambush-feeding detritivores and carnivores (FG2, 4 and 5). The first set is associated with
conditions of stronger mixing and higher nutrient and chlorophyll-a concentrations (Appendix
S14). The species in these FGs also have larger niche widths, i.e. they display broader
tolerances to variations in environmental conditions. Therefore, these FGs are found more
frequently in high latitude environments, or boundary current systems, where either
seasonality or physical mixing lead to higher mean annual productivity (Sarmiento & Gruber,
2006; Roy 2018). In contrast, the FGs of the second set display narrower niches and are
associated with conditions typical of the warmer tropical oligotrophic gyres (i.e., weaker
water mixing and lower nutrient and chlorophyll-a concentrations; Appendix S14).
Meanwhile, the FGs that are in the center of the niche space (FG1, 8, 10 and 11) could not be

associated to any particular environment, likely because they include genera with a wide

latitudinal distributions.

FGs can also combine species that are known to inhabit different vertical layers of the water

column (e.g., FG9 which gathers epipelagic species with meso-bathypelagic ones). This is

because depth preferences cannot be considered as a functional trait per se (Litchman et al.,

2013). Rather, the depth range of a species reflects the outcome of the interactions between its

traits and the abiotic and biotic conditions. Therefore, our approach likely overlooks vertical

habitat partitioning that occur between or within FGs.

The continuum of FGs in niche space was also found in functional trait space. Indeed, the FGs

of sets (i) and (ii) were often found on opposite sides of the FAMD dimensions (Appendix

S12). The global distribution of the FGs and CWM trait values further support the statements

above (Figs. 3 and 4): larger myelinated free-spawning and current- and cruise- feeding

copepods occur more frequently near the poles, whereas smaller amyelinated sac-spawning

and ambush-feeding copepods tend to occur in tropical oligotrophic gyres. Similar

continuums of copepod functional traits and niches were found at regional (Table 1) to global

scales (Brun et al., 2016).

In summary, our results support the view that planktonic copepods display a continuum of

functional traits with a strong latitudinal gradient (Figs. 4 and 5; Appendix S9), driven by

global gradients in abiotic conditions. Consequently, such a continuum should be explicitly

represented in marine ecosystem models (e.g. similar to Serra-Pompei et al., 2020), which too

often rely on a few size classes only (Le Quéré et al., 2005). These need to include FGs that

are both distinct, and important in terms of their biomass. To rank the present FGs based on

their contribution to total copepod community abundance in the upper ocean, we implemented

a preliminary synthesis of copepod abundance observations from various data sources

(Appendix S16). Examining the relative contribution of the eleven FGs to mean annual

community abundance in the upper 200m (Fig. S16b) showed that three FGs emerge as the

most abundant, regardless of latitude and sampling gears: FG10 (Oithonids; small ambush-

feeding omnivores), FG2 (Oncaeids; small cruise-feeding detritivores) and FG1

(*Clausocalanus* spp.; small cruise-feeding omnivores-herbivores). These FGs are thus likely

key the development of future ecosystem models. Future work is needed to better assess the

contribution of the copepod FGs to biomass and clarify their relative priorities for inclusion in

ecosystem models.

4.2. Why do copepod functional traits show diverging biogeographic patterns in the ocean?

The largest dissimilarity in community trait expression was found between regions 2 and 4
(Fig. 5b-m) and was driven by differences in CWM body size, myelination, carnivory and
ambush-feeding vs. current-feeding (Appendices S9, S10 and S15). We found the boundaries
of the four main regions to overlap with the trajectories of well-known boundary currents as
well as equatorial counter currents and fronts (Fig. 5). Trait patterns emerge from
environment-driven distribution models, so they reflect the fact that certain trait combinations
are more competitive than others under varying temperature and food conditions (Barton et
al., 2013; van Someren Gréve et al., 2017; McGinty et al., 2021).

Median copepod body size decreased from the poles to the equator with a slight increase in
upwelling systems. Such a pattern is primarily driven by the strong negative relationship
between the body size of marine ectotherms and temperature, according to Bergmann's rule
(Brun et al., 2016; McGinty et al., 2018; Evans et al., 2020; Brandão et al., 2021; Campbell et
al., 2021). The processes underlying Bergmann's rule remain debated, but it is likely that
warmer temperatures (or a factor confounded with temperature) decrease growth efficiency
and/or promote the maturation of adults at smaller body sizes (Atkinson 1994; Isla et al.,
2008).

Bergmann's rule occurs at the intraspecific and interspecific levels, meaning that warming-
induced decreases in median body size can emerge if clades of small species replace large-
bodied ones along a latitudinal gradient. Such a turnover in size classes has been observed as
well (Evans et al., 2020; Brandão et al., 2021) and is supported by our SDM projections.
Indeed, functional traits that represent shifts in clade composition such as myelination,
spawning strategy, carnivory or feeding modes display latitudinal gradients that are positively
or negatively correlated with the body size gradient (Appendices S9 and S10). The latitudinal
patterns of these functional traits likely result from changes in food availability, quality, and
predation pressure (Horne et al., 2016; van Someren Gréve et al., 2017; Roy, 2018).

Lipid-rich myelin sheaths enable a faster conduction of nerve responses. This promotes faster
reaction times and thus more efficient feeding or escape behaviors (Lenz, 2012). Myelin
sheaths are cholesterol-rich so they require larger metabolic investments of dietary lipids
(Lenz, 2012), which helps explain why copepod communities show larger proportion of small
amyelinated taxa in tropical gyres where smaller, lipid-poor phytoplankton dominate (Roy,
2018). Therefore, the proportion of myelinated species follows the same spatial pattern as
body size and current-feeding (Fig. 5b,c,j,m) and peaks in productive environments
characterized by larger and lipid-rich plankton (Roy 2018).

Similarly, sac-spawning is an energy-conservative spawning strategy, as it reduces egg-
mortality at the cost of fecundity and hatching speed. The increased proportions of sac-
spawners in tropical gyres (Fig. 5d) could reflect an adaptation to limited food availability
(Kiørboe & Sabatini, 1994; Barton et al., 2013) and higher rates of carnivory (Fig. 5e;
Woodd-Walker et al., 2002) and egg cannibalism among copepods (Ohman & Hirche, 2001;
Segers & Taborsky, 2011). Ambush-feeding is a passive feeding mode that lowers predation
risk and energy costs compared to current- and cruise-feeding, but at the expense of feeding
efficiency (Kiørboe, 2011a; van Someren Gréve et al., 2017). Consequently, the increased
proportion of ambush-feeders in region 2 (Fig. 5i) may result from trade-offs in functional
traits expression driven by abiotic and biotic filtering (Litchman et al., 2013), as oligotrophic
gyres seem to promote food-webs with increased carnivorous predation (Fig. 5e) and resource
competition (Woodd-Walker et al., 2002; Prowe, Visser, Andersen, Chiba, & Kiørboe, 2019).
Since our approach is based on presence data and habitat suitability indices rather than
abundances, it likely underestimates the contribution of very abundant ambush-feeding
species like *Oithona similis* at high latitudes (Pinkerton et al., 2010; Prowe et al., 2019;
Becker et al., 2021). As a result, we probably underestimate the proportion of ambush-feeders
in regions such as the Southern Ocean compared to Prowe et al. (2019), although these
authors discarded other ambush-feeding copepods such as the Corycaeidae (Benedetti et al.,
2016; Brun et al., 2017). These elements support the fact that our modelled copepods
functional traits patterns emerge from interactions between environmental conditions and the
relative fitness resulting from different trait combinations.
598
4.3. Caveats and conclusion
Our results are sensitive to: (i) the way the species were positioned in a functional space and
how the FGs were defined from the latter, (ii) the quantity and quality of the functional trait
data considered, and (iii) the SDMs chosen to generate the CWM traits values. We
investigated the sensitivity of our FGs to the clustering approach (Appendix S4). We assessed
how using this alternative dimension reduction analysis affected the quality of the functional
trait space by computing the Area Under the Curve (AUC) criterion (Mouillot et al., 2021).
We found an AUC value of 0.81, which is substantially higher than the recommended 0.7
threshold and indicates a "high quality trait space". Plus, all the functional dendrograms
emerging from alternative clustering approaches showed fairly similar structure (mean
Baker's Gamma correlation coefficient was 0.75 ± 0.12), indicating that they all lead to
similar FGs. The traits chosen for this study only cover a fraction of the traits mentioned in

the literature (Litchman et al., 2013; Appendix S3). Therefore, our study likely underestimates

the true diversity of copepod functions in the ocean. More field observations and lab

experiments are required to document the existing traits of many copepod species (Barton et

al., 2013).

We found some regional differences between the mean annual HSI projections of the SDMs.

The GLM estimated higher mean HSI than the two other model types in the Southern Ocean

(Appendix S8). Such variability stems from how the various SDMs cope with limited

predictors and species data availability in winter conditions at the very high latitudes. Here,

the less complex response curves of GLMs (Merow et al., 2014) lead to higher average HSI in

very cold temperatures compared to GAMs. The HSI patterns obtained from the GAMs and

ANNs were closer to the latitudinal diversity gradients that were previously observed for

marine ectotherms (Tittensor et al., 2010; Benedetti et al., 2021). Interestingly, inter-SDMs

variability was much lower when looking at CWM traits projections rather than HSI patterns

(Appendix S8). The fact that CWM body size and CWM myelination show very similar

spatial patterns and ranges to the estimates of Brun et al. (2016) gives us further confidence in

our spatial projections.

To conclude, we recommend the inclusion of multiple FGs of copepods in marine ecosystem

models, based on our functional dendrogram (Fig. 2), with the level of complexity of copepod

representation reflecting not only the FGs dominating community biomass at the scale of the

study region(s), but also the traits of interest. Ideally, future global marine ecosystems, that

cannot efficiently include 11 copepod groups, will include a few (3-4) FGs that cover the

main gradients observed in trait space (Fig. 2; Appendix S12), niche space (Fig. 3),

geographical space (Figs. 4, 5; Appendices S9, S10) and that represent the majority of global

or regional biomass. For other ecological applications or to represent food-web interactions

across trophic levels (Serra-Pompei et al., 2020), these groups could be split into further FGs

with specific characteristics (e.g., size classes, high lipid content, grazing and mortality rates

etc.). Future studies are required to improve the coverage of functional trait data to include

other taxonomic groups and more quantitative traits (Appendix S3). Ongoing compilations of

zooplankton biomass data will also help rank the regional to global importance of the present

FGs, and to assess links between ecosystem function and service provision.

Data Availability Statement

The copepod species occurrences data used to train the species distribution models are

publicly available on Zenodo (https://doi.org/10.5281/zenodo.5101349). The species

functional traits table is available as Supplementary Appendix S2. All R codes are accessible
on the GitHub account of J.W. (https://github.com/jonas-wydler) and are also available on
Zenodo (https://doi.org/10.5281/zenodo.7050567).

References

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015).
spthin: an R package for spatial thinning of species occurrence records for use in ecological niche
models. *Ecography*, 38(5), 541–545.

Atkinson, D. (1994). Temperature and Organism Size—A Biological Law for Ectotherms? In
M. Begon & A. H. Fitter (Eds.), *Advances in Ecological Research* (Vol. 25, pp. 1-58): Academic
Press.

Audigier, V., Husson, F., & Josse, J. (2016). A principal component method to impute missing
values for mixed data. *Advances in Data Analysis and Classification*, 10(1), 5–26.

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences
for species distribution models: how, where and how many? *Methods in Ecology and Evolution, 3*(2),
327-338.

Barton, A. D., Pershing, A. J., Litchman, E., Record, N. R., Edwards, K. F., Finkel, Z. V., . . .
Ward, B. A. (2013). The biogeography of marine plankton traits. *Ecology Letters*, 16(4), 522–534.
doi: 10.1111/ele.12063

Beaugrand, G., Edwards, M., & Legendre, L. (2010). Marine biodiversity, ecosystem
functioning, and carbon cycles. *Proceedings of the National Academy of Sciences*, 107(22), 10120-
10124. doi:10.1073/pnas.0913855107

Becker, É. C., Mazzocchi, M. G., de Macedo-Soares, L. C. P., Costa Brandão, M., &
Santarosa Freire, A. (2021). Latitudinal gradient of copepod functional diversity in the South Atlantic
Ocean. *Progress in Oceanography*, 199, 102710. doi:https://doi.org/10.1016/j.pocean.2021.102710

Benedetti, F., Gasparini, S., & Ayata, S.-D. (2016). Identifying copepod functional groups
from species functional traits. *Journal of Plankton Research*, 38(1), 159-166.
doi:10.1093/plankt/fbv096

Benedetti, F., Vogt, M., Righetti, D., Guilhaumon, F., & Ayata, S.-D. (2018). Do functional
groups of planktonic copepods differ in their ecological niches? *Journal of Biogeography*, 45(3), 604–
616. doi: 10.1111/jbi.13166

Benedetti, F., Vogt, M., Elizondo, U. H., Righetti, D., Zimmermann, N. E., & Gruber, N.
(2021). Major restructuring of marine plankton assemblages under global warming. *Nature
Communications*, 12(1), 5226. doi:10.1038/s41467-021-25385-x

Boyer, T.P., J. I. Antonov, O. K. Baranova, C. Coleman, H. E. Garcia, A. Grodsky, D. R.
Johnson, R. A. Locarnini, A. V. Mishonov, T.D. O'Brien, C.R. Paver, J.R. Reagan, D. Seidov, I. V.

Smolyar, and M. M. Zweng, 2013: World Ocean Database 2013, NOAA Atlas NESDIS 72, S.
Levitus, Ed., A. Mishonov, Technical Ed.; Silver Spring, MD, 209 pp.,
http://doi.org/10.7289/V5NZ85MT

Brun, P., Payne, M. R., & Kiørboe, T. (2016). Trait biogeography of marine copepods–an
analysis across scales. *Ecology Letters*, 19(12), 1403–1413. doi: 10.1111/ele.12688

Brun, P., Payne, M. R., & Kiørboe, T. (2017). A trait database for marine copepods. *Earth*
System Science Data, 9(1), 99–113. doi:10.5194/essd-2016-30

Campbell, M. D., Schoeman, D. S., Venables, W., Abu-Alhaija, R., Batten, S. D., Chiba, S., . .
. Richardson, A. J. (2021). Testing Bergmann's rule in marine copepods. *Ecography*, 44(9), 1283-
1295. doi:https://doi.org/10.1111/ecog.05545

Cormen, T. H., Leiserson, C. E., Rivest, R. L., & Stein, C. (2009). Introduction to algorithms.
MIT and McGraw.

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., . . . others (2013).
Collinearity: a review of methods to deal with it and a simulation study evaluating their performance.
Ecography, 36(1), 27–46. doi: 10.1111/j.1600-0587.2012.07348.x

Evans, L. E., Hirst, A. G., Kratina, P., & Beaugrand, G. (2020). Temperature-mediated
changes in zooplankton body size: large scale temporal and spatial analysis. *Ecography, 43*(4), 581-
590. doi:https://doi.org/10.1111/ecog.04631

Flynn, K. J., St John, M., Raven, J. A., Skibinski, D. O. F., Allen, J. I., Mitra, A., & Hofmann,
E. E. (2015). Acclimation, adaptation, traits and trade-offs in plankton functional type models:
reconciling terminology for biology and modelling. Journal of Plankton Research, 37(4), 683-691.
doi:10.1093/plankt/fbv036

Guisan, A., Thuiller, W., & Zimmermann, N. E. (2017). Habitat suitability and distribution
models: with applications in R. Cambridge University Press.

Hansen, B., Bjornsen, P. K., & Hansen, P. J. (1994). The size ratio between planktonic
predators and their prey. *Limnolony & Oceanography*, 39(2), 395–403.

Hébert, M.-P., Beisner, B. E., & Maranger, R. (2017). Linking zooplankton communities to
ecosystem functioning: toward an effect-trait framework. *Journal of Plankton Research*, 39(1), 3–12.
doi:10.1093/plankt/fbw068

Hijmans, R. J. (2012). Cross-validation of species distribution models: removing spatial
sorting bias and calibration with a null model. *Ecology*, 93(3), 679-688.
doi:https://doi.org/10.1890/11-0826.1

Hofmann Elizondo, U., Righetti, D., Benedetti, F., & Vogt, M. (2021). Biome partitioning of
the global ocean based on phytoplankton biogeography. *Progress in Oceanography, 194*, 102530.
doi:https://doi.org/10.1016/j.pocean.2021.102530

Horne, C. R., Hirst, A. G., Atkinson, D., Neves, A., & Kiørboe, T. (2016). A global synthesis
of seasonal temperature– size responses in copepods. *Global Ecology and Biogeography*, 25(8), 988–
999. doi: 10.1111/geb.12460

Huys, R. & Boxshall, G.A. (1991). *Copepod Evolution*. London, UK: Ray Society
(Publications), 159. ISBN 0-903-87421-0. 468 pp.

Isla, J. A., Lengfellner, K., & Sommer, U. (2008). Physiological response of the copepod
Pseudocalanus sp. in the Baltic Sea at different thermal scenarios. *Global Change Biology*, 14(4),
895-906. doi:https://doi.org/10.1111/j.1365-2486.2008.01531.x

Jónasdóttir, S. H., Visser, A. W., Richardson, K., & Heath, M. R. (2015). Seasonal copepod
lipid pump promotes carbon sequestration in the deep North Atlantic. *Proceedings of the National
Academy of Sciences, 112*(39), 12122-12126. doi:10.1073/pnas.1512110112

Kenitz, K. M., Visser, A. W., Mariani, P., & Andersen, K. H. (2017). Seasonal succession in
zooplankton feeding traits reveals trophic trait coupling. *Limnology & Oceanography*, 62(3), 1184–
1197.

Kiørboe, T. (2011a). What makes pelagic copepods so successful? *Journal of Plankton
Research*, 33(5), 677– 685. doi:10.1093/plankt/fbq159

Kiørboe, T. (2011b). How zooplankton feed: mechanisms, traits and trade-offs. *Biological
Reviews*, 86(2), 311– 339. doi: 10.1111/j.1469-185X.2010.00148.x

Kiørboe, T., & Hirst, A. G. (2014). Shifts in mass scaling of respiration, feeding, and growth
rates across life-form transitions in marine pelagic organisms. *The American Naturalist*, 183(4), E118–
E130.

Kiørboe, T., & Sabatini, M. (1994). Reproductive and life cycle strategies in egg-carrying
cyclopoid and free-spawning calanoid copepods. *Journal of Plankton Research*, 16(10), 1353–1366.

Legendre, P., & Legendre, L. (2012). Numerical Ecology. Elsevier.

Le Quéré, C., Harrison, S. P., Colin Prentice, I., Buitenhuis, E. T., Aumont, O., Bopp, L., . . .
others (2005). Ecosystem dynamics based on plankton functional types for global ocean
biogeochemistry models. *Global Change Biology*, 11(11), 2016–2040. doi: 10.1111/j.1365-
2486.2005.01004.x

Le Quéré, C., Buitenhuis, E. T., Moriarty, R., Alvain, S., Aumont, O., Bopp, L., . . . Vallina,
S. M. (2016). Role of zooplankton dynamics for Southern Ocean phytoplankton biomass and global
biogeochemical cycles. *Biogeosciences*, 13(14), 4111-4133. doi:10.5194/bg-13-4111-2016

Lenz, P. H. (2012). The biogeography and ecology of myelin in marine copepods. *Journal of
Plankton Research*, 34(7), 575–589. doi:10.1093/plankt/fbs037

Litchman, E., Ohman, M. D., & Kiørboe, T. (2013). Trait-based approaches to zooplankton
communities. *Journal of Plankton Research*, 35(3), 473–484. doi:10.1093/plankt/fbt019

Longhurst, A. R. (2010). Ecological geography of the sea. Elsevier.

McGinty, N., Barton, A. D., Record, N. R., Finkel, Z. V., & Irwin, A. J. (2018). Traits
structure copepod niches in the North Atlantic and Southern Ocean. *Marine Ecology Progress Series*,
601, 109-126. https://doi.org/10.3354/meps12660

McGinty, N., Barton, A. D., Record, N. R., Finkel, Z. V., Johns, D. G., Stock, C. A., & Irwin,
A. J. (2021). Anthropogenic climate change impacts on copepod trait biogeography. *Global Change*
Biology, 27(7), 1431-1442. doi:https://doi.org/10.1111/gcb.15499

Merow, C., Smith, M. J., Edwards Jr, T. C., Guisan, A., McMahon, S. M., Normand, S., . . .
Elith, J. (2014). What do we gain from simplicity versus complexity in species distribution models?
Ecography, 37(12), 1267–1281. doi: 10.1111/ecog.00845

Mouillot, D., Loiseau, N., Grenié, M., Algar, A. C., Allegra, M., Cadotte, M. W., . . . Auber,
A. (2021). The dimensionality and structure of species trait spaces. *Ecology Letters*, 24(9), 1988-2009.
doi:https://doi.org/10.1111/ele.13778

Ohman, M. D., & Hirche, H. J. (2001). Density-dependent mortality in an oceanic copepod
population. *Nature, 412*(6847), 638-641. doi:10.1038/35088068

Ohman, M. D., & Romagnan, J.-B. (2016). Nonlinear effects of body size and optical
attenuation on diel vertical migration by zooplankton. *Limnology & Oceanography*, 61(2), 765–770.
doi: 10.1002/lno.10251

Pagès, J. (2004). Analyse factorielle de données mixtes: principe et exemple d'application.
Montpellier SupAgro, http://www. agro-montpellier. fr/sfds/CD/textes/pages1.pdf

Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J., & Ferrier, S.
(2009). Sample selection bias and presence-only distribution models: implications for background and
pseudo-absence data. *Ecological Applications*, 19(1), 181–197.

Pinkerton, M. H., Smith, A. N. H., Raymond, B., Hosie, G. W., Sharp, B., Leathwick, J. R., &
Bradford-Grieve, J. M. (2010). Spatial and seasonal distribution of adult Oithona similis in the
Southern Ocean: Predictions using boosted regression trees. *Deep Sea Research Part I:*
Oceanographic Research Papers, 57(4), 469-485. doi:https://doi.org/10.1016/j.dsr.2009.12.010

Pomerleau, C., Sastri, A. R., & Beisner, B. E. (2015). Evaluation of functional trait diversity
for marine zooplankton communities in the northeast subarctic pacific ocean. *Journal of Plankton*
Research, 37(4), 712–726. doi:10.1093/plankt/fbv045

Prowe, A. E. F., Pahlow, M., Dutkiewicz, S., Follows, M., & Oschlies, A. (2012). Top-down
control of marine phytoplankton diversity in a global ecosystem model. *Progress in Oceanography,*
101(1), 1-13. doi:https://doi.org/10.1016/j.pocean.2011.11.016

Prowe, A. E. F., Visser, A. W., Andersen, K. H., Chiba, S., & Kiørboe, T. (2019).
Biogeography of zooplankton feeding strategy. *Limnology & Oceanography*, 64(2), 661–678. doi:
10.1002/lno.11067

Razouls C., Desreumaux N., Kouwenberg J., & de Bovée F. (2005-2022). Biodiversity of
Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne
University, CNRS. Available at http://copepodes.obs-banyuls.fr/en

Reygondeau, G., Guieu, C., Benedetti, F., Irisson, J.-O., Ayata, S.-D., Gasparini, S., &
Koubbi, P. (2017). Biogeochemical regions of the Mediterranean Sea: An objective multidimensional
and multivariate environmental approach. *Progress in Oceanography*, 151, 138-148.
doi:https://doi.org/10.1016/j.pocean.2016.11.001

Ricotta, C. (2005). A note on functional diversity measures. *Basic and Applied Ecology*, 6(5),
479–486.

Righetti, D., Vogt, M., Gruber, N., Psomas, A., & Zimmermann, N. E. (2019). Global pattern
of phytoplankton diversity driven by temperature and environmental variability. *Science Advances*,
5(5), eaau6253. doi:10.1126/sciadv.aau6253

Roy, S. (2018). Distributions of phytoplankton carbohydrate, protein and lipid in the world
oceans from satellite ocean colour. *The ISME Journal*, 12(6), 1457-1472. doi:10.1038/s41396-018-
0054-8

Sailley, S., Vogt, M., Doney, S., Aita, M., Bopp, L., Buitenhuis, E., ... Yamanaka, Y. (2013).
Comparing food web structures and dynamics across a suite of global marine ecosystem models.
Ecological Modelling, 261, 43–57.

Sano, M., Maki, K., Nishibe, Y., Nagata, T., & Nishida, S. (2013). Feeding habits of
mesopelagic copepods in Sagami Bay: Insights from integrative analysis. *Progress in Oceanography,*
110, 11-26. doi:https://doi.org/10.1016/j.pocean.2013.01.004

Sarmiento, J. L. & Gruber, N. (2006), *Ocean Biogeochemical Dynamics*, Princeton Univ.
Press, Princeton, N. J.

Segers, F. H. I. D., & Taborsky, B. (2011). Egg size and food abundance interactively affect
juvenile growth and behaviour. *Functional Ecology*, 25(1), 166-176.
doi:https://doi.org/10.1111/j.1365-2435.2010.01790.x

Serra-Pompei, C., Soudijn, F., Visser, A. W., Kiørboe, T., & Andersen, K. H. (2020). A
general size- and trait-based model of plankton communities. *Progress in Oceanography*, 189,
102473. doi:https://doi.org/10.1016/j.pocean.2020.102473

Stamieszkin, K., Pershing, A. J., Record, N. R., Pilskaln, C. H., Dam, H. G., & Feinberg, L. R.
(2015). Size as the master trait in modeled copepod fecal pellet carbon flux. *Limnology &*
Oceanography, 60(6), 2090–2107. doi: 10.1002/lno.10156

Steinberg, D. K., & Landry, M. R. (2017). Zooplankton and the ocean carbon cycle. *Annual*
Review of Marine Science, 9, 413–444. doi: 10.1146/annurev-marine-010814-015924

Stocker, T. (2014). Climate change 2013: the physical science basis: Working group i
contribution to the fifth assessment report of the intergovernmental panel on climate change.
Cambridge University Press.

Thuiller, W., Georges, D., Engler, R., Georges, M. D., & Breiner F. (2020). biomod2:
Ensemble Platform for Species Distribution Modeling. R package version 3.4.6. https://CRAN.R-
project.org/package=biomod2

Tittensor, D. P., Mora, C., Jetz, W., Lotze, H. K., Ricard, D., Berghe, E. V., & Worm, B.
(2010). Global patterns and predictors of marine biodiversity across taxa. *Nature*, 466(7310), 1098-
1101. doi:10.1038/nature09329

Turner, J. T. (2015). Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's
biological   pump.   *Progress   in   Oceanography*,   130,   205–248.
http://dx.doi.org/10.1016/j.pocean.2014.08.005

Vallina, S. M., Follows, M. J., Dutkiewicz, S., Montoya, J. M., Cermeno, P., & Loreau, M.
(2014). Global relationship between phytoplankton diversity and productivity in the ocean. *Nature*
Communications, 5(1), 4299. doi:10.1038/ncomms5299

van Someren Gréve, H., Almeda, R., & Kiørboe, T. (2017). Motile behavior and predation risk
in planktonic copepods. *Limnology & Oceanography*, 62(5), 1810– 1824. doi: 10.1002/lno.10535

Veloz, S. D. (2009). Spatially autocorrelated sampling falsely inflates measures of accuracy
for presence-only niche models. *Journal of Biogeography*, 36(12), 2290–2299.

Villéger, S., Mason, N. W., & Mouillot, D. (2008). New multidimensional functional diversity
indices for a multifaceted framework in functional ecology. *Ecology*, 89(8), 2290-2301.

Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E.
(2007). Let the concept of trait be functional! *Oikos*, 116(5), 882–892. doi: 10.1111/j.2007.0030-
1299.15559.x

Visser, A. W., Grønning, J., & Jónasdóttir, S. H. (2017). *Calanus hyperboreus* and the lipid
pump. *Limnology & Oceanography*, 62(3), 1155-1165. doi:10.1002/lno.10492

Woodd-Walker, R. S., Ward, P., & Clarke, A. (2002). Large-scale patterns in diversity and
community structure of surface water copepods from the Atlantic ocean. *Marine Ecology Progress*
Series, 236, 189–203.

Yamaguchi, A., Watanabe, Y., Ishida, H., Harimoto, T., Furusawa, K., Suzuki, S., . . . Mac
Takahashi, M. (2002). Community and trophic structures of pelagic copepods down to greater depths
in the western subarctic Pacific (WEST-COSMIC). *Deep Sea Research Part I: Oceanographic*
Research Papers, 49(6), 1007-1025.

Biosketch

Fabio Benedetti is a postdoctoral researcher and **Meike Vogt** a senior research scientist in
the Environmental Physics (UP) group of ETH Zürich. Both share broad interests in plankton
biogeography and functional diversity and theirs links with biodiversity, ecosystem function
and biogeochemical cycles in the global ocean. F.B. is a macroecologist specialized in trait-