Cryo-electron tomography of Birbeck granules reveals the

molecular mechanism of langerin lattice formation

Authors: Toshiyuki Oda[1*], Haruaki Yanagisawa[2], Hideyuki Shinmori [3], Youichi Ogawa[4],

and Tatsuyoshi Kawamura[4]

Affiliations: [1] Department of Anatomy and Structural Biology, Graduate School of

Medicine, University of Yamanashi, 1110 Shimokato, Chuo, Yamanashi, 409-3898, Japan. [2]

Department of Cell Biology and Anatomy, Graduate School of Medicine, the University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.

[3] Faculty of Life and Environmental Science, University of Yamanashi, 4-4-37 Takeda,

Kofu, Yamanashi, 400-8510, Japan

[4] Department of Dermatology, Faculty of Medicine, University of Yamanashi, 1110

Shimokato, Chuo, Yamanashi, 409-3898, Japan.

*Correspondence to: toda@yamanashi.ac.jp

Abstract

Langerhans cells are specialized antigen-presenting cells localized within the epidermis and

mucosal epithelium. Upon contact with Langerhans cells, pathogens are captured by the C-

type lectin langerin and internalized into a structurally unique vesicle known as a Birbeck

granule. Although the immunological role of Langerhans cells and Birbeck granules have

been extensively studied, the mechanism by which the characteristic zippered membrane

structure of Birbeck granules is formed remains elusive. In this study, we observed isolated

Birbeck granules using cryo-electron tomography and reconstructed the 3D structure of the

repeating unit of the honeycomb lattice of langerin at 6.4 Å resolution. We found that the

interaction between the two langerin trimers was mediated by docking the flexible loop at

residues 258-263 into the secondary carbohydrate-binding cleft. Mutations within the loop

inhibited Birbeck granule formation and the internalization of HIV pseudovirus. These

findings suggest a molecular mechanism for membrane zippering during Birbeck granule

biogenesis and provide insight into the role of langerin in the defense against viral infection.

Introduction

The epidermis and mucosa provide the first line of defense against pathogens. In these tissues,

specialized antigen-presenting cells called Langerhans cells capture pathogens by

recognizing their surface mannose-containing oligosaccharides via the C-type lectin langerin

(Valladeau et al., 2003; Doebel et al., 2017). Pathogens internalized via langerin are enclosed

in a specialized organelle called a Birbeck granule, which has a central striated lamella

between closely apposed membranes (Birbeck et al., 1961; Valladeau et al., 2000;

McDermott et al., 2004; Valladeau et al., 1999). Once pathogens are captured, Langerhans

cells undergo maturation and migrate to peripheral lymph nodes, where they present

processed antigens to T cells (Merad et al., 2008; de_Jong MA, 2010; Rhodes et al., 2021).

Langerin is a type II transmembrane protein that is composed of a cytoplasmic tail, coiled-

coil neck region, and carbohydrate recognition domain (CRD) (Stambach and Taylor, 2003;

Valladeau et al., 1999; Valladeau et al., 2000) (Figure 1A). Through the coiled-coil neck

region, langerin forms a trimer, which is functionally important for oligosaccharide binding

(Stambach and Taylor, 2003; Weis and Drickamer, 1996; Weis and Drickamer, 1994). The

calcium-dependent carbohydrate-binding capacity of langerin CRD is essential for the

formation of Birbeck granules because the removal of calcium ions causes them to unzip and

disintegrate (Stambach and Taylor, 2003; Bartosik, 1992). It has been proposed that the "face-

to-face" interaction between the langerin trimers via CRDs confers the characteristic zippered

membranes with a central striation of Birbeck granules (Thepaut et al., 2009) (Figure 1B and

1C). However, the precise molecular mechanism of Birbeck granule formation is unknown

because of the lack of a high-resolution structure.

Langerin's high affinity for mannose and oligo-mannose allows Langerhans cells to capture

a wide range of pathogens, including *Saccharomyces cerevisiae, Candida albicans,*

Mycobacterium leprae, herpes simplex virus, and human immunodeficiency virus-1 (HIV-1)

(Tateno et al., 2010; Hunger et al., 2004; de Jong et al., 2010; de Witte et al., 2007; Turville et al.,

2002). Although Langerhans cells in the mucosa are the initial target of HIV-1 in sexually

transmitted infections, the efficiency of Langerhans cell infection is low, requiring high viral

loads of HIV-1 to infect cells (Kawamura et al., 2001; Kawamura et al., 2000; Ogawa et al., 2013).

The resistance of Langerhans cells to HIV-1 infection is mediated by the efficient internalization

of HIV-1 into Birbeck granules and subsequent autophagic degradation (de Witte et al., 2007;

Ribeiro et al., 2016). Therefore, elucidating the mechanism by which HIV-1 is internalized into

Birbeck granules is essential to understanding the role of Langerhans cells as a barrier to HIV-1

infection.

In the present study, we reconstructed the 3D structure of isolated Birbeck granules using cryo-

electron tomography and identified a flexible loop region that is crucial for Birbeck granule

formation and virus internalization.

Results

Cryo-electron microscopy of isolated Birbeck granules

Exogenous expression of langerin in mammalian cell lines readily reproduces the formation

of Birbeck granules (McDermott et al., 2004). We found that the addition of yeast mannan to

langerin-expressing cells enhances the formation of large Birbeck granules (Figure 1C). To

purify these Birbeck granules, we expressed SNAP-tagged human langerin in 293T cells,

biotinylated the tag by incubation with benzylguanine-biotin, and immunoprecipitated the

Birbeck granules using streptavidin agarose (Figure 1D and S1A). Under cryo-electron

microscopy, the isolated Birbeck granules were observed to be thin lamellar vesicles of

assorted sizes, similar to those reported previously (Figure 1E, 1F, and S1B) (Sagebiel and

Reed, 1968; Lenormand et al., 2013). Unlike regular 2D crystals of proteins, Birbeck

granules appear wavy, suggesting structural heterogeneity. The 2D classification of the

segmented images extracted from the micrographs showed nearly three-fold symmetrical

arrangements of pores with a diameter of approximately 10 nm; however, no clear crystalline

lattice was observed (Figure 1G). Although most Birbeck granules were oriented in

approximately the same direction with respect to the beam axis, some granules were twisted,

providing different views that were sufficient for 3D reconstruction (Figure S1C).

Subtomogram averaging of the repeating unit of Birbeck granules

We conducted cryo-electron tomography of the isolated Birbeck granules and reconstructed

the repeating units by averaging the subtomograms (Figure 2 and S2A). The initial density

map (Figure 2A) shows a langerin trimer (blue) with three inverted trimers (red) attached to

the CRDs. However, because of the flexibility between each langerin trimer, the resolution

was limited to ~24 Å. To improve the resolution, we generated a mask enclosing the central

langerin trimer and three attached CRDs and performed focused refinement and 3D

classification (Figure 2B). The resolution was improved to 6.4 Å (Figure S2B), which clearly

visualized the alpha helices and enabled us to generate a model based on the previously

reported crystal structure of the langerin trimer (Feinberg et al., 2010). We then performed a

two-body refinement to visualize the spatial relationship between the two opposing langerin

trimers (Nakane et al., 2018). Based on this refinement (Figure 2C), we built a honeycomb

lattice model (Figure 2D and Movie S1), which agrees with the striation of the Birbeck

granules observed by ultra-thin section electron microscopy and the ~10 nm pores visualized

in the 2D classification images (Figure 1C and 1G) (Birbeck et al., 1961). However, in reality,

the honeycomb lattice is non-uniform because of the structural heterogeneity.

Interaction between the two langerin trimers via flexible loops

An in-depth analysis of the interaction region between the two CRDs (Figure 3) revealed that

the interaction is mediated by a loop at residues 258-263, which is highly flexible in the

crystal structure (Feinberg et al., 2010). The interaction between the two CRDs appears to be

maintained by docking the loops of two langerin molecules into each other's lateral cleft,

which constitutes the secondary carbohydrate-binding site (Figure 3A, pink circles)

(Chatwell et al., 2008). The interaction between the two CRDs is flexible because the first

eigenvector of the multi-body analysis represents an approximately 30° tilting motion with

the loop as the fulcrum (Figure 3B, and Movie S2). Interestingly, loop 258-263 of the inverted

molecule appears to participate in the secondary carbohydrate-binding site of the upright

molecule (Figure 3A). A previous study found that the secondary calcium-independent

binding site has a lower affinity for carbohydrate ligands than the primary calcium-dependent

binding site (Chatwell et al., 2008). The docked loop 258-263 may provide additional

interactions with the hydroxyl group of mannose via Glu261 and Asp263, thus increasing the

affinity of the secondary binding site when two langerin trimers form a complex. Furthermore,

the Trp264Arg mutation is associated with a lack of Birbeck granules (Verdijk et al., 2005;

Figdor et al., 2002). Trp264 may also be involved in docking the loop into the cleft; however,

it is more likely that disruption of the hydrophobic core of the CRD is responsible for this

defect in Birbeck granule formation (Chatwell et al., 2008).

Loop 258-263 is essential for Birbeck granule formation

To investigate the importance of loop 258-263 in the biogenesis of Birbeck granules, we

mutated three residues in the loop: Met260Ala, Glu261Arg, and Asp263Lys, and generated

three mutated langerin constructs, each carrying one (MRGD), two (MRGK), or three

mutations (ARGK). We expressed these three langerin mutants in 293T cells and induced

Birbeck granule formation by adding yeast mannan (Figure 4A). Under ultra-thin section

microscopy, wild-type langerin produced Birbeck granules a few micrometers in length in

the presence of mannan (Figure 4A, MEGD WT). As the number of mutations increased, the

length of Birbeck granules decreased. With two point mutations (MRGK), some cross-

sections became spherical rather than rod-shaped, suggesting that the Birbeck granules failed

to extend. With three mutations (ARGK), clusters of Birbeck granules were not observed,

with isolated individual granules observed instead. The length of individual Birbeck granules

and the total length of granules per area were measured to confirm the above observations

(Figure 4B and 4C). In contrast, the number of the Birbeck granules showed less significant

differences because numerous short fragments were present in the mutant specimen (Figure

4D). These results demonstrate the importance of loop 258-263 in the biogenesis of Birbeck

granules.

Loop 258-263 is crucial for the internalization of HIV pseudovirus

To investigate the impact of the mutations in loop 258-263 on virus internalization by Birbeck

granules, we added lentiviruses pseudo-typed with the HIV-1 env protein to langerin-

expressing 293T cells. Ultra-thin section electron microscopy confirmed internalization of

viruses into Birbeck granules (Figure S3A). After 30 min of incubation, we retrieved viruses

attached to the surface langerin by washing with EDTA buffer, removed non-specifically

attached viruses by trypsinization, and isolated the Birbeck granules using streptavidin-

agarose (Figure 5). A langerin mutant that lost its calcium-binding capacity (Stambach and

Taylor, 2003) was used as the negative control (Figure 5, lectin (-)). The HIV pseudoviruses

attached to both wild-type and ARGK mutant langerin and the attachment was abolished by

the addition of mannan (Figure 5A). This indicated that the mutation of the loop did not

disrupt the binding of the glycosylated envelope proteins to the primary carbohydrate-

binding sites. In contrast, the internalization of the pseudovirus was significantly inhibited

by the ARGK mutation (Figure 5B and 5C). These results suggest that disruption of Birbeck

granule formation by the mutation of loop 258-263 inhibited the internalization of HIV

pseudovirus.

Discussion

Virus internalization mechanism of Birbeck granules

In the present study, we reconstructed the 3D structure of isolated Birbeck granules and found

that docking of the flexible loop 258-263 into the secondary carbohydrate-binding cleft is

essential for assembly of the langerin lattice. In agreement with this observation, we showed

that mutations introduced into the loop inhibited Birbeck granule formation and

internalization of HIV pseudovirus. Based on the structure of the glycosylated HIV env

protein (Stewart-Jones et al., 2016; Pan et al., 2020), we propose a model of HIV

internalization by langerin (Figure 6A and 6B). A previous study proposed that N-linked

high-mannose oligosaccharides bind to langerin at both primary and secondary carbohydrate-

binding sites (Chatwell et al., 2008). Our model requires bending of the coiled-coil necks of

the two langerin trimers to form a complex with HIV-1 env via the high-mannose

oligosaccharide (Figure 6B). Although the triple coiled-coil structure of the neck domain

does not seem to bend strongly, the coiled-coil prediction profile suggests that the N-terminal

part of the neck does not necessarily form a rigid triple coiled coil (Lupas et al., 1991) (Figure

S1D). Thus, we speculate that the neck region of the langerin trimer is flexible and swings

on the cell surface, allowing loop 258-263 of one trimer to dock into the cleft of an adjacent

trimer. Once the triad structure, consisting of two langerin trimers and one env protein, is

formed, the elasticity of the neck region can deform the cell membrane and initiate

invagination. The formation of clathrin pits may also be involved in membrane invagination,

because the cytoplasmic ends of Birbeck granules are coated with clathrin (Mc et al., 2002).

Additionally, it is possible that other intracellular mechanisms such as the actin cytoskeleton,

are involved in this membrane deformation process because annexin A2 is required for the

proper formation of Birbeck granules (Thornton et al., 2020).

Significance of langerin-specific secondary carbohydrate-binding cleft

It is noteworthy that the docking of loop 258-263 into the secondary calcium-independent

carbohydrate-binding cleft, which is not found in other C-type lectins (Chatwell et al., 2008),

mediates the assembly of the langerin lattice. In DC-SIGN, which is another C-type lectin

expressed in dendritic cells, and is closely related to langerin, the region corresponding to the

secondary carbohydrate-binding site of langerin forms a shallow pocket, that is occupied by

two calcium ions (Figure 7A) (Guo et al., 2004; Geijtenbeek et al., 2000; Geijtenbeek et al.,

2000). This structural difference between langerin and DC-SIGN may account for the

inability of DC-SIGN to form Birbeck granules despite its high sequence similarity to

langerin (Cambi et al., 2009). Another characteristic of the secondary carbohydrate-binding

site of langerin is its weak affinity for mannose and maltose (Chatwell et al., 2008). This may

reflect that the sugar-binding surface of langerin CRD is less negatively charged than that of

DC-SIGN CRD (Figure 7A) (Chatwell et al., 2008). Moreover, the interaction between the

secondary binding site and mannose/maltose is mainly mediated by backbone amide and

carbonyl groups (Chatwell et al., 2008). It is possible that the side chains of Glu261 and

Asp263 of loop 258-263 docked into the cleft of an adjacent langerin CRD provide the

negative charge required for stabilizing oligo-mannose bound to the secondary binding site

(Figure 7A and B). In agreement with this hypothesis, in the present study the Asp263Lys

mutation caused a significant defect in Birbeck granule formation. The secondary

carbohydrate-binding site may only be fully functional when two langerin trimers are bound

to each other (Figure 6 and 7B). The effect of the Met260Ala mutation on Birbeck granule

formation was unexpected, because the mutation was supposed to be silent in terms of side

chain charge. The observed structural variation of the langerin trimer complex suggests that

the side chain of Met260 acts as a hook to stabilize the interface between the two intersecting

loops.

Our model of Birbeck granule formation suggests that oligosaccharide ligands are captured

by the lateral surface of langerin (Figure 6). In contrast, DC-SIGN binds to dengue virus on

its upper surface (Pokidysheva et al., 2006). The difference in the ligand-binding manner

between langerin and DC-SIGN may contribute to their different carbohydrate recognition

preferences (Valverde et al., 2020; Takahara et al., 2004). Further studies are needed to

investigate the physiological significance of the unique carbohydrate-binding mechanism of

langerin and the honeycomb lattice structure of Birbeck granules.

Methods

Isolation of Birbeck granules

The cDNA of human langerin was obtained from the Mammalian Gene Collection library

(Dnaform Inc., Yokohama, Japan) and was subcloned into pcDNA3.1-IRES-GFP (Schaefer

et al., 2008). A codon-optimized sequence coding SNAP-3×HA-HRV3C cleavage site

(obtained from ThermoFisher Scientific, Waltham, MA) was inserted immediately before the

start codon of the langerin sequence (Cole, 2013). O^6-benzylguanine-biotin was obtained by

reacting 16 mM O^6-benzylguanine-amine (Accela ChemBio, Shanghai, China) with 50 mM

biotin-AC$_5$-Osu (Dojindo, Kumamoto, Japan) and 30 mM triethylamine in

dimethylformamide for 24 hours at 30℃. Lenti-X 293T cells (TakaraBio, Tokyo, Japan) were

transfected with the langerin-expression plasmid by electroporation using ECM630

electroporator (BTX, Holliston, MA) and seeded onto the polyethylenimine-coated dish (PEI,

obtained from Fujifilm Wako Chemicals, Osaka, Japan). After 48-hour culturing in

DMEM+10% fetal bovine serum (FBS), the culture medium was replaced with serum-free

DMEM containing 10 μM O^6-benzylguanine-biotin and 10 μg/ml yeast mannan (Nacalai

tesque, Kyoto, Japan) and the cells were incubated for 30 min at 37 ℃. The cells were then

collected by scraping and washed five times with isotonic HMC buffer (Hepes 30 mM, pH

7.2, 150 mM NaCl, 2 mM $MgCl_2$, 2 mM $CaCl_2$, protease inhibitor cocktail (Nacalai tesque)).

The washed cells were homogenized with a Dounce glass homogenizer and were further

disrupted by sonication for 8 seconds using a Q125 sonicator (Qsonica, Newtown, CT). The

cell homogenate was loaded on top of a discontinuous Opti-prep density gradient (10-20-

40% layers, ThermoFisher Scientific) and ultracentrifuged for 1 hour at 100,000×g. The

fraction at the interface of 20/40% layers was collected and diluted 10-fold with HMC buffer

and was again disrupted by sonication for 2 min (120 cycles of 1-sec pulse and 3-sec wait)

at 4℃. Large debris was removed by filtration through a 20 µm-pore polyethylene mesh (GL

Science, Tokyo, Japan) and the cell lysate was incubated with a 30 µl slurry of streptavidin

agarose (Solulink, San Diego, CA) overnight at 4℃. The streptavidin agarose beads were

washed five times with HMC buffer and the bound Birbeck granules were eluted by overnight

digestion using home-made E. coli overexpressed His×6-HRV3C protease in HMC buffer at

4℃. After digestion and elution, HRV3C protease was removed by incubating the elute with

Ni-NTA resin (Qiagen, Düsseldorf, Germany). About 50 µl of elute containing released

Birbeck granules (approximately 0.05 mg/ml) were obtained from twelve 100-mm dishes of

cell culture.

Cryo-electron microscopy of isolated Birbeck granules

5 µl of the purified Birbeck granules was applied onto freshly glow-discharged holey carbon

grids, Quantifoil R1.2/1.3 Cu/Rh 200 mesh (Quantifoil Micro Tools GmbH, Großlöbichau,

Germany), blotted from the backside to concentrate the granules by filtration through the

holes, and 4 µl of HMC buffer containing cytochrome *c*-stabilized 15-nm colloidal gold (BBI

Solutions) was applied for attachment of fiducial markers (Oda et al., 2015). Grids were

blotted from the backside for 10 seconds at 4°C under 99% humidity and plunge frozen in

liquid ethane using Vitrobot Mark IV (Thermo Fisher Scientific). We removed one of the

blotting arms for one-side blotting because Birbeck granules would be absorbed onto the

filter paper. Images were recorded using a Titan Krios G3i microscope at the University of

Tokyo (Thermo Fisher Scientific) at 300 keV, a Gatan Quantum-LS Energy Filter (Gatan,

Pleasanton, CA) with a slit width of 20 eV, and a Gatan K3 Summit direct electron detector

in the electron counting mode. The nominal magnification was set to 33,000× with a physical

pixel size of 2.67 Å/pixel. Movies were acquired using the SerialEM software (Mastronarde,

2005), and the target defocus was set to 3-6 µm. The angular range of the tilt series was from

-60° to 60° with 3.0° increments. Each movie was recorded for 0.74 sec with a total dose of

1.24 electrons/Å^2 and subdivided into 20 frames. The total dose for one tilt series acquisition

is thus 50 electrons/$Å^2$.

Data processing

Movies were subjected to beam-induced motion correction using MotionCor2 (Zheng et al.,

2017), and tilt series images were aligned, CTF corrected, and back-projected to reconstruct

3D tomograms using the IMOD software package (Kremer et al., 1996). For the subsequent

analysis using Relion-4.0 (Kimanius et al., 2021), the tomogram positioning step was skipped

to keep the angle offset and the Z-shift zero. Tomograms were 4×binned to reduce loads of

the calculation. The areas of the Birbeck granules were manually selected using the 3dmod

model tool (Kremer et al., 1996), and the center coordinates for the subtomograms were

determined by placing points at 8-pixel (~8.5 nm) intervals within the selected area using a

custom-made Ruby script (Metlagel et al., 2007). For twisting Birbeck granules (Figure S1C),

center points were picked manually by selecting the mid-portion of the lamella. Volumes with

32×32×32 pixel-dimensions were extracted from 8×binned tomograms and were averaged

using the PEET software suite (Nicastro et al., 2006). A randomly selected subtomogram was

used for the initial reference. Alignments were repeated five times for 8×binned and twice

for 4×binned tomograms with 64×64×64 pixel-dimensions. The coordinates and the rotation

angles of the aligned subtomograms were exported to Relion-4.0. Initial

pseudosubtomograms were generated with the box size of 128 pixels, the cropped box size

of 64 pixels, and the binning factor of 4. The subtomograms were 3D refined using a spherical

mask of 320-Å diameter. After 3D classification, 2×binned pseudosubtomograms were

generated with the box size of 256 pixels and the cropped box size of 128 pixels based on the

refined coordinates and angles and were 3D refined using the same spherical mask. Finally,

un-binned pseudosubtomograms were generated with a box size of 256 pixels and a cropped

box size of 128 pixels. 3D refinement was done using a tighter mask enclosing the central

upright langerin molecule and three bound CRDs, and low-resolution subtomograms were

removed by 3D classification. The alignment was further refined using CTF refinement and

frame alignment. We initially refined the surrounding inverted langerin molecule by 3D

refinement masking out the central upright molecule, but the refinement was biased toward

the center. To improve the map quality of the inverted molecule, we applied 3D multi-body

refinement (Nakane et al., 2018), which successfully compensated for the flexibility between

the upright and the inverted molecules. For model building, the crystal structure of langerin

trimer (Feinberg et al., 2010) (PDB ID: 3KQG) was fitted into the map using UCSF Chimera

(Pettersen et al., 2004). Then, the fitted model was further refined using PHENIX real-space

refinement tool, and the refined model was validated using the comprehensive validation tool

(Adams et al., 2010). The Coulombic surfaces rendering was conducted using UCSF

Chimera and other model presentations, surface rendering, and morph movie generation were

conducted using ChimeraX (Goddard et al., 2018). The data processing and the statistics of

the model validation are summarized in Figure S2A and Table S1.

Ultrathin section electron microscopy of Birbeck granules

Pellets of 293T cells with mannan-induced Birbeck granules were fixed with 4%

paraformaldehyde for 1 hour at 4℃ and stained with 1% osmium tetroxide and subsequently

with 1% uranium acetate. After dehydration in ethanol and acetone, the samples were

embedded in Quetol 812 resin (Nissin EM, Tokyo, Japan). 60-nm ultrathin sections were cut

using a ULTRACUT microtome (Reichert Leica) and mounted onto Formvar-coated copper

grid. Images were recorded using a JEM-2100F microscope (JEOL, Tokyo, Japan) at

University of Yamanashi operated at 200 keV equipped with an F216 CMOS camera (TVIPS

GmbH, Gauting, Germany). Length of Birbeck granules were measured using Fiji software

(Schindelin et al., 2012), and plots were generated using PlotsOfData web tool (Postma and

Goedhart, 2019). For total length of Birbeck granules per 100 μm^2, cross sections of cells

with nuclei of > 5 μm in diameter were selected, and the sum of the lengths of Birbeck

granules within the cells were measured. The areas of the cross sections were also measured

using Fiji software. 100 μm^2 corresponds to the approximate mean of the cross-sectional area

of one cell.

Lentivirus preparation

Lentiviruses pseudotyped with HIV-1 env were prepared by transfecting the Lenti-X 293T

cells with pCMV-dR8.3 Δvpr (Addgene plasmid #8455), pLOX-CW-tdTomato, and pCEP4-

AD8gp160 (Adgene plasmid #123260) (Stewart et al., 2003; Heredia et al., 2019). The vector

pLOX-CW-tdTomato was generated by replacing the GFP sequence of pLOX-CWgfp

(Addgene plasmid #12241) (Salmon et al., 2000) with the tdTomato sequence derived from

pCDH-EF1-Luc2-P2A-tdTomato, a gift from Kazuhiro Oka (Addgene plasmid # 72486). The

transfections were carried out using the PEI-method with the ratio at PEI: dR8.3: tdTomato:

gp160 = 4:1:1:1. 16 hours post-transfection, the cells were washed three times with PBS and

incubated in a fresh medium for another 32 hours. The virus-containing medium was

collected and removed cell debris by a 1,000×g centrifuge and a 0.45 μm filtration (Millipore).

The obtained supernatants were immediately used for subsequent assays.

Virus attachment and internalization assay

293T cells were transfected with wild type langerin, the ARGK mutant, and, as a negative

control, a lectin (-) mutant lacking calcium-binding ability. The lectin (-) langerin mutant was

generated by introducing three point-mutations at the calcium-binding site: Glu285Lys;

Asn287Lys; and Glu293Lys (Chatwell et al., 2008). We conducted surface labeling of

langerin-expressing cells and confirmed that these mutations did not alter the surface

expression of langerin (Figure S3B). 48 hours post-transfection, the cells were pre-treated

with or without 10 µg/ml yeast mannan for 5 min at 37°C. The cells were then incubated in

the pseudovirus-containing supernatants plus 10 µM O^6-benzylguanine-biotin and 10 µg/ml

DEAE-dextran (Sigma Aldrich, St. Louis, MO) with or without 10 µg/ml yeast mannan for

30 min at 37°C. After the incubation, supernatant containing unbound viruses were collected.

The cells were then washed with TBS-Ca buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 5

mM $CaCl_2$) three times, and the viruses attached to langerin at the cell surface was retrieved

by washing with TBS-EDTA buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 10 mM EDTA).

The collected supernatant of the medium and the TBS-EDTA buffer were centrifuged at

1,000×g for 5 min to remove cell debris, and the viruses were sedimented by ultra-

centrifugation at 100,000×g for 3 hours. After the TBS-EDTA washing, non-specifically

bound viruses were removed by trypsinization using 0.25 % (w/v) trypsin and 1mM EDTA

for 5 min at 37℃ as previously described (Burkard et al., 2014). Detached cells were washed

twice with DMEM+10 % FBS, and five times with TBS-Ca. Internalized viruses within

Birbeck granules were isolated from the washed cells by streptavidin-agarose precipitation

as described above. Viruses, langerin, and tubulin were detected by immunoblotting using

HIV-1 gag p24 antibody (1:1000 dilution, MAB7360, mouse monoclonal, R&D systems,

Minneapolis, MN), human CD207 antibody (1:1000 dilution, 11841-1-AP, rabbit polyclonal,

Proteintech Group Inc., Rosemont, IL), and tubulin antibody (1:3000 dilution, B-512, mouse

monoclonal, Sigma-Aldrich), respectively. ELSA was carried out using HIV Type 1 p24

Antigen ELISA kit (ZeptoMatrix, Buffalo, NY). The p24 antibody detected unprocessed

(Pr55^{gag}) and fully processed (p24) capsid proteins (Gonelli et al., 2019). Band intensities

were measured using ImageJ (Schneider et al., 2012). Surface labeling was carried out by

treating the cells with 0.5mM biotin-sulfo-osu (Dojindo, Osaka, Japan) for 30 min at 37℃.

Cells were washed three times with TBS and collected by scraping. Surface-expressed

proteins were extracted by incubating the cells in TBS plus 1% Triton X-100 for 1 hour at

4℃, and immunoprecipitated using streptavidin agarose. ELISA using p24 antibody showed

that >99% of the viruses remained in the supernatant (Figure S3C).

Isolation of langerin-expressing stable cell lines

We isolated a cell line stably expressing langerin by selecting HEK293 cells using G418

(Sigma Aldrich). However, the expression levels of the isolated cell line was 10~20% of that

of the transiently expressing cells (Figure S3D). These cells can form Birbeck granules in the

presence of yeast mannan, but with lower efficiency (Figure S3E and F). Therefore, we chose

to use the transient expression system in the main experiments.

Author contributions

T.O. conceived the project and designed the experiments. T.O., H.Y., and H.S. performed the experiments. T.O., and H.Y. analyzed the data. T.O., H.Y., Y.O., and T.K. wrote the manuscript.

Data availability

The map and the model will be available on the EMDB upon publication under the following accession numbers: EMD-32906, and PDB 7WZ8.

Acknowledgements

We thank Mrs. Natsuko Maruyama (University of Yamanashi) for technical assistance. This research is partially supported by Platform Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from Japan Agency for Medical Research and Development (AMED) under Grant Number JP19am0101115. This research was conducted using the Fujitsu PRIMERGY CX400M1/CX2550M5 (Oakbridge-CX) and the SGI Rackable C2112-4GP3/C1102-GP8 (Reedbush-H/L) in the Information Technology Center, The University of

Tokyo. This work was supported by the Takeda Science Foundation (to T.O.), the Daiichi

Sankyo Foundation of Life Science (to T.O.), the Japan Society for the Promotion of Science

(KAKENHI Grant number JP21H02654 to T.O.), and the Naito Foundation (to T.O.).

Molecular graphics and analyses performed with UCSF ChimeraX, developed by the

Resource for Biocomputing, Visualization, and Informatics at the University of California,

San Francisco, with support from National Institutes of Health R01-GM129325 and the

Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and

Infectious Diseases.

References

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W. et al. 2010. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta. Crystallogr. D. Biol. Crystallogr.* **66**, 213-221.

Bartosik, J. 1992. Cytomembrane-derived Birbeck granules transport horseradish peroxidase to the endosomal compartment in the human Langerhans cells. *J. Invest. Dermatol.* **99**, 53-58.

Birbeck, M.S., Breathnach, A.S., and Everall, J.D. 1961. An electron microscope study of basal melanocytes and high-level clear cells (Langerhans cells) in vitiligo. *J. Invest. Dermatol.* **37**, 51-63.

Burkard, C., Bloyet, L.M., Wicht, O., van, Kuppeveld F.J., Rottier, P.J., de, Haan C.A., and Bosch, B.J. 2014. Dissecting virus entry: replication-independent analysis of virus binding, internalization, and penetration using minimal complementation of beta-galactosidase. *PLoS. One.* **9**, e101762.

Cambi, A., Beeren, I., Joosten, B., Fransen, J.A., and Figdor, C.G. 2009. The C-type lectin DC-SIGN internalizes soluble antigens and HIV-1 virions via a clathrin-dependent mechanism. *Eur. J. Immunol.* **39**, 1923-1928.

Chatwell, L., Holla, A., Kaufer, B.B., and Skerra, A. 2008. The carbohydrate recognition domain of Langerin reveals high structural similarity with the one of DC-SIGN but an additional, calcium-independent sugar-binding site. *Mol. Immunol.* **45**, 1981-1994.

Cole, N.B. 2013. Site-specific protein labeling with SNAP-tags. *Curr. Protoc. Protein. Sci.* **73**, 30.1.1-30.1.16.

de Jong, M.A., de, Witte L., Taylor, M.E., and Geijtenbeek, T.B. 2010. Herpes simplex virus type 2 enhances HIV-1 susceptibility by affecting Langerhans cell function. *J. Immunol.* **185**, 1633-1641.

de_Jong MA, Geijtenbeek T.B. 2010. Langerhans cells in innate defense against pathogens. *Trends. Immunol.* **31**, 452-459.

de Witte, L., Nabatov, A., Pion, M., Fluitsma, D., de Jong M.A., de, Gruijl T., Piguet, V., van, Kooyk Y., and Geijtenbeek, T.B. 2007. Langerin is a natural barrier to HIV-1

transmission by Langerhans cells. *Nat. Med.* **13**, 367-371.

Doebel, T., Voisin, B., and Nagao, K. 2017. Langerhans Cells - The Macrophage in

Dendritic Cell Clothing. *Trends. Immunol.* **38**, 817-828.

Feinberg, H., Powlesland, A.S., Taylor, M.E., and Weis, W.I. 2010. Trimeric structure of

langerin. *J. Biol. Chem.* **285**, 13285-13293.

Figdor, C.G., van, Kooyk Y., and Adema, G.J. 2002. C-type lectin receptors on dendritic

cells and Langerhans cells. *Nat. Rev. Immunol.* **2**, 77-84.

Geijtenbeek, T.B., Kwon, D.S., Torensma, R., van, Vliet S.J., van, Duijnhoven G.C.,

Middel, J., Cornelissen, I.L., Nottet, H.S., KewalRamani, V.N., Littman, D.R. et al. 2000.

DC-SIGN, a dendritic cell-specific HIV-1-binding protein that enhances trans-infection of

T cells. *Cell.* **100**, 587-597.

Geijtenbeek, T.B., Krooshoop, D.J., Bleijs, D.A., van, Vliet S.J., van, Duijnhoven G.C.,

Grabovsky, V., Alon, R., Figdor, C.G., and van, Kooyk Y. 2000. DC-SIGN-ICAM-2

interaction mediates dendritic cell trafficking. *Nat. Immunol.* **1**, 353-357.

Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., and

Ferrin, T.E. 2018. UCSF ChimeraX: Meeting modern challenges in visualization and

analysis. *Protein. Sci.* **27**, 14-25.

Gonelli, C.A., Khoury, G., Center, R.J., and Purcell, D.F.J. 2019. HIV-1-based Virus-like

Particles that Morphologically Resemble Mature, Infectious HIV-1 Virions. *Viruses.* **11**,

507.

Guo, Y., Feinberg, H., Conroy, E., Mitchell, D.A., Alvarez, R., Blixt, O., Taylor, M.E.,

Weis, W.I., and Drickamer, K. 2004. Structural basis for distinct ligand-binding and

targeting properties of the receptors DC-SIGN and DC-SIGNR. *Nat. Struct. Mol. Biol.* **11**,

591-598.

Heredia, J.D., Park, J., Choi, H., Gill, K.S., and Procko, E. 2019. Conformational

Engineering of HIV-1 Env Based on Mutational Tolerance in the CD4 and PG16 Bound

States. *J. Virol.* **93**, e00219-00219.

Hunger, R.E., Sieling, P.A., Ochoa, M.T., Sugaya, M., Burdick, A.E., Rea, T.H., Brennan,

P.J., Belisle, J.T., Blauvelt, A., Porcelli, S.A. et al. 2004. Langerhans cells utilize CD1a and

langerin to efficiently present nonpeptide antigens to T cells. *J. Clin. Invest.* **113**, 701-708.

Kawamura, T., Cohen, S.S., Borris, D.L., Aquilino, E.A., Glushakova, S., Margolis, L.B.,

Orenstein, J.M., Offord, R.E., Neurath, A.R., and Blauvelt, A. 2000. Candidate

microbicides block HIV-1 infection of human immature Langerhans cells within epithelial

tissue explants. *J. Exp. Med.* **192**, 1491-1500.

Kawamura, T., Qualbani, M., Thomas, E.K., Orenstein, J.M., and Blauvelt, A. 2001. Low

levels of productive HIV infection in Langerhans cell-like dendritic cells differentiated in

the presence of TGF-beta1 and increased viral replication with CD40 ligand-induced

maturation. *Eur. J. Immunol.* **31**, 360-368.

Kimanius, D., Dong, L., Sharov, G., Nakane, T., and Scheres, S.H.W. 2021. New tools for

automated cryo-EM single-particle analysis in RELION-4.0. *Biochem. J.* **478**, 4169-4185.

Kremer, J.R., Mastronarde, D.N., and McIntosh, J.R. 1996. Computer visualization of

three-dimensional image data using IMOD. *J. Struct. Biol.* **116**, 71-76.

Lenormand, C., Spiegelhalter, C., Cinquin, B., Bardin, S., Bausinger, H., Angenieux, C.,

Eckly, A., Proamer, F., Wall, D., Lich, B. et al. 2013. Birbeck granule-like "organized

smooth endoplasmic reticulum" resulting from the expression of a cytoplasmic YFP-tagged

langerin. *PLoS. One.* **8**, e60813.

Lupas, A., Van, Dyke M., and Stock, J. 1991. Predicting coiled coils from protein

sequences. *Science.* **252**, 1162-1164.

Mastronarde, D.N. 2005. Automated electron microscope tomography using robust

prediction of specimen movements. *J. Struct. Biol.* **152**, 36-51.

Mc, Dermott R., Ziylan, U., Spehner, D., Bausinger, H., Lipsker, D., Mommaas, M.,

Cazenave, J.P., Raposo, G., Goud, B., de, la Salle H. et al. 2002. Birbeck granules are

subdomains of endosomal recycling compartment in human epidermal Langerhans cells,

which form where Langerin accumulates. *Mol. Biol. Cell.* **13**, 317-335.

McDermott, R., Bausinger, H., Fricker, D., Spehner, D., Proamer, F., Lipsker, D.,

Cazenave, J.P., Goud, B., De, La Salle H., Salamero, J. et al. 2004. Reproduction of

Langerin/CD207 traffic and Birbeck granule formation in a human cell line model. *J.*

Invest. Dermatol. **123**, 72-77.

Merad, M., Ginhoux, F., and Collin, M. 2008. Origin, homeostasis and function of

Langerhans cells and other langerin-expressing dendritic cells. *Nat. Rev. Immunol.* **8**, 935-

947.

Metlagel, Z., Kikkawa, Y.S., and Kikkawa, M. 2007. Ruby-Helix: an implementation of

helical image processing based on object-oriented scripting language. *J. Struct. Biol.* **157**,

95-105.

Nakane, T., Kimanius, D., Lindahl, E., and Scheres, S.H. 2018. Characterization of

molecular motions in cryo-EM single-particle data by multi-body refinement in RELION.

Elife. **7**, e36861.

Nicastro, D., Schwartz, C., Pierson, J., Gaudette, R., Porter, M.E., and McIntosh, J.R. 2006.

The molecular architecture of axonemes revealed by cryoelectron tomography. *Science.*

313, 944-948.

Oda, T., Yanagisawa, H., and Kikkawa, M. 2015. Detailed structural and biochemical

characterization of the nexin-dynein regulatory complex. *Mol. Biol. Cell.* **26**, 294-304.

Ogawa, Y., Kawamura, T., Matsuzawa, T., Aoki, R., Gee, P., Yamashita, A., Moriishi, K.,

Yamasaki, K., Koyanagi, Y., Blauvelt, A. et al. 2013. Antimicrobial peptide LL-37

produced by HSV-2-infected keratinocytes enhances HIV infection of Langerhans cells.

Cell. Host. Microbe. **13**, 77-86.

Pan, J., Peng, H., Chen, B., and Harrison, S.C. 2020. Cryo-EM Structure of Full-length

HIV-1 Env Bound With the Fab of Antibody PG16. *J. Mol. Biol.* **432**, 1158-1168.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C.,

and Ferrin, T.E. 2004. UCSF Chimera--a visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25**, 1605-1612.

Pokidysheva, E., Zhang, Y., Battisti, A.J., Bator-Kelly, C.M., Chipman, P.R., Xiao, C.,

Gregorio, G.G., Hendrickson, W.A., Kuhn, R.J., and Rossmann, M.G. 2006. Cryo-EM

reconstruction of dengue virus in complex with the carbohydrate recognition domain of

DC-SIGN. *Cell.* **124**, 485-493.

Postma, M., and Goedhart, J. 2019. PlotsOfData-A web app for visualizing data together

with their summaries. *PLoS. Biol.* **17**, e3000202.

Rhodes, J.W., Botting, R.A., Bertram, K.M., Vine, E.E., Rana, H., Baharlou, H., Vegh, P.,

O'Neil, T.R., Ashhurst, A.S., Fletcher, J. et al. 2021. Human anogenital monocyte-derived

dendritic cells and langerin+cDC2 are major HIV target cells. *Nat. Commun.* **12**, 2147.

Ribeiro, C.M., Sarrami-Forooshani, R., Setiawan, L.C., Zijlstra-Willems, E.M., van,

Hamme J.L., Tigchelaar, W., van, der Wel N.N., Kootstra, N.A., Gringhuis, S.I., and

Geijtenbeek, T.B. 2016. Receptor usage dictates HIV-1 restriction by human TRIM5alpha

in dendritic cell subsets. *Nature.* **540**, 448-452.

Sagebiel, R.W., and Reed, T.H. 1968. Serial reconstruction of the characteristic granule of

the Langerhans cell. *J. Cell. Biol.* **36**, 595-602.

Salmon, P., Oberholzer, J., Occhiodoro, T., Morel, P., Lou, J., and Trono, D. 2000.

Reversible immortalization of human primary cells by lentivector-mediated transfer of

specific genes. *Mol. Ther.* **2**, 404-414.

Schaefer, M.R., Williams, M., Kulpa, D.A., Blakely, P.K., Yaffee, A.Q., and Collins, K.L.
2008. A novel trafficking signal within the HLA-C cytoplasmic tail allows regulated
expression upon differentiation of macrophages. *J. Immunol.* **180**, 7804-7817.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B. et al. 2012. Fiji: an open-source platform
for biological-image analysis. *Nat. Methods.* **9**, 676-682.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. 2012. NIH Image to ImageJ: 25 years
of image analysis. *Nat. Methods.* **9**, 671-675.

Stambach, N.S., and Taylor, M.E. 2003. Characterization of carbohydrate recognition by
langerin, a C-type lectin of Langerhans cells. *Glycobiology.* **13**, 401-410.

Stewart, S.A., Dykxhoorn, D.M., Palliser, D., Mizuno, H., Yu, E.Y., An, D.S., Sabatini,
D.M., Chen, I.S., Hahn, W.C., Sharp, P.A. et al. 2003. Lentivirus-delivered stable gene
silencing by RNAi in primary cells. *RNA.* **9**, 493-501.

Stewart-Jones, G.B., Soto, C., Lemmin, T., Chuang, G.Y., Druz, A., Kong, R., Thomas,
P.V., Wagh, K., Zhou, T., Behrens, A.J. et al. 2016. Trimeric HIV-1-Env Structures Define
Glycan Shields from Clades A, B, and G. *Cell.* **165**, 813-826.

Takahara, K., Yashima, Y., Omatsu, Y., Yoshida, H., Kimura, Y., Kang, Y.S., Steinman,
R.M., Park, C.G., and Inaba, K. 2004. Functional comparison of the mouse DC-SIGN,
SIGNR1, SIGNR3 and Langerin, C-type lectins. *Int. Immunol.* **16**, 819-829.

Tateno, H., Ohnishi, K., Yabe, R., Hayatsu, N., Sato, T., Takeya, M., Narimatsu, H., and
Hirabayashi, J. 2010. Dual specificity of Langerin to sulfated and mannosylated glycans via
a single C-type carbohydrate recognition domain. *J. Biol. Chem.* **285**, 6390-6400.

Thepaut, M., Valladeau, J., Nurisso, A., Kahn, R., Arnou, B., Vives, C., Saeland, S., Ebel,
C., Monnier, C., Dezutter-Dambuyant, C. et al. 2009. Structural studies of langerin and
Birbeck granule: a macromolecular organization model. *Biochemistry.* **48**, 2684-2698.

Thornton, S.M., Samararatne, V.D., Skeate, J.G., Buser, C., Luhen, K.P., Taylor, J.R., Da,
Silva D.M., and Kast, W.M. 2020. The Essential Role of anxA2 in Langerhans Cell
Birbeck Granules Formation. *Cells.* **9**, 974.

Turville, S.G., Cameron, P.U., Handley, A., Lin, G., Pohlmann, S., Doms, R.W., and
Cunningham, A.L. 2002. Diversity of receptors binding HIV on dendritic cell subsets. *Nat.*
Immunol. **3**, 975-983.

Valladeau, J., Duvert-Frances, V., Pin, J.J., Dezutter-Dambuyant, C., Vincent, C.,
Massacrier, C., Vincent, J., Yoneda, K., Banchereau, J., Caux, C. et al. 1999. The

monoclonal antibody DCGM4 recognizes Langerin, a protein specific of Langerhans cells,

and is rapidly internalized from the cell surface. *Eur. J. Immunol.* **29**, 2695-2704.

Valladeau, J., Ravel, O., Dezutter-Dambuyant, C., Moore, K., Kleijmeer, M., Liu, Y.,

Duvert-Frances, V., Vincent, C., Schmitt, D., Davoust, J. et al. 2000. Langerin, a novel C-

type lectin specific to Langerhans cells, is an endocytic receptor that induces the formation

of Birbeck granules. *Immunity.* **12**, 71-81.

Valladeau, J., Dezutter-Dambuyant, C., and Saeland, S. 2003. Langerin/CD207 sheds light

on formation of birbeck granules and their possible function in Langerhans cells. *Immunol.*

Res. **28**, 93-107.

Valverde, P., Martinez, J.D., Canada, F.J., Arda, A., and Jimenez-Barbero, J. 2020.

Molecular Recognition in C-Type Lectins: The Cases of DC-SIGN, Langerin, MGL, and L-

Sectin. *Chembiochem.* **21**, 2999-3025.

Verdijk, P., Dijkman, R., Plasmeijer, E.I., Mulder, A.A., Zoutman, W.H., Mieke,

Mommaas A., and Tensen, C.P. 2005. A lack of Birbeck granules in Langerhans cells is

associated with a naturally occurring point mutation in the human Langerin gene. *J. Invest.*

Dermatol. **124**, 714-717.

Weis, W.I., and Drickamer, K. 1994. Trimeric structure of a C-type mannose-binding

protein. *Structure.* **2**, 1227-1240.

Weis, W.I., and Drickamer, K. 1996. Structural basis of lectin-carbohydrate recognition.

Annu. Rev. Biochem. **65**, 441-473.

Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. 2017.

MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron

microscopy. *Nat. Methods.* **14**, 331-332.

Figure 1

Figure 1: Isolation of Birbeck granules. (A) Domain organization of langerin. Human

langerin is a 328-amino acids protein composed of an N-terminal cytoplasmic tail, a

transmembrane domain (TM), a coiled-coil neck, and a C-terminal carbohydrate-recognition

domain (CRD). We fused a SNAP-tag at the N-terminus of langerin and inserted an HRV3C

protease cleavage site between the tag and the langerin sequences. For streptavidin-mediated

precipitation of Birbeck granules, the SNAP-tag was biotinylated (star). (B) Model of

langerin oligomerization within Birbeck granules. Langerin trimers bind to each other face-

to-face via the CRDs, bringing the two layers of the plasma membrane closer together. (C)

Birbeck granules formed in 293T cells overexpressing langerin. Addition of yeast mannan

induced the formation of Birbeck granules a few micrometers long. Inset shows a magnified

view of Birbeck granules. (D) SDS-PAGE of isolated Birbeck granules. Purified langerin

(arrowhead) was released from streptavidin-agarose by HRV3C digestion. M: molecular

weight marker; and BG: isolated Birbeck granules. (E) Cryo-electron microscopy of isolated

Birbeck granules. Black square indicates the position of the close-up view shown in F. (F)

Wavy lamellar structure of the Birbeck granule. Black dots were gold nanoparticles used as

fiducial markers. (G) Class averages of the projection images of Birbeck granules. The image

dimension is 34 nm^2. Although 2D classification did not converge well due to the continuity

617   of the structure, some classes showed a porous structure with a honeycomb-like lattice.

A

B

C

5 nm

D

10 nm

Figure 2

Figure 2: Averaged subtomograms of Birbeck granules. (A) Result of the initial 3D

refinement. The central langerin trimer (blue) binds to the three inverted langerin trimers

(red). (B) Result of the focused refinement. Improved resolution (6.4 Å) allowed secondary

structure modeling. The gray mesh and the cyan surface indicate low and high threshold

isosurfaces, respectively. (C) Result of the two-body refinement. The second body (orange)

was refined with respect to the first body (cyan). (D) Honeycomb model of the langerin lattice

within Birbeck granules. This honeycomb lattice is an ideal model,   with the assumption of

structural uniformity. (A-D) left, top views; right, side views.

Figure 3

Figure 3: Binding interface between two langerin trimers. (A) Two langerin trimers

interact via the loops at residues 258-263. Amino acid codes and residue numbers are

indicated. Yellow and pink circles indicate the primary and secondary carbohydrate-binding

sites. The orientations of the side chains are speculative because the resolution of these maps

(6.4 Å) was insufficient for precise modeling. Calcium ions (green) were placed based on the

crystal structure (PDB ID: 3KQG). Top and side views are displayed in the top and bottom

rows, respectively. Broken squares indicate the positions of the magnified views on the right.

(B) Flexibility between the two trimers. The first eigenvector of the structural variation of

the inverted trimer (red) relative to the upright trimer (blue) shows an approximately 30°

rotation about the loop 258-263 (see Movie S2).

Figure 4

Figure 4: Mutations to the loop 258-263 disrupted Birbeck granule formation. (A) Ultra-

thin electron microscopy of Birbeck granules with and without yeast mannan in langerin-

overexpressing 293T cells. Residues at 260-263 (Met-Glu-Gly-Asp, MEGD) were mutated

to Met-Arg-Gly-Asp (MRGD), Met-Arg-Gly-Lys (MRGK), or Ala-Arg-Gly-Lys (ARGK).

The Mannan (+) WT image has a lower magnification than the others in order to show the

entire Birbeck granules. (B-D) Quantification of Birbeck granule formation. Horizontal lines

indicate the median values. Single asterisks and the double asterisk indicate statistically

significant differences with $p < 0.01$ and $p = 0.0101$, respectively. NS indicates no statistically

significant differences. p values were calculated using Bonferroni-corrected Student's t-tests.

(B) Lengths of individual Birbeck granules were measured. N = 337 (WT mannan-), 386

(WT mannan+), 348 (MRGD), 233 (MRGK), and 70 (ARGK). (C) Sum of the length of

Birbeck granules per 100 μm^2 cell area. Cross-sections of cells with nuclei of > 5 μm in

diameter were selected and the total lengths of the Birbeck granules were measured within

the cells. N = 20 for all the samples. (D) Number of the Birbeck granules per 100 μm^2 cell

area. N = 20 for all the samples.

Figure 5

Figure 5: Mutation to the loop 258-263 inhibited the internalization of pseudoviruses.

HIV-1 pseudoviruses were added to langerin-expressing 293T cells. Yeast mannan (10

µg/ml) was added to block the lectin-dependent binding of pseudoviruses. A langerin mutant

lacking calcium binding ability (lectin (-)) was used as the negative control. (A) Immunoblots

of pseudoviruses attached to the cell surface. Unbound and attached viruses were collected

from the supernatant of the culture medium and TBS-EDTA buffer, respectively. Samples of

unbound viruses were diluted 50-fold to adjust the band intensities. The expression levels of

SNAP-tagged langerin show that the numbers of transfected cells were approximately the

same in each experiment. $Pr55^{gag}$ and p24 indicate unprocessed and fully-processed capsid

proteins, respectively. (B) Immunoblots of internalized pseudoviruses. Birbeck granules were

isolated by precipitation using streptavidin-agarose, and intracellular viruses and langerin

were detected by their respective antibodies. Tubulins in the whole-cell lysates were detected

for loading controls. (C) Quantification of internalized viruses using p24 ELISA. Horizontal

lines indicate the mean. NS and Asterisk indicate no significant difference and statistically

significant differences ($p < 0.001$) calculated using Bonferroni-corrected Student's *t*-tests

(N=4), respectively.

A

B

HIV

env

Langerin

Plasma membrane

Figure 6

Figure 6: Model of HIV internalization by langerin. (A) Glycosylated structure of HIV-1

envelope (env, gray) (Constructed based on PDB ID: 5FYJ and 6ULC) bound to the langerin

trimer complex. Branches of the high-mannose oligosaccharide interacting with the primary

(yellow) and secondary (pink) carbohydrate-binding sites. The broken square indicates the

position of the close-up view on the right. (B) Schematics of HIV internalization by langerin.

The inset on the top shows an overview of antigen processing within a Langerhans cell.

Viruses are captured by langerin, internalized into a Birbeck granule, and degraded by the

autophagic system. The red square indicates the viral entry process described in the lower

panels. The high-mannose oligosaccharide on HIV-1 env first binds to the primary

carbohydrate-binding site, and one of the branches of the oligo-mannose weakly interacts

with the secondary binding site. The flexible neck region allows docking of the loop 258-263

of an adjacent trimer into the secondary carbohydrate-binding cleft, reinforcing the binding

of the oligo-mannose. As the elastic neck regions straighten, the plasma membrane is

deformed into a lamellar invagination, which internalizes the virus.

A

Langerin

DC-SIGN

B

Langerin

Figure 7

Figure 7: Coulombic surfaces of the carbohydrate-binding sites. (A) Comparison of the

surface charge between the CRDs of langerin and DC-SIGN (PDB ID: 1SL4). Blue and red

coloring indicates positive and negative charges, respectively. The primary and secondary

carbohydrate-binding sites are indicated by yellow and pink arrowheads, respectively. The

region of DC-SIGN corresponding to the secondary carbohydrate-binding sites of langerin is

occupied by calcium ions (green) and does not form a cleft. (B) Surface model of the langerin

trimer complex. Mannose ligands are placed based on a previously proposed model (Chatwell

et al., 2008). (A-B) Broken squares indicate the position and the orientation of the side (left)

and the top (right) views. Oligosaccharide ligands are shown in gray.