Phenotypic and transcriptional response of *Daphnia pulicaria* to the combined effects of

temperature and predation

Aaron Oliver[1]¶, Hamanda B. Cavalheri[2]¶, Thiago G. Lima[2], Natalie T. Jones[2], Sheila

Podell[1], Daniela Zarate[2], Eric Allen[1], Ronald S. Burton[3], Jonathan B. Shurin[2]*

[1] Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,

University of California, San Diego, La Jolla, California, United States of America

[2] Department of Biological Sciences, Ecology Behavior and Evolution Section, University

of California, San Diego, La Jolla, California, United States of America

[3] Marine Biology Research Division, Scripps Institution of Oceanography, University of

California, San Diego, La Jolla, California, United States of America

¶These authors contributed equally to this work

* Corresponding author

Email: jshurin@ucsd.edu (JBS)

Abstract

    Daphnia, an ecologically important zooplankton species in lakes, shows both genetic adaptation and phenotypic plasticity in response to temperature and fish predation, but little is known about the molecular basis of these responses and their potential interactions. We performed a factorial experiment exposing laboratory-propagated *Daphnia pulicaria* clones from two lakes in the Sierra Nevada mountains of California to normal or high temperature (15°C or 25°C) in the presence or absence of fish kairomones, then measured changes in life history and gene expression. Exposure to kairomones increased upper thermal tolerance limits for physiological activity in both clones. Cloned individuals matured at a younger age in response to higher temperature and kairomones, while size at maturity, fecundity and population intrinsic growth were only affected by temperature. At the molecular level, both clones expressed more genes differently in response to temperature than predation, but specific genes involved in metabolic, cellular, and genetic processes responded differently between the two clones. Although gene expression differed more between clones from different lakes than experimental treatments, similar phenotypic responses to predation risk and warming arose from these clone-specific patterns. Our results suggest that phenotypic plasticity responses to temperature and kairomones interact synergistically, with exposure to fish predators increasing the tolerance of *Daphnia pulicaria* to stressful temperatures, and that similar phenotypic responses to temperature and predator cues can be produced by divergent patterns of gene regulation.

Introduction

Many species are at risk of extinction as the environment changes at an unprecedented pace. Ecological and evolutionary research aims to predict species' responses to anthropogenic environmental change, such as global warming and introduction of predators [1,2]. To cope with such stressors, species can either migrate to more suitable habitats or adapt to new conditions. Understanding the potential and limits of genetic adaptation and phenotypic plasticity to maintain fitness is critical in order to predict persistence of populations and changes in biodiversity under environmental stresses [3]. The mechanisms by which organisms are adapted to present day conditions throughout their ranges therefore provide a basis for predicting extinction, persistence, and changes in biodiversity in a rapidly changing environment [4,5].

Plasticity and genetic adaptation impact organismal fitness in response to environmental change. Plasticity allows a genotype to express multiple phenotypes in different environments whereas genetic adaptation arises from environment-dependent variation in fitness of different genotypes [6]. Evidence for rapid evolution in natural systems, particularly in response to human-induced environmental change, indicates that most of the observed changes are not genetically based, but rather a consequence of plasticity [7]. Plasticity at level of gene expression is one of the most important mechanisms for coping with stress [8,9], yet the magnitude and genetic basis of such plasticity remains largely unknown.

Plasticity arises from differential gene expression patterns in response to environmental cues [10]. Variable levels of plasticity may evolve in a population if reaction norms differ across genotypes and the slope of the reaction norm is correlated with fitness

[11]. Either decreases or increases in phenotypic plasticity could contribute to adaptation

to variable environments depending on the rate and predictability of environmental change

[6,12].

Several factors could constrain the evolution of plasticity at the transcriptome level.

First, genetic variation potential for plasticity could be limited or absent. Alternatively,

even when genetic variation is present, its evolution could be constrained by costs of the

mechanisms underlying plastic responses [13]. A trade-off is expected where enhanced

plasticity would be beneficial in more spatially or temporally variable environments, but

detrimental in a stable environment [14,15]. In addition, if environmental variation is

unpredictable, then plasticity may fail to match phenotypes to the environment to produce

higher fitness.

Daphnia is an ecologically important genus of crustacean zooplankton in lakes that

transfers energy from phytoplankton to fish and invertebrate predators and exerts top-down

grazing control on primary production [16]. *Daphnia* shows both genetic adaptation and

plasticity at a phenotypic level in tolerance to two important determinants of fitness:

temperature and predation. Previous studies have documented that the response of *Daphnia*

populations to thermal stress is strongly correlated with environmental conditions and can

be either genetic, plastic or both [17–19]. Although these studies show plastic and

evolutionary changes, the co-occurrence of other environmental stressors, such as

predation, with temperature might interact in natural populations. Fish predation and

temperature impose selection on many of the same traits, and in the same direction [18,20].

For instance, *Daphnia* often mature at smaller sizes and younger ages in warmer water and

when fish are present, yet little is known about the molecular basis of this response among

populations or their potential interactive effects.

We collected *Daphnia pulicaria* from two lakes in the Sierra Nevada Mountains,

California, USA with different thermal conditions to conduct an experiment to evaluate the

degree to which life history, thermal tolerance and gene expression are influenced by

temperature and predation. We predicted that warm temperatures and exposure to predators

should produce similar responses in life history traits, then used transcriptomics to ask

whether the same genes underlie the plastic response to fish kairomones and warming, or

if the similar phenotypic effects arise from pleiotropic effects where altered gene regulation

affects multiple phenotypic traits. Our goal was to ask whether *D. pulicaria's* plastic

response to one selective agent (fish or warming) magnified or dampened the response to

the other in terms of both phenotype and gene expression.

Materials and methods

Lake sampling

Gardisky (37.955774, -119.251198) and Blue Lakes (38.051164, -119.270342),

located in Inyo National Forest, California, USA, were sampled in late August and early

September 2017, respectively. In both lakes we measured temperature throughout the water

column at each meter using a field probe (YSI Incorporated, Yellow Springs, Ohio, USA).

We also collected live zooplankton from the deepest point of the lake using a 30 cm

diameter, 63 μm mesh conical net with a 1 m length through the water column, starting 1

m above the lake bottom. These samples were kept cold until returning to the laboratory,

where we searched for *Daphnia pulicaria*. When present, 30 *D. pulicaria* females carrying

eggs in the brood pouch were separated in 50 ml centrifuge tubes filled with COMBO

medium [21]. Each *Daphnia* individual was considered a maternal line. Each maternal line

was cultured for at least twelve generations in separate 50 mL tubes filled with COMBO

medium under ambient lab conditions. All *D. pulicaria* were fed non-viable cells of the

green alga *Nannochloropsis* sp. (Brine Shrimp Direct, Ogden, Utah, USA) at a constant

high rate of 24×10^6 cells every two days.

Experimental design

We randomly picked one clonal line from each lake to propagate. *Daphnia*

pulicaria reproduces asexually under benign conditions. Thus, all individuals propagated

asexually in the experiment. We started by randomly picking one mature female from each

clone. Upon the release of the second clutch, we isolated three neonates that were separated

and moved to separate 100 mL containers containing the same media and algae. All

individuals were transferred to fresh media and algae three times a week and reared at 15

$\pm$ 1°C and photoperiod 12:12 h light/dark. Propagation continued until 100 neonates were

collected for each of the clonal lines, with all individuals used for further study born within

a 12-hour period. All neonates were placed into 100 ml jars containing COMBO medium

at a density of 3 *Daphnia*/jar [21]. Each jar was randomly allocated to one of four

treatments: (1) optimum temperature (15°C) without kairomones (fish cues), (2) high

temperature (25°C) without fish cues, (3) optimum temperature with fish cues, and (4) high

temperature with fish cues. An optimum temperature near 15°C is consistent with previous

reports of thermal preference in *Daphnia pulicaria* [24], although there is variation in

tolerable temperatures between *D. pulicaria* genotypes [25]. All *Daphnia* were transferred

to fresh medium, with algae (and kairomones in the fish cue treatment) daily. We

monitored jars daily for maturation (i.e., release of first clutch into the brood chamber).

Upon reaching maturity, individuals were either preserved in RNAlater (Qiagen, USA) and

kept in -20°C for subsequent RNAseq analyses or assigned to phenotypic assay (see below).

Since RNA was extracted from whole individuals, no food was added during the last 12

hours before sampling to minimize algal RNA contamination (most of which will be

digested and hence degraded after 12 hours). The period without food was kept short to

minimize starvation-dependent gene regulation.

Kairomone collection

COMBO medium conditioned by the presence of planktivorous fish was collected

daily from a tank containing 5 juvenile rainbow trout (*Oncorhynchus mykiss*; ~ 5 cm in

total length) in 72 L of COMBO. Each day, media containing fish chemical cues was

filtered through 0.7 μm mesh membrane filters and added at a concentration of 0.007 fish/L

to the predator treatments. This concentration of kairomone media is above the threshold

used by other studies to elicit a phenotypic response, such as the concentration of 5 μl of

kairomone media per 100 ml of solution [26].

D. pulicaria predation by *O. mykiss* in artificially stocked lakes is well established,

specifically predation by juvenile rainbow trout [27,28], *O. mykiss* is one of the most

abundant species for stocking previously fishless lakes in the Sierra Nevada mountains, the

source of our *Daphnia* samples [29]. *O. mykiss* were fed TetraMin Tropical Flakes (Tetra,

Blacksburg, Virginia, USA) prior to kairomone collection, rather than juvenile conspecific

daphniids as in some other studies [30,31]. We selected this method to minimize the

presence of conspecific alarm cues in the isolated kairomone media, so that we could

measure the phenotypic and genetic responses to only predator-released, diet agnostic

kairomone cues. Such research strategies are supported by Mitchell et al. [32], who call for

further research to determine the relationship between predator odor and diet-based cues

in aquatic systems. Following their nomenclature recommendations, we define the

kairomones used in this study to be predator odor cues, as opposed to alarm cues released

by injured conspecifics or diet cues released by predators after digesting conspecifics. All

procedures involving animals were reviewed and approved by the Institutional Animal

Care and Use Committee of the University of California San Diego (IACUC protocol

number: S14140).

Phenotype assays

Individuals from the experimental generation assigned to the phenotypic assay were

scored for the following life-history variables: age and size at maturity, and age and number

of offspring from the first three clutches. These data were used to calculate intrinsic

population growth rate for each maternal line following the Lotka–Euler equation [33]. We

also measured critical maximum temperature (CT_{max}) using a heat ramping assay. After a

30-minute resting period at ambient temperature, *Daphnia* from the four treatments were

transferred to 0.5ml Eppendorf tubes and placed in a thermal heater (4x6 thermal heater,

Corning Digital Dry Bath Heater Dual Block). The water temperature increased 0.1°C

every 20 seconds. We continuously monitored the state of individuals and recorded the

temperature when each individual *Daphnia* lost swimming ability and sank to the bottom

of the tube. The temperature when *Daphnia* became immobilized was used as a proxy for

CT_{max} [17]. We measured life history traits of 11 to 13 individuals per treatment per lake

and CT_{max} of 10 to 12 individuals per treatment per lake. In total, we scored phenotypes of

186 individuals.

Statistical analysis for phenotype

We log-transformed all continuous variables, except intrinsic growth rate, after

visually assessing the probability distribution that best fit the data. We analyzed the effects

of the treatments using generalized linear models for each trait using the statistical software

R [34]. Temperature treatment, fish cue treatment, source lake and the interactions of these

variables were modeled as fixed effects for age and size at maturity, average number of

offspring, intrinsic growth rate and CT_{max}. Since we expected that CT_{max} would be

influenced by body size we included an additive effect of size.

Additionally, to test the effects of slopes of the reaction norms, we calculated the

pairwise differences for each dependent variable between each treatment combination

within a lake. We computed pairwise differences using the least square mean values based

on the generalized linear models for each trait using *lsmeans* function in the *lmerTest* R

package [35]. The pairwise difference test was corrected using Tukey's adjustment for

multiple comparisons [36].

RNA isolation, library preparation and sequencing

We extracted RNA using the TRI Reagent Protocol (Sigma-Aldrich, USA) from

samples preserved in RNAlater (Qiagen, USA). Any remaining genomic DNA was

removed using the TURBO DNA-free Kit (Invitrogen, USA). Each treatment included five

biological replicates, with each replicate comprised of 35 clonal *Daphnia* individuals.

Extracted RNA was stored in -80°C until sequencing. Quality of the isolated RNA was

assessed using RNA Nano 6000 Assay Kit of the Agilent Bioanalyzer 2100 system (Agilent

Technologies, CA, USA) and mRNA-seq libraries were constructed using NEBNext

UltraTM RNA Library Prep Kit for Ilumina (NEB, USA) following manufacturer's

recommendations. A total amount of 1 μg RNA per replicate was used as input material

for RNA-seq library preparations. Samples that did not reach 1 μg RNA were pooled with

a biological replicate grown under the same conditions, a process described by Takele

Assefa et al. [37], leaving 18 replicates from Blue Lake and 16 replicates from Gardisky

Lake. Index codes were added to attribute sequences to each sample and all 34 libraries

were clustered on a cBot System using the PE Cluster kit cBot-HS (Illumina). After

generating the clusters, libraries were sequenced using the Illumina HiSeq 2000 platform

and 150 bp paired-end reads were generated.

Read cleaning and transcriptome comparison

The raw reads were preprocessed using fastp v. 0.23.1 [38] with the command line

options "-x -f 18 -F 18". These options enabled polyX tail trimming and removed 18 low-

quality bases from the beginning of all forward and reverse reads, respectively. To check

for contamination in our read set, we used Kraken2 v. 2.0.9 [39] with a taxonomy database

constructed from the NCBI nr database [40] as of August 2021. A random subset of

1,000,000 read pairs was used for each Kraken2 run per sample. All downstream analyses

were based on the remaining cleaned, validated data.

Separate *de novo* assemblies for were performed on the Blue and Gardisky samples

using Trinity v. r20140413p1 [41] with parameter min_kmer_cov set to 2. Transcriptome

completeness was assessed using BUSCO v. 5.2.2 [42] with the arthopoda_odb10 lineage

dataset, and also by comparison to the following published, annotated genomes: *D.*

pulicaria LK16 [43], *D. pulex* TCO [44], *D. galeata* M5 [45], *D. magna* SK [46] and

Eulimnadia texana JT4(4)5-L [47].

Differential expression analysis

Relative expression levels of annotated genes were estimated by independently

mapping RNA-Seq reads from each lake sample back to the annotated *D. pulicaria* LK16

gene set using STAR v. 2.7.9a [48]. A read count matrix was generated from these

alignments using featureCounts v. 2.0.1 [49]. Genes with read counts below 10 across all

conditions were filtered from the remaining pipeline. The significance of difference in gene

expression of each treatment for each genotype was determined with the R package

DESeq2 v. 1.32.0 [50], using a model based on negative binomial distribution. Statistical

significance (p-value) was adjusted using the q-value obtained from the false discovery

rate [51], with a q-value < 0.05 and |log2(foldchange)| > 1 set as the threshold for

significant differential expression. Counts of differentially expressed genes across

treatments were visualized using the R package VennDiagram v. 1.7.1 [52]. A principal

component analysis (PCA) was visualized using read counts corrected through the variance

stabilizing transformation [53] function of DESeq2, The PCA was generated using

corrected read counts from every gene, rather than the DESeq2 default of using only the

500 most variable genes. These correct read counts were also used to perform a

PERMANOVA test [54] through the R package vegan v. 2.5-7 [55] with 999 permutations.

The groups tested with PERMANOVA were based on lake, temperature, fish cues, and

their interactions terms.

Gene Ontology (GO) enrichment analyses were performed to determine whether

significantly differentially expressed gene sets were enriched for any biologically relevant

functional categories [56]. GO term enrichment lists were generated using GOseq v. 1.44

[57] for the following one-factor comparisons: Blue, temperature rise with kairomones;

Blue, temperature rise without kairomones; Gardisky, temperature rise with kairomones;

Gardisky, temperature rise without kairomones. Further control of the False Discovery

Rate was performed using the Benjamini–Hochberg method [58], and the cutoff for

enriched GO terms was an adjusted p-value less than 0.05.

Orthology of differentially expressed genes

Daphnia pulex genes previously demonstrated to be differentially expressed in

response to temperature and/or predator kairomones were accessed from the *Daphnia*

Stressor Database [59] as of September 2021. Predicted protein sequences for these genes

were downloaded from wFleaBase [60]. These genes were compared to the annotated

Daphnia pulicaria LK16 genome using OrthoVenn2 [61] with an e-value of 1e-5.

OrthoVenn2 results were visualized using the R package VennDiagram v. 1.7.1 [52].

Results

Phenotype

Several traits in the two clones responded differently to temperature and fish cues

(Table 1). The Gardisky Lake clone matured at a younger age than Blue Lake (Lake: $p <$

0.001, Table 1), but age at maturity was earlier for both clones in the presence of fish cues

at high temperature (Temp x Fish: p-value = 0.006, Table 1, Fig 1). The post-hoc test (least

square means) showed that fish had an effect only on the Blue Lake clone reared at 25°C.

At 25°C, Blue Lake replicates reared with fish cues matured 0.55 ± 0.16 days earlier

compared to replicates without fish cues (Fig 1).

Factor	Df	Deviance	Deviance residuals	Pr (>Chi)
Age at maturity				
Temperature	1, 184	16.0918	3.1609	**< 0.001**
Fish cues	1, 183	0.1340	3.0270	**0.002**
Lake	1, 182	0.3961	2.6309	**< 0.001**
Temp x Fish	1, 181	0.1036	2.5273	**0.006**
Temp x Lake	1, 180	0.0020	2.5253	0.7098
Fish x Lake	1, 179	0.0147	2.5106	0.3074
Temp x Fish x Lake	1, 178	0.0011	2.5096	0.7800
Size at maturity				
Temperature	1, 182	0.1078	0.9068	**< 0.001**
Fish cues	1, 181	0.00002	0.9068	0.9349
Lake	1, 180	0.2726	0.6341	**< 0.001**
Temp x Fish	1, 179	0.0054	0.6286	0.1989
Temp x Lake	1, 178	0.0382	0.5904	**< 0.001**
Fish x Lake	1, 177	0.0046	0.5858	0.2368
Temp x Fish x Lake	1, 176	0.0024	0.5833	0.3917
Average number of offspring				
Temperature	1, 87	2.7408	9.7072	**< 0.001**

Fish cues	1, 86	0.0403	9.6669	0.3436
Lake	1, 85	5.7215	3.9454	**< 0.001**
Temp x Fish	1, 84	0.0105	3.9349	0.6290
Temp x Lake	1, 83	0.0826	3.8524	0.1751
Fish x Lake	1, 82	0.0030	3.8494	0.7963
Temp x Fish x Lake	1, 81	0.2120	3.6374	**0.0297**
Intrinsic growth rate				
Temperature	1, 87	0.6194	0.0962	**< 0.001**
Fish cues	1, 86	0.0007	0.0954	0.3087
Lake	1, 85	0.0327	0.0627	**< 0.001**
Temp x Fish	1, 84	0.0000	0.0627	0.9713
Temp x Lake	1, 83	0.0003	0.0624	0.5032
Fish x Lake	1, 82	0.0003	0.0620	0.4853
Temp x Fish x Lake	1, 81	0.0014	0.0606	0.1665
Critical maximum temperature (CT$_{max}$)				
Size at maturity	1, 91	0.0003	0.0256	0.1655
Temperature	1, 90	0.0082	0.0173	**< 0.001**
Fish cues	1, 89	0.0009	0.0163	**0.0125**
Lake	1, 88	0.0004	0.0159	0.1112
Temp x Fish	1, 87	0.0015	0.0144	**0.0019**
Temp x Lake	1, 86	0.0006	0.0138	0.0523
Fish x Lake	1, 85	0.0003	0.0134	0.1356
Temp x Fish x Lake	1, 84	0.0000	0.0134	0.5433

Table 1. Results of the generalized linear model analysis of the life-history traits of

Daphnia pulicaria.

Clonal lines were collected from Blue and Gardisky Lakes and reared under different

temperature (15°C or 25°C) and fish cue treatments (presence or absence).

Fig 1. Predation cues influence critical maximum temperature of Daphnia pulicaria.

Means ± 1 S.E.M. of age at maturity (days), size at maturity (mm), average number of

offspring of the first three clutches, intrinsic growth rate (r), and critical maximum

temperature (CTmax, °C) of Daphnia pulicaria collected at Blue and Gardisky Lakes

during summer 2017 in response to temperature (15°C and 25°C, x-axis) and predation

cues treatments. Circles show response without fish cues and triangles response to fish

cues.

Temperature affected size at maturity differently in the two populations (Temp x

Lake: $p < 0.001$, Table 1) but fish cues had no main effect across clones (Fish: $p = 0.093$,

Table 1, Fig 1). Post hoc tests revealed that the Blue Lake clone matured at 1.26 ± 0.006

mm regardless of temperature, while Gardisky matured at 15°C with 1.31 ± 0.01 mm

compared to 1.41 ± 0.01 mm at 25°C.

The average number of offspring for the first three clutches showed a significant

three-way interaction (Temp x Fish x Lake: $p = 0.029$, Table 1, Fig 1). For the Blue Lake

clone, offspring number only varied with temperature; on average Blue had 1.1 ± 0.11 more

offspring at 25°C compared to 15°C. In contrast, the effect of temperature on fecundity

was greater for the Gardisky Lake clone when fish cues were present. Individuals had 1.67

± 0.30 more offspring at 25°C than 15°C in the presence of fish cues, while without fish

cues the difference between 25°C and 15°C response was 0.98 ± 0.23 offspring.

The intrinsic growth rate was significantly affected by the main effects of

temperature and lake (Temp: $p < 0.001$ and Lake: $p < 0.001$, Table 1, Fig 1). At each

temperature, the Gardisky Lake clone had higher intrinsic growth rates compared to the

Blue Lake clone. For instance, the Gardisky clone had an $r = 0.15 \pm 0.001*day^{-1}$ at 15°C,

while Blue had $r = 0.11 \pm 0.004*day^{-1}$. At 25°C, the intrinsic growth rate was $0.32 \pm$

$0.006*day^{-1}$ for the Gardisky Lake clone, while the growth rate for the Blue Lake clone

was $0.28 \pm 0.007*day^{-1}$.

CT_{max} increased significantly when *Daphnia* were reared at 25°C compared to

15°C. CT_{max} was significantly affected by the interaction between temperature and fish

cues (Temp x Fish: $p = 0.002$, Table 1) and showed a marginally significant interaction

between temperature and lake (Temp x Lake: $p = 0.052$, Table 1), but no main effect of

lake. Post hoc tests showed that at 15°C, fish cues had no effect on CT_{max}. The thermal

tolerance of the Gardisky clone increased by 0.8 ± 0.12°C in the presence of fish cues,

while the Blue Lake clone increased by 0.3 ± 0.14°C at 25°C.

Gene expression

For the Blue Lake clone, sequencing produced on average 42,306,315 raw pair-ended reads

for each sample. Across all Blue Lake samples, 79.4% of the cleaned reads (after removing

adaptor sequences, low-quality and ambiguous sequences) were mapped to the D. pulicaria

LK16 primary gene set. The Gardisky clone produced on average 44,025,859 raw pair-

ended reads for each sample and 80.0% of the cleaned reads mapped to the D. pulicaria

LK16 primary gene set. A breakdown of read mapping rate and estimated read set

contamination by sample is visualized in S1 Fig. *De novo* assemblies of the Blue and

Gardisky clones show sequence similarity to previously annotated *Daphnia* genomes (S2

Fig) and high levels of completeness consistent with previously annotated arthropod

genomes (S3 Fig). A PCA plot generated using DESeq2-corrected read counts (Fig 2)

shows clear divergence in the gene expression between clones under all growth conditions.

A PERMANOVA test of gene expression data shows that gene expression was affected by

both source lake (Pseudo F = 9.6582, R^2 = 0.22284, p = 0.001) and temperature treatments

(Pseudo F = 4.367, R^2 = 0.10078, p = 0.001), but gene expression was not significantly

different between predation treatments (Pseudo F = 0.6706, R^2 = 0.01547, p = 0.746).

Fig 2. Genotype, not treatment group, is the dominant source of differential gene

expression.

Principal component analysis of gene expression patterns generated through DESeq2

prominently separates individuals by source lake along the first principal component.

Temperature treatments (25N vs. 15N and 25Y vs. 15Y) produced the greatest

number of differentially expressed genes (DEGs) for both the Blue and Gardisky clones

(Fig 3). A total of 320 genes were differentially expressed between one-factor comparisons

for the Blue Lake clone: 95 due to temperature differences with predation cues, 292 due to

temperature differences without predation cues, and 4 due to fish cues at 25°C (Fig 3a).

For the Gardisky Lake clone, we identified 575 DEGs. In total, 396 DEGs were specific to

25N vs. 15N and 52 DEGs were specific to 25Y vs. 15Y. These two treatment comparisons

shared 127 genes (Fig 3b). Predator cues alone had no measurable effect on gene

expression when comparing between Gardisky clones grown under the same temperature.

Tables listing all differentially expressed genes are presented in Supplementary Tables 1a

and 1b.

Fig 3. Many genes are differentially expressed in response to temperature changes,

while kairomones have minimal observable effect at the level of transcript

expression.

Venn diagram of differentially expressed genes for *Daphnia pulicaria* from **(a)** Blue and

(b) Gardisky Lakes in response to temperature and predation cues. Cyan and purple

sections illustrate differential expression from the introduction of predator cues (Y vs. N

for the presence and absence of fish cues), while the green and orange sections illustrate

differential expression due to temperature change. Treatment abbreviations: 15Y, 15°C

with kairomones; 15N, 15°C without kairomones; 25Y, 25°C with kairomones; 25N,

25°C without kairomones.

Comparisons of significant GO terms for differentially expressed genes were

difficult to interpret due to the limited specificity of the functional information they provide.

Genes related to proteolysis, chitin turnover, and vision appeared to be upregulated at

higher water temperatures, but no significantly over- or under-represented GO terms were

detected in response to kairomones. GO terms for phototransduction and visual perception

were enriched in all temperature comparisons, while protease and peptidase activity were

enriched in every comparison but 25Y vs. 15Y in the Blue clone. Meanwhile, terms

associated with chitin binding were enriched in every comparison but 25Y vs. 15Y in the

Gardisky clone. Complete lists of all enriched GO terms are given as Supplementary Tables

2a-2d.

Gene orthology

From the Daphnia Stressor Database, we retrieved 386 genes that responded to

kairomones in *D. pulex* across 7 studies, and 2660 genes that responded to temperature

changes in *D. pulex* across 4 studies. 315 of these previously documented kairomone-

responsive genes (81.6%) have orthologs in the *D. pulicaria* LK16 genome, while 1796 of

the temperature-responsive genes (67.5%) are orthologous to genes in *D. pulicaria* (Fig

4a). 35% of the temperature responsive *D. pulicaria* DEGs in this study are orthologous to

previously reported temperature-responsive genes from *D. pulex*, but only one of the four

genes in the Blue clone that responded to kairomones was orthologous to a previously

described kairomone-sensitive gene (Fig 4b). On the other hand, we observed that 18% of

described kairomone-responsive genes in *D. pulex* have orthologs that respond to

temperature in *D. pulicaria*, consistent with earlier phenotypic observations (Fig 1)

showing an interactive relationship between these variables. Lists of gene identifiers for

orthologous clusters and singleton genes are available in Supplementary Files 1 and 2.

Fig 4. Gene orthology analysis reveals similarities between stress-responsive genes in

***D. pulicaria* and *D. pulex*.**

Venn diagrams of orthologous gene clusters between *D. pulex* DEGs from the *Daphnia*

Stressor Database and **(a)** the *D. pulicaria* genome or **(b)** DEGs from *Daphnia pulicaria.*

Numbers indicate protein families with at least two members, excluding "singletons" that

have no orthologous or paralogous matches. Gene identifiers for orthologous clusters and

singleton genes are available in **(a)** Supplementary File 1 and **(b)** Supplementary File 2.

Discussion

Daphnia clonal lines that were stressed by temperature and/or predator cues

exhibited similar phenotypic responses associated with distinctly different patterns of gene

regulation. Temperature was the most important factor driving changes in gene expression,

as most of the DEGs in both clones responded to this variable, versus only a handful to

kairomone cues. Yet fish cues magnified the effect of 10°C of warming on age of maturity

and increased the critical maximum temperature at which *D. pulicaria* lose motor control

when individuals were reared at high temperatures, but not at their optimum temperature.

Genetic pathways that underlie the plastic response to predators and temperature may be

subject to different regulatory pressures, but phenotypic responses suggest that exposure

to predator cues may increase thermal tolerance and affect life history response to warming.

Phenotypic responses

Both clones matured at a younger age and produced more offspring when reared

in high temperature, thus having a higher intrinsic growth rate compared to lower

temperature conditions (Fig 1). Indeed, high temperature increases metabolic rate, often

leading to a larger reproductive investment and faster development [62,63]. In addition, we

found an interactive effect of temperature and predation on age at maturity and critical

maximum temperature.

Our finding that individuals reared with predator cues had higher thermal

tolerance and matured younger at 25°C than those reared without predator cues agree

with other studies that also found synergistic effect of these two stressors in *Daphnia* and

other aquatic organisms [64–69]. In *D. pulex*, after seven generations, Tseng and

O'Connor observed increased thermal plasticity in individuals reared in higher

temperatures only when they were also reared with predators [64]. Zhang et al. tested

resurrected *D. magna* population from time periods without fish and from a high-density

fish period [65]. They found that *Daphnia* that coexisted with fish exhibited earlier

maturation in high temperatures compared to *Daphnia* from the pre-fish period. Earlier

maturation under predation risk is common in size-dependent predation of *Daphnia*

where predators, such as fish, prefer larger prey items, inducing a smaller size at maturity

and earlier age of first reproduction [63]. Our results agree with these studies showing

that fish cues magnify the life history response to temperature and show that exposure to

fish predation can increase the tolerance of stressfully high temperatures.

Genetic responses to temperature

We observed upregulation of serine peptidases in response to high temperature

treatment for both clones. These enzymes are the most important digestive proteases in *D.*

magna [70]. They are involved in proteolysis in the digestive system, the process by which

peptide bonds in proteins are broken generating free amino acids. This upregulation may

indicate a necessity to accommodate higher feeding rates caused by increased metabolism

in high temperatures [62,63]. Moreover, the *D. pulex* genome contains many peptidase

genes, which might indicate adaptation to high variation in food availability in aquatic

environment [71].

Genes related to visual perception were also upregulated in response to high

temperature treatment in both clones. Vision has previously been associated with vertical

migration, because visible light is a proxy for both *Daphnia's* visibility to predators and

the amount of incoming photodamage due to UV radiation at near-surface depths [72].

Vertical migration can serve as either a predator avoidance mechanism induced directly

from kairomones or a diel behavior triggered by temperature change [72]. Higher

temperatures can serve as a proxy for shallower depths, and therefore more incoming

radiation. This may explain why our dataset shows enrichment of phototransduction

genes as a response to temperature while previous studies have observed

phototransduction gene enrichment in response to kairomones [31].

The missing kairomone response: behavioral changes and proximate cues

In total, we only detected 4 DEGs across two *Daphnia pulicaria* clones in response

to kairomones from *Oncorhynchus mykiss.* Such a low number of DEGs related to

kairomone exposure is unexpected, but not unprecedented in previous *Daphnia* studies.

One study on *Daphnia galeata* M6 by Tams et al. also found 4 differentially expressed

transcripts in response to kairomones from the fish *Leuciscus idus* using a two-factor

analysis [73], and another study found no DEGs in *Daphnia magna* Iinb1 when exposed to

kairomones from stickleback fish [74]. Tams et al. hypothesize that life history changes

might be associated with only a few genes compared to morphological defenses, or that

regulation could be post-translational and therefore would not appear in an RNA

sequencing study [73]. Defense strategies can also include behavioral changes such as

vertical migration [75,76] and the transition into a state of alertness [77], which RNA

expression data may not fully capture. In contrast, morphological changes such as

neckteeth formation have been associated with far more genes, such as 230 DEGs found in

D. pulex in response to *Chaoborus* kairomones [78].

Miehls et al. find that the spiny water flea *Bythotrephes longimanus* displays no

changes in life history or morphology when exposed to kairomones from the predator

Perca flavescens, but increased temperature does induce morphologic defense against

predation [79]. The authors hypothesize that temperature is a stronger predictor of

predation risk than kairomones. This is because adult *P. flavescens* are present year-round

but only juveniles are gape-limited in their consumption of *B. longimanus,* such that the

morphologic response of increased tail length is only effective in discouraging predation

from juveniles and not adult fish. If there is no difference between the kairomones released

from juvenile and adult *P. flavescens,* then temperature could serve as a proximate cue for

predation risk that could be damped through some phenotypic response.

The interactions between *B. longimanus* and *P. flavescens* closely mirror predation

of *D. pulicaria* by *O. mykiss*, as *O. mykiss* are among the most abundant stocked fished in

the Sierra Nevada lakes [29] and thus *D. pulicaria* are exposed to their kairomones year-

round. Juvenile *O. mykiss* prey on *Daphnia* as a large percentage of their diet, while trout

larger than 350 mm are primarily piscivorous [27]. A morphological defense response

targeted solely at juvenile trout might require either a juvenile-specific odor, a dynamic

response based on cue concentration, or some additional proxy for *O. mykiss* life stage that

would suggest a high concentration of juveniles. Riessen and Gilbert propose that *D. pulex*

may use water temperature to serve as a proximate cue for predator densities [80], so water

temperature as a proxy for predator life stage is not implausible. Our dataset found some

overlap between kairomone-responsive genes in *D. pulex* and temperature-responsive

genes in *D. pulicaria,* so an intermediate defensive response may be inducible through heat

alone as found by Suppa et al. [81].

However, behavior-based predator avoidance strategies may not require such

specificity. *O. mykiss* can produce the kairomone 5α-cyprinol sulfate in their bile without

consuming *Daphnia*, and this compound induces diel vertical migration in *D. pulex* [75].

Vertical migration still has an associated tradeoff in fitness [82], but this defense might be

preferred when food sources are abundant and temperature gradients are tolerable [83].

Another study found that *D. pulicaria* lineages that coexist with *O. mykiss* vertically

migrate without kairomone exposure, while *D. pulicaria* lineages without prior contact

with *O. mykiss* require kairomone exposure to induce diel vertical migration [84]. Thus,

some environments with high predation might select for *Daphnia* genotypes that respond

to predation primarily through behavioral responses, such as vertical migration.

The missing kairomone response: predator- and diet-specific cues

An alternative explanation for the lack of DEGs may lie in the subtle differences

between kairomone sources. Some kairomones are innately released by the predator

(predator odors), from damaged prey (alarm cues), and from digested prey excreted by the

predator (dietary cues). This study and Tams et al. explicitly avoided the addition of

conspecific alarm cues in our kairomone mixture as to measure only the effect of predator

odors [73]. Other studies, such as [30,31], tested responses to a mixture of predator-specific

odors and alarm cues released by consumed conspecifics. Mitchell et al. call for the

separation of nomenclature between cues released from the predator regardless of diet and

diet-specific cues [32], so that researchers can more accurately discern the independent

effects of these different kairomone compounds.

Previous research on the interaction between predator odors and alarm cues is

mixed. Some studies propose that defense systems rely on both types of compounds to

induce measurable defense responses [85,86], while others observe that alarm cues by

themselves can induce separate responses than predator odors alone [87,88]. Alarm cues

are thought to be nonspecific, as *Daphnia* respond differently to different predators [89]

and alarm cues alone might be unable to induce defense responses effective against a

specific predator [90]. The most effective defense responses may therefore require predator

odors to identify the type of predator combined with alarm cues to induce an immediate

response. A required mixture of predator odors and alarm cues for maximum observed

response in *D. pulicaria* would succinctly explain why we observed such a minor response

in our experiment. Results from this study support the conclusion that the detection of

predator odors alone may be insufficient to induce defense responses at the level of gene

expression, at least in the context of *D. pulicaria – O. mykiss* interactions.

Conclusion

Our study highlights that expression patterns of genes differed between *Daphnia*

pulicaria clones while phenotypic responses and interactions were qualitatively similar,

suggesting that different transcriptomic responses can result in similar phenotypes. Other

studies have found distinct gene expression patterns among individuals from different

zooplankton species [91,92]. These results suggest that diverse genetic pathways can give

rise to similar phenotypic plastic responses to environmental stress. We also showed

synergistic interactions between temperature and predation for some traits. Our results

show that *Daphnia* can have similar phenotypes through distinct molecular mechanisms

and the synergistic effects of temperature and predation may represent an important

mechanism for organisms to adapt to a rapidly changing environment. As one of the first

studies incorporating the *Daphnia pulicaria* genome, we hope to lay the groundwork for

future differential expression studies through incorporation into knowledge bases such as

the *Daphnia* Stressor Database. Overall, we find that the interactions between stressors are

heavily genotype dependent, and echo the sentiment that novel research strategies are

necessary [32] to further investigate the chemical signals of predation in *Daphnia*.

Data Availability

548   Read sets generated and/or analyzed during the current study are available on the

Sequence Read Archive (SRA) under BioProject PRJNA698341. The SRA Accession

numbers for the individual paired sets of reads associated with Blue Lake are

SRR13591461,   SRR13591462,   SRR13591463,   SRR13591464,   SRR13591465,

SRR13591466,   SRR13591471,   SRR13591482,   SRR13591485,   SRR13591486,

SRR13591487,   SRR13591488,   SRR13591489,   SRR13591490,   SRR13591491,

SRR13591492, SRR13591493, and SRR13591494, The SRA Accession numbers for the

individual paired sets of reads associated with Gardisky Lake are SRR13591467,

SRR13591468,   SRR13591469,   SRR13591470,   SRR13591472,   SRR13591473,

SRR13591474,   SRR13591475,   SRR13591476,   SRR13591477,   SRR13591478,

SRR13591479, SRR13591480, SRR13591481 ,SRR13591483, and SRR13591484.

Phenotype data, used to generated Fig 1, is included as S3 File.

Acknowledgements

We are grateful to the following people for their assistance during the execution of

this study: Jennifer Leong, Didra Felix, Carol Blanchette, Brent Salzmann, Kim Rose, Scott

Forster, and Josh Kohn. Funding was provided by National Science Foundation DEB grant

(1457737) to JBS, Brazilian Federal Agency CAPES (13768-13-1) graduate scholarship to

HBC and research funding by Frontiers of Innovation Scholars Program of the University

of California San Diego (3-G3056) to HBC. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript. The work

was performed in part at the University of California Valentine Eastern Sierra Reserve.

References

1.   Carlson SM, Cunningham CJ, Westley PAH. Evolutionary rescue in a changing
world. Trends in Ecology & Evolution. 2014;29(9):521–30.

2.   Parmesan C. Ecological and Evolutionary Responses to Recent Climate Change.
Annual Review of Ecology, Evolution, and Systematics. 2006;37(1):637–69.

3.   Urban MC, Bocedi G, Hendry AP, Mihoub J-B, Pe'er G, Singer A, et al. Improving
the forecast for biodiversity under climate change. Science [Internet]. 2016 Sep 9
[cited 2022 Jan 21]; Available from:
https://www.science.org/doi/abs/10.1126/science.aad8466

4.   Gienapp P, Teplitsky C, Alho JS, Mills J a., Merilä J. Climate change and evolution:
Disentangling environmental and genetic responses. Molecular Ecology.
2008;17:167–78.

5.   Merilä J, Hendry AP. Climate change, adaptation, and phenotypic plasticity: The
problem and the evidence. Evolutionary Applications. 2014;7(1):1–14.

6.   DeWitt TJ, Sih A, Wilson DS. Cost and limits of phenotypic plasticity. Trends in
Ecology & Evolution. 1998;13(97):77–81.

7.   Hendry AP, Farrugia TJ, Kinnison MT. Human influences on rates of phenotypic
change in wild animal populations. Molecular Ecology. 2008;17(1):20–9.

8.   Hoffmann AA, Shirriffs J, Scott M. Relative importance of plastic vs genetic factors
in adaptive differentiation: Geographical variation for stress resistance in
Drosophila melanogaster from eastern Australia. Functional Ecology.
2005;19(2):222–7.

9.   Yampolsky LY, Zeng E, Lopez J, Williams PJ, Dick KB, Colbourne JK, et al.
Functional genomics of acclimation and adaptation in response to thermal stress in
Daphnia. BMC Genomics. 2014 Oct 4;15(1):859.

10.   Whitehead A, Crawford DL. Neutral and adaptive variation in gene expression.
2007;103(May).

11.   Aubin-Horth N, Renn SCP. Genomic reaction norms: using integrative biology to
understand molecular mechanisms of phenotypic plasticity. Molecular ecology.
2009 Sep;18(18):3763–80.

12.   Via S, Lande R. Genotype-Environment Interaction and the Evolution of Phenotypic
Plasticity. Evolution. 1985;39(3):505–22.

13.   Sørensen JG, Kristensen TN, Loeschcke V. The evolutionary and ecological role of
heat shock proteins. Ecology Letters. 2003;6(11):1025–37.

14.   Huang Y, Agrawal AF. Experimental Evolution of Gene Expression and Plasticity
in Alternative Selective Regimes. PLOS Genetics. 2016;12(9):e1006336.

15.   Kenkel C, Matz M V. Enhanced gene expression plasticity as a mechanism of
adaptation to a variable environment in a reef-building coral. Nature Ecology and
Evolution. 2016;1(3):059667.

16.   Miner BE, De Meester L, Pfrender ME, Lampert W, Hairston NG. Linking genes to
communities and ecosystems: Daphnia as an ecogenomic model. Proceedings of the
Royal Society B: Biological Sciences. 2012;279(1735):1873–82.

17.   Geerts AN, Vanoverbeke J, Vanschoenwinkel B, Van Doorslaer W, Feuchtmayr H,
614   Atkinson D, et al. Rapid evolution of thermal tolerance in the water flea Daphnia.
615   Nature Clim Change. 2015 Jul;5(7):665–8.

616   18.   Williams PJ, Dick KB, Yampolsky LY. Heat tolerance, temperature acclimation,
617   acute oxidative damage and canalization of haemoglobin expression in Daphnia.
618   Evolutionary Ecology. 2012;26(3):591–609.

619   19.   Yampolsky LY, Schaer TMM, Ebert D. Adaptive phenotypic plasticity and local
620   adaptation for temperature tolerance in freshwater zooplankton. Proceedings
621   Biological sciences / The Royal Society. 2014;281:20132744.

622   20.   Stoks R, Govaert L, Pauwels K, Jansen B, De Meester L. Resurrecting complexity:
623   The interplay of plasticity and rapid evolution in the multiple trait response to strong
624   changes in predation pressure in the water flea Daphnia magna. Ecology Letters.
625   2016;19(2):180–90.

626   21.   Kilham SS, Kreeger DA, Lynn SG, Goulden CE, Herrera L. COMBO: a defined
627   freshwater culture medium for algae and zooplankton. Hydrobiologia.
628   1998;377(1):147–59.

629   22.   Van Wagenen J, Miller TW, Hobbs S, Hook P, Crowe B, Huesemann M. Effects of
630   Light and Temperature on Fatty Acid Production in Nannochloropsis Salina.
631   Energies. 2012 Mar;5(3):731–40.

632   23.   Coggins BL, Pearson AC, Yampolsky LY. Does geographic variation in thermal
633   tolerance in Daphnia represent trade-offs or conditional neutrality? Journal of
634   Thermal Biology. 2021 May 1;98:102934.

635   24.   Glaholt SP, Kennedy ML, Turner E, Colbourne JK, Shaw JR. Thermal variation and
636   factors influencing vertical migration behavior in Daphnia populations. J Therm
637   Biol. 2016 Aug;60:70–8.

638   25.   Palaima A, Spitze K. Is a jack-of-all-temperatures a master of none? An
639   experimental test with Daphnia pulicaria (Crustacea: Cladocera). Evol Ecol Res.
640   2004;6(2):215–25.

641   26.   Christjani M, Fink P, von Elert E. Phenotypic plasticity in three Daphnia genotypes
642   in response to predator kairomone: evidence for an involvement of chitin
643   deacetylases. Journal of Experimental Biology. 2016 Jun 1;219(11):1697–704.

644   27.   Beauchamp DA. Seasonal and Diel Food Habits of Rainbow Trout Stocked as
645   Juveniles in Lake Washington. Transactions of the American Fisheries Society.
646   1990 May 1;119(3):475–82.

647   28.   Hembre LK. Effects of a rainbow trout stocking moratorium on the Daphnia species
648   composition and water quality of Square Lake (Minnesota). Lake and Reservoir
649   Management. 2019 Apr 3;35(2):127–39.

650   29.   Knapp RA, Matthews KR, Sarnelle O. Resistance and Resilience of Alpine Lake
651   Fauna to Fish Introductions. Ecological Monographs. 2001;71(3):401–21.

652   30.   An H, Do TD, Jung G, Karagozlu MZ, Kim C-B. Comparative Transcriptome
653   Analysis for Understanding Predator-Induced Polyphenism in the Water Flea
654   Daphnia pulex. International Journal of Molecular Sciences. 2018 Jul;19(7):2110.

655   31.   Zhang X, Blair D, Wolinska J, Ma X, Yang W, Hu W, et al. Genomic regions
656   associated with adaptation to predation in Daphnia often include members of
657   expanded gene families. Proceedings of the Royal Society B: Biological Sciences.
658   2021 Jul 28;288(1955):20210803.

659   32.   Mitchell MD, Bairos-Novak KR, Ferrari MCO. Mechanisms underlying the control
660   of responses to predator odours in aquatic prey. Journal of Experimental Biology.
661   2017 Jun 1;220(11):1937–46.

662   33.   Roff DA. Evolutionary quantitative genetics. New York, NY: Chapman and Hall;
663   1997.

34.   Core Team R. R: A language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria; 2018.

35.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear

mixed effects models. Journal of Statistical Software. 2017;82:1–26.

36.   Tukey JW. The Philosophy of Multiple Comparisons. Statistical Science. 1991

Feb;6(1):100–16.

37.   Takele Assefa A, Vandesompele J, Thas O. On the utility of RNA sample pooling to

optimize cost and statistical power in RNA sequencing experiments. BMC

Genomics. 2020 Apr 19;21(1):312.

38.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018 Sep 1;34(17):i884–90.

39.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2.

Genome Biology. 2019 Nov 28;20(1):257.

40.   Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database

resources of the national center for biotechnology information. Nucleic Acids Res.

2021 Dec 1;gkab1112.

41.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-

length transcriptome assembly from RNA-Seq data without a reference genome. Nat

Biotechnol. 2011 Jul;29(7):644–52.

42.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update:

Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic

Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Molecular

Biology and Evolution. 2021 Oct 1;38(10):4647–54.

43.   Jackson CE, Xu S, Ye Z, Pfrender ME, Lynch M, Colbourne JK, et al.

Chromosomal rearrangements preserve adaptive divergence in ecological speciation

[Internet]. 2021 Aug [cited 2021 Nov 2] p. 2021.08.20.457158. Available from:

https://www.biorxiv.org/content/10.1101/2021.08.20.457158v1

44. Colbourne JK, Pfrender ME, Gilbert D, Thomas WK, Tucker A, Oakley TH, et al.

The Ecoresponsive Genome of Daphnia pulex. Science. 2011 Feb 4;331(6017):555–

61.

45. Nickel J, Schell T, Holtzem T, Thielsch A, Dennis SR, Schlick-Steiner BC, et al.

Hybridization Dynamics and Extensive Introgression in the Daphnia longispina

Species Complex: New Insights from a High-Quality Daphnia galeata Reference

Genome. Genome Biology and Evolution. 2021 Dec 1;13(12):evab267.

46. Lee B-Y, Choi B-S, Kim M-S, Park JC, Jeong C-B, Han J, et al. The genome of the

freshwater water flea Daphnia magna: A potential use for freshwater molecular

ecotoxicology. Aquatic Toxicology. 2019 May 1;210:69–84.

47. Baldwin-Brown JG, Weeks SC, Long AD. A New Standard for Crustacean

Genomes: The Highly Contiguous, Annotated Genome Assembly of the Clam

Shrimp Eulimnadia texana Reveals HOX Gene Order and Identifies the Sex

Chromosome. Genome Biology and Evolution. 2018 Jan 1;10(1):143–56.

48. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

ultrafast universal RNA-seq aligner. Bioinformatics. 2013 Jan;29(1):15–21.

49. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014 Apr

1;30(7):923–30.

50. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome biology. 2014;15(12):550.

51. Storey JD, Tibshirani R. Statistical significance for genomewide studies. PNAS.

2003 Aug 5;100(16):9440–5.

52.   Chen H. VennDiagram: Generate High-Resolution Venn and Euler Plots [Internet].
2021 [cited 2022 Jan 12]. Available from: https://CRAN.R-
project.org/package=VennDiagram

53.   Anders S, Huber W. Differential expression analysis for sequence count data. Nat
Prec. 2010 Mar 15;1–1.

54.   Anderson MJ. Permutational Multivariate Analysis of Variance (PERMANOVA).
In: Wiley StatsRef: Statistics Reference Online [Internet]. John Wiley & Sons, Ltd;
2017 [cited 2022 Feb 8]. p. 1–15. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1002/9781118445112.stat07841

55.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al.
vegan: Community Ecology Package [Internet]. 2020. Available from:
https://CRAN.R-project.org/package=vegan

56.   Gene Ontology Consortium. The Gene Ontology resource: enriching a GOld mine.
Nucleic Acids Res. 2021 Jan 8;49(D1):D325–34.

57.   Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for
RNA-seq: accounting for selection bias. Genome biology. 2010;11(2):R14.

58.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society:
Series B (Methodological). 1995 Jan 1;57(1):289–300.

59.   Ravindran SP, Lüneburg J, Gottschlich L, Tams V, Cordellier M. Daphnia stressor
database: Taking advantage of a decade of Daphnia '-omics' data for gene
annotation. Sci Rep. 2019 Jul 31;9(1):11135.

60.   Colbourne JK, Singan VR, Gilbert DG. wFleaBase: the Daphnia genome database.
BMC Bioinformatics. 2005 Mar 7;6:45.

61.   Xu L, Dong Z, Fang L, Luo Y, Wei Z, Guo H, et al. OrthoVenn2: a web server for
whole-genome comparison and annotation of orthologous clusters across multiple
species. Nucleic Acids Res. 2019 Jul 2;47(W1):W52–8.

62.   Stibor H. Predator induced life-history shifts in a freshwater cladoceran. Oecologia.
1992;92(2):162–5.

63.   Riessen HP. Predator-induced life history shifts in Daphnia: a synthesis of studies
using meta-analysis. Canadian Journal of Fisheries and Aquatic Sciences.
1999;56:2487–94.

64.   Tseng M, O'Connor MI. Predators modify the evolutionary response of prey to
temperature change. Biology letters. 2015;11(12):20150798.

65.   Zhang C, Jansen M, De Meester L, Stoks R. Energy storage and fecundity explain
deviations from ecological stoichiometry predictions under global warming and
size-selective predation. The Journal of animal ecology. 2016 Nov;85(6):1431–41.

66.   Luhring TM, DeLong JP. Predation changes the shape of thermal performance
curves for population growth rate. Current zoology. 2016 Oct;62(5):501–5.

67.   Luhring TM, Vavra JM, Cressler CE, DeLong JP. Predators modify the temperature
dependence of life-history trade-offs. Ecology and Evolution. 2018 Sep
1;8(17):8818–30.

68.   Riessen HP. Water temperature alters predation risk and the adaptive landscape of
induced defenses in plankton communities. Limnology and Oceanography. 2015
Nov 1;60(6):2037–47.

69.   Miller LP, Matassa CM, Trussell GC. Climate change enhances the negative effects
of predation risk on an intermediate consumer. Global Change Biology.
2014;20:3834-3844.

70.   von Elert E, Agrawal MK, Gebauer C, Jaensch H, Bauer U, Zitt A. Protease activity
in gut of Daphnia magna: evidence for trypsin and chymotrypsin enzymes.

Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular

Biology. 2004;137(3):287–96.

71. Colbourne JK, Eads BD, Shaw J, Bohuski E, Bauer DJ, Andrews J. Sampling

Daphnia's expressed genes: preservation, expansion and invention of crustacean

genes with reference to insect genomes. BMC Genomics. 2007;8(1):217.

72. Williamson CE, Fischer JM, Bollens SM, Overholt EP, Breckenridge JK. Toward a

more comprehensive theory of zooplankton diel vertical migration: Integrating

ultraviolet radiation and water transparency into the biotic paradigm. Limnology

and Oceanography. 2011;56(5):1603–23.

73. Tams V, Nickel JH, Ehring A, Cordellier M. Insights into the genetic basis of

predator-induced response in Daphnia galeata. Ecology and Evolution.

2020;10(23):13095–108.

74. Orsini L, Brown JB, Shams Solari O, Li D, He S, Podicheti R, et al. Early

transcriptional response pathways in Daphnia magna are coordinated in networks of

crustacean-specific genes. Molecular Ecology. 2018;27(4):886–97.

75. Hahn MA, Effertz C, Bigler L, von Elert E. 5α-cyprinol sulfate, a bile salt from fish,

induces diel vertical migration in Daphnia. Pohnert G, Baldwin IT, Pohnert G,

editors. eLife. 2019 May 2;8:e44791.

76. Loose CJ, Elert E, Dawidowicz P. Chemically-induced diel vertical migration in

Daphnia: a new bioassay for kairomones exuded by fish. Archiv für Hydrobiologie.

1993;329–37.

77. Diel P, Kiene M, Martin-Creuzburg D, Laforsch C. Knowing the Enemy: Inducible

Defences in Freshwater Zooplankton. Diversity. 2020 Apr;12(4):147.

78. Rozenberg A, Parida M, Leese F, Weiss LC, Tollrian R, Manak JR. Transcriptional

profiling of predator-induced phenotypic plasticity in Daphnia pulex. Frontiers in

Zoology. 2015 Jul 25;12(1):18.

79. Miehls ALJ, McAdam AG, Bourdeau PE, Peacor SD. Plastic response to a proxy
cue of predation risk when direct cues are unreliable. Ecology. 2013;94(10):2237–
48.

80. Riessen HP, Gilbert JJ. Divergent developmental patterns of induced morphological
defenses in rotifers and Daphnia: Ecological and evolutionary context. Limnology
and Oceanography. 2019;64(2):541–57.

81. Suppa A, Caleffi S, Gorbi G, Marková S, Kotlík P, Rossi V. Environmental
conditions as proximate cues of predation risk inducing defensive response in
Daphnia pulex. Biologia. 2021 Feb 1;76(2):623–32.

82. Lampert W. The Adaptive Significance of Diel Vertical Migration of Zooplankton.
Functional Ecology. 1989;3(1):21–7.

83. Stich H-B, Lampert W. Growth and reproduction of migrating and non-migrating
Daphnia species under simulated food and temperature conditions of diurnal vertical
migration. Oecologia. 1984 Feb;61(2):192–6.

84. Hembre LK, Peterson LA. Evolution of predator avoidance in a Daphnia
population: evidence from the egg bank. Hydrobiologia. 2013 Jan 1;700(1):245–55.

85. Ślusarczyk M. Predator-induced diapause in Daphnia magna may require two
chemical cues. Oecologia. 1999 May 1;119(2):159–65.

86. Stabell OB, Ogbebo F, Primicerio R. Inducible Defences in Daphnia Depend on
Latent Alarm Signals from Conspecific Prey Activated in Predators. Chemical
Senses. 2003 Feb 1;28(2):141–53.

87. Schoeppner NM, Relyea RA. Interpreting the smells of predation: how alarm cues
and kairomones induce different prey defences. Functional Ecology.
2009;23(6):1114–21.

88. Ślusarczyk M, Rygielska E. Fish Faeces as the Primary Source of Chemical Cues
Inducing Fish Avoidance Diapause in Daphnia Magna. Hydrobiologia. 2004 Sep
1;526(1):231–4.

89. Boeing WJ, Ramcharan CW, Riessen HP. Multiple predator defence strategies in
Daphnia pulex and their relation to native habitat. Journal of Plankton Research.
2006 Jun 1;28(6):571–84.

90. Nagano M, Doi H. Ecological and evolutionary factors of intraspecific variation in
inducible defenses: Insights gained from Daphnia experiments. Ecology and
Evolution. 2020;10(16):8554–62.

91. Lima TG, Willett CS. Locally adapted populations of a copepod can evolve
different gene expression patterns under the same environmental pressures. Ecol
Evol. 2017 May 9;7(12):4312–25.

92. Roy Chowdhury P, Frisch D, Becker D, Lopez JA, Weider LJ, Colbourne JK, et al.
Differential transcriptomic responses of ancient and modern Daphnia genotypes to
phosphorus supply. Mol Ecol. 2015 Jan;24(1):123–35.

Supplementary Tables, Figures, and Files

S1 Tables 1a-1b. Data associated with Fig 3.

Differentially expressed genes in the **(a)** Blue Lake clone and **(b)** the Gardisky Lake

clone. The comparison column denotes if genes were responsive to temperature and/or

kairomone treatments.

S2 Tables 2a-2d. Gene Ontology of differentially expressed genes.

Upregulated Gene Ontology terms for comparisons between: **(a)** Blue Lake clone, 15Y

and 25Y treatments, **(b)** Blue Lake clone, 15N and 25N treatments, **(c)** Gardisky Lake

clone, 15Y and 25Y treatments, and **(d)** Gardisky Lake clone, 15N and 25N treatments.

S1 Fig. Read-based taxonomy assignment consistent with mapping rates to the

***Daphnia pulicaria* genome.**

Panel **(a)** shows Kraken2 read taxonomy on a random subset of one million cleaned

reads from each sample. Panel **(b)** plots the alignment rate of the cleaned reads from each

sample to the chromosomes of *Daphnia pulicaria* LK16 using the STAR aligner.

S2 Fig. Blue and Gardisky transcripts are highly similar to *Daphnia pulicaria* LK16.

Results of a two-way average nucleotide identity (ANI) matrix and associated

dendrogram for *the de novo* assembled transcriptomes from Blue and Gardisky Lakes,

predicted genes from *Daphnia* species with annotated genomes, and the annotated gene

set of the related arthropod *Eulimnadia texana*. Our transcriptome sets are closest to those

from *D. pulicaria*, within the well-described *Daphnia pulex-pulicaria* species complex.

S3 Fig. Blue and Gardisky transcriptomes are highly complete and highly

duplicated.

Bar graph of gene set completeness as determined by BUSCO using the arthropod

marker gene set. Our *de novo* assembled transcriptomes have completeness scores

comparable to or better than genome-derived transcript sets from other *Daphnia* studies.

However, Blue and Gardisky transcriptome sets have additional gene duplication in our

transcriptome due to the inclusion of gene isoforms from *de novo* assembly.

S1 File. Orthologous clusters associated with Fig 4a.

Orthologous clusters and singletons for the gene orthology analysis between the *D.*

pulicaria genome, temperature-responsive genes in *D. pulex*, and kairomone-responsive

genes in *D. pulex*. *D. pulex* genes were collected from the *Daphnia* Stressor Database.

Data is in Excel format.

S2 File. Orthologous clusters associated with Fig 4b.

Orthologous clusters and singletons for the gene orthology analysis between temperature-

responsive *D. pulicaria* genes, kairomone-responsive *D. pulicaria* genes, temperature-

responsive genes in *D. pulex*, and kairomone-responsive genes in *D. pulex*. *D. pulex*

genes were collected from the *Daphnia* Stressor Database. Data is in Excel format.

S3 File. Phenotype and life history data associated with Fig 1.

Figure1

Figure2

Figure3

Figure4