Title: Bimodal breeding phenology in the Parsley Frog *Pelodytes punctatus* as a bet-hedging

strategy in an unpredictable environment despite strong priority effects

Running title: breeding in an unpredictable habitat

Authors:

Hélène JOURDAN-PINEAU[1], Pierre-André CROCHET[2,*], Patrice DAVID[2,*]

Affiliations :

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France.

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France

[2] CEFE, UMR 5175 CNRS, Montpellier, France

*both authors contributed equally

Corresponding author: helene.jourdan@cirad.fr

Keywords: breeding phenology, bet-hedging, priority effects, stochastic environment, anuran

ABSTRACT

When environmental conditions are unpredictable, expressing alternative phenotypes spreads

the risk of failure, a mixed strategy called bet-hedging. In the southern part of its range, the

Parsley Frog *Pelodytes punctatus* breeds both in autumn and in spring. Our aim was to study

the breeding phenology and reproductive success associated with the use of those two seasonal

niches to understand how this breeding strategy can be maintained. Field surveys revealed that

breeding phenology was typically bimodal with a higher breeding effort in autumn. More

importantly, in spring, the survival rate of offspring was severely reduced by the presence of

autumn tadpoles, indicating a clear priority effect. However, the autumn cohort often failed to

survive over winter, in which case spring cohorts were often successful. Based on those results,

we constructed a model in which females can allocate a variable portion of eggs to each season

and added a priority effect. We conclude that the existence of the two breeding seasons may

indeed constitute a bet-hedging strategy.

INTRODUCTION

Breeding phenology is one of the key components of adaptation to temporally variable environments. Temporal dynamics of both the biotic and abiotic environment impose selective constraints on parental development and physiological state (to be able to reproduce) as well as offspring survival (at the various developmental stages until they reach sexual maturity and start to reproduce) (Rand 1973). There is a vast amount of literature on intraspecific variation of breeding patterns in relation to environmental conditions, in particular latitude, altitude and climate. In the context of current climate change, many species in temperate regions have advanced their breeding time (e.g. Brown et al., 2016; Frederiksen et al., 2004; Møller, 2008), as a result of microevolutionary changes and/or of phenotypic plasticity (Charmantier & Gienapp 2014). Most of these studies concern species with a single reproductive peak in the year, which has to match as precisely as possible a seasonal peak of resource availability in order to maximize reproductive success (e.g. caterpillar availability for tits). The exact date of the resource peak may vary from year to year and species usually rely on cues to anticipate it and plastically delay or advance the onset of reproduction every year. However, in some cases reproductive success depends on even more irregular and/or unpredictable conditions. In such situations, species face the risk of complete reproductive failure at any given breeding attempt, a regime that favors the expression of alternative phenotypes to spread the risk (Cohen 1970; Slatkin 1974; Philippi & Seger 1989; Leimar 2005; Venable 2007).

Theory predicts that in stochastic environments, selection favors life history traits that reduce temporal fitness variation even if they result in lowered arithmetic mean fitness (Philippi & Seger 1989). This risk-spreading strategy is called bet-hedging. In temporally variable environments, the long-term fitness of a genotype is measured by the geometric mean of the fitness contribution over successive generations for a particular genotype (Lewontin & Cohen 1969; Olofsson *et al.* 2009; Simons 2011; Yasui & Yoshimura 2018). This geometric mean

fitness is highly impacted by low values; hence, traits with lower fitness variation may have

higher long-term fitness. In principle, lower variation in fitness can be achieved either using the

same low-risk strategy (conservative bet-hedging), or displaying several strategies, either at

once or over several instances (diversified bet-hedging). While there is an abundant theoretical

literature on bet-hedging, empirical studies have provided limited evidence so far (Simons

2011), and the most comprehensive examples concern the timing of germination/diapause and

the fraction of dormant seeds/diapausing eggs (Venable 2007; Gremer *et al.* 2016; García-

Roger *et al.* 2017; Wang & Rogers 2018). Some studies even report experimental evolution of

bet-hedging traits in response to unpredictable environment (Beaumont *et al.* 2009; Graham *et*

al. 2014; Maxwell & Magwene 2017; Tarazona *et al.* 2017).

One of the best examples of stochastic, unpredictable environments is provided by temporary

ponds, alternating between inundation and drought where each breeding event is a bet as habitat

desiccation can occur before the end of the breeding cycle. Several examples of bet-hedging

occur in temporary ponds. Fairy shrimps (Anostraca) produce drought-resistant eggs showing

asynchronous hatching at different hydroperiods (Saiah & Perrin 1990; Simovich & Hathaway

1997; Wang & Rogers 2018). Similarly, rotifers produce diapausing eggs to overpass

unfavorable planktonic growing season and only a fraction of those eggs hatch when conditions

are suitable (García-Roger *et al.* 2017; Tarazona *et al.* 2017).

For amphibian species breeding in temporary ponds, drought can cause 100% mortality of eggs

or larvae, resulting in complete failure of one breeding event. One way to reduce the risk of

losing a breeding opportunity entirely is to spread this risk at a spatial scale, partitioning broods

into various pools as done by the Neotropical poison frog *Allobates femoralis* (Erich *et al.*

2015). Another bet-hedging strategy could be to split the breeding effort at a temporal scale and

exploit all suitable temporal windows.

From an ecological point of view, such temporal niche partitioning is expected to reduce inter

and intra-specific competition as well as resource depletion (Carothers & Jaksić 1984). For

example, species may share the same habitat but have opposite activity patterns (nocturnal

versus diurnal species), as is the case in Neotropical felid community or in grassland ants

(Albrecht & Gotelli 2001; Di Bitetti *et al.* 2010). The same type of temporal segregation at a

daily scale is also observed within species: brown trout *Salmo trutta* reduces competition for a

limiting resource by sequential use of foraging areas (Alanärä *et al.* 2001). Voltinism in insects

is another well-studied example of temporal partitioning at the annual scale which is an

adaptation to predictable seasonal cycles (Kivelä *et al.* 2013; Zeuss *et al.* 2017; Forrest *et al.*

2019).

However, if successive seasonal cohorts overlap, fitness gains may be asymmetric, because

progeny produced by late breeding may suffer from competition or even predation from earlier

cohorts (Morin 1987; Ryan & Plague 2004; Eitam *et al.* 2005). Those priority effects are often

difficult to disentangle from seasonal effects due to environmental differences experienced by

the temporal cohorts (Morin *et al.* 1990). If priority effects are strong, late breeders may select

breeding sites in order to limit the competition by conspecifics (Halloy & Fiaño 2000; Halloy

2006; Sadeh *et al.* 2009) and this may restrict late breeders to poorer sites (Crump 1991).

In amphibians, this temporal partitioning of breeding activity is thought to regulate community

dynamics through interspecific competition (Lawler & Morin 1993; Gottsberger & Gruber

2004; Richter-Boix *et al.* 2006b, 2007c). Similarly, community composition may depend on

species arrival and priority effects whereby species arriving earlier monopolize available

resources and gain a competitive advantage over late species (Morin *et al.* 1990; Blaustein &

Margalit 1996; Urban & De Meester 2009). In Mediterranean regions, climatic conditions are

characterized by dry, hot summers and mild winters, with the maximum rainfalls in autumn and

spring. This leads to large breeding asynchrony observed between and within amphibian species

(Diaz-Paniagua 1988; Jakob, Poizat et al. 2003; Richter-Boix, Llorente et al. 2006; Vignoli,

Bologna et al. 2007): whereas most species typically breed in spring, some species breed earlier

at the end of winter, and some even breed in autumn in addition to spring.

The Parsley Frog *Pelodytes punctatus* is a small sized anuran distributed in Spain and most of

France (locally reaching neighboring countries). It has a broad ecological niche but has poor

competitive abilities and is sensitive to fish predation (Morand & Pierre 1995; Crochet *et al.*

2004; Richter-Boix *et al.* 2007b); it thus prefers seasonally flooded habitats to large permanent

water bodies (Guyétant *et al.* 1999; Salvidio *et al.* 2004; Richter-Boix *et al.* 2007a). In Spain

the Parsley Frog shows a bimodal breeding pattern with higher reproductive effort in spring

than in autumn (Guyétant *et al.* 1999; Richter-Boix *et al.* 2006b). In France, in addition to spring

breeding, autumnal breeding is also observed in Mediterranean regions and areas with mild

oceanic climate (Guyétant *et al.* 1999; Jakob *et al.* 2003; Richter-Boix *et al.* 2006a; Cayuela *et*

al. 2012) but the importance of autumn versus spring reproduction has not been quantified. In

the rest of the range and at higher altitudes, only spring breeding occurs.

In the Mediterranean areas of southern France, the Parsley Frog uses temporary ponds that refill

in September and October but may dry during autumn or later in late spring. Adults thus have

to deal with very unpredictable environmental conditions for their future offspring. In addition

to this unpredictable risk, tadpoles hatched in autumn or spring are exposed to very different

environmental conditions, the most obvious being that the autumn tadpoles overwinter while

the spring ones do not. This should result in drastically different developmental trajectories but

also in different offspring survivals. Both seasonal cohorts may also interact, leading to a

potential competitive advantage to the earlier cohort over the later, i.e. a priority effect. The

relative success of each breeding period and the outcome of the interaction between cohorts are

key parameters to understand the persistence of this two-peaks breeding strategy. In fact,

several pieces of information are still lacking in order to understand the evolutionary basis of

this seasonally variable breeding strategy. Do we have a single protracted breeding season or a

really bimodal reproduction generated by the coexistence of alternative breeding timing? If so,

what is the relative importance of autumn versus spring reproduction? What is the survival of

offspring produced at the two breeding periods and how is it affected by the presence of

conspecifics? Once this basic knowledge is obtained, it can be fed into theoretical models for

the evolution of mixed breeding strategies.

In this paper, we characterize the breeding phenology (temporal dynamic, relative proportion

of each breeding period) of Parsley Frog in a French Mediterranean area based on results from

a 3-year field survey. We monitored the survival of offspring produced in each season to

estimate the success of this breeding strategy. We also investigated the factors influencing

breeding and tadpole survival, in particular whether there is a priority effect between seasonal

cohorts. Finally, using an analytical model adapted from Cohen (Cohen 1966) we tested

whether the coexistence of the two breeding periods can be interpreted as a bet-hedging

strategy.

MATERIAL AND METHODS

Field survey

The field study was carried out from September 2007 to August 2010 in 19 ponds situated

around Montpellier, southern France (Annex 1). Those ponds are man-made environments,

often dug out to provide drinking water for livestock (sheep and cows) or for game. The ponds

surveyed included temporary and permanent sites. We define here the autumn breeding season

as the period spanning from September to December and the spring breeding season from

January to April. We surveyed each pond twice each month. During each visit, we recorded the

depth of the pond.

Sampling methods

At every visit (mostly diurnal), we looked for newly deposited egg masses throughout the entire water body and classified the egg masses as small, medium and large, corresponding to an average of 75, 150 and 250 eggs per mass, respectively (Salvador & Paris 2001, and personal observation). The Parsley Frog's embryonic period ranges from 5 days at 15°C to 15 days at 10°C (Toxopeus *et al.* 1993). Moreover, embryos stay attached to the jelly for several additional days (Guyétant *et al.* 1999). Thus, with an interval of 15 days between two successive visits, we may have missed a few masses but we have avoided double-counting masses since 15-day old masses can readily be distinguished from new ones based on the developmental stages of the embryos. In only 2% of the larval cohorts produced were small larvae observed in ponds where we did not notice the presence of egg masses before. The probability of detection of an egg mass, even if not perfect, was similar in autumn and in spring.

 We estimated the number of amphibian larvae and invertebrates present in the ponds using 5 to 10 dipnet sweeps (depending on the pond size). The anuran community of the area consists of 7 species: *Pelodytes punctatus*, *Pelobates cultripes*, *Alytes obstetricans*, *Bufo bufo*, *Epidalea calamita*, *Hyla meridionalis*, and *Pelophylax sp*., (*P. ridibundus* and/or *P. perezi* & *P*. kl. *grafi*, depending on the sites). Potential predators of tadpoles are urodeles and aquatic invertebrates. Two urodele species (*Lissotriton helveticus* and *Triturus marmoratus*) were recorded in the ponds but due to the rare occurrence of *Triturus marmoratus*, only *Lissotriton helveticus* was included in subsequent analyses (as adults as well as larvae).

We also surveyed dragonfly larvae (Anisoptera) and backswimmers (Heteroptera, Notonectidae) that are potential predators of tadpoles (Richter-Boix *et al.* 2007a) except during the first year. Diving beetles (Coleoptera, Dytiscidae) are also known to prey on tadpoles but were very rare in the studied ponds and thus not considered for this study.

We divided the total counts for each amphibian larvae and invertebrate predators captured in

each pond by the number of dipnet sweeps taken in each pond. This procedure yielded a crude

proxy for density on the basis of catch per unit effort and could therefore be compared across

localities.

Reproductive effort and offspring survival

Reproductive strategy of Parsley Frog was described by two measures: the *presence of egg*

masses (binary variable: whether some eggs were laid or not when we visited a pond) and the

number of egg masses (integer, non-zero; applies only to cases where egg masses are present).

We normalized the number of egg masses by their size (e.g. a small egg mass equals ½ medium

egg mass).

For each breeding event, we estimated the *hatching rate* as the ratio of the number of small

tadpoles (Gosner stage 26, free swimming tadpole) to the number of eggs spawned. Similarly,

we quantified the *survival rate from egg to metamorph* as the ratio of the number of metamorphs

(Gosner stage 42-43) to the number of eggs spawned. Finally, we calculated the *survival rate*

during larval stage as the ratio of the number of metamorphs over the number of small tadpoles.

This index could only be estimated in about one third of the breeding events when hatching was

successful (i.e. the number of small tadpoles was not null).

The number of tadpoles in a pond was estimated using the mean number of tadpoles caught per

dipnet sweep scaled to a sampling surface of 1 m² (we estimated that one dipnet sweep sampled

a surface of 0.5 m², taking the dipnet size and the length of the haul into account) and then

multiplied by the surface of the pond. This should not be taken as an attempt to estimate

precisely the number of tadpoles present in a pond at a given time but as an index of abundance

that can be compared between ponds and between breeding events. It was sometimes impossible

to follow the larval development and metamorphosis of offspring from a particular breeding

event. Indeed, Parsley Frogs may breed three to four times during each seasonal breeding

In these cases, the successive sub-cohorts produced are indistinguishable after a few weeks, and

we summed the eggs counted in two or three successive visits to evaluate survival from a

combination of breeding events within a given season (and within a site). Survival measures

should be viewed as an index to assess the differences of reproductive success between seasons

as there is no reason to expect any seasonal bias in this index.

Explanatory variables

Explanatory variables for the breeding probability and breeding effort are the season, depth of

the pond as well as the presence of conspecific and inter-specific competitors (larvae of anuran

species) and predators (invertebrates and adult newts) in the pond. Except for the depth of the

pond, all those explanatory variables were also applied to explain the success (offspring

survival) of breeding events. We summed the density of competitors and similarly the density

of predators despite the differences in competitive performance and predation pressure of the

various species toward Parsley Frog tadpoles.

To assess the potential impact of predation and competition on survival rates, we evaluated the

mean density of predators and competitors encountered by Parsley Frog tadpoles during their

larval development. More precisely, data from literature indicates that only small tadpoles (<12

mm snout-vent length) have lower survival due to predation by aquatic invertebrates (Tejedo

1993). Above this size, the predators will only injure them or even fail to catch them. Larvae

laid in autumn reached this limit size in about 3 months, whereas only 1.5 month is necessary

for larvae laid in spring (personal observation). Thus, we used the mean density of predators

and competitors over a period of 3 months after spawning date for autumn tadpoles and 1.5

months for spring tadpoles.

Statistical analyses

All statistical analyses were performed on R 3.4.1 (R Core Team 2018). To assess if pond characteristics differed between seasons, we applied a linear model for the depth of the pond and generalized linear models with a quasi-Poisson family for all other variables to account for overdispersion. The *presence of egg masses* and the *number of egg masses* were analysed using a generalized mixed model with site as random effect, with a binomial family or a negative binomial family (to account for overdispersion), respectively. The hatching rate and survival rates from egg to metamorph were often zero hence we decided to analyse them as binary variables using a generalized mixed model with site as random effect and a binomial family. Those derived binary variables, called *hatching success* and the *metamorphosis success* are the probability of observing at least one hatchling or one metamorph in a pond where egg masses were observed. The significance of fixed effects were tested using Chi² tests to compare nested models (Zuur *et al.* 2009).

Bet-hedging model

Finally, we wondered if the coexistence of two breeding periods could result from a bet-hedging strategy, with the optimal strategy being to split the breeding effort between the two favourable seasons to spread the risk of complete failure (Seger & Brockman 1987). The following model is an ESS model derived from Saiah and Perrin (Saiah & Perrin 1990) on the hatching probability of fairy shrimp seasonal cohorts. Their model was primarily inspired by Cohen (1966), on the optimal reproduction strategy of an annual plant whose seeds can either germinate or remain dormant. In our case, there are two strategies: autumn breeding with initial success (i.e. the ability of offspring to persist until spring) depending on the environmental conditions, and spring breeding with success depending mainly on the presence of autumn

tadpoles, hence on the initial success of autumn breeding (as suggested by the results on success

of autumn and spring breeding events, see below).

Let c be the proportion of eggs laid in autumn (thus 1-c in spring) – we assume, in agreement

with our data (see results), that c represents a fixed strategy, i.e. the frogs cannot predict failure

in advance to avoid laying in autumn, nor can they avoid laying eggs in spring when an autumn

cohort is present. As mentioned above, the autumn cohort is assumed to succeed or fail, at

random, with probability q and 1-q respectively. When it succeeds, a fraction s_1 of the offspring

survive to reproductive age. The spring cohort completely fails whenever the autumn cohort

has survived in a pond (a reasonable simplification based on our survival rates estimates, see

below), otherwise a proportion s_2 of spring tadpoles survive. Overall, the expected number of

offspring reaching sexual maturity is $c\ s_1$ when the autumn cohort doesn't fail and

$(1-c)\ s_2$ when it does.

If each frog reproduced only during one year, the optimal strategy would maximize the

geometric mean of the annual reproductive outcome (Dempster 1955) which is

$$W = (c\ s_1)^q \left((1-c)\ s_2\right)^{1-q}$$

Or, equivalently

$$Log(W) = q\ Log(c\ s_1) + (1-q)\ Log\left((1-c)\ s_2\right)$$

However, reproductive life lasts more than one year in frogs (say, n years), which in itself is a

way to spread the risk of failure among successive cohorts of offspring – an uncertainty remains

however, for each frog, on how many (k) of the n breeding years will not allow the autumn

cohort to survive. For each individual, k is distributed binomially with probability 1-q so that

$$Log(W) = \sum_{k=0}^{n} \binom{n}{k} (1-q)^k\ q^{n-k}\ Log(c\ s_1\ (n-k) + (1-c)\ k\ s_2)$$

where $\binom{n}{k}$ represents the number of possible repartitions of the k years with autumn failure

among the total number of breeding years n.

The selection gradient on c is the derivative of the function Log(W), which indicates whether

selection favours an increase in c (if positive) or a decrease (if negative):

$$G[c] = \sum_{k=0}^{n} \binom{n}{k} (1-q)^k q^{n-k} \frac{(s_1 + s_2)\, k - n\, s_1}{c(k\,(s_1 + s_2) - n\, s_1) - k\, s_2}$$

If some value of c within the authorized interval [0,1] results in $G[c]=0$ then it is considered an

evolutionary stable strategy (ESS) provided the second derivative is negative (i. e. $G[c]$ is

positive below the ESS and negative above).

We explored numerically the selection gradients in order to find potential ESS using

Mathematica (Wolfram Research Inc. 2018) based on the following parameter combinations.

We set survival probabilities based on our estimates of survival from egg to metamorphose: s_1

= 0.047 (estimated among breeding events producing offspring that survived until spring) and

$s_2 = 0.038$ (in the absence of autumn tadpoles). We assumed that survival and fecundity were

equal for both seasonal cohorts for the rest of the life cycle. We set the number of reproductive

years $n = 3$ to 5, according to a study of age structure of a breeding population in Spain (Esteban

et al. 2004). Note that this model applies at the individual level (as developed above) as well as

at the genotype level.

RESULTS

Characteristics of temporal niches

Pond depth was not significantly different between the autumn (here from September to

December) and spring (here from January to April) breeding seasons (Table 1). The densities

of amphibian larvae (other than Parsley Frog) were not significantly different. In autumn,

extreme densities of *Epidalea calamita* tadpoles were recorded in some small ponds whereas

the well-known spring breeding-species (*Hyla meridionalis*, *Pelophylax sp.*, *Triturus*

marmoratus, *Lissotriton helveticus*) reproduce later than the Parsley Frog, hence their larvae

are only present from April onwards. The density of potential invertebrate predators was higher

in autumn than in spring (χ^2_1 = 37.17, p-value = 0.005) with the lowest density being from

December to March. On the contrary, the number of adult newts (potential predators of Parsley

Frog tadpoles) was higher in spring than in autumn (χ^2_1 = 369.36, p-value = $2.2 * 10^{-16}$).

309   Breeding phenology

We registered 184 breeding events, 79 in autumn and 105 in spring. Note that in two sites, one

breeding event was recorded in May. The *number of egg masses* recorded in one pond showed

a bimodal pattern with a peak in October and another in February (Figure 1).

The *presence of egg mass*es (finding at least one egg mass when visiting a pond) was not

significantly different between the two seasons (0.18 ± 0.02, mean ± S.E per visit in autumn

and 0.22 ± 0.02 per visit in spring, (over 429 and 470 visits, respectively) χ^2_1 = 2.31, p-value=

0.128, see also Table 2). This variable was not affected by the presence of anurans from other

species (larvae), nor by the presence of predators (invertebrates or adult newts). It was

positively related to the depth of the pond (χ^2_1 = 20.40, p-value = $6.3 * 10^{-6}$). The *presence of egg*

masses observed in spring was not affected by the presence of autumn tadpoles (χ^2_1 = 0.03, p-

value = 0.875).

The *number of egg masses* was higher in autumn than in spring (23.0 ± 4.0 egg masses per

breeding event in autumn and 13.7 ± 2.4 in spring; χ^2_1=9.25, p-value = 0.002, Fig. 2, Table 2

and Annex 2). As a result, autumn breeding contributed slightly more than spring breeding to

the production of egg masses (57% versus 42.9%).

326   Breeding success

Hatching success (i.e. the percentage of breeding events producing at least one larvae) was

higher in autumn than in spring (68.4% and 43.8% respectively, χ^2_1 = 11.12, p-value = 0.001,

Table 2). *Metamorphosis success* (i.e. the percentage of breeding events producing at least one

metamorph) was not significantly different between the two seasons (34.2% in autumn and

29.8% in spring $\chi^2_1 = 0.39$, p = 0.531).

Drought (pond totally dried up) caused the total failure of 7 breeding events in autumn and of

5 breeding events in spring over the 3 year-survey and the 19 sites (representing 9% and 4.8%

of the breeding events, those percentages are not significantly different, $\chi^2_1 = 0.66$, p = 0.42).

Drought caused mortality of offspring at different developmental stages (mostly eggs for

autumn cohort and tadpoles for spring cohort). Neither *hatching success* nor *metamorphosis*

success were explained by interspecific competition (the density of other amphibian larvae) or

by predation (density of potential invertebrate predators or number of adult newts, see Table

2).

Survival rates are represented in Figure 3 and Annex 3. The *survival rates from egg to*

metamorph were similarly low (autumn: 2.24 % ± 0.61 and spring: 1.97 % ± 0.73, Table 2),

resulting in a higher contribution (74.6%) of autumn breeding in the overall production of

metamorphs per site and per year (due to the higher breeding effort in autumn, see above).

The autumn cohort persisted until spring in 34/79 breeding events (43%, corresponding to the

rate of initial success, *q,* see bet-hedging model). In those cases, tadpoles laid in spring

coexisted in their pond with tadpoles from the autumn cohort. The presence of an autumn cohort

of Parsley Frog tadpoles significantly reduced the *metamorphosis success* of spring cohorts to

18.4% in presence of autumn tadpoles, versus 40.0% in absence of autumn tadpoles, $\chi^2_1 = 10.60$,

p-value = 0.005). This reduction effect was not significant for the *hatching success* (32.6% in

presence of autumn tadpoles, versus 53.6% in absence of autumn tadpoles $\chi^2_1 = 4.63$, p-value

= 0.099). Accordingly, all three survival rates were reduced in the presence of autumn tadpoles

and this effect was most pronounced for the *survival from egg to metamorphs* (3.77% ± 1.4

versus 0.16% ± 0.08 in absence versus presence of autumn tadpoles, Figure 4 and Annex 4).

We also tested if the *metamorphosis success* of autumn tadpoles might be affected by the

presence of spring tadpoles, but this was not the case ($\chi^2_1 = 2.75$, p-value = 0.097).

Finally, figure 5 summarizes the breeding strategy of Parsley Frog showing the presence of egg

masses, tadpoles and the metamorphs in each studied site, over the three years of survey. It

illustrates the quasi-exclusion between the two cohorts: there were only 4 cases in total (out of

47 observations) where metamorphs from the two seasonal cohorts emerged in spring in the

same pond during the same year.

Maintenance of spring breeding

Selection gradients based on our bet-hedging model predict that a mixed strategy is maintained

when the rate of initial success of the autumn cohort (*q*) is between 0.2 and 0.8 for a number of

reproductive years n = 3. In this condition, a pure autumn strategy is predicted above 0.8, and

a pure spring strategy below 0.2. (Figure 6). The maintenance of this strategy is less probable

if the number of reproductive years increases (n = 5 years of breeding), with a reduced range of

q leading to a stable mixed strategy.

DISCUSSION

Cost and benefits of a bimodal breeding phenology

We used field surveys to describe the breeding phenology of the Parsley Frog in the French

Mediterranean region but also to quantify the relative contribution and success of each seasonal

reproduction (autumn and spring reproduction). This quantification, rarely achieved in the wild

(but see Licht 1974; Banks & Beebee 1988; Gascon 1992; Wheeler *et al.* 2015), is essential to

understand the evolution of this bimodal breeding strategy. We confirmed the existence of two

distinct seasonal peaks in breeding activity, probably mediated by cold temperature in

December and January as adult Parsley Frogs tend to breed in mild and rainy periods as was

previously observed (Toxopeus *et al.* 1993; Guyétant *et al.* 1999; Jakob *et al.* 2003). However

breeding episodes occurred even in the absence of rainfall as long as ponds were filled with

water (personal observations and Richter-Boix *et al.* 2006b).

The *number of egg masses* in our population was higher in autumn than in spring. This is in

apparent contradiction with Richter-Boix et al. (2006b) who found that a four-fold higher

breeding effort in spring than in autumn in the northeast of the Iberian Peninsula. While we

don't have a definitive explanation for this difference, we suggest it could be related to higher

competition among anuran larvae in autumn in north-east Spain compared to France. In our

study area in southern France, larvae of *Pelodytes punctatus* are typically the only anuran larvae

found after the summer drought in the ponds in autumn. In contrast, four other species of Anura

(*Hyla meridionalis*, *Epidalea calamita*, *Alytes obstetricans* and *Pelophylax perezi*) have

tadpoles in autumn and three of them (i.e. all except *E. calamita*) can have overwintering

tadpoles in Spanish ponds (Richter-Boix *et al.* 2006b). These authors also showed that

Pelodytes punctatus tadpoles suffer from interaction with *Hyla meridionalis* (Richter-Boix *et*

al. 2007b). It is thus possible that increased competition for *Pelodytes punctatus* larvae in

autumn and winter makes the autumn niche less favourable in north-eastern Spain compared to

our study area in southern France and reduce Parsley Frog investment in autumn breeding there.

The *survival rate from egg to metamorph* was low in both seasons. The combination of *numbers*

of egg masses and survival rates eventually resulted in a higher contribution of autumn breeding

to the overall production of metamorphs. The overall low survival rates of offspring that we

found is in line with previous field studies in anurans (e.g. Licht 1974; Banks & Beebee 1988)

and can be caused by pond desiccation, predation, inter and intra-specific competition for food

and parasitism or pathogen infections. Our study revealed no obvious effect of variation in

predation on tadpole survival even if the predation pressure encountered by tadpoles at the

beginning of their development varies from site to site (but not between seasons). This may

seem surprising since many studies experimentally demonstrated that predation cause

substantial mortality to tadpole populations (e.g. Tejedo 1993; Van Buskirk & Arioli 2005).

This may be due to the lack of information about predation during the first year of survey which

reduced our statistical power or to the fact that causative factors are numerous and more

complex to identify in the field. However, other studies reported no effect of predation on

tadpole survival (Hartel *et al.* 2007) or even a positive effect (Barandun & Reyer 1997),

probably due to predator-induced phenotypic plasticity. Nevertheless, our results suggest that

the predation pressure is probably not a stronger constraint in one season than in the other.

Spring tadpoles should be exposed to more competitors during their development than autumn

tadpoles since the majority of amphibian species in the local community breed in March and

April. Nevertheless, we found no effect of interspecific competition on survival for any of the

two seasonal tadpole cohorts. This seems surprising since Parsley Frog is a poor competitor as

a tadpole compared to most species of the anuran community, in particular *Hyla meridionalis*

and *Rana perezi*, present in spring in permanent ponds (Richter-Boix *et al.* 2007b). On the

contrary, in small temporary ponds and during autumn and winter season, Parsley Frog tadpoles

encounter mostly *Bufo bufo* and *Epidalea calamita* with even lower competitive abilities

(Richter-Boix *et al.* 2007b). We hypothesize that interspecific competition effect was not

detected in our study due to numerous uncontrolled sources of variation.

Priority effects

We revealed a striking negative effect of the presence of conspecific autumn tadpoles on the

survival of spring tadpoles in the Parsley Frog. Previous studies have demonstrated the

occurrence of such intraspecific priority effects in amphibians in experimental settings (Morin

et al. 1990; Eitam *et al.* 2005; Murillo-Rincón *et al.* 2017) but as far as we know, our study is

the first evidence for intraspecific, inter-cohort competition in amphibians in nature. In the field,

we observed in several occasions that large autumn tadpoles were eating freshly laid eggs of

their own species, which could partly explain the lower hatching rate of spring eggs in presence

of autumn tadpoles. Moreover, (Tejedo 1991) previously described how Parsley Frog tadpoles

preys on *Epidalea calamita* eggs. In this latter study, predatory tadpoles were exclusively old

tadpoles and they could cause a loss of 50 to 100% of the eggs. Oophagy has also been

demonstrated to be responsible for interspecific priority effects between *Scaphiosus couchii*

and *Bufo speciosus* (Dayton & Fitzgerald 2005). Intraspecific oophagy has been described in

some anuran species (Summers 1999; Dayton & Wapo 2002) and has been proposed as an

energetic opportunistic response in food shortage in temporary ponds.

However, the presence of autumn tadpoles also affects the larval survival (post-hatching) of

spring tadpoles. This may reflect competition for resources between large autumn and small

spring tadpoles as shown in *Rana arvalis* (Murillo-Rincón *et al.* 2017). Interference competition

mediated by microorganisms may also play a role: smaller tadpoles could display coprophagy

instead of feeding on higher quality resources (Beebee & Wong 1992; Baker & Beebee 2000).

This large priority effect between the two seasonal tadpole cohorts of Parsley Frog has a great

impact on the overall efficiency of breeding: in most ponds, there could be only one successful

breeding period, autumn or spring. Nonetheless, we found no indication that spring breeders

select their oviposition site to avoid conspecifics, as other amphibian species sometimes do

(Sadeh *et al.* 2009). Accordingly, the *presence of egg masses* was also unaffected by the

presence of potential competitors or predators.

Seasonal partitioning of breeding: a bet-hedging strategy?

The temporal partitioning of breeding activity could reflect several evolutionary processes: 1)

the existence of two specialized phenotypes either genetically determined (in which case we

would expect temporal genetic differentiation between cohorts) or set by early environmental

cues (phenotypic plasticity); 2) a use of alternative strategies by some or all individuals (bet-

hedging). We previously demonstrated that the two temporal cohorts do not reflect two

genetically distinct temporal populations (Jourdan-Pineau *et al.* 2012) but breeding phenology

may still be set once for good for each individual. In this case, breeding in autumn or in spring

could be determined by the physiological state (and sexual maturity) of the breeder and

maintained year after year, by physiological constraints (typically the case for a capital breeder

species which stores energy for future reproduction (*e.g.* Lardner & Loman 2003). In a

diversified bet-hedging strategy, individual breeding activities could vary from year to year

(each year, individuals would "choose" one breeding season) or individuals could split their

breeding effort between the two seasons in some or most years. There is no individual data

available for this species and our only attempt to mark adults with visible implant alpha tags was

not successful. Preliminary results based on genotyping of eggs, spawned in the same pond at

different periods, suggests that females could breed several times in one year but this has to be

confirmed (unpublished data). Clearly, this is a line of research to develop in the future if we

want to fully understand the evolution of reproduction in this system.

Based on our field survey, it appears that the bimodal breeding phenology of Parsley Frog is a

typical diversified bet-hedging strategy. The large priority effect between the two seasonal

cohorts, combined with high unpredictability of conditions that result in failure or success of

entire cohorts, results in frequency dependent-selection and favour risk-spreading strategies:

the best option is to develop in ponds with the smaller number of conspecific competitors. These

conditions are found partly in autumn, when the habitat becomes favourable after the dry

summer period, or in spring, as some of the autumn cohorts have died in the winter, leaving the

habitat free. Poethke et al. (2016) developed a theoretical model in which they outlined this

impact of competition on the evolution of bet-hedging strategy. Using a model for optimal

germination fraction, based on field data on desert plants, Gremer and Venable (Gremer &

Venable 2014) also showed that density-dependence could explain the observed bet-hedging

strategy of germination spread in time (i.e. not all seeds at once). Density-dependence was not

included in our model and we do not have field data to assess its effect in our populations. This

would be a fruitful line of research to improve our understanding of this breeding system.

In the Parsley Frog, our model shows that the observed mixed breeding strategy ($0 < c < 1$) is

maintained if the rate of initial success of the autumn cohort (q) is between 0.2 and 0.8 (if

females have on average 3 years of breeding in their lifetime) or between 0.35 and 0.65 (for 5

years of breeding). Those conditions are fulfilled according to our field estimates ($q = 0.43$).

We estimated the proportion of eggs laid in autumn by all breeders ($c = 0.57$) but could not

estimate this proportion at the individual level. Survival rates set in the model were based on

our field estimates of survival from egg to metamorphosis; hence, we assumed similar survival

after metamorphosis of the two cohorts. Unfortunately, we have no information about survival

of Parsley Frog during its adult terrestrial life. However, the adult survival is an important

parameter in our model since it determines the number of reproductive years. The mixed

breeding strategy is less stable when the number of breeding opportunities per lifetime increases

– as the risk is now spread over several successive years. Indeed, experiencing variation in

reproductive success among those opportunities is less harmful when it is possible to try again

the next year. A skeletochronology study conducted in a upland population in Spain indicated

that the mean age of sampled Parsley Frog females was 5.01 years (with a standard deviation

of 1.99) (Esteban *et al.* 2004). Assuming a minimal age at first reproduction of 1 year (as done

by Esteban *et al.* 2004), this translates into an average number of reproductive years or females

of n = 4. Our evaluation of the bet-hedging strategy with n = 5 is thus probably conservative.

We previously showed that the Parsley Frog successfully exploits two temporal niches in the

Mediterranean region thanks to a high phenotypic plasticity of tadpole development to face

very different seasonal environments (Jourdan-Pineau *et al.* 2012). Recently, the combination

of phenotypic plasticity and bet-hedging has been theoretically investigated, suggesting that phenotypic plasticity could further minimize fitness variances caused by mismatches between phenotype and environment (Rádai 2020; Haaland *et al.* 2021). Interestingly, in the wolf spider, temperature and day length leads to alternative developmental types within broods. This cohort splitting is both probabilistic and sensitive to environment, a phenomenon proposed as being a plastic bet-hedging strategy by Rádai (2020). In this case, the various plastic phenotypes, triggered by environmental variations, constitute a bet-hedging response to grassland habitats with substantial and unpredictable year-to-year variation.

In the Parsley Frog, priority effects are not the only factor influencing the relative success of the spring and autumn strategies in terms of future recruitment: autumn tadpoles metamorphose earlier and at a much larger size than spring tadpoles (Jourdan-Pineau *et al.* 2012, unpublished data). This should confer to them a significant advantage in terms of survival to adulthood (Smith 1987; Altwegg & Reyer 2003; Székely *et al.* 2020) even if we don't know whether size and date of metamorphosis affects survival and ultimately fitness in our model. In addition, there is no significant difference in cohort survival (the probability to produce at least one metamorph) between spring and autumn, in spite of a slightly higher risk of drought (and hence complete disappearance of the cohort) in autumn. Density-dependence (on which we have no information) might partly explain why autumn cohorts do as well as spring cohorts in spite of higher drought risk. Additionally, our measures of breeding success are very rough because counting precisely the number of larvae from each cohort in the ponds over the course of the season is extremely difficult. There is thus still much to learn to fully understand the advantages and disadvantages of autumn and spring strategies in this species.

Lastly, as explained above, we still don't know if individual females usually breed once a year (either spring or autumn) or several times a year (potentially spring and autumn of the same year). Capture-mark-recapture of adults and larvae would alleviate some of these limitations

but would be highly challenging. However, our results remain valid for a large range of

parameters, and these uncertainties should not affect our conclusion that the breeding strategy

of Parsley Frogs in southern France constitutes an original example of a bet-hedging strategy

driven by high environmental stochasticity and large inter-cohort priority effect.

ACKNOWLEDGMENTS

This research was supported by a grant from Agence Nationale de la Recherche (SCOBIM

JCJC 0002). We are grateful to Vincent Mouret, Alain Fizesan, Denis Rey, Simon Russeil and

Jérémy Aubain for help in field work. Virginie Ravigné provided valuable help with the bet-

hedging model.

CONFLICT OF INTEREST DISCLOSURE

The authors of this article declare that they have no financial conflict of interest with the content

of this article. HJ-P is one of the PCIEvolBiol recommenders.

REFERENCES

Alanärä, A., Burns, M.D. & Metcalfe, N.B. (2001). Intraspecific resource partitioning in brown

trout: The temporal distribution of foraging is determined by social rank. *J. Anim. Ecol.*,

70, 980–986.

Albrecht, M. & Gotelli, N.J.J. (2001). Spatial and temporal niche partitioning in grassland ants.

Oecologia, 126, 134–141.

Altwegg, R. & Reyer, H.-U.U. (2003). Patterns of natural selection on size at metamorphosis

in water frogs. *Evolution (N. Y).*, 57, 872–882.

Baker, G.C. & Beebee, T.J.C. (2000). Evidence for the induction of interference competition

between anuran larvae in plastic pond cages. *Amphib. Reptil.*, 21, 25–37.

Banks, B. & Beebee, T. (1988). Reproductive success of natterjack toads Bufo calamita in two

contrasting habitats. *J. Anim. Ecol.*, 57, 475–492.

Barandun, J. & Reyer, H. (1997). Reproductive ecology of Bombina variegata: development of

eggs and larvae. *J. Herpetol.*, 31, 107–110.

Beaumont, H.J.E., Gallie, J., Kost, C., Ferguson, G.C. & Rainey, P.B. (2009). Experimental

evolution of bet hedging. *Nature*, 462, 90–93.

Beebee, T.J.C. & Wong, A.L.C. (1992). Prototheca-mediated interference competition between

anuran larvae operates by resource diversion. *Physiol. Zool.*, 65, 815–831.

Di Bitetti, M.S., De Angelo, C.D., Di Blanco, Y.E. & Paviolo, A. (2010). Niche partitioning

and species coexistence in a Neotropical felid assemblage. *Acta Oecologica*, 36, 403–412.

Blaustein, L. & Margalit, J. (1996). Priority effects in temporary pools: nature and outcome of

mosquito larva-toad tadpole interactions depend on order of entrance. *J. Anim. Ecol.*, 65,

77–84.

Brown, C.J., O'Connor, M.I., Poloczanska, E.S., Schoeman, D.S., Buckley, L.B., Burrows,

M.T., *et al.* (2016). Ecological and methodological drivers of species' distribution and

phenology responses to climate change. *Glob. Chang. Biol.*, 22, 1548–1560.

Van Buskirk, J. & Arioli, M. (2005). Habitat specialization and adaptive phenotypic divergence

of anuran populations. *J. Evol. Biol.*, 18, 596–608.

Carothers, J.H. & Jaksić, F.M. (1984). Time as a Niche Difference : The Role of Interference

Competition. *Oikos*, 42, 403–406.

Cayuela, H., Besnard, A., Béchet, A., Devictor, V. & Olivier, A. (2012). Reproductive

dynamics of three amphibian species in Mediterranean wetlands: The role of local

precipitation and hydrological regimes. *Freshw. Biol.*, 57, 2629–2640.

Charmantier, A. & Gienapp, P. (2014). Climate change and timing of avian breeding and

migration: Evolutionary versus plastic changes. *Evol. Appl.*, 7, 15–28.

Cohen, D. (1966). Optimizing reproduction in a randomly varying environment. *J. Theor. Biol.*,

12, 119–129.

Cohen, D. (1970). A Theoretical Model for the Optimal Timing of Diapause. *Am. Nat.*, 104,

389–400.

Crochet, P.A., Chaline, O., Cheylan, M. & Guillaume, C.P. (2004). No evidence of general

decline in an amphibian community of Southern France. *Biol. Conserv.*, 119, 297–304.

Crump, M.L. (1991). Choice of Oviposition Site and Egg Load Assessment by a Treefrog.

Herpetologica, 47, 308–315.

Dayton, G.H. & Fitzgerald, L.A. (2005). Priority effects and desert anuran communities. *Can.*

J. Zool., 83, 1112–1116.

Dayton, G.H. & Wapo, S.D. (2002). Cannibalistic Behavior in Scaphiopus couchii: More

Evidence for Larval Anuran Oophagy. *J. Herpetol.*, 36, 531–532.

Dempster, E.R. (1955). Maintenance of genetic heterogeneity. In: *Cold Spring Harbor*

Symposia on Quantitative Biology. pp. 25–31.

Eitam, A., Blaustein, L. & Mangel, M. (2005). Density and intercohort priority effects on larval

Salamandra salamandra in temporary pools. *Oecologia*, 146, 36–42.

Erich, M., Ringler, M., Hödl, W. & Ringler, E. (2015). Brood-partitioning behaviour in

unpredictable environments: hedging the bets? *Behav. Ecol. Sociobiol.*, 69, 1011–1017.

Esteban, M., Sanchez-Herraiz, M.J., Barbadillo, L.J. & Castanet, J. (2004). Age structure and

growth in an isolated population of Pelodytes punctatus in northern Spain. *J. Nat. Hist.*,

38, 2789–2801.

Forrest, J.R.K., Cross, R. & Caradonna, P.J. (2019). Two-year bee, or not two-year bee? How

voltinism is affected by temperature and season length in a high-elevation solitary bee.

Am. Nat., 193.

Frederiksen, M., Harris, M.P., Daunt, F., Rothery, P. & Wanless, S. (2004). Scale-dependent

604 climate signals drive breeding phenology of three seabird species. *Glob. Chang. Biol.*, 10,

605 1214–1221.

606 García-Roger, E.M., Carmona, M.J. & Serra, M. (2017). Modes, mechanisms and evidence of

607 bet hedging in rotifer diapause traits. *Hydrobiologia*, 796, 223–233.

608 Gascon, C. (1992). Aquatic predators and tadpole prey in central Amazonia: field data and

609 experimental manipulations. *Ecology*, 73, 971–980.

610 Gottsberger, B. & Gruber, E. (2004). Temporal partitioning of reproductive activity in a

611 neotropical anuran community. *J. Trop. Ecol.*, 20, 271–280.

612 Graham, J.K., Smith, M.L. & Simons, A.M. (2014). Experimental evolution of bet hedging

613 under manipulated environmental uncertainty in Neurospora Crassa. *Proc. R. Soc. B Biol.*

614 *Sci.*, 281.

615 Gremer, J.R., Kimball, S. & Venable, D.L. (2016). Within-and among-year germination in

616 Sonoran Desert winter annuals: bet hedging and predictive germination in a variable

617 environment. *Ecol. Lett.*, 19, 1209–1218.

618 Gremer, J.R. & Venable, D.L. (2014). Bet hedging in desert winter annual plants: Optimal

619 germination strategies in a variable environment. *Ecol. Lett.*, 17, 380–387.

620 Guyétant, R., Temmermans, W. & Avrillier, J.N. (1999). Phénologie de la reproduction chez

621 Pelodytes punctatus Daudin, 1802 (Amphibia, Anura). *Amphibia-Reptilia*, 20, 149–160.

622 Haaland, T.R., Wright, J. & Ratikainen, I.I. (2021). Individual reversible plasticity as a

623 genotype-level bet-hedging strategy. *J. Evol. Biol.*, 34, 1022–1033.

624 Halloy, M. (2006). Choice of oviposition site in Pleurodema borellii (Leptodactylidae):

625 importance of conspecific tadpole size. *South Am. J. Herpetol.*, 1, 72–78.

626 Halloy, M. & Fiaño, J.M. (2000). Oviposition Site Selection in Pleurodema borellii (Anura:

627 Leptodactylidae) May Be Influenced by Tadpole Presence. *Copeia*, 2000, 606–609.

628 Hartel, T., Nemes, S. & Mara, G. (2007). Breeding phenology and spatio-temporal dynamics

of pond use by the yellow-bellied toad (bombina variegata) population: The importance of

pond availability and duration. *Acta Zool. Litu.*, 17, 56–63.

Jakob, C., Poizat, G., Veith, M., Seitz, A. & Crivelli, A.J. (2003). Breeding phenology and

larval distribution of amphibians in a Mediterranean pond network with unpredictable

hydrology. *Hydrobiologia*, 499, 51–61.

Jourdan-Pineau, H., David, P. & Crochet, P.-A. (2012). Phenotypic plasticity allows the

Mediterranean parsley frog Pelodytes punctatus to exploit two temporal niches under

continuous gene flow. *Mol. Ecol.*, 21, 876–86.

Kivelä, S.M., Välimäki, P. & Gotthard, K. (2013). Seasonality maintains alternative life-history

phenotypes. *Evolution*, 67, 3145–60.

Lardner, B. & Loman, J. (2003). Growth or reproduction? Resource allocation by female frogs

Rana temporaria. *Oecologia*, 137, 541–546.

Lawler, S.P. & Morin, P.J. (1993). Temporal Overlap , Competition , and Priority Effects in

Larval Anurans. *Ecology*, 74, 174–182.

Leimar, O. (2005). The evolution of phenotypic polymorphism: randomized strategies versus

evolutionary branching. *Am. Nat.*, 165, 669–681.

Lewontin, R.C. & Cohen, D. (1969). On population growth in a randomly varying environment.

Proc. Natl. Acad. Sci. U. S. A., 62, 1056–1060.

Licht, L.E. (1974). Survival of embryos, tadpoles, and adults of the frogs Rana aurora aurora

and Rana pretiosa pretiosa sympatric in southwestern British Columbia. *Can. J. Zool.*, 52,

613–627.

Maxwell, C.S. & Magwene, P.M. (2017). When sensing is gambling: An experimental system

reveals how plasticity can generate tunable bet-hedging strategies. *Evolution (N. Y).*, 71,

859–871.

Møller, A.P. (2008). Climate change and micro-geographic variation in laying date. *Oecologia*,

155, 845–857.

Morand, A. & Pierre, J. (1995). Habitat variability and space utilization by the amphibian

communities of the French upper-rhone floodplain. *Hydrobiologia*, 300–301, 249–257.

Morin, P. (1987). Predation, breeding asynchrony, and the outcome of competition among

treefrog tadpoles. *Ecology*, 68, 675–683.

Morin, P.J., Lawler, S.P. & Johnson, E.A. (1990). Ecology and Breeding Phenology of Larval

Hyla-Andersonii - the Disadvantages of Breeding Late. *Ecology*, 71, 1590–1598.

Murillo-Rincón, A.P., Kolter, N.A., Laurila, A. & Orizaola, G. (2017). Intraspecific priority

effects modify compensatory responses to changes in hatching phenology in an amphibian.

J. Anim. Ecol., 86, 128–135.

Olofsson, H., Ripa, J. & Jonzén, N. (2009). Bet-hedging as an evolutionary game: the trade-off

between egg size and number. *Proc. Biol. Sci.*, 276, 2963–9.

Philippi, T. & Seger, J. (1989). Hedging one's evolutionary bets, revisited. *Trends Ecol. Evol.*,

4, 41–44.

Poethke, H.J., Hovestadt, T. & Mitesser, O. (2016). The evolution of optimal emergence times:

bet hedging and the quest for an ideal free temporal distribution of individuals. *Oikos*, 125,

1647–1656.

R Core Team. (2018). R: A Language and Environment for Statistical Computing.

Rádai, Z. (2020). Cohort splitting from plastic bet-hedging: insights from empirical and

theoretical investigations in a wolf spider. *Theor. Ecol.*

Rand, W.M. (1973). A Stochastic Model of the Temporal Aspect of Breeding Strategies. *J.*

theor. Biol., 40, 337–351.

Richter-Boix, A., Llorente, G.A. & Montori, A. (2006a). A comparative analysis of the adaptive

developmental plasticity hypothesis in six Mediterranean anuran species along a pond

permanency gradient. *Evol. Ecol. Res.*, 8, 1139–1154.

Richter-Boix, A., Llorente, G.A. & Montori, A. (2006b). Breeding phenology of an amphibian

community in a Mediterranean area. *Amphibia-Reptilia*, 27, 549–559.

Richter-Boix, A., Llorente, G.A. & Montori, A. (2007a). A comparative study of predator-

induced phenotype in tadpoles across a pond permanency gradient. *Hydrobiologia*, 583,

43–56.

Richter-Boix, A., Llorente, G.A. & Montori, A. (2007b). Hierarchical competition in pond-

breeding anuran larvae in a Mediterranean area. *Amphibia-Reptilia*, 28, 247–261.

Richter-Boix, A., Llorente, G.A. & Montori, A. (2007c). Structure and dynamics of an

amphibian metacommunity in two regions. *J. Anim. Ecol.*, 76, 607–618.

Ryan, T.J. & Plague, G.R. (2004). Hatching asynchrony, survival, and the fitness of alternative

adult morphs in Ambystoma talpoideum. *Oecologia*, 140, 46–51.

Sadeh, A., Mangel, M. & Blaustein, L. (2009). Context-dependent reproductive habitat

selection: the interactive roles of structural complexity and cannibalistic conspecifics.

Ecol. Lett., 12, 1158–64.

Saiah, H. & Perrin, N. (1990). Autumnal vs spring hatching in the fairy shrimp Siphonophanes

grubii (Dybowski)(Crustacea, Anostraca): diversified bet-hedging strategy? *Funct. Ecol.*,

4, 769–775.

Sakai, S. & Harada, Y. (2005). Production of offspring using current income and reserves: size-

number trade-off and optimal offspring, size. *J. Theor. Biol.*, 233, 65–73.

Salvador, A. & Paris, M.G. (2001). *Anfibios Espanoles*.

Salvidio, S., Lamagni, L., Bombi, P. & Bologna, M.A. (2004). Distribution, ecology and

conservation of the parsley frog (Pelodytes punctatus) in Italy (Amphibia, Pelodytidae) .

Ital. J. Zool., 71, 73–81.

Seger, J. & Brockman, H.J. (1987). What is bet-hedging? *Oxford Surv. Evol. Biol.*, 4, 182–211.

Simons, A.M. (2011). Modes of response to environmental change and the elusive empirical

evidence for bet hedging. *Proc. R. Soc. B Biol. Sci.*, 278, 1601–1609.

Simovich, M. & Hathaway, S. (1997). Diversified bet-hedging as a reproductive strategy of

some ephemeral pool anostracans (Branchiopoda). *J. Crustac. Biol.*, 17, 38–44.

Slatkin, M. (1974). Hedging one's evolutionary bets. *Nature*, 250, 704–705.

Smith, D.C. (1987). Adult Recruitment in Chorus Frogs : Effects of Size and Date at

Metamorphosis Author (s): David C . Smith Published by : Ecological Society of America

Stable URL : http://www.jstor.org/stable/1939265 ADULT RECRUITMENT IN

CHORUS FROGS : EFFECTS OF SIZE A. *Ecol. Soc. Am.*, 68, 344–350.

Summers, K. (1999). The effects of cannibalism on Amazonian poison frog egg and tadpole

deposition and survivorship in Heliconia axil pools. *Oecologia*, 119, 557–564.

Székely, D., Cogălniceanu, D., Székely, P., Armijos-Ojeda, D., Espinosa-Mogrovejo, V. &

Denoël, M. (2020). How to recover from a bad start: Size at metamorphosis affects growth

and survival in a tropical amphibian. *BMC Ecol.*, 20, 1–8.

Tarazona, E., García-Roger, E.M. & Carmona, M.J. (2017). Experimental evolution of bet

hedging in rotifer diapause traits as a response to environmental unpredictability. *Oikos*,

126, 1162–1172.

Tejedo, M. (1991). Effect of Predation by 2 Species of Sympatric Tadpoles on Embryo Survival

in Natterjack Toads (Bufo-Calamita). *Herpetologica*, 47, 322–327.

Tejedo, M. (1993). Size-dependent vulnerability and behavioral responses of tadpoles of two

anuran species to beetle larvae predators. *Herpetologica*, 49, 287–294.

Toxopeus, A., Ohm, M. & Arntzen, J. (1993). Reproductive biology of the parsley frog,

Pelodytes punctatus, at the northernmost part of its range. *Amphibia-reptilia*, 14, 131–147.

Urban, M.C. & De Meester, L. (2009). Community monopolization: Local adaptation enhances

priority effects in an evolving metacommunity. *Proc. R. Soc. B Biol. Sci.*, 276, 4129–4138.

Venable, D.L. (2007). Bet hedging in a guild of desert annuals. *Ecology*, 88, 1086–1090.

Wang, C.-C. & Rogers, D.C. (2018). Bet hedging in stochastic habitats: an approach through

large branchiopods in a temporary wetland. *Oecologia*, 188, 1081–1093.

Wheeler, C.A., Bettaso, J.B., Ashton, D.T. & Welsh, J. (2015). Effects of water temperature on

breeding phenology, growth, and metamorphosis of foothill yellow-legged frogs (Rana

boylii): a case study of the regulated mainstem and unregulated tributaries of California's

Trinity River. *River Res. Appl.*, 31, 132-1276–1286.

Wolfram Research Inc. (2018). Mathematica, Version 11.3.

Yasui, Y. & Yoshimura, J. (2018). Bet-hedging against male-caused reproductive failures may

explain ubiquitous cuckoldry in female birds. *J. Theor. Biol.*, 437, 214–221.

Zeuss, D., Brunzel, S. & Brandl, R. (2017). Environmental drivers of voltinism and body size

in insect assemblages across Europe. *Glob. Ecol. Biogeogr.*, 26, 154–165.

Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009). *Mixed effects models*

and extensions in ecology with R. Springer Science & Business Media.

Table 1: Description of the seasonal niches regarding adult breeding (upper part) and larval

development (lower part) of the Parsley Frog. Mean and standard error (se) of the explanatory

variables depending on the season (autumn and spring). N: sample size. *P.p* is *Pelodytes*

punctatus. Depth is in centimeters. Density of amphibian larvae or invertebrate predators is the

mean number of individuals sampled in one dipnet sweep. Parsley Frog is excluded from

calculations indicating "amphibian larvae" or "anuran adults". * denotes significant difference

between season for the considered variable.

Variable	Autumn			Spring		
	N	mean	se	N	mean	se
Adult breeding						
Depth	416	58.03	2.16	459	62.86	2.17
Density of amphibian larvae	429	3.25	1.40	470	1.17	0.54
Density of invertebrate predators*	279	0.81	0.13	275	0.41	0.08
Number of adult newts*	429	0.13	0.03	464	1.08	0.12
Tadpole development						
Density of amphibian larvae during development	52	0.67	0.26	54	3.97	1.93
Density of invertebrates predators during development	39	0.66	0.15	46	0.79	0.17
Number of adult newts during development	52	0.34	0.15	54	0.49	0.13

754   Table 2: Results of the statistical analyses performed to explain the variation of *presence of egg*
755   *masses*, *number of egg masses*, *hatching success* and *metamorphosis success*. Bold letters
756   indicate a significant test (p-value < 0.05).

Variable	Tested effect	χ^2 value	p-value
Presence of egg masses	Season	2.3101	0.1284
	Depth	**20.398**	**6.289e-06**
	Density of invertebrates	0.1869	0.6655
	Number of adult newts	0.2419	0.6229
	Density of amphibian larvae	0.1292	0.7192
	Number of adult anurans	0.3972	0.5286
Number of egg masses	Season	**9.25**	**0.0023**
Hatching success	Season	**11.119**	**0.00085**
	Density of invertebrates	0.1549	0.6939
	Number of adult newts	0.8315	0.3618
	Density of amphibian larvae	0.1501	0.6984
Metamorphosis success	Season	0.3932	0.5306
	Density of invertebrates	0.6914	0.4057
	Number of adult newts	0.4911	0.4835
	Density of amphibian larvae	1.8085	0.1787

763
764 Figure 1: Mean monthly number of egg masses produced by the Parsley Frog for each recorded
765 breeding event. Error bars are standard errors among sites among years.
766

768   Figure 2: Mean number of egg masses produced per season by the Parsley Frog for each

769   recorded breeding event. Error bars are standard errors, among sites, among year. Autumn in

770   grey and spring in black.

Figure 3: Mean survival rates during embryonic stage (hatching rate, n = 159), larval stage (n

= 79) and from eggs to metamorphs (n = 163). Error bars are standard errors, among sites,

among year. Autumn in grey and spring in black.

778   Figure 4: Mean survival rates during embryonic stage (hatching rate n = 86), larval stage (n =

779   27) and from eggs to juveniles (n = 90) of spring cohorts, in presence (black) or absence (white)

780   of older tadpoles laid in autumn. Error bars are standard errors, among sites, among year.

782
783 Figure 5: Summary of breeding strategy of the Parsley Frog in the 19 studied sites (vertical
784 lines) in the three successive years: presence of egg masses (stars), presence of tadpoles
785 (squares) and presence of metamorphs (frogs). Grey is indicative of autumn events and black
786 is indicative of spring events.
787

789
790

Figure 6: Evolutionary stable strategy (black lines), based on a bet-hedging model, predicting

the proportion of eggs laid in autumn (c, x-axis) in relation to the rate of success of the autumn

cohort (q, y-axis), depending on the number of breeding years (n). We set the survival rate of

autumn tadpoles to 4.7% (estimated among breeding events producing offspring that survived

until spring) and the survival rate of spring tapdoles to 3.8%, (estimated in absence of autumn

cohort). The horizontal and vertical lines indicate the field estimates of c and q.

797
798