The decision to develop into αβ or γδ T cells is pre-programmed in distinct subpopulations of DN1 thymocytes

Seungyoul Oh[1,2], Xin Liu[1], Sara Tomei[3,4], Shalin H. Naik[3,4], Daniel H.D. Gray[3,4] & Mark M.W. Chong[1,2*]

1. St Vincent's Institute of Medical Research, Fitzroy, VIC, Australia

2. Department of Medicine (St Vincent's), University of Melbourne, Fitzroy, VIC, Australia

3. The Walter and Eliza Hall Institute of Medical Research, Melbourne, VIC, Australia

4. Department of Medical Biology, University of Melbourne, Melbourne, VIC, Australia

*Address correspondence to Mark M.W. Chong: mchong@svi.edu.au

Abstract

T cells are divided into the αβ and γδ lineages. It is currently thought that these lineages differentiate in the thymus from uncommitted progenitor thymocytes. However, in this study we show that the decision to differentiate into one or either lineage is in fact already fixed in these apparent uncommitted progenitors. Using single-cell RNA sequencing, we reassemble *de novo* a model of early T cell development based on the transcriptional profiles of individual CD4$^-$CD8$^-$ double negative and γδ thymocytes. We show that the earliest thymocyte stage, known as CD4$^-$CD8$^-$ double negative 1 (DN1), is actually comprised of a mixture of transcriptionally distinct subpopulations. Although not yet expressing definitive markers of αβ and γδ lineages, such as the lineage-defining T cell receptors, specific DN subpopulations exhibit restricted developmental potential for either αβ or γδ lineage. Furthermore, specific γδ-primed DN1 subpopulations preferentially develop into IL-17 or IFNγ-producing γδ T cells. Thus, T cell lineage decisions are hardwired from the earliest stages of T cell development.

Introduction

T cells are divided into two principal lineages that are distinguished by the composition by their T cell receptors (TCRs) (Carpenter & Bosselut, 2010). Those that express an αβ TCR include conventional CD4$^+$ and CD8$^+$ T cells and these can be further divided into diverse subsets that have distinct immunologic functions (Harrington et al., 2005; Rothenberg, Moore, & Yui, 2008). T cells that express a γδ TCR are mainly divided into two subtypes defined by their secretion of IL-17A or IFNγ. IL-17A-producing γδ T cells are important for immunity to intracellular bacteria, while IFNγ-producing γδ T cells are well-known for their anti-tumor function (Bai et al., 2017; Gao et al., 2016; Ribot, Lopes, & Silva-Santos, 2021).

T cell development occurs in the thymus, and it is thought that αβ and γδ T cells both arise from common progenitors that are seeded from the bone marrow (Ciofani & Zuniga-Pflucker, 2010; Taghon, Yui, Pant, Diamond, & Rothenberg, 2006). In mice, the earliest stages of T cell development are termed double-negative (DN) because they lack CD4 and CD8 expression. These are further subdivided into DN1 (CD44$^+$CD25$^-$), DN2 (CD44$^+$CD25$^+$), DN3 (CD44$^-$CD25$^+$) and DN4 (CD44CD25$^-$) stages. αβ and γδ development is thought to bifurcate at the DN2b>DN3 transition, when *TCRB/G/D* gene rearrangement occurs to generate a functional pre-TCR or γδ TCR (Ciofani, Knowles, Wiest, von Boehmer, & Zuniga-Pflucker, 2006; Fehling, Krotkova, Saint-Ruf, & von Boehmer, 1995; Kreslavsky, Gleimer, Garbe, & von Boehmer, 2010; Oh, Gray, & Chong, 2021). Evidence also suggests that effector outcomes of γδ T cells may also be determined in the thymus rather than in the periphery because the expression of IL-17A and IFNγ by mature cells correlates with Vγ chain usage. Vγ2$^+$ γδ T cells tend to produce IL-17A while Vγ1$^+$γδ T cells tend to produce IFNγ (Fahl, Coffey, & Wiest, 2014; Narayan et al., 2012).

The prevailing model suggests that TCR signaling plays a deterministic role in lineage commitment and subsequent γδ effector differentiation (Munoz-Ruiz, Sumaria, Pennington, & Silva-Santos, 2017). Strong signals have been shown to activate ERK and induce EGR-ID3 and this appears to drive γδ differentiation, whereas weak signals appear to promote an αβ fate (Hayes, Li, & Love, 2005; Lauritsen et al., 2009; Munoz-Ruiz et al., 2017).

Additionally, weak TCR signaling may promote an IL-17A phenotype in γδ T cells, while strong signaling

promotes an IFNγ phenotype (Sumaria, Grandjean, Silva-Santos, & Pennington, 2017).

However, the role of the TCR remains debatable because a lack of Notch signaling at the DN2 stage compromises

progression to DN3 and impairs αβ development but not γδ development (Tanigaki et al., 2004; Wolfer, Wilson,

Nemir, MacDonald, & Radtke, 2002). How can the stage prior to the apparent γδ branching be compromised and

yet γδ T cell development still proceed? Evidence suggest that this Notch-independence is at least partially due to

the transcription factor ID3 because a weak TCR signal in cooperation with Notch signaling is sufficient to inhibit

E-protein activity (Lauritsen et al., 2009). However, ID3 activity could explain the development of only a subset

of γδ T cells. This murkiness surrounding the αβ versus γδ decision has arisen partly due to a lack of markers to

define the development of thymocytes that have been selected to mature into γδ T cells (Pellicci, Koay, & Berzins,

2020). Together, these ambiguities raise the question as to whether the early stages of T cell development are

really bipotent homogenous populations that can give rise to both αβ and γδ T cells. Emerging evidence in other

developmental pathways suggests that progenitors can be heterogeneous with biases in lineage potential (Naik et

al., 2013). In this study, we perform single-cell RNA sequencing (scRNA-seq) of early thymocytes from mouse

thymus to determine cellular heterogeneity at single-cell resolution and to assemble a model of T cell development

de novo in order to address when the αβ versus γδ lineage decision is actually made.

Results

Differences in the kinetics of αβ versus γδ T cell development from DN1 thymocytes

To better understand the development of the γδ lineage, we began by investigating the kinetics of αβ versus γδ T

cell development in OP9-DL1 cultures, which supports the development of hematopoietic progenitors *in vitro*

into either γδ T cells or αβ cells up to the $CD4^+CD8^+$ double positive (DP) stage (Holmes & Zuniga-Pflucker,

2009). Total DN1 ($CD44^+CD25^-Lin^-$) and DN2 ($CD44^+CD25^+Lin^-$) thymocytes were sorted from C57BL/6 mice

and cocultured on OP9-DL1 cells. The cultures were then analyzed after 9-23d. DP cells were detected by CD8α

(and CD4) expression while γδ cells were detected by TCRγδ expression (Supplementary Fig. 1a, b).

While DN1 and DN2 thymocytes gave rise to both lineages, surprisingly, there was a significant difference in the

kinetics in the development of DN1 thymocytes into γδ cells compared to αβ cells. Starting from DN1 thymocytes,

γδ cells were already present at 9d, with the percentage and cell number peaking at 12d, waning and then

increasing again at 20d (Fig. 1a, b). In contrast, αβ cells only started appearing after 14d and peaked at 20d. The

earlier γδ peak suggests that either γδ development is either simply more rapid than αβ development or that there

may be different developmental pathways for the two lineages, with γδ cells potentially produced in two waves.

Unlike the discordant kinetics of the DN1 cultures, culturing DN2 thymocytes produced both αβ and γδ lineage

cells from 9d and in similar proportions throughout (Fig. 1a, b). γδ cells were produced at a consistently low

frequency of 2-6%, while between one-third to half the cells were αβ cells at each timepoint. Given that DN2

thymocytes are only one stage along from DN1, the consistent rate of both αβ and γδ production argues against

γδ development being simply more rapid.

If the αβ and γδ lineages develop independently, we might expect to see each lineage being derived from distinct

DN1 thymocytes. To investigate this, we tagged individual DN1 thymocytes with unique genetic heritable

barcodes (Naik et al., 2013) and tracked the propagation of these barcodes in OP9-DL1 cultures. If an αβ cell and

γδ cell inherits the same barcode sequences, it means that they were derived from the same progenitor. CD8α$^+$

and TCRγδ$^+$ cells were sorted after 14d and 20d of culture and analyzed for barcode composition (Supplementary

Fig. 2a). While there was some overlap in barcode usage between the four populations, the barcode profiles were

largely non-overlapping, with Pearson's correlations of $r<0.6$ for the first experiment (Fig. 1d, e) and even lower

in the second experiment (Supplementary Fig. 2c, d). Together with the kinetic analysis, this suggest that the DN1

thymocytes may in fact be a heterogenous mixture consisting of distinct subpopulations that developed into either

αβ cells or γδ cells.

Fig 1. Potential heterogeneity in the lineage outcomes of DN1 thymocytes. **(a)** Total DN1

(CD44+CD25-Lin-) and DN2 (CD44+CD25+Lin-) thymocytes were sorted from the thymus of

C57BL/6 mice then analyzed for $\alpha\beta$ versus $\gamma\delta$ lineage differentiation in OP9-DL1 cultures. The

cultures were analyzed at the indicated time points by flow cytometry. $\alpha\beta$ lineage cells were identified

as CD8α^+ while $\gamma\delta$ lineage cells were identified as TCR$\gamma\delta^+$. CD4 expression was also analyzed but

was concomitant with CD8α expression (not shown). Lin = CD4, CD8, B220, CD11b and CD11c.

Representative flow cytometric plots are shown. **(b)** The mean percentages ± S.E.M. of 4 to 7

replicates are shown. **(c)** The mean cell numbers ± S.E.M. are shown. **(d, e)** Barcoded total DN1

thymocytes were differentiated on OP9-DL1 cells. $\alpha\beta$ and $\gamma\delta$ cells were then sorted from the culture

at 14d and remaining culture was sorted at 20d. The barcode composition of the four populations was

then analyzed. Heatmaps are comparing by individual barcodes (rows) across the four populations

the (column) are shown. Heatmap with normalization of barcodes within each population is shown in

(d), while normalization of each barcode usage across all four populations is shown in **(e)**. Indicated

under **(d)** are Pearson's correlation (*r*) analysis between each pair of populations. Shown is one

replicate of two independent experiments. The second replicate is shown in Figure S2C-D and the

corresponding t-SNE clustering to group the barcodes information is shown in the Supplementary

Figure 2b.

A comprehensive transcriptional map of early T cell development at single-cell resolution

To better understand the potential heterogeneity of the early stages of T cell development, we next characterized

the transcriptional landscape of DN and $\gamma\delta$ thymocytes at single-cell resolution. Cells were sorted from the thymus

of C57BL/6 mice for analysis by 10X scRNA-seq over three independent runs (Fig. 2a, b and Supplementary

Table 1). The first run consisted of total DN and TCR$\gamma\delta^+$ cells, the second consisted only of DN1 and DN2 cells

and the third involved sorting DN1+2, DN3 and TCR$\gamma\delta^+$ cells separately and mixing back together post-sort at a

ratio of 55% to 30% to 15%, respectively (Fig. 2a and Supplementary Table 1). This was to ensure that a sufficient

number of DN1 and DN2 cells were captured for high-resolution analysis that would not have been achieved by

analyzing total DN cells.

Fig. 2. Single-cell RNA sequencing analysis of early T cell development. (a) Shown are the gating

strategies used to sort DN and gd thymocytes from C57BL/6 mice for 10X scRNA-seq. Three

separate runs were completed. The first run was total DN and TCRγδ+ cells. The second run consisted

only of DN1 and DN2 cells. The third run involved sorting DN1 plus DN2, DN3 and TCRγδ$^+$ cells

separately, which were then mixed back together post-sort at a ratio of 55% to 30% to 15%

respectively. Dump = CD4, CD8, B220, CD11b, CD11c, TCRβ and TCRγδ. **(b)** Following processing

of the 10X data on CellRanger, each dataset was analyzed for clustering based on the first 12 principal

components in Seurat. Shown is the t-SNE plot of each run color-coded to DN developmental stage,

TCRγδ$^+$ thymocytes or other (non-thymocytes). **(c)** The three datasets, totaling 22,094 cells, were

integrated with SCTransform, then the cells were clustered with Seurat. This yielded 30 distinct

clusters (left plot), which were then annotated to DN developmental stage, TCRγδ$^+$ thymocytes or

other (non-thymocytes) (right plot). **(d)** Violin plots showing expression of selected markers used to

assign individual clusters to DN development stage.

A total of 22,094 high quality cells passed quality control checks across the three 10X datasets. The three datasets

were first integrated to assemble a global view of early T cell development. To recover biological distinction from

these different replicates and minimize batch-associated variability, the pooled data was normalized using

SCTransform. Following dimensional reduction, unsupervised clustering was performed using the first 12

principal components (PC). This identified 30 distinct clusters (Fig. 2c), which were assigned to a DN stage or

mature γδ thymocytes based on the expression of canonical markers (Supplementary Fig. 1a, b). High *Cd44* and

Il7r expression but low T lineage gene expression, including *Il2ra, Tcf7, Cd24a, Notch1 and Bcl11b,* identified

DN1 cells (Fig. 2d). DN2 cells were identified by upregulation of T lineage genes and downregulation of *Il7r*.

DN3 cells were identified by *Ptcra* and further upregulation of T lineage genes. Low *Cd8b1* and loss of *Il2ra*

distinguished DN4 from DN3 cells (Fig. 2d). High *Trdc, Id3, Sox13* and *Rorc* identified mature γδ thymocytes

(Fig. 2d). This analysis revealed that multiple distinct populations correspond to each of the canonical DN stages

and suggests that DN thymocytes are much more heterogenous than previously thought.

We also tested the effect of cell cycle gene expression on the clustering. Cell cycling scores were calculated from

the integrated data and regressed using Seurat's built-in regression model (Supplementary Fig. 3a, b). There was

a slight variation in the number of output clusters, but we did not observe any biological variability that could be

accounted for by cell cycle status, with the genetic profiles nearly identical between output with cell cycle genes

left in or regressed out (Supplementary Fig. 3c). Therefore, exclusion of cell cycle genes was not necessary for

downstream analyses.

Trajectory analysis suggests that αβ and γδ T cell development occurs in parallel

To understand how the single cell transcriptomic map of DN and γδ thymocytes aligned with the established

understanding of early T cell development, we reconstructed a developmental pathway *de novo* from the identified

clusters by ordering the cells based on tracking gene expression in pseudotime analysis.

Pseudotime inference of the integrated data with Slingshot(Street et al., 2018) ordered the cells in the expected

sequence from DN1 through to DN4, corresponding to αβ development (Germain, 2002). There were also two

branch points predicted, one corresponding to the γδ lineage branching from DN1 cells and another branch point

at DN2 and ending with a subpopulation of DN3 cells (Fig. 3a).

We also performed pseudotime inference using Monocle 2 (Qiu et al., 2017). This assembled the cells along an

asymmetric trajectory that divided into six states (Fig. 3b), which were then analyzed for signature gene

expression (Supplementary Fig. 4) to assign the cells to the conventional DN development stages (Fig. 3c, d).

State 2 comprising DN1 and some DN2a cells, was identified as the starting point. State 3 corresponded to the

main αβ pathway, with DN4 as the endpoint. States 5 and 6 terminated with DN3 and corresponded to β-selection

failure (Fig. 3b, d). State 1 corresponded to the γδ branch, and like Slingshot, Monocle 2 inferred that γδ cells

develop directly from DN1. Interestingly, while *Rag* expression was highest in DN3 cells, which is expected,

expression could already be detected in some DN1 cells (Fig. 3e, f). Thus, trajectory analyses suggest that the γδ

lineage decision may be much earlier than the current paradigm of branching at DN2b>DN3 (Supplementary Fig.

1).

Fig. 3. Trajectory analyses predict αβ and γδ T cell development occurs in parallel from the

DN1 stage. (a) Pseudotime analysis of the integrated 10X dataset with Slingshot characterizing the

course of development from DN1 to DN4 (αβ lineage) and γδ T cells, visualized by UMAP. Cells

are color-coded by developmental stage. **(b)** Pseudotime analysis of total DN and γδ thymocytes

(first 10X run) with Monocle 2. The six states identified in the asymmetric trajectory are indicated.

(c) Color-coding of the Monocle 2 analysis by thymocyte development stage based on expression of

marker genes by the individual clusters. **(d)** Shown are the proportions of DN developmental stage

or TCRγδ⁺ cells that contributed to each state of the Monocle 2 analysis. **(e)** Expression of *Rag1* and

Rag2 overlaid onto the Monocle 2 analysis. **(f)** Expression of *Rag1* and *Rag2* overlaid onto the

Slingshot analysis.

Extensive heterogeneity within DN1 and DN2 thymocytes

The trajectory analyses, as well as the OP9-DL1 kinetic studies, predicted that γδ T cell development might be

occurring in parallel with αβ development, rather than branching off at the later DN2b>DN3 point. To better

delineate the earlier stages of the developmental model, we analyzed the second 10X run that was performed

specifically on DN1 and DN2 thymocytes. Unsupervised clustering of the 8,851 cells that pass quality control checks

identified 26 distinct subpopulations (Fig. 4a). Analysis for marker genes identified eight DN1, five DN2a and 10

DN2b subpopulations (Fig. 4a). There were also three small clusters of non-T cells consisting of doublets and B

cells that, for simplicity, were excluded from downstream analysis. DN1s expressed high levels of progenitor

markers, including *Hhex, Il7r* and *Cd44* but low levels of T-commitment markers, such as *Il2ra, Tcf7, Notch1,*

Cd24a and *Myb* (Supplementary Fig. 5a, b). Late T-lineage commitment genes, including *Rag1, Rag2, Notch3*

and *Ptcra,* were used to separate DN2a and DN2b subpopulations (Supplementary Fig. 5a, b). While heterogeneity

in DN1 thymocytes was reported previously (Porritt et al., 2004), the degree of heterogeneity among both DN1

and DN2 thymocytes was far more extensive than expected, which could be clearly seen by the differential

expression of many marker genes (Supplementary Fig. 5b).

Identification of novel cell surface markers for delineating DN1 and DN2 subpopulations

In order to study these novel subpopulations of DN1 and DN2 thymocytes, we needed to identify useful cell

surface markers that could be used for flow cytometry. Differential expression analysis was performed to select

features (p<0.05 and 2-fold difference) that were cross-referenced to GO terms for cell surface proteins

(Supplementary Fig. 6a, b). We then tested commercial antibodies against these candidates. While not all

 antibodies produced a staining pattern consistent with mRNA expression levels, we were able to identify a

 minimal panel to delineate both DN1 and DN2 thymocytes.

Fig. 4. Identification of cell surface markers for delineating DN1 and DN2 the subpopulations

inferred from scRNA-seq. (a) t-SNE visualization of 8,851 DN1 and DN2 thymocytes from the

second 10X run clustered with Seurat. This yielded 26 distinct clusters (left plot), which were then

annotated as DN1a, DN1b, DN1c, DN1d, DN1e, DN2a or DN2b (right plot). **(b)** Genes found to be

differentially expressed between DN1 subpopulations were annotated against Gene Ontology (GO)

terms for cell surface markers (Figure S6). Shown is a heatmap showing the average expression of

the genes encoding cell surface proteins selected to delineate the DN1 subpopulations. **(c)** Flow

cytometric validation of the selected markers used to identify DN1 subpopulations. Total DN1 cells

(CD4$^-$CD8$^-$ CD44$^+$CD25$^-$Lin$^-$) were first divided into five populations based on c-Kit and CD24. The

c-Kit$^-$CD24$^-$ cells (total DN1e) was then divided into three populations based on Ly6a and Bst2.

Finally, the Ly6a$^-$Bst2$^-$ cells were divided into two populations based on Klrk1. Lin = B220, CD11b,

CD11c, TCRβ and TCRγδ. **(d)** Heatmap showing the average expression of genes encoding cell

surface proteins selected to delineate the DN2 subpopulations. **(e)** Flow cytometric validation of the

selected markers used to identify DN2 subpopulations. Total DN2 cells (CD4$^-$CD8$^-$ CD44$^+$CD25$^+$Lin$^-$)

were first divided into two populations based on c-Kit and Thy1. The c-Kit$^+$Thy1^{int} cells (total DN2a)

were divided into three populations based on Ly6d and CD53. Then, the c-Kit$^+$Thy1^{int}Ly6d^{int}CD53$^-$

cells were divided into two populations based on CD53 and CD3e. The c-Kit$^-$Thy1$^+$ cells (total DN2b)

were divided into three populations based on Ly6d and CD53. Then, the c-Kit$^+$Thy1^{int}Ly6d^{int}CD53$^{int/+}$

cells were divided into five populations based on CD53 and CD3e.

The combination of five markers encoded by the genes *Cd24a, Kit, Bst2, Ly6a* and *Klrk1* distinguished the eight

DN1 subpopulations (Fig. 4b). CD24 and c-Kit was previously shown to separate DN1 thymocytes into five

subpopulations, known as DN1a, DN1b, DN1c, DN1d and DN1e(Porritt et al., 2004). DN1a to DN1d each

corresponded to a single cluster (Fig. 4a). Bst2, Ly6a and Klrk1 were then used to delineate DN1e-1 to DN1e-4

(Fig. 4c).

The pair of DN2a clusters 7/15 could not be separated by cell surface markers as well as the pairs of DN2b clusters

4/23, 9/13 and 8/19 due to very similar gene expression profiles. Thus, these cluster pairs were combined. The

resulting 11 DN2 clusters were annotated as DN2a-1 to 4 and DN2b-1 to 7 based on the expression of genes

encoded by *Thy1, Kit, Cd53, Ly6d* and *Cd3e* (Fig. 4d, e).

Distinct DN1 subpopulations are primed to develop into either αβ or γδ lineage

The heterogeneity of the DN1 thymocytes could potentially represent different progenitors of αβ and γδ lineages,

which would explain the early branch point inferred by pseudotime trajectory analyses (Fig. 3a-c). To test this,

we sorted the DN1 and DN2 subpopulations using our antibody panels and accessed their αβ versus γδ lineage

outcomes in OP9-DL1 cultures after 14d and 20d (Fig. 5a-f).

At 14d, we observed strong αβ lineage restriction in DN1a, DN1b and DN1c cultures, whereas DN1d and DN1e

cultures only generated TCRγδ[+] cells (Fig. 5a-c and Supplementary Table 2). At 20d, there was significant

increase in the percentage and number of αβ cells produced from DN1a and DN1b, while DN1c had given rise to

some γδ cells as well as αβ cells (Fig. 5a-c and Supplementary Table 3). DN1d and DN1e subpopulations

continued to exhibit γδ restriction, with DN1e-1 and DN1e-2 producing the highest percentage and number of γδ

T cell among DN1 subpopulations (Fig. 5a-c and Supplementary Table 2).

At 14d, all DN2b subpopulations and DN2a-1 produced a high percentage and number of αβ cells and few TCRγδ[+]

cells, while the rest of DN2a subpopulations produced mostly TCRγδ[+] cells and few αβ lineage cells (Fig. 5d, e,

Supplementary Fig. 7 and Supplementary Table 4). By day 20, all DN2 subpopulations had produced αβ lineage

cells, while TCRγδ[+] cells had been lost from the DN2a cultures (Fig. 5d, e, Supplementary Fig. 7 and

Supplementary Table 5). There was also a dramatic reduction in cell numbers in the DN2b cultures, which may

be a result of the cells undergoing cell death after reaching the DP stage (Fig. 5e).

Fig. 5. Restricted αβ or γδ potential of DN1 subpopulations. (a) Sorted DN1 subpopulations were

cultured on OP9-DL1 cells to assess their lineage potential. The cultures were analyzed at 14d and

20d of culture by flow cytometry. αβ lineage cells were identified as CD8α$^+$ while γδ lineage cells

were identified as TCRγδ$^+$. Representative flow cytometric plots are shown. **(b, c)** Pooled data

analyzing αβ versus γδ differentiation from sorted DN1 subpopulations. The means ± S.E.M. of four

to nine replicates performed over four independent experiments are shown. See Table S2 and S3 for

P-value calculations. **(d, e)** Pooled data analyzing αβ versus γδ differentiation from sorted DN2

subpopulations. The means ± S.E.M. of four to nine replicates performed over four independent

experiments are shown. See Supplementary Table 4 and 5 for *P*-value calculations. **(f)** Repopulation

of dGuo-depleted fetal FTOCs with select DN1 subpopulations. The lobes were then analyzed by

flow cytometry for CD8α versus TCRγδ expression after 14d. Shown is a representative from three

independent experiments.

We also tested the differentiation of select DN1 subpopulations in fetal thymic organ cultures (FTOCs). Like in

the OP9-DL1 co-cultures, DN1b cells only produced αβ cells, DN1c cells primary produced αβ cells and some

γδ cells, while DN1d and DN1e-4 cells were entirely restricted to the γδ lineage (Fig. 5f). Thus, both OP9-DL1

and FTOC systems support the model that the decision to develop into αβ versus γδ lineages is largely hardwired

from the DN1 stage rather than a decision made later in T cell development. However, there remains some

flexibility because small numbers of TCRγδ⁺ cells can also differentiate from DN1c and DN2a subpopulations.

Gene expression analysis of DN1 subpopulations suggest potential relationships with distinct mature γδ

subsets

In our initial clustering analyses of the DN and γδ thymocyte scRNA-seq data, γδ thymocytes were found to

segregate into two distinct but closely related clusters (Fig. 2c). We therefore wanted to determine how these

related to the different DN1 subpopulations. Firstly, differential expression analysis revealed substantial

differences between two γδ clusters, with 93 genes expressed at significantly higher levels by the γδ-1 cluster

while 117 genes were expressed at significantly higher levels by γδ-2 cluster (Fig. 6a). Genes that were highly

expressed by the γδ-1 cluster included *Gzma, Blk, Maf, Sox13, Etv5, Gata3, Ccr9, Rorc, Sox4, Tcf12, Lgals9,*

Cmak4 and *Bcl11b*, which are all associated with IL-17A-producing γδ T cell subset (Sagar et al., 2020). This

suggests that the γδ-1 cluster is likely to be γδ thymocytes that go on to mature into the IL-17-producing subset,

while the γδ-2 cluster are those that are likely to mature into the IFNγ producing γδ T cell subset. This is consistent

with a previous study that suggested the eventual effector function of γδ T cells, may already be acquired in the

thymus (Sumaria et al., 2017). The unique transcriptional profiles of γδ-1 and γδ-2 thymocytes appears to be

evidence of this.

Fig. 6. The gene expression profiles of the different DN1 subpopulations correlate with distinct

γδ effector subsets. (a) Heatmap showing genes differentially expressed (*P*-value <0.05 and 2-fold

difference) between the γδ-1 and γδ-2 thymocyte subpopulations identified in the cluster analysis

(clusters 11 and 19) of DN and γδ thymocyte scRNA-seq data from Figure 2c. Each column is an

individual cell in the dataset while each row is a differentially expressed gene. Genes associated with

either IL-17A (left side) or IFNγ production (right side) are indicated. **(b)** Analysis of DN1

subpopulations for expression of the 210 genes differentially expressed between the two γδ thymocyte

populations. Transcription factors associated with either IL-17A or IFNγ are indicated.

Next, to investigate the relationship between the mature γδ thymocyte subpopulations and the DN1

subpopulations that differentiate into TCRγδ⁺ cells, the DN1c, d and e subpopulations were analyzed

for the expression of the 210 differentially genes that distinguished the two mature γδ thymocyte

populations. This revealed a similar transcriptional profile between DN1d and γδ-1 thymocytes, while

DN1c and all the DN1e subpopulations overlapped significantly with γδ-2 thymocytes (Fig. 6b).

However, there were also clearly differences between the DN1e subpopulations.

We then focused on transcription factors because they are regulators of gene expression and could potentially play

key roles in the hardwiring γδ effector outcomes in DN1 thymocytes. *Sox13* was highly expressed by DN1d cells

(Fig. 6b), which was previously shown to be important for the differentiation of a subset of DN1 thymocytes into

Tγδ17 cells (Spidale et al., 2018). Interestingly, DN1e subpopulations also express transcription factors associated

with IL-17-producing γδ T cells. Notably, DN1e-1 and DN1e-2 cells expressed high levels of *Maf*, while *Gata3*

was highly expressed by both DN1e-1 and DN1d cells (Fig. 6b). We thus predict that γδ effector outcomes could

be pre-wired from the DN1 stage, with DN1d going on to develop into IL-17-producing cells and DN1e potentially

producing both γδ effector subsets.

Hardwiring of γδ subset outcomes in DN1 subpopulations.

To determine if different DN1 subpopulations give rise to different γδ T cell subsets, the TCRγδ⁺ cells that

developed from the different DN1 subpopulations in OP9-DL1 co-cultures were analyzed for effector phenotype

by staining for intracellular IL-17A and IFNγ expression and for expression of specific Vγ chains (Fig. 7a).

Cytokine production is highly associated with specific Vγ chain usage, with Vγ1⁺ cells enriched for IFNγ and

Vγ2⁺ cells enriched for IL-17A production (Narayan et al., 2012; O'Brien & Born, 2010).

DN1c thymocytes generated both Vγ1⁺ and Vγ2⁺ cells, with only a low percentage of IFNγ expression by the

Vγ1⁺ cells (Fig. 7b-d and Supplementary Table 6). DN1d primarily produced Vγ2⁺ cells that secreted IL-17A.

Similarly, DN1e-1 and DN1e-2 were restricted to a Vγ2$^+$IL-17A$^+$γδ. On the other hand, DN1e-3 and DN1e-4

exhibited plasticity and could generate both IL-17A and IFNγ producing cells (Fig. 7b-d and Supplementary Table

6). This strongly suggests that not only is the αβ versus γδ lineage decision largely determined by the DN1 stage

in T cell development, but even γδ effector outcomes are already hardwired.

Fig. 7. Restriction of γδ effector subset outcomes in different DN1 subpopulations. (a) Gating

strategy for analyzing the phenotype of γδ T cells generated from DN1 subpopulations after co-

culturing with OP9-DL1 cells for 14 days. TCRγδ$^+$ (CD4$^-$CD8$^-$TCRβ$^-$) were first divided based on

Vγ1 versus Vγ2 expression. The three subpopulations, including Vγ1$^-$Vγ2$^-$ double negative (DN) cells

were then analyzed for intracellular IL-17A and IFNγ expression. **(b)** Shown are the flow cytometric

plots from a representative of three independent experiments for cytokine expression by γδ T cells

derived from DN1 subpopulations. The top row shows the Vγ1 versus Vγ2 expression. Gated Vγ1$^+$,

Vγ2$^+$ and double negative (DN) cells were then analyzed for IL-17A versus IFNγ expression in the

bottom three rows. **(c)** Pooled data analyzing the percentage of Vγ1 versus Vγ2 cells differentiated

from sorted DN1 subpopulations. The means ± S.E.M of four to six replicates performed over three

independent experiments is shown. See Supplementary Table 6 for *P*-value calculations. **(d)** Pooled

data analyzing percentage of Vγ1$^+$IFNγ$^+$ (left) and Vγ2$^+$IL-17A$^+$ (right) cells out of total TCRγδ$^+$ cells.

The means ± S.E.M is shown (*P<0.05, **P<0.01).

Discussion

A precise understanding of the earliest events in T cell development has until now been complicated by two factors.

First, it has been unclear if the earliest thymocytes constitute a truly bipotent homogenous population that can

differentiate into both αβ and γδ lineages. Second, previous studies that analyzed whole populations by flow

cytometry largely lacked the resolution to reveal the asynchronous development of rare subpopulations.

Leveraging the power of scRNA-seq for rare cell type identification, our study has revealed the extensive

heterogeneity within the early stages of T cell development. Significantly, we show that the αβ versus γδ lineage

decision is already hardwired within DN1 stage. This earliest stage of T cell development in the thymus, instead

of being an uncommitted T cell progenitor population, in fact consists of distinct subpopulations that primed to

develop into αβ T cells or γδ T cells. This is well before the expression of the lineage-defining TCRs.

The concept that the early stages of T cell development may consist of distinct subpopulation has previously been

considered. It was shown that DN1 thymocytes can be subdivided into five subpopulations (DN1a-e) based on

CD24 and c-Kit expression (Porritt et al., 2004), while neonatal IL-17-producing cells were shown to develop

from a subset of DN1d thymocytes that is determined by a cell intrinsic program prior to TCR signaling (Spidale

et al., 2018). Despite these findings, the cellular heterogeneity of DN1e thymocytes and their lineage specification

remained unclear. We have now defined eight DN1 subpopulations, including four novel DN1e subpopulations

and showed that DN1a and DN1b are restricted to $\alpha\beta$ lineage whereas DN1d and DN1e subpopulations are

restricted to $\gamma\delta$ lineage. Therefore, lineage commitment is already determined by the DN1 stage, which is well

before the apparent lineage bifurcation at the DN2b>3a stage.

Unlike $\alpha\beta$ T cells, $\gamma\delta$ T cells are thought to acquire their effector potential in the thymus rather than upon antigen

exposure in secondary lymphoid organs (Ribot et al., 2009). Indeed, a previous study showed that some IL-17-

producing $\gamma\delta$ T cells develop from a distinct subset of DN1 thymocytes (Spidale et al., 2018). Although none of

the DN1 subpopulations expressed any definitive markers of specific mature T cell developments, we observed

substantial transcriptomic overlap between the IL-17-primed subpopulations with mature IL-17-expressing $\gamma\delta$

thymocytes and between the IFNγ-primed subpopulations with mature IFNγ-expressing $\gamma\delta$ thymocytes. Notably,

we observed expression of many important transcription factors. Thus, critical components of the IL-17 or IFNγ

$\gamma\delta$ T cell transcriptional programs are already in place within distinct DN1 subpopulations.

We showed that DN1d thymocytes express many of transcription factors that are expressed by mature IL-17-

producing $\gamma\delta$ thymocytes but not IFNγ-producing thymocytes like *Bcl11b, Etv5, Sox13, Rcf7, Rorc and Maf*.

S*ox13* has previously be shown to be an important lineage determining factor for the neonatal IL-17A producing

cells (Spidale et al., 2018). It is thought to act in concert with other transcription factors like *Bcl11b, Tcf7, Rorc*

and *Maf* to specify the IL-17A-effector program in $\gamma\delta$ T cells (Malhotra et al., 2013; Spidale et al., 2018).

We also showed that DN1e-1 and DN1e-2 are also restricted to Vγ2$^+$ IL-17A- producing γδ T cell fate, while only

DN1e-3 and DN1e-4 cell display the plasticity to also develop into Vγ1$^+$ IFNγ$^+$ cells. This plasticity is likely to

involve the differential expression of transcription factors that contribute to distinct effector fates. Although all

four DN1e subpopulations expressed *Stat1*, which regulates IFNγ stimulated genes (Qiao et al., 2013), DN1e-1

and DN1e-2 also expressed *Maf*. cMaf is known to positively regulate IL-17A- producing γδ T cell development

(Zuberbuehler et al., 2019), whereas a lack of cMaf expression by γδ T cells correlates with increased IFNγ

expression (Gabrysova et al., 2018; Iwata et al., 2017; Zuberbuehler et al., 2019). Furthermore DN1e-1 cells were

found to express *Gata3*, an important regulator of IFNγ expression (Yagi et al., 2010). The interplay between

transcription factors may thus be key to the lineage decisions of DN1 subpopulations. Further analysis of

chromatin states and epigenetic mechanisms associated with these transcription factors will likely to be valuable

to reveal the intricacies of the regulatory cascades driving the different γδ effector outcomes.

While it is clear that that the majority of DN1 thymocytes exhibited a restricted lineage outcome, the fact that

DN1e-3 and DN1e-4 can differentiate into both IL-17A and IFNγ-producing cells, and that DN1c and DN2a still

displayed some γδ potential in addition to their dominant αβ outcome, means that we cannot entirely rule out that

at least some decisions could be made at the DN2b>3 stage. While largely hardwired, there remains some lineage

flexibility and there must be other pathways that can contribute to the αβ versus γδ lineage decision. TCR

signaling may have a role in these remaining undecided precursors. Moreover, appropriate TCR signaling may

still be required for maturation of early progenitors even if they are already hardwired to develop into either αβ

or γδ cells.

Current models for the αβ versus γδ lineage decision frequently center on the role of TCR signals at the DN2b>3

stage. The evidence for a role of differing TCR signals comes from studies that have employed genetic mouse

models to manipulate TCR signaling at the DN3 stage. This revealed that strong signals activate the ERK-EGR-

ID3 pathway to drive γδ T cell differentiation, whereas a weak signal promotes adoption of the αβ fate (Hayes et

al., 2005; Lauritsen et al., 2009; Munoz-Ruiz et al., 2017). TCRγδ signaling is also thought to play a role in

subsequent γδ T cell effector differentiation (Munoz-Ruiz et al., 2017). Weak TCR signaling appears to promote

an IL-17A effector phenotype, while strong signaling promotes IFNγ phenotype (Sumaria et al., 2017). However,

it is still unclear whether TCR signal strength is dependent on instructive extrinsic signals or simply a result of

stochastic selection of TCR chains. The instructive model proposes that TCRγδ signal competes with pre-TCR

signal and the lineage decision is determined by cell specific interactions that activate key transcription factors,

which in turn instructs gene expression program (Carpenter & Bosselut, 2010). In contrast, the stochastic model

suggests that there is random rearrangement of TCR genes, where thymocytes that successfully assemble a β

chain progress along the αβ lineage, while those that assemble specific Vγ chains differentiate into a specific

subset of γδ T cells following appropriate TCR signaling. It is also possible that there may a component of both

instruction and random selection to these decisions. Further analysis of whether TCR signaling occurs in a

deterministic manner or in a random manner will be important to provide mechanistic insight into lineage

commitment and subsequent γδ effector differentiation of those DN1c and DN2a. Moreover, as mentioned,

appropriate TCR signaling may still be important for maturation of the γδ-restricted DN1 subpopulations, even if

lineage outcome has already been determined.

Our novel finding of highly heterogenous DN1 and DN2 subpopulations highlights the concept that progenitor

populations may actually not be multi/bipotent cells at all that are. Progenitors may in fact be a grouping of

lineage-primed cells at a similarly early stage in differentiation. On the surface, they may lack lineage defining

markers and they are similar enough to appear to constitute a population, but individually, these cells have unique

transcriptional programs that already specify a lineage-fate. Such heterogeneity in progenitor populations have

been recently describe in the neural cortex (Johnson et al., 2015), prostate epithelium (Crowley et al., 2020) and

in the heart (Churko et al., 2018). Within the hematopoietic system, lymphoid-primed multipotent progenitors in

the bone marrow also appear to exhibit substantial restriction to specific lineage fates (Naik et al., 2013). More

specifically, myeloid progenitors can be divided into subpopulations that are primed towards neutrophil, basophil,

eosinophil, monocyte, dendritic cell, erythrocyte or megakaryocyte fate (Paul et al., 2016).

In conclusion, we have made the novel discovery that the αβ versus γδ T lineage decision, as well as the effector

outcomes of γδ T cells, are already hardwired in DN1 thymocytes. Such hardwiring in apparently uncommitted

progenitor and prior to the apparent lineage decision, at DN2b>3 in the case of T cell development, could

potentially be a common feature of many cellular developmental pathways. It will thus be of interest to perform

a similar analysis of other immune developmental pathways, including other lymphocytes, in the future.

Materials and Methods

Mice and thymocyte preparations

Thymuses was harvested from C57BL/6 mice at around 6 weeks of age. All experiments were approved by the St

Vincent's Hospital Animal Ethics Committee and performed under the Australian code for the care and use of

animals for scientific purposes.

Total thymocytes were obtained by crushing the thymus through a metal sieve to generate a single cell suspension.

The cells were washed with PBS and filtered through a 70 μm sieve to remove any clumps. CD4$^+$ and CD8$^+$

expressing thymocytes were depleted using anti-CD4 and CD8 magnetic-activated cell-sorting (MACS) beads

(Miltenyi Biotec), according to the manufacturer's instructions. The depleted thymocyte preparation was stained

with surface antibodies for sorting on a FACSAria (BD Biosciences).

Flow cytometry

For the analysis of cell surface phenotype, the cells were simply stained with antibodies. For the analysis of

intracellular cytokine expression, cells were first restimulated *in vitro* with 50ng/mL phorbol 12-myristate 13-

acetate (PMA) + 2 μg/mL ionomycin (both Sigma-Aldrich) in the presence of Monensin (BD Biosciences) at

37°C for 2.5h before staining with cell surface antibodies. The cells were then fixed with the Intracellular Fixation

and Permeabilization buffer set (eBioscience) and stained with antibodies against cytokines. All antibodies were

purchased from eBiosciences or BD Biosciences. Flow cytometry data were then acquired on a BD LSR Fortessa

III (BD Biosciences) and analyzed with FlowJo software v10.7.0 (Treestar). When only analyzing cell surface

phenotype, dead cells were excluded by DAPI. For intercellular cytokine analyses the cells were not stained with

DAPI but were gated on live cells determined by size.

OP9-DL1 co-cultures

Thymocyte subpopulations for interest were purified by MACS depletion and cell sorting then plated onto OP9-

DL1 monolayers (Holmes & Zuniga-Pflucker, 2009). The OP9-DL1 cells were inactivate with Mitomycin C

(Stem Cell) immediately prior to use. 5×10^2 to 3×10^3 sorted thymocytes were seeded per well of a 96-well plate

in aMEM (Life Technologies) supplemented with 20% FCS (Bovogen Biologicals),

penicillin/streptomycin/gentamycin (Gibco), 2ng/mL murine IL-7 (Peprotech) and 5ng/mL human FLT3L

(Peprotech) and were incubated at 37°C and 5% CO_2. The media was refreshed every 2d and freshly inactivated

OP9-DL1 cells were added every 4d. The cells were analyzed by flow cytometry after 9-26d of culture.

Fetal thymic organ culture (FTOC)

Fetal thymic lobes were isolated from embryos at gestational 14d following timed pregnancies of C57BL/6 female

mice. They were cultured for 5-6d on 0.8mm isopore membranes (Millipore) atop surgical gelfoam sponge

(Ferrosan Medical Devices) soaked in RPMI-1640 (Sigma) supplemented with 10% FCS (Bovogen Biologicals),

20mM HEPES (Sigma-Aldrich), 50mM 2-mercaptoethanol (Sigma) and 1.35mM 2'-deoxygyanosine (dGuo,

Sigma) to deplete endogenous thymocytes. The depleted thymic lobes were then transferred onto new sponges

soaked in fresh media with supplements but without dGuo for 2d before repopulation with thymocyte progenitors.

To repopulate thymic lobes, they were placed in 20mL hanging drop cultures on Terasaki plates (Sigma-Aldrich)

containing 5×10^2 to 2×10^3 sorted thymocytes for 24h before returning to fresh sponges. The media was refreshed

every 3-4d. Single cell suspensions of the thymic lobes were generated by passing through a 70mm sieve for

analysis by flow cytometry.

Barcode transduction

Sorted total DN1 thymocytes were pre-cultured on OP9-DL1 for 24h. The cells were then transduced with barcode

lentivirus library(Naik et al., 2013) in StemSpan medium (Stem Cell Technologies) by centrifugation at 900 ×g

for 1.5h at room temp. A viral titer pre-determined to give 10% transduction efficiency was used to ensure that

the cells are not transduced with multiple barcodes. 2ng/mL murine IL-7 and 5ng/mL human FLT3L was then

added to each well and the cells were returned to the incubator. The following day, fresh αMEM with supplements

was added. αβ (CD90.2$^+$ CD8α$^+$ TCRγδ$^-$) and γδ (CD90.2$^+$ CD8α$^-$ TCRγδ$^+$) lineage cells were sorted after 14d

and 20d OP9-DL1 co-culture.

Barcode amplification and sequencing

Barcode library construction was performed as described previously(Naik et al., 2013). The cells were lysed in

0.5mg/ml Proteinase K (Invitrogen) in Direct PCR Lysis Buffer (Viagen) at 55° C for 2h. The Proteinase K was

then inactivated at 85°C for 30min and 95° C for 5min. The lysate was split into 2 wells for technical replicate

PCRs. A first round of PCR was performed using 1× Standard-Taq magnesium free reaction buffer pack (NEB)

with 2 mM MgCl2 (NEB), 0.2mM dNTPs (made in house), 0.5μM TopLiB forward primer

(TGCTGCCGTCAACTAGAACA) and 0.5μM of BotLiB reverse primer (GATCTCGAATCAGGCGCTTA) for

32 cycles (1 cycle at 95 °C for

5min, 30 cycles at 95°C for 15sec, 57.2°C for 15sec and 72°C for 15sec followed by 1

cycle at 72°C for 10min). A second round of PCR was then performed to add different Illumina index to each

sample by amplifying the first round PCR product with a sample specific Illumina forward index primer and a

common Illumina reverse index primer for 32 cycles (1 cycle at 95°C for 5min, 30 cycles at 95°C for 15sec,

57.2°C for 15sec and 72°C for 15sec followed by 1 cycle at 72°C for 10min). An aliquot of the PCR product was

run on a 2% agarose gel to check for barcode amplification, then the samples were pooled and the DNA was

cleaned using NucleoMag SPRI beads (Machery-Nagel). A 75-cycle sequencing run was performed on a NextSeq

instrument (Illumina).

Cellular barcode data processing and analysis

The data was demultiplexed and aligned to the reference barcode library using *processAmplicons* function from edgeR package (Robinson, McCarthy, & Smyth, 2010). The barcode counts were then processed in the following steps: 1) barcodes with 0 read counts in all sample/cell types were excluded from the analysis; 2) Barcodes that were detected in the water control were also removed from the analysis; 3) The read count between technical replicates of the same sample was averaged and the total read count in each sample was then normalized to 100%.

t-distributed stochastic neighbor embedding (t-SNE) was used to the cluster the normalized barcode profiles and visualize any lineage biases of individual barcodes. DBSCAN (version 1.1-8) was then used to classify barcodes based on their corresponding t-SNE coordinates. Heatmaps were plotted to visualize lineage output of barcodes that were identified by transforming the normalized reads value for each barcode using logarithmic transformation. Two-tailed Pearson's correlation analysis with 95% confidence interval was calculated to determine the correlation between each lineage outcomes using Prism v9 (GraphPad).

Single-cell RNA sequencing

Sorted thymocytes were counted and checked for viability, then loaded onto the 10X Chromium platform (10X Genomics) for scRNA-seq library construction using the Single Cell V2 or V3.1 Reagent Kit according to the manufacturer's instructions. Libraries were sequenced using 150-cycle/150bp-reads NextSeq500 (Illumina) or 300-cycle/150bp-reads Novaseq (Illumina). Sequencing files were demultiplexed and aligned to *Mus musculus* transcriptome reference (mm10), and count matrix was extracted using CellRanger Single Cell software v2.1.1 or v4.0.0 (10X Genomics) (Zheng et al., 2017) (Supplementary Table 1).

The Illumina sequencing output was pre-processed with Seurat (v2.3 or v.3.2.2) on R (v3.6.3 and 4.1.0). Cells with <500 genes, >5000 genes or >7% mitochondria gene expression were filtered out as low-quality cells. Following normalization and removal of confounders, highly variable genes were identified and selected using VST selection method (Zwiener, Frisch, & Binder, 2014). Unsupervised linear principal component analysis (PCA) was performed on these highly variable genes to grouped them into 20 principal components. Cell

clustering was implemented using the number of components that retain >90% of variance of gene expression in

the data (stated in Supplementary Table 1).

DoubletFinder (v2.0.3) was applied to remove likely sequencing doublets before downstream analyses (McGinnis,

Murrow, & Gartner, 2019). The expected number of doublets was calculated as 0.75% of every 1,000 targeted

cells of recovery. The remaining cells were re-clustered and were visualized with t-SNE or uniform manifold

approximation and projection (UMAP) dimensional reduction with a resolution of 2.0 (Table S1). Differential

expression between subclusters was carried out using FindAllMarkers function, with default parameters;

differentially expressed genes with adjusted p-value <0.05 and fold change >0.5 or <-0.5 (log2FC) were

considered unless otherwise stated. The heatmaps was generated using function DoHeatmap. Hierarchical

clustering heatmaps were generated with the R package ComplexHeatmap (v2.8.0) using Euclidean distance

measures (Gu, Eils, & Schlesner, 2016).

Cell cycle genes specific to either the G1, S, G2M stages was used to perform cell cycle scoring and assign cells

to their respective stage of the cell cycle (Macosko et al., 2015). Cell cycle genes were regressed using Seurat's

built-in regression model.

Merging of multiple scRNA-seq datasets

To compare cell types and proportions across three independent sequencing runs, the datasets were integrated as

described at https://satijalab.org/seurat/archive/v3.0/integration.html (Stuart et al., 2019). The Seurat package

(v.3.2.2) was used to assemble multiple distinct scRNA-seq datasets into an integrated dataset and cell cycling

scores were calculated. To remove technical variability, the datasets were pre-processed and normalized using

SCTransform (Hafemeister & Satija, 2019). To correct for experimental batch effect, integration anchors were

identified between the experiments then merged using canonical correlation analysis. Linear dimensional

reduction was applied and principal components that retain >90% of variance of gene expression in the data were

included for downstream analysis. Unsupervised clustering was implemented on the integrated data and clustered

the cells using a resolution of 2.0.

Pseudotime trajectory construction

Filtered 10X data was imported into Monocle 2 by generating a cell dataset from the raw counts slot of the Seurat

object. Cells were ordered into a branch pseudotime trajectory according to the procedure recommended in the

Monocle 2 documentation (Qiu et al., 2017). The highly variable genes identified by Seurat were chosen as

ordering genes to recover pseudospatial trajectories using the setOrderingFilter, reduceDimension and orderCells

functions in Monocle 2 using default parameters. Differential expression between pseudotime states were

determined using the Seurat function FindAllMarkers.

Slingshot trajectory construction

Cluster information from merged 10X dataset were used as input for Slingshot package (v2.0.0) to infer cell

trajectory (Street et al., 2018). Default parameters was applied and DN1b was set as the starting point for the

minimum spanning tree. The previously generated UMAP clustering from the integrated data was set as the

cellular embedding on which Slingshot performed trajectory inference computation.

GO term enrichment analysis

The biomaRt package (v3.13) was loaded from the Bioconductor using R studio(Amezquita et al., 2020). The

Gene Ontology (GO) terms Cell surface markers (0009986) and Transcription factors (0003700, 0008134,

0140110) were obtained from 'mmusculus_gene_ensembl' and applied to the input (differentially expressed) gene

lists. Gene ontology enrichment analysis for GO Biological Process was conducted using ShinyGo (v0.66) (Ge,

Jung, & Yao, 2020).

Statistical analysis

Statistical testing was performed with one-way analyses of variance (ANOVA) using Prism v9 (GraphPad). *P*

values are shown as * < 0.05, ** < 0.01, *** < 0.001, and **** < 0.0001 where each statistical significance was

found, and all data are represented as means ± S.E.M.

Data availability

All scRNA-seq datasets have been deposited in the NCBI Gene Expression Omnibus repository under accession

number GSE188913.

Acknowledgments

We thank Lucy Kloboucnik in the St Vincent's Hospital Melbourne's BioResources Centre for expert animal

husbandry, and Anthony Di Carluccio from St Vincent's Institute's Cytometry Facility and Casey Anttila from

Walter and Eliza Hall Institute's Cytometry Facility for assistance with cell sorting and the scRNA-seq libraries.

We also thank Dr David Izon (St Vincent's Institute) for the OP9-DL1 cell line, who originally obtained them

from Dr Juan Carlos Zúñiga-Pflücker (University of Toronto).

Competing interests

The authors declare that they have no competing interests.

References

Amezquita, R. A., Lun, A. T. L., Becht, E., Carey, V. J., Carpp, L. N., Geistlinger, L., . . . Hicks, S. C. (2020).

Orchestrating single-cell analysis with Bioconductor. *Nat Methods, 17*(2), 137-145. doi:10.1038/s41592-

019-0654-x

Bai, H., Gao, X., Zhao, L., Peng, Y., Yang, J., Qiao, S., . . . Yang, X. (2017). Respective IL-17A production by

gammadelta T and Th17 cells and its implication in host defense against chlamydial lung infection. *Cell*

Mol Immunol, 14(10), 850-861. doi:10.1038/cmi.2016.53

Carpenter, A. C., & Bosselut, R. (2010). Decision checkpoints in the thymus. *Nat Immunol, 11*(8), 666-673.

doi:10.1038/ni.1887

Churko, J. M., Garg, P., Treutlein, B., Venkatasubramanian, M., Wu, H., Lee, J., . . . Wu, J. C. (2018). Defining

human cardiac transcription factor hierarchies using integrated single-cell heterogeneity analysis. *Nat*

Commun, 9(1), 4906. doi:10.1038/s41467-018-07333-4

Ciofani, M., Knowles, G. C., Wiest, D. L., von Boehmer, H., & Zuniga-Pflucker, J. C. (2006). Stage-specific and

differential notch dependency at the alphabeta and gammadelta T lineage bifurcation. *Immunity, 25*(1),

105-116. doi:10.1016/j.immuni.2006.05.010

Ciofani, M., & Zuniga-Pflucker, J. C. (2010). Determining gammadelta versus alphass T cell development. *Nat*

Rev Immunol, 10(9), 657-663. doi:10.1038/nri2820

Crowley, L., Cambuli, F., Aparicio, L., Shibata, M., Robinson, B. D., Xuan, S., . . . Shen, M. M. (2020). A single-

cell atlas of the mouse and human prostate reveals heterogeneity and conservation of epithelial progenitors.

Elife, 9. doi:10.7554/eLife.59465

Fahl, S. P., Coffey, F., & Wiest, D. L. (2014). Origins of gammadelta T cell effector subsets: a riddle wrapped in

an enigma. *J Immunol, 193*(9), 4289-4294. doi:10.4049/jimmunol.1401813

Fehling, H. J., Krotkova, A., Saint-Ruf, C., & von Boehmer, H. (1995). Crucial role of the pre-T-cell receptor

alpha gene in development of alpha beta but not gamma delta T cells. *Nature, 375*(6534), 795-798.

doi:10.1038/375795a0

Gabrysova, L., Alvarez-Martinez, M., Luisier, R., Cox, L. S., Sodenkamp, J., Hosking, C., . . . O'Garra, A. (2018).

c-Maf controls immune responses by regulating disease-specific gene networks and repressing IL-2 in

CD4(+) T cells. *Nat Immunol, 19*(5), 497-507. doi:10.1038/s41590-018-0083-5

644 Gao, J., Shi, L. Z., Zhao, H., Chen, J., Xiong, L., He, Q., . . . Sharma, P. (2016). Loss of IFN-gamma Pathway

645   Genes in Tumor Cells as a Mechanism of Resistance to Anti-CTLA-4 Therapy. *Cell, 167*(2), 397-404

646   e399. doi:10.1016/j.cell.2016.08.069

647 Ge, S. X., Jung, D., & Yao, R. (2020). ShinyGO: a graphical gene-set enrichment tool for animals and plants.

648   *Bioinformatics, 36*(8), 2628-2629. doi:10.1093/bioinformatics/btz931

649 Germain, R. N. (2002). T-cell development and the CD4-CD8 lineage decision. *Nat Rev Immunol, 2*(5), 309-322.

650   doi:10.1038/nri798

651 Gu, Z., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in multidimensional

652   genomic data. *Bioinformatics, 32*(18), 2847-2849. doi:10.1093/bioinformatics/btw313

653 Hafemeister, C., & Satija, R. (2019). Normalization and variance stabilization of single-cell RNA-seq data using

654   regularized negative binomial regression. *Genome Biol, 20*(1), 296. doi:10.1186/s13059-019-1874-1

655 Harrington, L. E., Hatton, R. D., Mangan, P. R., Turner, H., Murphy, T. L., Murphy, K. M., & Weaver, C. T.

656   (2005). Interleukin 17-producing CD4+ effector T cells develop via a lineage distinct from the T helper

657   type 1 and 2 lineages. *Nat Immunol, 6*(11), 1123-1132. doi:10.1038/ni1254

658 Hayes, S. M., Li, L., & Love, P. E. (2005). TCR signal strength influences alphabeta/gammadelta lineage fate.

659   *Immunity, 22*(5), 583-593. doi:10.1016/j.immuni.2005.03.014

660 Holmes, R., & Zuniga-Pflucker, J. C. (2009). The OP9-DL1 system: generation of T-lymphocytes from embryonic

661   or hematopoietic stem cells in vitro. *Cold Spring Harb Protoc, 2009*(2), pdb prot5156.

662   doi:10.1101/pdb.prot5156

663 Iwata, S., Mikami, Y., Sun, H. W., Brooks, S. R., Jankovic, D., Hirahara, K., . . . Kanno, Y. (2017). The

664   Transcription Factor T-bet Limits Amplification of Type I IFN Transcriptome and Circuitry in T Helper

665   1 Cells. *Immunity, 46*(6), 983-991 e984. doi:10.1016/j.immuni.2017.05.005

666 Johnson, M. B., Wang, P. P., Atabay, K. D., Murphy, E. A., Doan, R. N., Hecht, J. L., & Walsh, C. A. (2015).

667   Single-cell analysis reveals transcriptional heterogeneity of neural progenitors in human cortex. *Nat

668   Neurosci, 18*(5), 637-646. doi:10.1038/nn.3980

Kreslavsky, T., Gleimer, M., Garbe, A. I., & von Boehmer, H. (2010). alphabeta versus gammadelta fate choice: counting the T-cell lineages at the branch point. *Immunol Rev, 238*(1), 169-181. doi:10.1111/j.1600-065X.2010.00947.x

Lauritsen, J. P., Wong, G. W., Lee, S. Y., Lefebvre, J. M., Ciofani, M., Rhodes, M., . . . Wiest, D. L. (2009). Marked induction of the helix-loop-helix protein Id3 promotes the gammadelta T cell fate and renders their functional maturation Notch independent. *Immunity, 31*(4), 565-575. doi:10.1016/j.immuni.2009.07.010

Macosko, E. Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., . . . McCarroll, S. A. (2015). Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell, 161*(5), 1202-1214. doi:10.1016/j.cell.2015.05.002

Malhotra, N., Narayan, K., Cho, O. H., Sylvia, K. E., Yin, C., Melichar, H., . . . Immunological Genome Project, C. (2013). A network of high-mobility group box transcription factors programs innate interleukin-17 production. *Immunity, 38*(4), 681-693. doi:10.1016/j.immuni.2013.01.010

McGinnis, C. S., Murrow, L. M., & Gartner, Z. J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst, 8*(4), 329-337 e324. doi:10.1016/j.cels.2019.03.003

Munoz-Ruiz, M., Sumaria, N., Pennington, D. J., & Silva-Santos, B. (2017). Thymic Determinants of gammadelta T Cell Differentiation. *Trends Immunol, 38*(5), 336-344. doi:10.1016/j.it.2017.01.007

Naik, S. H., Perie, L., Swart, E., Gerlach, C., van Rooij, N., de Boer, R. J., & Schumacher, T. N. (2013). Diverse and heritable lineage imprinting of early haematopoietic progenitors. *Nature, 496*(7444), 229-232. doi:10.1038/nature12013

Narayan, K., Sylvia, K. E., Malhotra, N., Yin, C. C., Martens, G., Vallerskog, T., . . . Immunological Genome Project, C. (2012). Intrathymic programming of effector fates in three molecularly distinct gammadelta T cell subtypes. *Nat Immunol, 13*(5), 511-518. doi:10.1038/ni.2247

O'Brien, R. L., & Born, W. K. (2010). gammadelta T cell subsets: a link between TCR and function? *Semin Immunol, 22*(4), 193-198. doi:10.1016/j.smim.2010.03.006

Oh, S., Gray, D. H. D., & Chong, M. M. W. (2021). Single-Cell RNA Sequencing Approaches for Tracing T Cell Development. *The Journal of Immunology, 207*(2), 363-370. doi:10.4049/jimmunol.2100408

Paul, F., Arkin, Y., Giladi, A., Jaitin, D. A., Kenigsberg, E., Keren-Shaul, H., . . . Amit, I. (2016). Transcriptional Heterogeneity and Lineage Commitment in Myeloid Progenitors. *Cell, 164*(1-2), 325. doi:10.1016/j.cell.2015.12.046

Pellicci, D. G., Koay, H. F., & Berzins, S. P. (2020). Thymic development of unconventional T cells: how NKT cells, MAIT cells and gammadelta T cells emerge. *Nat Rev Immunol, 20*(12), 756-770. doi:10.1038/s41577-020-0345-y

Porritt, H. E., Rumfelt, L. L., Tabrizifard, S., Schmitt, T. M., Zuniga-Pflucker, J. C., & Petrie, H. T. (2004). Heterogeneity among DN1 prothymocytes reveals multiple progenitors with different capacities to generate T cell and non-T cell lineages. *Immunity, 20*(6), 735-745. doi:10.1016/j.immuni.2004.05.004

Qiao, Y., Giannopoulou, E. G., Chan, C. H., Park, S. H., Gong, S., Chen, J., . . . Ivashkiv, L. B. (2013). Synergistic activation of inflammatory cytokine genes by interferon-gamma-induced chromatin remodeling and toll-like receptor signaling. *Immunity, 39*(3), 454-469. doi:10.1016/j.immuni.2013.08.009

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A., & Trapnell, C. (2017). Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods, 14*(10), 979-982. doi:10.1038/nmeth.4402

Ribot, J. C., deBarros, A., Pang, D. J., Neves, J. F., Peperzak, V., Roberts, S. J., . . . Silva-Santos, B. (2009). CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat Immunol, 10*(4), 427-436. doi:10.1038/ni.1717

Ribot, J. C., Lopes, N., & Silva-Santos, B. (2021). gammadelta T cells in tissue physiology and surveillance. *Nat Rev Immunol, 21*(4), 221-232. doi:10.1038/s41577-020-00452-4

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics, 26*(1), 139-140. doi:10.1093/bioinformatics/btp616

Rothenberg, E. V., Moore, J. E., & Yui, M. A. (2008). Launching the T-cell-lineage developmental programme. *Nat Rev Immunol, 8*(1), 9-21. doi:10.1038/nri2232

Sagar, Pokrovskii, M., Herman, J. S., Naik, S., Sock, E., Zeis, P., . . . Grun, D. (2020). Deciphering the regulatory landscape of fetal and adult gammadelta T-cell development at single-cell resolution. *EMBO J, 39*(13), e104159. doi:10.15252/embj.2019104159

Spidale, N. A., Sylvia, K., Narayan, K., Miu, B., Frascoli, M., Melichar, H. J., . . . Kang, J. (2018). Interleukin-17-Producing gammadelta T Cells Originate from SOX13(+) Progenitors that Are Independent of gammadeltaTCR Signaling. *Immunity, 49*(5), 857-872 e855. doi:10.1016/j.immuni.2018.09.010

Street, K., Risso, D., Fletcher, R. B., Das, D., Ngai, J., Yosef, N., . . . Dudoit, S. (2018). Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics, 19*(1), 477. doi:10.1186/s12864-018-4772-0

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, . . . Satija, R. (2019). Comprehensive Integration of Single-Cell Data. *Cell, 177*(7), 1888-1902 e1821. doi:10.1016/j.cell.2019.05.031

Sumaria, N., Grandjean, C. L., Silva-Santos, B., & Pennington, D. J. (2017). Strong TCRgammadelta Signaling Prohibits Thymic Development of IL-17A-Secreting gammadelta T Cells. *Cell Rep, 19*(12), 2469-2476. doi:10.1016/j.celrep.2017.05.071

Taghon, T., Yui, M. A., Pant, R., Diamond, R. A., & Rothenberg, E. V. (2006). Developmental and molecular characterization of emerging beta- and gammadelta-selected pre-T cells in the adult mouse thymus. *Immunity, 24*(1), 53-64. doi:10.1016/j.immuni.2005.11.012

Tanigaki, K., Tsuji, M., Yamamoto, N., Han, H., Tsukada, J., Inoue, H., . . . Honjo, T. (2004). Regulation of alphabeta/gammadelta T cell lineage commitment and peripheral T cell responses by Notch/RBP-J signaling. *Immunity, 20*(5), 611-622. doi:10.1016/s1074-7613(04)00109-8

Wolfer, A., Wilson, A., Nemir, M., MacDonald, H. R., & Radtke, F. (2002). Inactivation of Notch1 impairs VDJbeta rearrangement and allows pre-TCR-independent survival of early alpha beta Lineage Thymocytes. *Immunity, 16*(6), 869-879. doi:10.1016/s1074-7613(02)00330-8

Yagi, R., Junttila, I. S., Wei, G., Urban, J. F., Jr., Zhao, K., Paul, W. E., & Zhu, J. (2010). The transcription factor

GATA3 actively represses RUNX3 protein-regulated production of interferon-gamma. *Immunity, 32*(4),

507-517. doi:10.1016/j.immuni.2010.04.004

Zheng, G. X., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., . . . Bielas, J. H. (2017). Massively

parallel digital transcriptional profiling of single cells. *Nat Commun, 8*, 14049. doi:10.1038/ncomms14049

Zuberbuehler, M. K., Parker, M. E., Wheaton, J. D., Espinosa, J. R., Salzler, H. R., Park, E., & Ciofani, M. (2019).

The transcription factor c-Maf is essential for the commitment of IL-17-producing gammadelta T cells.

Nat Immunol, 20(1), 73-85. doi:10.1038/s41590-018-0274-0

Zwiener, I., Frisch, B., & Binder, H. (2014). Transforming RNA-Seq data to improve the performance of

prognostic gene signatures. *PLoS One, 9*(1), e85150. doi:10.1371/journal.pone.0085150

756