1

Ex Vivo Herpes Simplex Virus Reactivation Involves a DLK-Dependent Wave of Lytic Gene

Expression that is Independent of Histone Demethylase Activity and Viral Genome Synthesis

Abigail L. Whitford[1], Corinne A. Clinton[1], E.B. Lane Kennedy[1], Sara A. Dochnal[1], Jon B.

Suzich[1] and Anna R. Cliffe[1,*]

1. Department of Microbiology, Immunology and Cancer Biology, University of Virginia,

Charlottesville, VA, 22908.

* Correspondence to Anna R. Cliffe, cliffe@virginia.edu

Running title: HSV-1 DLK-dependent Phase I reactivation ex vivo

Abstract word count: 211

Text word count: 4833

Abstract

Herpes Simplex virus-1 (HSV-1) maintains a lifelong latent infection in neurons and periodically reactivates, resulting in the production of infectious virus. The exact cellular pathways that induce reactivation are not understood. In primary neuronal models of HSV latency, the cellular protein Dual Leucine Zipper kinase (DLK) has been found to initiate a wave of viral gene expression known as Phase I. Phase I occurs independently of both viral DNA replication and the activities of histone demethylase enzymes required to remove repressive heterochromatin modifications associated with the viral genome. Here we investigated whether Phase-I like gene expression occurs in ganglia reactivated from infected mice. Using the combined trigger of explant-induced axotomy and inhibition of PI3K signaling, we found that HSV lytic gene expression was induced rapidly from both sensory and sympathetic neurons. *Ex vivo* reactivation involved a wave of viral late gene expression that occurred independently of viral genome synthesis and histone demethylase activity, and preceded the detection of infectious virus. Importantly, we found that DLK was required for the initial induction of lytic gene expression. These data confirm the essential role of DLK in inducing HSV-1 gene expression from the heterochromatin associated genome and further demonstrate that HSV-1 gene expression during reactivation occurs via mechanisms that are distinct from lytic replication.

Introduction

The ubiquitous human pathogen Herpes Simples Virus persists for life in the form of a latent infection in neurons. In response to a variety of different stimuli, the virus can reactivate from the latent state, resulting in the release of infectious virus and subsequent replication in the surrounding tissue. Clinically, reactivation of the virus can manifest as a variety of disease states including lesions at the body surface, keratitis, and encephalitis. In addition, there is growing evidence of a link between HSV infection and the development of late onset Alzheimer's disease, particularly in individuals with the ApoE4 variant (1-9). There are potentially different stimuli that can induce HSV to reactivate from latency. These stimuli may converge on single cellular pathways or operate via distinct mechanisms to induce reactivation (10). Identifying the cellular pathways important in reactivation is required to understand how viral gene expression initiates from the latent genome and to ultimately develop therapeutics to prevent reactivation occurring.

During a latent infection of neurons, the HSV genome is assembled into repressive heterochromatin. This has been characterized by the enrichment of post-translational modifications on histone H3; namely di- and tri-methyl lysine 9 (H3K9me2/3) and tri-methyl lysine 27 (H3K27me3) on lytic promoters (11-17). By assembling into heterochromatin, viral lytic transcripts are maintained in a silent state. In addition, both host and viral miRNAs target lytic mRNAs (18-22). Therefore, the action of transcriptional and translational silencing results in limited synthesis of viral lytic proteins. This lack of lytic proteins suggests that HSV is reliant upon host signaling to initiate gene expression and reactivation. To understand the mechanism of HSV reactivation, it is important to determine how activation of host cell pathways ultimately converge on the repressed viral genome to induce lytic gene expression.

There is evidence that the initial induction of HSV-1 lytic gene expression following

reactivation occurs in a manner that is distinct from the mechanisms of viral gene expression

during lytic replication. HSV lytic genes can be divided into groups characterized by their

requirements for viral protein synthesis and viral DNA replication during lytic replication (23).

Immediate early (IE) genes are expressed independently viral protein synthesis during lytic

infection and instead require the viral tegument protein VP16 for maximal expression. Early

(E) genes are expressed following the production of IE proteins. Certain IE proteins (including

ICP4, ICP0 and ICP27) stimulate viral E protein synthesis. Late (L) genes require viral DNA

synthesis and are subdivided into genes that are expressed at low levels even prior to DNA

replication but increased with genome synthesis (leaky L) and those that are fully dependent

on viral DNA replication (true L). The dependence on DNA replication for L gene expression

is not fully understood but likely involves a shift in genome accessibility and increased binding

of host transcriptional machinery (RNA Pol II, TBP, and TAF1) (24). In contrast to this

regulated cascade, in models of HSV reactivation, an initial burst of lytic gene expression,

named Phase I of reactivation, has been observed where inhibition of protein synthesis prior

to the accumulation of IE transcripts does not prevent E gene expression (25). In addition, L

gene expression is unaffected by inhibition of viral DNA replication (25). Together, these

data indicate that initial expression of lytic transcripts during the early stages of reactivation

does not resemble the early stages of *de novo* infection in non-neuronal cells.

Phase I of HSV reactivation has largely been identified in primary neuronal models of

HSV latency (26). In these experimental models, a variety of stimuli have been found to

induce HSV to reactivate from a latent infection, including loss of neurotrophic factor support

(25, 27-29), increased neuronal excitation (30), modulation of DNA damage/repair (31) and

exposure to corticosteroids (27). Precisely how activation of these pathways permits

expression of the viral gene transcripts for reactivation to occur is not fully understood.

Previously, we have identified a role for the cell stress protein, dual leucine zipper kinase

(DLK), in inducing Phase I of HSV reactivation (27). DLK is activated by the loss of

neurotrophic factor support, which can be mimicked by inhibition of Phosphoinositide 3 (PI3)-

kinase activity (27), and during heightened neuronal excitation and interleukin-1 (IL-1)

treatment (30). DLK is a master regulator of axonal responses to stress and can mediate a

variety of responses including Wallerian degeneration, axon regeneration, apoptosis, and

axon pruning (32). Upon activation, DLK is known to re-direct the cell stress protein, c-Jun N-

terminal kinase (JNK), from its physiological role in neurons, maintaining synaptic

arborization, to its cell stress function (33). Accordingly, we and others have identified a role

for JNK in reactivation of HSV from latency (27, 30, 31). JNK is also important in reactivation

of the related Alpha herpesvirus, Varicella-Zoster virus, from a latent infection (34)

highlighting its potential central role in reactivation of human Alpha herpesviruses.

A role for DLK and JNK in HSV reactivation has been mostly widely studied on primary

neuronal models of latency in murine sympathetic neurons (27, 30). In these neurons, latency

is established in the presence of the HSV DNA replication inhibitor, acyclovir. After the

removal of acyclovir, reactivation can be induced by PI3-kinase inhibition or forskolin (25, 27,

30, 35, 36). In these systems, full reactivation occurs around 48h post-stimulus, which

requires the activities of histone demethylase enzymes, indicating that reactivation requires

removal of repressive heterochromatin (27, 30). However, the DLK-dependent Phase I peaks

around 18h post-stimuli and (27, 30, 37), importantly, Phase I occurs independently of lysine

9 and lysine 27 histone demethylase activity (27, 30). Instead, JNK activation induces histone

phosphorylation on H3Serine10 (H3S10) on histones that maintain the H3K9me3

modification (27); this is known as a histone methyl/phospho switch and presumably permits

transcription by overriding the repressive H3K9me3 modification. However, although

demonstrated *in vitro*, the possibility of a DLK/JNK-dependent wave of gene expression,

characteristic of Phase I, has not been explored following *in vivo* infections.

A robust mode of HSV reactivation from infected mice is explanation of the sensory

trigeminal ganglia (TG). The action of severing the axon (axotomy), is thought to be the

trigger that induces HSV to reactivate. In this model, whether a Phase I-like wave of gene

expression occurs has not been fully explored. Data from a previous study suggests that a

H3 lysine 9 demethylase functions to promote lytic gene expression at an early time-point (6

hours post-explant) (38). This indicates that wave of gene expression that is independent of

histone demethylase activity occurs at an earlier time-point, or that explant induced

reactivation does not involve a Phase-I like wave of viral gene expression. There are multiple

differences between the *in vivo* experiments and *in vitro* latency models, including the

presence of host immune system *in vivo*. In addition, explant-induced reactivation has been

investigated mostly in sensory neurons whereas models for *in vitro* infection often use

sympathetic neurons; both neuronal types are targets of HSV latent infection in humans (2,

39-41). *In vitro* models also use acyclovir to promote the establishment of latency. Finally,

potential differences resulting from viral strains used cannot be ruled out. In experiments

investigating the role for histone demethylases *ex vivo*, HSV strain F was used (16, 38),

whereas the in vitro models have been performed with the KOS and Patton strains (25, 27).

To determine whether previous observations of a DLK/JNK triggered Phase I

reactivation *in vitro* were recapitulated e*x vivo,* we dissected trigeminal ganglia from latently

infected mice and determined whether we observed lytic gene expression in response to PI3-

kinase inhibition. We found that after only five hours post excision, in treated ganglia, there

was robust expression of IE, E, and L viral genes and this gene expression was independent

of viral DNA replication. Supporting previous findings, we also found that this initial burst of

lytic gene expression was not dependent upon LSD1 (H3K9-demethylase) nor JMJD3 and

UTX (H3K27-demethylases). Therefore, Phase I of reactivation was observed *ex vivo* and

was not reliant upon the removal of repressive heterochromatic marks. We found that this

Phase 1 was dependent upon DLK, thus indicating that neuronal stress pathways can trigger

bi-phasic reactivation of HSV-1 in ganglia *ex vivo*.

Materials Methods

Cells and Viruses

HSV-1 stocks of KOS were grown and titrated on Vero cells obtained from the American Type Culture Collection (Manassas, VA) as described previously (11). Cells were maintained in Dulbecco's Modified Eagle's Medium (Gibco) supplemented with 10% FetalPlex (Gemini Bio-Products) and 2 mM L-Glutamine. KOS was kindly provided by Dr. David Knipe, Harvard Medical School.

Reagents

Compounds used in the study are as follows: Acycloguanosine (acyclovir; ACV), GSK-J4 and GNE-3511(Millipore Sigma) SP600125 (Thermo Fisher Scientific), OG-L002 (Tocris); Nerve Growth Factor 2.5S (Alomone Labs).

Mouse Infections

Six-week-old male and female CD-1 mice (Charles River Laboratories) were anesthetized by intraperitoneal injection of ketamine hydrochloride (80mg/kg) and xylazine hydrochloride (10mg/kg) and inoculated with 1.5×10^6 PFU/eye of virus (in a 5µl volume) onto scarified corneas, as described previously (11). Mice were housed in accordance with institutional and National Institutes of Health guidelines on the care and use of animals in research, and all procedures were approved by the Institutional Animal Care and Use Committee of the University of Virginia. Criteria used for clinical scoring based on the formation of lesions, neurological and eye symptoms is shown in table 1 and based on a previously establishing scoring scale (42). Mice were randomly assigned to groups and all experiments included biological repetitions from independent litters.

Table 1. Clinical scoring scale for HSV infected mice.

Score	Description of score by type		
	Lesion	Eye	Neurological
0	Lesion free	Symptom free	Symptom free
1	Small area of broken skin <0.5cm	Pus around edges	Hunched posture, normal movement
2	Area of broken skin 0.5-1 cm	Pus and squint	Hunched posture, slow movement
3	Bleeding, scabbing or pustules	Closed	Hunched posture, labored breathing and/or hind leg paralysis
4	Broken skin >cm with multiple pustules or scabbing	Scab formation	Hunched, ruffled fur, little to no movement
5	Sever scabbing or bleeding with pustules	Sever scabbing	Moribund or dead

179
180

Explant Induced Reactivation

Trigeminal ganglia (TG) and superior cervical ganglia (SCG) were removed at least 28 days

post-infection. The ganglia were maintained intact and immediately placed in reactivation

media containing DMEM/F12 (Gibco) supplemented with 10% Fetal Bovine Serum and

Mouse NGF 2.5S (50 ng/mL). Compounds were added at the following concentrations;

LY294002 40μM, GNE-3511 8μM, ACV 100μM, GSK-J4 10μM, OG-L002 50μM. For

treatments with GNE-5311, ACV, GSK-J4 and OG-L002, ganglia were treated with the

compounds 3-60 minutes prior to the addition of LY294002. Ganglia were placed on a

shaking platform at 50rpm at 37°C.

Quantification of Viral Transcripts and Genome Copy Number

At the required time point, the reactivation media was removed and ganglia snap-frozen in

liquid nitrogen. Lysis was carried out by addition of the excised ganglia to BeadBug^{TM}

homogenization microtubules then homogenized for 60 seconds using the BeadBug^{TM}

microtube homogenizer. DNA and RNA were isolated from the homogenized mixture using

the *Quick* DNA/RNA miniprep kit. Following RNA isolation TURBO DNA-free Kit (Invitrogen)

was used to remove any contaminant DNA. mRNA was reverse transcribed into cDNA using

MaximaRT (ThermoFisher) using random hexamers for first strand synthesis. Equal amounts

of mRNA were used for each reverse transcriptase experiment (20-30 ng/reaction). Power

SYBR Green PCR Master Mix was used for qPCR (Applied Biosystems). Standard Curves

were used to calculate the relative mRNA or DNA copy number per genome and were

standardized to cellular GAPDH or 18s. All RNA samples were run in triplicate and all DNA

samples were run in duplicate on an Applied Biosystems™ QuantStudio™ 6 Flex Real-Time

PCR System. The primers that were used are described in Table 2.

Table 2. Oligonucleotides used in this study

Primer	Sequence 5′-3′
GAPDH F	CAT GGC CTT CCG TGT TCC TA
GAPDH R	GCG GCA CGT CAG ATC CA
18s F	CAC GGA CAG GAT TGA CAG ATT
18s R	GCC AGA GTC TCG TTC GTT ATC
ICP27 F	GCA TCC TTC GTG TTT GTC ATT CTG
ICP27 R	GCA TCT TCT CTC CGA CCC CG
ICP8 F	GGA GGT GCA CCG CAT ACC
ICP8 R	GGC TAA AAT CCG GCA TGA AC
gC F	GAG TTT GTC TGG TTC GAG GAC
gC R	ACG GTA GAG ACT GTG GTG AA

209 <u>Preformed virus titration</u>

210 Sterile milk was added to media containing reactivated ganglia and sample snap frozen.

211 Ganglia were thawed at 37°C then homogenized using a BeadBug microtube homogenizer.

212 The homogenized ganglia were then sonicated at 25% for 30 seconds three times on ice in a

213 Fisherbrand™ Model 120 Sonic Dismembrator. After sonication, ganglia homogenates were

214 frozen on dry ice and thawed at 37°C. Homogenates from all time points were simultaneously

215 titrated on Vero cells.

Results

Explant combined with PI3-kinase inhibition triggers robust HSV-1 lytic gene expression

We first set out to determine whether a wave of lytic gene expression occurs following

reactivation of explanted ganglia, similar to what has been observed in primary neuronal

cultures. Female mice were infected via the ocular route and reactivation studies were

carried out at least 28 days post-infection. Previously, studies from our lab and others have

shown that addition of a reactivation stimulus to sympathetic neurons infected *in vitro* yields

induction of lytic viral gene expression around 15-20 hours post-stimulus (25, 27). In addition,

a previous study examining explant induced reactivation combined with deprivation of nerve

growth factor (NGF) also resulted in lytic gene induction around 12-15 hours post-explant

(43). Therefore, we initially examined the effect of PI3-K inhibition (using LY294002; 40 µM)

at 20 hours post-reactivation to determine whether loss of the PI3K/AKT branch of the NGF-

signaling pathway also promotes reactivation *ex vivo*. Acyclovir (ACV; 100 µM) was also

introduced alongside LY294002 because late gene expression during the initial activation of

viral gene expression in *in vitro* models has been reported to proceed independently of DNA

replication (25).

In ganglia that were explanted and maintained in NGF to provide continued

neurotrophin support, very little induction of lytic gene expression was observed at 20h post-

explant, especially for representative IE and E transcripts. A slight increase in *gC* mRNA was

observed, although this increase was not statistically significant (P=0.1255, Mann Whitney U

Test). However, the addition of the PI3-kinase inhibitor (LY294002) increased viral gene

expression by 100-1000-fold. RT-qPCRs were carried out using primers against IE (ICP27,

Fig.1 B), E (ICP8, Fig. 1C) and L (gC, Fig. 1D) genes. All three genes were significantly

induced at 20h post excision. The addition of ACV to inhibit viral DNA replication did not

inhibit the induction of IE or E gene expression, as expected. However, expression of the gC

mRNA was significantly reduced in the presence of ACV, indicating that its maximal

expression was dependent on viral genome synthesis. However, gC mRNA levels with ACV

were still significantly increased compared to the latent samples. At 20h post-reactivation, an

increase in viral genomes was also observed (Fig. 1A). Together, these data show that at

20h post-reactivation viral gene expression does not resemble the Phase I observed in

primary neuronal cultures, as genome synthesis has occurred, and full late gene expression

is dependent on viral DNA replication. However, this did not rule out the possibility of a Phase

I-like wave of gene expression occurring prior to 20h, especially as *gC* mRNA was still

induced compared to the 0h (latent) samples, indicating that there may still be some *gC*

expression occurred even from non-replicated viral genomes.

<u>PI3K inhibition/explant induces rapid late gene expression in the absence of detectible</u>

<u>genome synthesis in trigeminal ganglia</u>

To determine whether viral lytic gene expression occurred prior to 20h post-stimulus in

explanted ganglia with features characteristic of Phase I gene expression, we decided to

undertake both male and female mice and determine changes in lytic gene expression

following PI3-kinase inhibition of explanted ganglia. Because we infected both male and

female mice, we assessed if sex impacted clinical manifestations of infection, mice were

monitored throughout the period of infection. Both male and female mice had minimal

mortality with only 92 and 88 percent mortality respectively. The mortality rates between

sexes were not found to be significant by Kaplan-Meier survival analysis (Fig. 2A). To further

analyze clinical symptoms, mice were evaluated by scoring lesions, neurological symptoms

and eye health based on a previously described scoring metric (42). Infection of female

mouse resulted in more severe clinical manifestations at days 7 and 9 post-infection (Fig.

2B). Despite these differences, there was no significant differences in the percentage of

weight loss following infection between male and female mice (Fig. 2C). Based on these

criteria, we concluded there were only minor sex-dependent differences upon HSV infection.

After at least 28 days post-infection, ganglia were explanted and incubated in the

presence of the PI3-kinase inhibitor, LY294002, with and without ACV. Values from both

male and female mice were combined (Fig. 3A-D). Quantification of viral DNA loads showed

that the copy number of viral genomes stayed constant up to 15 hours post-stimuli and were

not affected by the presence of ACV, indicating that detectible viral genome synthesis did not

occur in this time period. In contrast, a robust induction of viral lytic gene expression occurred

by 5h post-stimuli, as inducted by an increase in IE (ICP27), E (ICP8) and late (gC) mRNA

copy number. For all gene classes examined, the increase in copy number was 100-fold for

IE mRNA (Fig. 3B), 20-fold for E mRNA (Fig. 3C), and 10-fold for L mRNA (Fig. 3D) 5 hours

post explant. The robust increase in late gene expression in the absence of detectible viral

genome synthesis suggested that late gene expression could occur even prior to DNA

replication. To further support this conclusion, late gene induction occurred to equivalent

levels even in the presence of ACV. The inclusion of ACV did prevent genome synthesis at

20h post-stimuli (Fig. 1A), indicating that the ACV was capable of acting on the explanted

ganglia. Therefore, these data indicate that the induction of lytic gene expression following

PI3-kinase inhibition in explanted sensory neurons resembles at least one feature of Phase I

gene expression as viral late gene expression occurred independently of viral DNA

replication.

To confirm that reactivation had no sex-dependent effects we performed quantification

of viral lytic gene expression in both male and female mice. We found no difference in viral

genome copy number (Fig. 3E) or ICP27 lytic gene expression (Fig. 3F) at any time point. We

therefore concluded that the observed phenotypes in lytic gene expression were sex-

independent.

<u>PI3-kinase Inhibition in Combination with Axotomy Triggers Rapid Lytic Gene Expression in</u>

<u>Sympathetic Neurons *Ex Vivo*</u>

Previous studies investigating Phase I gene expression *in vitro* have largely used

sympathetic neurons, where the induction of lytic gene expression has been found to occur

15-20h post stimuli (25-27). However, we found that lytic gene expression induced ex vivo

from sensory neurons was robustly induced by 5h post-stimulus (Fig 2). Therefore, to

determine whether the enhanced kinetics of reactivation observed *ex vivo* from sensory

neurons in the trigeminal ganglia could result from the use of different neuronal types, we

investigated reactivation *ex vivo* from the superior cervical ganglia (SCGs). Latently infected

SCGs were explanted and incubated in the presence of the PI3-kinase inhibitor, LY294002.

Quantification of viral DNA loads revealed that the copy number of viral genomes remained

constant at 5 hours after stimulation, indicating that no detectable viral genome synthesis

occurred during this time (Fig. 4A). In concordance with our sensory ganglia data (Fig. 1),

there was a significant 10-fold increase in viral genome copy number at 20 hours post

excision (Fig. 4A). By 5h post-stimuli a robust stimulation of viral lytic gene expression had

occurred, as evidenced by a 10-fold increase in IE (*ICP27,* Fig 4B) and a 20-fold increase in

late (*gC*) mRNA copy numbers (Fig. 4C). This indicates that the increased kinetics observed

ex vivo in sensory neurons was not due to different neuronal subtypes and instead likely

results from either the combination of triggers or as a result of *in vivo* infection.

Analysis of preformed virus production in sensory neurons *ex vivo*

Based on our data that viral late gene expression was detectible by 5 hours and robustly expressed by 10 hours post-explant, we investigated when preformed infectious virus could be detected in the explanted ganglia. The rationale for this was that Phase I lytic gene expression occurs in cultured neurons before the production of *de novo* virus (25). To ensure that we detected only preformed virus and not virus produced from remaining intact cells, the ganglia were homogenized and subjected to three rounds of sonication and two cycles of freezing and thawing. Using a viral stock of known titer, we confirmed that that this procedure did not result in a detectible loss in viral titer (data not shown). Infectious virus was robustly detected at 20h post-explant in seven out of the eight ganglia tested (Fig. 5). At 15 hours, virus was only detected in four out of eight ganglia tested and at 10 hours only two ganglia had detectible virus. No infectious virus was detected at 7 hours post-explant. These findings suggest that a detectable increase in late gene expression occurs 10 to 15 hours before infectious virus production, observed at 20 hours, in a population of reactivating ganglia.

DLK activity is required for reactivation *ex vivo*

Previously, we have found that the neuronal cell stress protein dual leucine zipper kinase (DLK) is required for HSV reactivation and acts to induce Phase I lytic gene expression. To determine whether DLK was also required for induction of lytic gene expression in sensory neurons reactivated *ex vivo* by PI3K-inhibition/axotomy, the DLK inhibitor GNE-3551 was added to the explanted ganglia and viral RNA was extracted at 5h post-reactivation. Inclusion of the DLK-inhibitor resulted in *ICP27*, *ICP8*, and *gC* mRNA (Fig. 6 A-C) levels that were equivalent to the un-reactivated samples and significantly decreased

compared to the reactivated ganglia that were not treated with the DLK-inhibitor. These data

indicate that DLK is required for reactivation from sensory neurons induced by the combined

trigger of explant and PI3-kinase inhibition.

<u>Lytic gene induction upon axotomy along with PI3-kinase inhibition is independent of histone</u>

<u>H3 histone lysine 9 and lysine 27 demethylase inhibitors</u>

HSV promoters are known to be enriched with histone H3 di- and trimethyl at lysine 9

(H3K9me2/3) and histone H3 trimethyl at lysine 27 (H3K27me3) during latency. The removal

of restrictive histone modifications, specifically the aforementioned methylation marks, was

previously shown to be essential for full reactivation of HSV *in vitro,* yet was not required for

initial lytic gene expression during Phase I of reactivation (27, 30). To determine if initial lytic

gene expression was independent of histone demethylation *ex vivo,* we used OG-L002 (60

µM), a drug that inhibits the histone lysine 9 demethylase LSD1, and GSK-J4 (20 µM), a

molecule that inhibits the histone lysine 27 demethylases UTX and JMJD3. HSV reactivation

has been demonstrated to be inhibited by both of these inhibitors (27, 30, 44, 45). Five hours

post excision, during Phase I of reactivation, the addition of OG-L002 and GSK-J4 had no

effect on the induction of ICP27, ICP8, or gC mRNA (Fig. 7A-C). These results indicate that

histone demethylase activity is not required for the initial induction of lytic gene expression

induced by axotomy and PI3-kinase inhibition *ex vivo.*

Together, our results demonstrate that features of Phase I gene expression, namely

late gene expression in the absence of viral DNA replication and lytic gene expression in the

presence of histone demethylase inhibitors, can occur from ganglia isolated from infected

mice. In addition, *ex vivo* reactivation was dependent on DLK activity.

364
<u>Discussion</u>

In vitro models of HSV latency are incredibly powerful for the study of molecular

mechanisms HSV latency and reactivation as well as the contribution of host factors and viral

factors during specific stages of infection. *In vitro* models can be used to investigate how the

host immune response can modulate latent infection and reactivation. Using these models,

unique aspects regarding entry into lytic gene expression during reactivation have been

uncovered, including the dependence on DLK/JNK for reactivation, the ability of late gene

expression to occur in the absence of DNA replication, and transcription despite the presence

of histone demethylase inhibitors. Therefore, it was highly important to validate these patterns

of gene expression from ganglia infected *in vivo*. Here we have confirmed that the features of

gene expression observed during Phase I reactivation *in vitro* also occur following

reactivation *ex vivo* when PI3k-inhibition and axotomy is used as a trigger.

There are certain caveats to our study, especially relating to the ability of inhibitors to

act on intact ganglia. However, we show that the DLK inhibitor, PI3-kinase inhibitor and

acyclovir can all act on intact ganglia. Therefore, we think it unlikely that the histone

demethylase inhibitors used were unable to penetrate the tissue and have an effect on viral

lytic gene induction. The concentrations of histone demethylase inhibitors were higher than

what we have previously observed to inhibit full HSV reactivation in primary neuronal models

(27, 30). We have shown that OG-L002 can inhibit forskolin mediated reactivation at 30 μM,

here we used 60 μM. Similarly, we have shown that GSK-J4 can inhibit both forskolin and

LY294002 induced reactivation at 3 μM; here we used it at 20 μM. Therefore, we think it

unlikely that the concentrations used were too low to inhibit the histone demethylases. We did

try higher concentrations that resulted in loss of viral genomes at 5h, indicating potential

neuronal toxicity at these higher concentrations.

Our study did result in the discovery of a unique phenotype of Phase I of reactivation

when observed *ex vivo*. Unlike our previous *in vitro* studies, we found that Phase I of

reactivation that occurred *ex vivo* had enhanced kinetics of lytic gene induction upon axotomy

and PI3-kinase stimulation. We observed this enhanced phenotype in both sensory and

sympathetic ganglia. A paralleled kinetic phenotype in both neuronal subtypes was validated

in a novel model of latency and reactivation utilizing a virus defective in cell spread, in which

our lab discovered that reactivation kinetics *in vitro* were equivalent in sensory and

sympathetic neurons (Dochnal et al, manuscript in preparation). Therefore, we posit that the

most likely explanation of the difference in reactivation kinetics between *in vitro* and *ex vivo*

studies is the combined trigger of axotomy and PI3-kinase inhibition. It has previously been

demonstrated that when nerve growth factor deprivation is paired with axotomy, treated

neurons undergo cell death more quickly than untreated neurons. (46). In addition, both PI3-

kinase inhibition and axotomy result in activation of DLK (32, 47). However, there are some

differences in how DLK is activated and the resulting downstream response in response to

the two stimuli. Following axotomy, DLK is rapidly activated at the proximal segment of the

severed axon in response to loss of cytoplasmic integrity and a calcium influx results from

cytoplasmic membrane rupture to ultimately result in transcription of genes involved in axon

regeneration (48). The exact mechanisms of DLK activation upon loss of NGF signaling that

mediates neuronal cell death or axon pruning is not fully understood. It is known that in

response to NGF deprivation, DLK protein levels are stabilized, DLK is phosphorylated (49),

and this phosphorylation results in the downstream activation of JNK, c-Jun, and other

transcription factors. The activation of these transcription factors promotes the expression of

pro-apoptotic genes (50), although in mature neurons there are multiple brakes downstream

that prevent apoptosis (51). It is therefore conceivable that these two independent pathways

to DLK activation may converge follow the dual trigger of axotomy/PI3-kinase inhibition to

result in more rapid and robust DLK activation and HSV reactivation.

Consistent with cellular pathways converging the enhance DLK activity, we observed a

robust induction of lytic gene expression at 5h post reactivation. For the IE gene ICP27 this

was an approximate 100-fold induction, compared to the 5-20-fold increase often observed in

an *in vitro* mouse model with PI3-kinase inhibition (27). The use of an *ex vivo* model system

will therefore be incredibly powerful for studying the mechanisms of gene expression that

occur during Phase I. We have previously observed a histone methyl/phospho switch on lytic

promoters during Phase I reactivation (27), which permits lytic gene expression

independently of recruitment of histone demethylase enzymes. There is evidence that the

methyl/phospho switch permits gene expression because repressive histone readers (for

example HP1) are no longer capable of interacting with the methylated residue on histone

H3K9 because phosphorylation at Serine 10 occludes binding (52, 53). Because of the robust

lytic gene induction *ex vivo*, and because we have now shown that the gene expression both

in vitro and *ex vivo* is DLK-dependent and histone demethylase independent, the

complementary models can now be used answer key questions on the mechanism of viral

lytic gene induction during Phase I reactivation.

<u>Acknowledgements</u>

We thank David Knipe (Harvard Medical School) for the KOS virus. This work was supported

by R01NS105630 (ARC), The Owens Family Foundation (ARC), NIH/NIAID T32AI007046

(ALW, JBS), NIH/NEI F30EY030397 (JBS) and NIH/NIGMS T32GM008136 (SAD).

References

1.   Jamieson GA, Maitland NJ, Wilcock GK, Craske J, Itzhaki RF. 1991. Latent herpes

simplex virus type 1 in normal and Alzheimer's disease brains. J Med Virol 33:224-7.

2.   Baringer JR, Pisani P. 1994. Herpes simplex virus genomes in human nervous system

tissue analyzed by polymerase chain reaction. Ann Neurol 36:823-9.

3.   Itzhaki RF, Lin WR, Shang D, Wilcock GK, Faragher B, Jamieson GA. 1997. Herpes

simplex virus type 1 in brain and risk of Alzheimer's disease. Lancet 349:241-4.

4.   Mori I, Kimura Y, Naiki H, Matsubara R, Takeuchi T, Yokochi T, Nishiyama Y. 2004.

Reactivation of HSV-1 in the brain of patients with familial Alzheimer's disease. J Med

Virol 73:605-11.

5.   Piacentini R, De Chiara G, Li Puma DD, Ripoli C, Marcocci ME, Garaci E, Palamara

AT, Grassi C. 2014. HSV-1 and Alzheimer's disease: more than a hypothesis. Front

Pharmacol 5:97.

6.   Itzhaki RF. 2018. Corroboration of a Major Role for Herpes Simplex Virus Type 1 in

Alzheimer's Disease. Front Aging Neurosci 10:324.

7.   De Chiara G, Piacentini R, Fabiani M, Mastrodonato A, Marcocci ME, Limongi D,

Napoletani G, Protto V, Coluccio P, Celestino I, Li Puma DD, Grassi C, Palamara AT.

2019. Recurrent herpes simplex virus-1 infection induces hallmarks of

neurodegeneration and cognitive deficits in mice. PLoS Pathog 15:e1007617.

8.   Tzeng NS, Chung CH, Lin FH, Chiang CP, Yeh CB, Huang SY, Lu RB, Chang HA,

Kao YC, Yeh HW, Chiang WS, Chou YC, Tsao CH, Wu YF, Chien WC. 2018. Anti-

herpetic Medications and Reduced Risk of Dementia in Patients with Herpes Simplex

Virus Infections-a Nationwide, Population-Based Cohort Study in Taiwan.

Neurotherapeutics 15:417-429.

9.   Readhead B, Haure-Mirande JV, Funk CC, Richards MA, Shannon P, Haroutunian V,

Sano M, Liang WS, Beckmann ND, Price ND, Reiman EM, Schadt EE, Ehrlich ME,

Gandy S, Dudley JT. 2018. Multiscale Analysis of Independent Alzheimer's Cohorts

Finds Disruption of Molecular, Genetic, and Clinical Networks by Human Herpesvirus.

Neuron 99:64-82 e7.

10.   Suzich JB, Cliffe AR. 2018. Strength in diversity: Understanding the pathways to

herpes simplex virus reactivation. Virology 522:81-91.

11.   Cliffe AR, Garber DA, Knipe DM. 2009. Transcription of the herpes simplex virus

latency-associated transcript promotes the formation of facultative heterochromatin on

lytic promoters. J Virol 83:8182-90.

12.   Knipe DM, Cliffe A. 2008. Chromatin control of herpes simplex virus lytic and latent

infection. Nat Rev Microbiol 6:211-21.

13.   Wang QY, Zhou C, Johnson KE, Colgrove RC, Coen DM, Knipe DM. 2005.

Herpesviral latency-associated transcript gene promotes assembly of heterochromatin

on viral lytic-gene promoters in latent infection. Proc Natl Acad Sci U S A 102:16055-9.

14.   Kwiatkowski DL, Thompson HW, Bloom DC. 2009. The polycomb group protein Bmi1

binds to the herpes simplex virus 1 latent genome and maintains repressive histone

marks during latency. J Virol 83:8173-81.

15.   Nicoll MP, Hann W, Shivkumar M, Harman LE, Connor V, Coleman HM, Proenca JT,

Efstathiou S. 2016. The HSV-1 Latency-Associated Transcript Functions to Repress

Latent Phase Lytic Gene Expression and Suppress Virus Reactivation from Latently

Infected Neurons. PLoS Pathog 12:e1005539.

16.   Liang Y, Vogel JL, Narayanan A, Peng H, Kristie TM. 2009. Inhibition of the histone

demethylase LSD1 blocks alpha-herpesvirus lytic replication and reactivation from

latency. Nat Med 15:1312-7.

17.   Dochnal SA, Francois AK, Cliffe AR. 2021. De Novo Polycomb Recruitment: Lessons

from Latent Herpesviruses. Viruses 13.

18.   Umbach JL, Kramer MF, Jurak I, Karnowski HW, Coen DM, Cullen BR. 2008.

MicroRNAs expressed by herpes simplex virus 1 during latent infection regulate viral

mRNAs. Nature 454:780-3.

19.   Jurak I, Kramer MF, Mellor JC, van Lint AL, Roth FP, Knipe DM, Coen DM. 2010.

Numerous conserved and divergent microRNAs expressed by herpes simplex viruses

1 and 2. J Virol 84:4659-72.

20.   Kramer MF, Jurak I, Pesola JM, Boissel S, Knipe DM, Coen DM. 2011. Herpes

simplex virus 1 microRNAs expressed abundantly during latent infection are not

essential for latency in mouse trigeminal ganglia. Virology 417:239-47.

21.   Jurak I, Hackenberg M, Kim JY, Pesola JM, Everett RD, Preston CM, Wilson AC,

Coen DM. 2014. Expression of herpes simplex virus 1 microRNAs in cell culture

models of quiescent and latent infection. J Virol 88:2337-9.

22.   Pan D, Flores O, Umbach JL, Pesola JM, Bentley P, Rosato PC, Leib DA, Cullen BR,

Coen DM. 2014. A neuron-specific host microRNA targets herpes simplex virus-1

ICP0 expression and promotes latency. Cell Host Microbe 15:446-56.

23.   Roizman B, Knipe DM, Whitley R. 2013. Herpes simplex viruses, 6th ed. Lippincott

Williams & Wilkins, Philadelphia.

24.   Dremel SE, DeLuca NA. 2019. Genome replication affects transcription factor binding

mediating the cascade of herpes simplex virus transcription. Proc Natl Acad Sci U S A

116:3734-3739.

25.   Kim JY, Mandarino A, Chao MV, Mohr I, Wilson AC. 2012. Transient reversal of

episome silencing precedes VP16-dependent transcription during reactivation of latent

HSV-1 in neurons. PLoS Pathog 8:e1002540.

26.   Cliffe AR, Wilson AC. 2017. Restarting Lytic Gene Transcription at the Onset of

Herpes Simplex Virus Reactivation. J Virol 91:e01419-16-6.

27.   Cliffe AR, Arbuckle JH, Vogel JL, Geden MJ, Rothbart SB, Cusack CL, Strahl BD,

Kristie TM, Deshmukh M. 2015. Neuronal Stress Pathway Mediating a Histone

Methyl/Phospho Switch Is Required for Herpes Simplex Virus Reactivation. Cell Host

Microbe 18:649-58.

28.   Wilcox CL, Johnson EM, Jr. 1987. Nerve growth factor deprivation results in the

reactivation of latent herpes simplex virus in vitro. J Virol 61:2311-5.

29.   Yanez AA, Harrell T, Sriranganathan HJ, Ives AM, Bertke AS. 2017. Neurotrophic

Factors NGF, GDNF and NTN Selectively Modulate HSV1 and HSV2 Lytic Infection

and Reactivation in Primary Adult Sensory and Autonomic Neurons. Pathogens 6:5-

13.

30.   Cuddy SR, Schinlever AR, Dochnal S, Seegren PV, Suzich J, Kundu P, Downs TK,

Farah M, Desai BN, Boutell C, Cliffe AR. 2020. Neuronal hyperexcitability is a DLK-

dependent trigger of herpes simplex virus reactivation that can be induced by IL-1.

Elife 9.

31.   Hu HL, Shiflett LA, Kobayashi M, Chao MV, Wilson AC, Mohr I, Huang TT. 2019.

TOP2beta-Dependent Nuclear DNA Damage Shapes Extracellular Growth Factor

Responses via Dynamic AKT Phosphorylation to Control Virus Latency. Mol Cell

74:466-480 e4.

32.   Tedeschi A, Bradke F. 2013. The DLK signalling pathway--a double-edged sword in

neural development and regeneration. EMBO Rep 14:605-14.

33.   Sengupta Ghosh A, Wang B, Pozniak CD, Chen M, Watts RJ, Lewcock JW. 2011.

DLK induces developmental neuronal degeneration via selective regulation of

proapoptotic JNK activity. The Journal of Cell Biology 194:751-764.

34. Kurapati S, Sadaoka T, Rajbhandari L, Jagdish B, Shukla P, Ali MA, Kim YJ, Lee G,
Cohen JI, Venkatesan A. 2017. Role of the JNK Pathway in Varicella-Zoster Virus
Lytic Infection and Reactivation. J Virol 91.
35. Camarena V, Kobayashi M, Kim JY, Roehm P, Perez R, Gardner J, Wilson AC, Mohr
I, Chao MV. 2010. Nature and duration of growth factor signaling through receptor
tyrosine kinases regulates HSV-1 latency in neurons. Cell Host Microbe 8:320-30.
36. Suzich JB, Cuddy SR, Baidas H, Dochnal S, Ke E, Schinlever AR, Babnis A, Boutell
C, Cliffe AR. 2021. PML-NB-dependent type I interferon memory results in a restricted
form of HSV latency. EMBO Rep 22:e52547.
37. Hu HL, Srinivas KP, Wang S, Chao MV, Lionnet T, Mohr I, Wilson AC, Depledge DP,
Huang TT. 2022. Single-cell transcriptomics identifies Gadd45b as a regulator of
herpesvirus-reactivating neurons. EMBO Rep 23:e53543.
38. Liang Y, Vogel JL, Arbuckle JH, Rai G, Jadhav A, Simeonov A, Maloney DJ, Kristie
TM. 2013. Targeting the JMJD2 histone demethylases to epigenetically control
herpesvirus infection and reactivation from latency. Sci Transl Med 5:167ra5.
39. Baringer JR, Swoveland P. 1973. Recovery of herpes-simplex virus from human
trigeminal ganglions. N Engl J Med 288:648-50.
40. Richter ER, Dias JK, Gilbert JE, 2nd, Atherton SS. 2009. Distribution of herpes
simplex virus type 1 and varicella zoster virus in ganglia of the human head and neck.
J Infect Dis 200:1901-6.
41. Warren KG, Brown SM, Wroblewska Z, Gilden D, Koprowski H, Subak-Sharpe J.
1978. Isolation of latent herpes simplex virus from the superior cervical and vagus
ganglions of human beings. N Engl J Med 298:1068-9.
42. Riccio RE, Park SJ, Longnecker R, Kopp SJ. 2019. Characterization of Sex
Differences in Ocular Herpes Simplex Virus 1 Infection and Herpes Stromal Keratitis

Pathogenesis of Wild-Type and Herpesvirus Entry Mediator Knockout Mice. mSphere

4.

43.   Du T, Zhou G, Roizman B. 2011. HSV-1 gene expression from reactivated ganglia is

disordered and concurrent with suppression of latency-associated transcript and

miRNAs. Proc Natl Acad Sci U S A 108:18820-4.

44.   Messer HG, Jacobs D, Dhummakupt A, Bloom DC. 2015. Inhibition of H3K27me3-

specific histone demethylases JMJD3 and UTX blocks reactivation of herpes simplex

virus 1 in trigeminal ganglion neurons. J Virol 89:3417-20.

45.   Liang Y, Quenelle D, Vogel JL, Mascaro C, Ortega A, Kristie TM. 2013. A novel

selective LSD1/KDM1A inhibitor epigenetically blocks herpes simplex virus lytic

replication and reactivation from latency. mBio 4:e00558-12.

46.   Fletcher GC, Xue L, Passingham SK, Tolkovsky AM. 2000. Death commitment point is

advanced by axotomy in sympathetic neurons. J Cell Biol 150:741-54.

47.   Wu CC, Wu HJ, Wang CH, Lin CH, Hsu SC, Chen YR, Hsiao M, Schuyler SC, Lu FL,

Ma N, Lu J. 2015. Akt suppresses DLK for maintaining self-renewal of mouse

embryonic stem cells. Cell Cycle 14:1207-17.

48.   Asghari Adib E, Smithson LJ, Collins CA. 2018. An axonal stress response pathway:

degenerative and regenerative signaling by DLK. Curr Opin Neurobiol 53:110-119.

49.   Huntwork-Rodriguez S, Wang B, Watkins T, Ghosh AS, Pozniak CD, Bustos D,

Newton K, Kirkpatrick DS, Lewcock JW. 2013. JNK-mediated phosphorylation of DLK

suppresses its ubiquitination to promote neuronal apoptosis. J Cell Biol 202:747-63.

50.   Watkins TA, Wang B, Huntwork-Rodriguez S, Yang J, Jiang Z, Eastham-Anderson J,

Modrusan Z, Kaminker JS, Tessier-Lavigne M, Lewcock JW. 2013. DLK initiates a

transcriptional program that couples apoptotic and regenerative responses to axonal

injury. Proc Natl Acad Sci U S A 110:4039-44.

51.   Kole AJ, Annis RP, Deshmukh M. 2013. Mature neurons: equipped for survival. Cell

588   Death Dis 4:e689.

589   52.   Fischle W, Wang Y, Allis CD. 2003. Binary switches and modification cassettes in

590   histone biology and beyond. Nature 425:475-9.

591   53.   Fischle W, Tseng BS, Dormann HL, Ueberheide BM, Garcia BA, Shabanowitz J, Hunt

592   DF, Funabiki H, Allis CD. 2005. Regulation of HP1-chromatin binding by histone H3

593   methylation and phosphorylation. Nature 438:1116-22.

Figure Legends

Figure 1. Explant combined with PI3-kinase inhibition triggers robust HSV-1 lytic gene

expression.

Mice were infected via corneal scarification and at least 28 days post infection trigeminal

ganglia (TG) were excised. Ganglia were either snap frozen at 0 hours, or reactivated for 20

hours in media alone, with LY294002 (40 μM) or LY294002 with acyclovir (ACV; 100μM). (A)

Viral genome copy number was quantified by qPCR. (B) Viral gene expression was quantified

by RT-qPCR for immediate early (*ICP27*), (C) early (*ICP8*) and (D) and late (*gC*) genes.

Transcript copy number was normalized to cellular control (*GAPDH*). n=6 biological

replicates. Mann-Whitney U (A-D). *p<0.05, **p<0.01. Individual biological replicates along

with the means and SEMs are represented.

Figure 2: Sex dependent phenotypes during HSV-1 *in vivo* infection.

Mice were infected via corneal scarification and monitored post infection. (A) Survival over

time was observed and the significance of the percentage survival between female and male

mice were analyzed using Kaplan-Meier survival analysis. (B) Clinicals scores were

calculated by scoring lesion, neurological, and eye phenotypes as outlined in table 1. (C)

Percent weight compared to pre-infection weights. Student T test (B-C). *p<0.05. The mean

and SEM are represented. N= 13 females, 8 males.

Figure 3: PI3K inhibition of explanted trigeminal ganglia induces rapid lytic gene expression in

the absence of detectible genome synthesis

Latently infected TGs from male and female mice were reactivated for 5,10 or 15 hours with

LY294002 in the presence and absence of acyclovir. (A) Viral genome copy number was

quantified by PCR. Viral gene expression also was quantified by RT-qPCR for (B) immediate

early (*ICP27*), (C) early (*ICP8*), and (D) late (*gC*) genes. The average genome copy number

(E) and ICP27 mRNA transcripts (F) for male and female mice were calculated. Transcript

copy number was normalized to cellular control (18s rRNA). Limit of detection indicated by

black dashed line. N≥19 biological replicates (A-D). Mann-Whitney U (**A-F**). *p<0.05,

**p<0.01, ***p<0.001. ****p<0.0001. The means and SEMs are represented.

<u>Figure 4: PI3K inhibition of explanted superior cervical ganglia induces rapid lytic gene</u>

<u>expression in the absence of detectible genome synthesis.</u>

Latently infected SCGs from female mice were reactivated for 5 or 20 hours with LY294002.

(A) Viral genome copy number was quantified by PCR. Viral gene expression was quantified

by RT-qPCR for (B) immediate early (*ICP27*), and (C) late (*gC*) genes. Transcript copy

number was normalized to cellular control (18s rRNA). Limit of detection indicated by black

dashed line. N≥12 biological replicates (A-C). Mann-Whitney U (**A-C**). *p<0.05, **p<0.01,

***p<0.001. ****p<0.0001. The means and SEMs are represented.

<u>Figure 5: Robust detection of pre-formed virus occurs 20 hours post excision</u>

Latently infected TGs from female mice were reactivated for 0, 7, 10, 15, or 20 hours with

LY294002 and titers of pre-formed virus quantified on Vero cells. Number of ganglia with

detectible virus is displayed beneath the X axis. N≥7 biological replicates. The median titer is

represented.

<u>Figure 6: DLK activity is required for the induction of lytic gene expression following</u>

<u>explant/PI3-kinase inhibition of latently infected TG.</u>

Latently infected TG were explanted and incubated with LY294002 and the DLK inhibitor

GNE-3511 (8 μM) for 5 hours post excision. Viral gene expression was quantified by RT-

qPCR for immediate early (*ICP27*) (A), early (*ICP8*) (B), and late (*gC*) (C) genes. Limit of

detection indicated by dashed line. n=12 biological replicates. Mann-Whitney U test. *p<0.05,

**p<0.01. Individual biological replicates along with the means and SEMs are represented.

<u>Figure 7: Lytic gene induction following axotomy/PI3-kinase inhibition is unaffected by histone</u>

<u>demethylase inhibitors.</u>

Latently infected TG were explanted and incubated with LY294002 along with the LSD1

inhibitor OG-L1002 (50 μM) or GSK-J4 (20 μM). Viral gene expression was quantified by RT-

qPCR for immediate early (*ICP27*) (A), early (*ICP8*) (B), and late (*gC*) (C) genes. Mann-

Whitney U Test, **p<0.01, ***p<0.001. Individual biological replicates along with the means

and SEMs are represented.

658
659

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7