Molecular and mineral responses of corals grown under artificial Calcite Sea conditions

Nicola Conci [1], Erika Griesshaber [1,4], Ramón E. Rivera-Vicéns [1], Wolfgang W. Schmahl [1,2,4], Sergio Vargas [1], Gert Wörheide [1,3,4] *

[1] Department of Earth and Environmental Sciences, Paleontology & Geobiology, Ludwig-Maximilians-Universität, Munich, Germany.
[2] SNSB - Mineralogische Staatssammlung, Munich, Germany
[3] SNSB - Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany
[4] GeoBio-Center LMU, Ludwig-Maximilians-Universität, Munich, Germany

* corresponding author, email: woerheide@lmu.de

Abstract

The formation of skeletal structures composed of different calcium carbonate polymorphs (aragonite and calcite) appears to be regulated both biologically and environmentally. Among environmental factors influencing aragonite and calcite precipitation, changes in seawater conditions – primarily in the molar ratio of magnesium and calcium during so-called "Calcite" (mMg:mCa below 2) or "Aragonite" seas (mMg:mCa above 2) – have had profound impacts on the distribution and performance of marine calcifiers throughout the Earth's history. Nonetheless, the fossil record shows that some species appear to have counteracted such changes and kept their skeleton polymorph unaltered. Here, the aragonitic octocoral *Heliopora coerulea* and the aragonitic scleractinian *Montipora digitata* were exposed to Calcite Sea-like mMg:mCa with various levels of changes in magnesium and calcium concentration, and both mineralogical (i.e., $CaCO_3$ polymorph) and gene expression changes were monitored. Both species maintained aragonite deposition at lower mMg:mCa ratios, while concurrent calcite presence was only detected in *M. digitata*. Despite a strong variability between independent experimental replicates for both species, the expression for a set of putative calcification-related genes, including known components of scleractinian skeleton organic matrix, was found to consistently change at lower mMg:mCa. These results support previously proposed involvements of the skeleton organic matrix in counteracting decreases in seawater mMg:mCa. Although no consistent changes in expression for calcium and magnesium transporters were observed, down-regulation calcium channels in *H. coerulea* in one experimental replicate and at an mMg:mCa of 2.5 might indicate the possibility of active calcium uptake regulation by the corals under altered mMg:mCa.

Introduction

Biomineralization refers to the process by which organisms secrete and form their skeletons (Lowenstam and Weiner, 1989). Anthozoan (Phylum Cnidaria) biomineralization holds a particularly important ecological role, as it represents the primary mechanism sustaining the growth of coral reefs. In corals, skeletons are composed of different calcium carbonate ($CaCO_3$) polymorphs. Members of the order Scleractinia produce aragonite exoskeletons, except *Paraconotrochus antarcticus*, a recently discovered species that exhibits a two-component (i.e., aragonite and calcite) skeleton (Stolarski et al., 2021). In the subclass Octocorallia, an opposite pattern occurs: the vast majority of species produce calcitic endoskeletal structures, while aragonite exoskeletons are deposited by members of the order Helioporacea (i.e., the blue corals).

As observed in other organisms, including mollusks, arthropods, and chordates, biomineralization in corals is a *biologically controlled* process (Lowenstam and Weiner, 1989). During biomineralization, corals control the availability and concentration of ions required for calcification (e.g., Ca^{2+} and HCO_3^-) in the extracellular calcifying medium (ECM) (Moya et al., 2008; Bertucci et al., 2011; Zoccola et al., 2015; Le Goff et al., 2016), and the production of various macromolecules putatively involved in calcium carbonate deposition, collectively referred to as the skeleton organic matrix (SkOM). SkOM components are secreted into the calcification space and are eventually occluded within the skeleton mineral. Several regulatory functions have been attributed to the SkOM of corals and other marine calcifiers. These include crystal nucleation, formation, and inhibition (Wheeler et al., 1981; Peled-Kamar et al., 2002; Puverel et al., 2005; Von Euw et al., 2017), and controlling the calcium carbonate ($CaCO_3$) polymorph of the skeleton (Thompson et al., 2000; Goffredo et al., 2011).

In corals, the effect of different organic compounds on $CaCO_3$ polymorphs (aragonite, calcite) has been investigated mostly through *in vitro* precipitation experiments. The addition of total SkOM extracts (Hohn and Reymond, 2019), or proteins (Rahman and Oomori, 2009; Goffredo et al., 2011) and lipids (Reggi et al., 2016) isolated from coral skeletons to supersaturated $CaCO_3$ solutions, has been shown to promote or inhibit the formation of specific $CaCO_3$ polymorphs and to affect the shape of $CaCO_3$ crystals. Recently, Laipnik et al. (2019) observed that protein-driven *in vitro* precipitation of different $CaCO_3$ polymorphs is also related to the magnesium concentration ($[Mg^{2+}]$) in the solution used. In those experiments, the absence of aragonite formation – the naturally occurring polymorph in

scleractinian skeletons – at low magnesium concentrations led the authors to propose a key role for seawater [Mg^{2+}] in the functioning of coral skeletogenic proteins.

Seawater chemistry has in fact been long included among the main drivers of selective inorganic precipitation of different $CaCO_3$ polymorphs, with the molar ratio of magnesium and calcium (mMg:mCa) likely representing one of the key factors leading to differential $CaCO_3$ precipitation (Morse and Mackenzie, 1990; Morse et al., 1997; Balthasar and Cusack, 2015). From an evolutionary perspective, the mMg:mCa is particularly interesting due to its estimated fluctuations in the last 600 million years, which caused alternating periods of calcite- and aragonite-favoring oceanic environments (Sandberg, 1983) that robustly correlate with the preferred $CaCO_3$ skeleton polymorph of the dominant reef builders during those geological periods (Stanley and Hardie, 1998, 1999). During the Cretaceous, for example, when the mMg:mCa dropped to ~ 1 (compared to the modern value of 5.2), scleractinian corals were replaced by calcitic bivalves (rudists, Phylum Mollusca) as main reef builders. Remarkably, some calcitic scleractinians have been reported from this period (Stolarski et al., 2007). However, fossils of cretaceous aragonitic corals indicate that some coral species could deposit aragonite despite the lower, calcite-favoring mMg:mCa marine environment (Janiszewska et al., 2017). Aragonitic blue coral fossils are also known from the Cretaceous (Eguchi, 1948; Colgan, 1984), further reinforcing the idea that the effect of different mMg:mCa ratios on the skeleton polymorph of corals differ between species and, likely, reflects the ability of some corals to deposit specific $CaCO_3$ polymorphs under chemically unfavorable conditions.

In vivo experiments exposing animals to different mMg:mCa ratios provide support for the ability of corals to counteract the putative decisive influence of seawater chemistry on polymorph formation. In an early work, Ries (2006) observed an increase in the amount of calcite present in the skeleton of three scleractinian corals, with the amount of calcite found correlating inversely to the seawater mMg:mCa ratio. These observations were later corroborated by Higuchi et al (2014), who reported significant amounts of calcite in the skeleton of juvenile *Acropora tenuis*, albeit at lower mMg:mCa ratios compared to Ries (2006). More recently, Yuyama et al. (2019) reported changes in gene expression in the scleractinian coral *A. tenuis* grown in Mg-depleted seawater. These included the up-regulation of several putative skeletogenic genes, suggesting that lower mMg:mCa elicits a response in the corals' calcification machinery. However, since Yuyama et al (2019) only manipulated the Mg-content of the water leaving the calcium levels constant, the molecular

response of corals to mMg:mCa ratios comparable to those observed during geological periods corresponding to Calcite Seas, when both the concentration of calcium and magnesium changed, remain unknown.

Here, we exposed the branching stony coral *Montipora digitata* (Scleractinia) and the blue coral *Heliopora coerulea* (Octocorallia) to both calcite and aragonite-inducing seawater and used gene expression analysis (RNA-seq), electron backscatter diffractometry (EBSD), and energy dispersive spectroscopy (EDS) to investigate the molecular and mineralogical response of these corals to different mMg:mCa ratios resembling past oceanic conditions.

Materials and Methods

Fragments of *H. coerulea* and *M. digitata* (ca. 1cm^3) were mechanically obtained (and attached to the substrate) from colonies cultured for more than a decade in the aquarium facilities of the Chair for Geobiology & Palaeontology of the Department of Earth and Environmental Sciences at Ludwig-Maximilians-Universität München (Munich, Germany). Following fragmentation, corals were allowed to recover for about one month prior to the start of each experiment. Two days before the start of each experiment, coral samples were incubated for ca. 36 hours in a 10% alizarin red solution to mark the skeleton deposited before the experiment.

Experimental Design

Two independent experiments were performed for each target species. Each experiment lasted ca. six weeks (22.07.-10.09.2018 and 01.11.-12.12.2018 for *M. digitata* and 04.01.-28.02.2019 and 01.03.-15.04.2019 for *H. coerulea)*. The experimental setup is illustrated in Figure 1. It consisted of three 8 L aquarium tanks: one (control tank) containing seawater with an mMg:mCa of 5.2 (10mM Ca^{2+} - 52 mM Mg^{2+}) and two treatment tanks with ratios 2.5 (17mM Ca^{2+}- 47 mM Mg^{2+}) and 1.5 (25mM Ca^{2+}- 37 mM Mg^{2+}). In all three tanks, the sum of [Ca^{2+}] and [Mg^{2+}] was equal and set at 62 mmol. A single source of Mg^{2+} and Ca^{2+}-free seawater was used for all three tanks and the mMg:mCa ratio was adjusted by adding different amounts CaCl$_2$xH$_2$O and MgCl$_2$x6H$_2$O to each tank. Water was replaced in each tank every two days. The composition of control and treatment artificial seawater used is provided in S.Mat1 (available online at https://github.com/PalMuc/CalciteSea).

Figure 1 Schematic diagram of the experimental design. For each species the experiment was repeated twice: 22.07.-10.09.2018 and 01.11.-12.12.2018 for *M. digitata*, and 04.01.-28.02.2019 and 01.03.-15.04.2019 for *H. coerulea.*

To allow for acclimatization, corals in treatment tanks were initially exposed for two days to an mMg:mCa ratio of 5.2, before lowering the mMg:mCa ratio to 4.0 and 3.0 for two days, respectively. After the acclimation phase, the mMg:mCa ratio was adjusted to the final experimental ratios. At the end of each experiment, one coral fragment was bleached for 24 hrs in 5% NaOCl, washed several times in ultrapure water, air dried, and stored for mineralogical analysis. Three corals from each tank were flash-frozen in liquid nitrogen and stored at -80° C until RNA extraction.

Mineralogical Analysis

To prepare coral samples for EBSD measurements, skeletons were first thoroughly washed with deionized water and then air dried. Samples were embedded in polypropylene moulds using ca. 30 ml of Epofix Resin (Struers) and 4 ml of Resin Hardener. Samples were placed in a vacuum desiccator to degas and polymerize for 24 hours. Embedded skeletons surfaces were ground using P320, P600, and P1200 silicon carbide paper (Buehler), and flushed with deionized water to remove resin and silicon residues. The first round of polishing was done with a 3 μm polycrystalline diamond suspension for ca. 10-15 minutes. The samples were then attached to cylinder weights with Thermoplastic Cement at 100-150°C. After reaching

room temperature, samples were finally polished (for ca. three hours) with MicroFloc in a Vibromet using a 50 nm Alumina suspension and washed. Prior to EBSD analysis, the polished samples were coated with 4-8 nm of carbon and mounted on a sample holder with a 70° tilt with respect to the beam. For each sample, an 'inner' and 'outer' (growing edge) area were analyzed to obtain mineralogical information about the skeleton deposited before and after the start of the experiment. EBSD measurements were performed with a Hitachi SU5000 field emission SEM operated at 20kV. The microscope is equipped with an electron backscatter diffraction (EBSD) detector (Oxford Instruments). Measurements were conducted using the AZtech Suite (Oxford Instruments), while phase and orientation maps were produced with the CHANNEL 5 HKL software (Schmidt and Olesen, 1989; Randle and Engler, 2000). Electron backscatter diffraction measurements give crystal orientation and phase composition (Nowell and Wright, 2005). Crystal orientation is shown color-coded in EBSD. The used color code is shown with each EBSD map.

Grain parameters for all samples were determined with Tango 5.12. For each mapped skeleton portion, the relative abundance of grain area and major axis was computed. In an effort to account for differences possibly caused by sectioning plane and natural spatial heterogeneity of grain size, an additional variable "*minor axis*" was calculated by dividing each grain area by its major axis. For data plotting all three measures were log-transformed. Mineralogical datasets (both raw files and datasets with additional variables (minor axis)) - alongside the R script used to generate the grain statistics plots are available at https://github.com/PalMuc/CalciteSea. All SEM raw files used to generate phase images, band contrast images, and orientation maps have been deposited in the Open Data LMU repository (https://data.ub.uni-muenchen.de; DOI pending).

Transcriptome Sequencing

Whole coral samples were homogenized in 1-2 ml of Trizol (Thermofisher) using a Polytron PT Homogenizer (Kinematica) and centrifuged at 15,000 g for 10 minutes to remove residual skeleton powder. RNA was extracted according to a modified TriZol protocol (Chomczynski and Mackey, 1995). Modifications included the substitution of 50% of the isopropanol with a high-salt solution (1.2 M sodium chloride and 0.8 M sodium citrate) in order to avoid the coprecipitation of polysaccharides and other potential contaminants. RNA purity and integrity were assessed on a NanoDrop 2100 spectrophotometer and a Bioanalyzer 2100 (Agilent), respectively. For sequencing, only samples with a RIN value > 8.0 were considered (N=17 *H. coeruela* and N=16 *M. digitata*). Strand-specific libraries were prepared with the SENSE

mRNA-Seq Library Prep Kit V2 for Illumina (Lexogen) and paired-end sequenced (50 bp) on an Illumina HiSeq1500 at the Gene Center of Ludwig-Maximilians-Universität München. Sequences were quality controlled with FastQC (www.bioinformatics.babraham.ac.uk) and low-quality reads were removed with the Filter Illumina program from the Agalma-Biolite transcriptome package (Q value cutoff: 28) (Dunn et al., 2013). Reads generated as part of these experiments have been deposited at the European Nucleotide Archive (www.ebi.ac.uk/ena) under Bioproject Number PRJEB36989 (*M. digitata*) and PRJEB36990 (*H. coerulea*). For the assembly of *H. coerulea,* previously sequenced reads (PRJEB30452) were also used.

Transcriptomes were assembled with TransPi (Rivera-Vicéns et al., 2022). Contigs of length < 300 bp were discarded. Host and symbiont contigs were separated with psytrans (https://github.com/sylvainforet/psytrans) using the *Dendronephthya gigantea* (Jeon et al. 2019) and *Acropora digitifera* (Shinzato et al., 2011) sequences for assigning host contigs of *H. coerulea* and *M. digitifera* sequences, respectively. Transcriptome completeness was assessed with BUSCO 3.0.2 (metazoa database) (Simão et al., 2015). Summary statistics of the transcriptomes are provided in Table 1.

Species	Contigs	N50 - Mean length	BUSCO [C-F-M]
H. coerulea	110,535	819 - 716	[97.0 - 1.4 - 1.6]
M. digitata	71,676	819 -705	[90.4 - 4.3 - 5.3]

Table.1 Summary statistics for the *H. coerulea* and *M. digitata* transcriptomes assembled with Transpi. Parameters refer to host transcriptome CDS only after contigs assignment with psytrans. For BUSCO scores, percentages of complete (C), fragmented (F), and missing (M) are provided.

Gene Expression Analysis
For gene expression analysis, read files of each species were mapped against their metatranscriptome (host and symbiont contigs) using Salmon 0.11.2 (Patro et al., 2017) via our local Galaxy server (galaxy.palmuc.org). Read quantification files for each sample were combined into a count matrix with the abundance_estimates_to_matrix.pl script provided with Trinity 2.8.6 (Grabherr et al., 2011), and count data for host (coral) sequences was filtered based on psytrans results. Differential expression analysis was performed using DESeq2 v.1.24.0 (Love et al., 2014). To account for the observed batch effect between experimental replicates (Figures S5 and S6), the experimental replicate was included as a factor in the model for DESeq2 analysis using the design formula "*~experiment + condition*".

For each comparison (i.e., mMg:Ca 5.2 vs 2.5 and 5.2 vs 1.5), differentially expressed genes with adjusted p-value < 0.01 and log-fold changes - 2 > and > 2 were considered. The complete DESeq2 R-script workflow used, along with the count matrices, expression tables, sequences of differentially expressed genes, and the transcriptomic assemblies of *H. coerulea* and *M. digitata* are available as supplementary materials at https://github.com/PalMuc/CalciteSea.

Results

Mineralogical Analysis

EBSD measurements of *Heliopora coerulea* and *Montipora digitata* skeletons showed a possible species-specific response to variations in seawater mMg:mCa (Figures 2–4). Under control conditions (mMg:mCa = 5.2) and mMg:mCa of 2.5, calcite was not detected in both *H. coerulea* or *M. digitata* (Figure 3–4). This pattern was consistent between the two experimental replicates. Contrary, the presence of calcite was detected in one *M. digitata* coral sample maintained under calcite-inducing conditions (mMg:mCa of 1.5) (Figure 2), whereas only aragonite was observed in the *H. coerulea* skeleton areas surveyed under these conditions.

Figure 2a: Light microscopy (left) and SEM (right) image of *Montipora digitata* skeleton with calcite. Red and blue squares indicate two areas deposited during the experiment and analyzed with EBSD. Data for the area deposited prior to the experiment is available in Figure S1 **b)** Phase map, band contrast, and orientation map of the area where calcite was observed (1-3) (blue rectangle) and the area composed only of aragonite (4-6) (red rectangle).

Figure 3 Calcium carbonate phase map, band contrast images, and EBSD orientation map of *Montipora digitata* skeletons grown under each experimental condition (*m*Mg:*m*Ca 5.2, 2.5, and 1.5) and during experimental replicate. Information for the *M. digitata* sample in which calcite was observed is available in Figure 2. In the phase map figures, red indicates aragonite and blue indicates calcite.

Figure 4 Calcium carbonate phase map, band contrast images, and EBSD orientation map of *Heliopora coerulea* skeletons grown under each experimental condition (*m*Mg:*m*Ca 5.2, 2.5 and 1.5) and experimental replicate. In the phase map figures, red indicates aragonite and blue indicates calcite.

Microscopy images (Figure 3a, 3b) show that the calcite did not form homogeneously within the newly deposited skeleton, but it was rather found to be restricted within mineralized circular structures. These appear in direct contact with the coral skeleton. Phase analysis (Figure 4 b1) revealed a mixed composition within these areas with the co-presence of both aragonite and calcite. Neither these circular structures nor calcite could be detected in the portion of the skeleton grown prior to the experiment (Figure S1). Analysis of grain area and diameters showed differences between species, experimental conditions, and replicates (Figures S3 and S4). In *M. digitata*, no difference in terms of grain area or both major and minor grain axes in corals exposed to an mMg:mCa of 2.5, while in one coral grown at an mMg:mCa of 1.5 a decrease in grain size - compared to control conditions - could be observed (Figure S3 and S4). On the contrary, in one experiment replicate both corals grown under treatment conditions (2.5 and 1.5) exhibited larger grain sizes (Figures S3 and S4). For both species, differences between control and treatment ratios were larger when grain areas were compared, while a higher overlap could be observed when grain minor axes were compared.

Gene Expression Analysis

Exposure to mMg:mCa ratios of 2.5 and 1.5 triggered changes in gene expression (p-value < 0.001; log fold change > |2|) in both species. The octocoral *H. coeruela* showed overall higher numbers of differentially expressed genes (DEGs) compared to *M. digitata* under both treatments. However, principal component analysis (PCA), based on log-transformed expression values relative to control, revealed a high dispersion of control and treatment samples for both species, and strong differences between the two experimental replicates for both species (Figures S5 and S6). In *M. digitata*, expression changes of known putative calcification-related genes were restricted to mMg:mCa of 1.5 (Figure 5). These included the over-expression of one galaxin-like 2 (AD50285.1) homolog, one matrix metalloproteinase-like protein, and one uncharacterized skeletal matrix protein (B3EX00.1) previously identified in the skeleton of scleractinian corals (Ramos-Silva et al., 2013; Conci et al., 2020). In *H. coerulea*, the only putative calcification DEG found coded for a cartilage-intermediate layer like-protein and was consistently down-regulated in blue corals exposed to mMg:mCa of 2.5.

a)

Color Key and Histogram

Row Z-Score

b)

Sequence ID	Annotation	Species	*m*Mg:*m*Ca	LFC / Padj
TransAbySS.k25.R164 4600.p1	Unch. skeletal matrix protein 1-like	*M. digitata*	1.5	2.57 / < 0.01
Trinity_DN1436_c0_g 1_i9.p1	Galaxin-like 2	*M. digitata*	1.5	2.00 / < 0.01
Trinity_DN121764_c0 _g1_i3.p1	Matrix Metalloproteinase 25-like	*M. digitata*	1.5	2.11 / < 0.01
SOAP.k25.C2929031. p1	Cartilage-intermediate-layer protein-like	*H.coerulea*	2.5	-10 / < 0.01

Fig 5: a) Gene expression heatmap of potential calcification-related genes in *M. digitata* under different *m*Mg:*m*Ca ratios. **b)** Table providing accession IDs, annotation, and log fold-change (LFC) and adjusted p-value (Padj). The table also includes the only potential calcification-related gene we observe changing across experiments in *H. coerulea.*

Analysis of single experimental replicates showed up-regulation for additional potential calcification-related genes. In *M. digitata*, these included, coadhesin-like and protocadherin-like reported in the *A. millepora* and *S. pistillata* skeletons (Ramos-Silva et al., 2013; Drake et al., 2013). Moreover, in one experimental replicate, galaxin was also up-regulated at *m*Mg:*m*Ca of 2.5. Finally, an uncharacterized skeletal organic matrix protein 5-like protein was found down-regulated at *m*Mg:*m*Ca of 1.5.

Although changes in the expression of calcium or magnesium transporters were not detected in both experimental replicates, different voltage-dependent calcium channel alpha subunits were down-regulated in *H. coerulea* under *m*Mg:*m*Ca of 2.5 in Experiment 1.

Discussion

The formation of coral skeletons is intimately linked to both cellular/molecular processes, and the surrounding environmental conditions. Among the latter, the magnesium/calcium molar ratio of seawater (mMg:mCa) appears to have influenced the polymorph of calcium carbonate ($CaCO_3$) of major marine calcifiers, and the polymorph initially adopted by marine organisms during the last 500 myr (Zhuravlev and Wood, 2008). However, organisms also appear to be less affected by subsequent changes in seawater chemistry promoting the precipitation of a different polymorph (Porter, 2010). Manipulative experiments conducted in artificially recreated "Calcite Seas" have, however, highlighted the inability of modern scleractinian corals to fully counteract the effects of mMg:mCa changes, leading to significant amounts of calcite being deposited within their skeletons (Ries et al., 2006; Higuchi et al., 2014). The present study combined mineralogical (electron backscattered diffraction, EBSD) and molecular (gene expression) analyses to provide a first insight into the effect of different Mg:Ca molar ratios on aragonitic octocoral skeletons, namely those of *Heliopora coerulea*, and compared the results with the scleractinian coral *Montipora digitata*. To examine and compare their responses at the polymorph and gene expression level, both species were exposed to three different mMg:mCa ratios: the modern value of 5.2 (control group), 2.5 and 1.5.

As previously shown for scleractinian species (Ries et al., 2006; Higuchi et al., 2014), our experiments highlighted the ability of *M. digitata* and *H. coerulea* to deposit aragonite when exposed to calcite-inducing mMg:mCa ratios down to 1.5. Albeit not being detected in both samples, the presence of calcite in the *M. digitata* samples also supports the observations from the two aforementioned studies. Although we could not detect calcite in corals gown at mMg:mCa ratios higher than 2, as previously described in *M. digitata* (Ries et al., 2006)

In this regard, differences in the experimental setup, analytical methods, and both coral target species, as well as growth stages have to be considered when comparing results across different studies. Although the characterization of the $CaCO_3$ polymorph with EBSD allowed us to obtain information about crystal size and orientation, it was here restricted to delimited areas of a limited number of coral colonies. This might 1) represent one of the reasons for the variability observed between experimental replicates for several of the parameters investigated, and 2) have possibly caused to have missed and/or underestimated calcite content in the skeletons. Future research should thus include additional methodologies (e.g., X-ray diffraction or Raman spectroscopy) to more

comprehensively assess the presence of polymorph changes and allow the investigation of higher sample numbers.

The origin and formation mechanisms of the calcite observed at low mMg:mCa ratios in *M.digitata* remain elusive. The distribution analysis of the detected calcite grains points to the calcite-inducing effects of the mMg:mCa ratio not being limited to polymorph substitutions (same structure but different composition). On the contrary, calcite was only observed in areas that also exhibit distinct macro-morphological differences. Compared to the rest of the skeleton – which exhibits branching-like structures – calcite in *M. digitata* appears as spherical bodies surrounded by (and potentially in direct contact with) the aragonitic skeleton.

In this regard, although the observed morphological variation could be related to the effects of mMg:mCa on the biological calcification machinery, we also consider the possibility that the observed structures originate in enclosed areas of *M. digitata*'s skeleton when corals are exposed to calcite-inducing conditions. In these chambers, the composition of calcification fluids might experience strong shifts and promote the precipitation of calcite alongside aragonite. Finally, it cannot be excluded that the presence of calcite could also be possibly related to other mechanisms, such as the random presence of foreign mineral material, that has been previously observed in scleractinian skeletons (Goffredo et al., 2012).

Variability was observed between experiments aiming to investigate the molecular mechanisms behind the effects of mMg:mCa on coral calcification. Consistent across experiments, expression data from *H. coerulea* highlighted expression changes for only one putative calcification-related gene: one cartilage-intermediate layer protein homolog underexpressed under mMg:mCa of 2.5. No expression changes for components of the known skeleton proteome of this species (Conci et al., 2020) were detected. On the other hand, in *M. digitata*, a set of calcification-related genes, including a galaxin-related protein, was found upregulated in both experimental replicates, when the mMg:mCa was lowered to 1.5. The response of SkOM genes to calcite-inducing mMg:mCa described here might nonetheless have been underestimated, as the high variability observed between and within experiments might have prevented the detection of changes in the expression of additional genes. Overexpression of multiple SkOM proteins (including galaxin-related proteins) under calcite-inducing mMg:mCa has nonetheless been previously reported in *A. tenuis* (grown under mMg:mCa of 0.5), and was linked to active initiation and maintenance of aragonite deposition under adverse environmental conditions (Yuyama and Higuchi, 2019). Generally,

the overexpression of calcification-related genes in corals exposed to calcite-inducing environments may indicate an increased energetic expenditure by the corals and explain the decline in calcification rates observed in different coral species at lower mMg:mCa ratios (Higuchi et al., 2014; Ries et al., 2006). The reduction in grain sizes observed here in *M. digitata* may also represent a consequence of a higher energetic cost of calcification under low mMg:mCa ratios. However, as for the gene expression analyses, the level of variation observed across experiments and possible effects of biological variability and sample preparation on the measured grain parameters prompt the need for additional research.

In addition to SkOM components, variation in mMg:mCa ratios could also potentially affect ion transport and/or chemistry of the calcification fluids. Although inconsistent between experiments, observed changes in the expression of calcium transporters in *H. coerulea* could be related to a response of the ion transport machinery used by corals for calcification. An involvement of calcium transporters has been excluded for *A. tenuis,* as no change in expression was detected (Yuyama and Higuchi, 2019). However, in that study the seawater mMg:mCa was manipulated by decreasing $[Mg^{2+}]$, while keeping $[Ca^{2+}]$ at normal levels (ca. 10mM). Thus, information on the putative response of calcium and other ion transporters in corals concurrently exposed to low $[Mg^{2+}]$ and increased $[Ca^{2+}]$ is currently lacking. In corals, calcium is supposedly transported to the calcification sites either paracellularly (Gagnon et al., 2012) or via active transcellular transport (Tambutté et al., 1996; Hohn and Merico, 2015). Different transporters have been associated with transcellular calcium transport including calcium channels, Ca^{2+}-ATPases, and proton exchangers (Zoccola et al., 1999, 2004; Barott et al., 2015; Barron et al., 2018). The regulation of calcium transport allows corals to maintain higher calcium concentrations in calcifying fluids ($[Ca^{2+}]_{cf}$) with respect to seawater (Sevilgen et al., 2019). Up-regulation of $[Ca^{2+}]_{cf}$ can also be enhanced in some coral species to counteract decreases in the saturation state of the calcifying medium caused by decreases in seawater pH (DeCarlo et al., 2018). Different studies have in fact examined the effects of ocean acidification on carbonate and calcium chemistry at calcification sites (Allison et al., 2018; Mollica et al., 2018). However, to the best of our knowledge, research examining the response of ion transport to the calcification site when the environmental calcium concentration is higher than $[Ca^{2+}]_{cf}$ is currently lacking. Further characterizations of the calcification-related calcium transport machinery in octocorals and scleractinians could provide additional insights into the localization and role of single calcium channels, and possibly corroborate the hypothesis that corals might control aragonite

precipitation, when exposed to high environmental $[Ca^{2+}]$, through the regulation of calcium transport.

Here, we examined possible changes in skeleton polymorph and in the expression of calcification-related genes in the aragonitic octocoral *H. coerulea* and the scleractinian *M. digitata* exposed to calcite-inducing *m*Mg:*m*Ca. This work provides the first comparative study of octocoral and scleractinian responses to changes in *m*Mg:*m*Ca. The two species were both found to continue producing aragonite, while the presence of calcite could only be detected in the skeleton of *M. digitata.* Despite variability both within and between experiments, the observed changes in gene expression support previously hypothesized molecular responses of SkOM genes (Yuyama and Higuchi, 2019) when the *m*Mg:*m*Ca is lowered.

Acknowledgments:

The study was supported by the German Research Foundation (DFG) grant "MINORCA" to S.V. (Project Number Va1146-2/1) and G.W. (Project Number Wo896/18-1). G.W. acknowledges support by LMU Munich's Institutional Strategy LMUexcellent within the framework of the German Excellence Initiative. We thank Peter Naumann for technical assistance and maintenance of the aquaria facilities and coral culturing, as well as Simone Schätzle and Gabriele Büttner for assistance in the laboratory and Moritz Zenkert for EBSD samples preparation.

References

Allison, N., Cole, C., Hintz, C., Hintz, K., Rae, J., and Finch, A.: The effect of ocean acidification on tropical coral calcification: Insights from calcification fluid DIC chemistry, Chem. Geol., 497, 162–169, 2018.

Balthasar, U. and Cusack, M.: Aragonite-calcite seas—Quantifying the gray area, Geology, 43, 99–102, 2015.

Barott, K. L., Perez, S. O., Linsmayer, L. B., and Tresguerres, M.: Differential localization of ion transporters suggests distinct cellular mechanisms for calcification and photosynthesis between two coral species, Am. J. Physiol. Regul. Integr. Comp. Physiol., 309, R235–46, 2015.

Barron, M. E., Thies, A. B., Espinoza, J. A., Barott, K. L., Hamdoun, A., and Tresguerres, M.: A vesicular Na+/Ca2+ exchanger in coral calcifying cells, PLoS One, 13, e0205367, 2018.

Bertucci, A., Tambutté, S., Supuran, C. T., Allemand, D., and Zoccola, D.: A new coral carbonic anhydrase in *Stylophora pistillata*, Mar. Biotechnol., 13, 992–1002, 2011.

Chomczynski, P. and Mackey, K.: Short technical reports. Modification of the TRI reagent procedure for isolation of RNA from polysaccharide- and proteoglycan-rich sources, Biotechniques, 19, 942–945, 1995.

Colgan, M. W.: The Cretaceous Coral *Heliopora* (Octocorallia, Coenothecalia)—a Common Indo-Pacific Reef Builder, in: Living Fossils, edited by: Eldredge, N. and Stanley, S. M., Springer New York, New York, NY, 266–271, 1984.

Conci, N., Lehmann, M., Vargas, S., and Wörheide, G.: Comparative proteomics of octocoral and scleractinian skeletomes and the evolution of coral calcification, Genome Biol. Evol., 12, 1623–1635, 2020.

DeCarlo, T. M., Comeau, S., Cornwall, C. E., and McCulloch, M. T.: Coral resistance to ocean acidification linked to increased calcium at the site of calcification, Proc. Biol. Sci., 285, 2018.

Drake, J. L., Mass, T., Haramaty, L., Zelzion, E., Bhattacharya, D., and Falkowski, P. G.: Proteomic analysis of skeletal organic matrix from the stony coral *Stylophora pistillata*, Proc. Natl. Acad. Sci. U. S. A., 110, 3788–3793, 2013.

Dunn, C. W., Howison, M., and Zapata, F.: Agalma: an automated phylogenomics workflow, BMC Bioinformatics, 14, 330, 2013.

Eguchi, M.: Fossil Helioporidae from Japan and the South Sea Islands, J. Paleontol., 22, 362–364, 1948.

Gagnon, A. C., Adkins, J. F., and Erez, J.: Seawater transport during coral biomineralization, Earth Planet. Sci. Lett., 329-330, 150–161, 2012.

Goffredo, S., Vergni, P., Reggi, M., Caroselli, E., Sparla, F., Levy, O., Dubinsky, Z., and Falini, G.: The skeletal organic matrix from Mediterranean coral *Balanophyllia europaea* influences calcium carbonate precipitation, PLoS One, 6, e22338, 2011.

Goffredo, S., Caroselli, E., Mezzo, F., Laiolo, L., Vergni, P., Pasquini, L., Levy, O., Zaccanti, F., Tribollet, A., Dubinsky, Z., and Falini, G.: The puzzling presence of calcite in skeletons of modern solitary corals from the Mediterranean Sea, Geochim. Cosmochim. Acta, 85, 187–199, 2012.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., and Regev, A.: Full-length transcriptome assembly from RNA-Seq data without a reference genome, Nat. Biotechnol., 29, 644–652, 2011.

Higuchi, T., Fujimura, H., Yuyama, I., Harii, S., Agostini, S., and Oomori, T.: Biotic control of skeletal growth by scleractinian corals in Aragonite–Calcite Seas, PLoS One, 9, e91021, 2014.

Hohn, S. and Merico, A.: Quantifying the relative importance of transcellular and paracellular ion transports to coral polyp calcification, Front. Earth Sci., 2, 1129, 2015.

Hohn, S. and Reymond, C. E.: Coral calcification, mucus, and the origin of skeletal organic molecules, Coral Reefs, 38, 973–984, 2019.

Janiszewska, K., Mazur, M., Escrig, S., Meibom, A., and Stolarski, J.: Aragonitic scleractinian corals in the Cretaceous calcitic sea, Geology, 45, 319–322, 2017.

Laipnik, R. 'anan, Bissi, V., Sun, C.-Y., Falini, G., Gilbert, P. U. P. A., and Mass, T.: Coral acid rich protein selects vaterite polymorph in vitro, J. Struct. Biol., 107431, 2019.

Le Goff, C., Ganot, P., Zoccola, D., Caminiti-Segonds, N., Allemand, D., and Tambutté, S.: Carbonic Anhydrases in Cnidarians: Novel Perspectives from the octocorallian *Corallium rubrum*, PLoS One, 11, e0160368, 2016.

Love, M. I., Huber, W., and Anders, S.: Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2, Genome Biol., 15, 550, 2014.

Lowenstam, H. A. and Weiner, S.: On biomineralization, 1989.

Mollica, N. R., Guo, W., Cohen, A. L., Huang, K.-F., Foster, G. L., Donald, H. K., and Solow, A. R.: Ocean acidification affects coral growth by reducing skeletal density, Proc. Natl. Acad. Sci. U. S. A., 115, 1754–1759, 2018.

Morse, J. W. and Mackenzie, F. T.: Geochemistry of Sedimentary Carbonates, Elsevier, 706 pp., 1990.

Morse, J. W., Wang, Q., and Tsio, M. Y.: Influences of temperature and Mg:Ca ratio on CaCO3 precipitates from seawater, Geology, 25, 85–87, 1997.

Moya, A., Tambutté, S., Bertucci, A., Tambutté, E., Lotto, S., Vullo, D., Supuran, C. T., Allemand, D., and Zoccola, D.: Carbonic anhydrase in the scleractinian coral *Stylophora pistillata*: characterization, localization, and role in biomineralization, J. Biol. Chem., 283, 25475–25484, 2008.

Nowell, M. M. and Wright, S. I.: Orientation effects on indexing of electron backscatter diffraction patterns, Ultramicroscopy, 103, 41–58, 2005.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C.: Salmon provides fast and bias-aware quantification of transcript expression, Nat. Methods, 14, 417–419, 2017.

Peled-Kamar, M., Hamilton, P., and Wilt, F. H.: Spicule matrix protein LSM34 is essential for biomineralization of the sea urchin spicule, Exp. Cell Res., 272, 56–61, 2002.

Porter, S. M.: Calcite and aragonite seas and the de novo acquisition of carbonate skeletons, Geobiology, 8, 256–277, 2010.

Puverel, S., Tambutté, E., Pereira-Mouriès, L., Zoccola, D., Allemand, D., and Tambutté, S.: Soluble organic matrix of two Scleractinian corals: partial and comparative analysis, Comp. Biochem. Physiol. B Biochem. Mol. Biol., 141, 480–487, 2005.

Rahman, M. A. and Oomori, T.: In Vitro Regulation of CaCO3 Crystal Growth by the Highly Acidic Proteins of Calcitic Sclerites in Soft Coral, *Sinularia polydactyla*, Connect. Tissue Res., 50, 285–293, 2009.

Ramos-Silva, P., Kaandorp, J., Huisman, L., Marie, B., Zanella-Cléon, I., Guichard, N., Miller, D. J., and Marin, F.: The skeletal proteome of the coral *Acropora millepora*: the evolution of calcification by co-option and domain shuffling, Mol. Biol. Evol., 30, 2099–2112, 2013.

Randle, V. and Engler, O.: Introduction to Texture Analysis: Macrotexture, Microtexture and

Orientation mapping, 125–151, 2000.

Reggi, M., Fermani, S., Samorì, C., Gizzi, F., Prada, F., Dubinsky, Z., Goffredo, S., and Falini, G.: Influence of intra-skeletal coral lipids on calcium carbonate precipitation, CrystEngComm, 18, 8829–8833, 2016.

Ries, J. B., Stanley, S. M., and Hardie, L. A.: Scleractinian corals produce calcite, and grow more slowly, in artificial Cretaceous seawater, Geology, 34, 525–528, 2006.

Rivera-Vicéns, R. E., Garcia-Escudero, C. A., Conci, N., Eitel, M., and Wörheide, G.: TransPi-a comprehensive TRanscriptome ANalysiS PIpeline for de novo transcriptome assembly, Mol. Ecol. Resour., https://doi.org/10.1111/1755-0998.13593, 2022.

Sandberg, P. A.: An oscillating trend in Phanerozoic non-skeletal carbonate mineralogy, Nature, 305, 19–22, 1983.

Schmidt, N.-H. and Olesen, N. O.: Computer-aided determination of crystal-lattice orientation from electron channeling patterns in the SEM, Can. Mineral., 27, 15–22, 1989.

Sevilgen, D. S., Venn, A. A., Hu, M. Y., Tambutté, E., de Beer, D., Planas-Bielsa, V., and Tambutté, S.: Full in vivo characterization of carbonate chemistry at the site of calcification in corals, Sci Adv, 5, eaau7447, 2019.

Shinzato, C., Shoguchi, E., Kawashima, T., Hamada, M., Hisata, K., Tanaka, M., Fujie, M., Fujiwara, M., Koyanagi, R., Ikuta, T., Fujiyama, A., Miller, D. J., and Satoh, N.: Using the *Acropora digitifera* genome to understand coral responses to environmental change, Nature, 476, 320–323, 2011.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M.: BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs, Bioinformatics, 31, 3210–3212, 2015.

Stanley, S. M. and Hardie, L. A.: Secular oscillations in the carbonate mineralogy of reef-building and sediment-producing organisms driven by tectonically forced shifts in seawater chemistry, Palaeogeogr. Palaeoclimatol. Palaeoecol., 144, 3–19, 1998.

Stanley, S. M. and Hardie, L. A.: Hypercalcification: paleontology links plate tectonics and geochemistry to sedimentology, GSA Today, 9, 1–7, 1999.

Stolarski, J., Meibom, A., Przenioslo, R., and Mazur, M.: A Cretaceous scleractinian coral with a calcitic skeleton, Science, 318, 92–94, 2007.

Stolarski, J., Coronado, I., and Murphy, J. G.: A modern scleractinian coral with a two-component calcite–aragonite skeleton, Proceedings of the National Academy of Sciences of the USA, 2021.

Tambutté, É., Allemand, D., Mueller, E., and Jaubert, J.: A compartmental approach to the mechanism of calcification in hermatypic corals, J. Exp. Biol., 199, 1029–1041, 1996.

Thompson, J. B., Paloczi, G. T., Kindt, J. H., Michenfelder, M., Smith, B. L., Stucky, G., Morse, D. E., and Hansma, P. K.: Direct observation of the transition from calcite to aragonite growth as induced by abalone shell proteins, Biophys. J., 79, 3307–3312, 2000.

Von Euw, S., Zhang, Q., Manichev, V., Murali, N., Gross, J., Feldman, L. C., Gustafsson, T., Flach, C., Mendelsohn, R., and Falkowski, P. G.: Biological control of aragonite formation in

stony corals, Science, 356, 933–938, 2017.

Wheeler, A. P., George, J. W., and Evans, C. A.: Control of calcium carbonate nucleation and crystal growth by soluble matrx of oyster shell, Science, 212, 1397–1398, 1981.

Yuyama, I. and Higuchi, T.: Differential gene expression in skeletal organic matrix proteins of scleractinian corals associated with mixed aragonite/calcite skeletons under low mMg/Ca conditions, PeerJ, 7, e7241, 2019.

Zhuravlev, A. Y. and Wood, R. A.: Eve of biomineralization: Controls on skeletal mineralogy, Geology, 36, 923–926, 2008.

Zoccola, D., Tambutté, E., Sénégas-Balas, F., Michiels, J. F., Failla, J. P., Jaubert, J., and Allemand, D.: Cloning of a calcium channel alpha1 subunit from the reef-building coral, *Stylophora pistillata*, Gene, 227, 157–167, 1999.

Zoccola, D., Tambutté, E., Kulhanek, E., Puverel, S., Scimeca, J.-C., Allemand, D., and Tambutté, S.: Molecular cloning and localization of a PMCA P-type calcium ATPase from the coral *Stylophora pistillata*, Biochim. Biophys. Acta, 1663, 117–126, 2004.

Zoccola, D., Ganot, P., Bertucci, A., Caminiti-Segonds, N., Techer, N., Voolstra, C. R., Aranda, M., Tambutté, E., Allemand, D., Casey, J. R., and Tambutté, S.: Bicarbonate transporters in corals point towards a key step in the evolution of cnidarian calcification, Sci. Rep., 5, 9983, 2015.