1

DLK-dependent biphasic reactivation of herpes simplex virus latency established in the

absence of antivirals

Sara Dochnal[1], Husain Merchant[2], Austin Schinlever[1,3], Aleksandra Babnis[1,4], Daniel P.

Depledge[2,5,6], Angus Wilson[2] and Anna R. Cliffe[1, *].

1. Department of Microbiology, Immunology and Cancer Biology, University of Virginia,

Charlottesville, VA, USA.

2. Department of Microbiology, New York University School of Medicine, New York, NY,

USA.

3. Present address: Department of Microbiology, New York University School of Medicine,

New York, NY, USA.

4. Present address: Technical University of Munich, School of Medicine, Institute of Virology,

Munich, Germany.

5. Institute of Virology, Hannover Medical School, Hannover, Germany.

6. German Center for Infection Research (DZIF), partner site Hannover-Braunschweig,

Hannover, Germany.

* Correspondence to Anna R. Cliffe, cliffe@virginia.edu

Abstract

An understanding of the molecular events that regulate entry of herpes simplex virus type 1 (HSV-1) into a latent infection of neurons and subsequent reactivation requires model systems. However, establishing a latent-like infection in cultured neurons is problematic as the release of any infectious virus can potentially superinfect the cultures. Here, we describe a new reporter virus, HSV-1 Stayput-GFP, that is defective for cell-to-cell spread. Use of this virus permits the establishment of a quiescent infection of neurons without the need for a viral DNA replication inhibitor. As such, Stayput-GFP provides a new model to study latency establishment and reactivation at the single neuron level. We demonstrate that reactivation involves an initial wave of viral gene expression consistent with the Phase I previously observed in primary neuron models where a viral DNA replication inhibitor is used during latency establishment. Phase I gene expression was dependent on the cell stress protein DLK but independent of histone demethylase activity and viral DNA replication. Progression into the later, Phase II wave of reactivation, characterized by detectable late viral protein synthesis and a requirement for histone demethylase activity, is also observed in the Stayput-GFP model system. These data demonstrate that the two waves of viral gene expression following HSV-1 reactivation are independent of secondary infection and occur when latent infections are established in the absence of a viral DNA replication inhibitor.

Importance

Herpes simplex virus-1 (HSV-1) enters a latent infection in neurons and periodically

reactivates. Reactivation manifests as a variety of clinical symptoms. Studying latency

and reactivation *in vitro* is invaluable to elucidating molecular mechanisms behind both

processes to be targeted by therapeutics and reduce clinical consequences. Here, we

describe a novel *in vitro* model system using a cell-to-cell spread defective HSV-1,

known as Stayput-GFP, which allows for the study of latency and reactivation at the

single neuron level. We anticipate this new model system will be an incredibly valuable

tool for studying the establishment and reactivation of HSV-1 latent infection *in vitro*.

Using this model, we find that initial reactivation events are dependent on cellular stress

kinase DLK, but independent of histone demethylase activity and viral DNA replication.

Our data therefore demonstrate the essential role of DLK in mediating a wave of lytic

gene expression unique to reactivation.

Introduction

Herpes simplex virus 1 (HSV-1) is a globally prevalent pathogen with the capacity to infect neurons, including both sensory and autonomic neurons (1-4) . Following neuronal infection, HSV-1 can enter a lytic replication cycle, establish a lifelong latent infection, or potentially undergo a limited amount of lytic gene expression even prior to latency establishment (5-13). While latency is largely asymptomatic, periodic reactivation of the virus can result in manifestations including cutaneous lesions, keratitis, and encephalitis. Epidemiological studies have also linked HSV infection with an increased risk of developing late onset Alzheimer's disease (LOAD), particularly in individuals with the ApoE4 variant (14-22).

The regulated expression of viral lytic transcripts has been well characterized following lytic infection (23-25) . The HSV-1 genome enters the host cell epigenetically naked (26-28) but becomes chromatinized by histones bearing transcriptionally permissive histone modifications on the viral genome (29-38). Viral gene expression is initiated from the genome in response to viral trans-activator and tegument protein VP16. VP16 and host cell factor 1 (HCF-1) bind together and are recruited to VP16-responsive enhancer elements in the immediate early (IE) viral promoters through cellular factor POU homeodomain protein Oct-1 (39). Together, this VP16-induced complex recruits additional factors involved in transcriptional activation, including general transcription factors, ATP-dependent chromatin remodelers, and histone modifying enzymes (32, 40-44). Synthesis of the IE proteins is required for early (E) mRNA transcription. Products of the early viral genes enable viral DNA replication. Viral genome synthesis is a pre-

requisite for true-late (TL) mRNA transcription, likely due to a shift in genome

accessibility and increased binding of host transcriptional machinery (45).

During HSV-1 latency, viral transcription is largely limited to a family of non-coding

RNAs known as the latency-associated transcripts (LATs) (46-48) . Viral lytic mRNAs

are largely transcriptionally repressed and promoters assemble into silent

heterochromatin marked by the tri-methylation of histone H3 lysine 27 (H3K27me3) and

di/tri-methylation of histone H3 on lysine 9 (H3K9me2/3) (49-54). The initiation of viral

gene expression during reactivation is induced from a heterochromatin-associated viral

genome and occurs in the absence of viral activators like VP16. Latent HSV-1 therefore

relies on host factors to act on the epigenetically silent viral genome and induce lytic

gene expression.

The mechanisms that regulate entry into lytic gene expression to permit

reactivation remain elusive. Using primary neuronal models of HSV-1 latent infection,

reactivation has been found to progress in a two-step or bi-phasic manner. Phase I is

characterized as a synchronous wave of lytic viral transcripts, occurring approximately

20 hours post-stimulus (25, 55-58). There is evidence that this initial induction of lytic

gene expression is not dependent on the lytic trans-activator VP16, or viral protein

synthesis (25, 59). Instead, cellular factors, including the stress kinases dual leucine

zipper kinase (DLK) and c-Jun N terminal kinase (JNK), are required for Phase I entry

(56, 58). Importantly, viral gene expression occurs despite the persistence of

heterochromatin on viral promoters. Instead, JNK-dependent histone phosphorylation of

histone H3S10 results in a methyl/phospho switch, which can permit gene expression

even while the repressive H3K9me3 histone modification is maintained (56). The

second wave of viral lytic gene expression, Phase II, occurs approximately 48 hours

post-stimulus. Phase II reactivation is characterized by the full transcriptional viral

cascade, including viral DNA replication, and ultimately infectious virus production (25).

In contrast to Phase I, viral protein synthesis, repressive heterochromatin removal,

VP16-mediated transactivation, and viral DNA replication are required for Phase II (25,

56, 58).

In vitro systems of HSV-1 latency are required to study the mechanisms of

latency establishment and reactivation that cannot be easily studied *in vivo* (60-63). *In*

vitro models more readily enable the use of small molecule inhibition and knockdowns

to perform functional studies, and components of the immune system can be included to

understand how the host immune response impacts latency and reactivation (64, 65).

However, there are complications involved in establishing latency *in vitro*. As also

observed in animal models, only a sub-population of infected neurons enter a latent

state whereas other neurons become lytically infected (11, 66-68). This leads to

superinfection of the cultures and an inability to establish a latent infection. Therefore,

many existing *in vitro* models have used viral DNA replication inhibitors, predominantly

acyclovir (ACV), to establish latency *in vitro* (56, 69-74). ACV is proposed to inhibit viral

DNA replication by incorporating into actively replicating viral genomes in lytic cells,

although there is also evidence from ACV-resistant strains that ACV can also inhibit the

viral DNA polymerase (75-77). There are some caveats associated with ACV use. Since

these inhibitors prevent complete genome synthesis, the process of DNA replication

and subsequent late gene expression cannot be studied in lytic neurons during latency

establishment. Moreover, whether ACV is incorporated into viral DNA and the fate of

any potential ACV-incorporated genome is unknown. Therefore, new model systems

are required in which latency establishment can be tracked without the need for viral

DNA replication inhibitors. Such a system can also be used to determine whether the

use of ACV in the cultures alters mechanisms of lytic gene expression during

reactivation.

To create a novel model of latency and reactivation in which viral late protein

synthesis can be tracked and that does not require viral DNA replication inhibitors, we

created Stayput-GFP. In this HSV-1 reporter virus, a late viral fusion protein Us11-GFP

was inserted into a previously characterized cell-to-cell spread deficient glycoprotein H

deletion virus (78) . The benefit of using a cell-to-cell spread defective virus is that viral

DNA replication and late gene expression can still be studied during the establishment

of latency, but superinfection does not occur. This is also advantageous for the study of

reactivation as it ensures that only reactivation in an individual neuron is investigated

without the risk of further *de novo* lytic replication by progeny virus. Using this system,

we found that there is heterogeneity following neuronal infection, with a subpopulation

of neurons that enter Late gene expression. This acute infection ultimately resolves over

30 days and quiescence is established without requiring viral DNA replication inhibitors.

Interestingly, we find that the longer timescale required for latency establishment than

that used in previous *in vitro* model systems results in an inability to induce reactivation

using only a single trigger. However, reactivation can be induced using a combination of

stimuli. Importantly, using this system, we still observed criteria for Phase I gene

expression, including the requirement for the cell stress-response kinase DLK, as well

as viral gene expression that is independent of histone demethylase activity and viral

DNA replication. Thus, Stayput-GFP provides a powerful new methodology to

investigate the establishment and reactivation from latent infection at the single neuron

level without the need for DNA replication inhibitors. This study also shows that Phase I

gene expression occurs even if a latent infection is established in the absence of

antivirals.

Results

<u>Construction of a gH-null US11-GFP HSV-1.</u>

To construct a recombinant HSV-1 that is deficient in cell-to-cell spread and can also be used to visualize cells containing detectable viral late protein, Us11 tagged with GFP (79) was inserted into an existing glycoprotein H (gH)-null virus, SCgHZ (78) (Fig. 1A). gH is essential for HSV-1 cell entry; it forms a heterodimer complex with gL, and together they activate the viral glycoprotein gB, which mediates fusion between the virus envelope and host cell membrane (80, 81). The GFP-tagged true-late protein is a useful indicator of both lytic infection and reactivation, and Us11-GFP wild-type virus has been used as such in several HSV-1 latency systems (55, 58, 62, 69, 82, 83). The tagged virus has previously been reported to express the full complement of HSV-1 genes, as wild-type Patton strain (79).

We first verified that the resulting virus (named Stayput-GFP) was deficient in cell-to-cell spread in non-neuronal and neuronal cells but otherwise undergoes gene expression and replication as a wild-type virus. The genome sequence and transcriptome of Stayput-GFP was validated by nanopore gDNA and direct RNA sequencing, respectively (Fig. 1B-C). As determined through plaque assay, the ability to produce infectious virus was perturbed in the gH-deletion strains but was rescued using previously constructed gH-complementing cell line, Vero-F6 (Fig. 1D, Supplemental Fig. 1). Importantly, replication in this cell line was indistinguishable from the parent SCgHZ and Us11-GFP Patton strain, which we chose for comparison as this virus has

Figure 1

FIG 1: Stayput-GFP replicates as wild-type but is unable to spread.

(A) Schematic overview of HSV-1 strain SC16, the gH-deletion mutant SCgHZ, and

Stayput-GFP. The gH deletion / LacZ insertion and Us11-GFP insertion sites are shown

by blue and green triangles respectively.

(B-C) Coverage plots derived from (B) nanopore gDNA sequencing and (C) nanopore

direct RNA sequencing of SCgHZ and Stayput-GFP. Sequence read data were aligned

against the SC16 reference genome and demonstrate a drop in coverage at the gH

locus.

(D) Vero F6 cells were infected with Stayput-GFP, SCgHZ, or Us11-GFP at an MOI of

5. Infectious virus was collected over time and titrated on F6 cells (n=3 biological

replicates).

(E-F) Neonatal sympathetic neurons were infected at an MOI of 0.5 PFU/cell with

Stayput-GFP or Us11-GFP in the absence of DNA replication inhibitors. Us11-GFP-

positive neurons were counted over time (n=3 biological replicates). Shapiro-Wilk

normality test. Unpaired student's t test between Us11-GFP and Stayput-GFP.

****p<0.0001. The means and SEMs are shown.

(G) Neonatal sympathetic neurons were infected with Stayput-GFP, SCgHZ, or Us11-

GFP at an MOI of 0.5 PFU/cell. Neurons were fixed at 6 hours post-infection and

stained for viral protein ICP4 (red) along controls β III tubulin (green) and Hoescht stain

(blue).

Vero cells

gH-complementing
Vero-F6 cells

Supplemental Figure 1

<u>Supplemental FIG 1: Stayput-GFP is unable to spread.</u>

Plaque forming assay of Stayput-GFP on Vero (left) or gH-complementing Vero F6

(right) cells 48 hours post-infection at a 10^{-7} dilution. GFP-positive area in example

image reported top left of each image. Titer of viral stock reported bottom left of each

image.

previously been used for latent infection studies in primary neurons.

To demonstrate cell-to-cell spread deficiency in neurons, we quantified GFP-

positive neurons over time. Infection of murine sympathetic neurons at an MOI of 0.5

PFU/cell resulted in detectable GFP-positive neurons, which remained constant after 24

hours post-infection (Fig. 1E), while the surrounding GFP-negative neurons remained

GFP-negative (Fig. 1F). This contrasts with the wild-type US11-GFP condition where

the number of lytic-infected neurons increased substantially over time. Therefore,

Stayput-GFP is unable to spread in neurons. At 6 hours post-infection, viral protein

ICP4-positive neurons were indistinguishable between SCgHZ, Stayput-GFP, or wild-

type Us11-GFP infected neurons. (Fig. 1G). Therefore, Stayput-GFP neuronal infection

was equivalent to Us11-GFP upon initial infection but failed to spread within the

neuronal culture.

Reactivation of Stayput-GFP in a primary neuronal model

We next wanted to confirm that Stayput-GFP undergoes reactivation in a manner

comparable to the backbone SCgHZ virus, in addition to the wild-type Us11-GFP virus

(Fig. 2A). Therefore, we infected mouse primary sympathetic neurons in the presence of

viral DNA replication inhibitor ACV (56, 69-72). ACV prevents the production of

infectious virus, and thus superinfection of the cultures. Following infection, ACV was

removed, and reactivation was triggered adding a PI3-kinase inhibitor LY294002, which

mimics loss of a branch of the NGF-signaling pathway in neurons (84) and has

previously been found to induce reactivation (56, 65, 69, 83). Following the addition of

the reactivation stimulus, viral gene expression increased uniformly between the

Stayput-GFP, SCgHZ, and wild-type Us11-GFP at 20 hours post-stimulus, the time-

point of Phase I gene expression (Fig. 2C). Viral gene expression continued to increase

from 20 to 48 hours for all three viruses, and GFP-positive neurons were visible for both

GFP-tagged viruses by 48 hours (Fig. 2B, 2D). This indicates that even in the absence

of cell-to-cell spread, reactivation progressed over a 48-hour period, initiating with viral

mRNA production and later detection of viral late protein.

By 72 hours-post stimulus, GFP and viral gene transcription were significantly

up-regulated in wild-type Us11-GFP in comparison to Stayput-GFP or SCgHZ,

suggesting that at this time-point, the readout of reactivation for wild-type Us11-GFP is

confounded by cell-to-cell spread (Fig. 2B, 2E). We are therefore unable to differentiate

between genuine reactivation and downstream cell-to-cell spread using a wild-type

virus. Previous attempts to reduce cell-to-cell spread include using pooled human

gamma globulin or the viral DNA packaging inhibitor WAY-150138 (69, 85-87).

However, using Stayput-GFP offers a built-in mechanism to prevent cell-to-cell spread

during reactivation and permit quantification of the progression to reactivation at the

single neuron level.

Figure 2

<u>FIG 2: Stayput-GFP in a latency and reactivation model using ACV to promote latency</u>

<u>establishment</u>

(A) The latency and reactivation model scheme. Neonatal sympathetic neurons were

infected with Stayput-GFP, parent virus SCgHZ, or wild-type Us11-GFP at an MOI of

7.5 PFU/cell in the presence of ACV (50 μM). 6 days later, ACV was removed, and 2

days later, cultures were reactivated with LY294002 (20 μM). The numbers of GFP

positive neurons in a single well (containing approximately 5,000 neurons) for Stayput-

GFP and wild-type Us11-GFP were counted over time (B). Viral gene expression also

was quantified by RT-qPCR for immediate early (*ICP27*), early (*ICP8*), and late (*gC*)

genes at 20 hours (C), 48 hours (D), and 72 hours (E) post-stimulus. Relative

expression to un-reactivated samples and cellular control (mGAPDH). n=6 biological

replicates from 3 litters. Normality determined by Kolmogorov-Smirnov test (B-E). Mann-

Whitney (B) or Kruskal-Wallis with comparison of means (C-E). *p<0.05, **p<0.01,

***p<0.001. The means and SEMs are represented. Individual biological replicates are

indicating in C-E.

<u>Stayput-GFP can be used to create a quiescence model of neuronal infection in the</u>

<u>absence of viral DNA replication inhibitors</u>

As demonstrated in Figure 1, when wild-type virus is used for neuronal infection,

superinfection of the cultures occurs, and a latent infection cannot be established. By

infecting with Stayput-GFP, we posited that we could create a model of latency

establishment without the use of DNA replication inhibitors. Following the infection of

neonatal sympathetic neurons at an MOI of 7.5 PFU/cell, the number of GFP-positive

neurons emerged by 1-day post-infection, increased until 3 days post-infection, and

then decreased until reaching zero by 30 days post-infection (Fig. 3A). The length of

time required for Us11-GFP to be lost from the cultures was surprising and may result

from the previously characterized restricted cell death in neurons (88), or a gradual

shut-off in protein synthesis in a sub-population of neurons that survive even late gene

expression. Single-cell tracking demonstrates that most neurons that become GFP-

positive also end up staining positive for cell death marker SYTOX™ Orange (Fig. 3B).

All classes of lytic viral gene expression emerged by 1-day post-infection, increased by

3 days post-infection, and then decreased over the span of 40-days post-infection (Fig.

3C-E). The latency associated transcript (LAT) was detectable by 1-day post-infection

and increased by 7 days post-infection (Fig. 3F). In contrast to the lytic transcripts, LAT

expression was maintained over the infection scheme of 30-40 days. In comparison to

lytic transcripts, LAT expression was approximately 400-fold higher from 30 days

onwards. This indicated that LAT-positive neurons persisted over this period, likely

reflecting entry into quiescence. In agreement with this hypothesis, at 40 days post-

infection there are approximately 200 viral DNA copies per cell, demonstrating that

Figure 3

A

Supplemental Figure 2

<u>Supplemental FIG 2: Stayput-GFP can be used to create a quiescence model in the</u>

<u>absence of viral DNA replication inhibitors in adult sensory neurons</u>

Neurons isolated from the trigeminal ganglia (TG) of female mice were infected with

Stayput-GFP at an MOI of 7.5 PFU/cell. The resolution of lytic infection was monitored

over time by imaging and counting GFP-positive neurons. n=12, 3 biological replicates.

The mean and SEM are shown.

viral genomes persist (Fig. 3H). Together, these data show that infection of sympathetic

neurons with a cell-to-cell defective virus results in a remaining population of neurons

containing viral genomes and the LAT transcript. Notably, this mimics events following

in vivo infection of mice.

We were also interested to determine whether a similar quiescent infection could

be established in adult sensory neurons, as these are also a commonly used model of

neuronal HSV-1 infection *in vitro* (66). Following infection with Stayput-GFP in primary

trigeminal ganglia (TG) neurons, GFP also increased by 1-day post-infection and then

was lost over time (Supplemental Fig. 2). Of interest, GFP was repeatedly lost within 15

days, a shorter period than that which is observed in neonatal sympathetic neurons.

Therefore, the Stayput-GFP virus can be used to establish a quiescent infection of both

sympathetic and sensory neurons *in vitro* without the use of viral DNA replication

inhibitors.

The ability of HSV-1 to undergo reactivation decreases with length of time infected

The presence of viral genomes and LAT transcripts suggested that HSV-1 had

established a quiescent infection 30 days post-infection; therefore, we hypothesized that

some genomes enter latency, which is defined by an ability to reactivate in response to

a stimulus. We thus attempted to reactivate cultures with LY294002 (20 μM). However,

we were unable to detect an increase in GFP-positive neurons after the addition of the

trigger (Fig. 4A). We were also unable to detect a change immediate early, early, or late

transcripts (data not shown). This was unexpected as LY294002 has repeatedly been

Figure 4

FIG 4: Reactivation decreases with length of time infected

Sympathetic neurons were infected with Stayput-GFP at an MOI of 7.5 PFU/cell and

were treated with LY294002 when GFP-positive neurons were no longer detected

(approximately 30 days post-infection). GFP-positive neurons were quantified over time;

peak GFP (48 hours post-stimulus) is represented (A). Neonatal SCGs were infected at

age postnatal day 8 (P8) with Stayput-GFP in the presence of ACV. ACV was removed

6 days post-infection and reactivation was triggered at the indicated times post-infection

(B). Neonatal SCGs were infected as above after different lengths of time *in vitro*,

representing indicated postnatal ages, and reactivated 8 days post-infection with

LY294002 (C). n=12 biological replicates from 3 litters. Normality determined by

Kolmogorov-Smirnov test. Unpaired student's t-test (B) or Mann-Whitney (A, C) based

on normality of data. **p<0.01, ***p<0.001, ****p<0.0001. Individual biological replicates

along with the means and SEMs are shown.

shown to elicit robust reactivation *in vitro* and was able to induce Stayput-GFP in a

model using ACV to promote latency establishment (Fig. 2). Therefore, we sought to

determine whether the inability to induce reactivation was due to the lack of ACV or the

more prolonged time between initial infection and the addition of the reactivation

stimulus. We infected neonatal cultures in the presence of ACV and reactivated over

increasing lengths of time. ACV was removed from all cultures 6 days post-infection.

We found that the number of GFP-positive neurons following addition of LY294002

decreased as the length of time infected increased (Fig. 4B). This was not likely due to

a loss of viral genomes, as viral genome copy number and LAT expression remained

constant over this period (Fig. 3F-G) and therefore instead reflected a more repressed

viral genome unable to undergo reactivation upon PI3-kinase inhibition.

Neurons are known to undergo intrinsic maturation, even in culture (89, 90) and

therefore the increased age of the neuron could have also impacted the ability of HSV-1

to undergo reactivation. Therefore, we investigated how reactivation changed with

increased neuronal maturation (Fig. 4C). We infected cultured neurons with Stayput-

GFP at an MOI of 7.5 at the postnatal (P) ages of P8, P16, and P24 and then

reactivated 8 days later. These postnatal ages of infection were chosen to reactivate at

the same ages of reactivation in Fig 4B. Importantly, we did not detect a decrease in

reactivation output as the age of the neuron increased. Together, these data indicate

that the decreased ability of Stayput-GFP to reactivate in a model that did not require

ACV to establish quiescence was due to the longer time frame of infection and not

associated with a lack of ACV in the cultures or increased age of the neuron.

Viral gene expression can be induced following long-term quiescent infection when multiple triggers are combined

We next sought to determine whether other known stimuli of HSV-1 reactivation

could induce Us11-GFP expression, indicative of entry into reactivation. We attempted a

number of triggers including forskolin (91-94) and heat-shock/hyperthermia (95-102),

which are both known inducers of HSV-1 reactivation. Alone these stimuli did not induce

Us11-GFP expression or viral lytic mRNA induction (data not shown). However, when

heat-shock (43°C for 3h), in addition to forskolin (30 μM) and LY294002 (both pulsed for

20 hours) were combined, Us11-GFP positive neurons were detected at 48 hours post-

stimulus, indicating progression to reactivation (Fig. 5A). Superinfection was also used

as this can induce rapid and robust reactivation, likely resulting in delivery of viral

tegument proteins. In comparison to superinfection, the combined heat-

shock/forskolin/LY294002 trigger resulted in reduced entry into reactivation/Us11-GFP

expression, indicating that only a sub-population of neurons undergo reactivation with

this combined trigger, which is consistent with previous studies investigating the

mechanism of HSV-1 reactivation both *in vivo* and *in vitro* (25, 70, 103, 104).

We went on to investigate whether viral mRNA expression was induced prior to

the detection of Us11-GFP positive neurons. Using the triple stimulus, we detected an

increase in viral lytic transcripts as early as 10 hours post-stimulus, which continued to

increase, peaking at 48 hours post-stimulus (Fig. 5B-D, Supplemental Fig. 3A).

Detection of immediate-early and early transcripts was slightly more robust than late

transcripts,

Figure 5

<u>FIG 5: Viral gene expression can be restarted following latency establishment</u>

Neonatal sympathetic neurons or adult sensory neurons were infected at an MOI of 7.5

PFU/cell with Stayput-GFP in the absence of viral DNA replication inhibitors. Following

the loss of GFP, signaling quiescence of the culture, wells were reactivated with a

variety of triggers, including combinations of LY-294002 (20 μM), forskolin (60 μM), and

heat shock (43°C for 3 hours), as well as a superinfection with untagged F strain at an

MOI of 10 PFU/cell. GFP was quantified over time, and the peak GFP, at 48 hours post-

stimulus, is depicted. N=12 biological replicates (A, E). Immediate early (B, F), early (C,

G) and late (D, H) viral transcripts were investigated over time following the stimulus.

Neonatal sympathetic neurons n=9 biological replicates from 3 litters. Mann-Whitney

against 0 hours (A-D). Adult sensory neurons n=9 biological replicates (E) or n=3

replicates (F, G, H). Normality determined by Kolmogorov-Smirnov test. *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001. Individual biological replicates along with the

means and SEMs are shown.

Supplemental Figure 3

although all late transcripts were significantly induced by 20 hours post-stimulus. (Fig.

5D, Supplemental Fig. 3B-E).

We also confirmed that the triple combinatorial stimulus elicits robust GFP re-

emergence in adult TGs (Fig. 5E). Similar to what we observed in the sympathetic

neurons, Us11-GFP positive neurons were detected by 48 hours post-stimulus.

Intriguingly, superinfection only induced GFP expression to equivalent levels as the

triple stimuli, which may be reflective of the repressive nature of sub-population of

mature sensory neurons to lytic replication and reactivation (66, 67). Together, these

data indicate that *in vitro* models of latency and reactivation can be established in

sympathetic and sensory neurons in the absence of ACV and reactivation can be

induced using a combination of triggers. In both models, a wave of lytic mRNA

expression was detected prior to the appearance of Us11-GFP positive neurons.

Neurons infected with Stayput-GFP undergo a DLK-dependent Phase I of reactivation

Phase I reactivation has largely been investigated using *in vitro* models in which

ACV has been used to promote latency establishment. In addition, Phase I has been

found to occur with the single triggers of forskolin or LY294002 (25, 56, 58). The

requirement for the multiple triggers for reactivation suggested that multiple cell-

signaling pathways converged to have a synergistic effect and induce reactivation even

in this more repressive model. Therefore, we were interested to determine whether the

characteristics of Phase I reactivation occurred in the model of quiescent infection

established in the absence of ACV and using the more robust trigger to induce

Figure 6

<u>FIG 6: Reactivation is dependent on DLK</u>

Cultures were infected with Stayput-GFP at an MOI of 7.5 PFU/cell in the absence of

ACV. Following loss of GFP, cultures were reactivated with a combination of LY294002,

forskolin, and heat shock in the presence of DLK inhibitor GNE-3511 (4 µM). Immediate

early (*ICP27*) and early (*ICP8/UL30*) viral genes (A-C) were investigated at 12.5 hours-

post stimulus, and GFP was counted over time (D). Peak GFP, consistently around 48

hours post-stimulus, is presented. n=9 biological replicates from 3 litters. Normality

determined by Kolmogorov-Smirnov. Mann-Whitney test. (*p<0.05, **p<0.01,

***p<0.001, ****p<0.0001). The mean and SEM are shown.

reactivation. Because viral gene expression was first detected at 10-15 hours post-

stimulus, potential Phase I viral transcription was investigated at 12.5 hours post-

stimulus by RT-qPCR and Phase II was investigated when Us11-GFP positive neurons

could be detected (48 hours post-stimulus). A characteristic of Phase I expression is the

requirement of the stress kinase dual leucine zipper kinase (DLK) (56, 58). Therefore,

using the DLK-specific inhibitor GNE3511 4 µM (105), we investigated the effect of DLK

on reactivation in our system. We found up-regulation of immediate early/early

transcripts 12.5 hours post-stimulus was eliminated with the addition of the DLK inhibitor

(Fig. 6A-C). Further, full reactivation, demonstrated by peak GFP expression at 48

hours, was also reduced to baseline levels upon the addition of the DLK inhibitor (Fig.

6D). Therefore, using our new model of HSV-1 established in the absence of ACV, our

data demonstrate that the initiation of viral lytic gene expression is dependent on DLK

activity.

In addition to the dependence on DLK activity, a further characteristic of Phase I

gene expression is the induction of viral mRNA transcripts independently of the activity

of histone de-methylase enzymes required for full reactivation. Therefore, we also

triggered reactivation in the presence of de-methylase inhibitors. GSK-J4 is known to

specifically inhibit the H3K27 histone demethylases UTX and JMJD3 (106) and has

previously been found to inhibit HSV-1 reactivation but not Phase I gene expression

(56, 58). OG-LOO2 is an LSD1 specific inhibitor. LSD1 has previously been shown to

be involved in removal of H3K9me2 from HSV-1 genomes and its activity is required for

full reactivation but not Phase I gene expression (42, 56, 58). However, because

histone de-methylase inhibitor independent gene expression in reactivation has only

been observed in models using ACV to establish latency, we investigated their role lytic

gene expression following stimulation of quiescent cultures with the triple trigger. Full

reactivation was reduced in the presence of OG-LOO2 or GSK-J4 as demonstrated by

GFP-positive neurons at 48 hours (Fig. 7D). However, the initial expression of lytic

transcripts at 12.5 hours post-stimulation was not inhibited by either OG-LOO2 or GSK-

J4 (Fig. 7A-C). Therefore, the initiation of gene expression in a manner that is

independent of histone demethylase activity occurs when reactivation is induced by a

triple stimulus and in a primary neuronal model in which latency was established without

ACV.

During Phase I, there is no detectable replication of viral genomes even though true late

gene expression occurs (56, 58). In addition, using ACV models to establish latency,

late gene expression has previously been found to occur to equivalent levels when viral

DNA replication is inhibited during reactivation. Although we did observe late gene

expression during the initial period of lytic gene induction (12.5-20 hours), the increase

was less robust than IE and E genes and appeared slightly delayed, not reaching

significance for all analyzed late transcripts until 24 hours post-stimulus (Fig. 5D,

Supplemental Fig. 3). Therefore, we investigated whether this late gene induction was

dependent on viral DNA replication by reactivating in the presence of ACV. The addition

of ACV inhibited entry in full reactivation, demonstrated by Us11-GFP positive neurons

at 48 hours post-stimulus, indicating that ACV was capable of blocking robust late gene

Figure 7

Figure 7: The early phase of lytic gene expression following a reactivation stimulus is
independent of demethylase activity.

Cultures were infected with Stayput-GFP at an MOI of 7.5 PFU/cell. Following loss of
GFP, cultures were reactivated with a combination of LY-294002, forskolin, and heat
shock in the presence of H3K27 demethylase inhibitor GSK-J4 (2 μM) or H3K9
demethylase inhibitor OG-L002 (20 μM). Immediate early (*ICP27*) and early
(*ICP8/UL30*) viral genes (A-C) were investigated at 12.5 hours-post stimulus and GFP
was counted over time (D). Peak GFP is presented. n=3 biological replicates from 3
litters. Normality determined by Kolmogorov-Smirnov. Mann-Whitney. *p<0.05,
**p<0.01, ***p<0.001, ****p<0.0001. The mean and SEM are shown.

expression at this late time-point (Fig. 8D). However, the addition of ACV did not inhibit

the induction of lytic transcripts at 22 hours post-stimulus. Importantly, we included

multiple true late genes in this analysis and all were induced to equivalent levels in the

presence and absence of ACV (Fig. 8A-C). Therefore, the initial expression of late

genes following a reactivation stimulus is independent of viral DNA replication. In

summary, using a model system in which a quiescent infection is established without

the need for ACV, all the previous characteristics of Phase I gene expression

(dependence on DLK and independence of histone demethylase activity and viral DNA

replication) were still observed.

Figure 8

Figure 8: Differential dependence on viral DNA replication between Phase I and II

reactivation.

Cultures were infected with Stayput-GFP at an MOI of 7.5 PFU/cell in the absence of

ACV. Following loss of GFP, cultures were reactivated with a combination of LY-

294002, forskolin, and heat shock in the presence of ACV (50 μM). Late (*VP16, gC,*

UL10) genes (A-C) were investigated at 22 hours-post stimulus. GFP was counted over

time and peak GFP is presented (D). n=12 biological replicates, normality determined

by Kolmogorov-Smirnov. Mann-Whitney. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

The mean and SEM are shown.

Discussion

We envision multiple potential uses for the Stayput-GFP virus model developed

here for investigating HSV-1 neuronal infection *in vitro*. The Stayput-GFP virus is

advantageous in models using DNA replication inhibitors to promote latency

establishment because it allows for the differentiation of *de novo* initial viral gene

expression/protein synthesis events and readouts from those that are a consequence of

downstream cell-to-cell spread. There have been previous attempts to control cell-to-

cell spread during reactivation in this model, for example, using the DNA packaging

inhibitor WAY150138 (69, 85-87). However, the use of Stayput-GFP means that

synthesis of WAY150138 would not be required. In addition, even in systems where

ACV is used, there can be low levels of lytic replication or spontaneous reactivation

after removal of ACV from cultures. This could impact studies looking at the

mechanisms of gene expression during reactivation. The use of Stayput-GFP helps limit

spontaneous reactivation events by inhibiting subsequent cell-to-cell spread, while at

the same time monitoring neurons that escape quiescence. Further, the GFP tag serves

as an imaging indicator in real time of when lytic infection is resolved and latency is

considered established. We are also able to track viral DNA replication and downstream

late viral transcription and protein synthesis during the latency establishment process *in*

vitro. The fate of lytic neurons, whether they undergo cell death or turn off gene

expression programs and enter the latency pool, can also be investigated by tracking

GFP and cell death at a single-cell level.

There are caveats and limitations to our system. Although there is some

discrepancy in what defines reactivation (107), it is ultimately defined by the production

of infectious virus. Due to the nature of the gH-deletion virus, *de novo* virus is by design

non-infectious and we are unable to demonstrate textbook reactivation. One could

theoretically resolve this issue by generating a cell line that expresses gH on its surface.

However, gH activates a fusogen and is prone to cellular synctia formation in a

constitutive expression system (81). While we are unable to demonstrate infectious

virus production, we do demonstrate the re-emergence of all classes of viral gene

transcripts and late viral protein synthesis.

An intriguing finding in our study is that reactivation output decreases as length of

infection increases. We find that, at least in the subtype of sympathetic neurons, this is

independent of neuronal age or length of time in culture within the 30-day infection

timeframe. The data demonstrating the persistence of viral DNA load from day 21-40

post-infection suggests this phenotype is also not due to a passive loss of viral

genomes over time in culture. We also demonstrate this phenotype is independent of

ACV use. A potential alternative explanation is that the viral genome becomes

increasingly chromatinized over time, leading to a more repressive phenotype. In

support of this hypothesis, the association of the facultative heterochromatin mark

H3K27me3 with the HSV-1 genome increases dramatically between 10- and 15-days

post-infection *in vivo* (50). The kinetics of H3K27me3 deposition remain to be

investigated *in vitro*, but if they are mirrored this could suggest that active

chromatinization and reinforcement of silencing continues even after initial shut-down of

viral gene expression. In the cellular context, H3K27me3 is linked with the recruitment

of canonical Polycomb repressor complex 1 (cPRC1), which may reinforce silencing

through long-range chromosomal interactions or 3D compaction (54, 108, 109). It is

therefore possible that even following H3K27me3 formation on the genome, there are

additional layers of protein recruitment that build up over time. In addition, it is also

possible that the accumulation of viral non-coding RNAs expressed in latency could

impact cellular pathways resulting in decreased signaling to the viral genome for

reactivation. The use of the Stayput-GFP model system will permit these different

avenues to be explored.

Our model system fits the criteria of Phase I of reactivation, which has previously

been explored in *in vitro* systems using a DNA replication inhibitor. These data are

interesting considering the discrepancies in conclusions drawn about reactivation

between *in vitro* and *in vivo* modeling. In our model system, as in other *in vitro* systems,

the progression to reactivation involves multiple waves of viral lytic gene expression.

We find that viral transcripts are upregulated in a synchronous fashion, approximately

10 hours post-stimulus. There is evidence *ex vivo* for a Phase I as all classes of viral

gene are expressed in a disordered, non-cascade fashion when a combination of

explant and nerve-growth factor deprivation are used (110). However, in other models

of reactivation *ex vivo*, there is evidence that Phase-I-like gene expression may not

occur, especially from studies investigating the requirement for histone demethylase

inhibitors. These latter experiments used explant (axotomy) to induce reactivation and

found that the earliest induction of lytic gene expression is dependent on lysine 9

demethylase activity (41, 42). These discrepancies between *in vitro* and *ex vivo* models,

or within *ex vivo* models, may result from the different trigger used to induce

reactivation, although additional differences include the subtype of neuron (sensory

versus sympathetic) or currently unknown effects of latency established *in vivo*.

Importantly, here we have demonstrated that any potential differences between *in vivo*

and *in vitro* observations on the mechanisms of reactivation do not result from the use

of ACV to establish a quiescent infection. Further work using the Stayput-GFP model

system will help elucidate how changes in the host immune response, neuronal

subtype, and stimulus used can potentially alter the mechanisms of viral gene

expression during reactivation.

Phase I, in addition to occurring synchronously and independently of histone

demethylases, is also proposed to occur in the absence of viral protein synthesis. In an

in vitro model employing ACV during latency establishment and stimulus LY294002

during reactivation, it is demonstrated that initial viral transcription occurs before the

appearance of viral late protein synthesis and, specifically, independently of viral

transactivator VP16 (25, 55). Therefore, cellular host factors must be responsible for

instigating the initial reactivation process. Evidence from an *in vitro* model system has

demonstrated these events are in fact navigated by cellular proteins JNK and DLK (56).

Interestingly, host cell proteins may also be implicated in restricting the full reactivation

process, including Gadd45b which appears to antagonize the HSV-1 late expression

program to prevent full reactivation (55). Interestingly, Gadd45b mRNA is increased in

response to LY294002 only in infected neurons, suggesting perhaps that a viral protein

may be mediating the gatekeeping from Phase I to Phase II reactivation.

In our system, Phase I gene expression was dependent on the neuronal

regulator of JNK activity, DLK, highlighting the central role of DLK in HSV-1 reactivation.

DLK is a cell protein implicated in neuronal stress signaling upstream of cellular protein

JNK (111). It has previously been found to be essential for HSV-1 reactivation following

PI3-kinase inhibition (56), as well as neuronal hyper-excitability through forskolin (58).

However, it has not, until now, been shown to be central to reactivation mediated by

heat shock. Although heat shock has been used as a trigger for HSV-1 reactivation (95-

102), the downstream molecular events following this stimulus are not well elucidated.

Heat shock has been shown to upregulate the expression of ICP0 and ICP22 in an NF-

Y site-dependent manner *in vitro* (98). Multiple studies have demonstrated that heat

shock during reactivation leads to the up-regulation of heat shock proteins, although

none of them relate to DLK. Following hyperthermia-induced reactivation *in vivo*, heat

shock protein H2P60 and HSP40 have been demonstrated to be up-regulated (102).

Components of the heat shock response pathway have also been demonstrated to be

up-regulated by LY294002 treatment in an *in vitro* system (55), including HSP70. In fact,

in this same system, treatment with cultures of heat shock factor 1 (HSF-1) activator

compound causes robust reactivation. Outside of the virological context, heat shock

protein chaperone HSP90 has been shown to bind and maintain DLK stability *in vivo*

and it is specifically required for DLK function following axon injury signaling (112). It is

a possibility that heat shock in our system is enhancing the function of DLK. Therefore,

multiple signals may converge on DLK, which is then able to activate JNK and protein

histone phosphorylation and to promote lytic gene expression from the heterochromatin-

associated viral genome for reactivation to occur. Indeed, synergy has been

demonstrated to enhance DLK activity in neurons (113). This central role for DLK is

especially important as it is largely a neuron-specific protein that regulates the response

to multiple forms of stress (114) and is therefore a potential target for novel therapeutics

that would prevent HSV-1 gene expression and ultimately reactivation.

Materials & Methods

Reagents

Compounds used in the study are as follows: Acycloguanosine, LY 294002, Forskolin,

GNE-3511, GSK-J4, OG-L002. Compound concentrations were used based on

previously published IC50s and assessed for neuronal toxicity using the cell body and

axon health and degeneration index as previously published (58).

Preparation of HSV-1 virus stocks

Stayput-GFP, as well as SCgHZ, is propagated and titrated on previously constructed

gH-complementing F6 cells, which contain copies of the gH gene under the control of

an HSV-1 gD promoter, as described in (78). Vero F6 cells were maintained in

Dulbecco's Modified Eagle's Medium (Gibco) supplemented with 10% FetaPlex (Gemini

BioProducts) and 250 μg/mL of G418/Geneticin (Gibco).

HSV-1 stocks of eGFP-Us11 Patton were grown and titrated on Vero cells obtained

from the American Type Culture Collection (Manassas, VA). Cells were maintained in

Dulbecco's Modified Eagle's Medium (Gibco) supplemented with 10% FetalPlex

(Gemini Bio-Products) and 2 mM L-Glutamine. eGFP-Us11 Patton (HSV-1 Patton strain

with eGFP reporter protein fused to true late protein Us11 (79) was kindly provided by

Dr. Ian Mohr at New York University.

<u>Construction of Stayput-GFP</u>

Stayput-GFP was created by inserting an eUs11-GFP tag into the previously created

gH-deficient HSV-1 SCgHZ virus (strain SC16) through co-transfection of SCgHZ viral

DNA and pSXZY-eGFP-Us11 plasmid (79) in Vero F6 cells, Transfections were carried

on in 6 well plates using 2 μg of viral DNA and 2 μg of pSXZY-eGFP-Us11 and 8ul of

jetPRIME® (Polyplus). GFP-positive plaques were subjected to 4 rounds of plaque

purification.

<u>Viral genomic DNA sequencing</u>

Genomic DNA was isolated from stocks of SCgHZ and Stayput-GFP using the Qiagen

DNeasy Blood and Tissue kit. 250 ng of gDNA was used as input for the SQK-LSK109

genomic DNA by ligation protocol (Oxford Nanopore Technologies Ltd.) and resulting

libraries were loaded onto individual R9.4.1 flongles for 20 hours of sequencing.

Resulting raw fast5 datasets were basecalled using Guppy v4.2.2 (-f FLO-MIN106 -k

SQK-RNA002) with only reads passing filter used for subsequent analyses. *De novo*

assembly was performed as follows. First, HSV-1 reads were extracted from the total

pool by aligning against the HSV-1 strain SC16 reference genome (MN159383.1) using

MiniMap2. HSV-1 reads shorter than 1,000 nt were subsequently filtered out prior to

performing de novo assembly with Canu (115). Canu yielded a single circular contig for

both SCgHZ and Stayput-GFP which was manually linearized at the canonical HSV-1

start site. Four rounds of genome polishing was performed using racon (116) before a

final round of SNP calling was achieved using medaka

(https://github.com/nanoporetech/medaka).

<u>Direct RNA sequencing</u>

For both SCgHZ and Stayput-GFP, NHDFs (passage 13) were infected at an MOI of 10.

At 6 hours post infection, the supernatant was removed and cells washed once with

PBS prior to lysing with 8 ml Trizol. Total RNA was extracted according to Trizol

manufacturer's instructions and eluted in nuclease-free water. Nanopore direct RNA-

Seq libraries were prepared for each sample using between ~500 ng of poly(A) RNA

that was previously isolated from 30 μg of total RNA using the Dynabeads™ mRNA

Purification Kit (Invitrogen, 61006). The poly(A) RNA was spiked with 0.5 μL of a

synthetic Enolase 2 (ENO2) calibration RNA (Oxford Nanopore Technologies Ltd.) and

direct RNA-Seq libraries prepared according to the standard SQK-RNA002 protocol

(Oxford Nanopore Technologies Ltd.). Raw fast5 datasets were then basecalled using

Guppy v4.2.2 (-f FLO-MIN106 -k SQK-RNA002) with only reads passing filter used for

subsequent analyses. Sequence reads were aligned against the Stayput-GFP genome

using MiniMap2 (Li, 2018) (-ax splice -k14 -uf --secondary=no), with subsequent parsing

through SAMtools and BEDtools (117, 118). Here sequence reads were filtered to retain

only primary alignments (Alignment flag 0 (top strand) or 16 (bottom strand)). Figures

associated with this study were generated using the R packages Gviz (119) and

GenomicRanges (120).

<u>Single-step growth curve</u>

1×10^5 Vero F6 cells were seeded per well in a 24-well plate. 24 hours later, when the

cells had reached approximately 90% confluency, cells were infected with Stayput-GFP,

SCgHZ, or wild-type Patton Us11-GFP at an MOI of 5 PFU/cell for 1 hour at 37°C with

gentle rocking (60 rpm). Inoculum was then removed and replaced with 400 µL of

media. At the specified time-points, 400 µL of 9% sterile milk was added to the well and

the plates were frozen in the -80C. Following three freeze thaw cycles, titrations were

carried out on Vero F6 cells.

<u>Primary neuronal cultures</u>

Sympathetic neurons from the superior cervical ganglia (SCG) of post-natal day 0–2

(P0-P2) CD1 Mice (Charles River Laboratories) were dissected as previously described

(Cliffe et al., 2015). Sensory neurons from the trigeminal ganglia (TG) or adult (P21-24)

CD1 Mice (Charles River Laboratories) were dissected as previously described (121).

Rodent handling and husbandry were carried out under animal protocols approved by

the Animal Care and Use Committee of the University of Virginia (UVA). Ganglia were

briefly kept in Leibovitz's L-15 media with 2.05 mM L-Glutamine before dissociation in

Collagenase Type IV (1 mg/mL) followed by Trypsin (2.5 mg/mL) for 20 min each at

37℃. Dissociated ganglia were triturated, and approximately 5,000 neurons per well

were plated onto rat tail collagen in a 24-well plate. Sympathetic neurons were

maintained in CM1 (Neurobasal Medium supplemented with PRIME-XV IS21 Neuronal

Supplement (Irvine Scientific), 50 ng/mL Mouse NGF 2.5S, 2 mM L-Glutamine, and

Primocin). Aphidicolin (3.3 mg/mL) was added to the CM1 for the first 5 days post-

dissection to select against proliferating cells. Sensory neurons were maintained in the

same media supplemented with GDNF (50 ng/ml; Peprotech 450-44), and more

Aphidicolin (6.6 mg/mL) was used for the first 5 days as more non-neuronal cells tend to

be dissected with this neuron type.

<u>Establishment and reactivation of latent HSV-1 infection in primary neurons using anti-</u>

<u>mitotic agents</u>

After being cultured for at least 24 hours without anti-mitotic agent aphidicolin, neonatal

SCGs were infected P6-P8 with either Us11-GFP, SCgHZ, or Stayput-GFP at an MOI of

7.5 PFU/cell assuming 5,000 cells/well in DPBS +CaCl2 +MgCl2 supplemented with 1%

Fetal Bovine Serum, 4.5 g/L glucose, and 10 mM Acyclovir (ACV) for 3 hr at 37 ℃.

Post-infection, inoculum was replaced with CM1 containing 50 mM ACV for 5–6 days,

followed by CM1 without ACV. Reactivation was carried out in BrainPhys (Stem Cell

Technologies) supplemented with 2 mM L-Glutamine, 10% Fetal Bovine Serum, Mouse

NGF 2.5S (50 ng/mL) and Primocin. Reactivation was quantified by counting number of

GFP-positive neurons or performing Reverse Transcription Quantitative PCR (RT-

qPCR) for HSV-1 lytic transcripts.

<u>Establishment and reactivation of latent HSV-1 infection in primary neurons without</u>

<u>using anti-mitotic agents</u>

After being cultured for at least 24 hours without anti-mitotic agent aphidicolin, neurons

were infected P6-P8 for neonatal SCGs and P30-32 for adult TGs. The cultures were

infected with Stayput-GFP at an MOI of 7.5 PFU/cell assuming 5,000 cells/well in DPBS

+CaCl2 +MgCl2 supplemented with 1% Fetal Bovine Serum, and 4.5 g/L glucose for 3

hr at 37 ℃. Post-infection, inoculum was replaced with CM1. Following the loss of GFP,

reactivation was carried out in BrainPhys (Stem Cell Technologies) supplemented with

2 mM L-Glutamine, 10% Fetal Bovine Serum, Mouse NGF 2.5S (50 ng/mL) and

Primocin with LY294002 20 μM and forskolin 60 μM (pulsed for 20 hours), as well as

and heat shock (3 hours at 43℃).

<u>Immunofluorescence</u>

Neurons were fixed for 15 min in 4% Formaldehyde and blocked in 5% Bovine Serum

Albumin and 0.3% Triton X-100 and incubated overnight in primary antibody. Following

primary antibody treatment, neurons were incubated for 1 hr in Alexa Fluor 488-, 555-,

and 647-conjugated secondary antibodies for multi-color imaging (Invitrogen). Nuclei

were stained with Hoechst 33258 (Life Technologies). Images were acquired using an

sCMOS charge-coupled device camera (pco.edge) mounted on a Nikon Eclipse Ti

Inverted Epifluorescent microscope using NIS-Elements software (Nikon). Images were

analyzed using ImageJ.

<u>Analysis of mRNA expression and viral genome copy number</u>

To assess expression of HSV-1 lytic mRNA, total RNA was extracted from

approximately 5,000 neurons using the Quick-RNA Miniprep Kit (Zymo Research) with

an on-column DNase I digestion. mRNA was converted to cDNA using the Maxima

cDNA synthesis kit (Thermo Fisher) using random hexamers for first-strand synthesis

and equal amounts of RNA (20–30 ng/ reaction). To assess viral DNA load, total DNA

was extracted from approximately 5,000 neurons using the Quick-DNA Miniprep Plus

Kit (Zymo Research). qPCR was carried out using PowerUp^{TM} SYBR Green Master Mix

(Thermoscientific). Relative mRNA/DNA copy numbers were determined using the

Comparative CT ($\Delta\Delta$CT) method normalized to mRNA levels in latently infected

samples. Viral RNAs were normalized to mouse reference gene 18s ribosomal RNA. All

samples were run in duplicate on an Applied Biosystems QuantStudio 6 Flex Real-Time

PCR System and the mean fold change compared to the reference gene calculated.

Exact copy numbers were determined by comparison to standard curves of known DNA

copy number of viral genomes or plasmid containing 18S rRNA (p18S.1 tag was a gift

from Vincent Mauro; Addgene plasmid # 51729 ; http://n2t.net/addgene:51729 ;

RRID:Addgene_51729, (122).

<u>Statistical Analysis</u>

All statistical analyses were performed using Prism V8.4. The normality test and

statistical test used for each figure are denoted in the figure legends. Individual

biological replicates are plotted, each corresponding to a single well of cells/neurons. All

neuronal experiments were repeated from pooled neurons from at least 3 litters.

Acknowledgements

We thank Dr. Ian Mohr at New York University for the gift of the Us11-GFP virus and the pSXZY-GFP-Us11 plasmid. We thank Gary Cohen at the University of Pennsylvania for SCgHZ and Vero F6 cells. This work was supported by National Institutes of Health grants NS105630 (ARC), T32GM008136 (SD), AI130618, AI147163 and AI151358 (ACW), The Owens Family Foundation (ARC), and a German Centre for Infection Research (DZIF) Associate Professorship (DPD).

References

1. Richter ER, Dias JK, Gilbert JE, 2nd, Atherton SS. 2009. Distribution of herpes simplex virus type 1 and varicella zoster virus in ganglia of the human head and neck. J Infect Dis 200:1901-6.
2. Baringer JR, Pisani P. 1994. Herpes simplex virus genomes in human nervous system tissue analyzed by polymerase chain reaction. Ann Neurol 36:823-9.
3. Baringer JR, Swoveland P. 1973. Recovery of herpes-simplex virus from human trigeminal ganglions. N Engl J Med 288:648-50.
4. Warren KG, Brown SM, Wroblewska Z, Gilden D, Koprowski H, Subak-Sharpe J. 1978. Isolation of latent herpes simplex virus from the superior cervical and vagus ganglions of human beings. N Engl J Med 298:1068-9.
5. Singh N, Tscharke DC. 2020. Herpes Simplex Virus Latency Is Noisier the Closer We Look. J Virol 94.
6. Bloom DC. 2016. Alphaherpesvirus Latency: A Dynamic State of Transcription and Reactivation. Adv Virus Res 94:53-80.
7. Speck PG, Simmons A. 1991. Divergent molecular pathways of productive and latent infection with a virulent strain of herpes simplex virus type 1. J Virol 65:4001-5.
8. Speck PG, Simmons A. 1992. Synchronous appearance of antigen-positive and latently infected neurons in spinal ganglia of mice infected with a virulent strain of herpes simplex virus. J Gen Virol 73 (Pt 5):1281-5.
9. Proenca JT, Nelson D, Nicoll MP, Connor V, Efstathiou S. 2016. Analyses of herpes simplex virus type 1 latency and reactivation at the single cell level using fluorescent reporter mice. J Gen Virol 97:767-777.
10. Maillet S, Naas T, Crepin S, Roque-Afonso AM, Lafay F, Efstathiou S, Labetoulle M. 2006. Herpes simplex virus type 1 latently infected neurons differentially express latency-associated and ICP0 transcripts. J Virol 80:9310-21.
11. Proenca JT, Coleman HM, Connor V, Winton DJ, Efstathiou S. 2008. A historical analysis of herpes simplex virus promoter activation in vivo reveals distinct populations of latently infected neurones. J Gen Virol 89:2965-2974.
12. Proenca JT, Coleman HM, Nicoll MP, Connor V, Preston CM, Arthur J, Efstathiou S. 2011. An investigation of herpes simplex virus promoter activity compatible with latency establishment reveals VP16-independent activation of immediate-early promoters in sensory neurones. J Gen Virol 92:2575-2585.
13. Chen SH, Kramer MF, Schaffer PA, Coen DM. 1997. A viral function represses accumulation of transcripts from productive-cycle genes in mouse ganglia latently infected with herpes simplex virus. J Virol 71:5878-84.
14. Tzeng NS, Chung CH, Lin FH, Chiang CP, Yeh CB, Huang SY, Lu RB, Chang HA, Kao YC, Yeh HW, Chiang WS, Chou YC, Tsao CH, Wu YF, Chien WC. 2018. Anti-herpetic Medications and Reduced Risk of Dementia in Patients with Herpes Simplex Virus Infections-a Nationwide, Population-Based Cohort Study in Taiwan. Neurotherapeutics 15:417-429.

15.   Letenneur L, Peres K, Fleury H, Garrigue I, Barberger-Gateau P, Helmer C, Orgogozo JM,
Gauthier S, Dartigues JF. 2008. Seropositivity to herpes simplex virus antibodies and risk of
Alzheimer's disease: a population-based cohort study. PLoS One 3:e3637.

16.   De Chiara G, Piacentini R, Fabiani M, Mastrodonato A, Marcocci ME, Limongi D, Napoletani
G, Protto V, Coluccio P, Celestino I, Li Puma DD, Grassi C, Palamara AT. 2019. Recurrent
herpes simplex virus-1 infection induces hallmarks of neurodegeneration and cognitive
deficits in mice. PLoS Pathog 15:e1007617.

17.   Itzhaki RF. 2014. Herpes simplex virus type 1 and Alzheimer's disease: increasing evidence
for a major role of the virus. Front Aging Neurosci 6:202.

18.   Itzhaki RF, Lin WR, Shang D, Wilcock GK, Faragher B, Jamieson GA. 1997. Herpes simplex
virus type 1 in brain and risk of Alzheimer's disease. Lancet 349:241-4.

19.   Koelle DM, Magaret A, Warren T, Schellenberg GD, Wald A. 2010. APOE genotype is
associated with oral herpetic lesions but not genital or oral herpes simplex virus shedding. Sex
Transm Infect 86:202-6.

20.   Steel AJ, Eslick GD. 2015. Herpes Viruses Increase the Risk of Alzheimer's Disease: A Meta-
Analysis. J Alzheimers Dis 47:351-64.

21.   Readhead B, Haure-Mirande JV, Funk CC, Richards MA, Shannon P, Haroutunian V, Sano M,
Liang WS, Beckmann ND, Price ND, Reiman EM, Schadt EE, Ehrlich ME, Gandy S, Dudley
JT. 2018. Multiscale Analysis of Independent Alzheimer's Cohorts Finds Disruption of
Molecular, Genetic, and Clinical Networks by Human Herpesvirus. Neuron 99:64-82 e7.

22.   Lopatko Lindman K, Hemmingsson ES, Weidung B, Brannstrom J, Josefsson M, Olsson J, Elgh
F, Nordstrom P, Lovheim H. 2021. Herpesvirus infections, antiviral treatment, and the risk of
dementia-a registry-based cohort study in Sweden. Alzheimers Dement (N Y) 7:e12119.

23.   Honess RW, Roizman B. 1974. Regulation of herpesvirus macromolecular synthesis. I.
Cascade regulation of the synthesis of three groups of viral proteins. J Virol 14:8-19.

24.   Honess RW, Roizman B. 1975. Regulation of herpesvirus macromolecular synthesis:
sequential transition of polypeptide synthesis requires functional viral polypeptides. Proc Natl
Acad Sci U S A 72:1276-80.

25.   Kim JY, Mandarino A, Chao MV, Mohr I, Wilson AC. 2012. Transient reversal of episome
silencing precedes VP16-dependent transcription during reactivation of latent HSV-1 in
neurons. PLoS Pathog 8:e1002540.

26.   Lentine AF, Bachenheimer SL. 1990. Intracellular organization of herpes simplex virus type 1
DNA assayed by staphylococcal nuclease sensitivity. Virus Res 16:275-92.

27.   Leinbach SS, Summers WC. 1980. The structure of herpes simplex virus type 1 DNA as
probed by micrococcal nuclease digestion. J Gen Virol 51:45-59.

28.   Pignatti PF, Cassai E. 1980. Analysis of herpes simplex virus nucleoprotein complexes
extracted from infected cells. J Virol 36:816-28.

29.   Conn KL, Hendzel MJ, Schang LM. 2008. Linker histones are mobilized during infection with
herpes simplex virus type 1. J Virol 82:8629-46.

30.   Conn KL, Hendzel MJ, Schang LM. 2011. Core histones H2B and H4 are mobilized during
infection with herpes simplex virus 1. J Virol 85:13234-52.

31.   Monier K, Armas JC, Etteldorf S, Ghazal P, Sullivan KF. 2000. Annexation of the
interchromosomal space during viral infection. Nat Cell Biol 2:661-5.

32.   Herrera FJ, Triezenberg SJ. 2004. VP16-dependent association of chromatin-modifying
coactivators and underrepresentation of histones at immediate-early gene promoters during
herpes simplex virus infection. J Virol 78:9689-96.

33.   Kent JR, Zeng PY, Atanasiu D, Gardner J, Fraser NW, Berger SL. 2004. During lytic infection
herpes simplex virus type 1 is associated with histones bearing modifications that correlate
with active transcription. J Virol 78:10178-86.

34.   Lacasse JJ, Schang LM. 2010. During lytic infections, herpes simplex virus type 1 DNA is in
complexes with the properties of unstable nucleosomes. J Virol 84:1920-33.

35.   Kutluay SB, Doroghazi J, Roemer ME, Triezenberg SJ. 2008. Curcumin inhibits herpes
simplex virus immediate-early gene expression by a mechanism independent of p300/CBP
histone acetyltransferase activity. Virology 373:239-47.

36.   Kutluay SB, Triezenberg SJ. 2009. Regulation of histone deposition on the herpes simplex
virus type 1 genome during lytic infection. J Virol 83:5835-45.

37.   Oh J, Fraser NW. 2008. Temporal association of the herpes simplex virus genome with
histone proteins during a lytic infection. J Virol 82:3530-7.

38.   Cliffe AR, Knipe DM. 2008. Herpes simplex virus ICP0 promotes both histone removal and
acetylation on viral DNA during lytic infection. J Virol 82:12030-8.

39.   Wysocka J, Herr W. 2003. The herpes simplex virus VP16-induced complex: the makings of a
regulatory switch. Trends Biochem Sci 28:294-304.

40.   Fan D, Wang M, Cheng A, Jia R, Yang Q, Wu Y, Zhu D, Zhao X, Chen S, Liu M, Zhang S, Ou
X, Mao S, Gao Q, Sun D, Wen X, Liu Y, Yu Y, Zhang L, Tian B, Pan L, Chen X. 2020. The
Role of VP16 in the Life Cycle of Alphaherpesviruses. Front Microbiol 11:1910.

41.   Liang Y, Vogel JL, Arbuckle JH, Rai G, Jadhav A, Simeonov A, Maloney DJ, Kristie TM. 2013.
Targeting the JMJD2 histone demethylases to epigenetically control herpesvirus infection and
reactivation from latency. Sci Transl Med 5:167ra5.

42.   Liang Y, Vogel JL, Narayanan A, Peng H, Kristie TM. 2009. Inhibition of the histone
demethylase LSD1 blocks alpha-herpesvirus lytic replication and reactivation from latency.
Nat Med 15:1312-7.

43.   Narayanan A, Ruyechan WT, Kristie TM. 2007. The coactivator host cell factor-1 mediates
Set1 and MLL1 H3K4 trimethylation at herpesvirus immediate early promoters for initiation
of infection. Proc Natl Acad Sci U S A 104:10835-40.

44.   Wysocka J, Myers MP, Laherty CD, Eisenman RN, Herr W. 2003. Human Sin3 deacetylase
and trithorax-related Set1/Ash2 histone H3-K4 methyltransferase are tethered together
selectively by the cell-proliferation factor HCF-1. Genes Dev 17:896-911.

45.   Dremel SE, DeLuca NA. 2019. Herpes simplex viral nucleoprotein creates a competitive
transcriptional environment facilitating robust viral transcription and host shut off. Elife 8.

46.   Stevens JG, Wagner EK, Devi-Rao GB, Cook ML, Feldman LT. 1987. RNA complementary to
a herpesvirus alpha gene mRNA is prominent in latently infected neurons. Science 235:1056-
9.

47.   Deatly AM, Spivack JG, Lavi E, Fraser NW. 1987. RNA from an immediate early region of the
type 1 herpes simplex virus genome is present in the trigeminal ganglia of latently infected
mice. Proc Natl Acad Sci U S A 84:3204-8.

48. Umbach JL, Kramer MF, Jurak I, Karnowski HW, Coen DM, Cullen BR. 2008. MicroRNAs
expressed by herpes simplex virus 1 during latent infection regulate viral mRNAs. Nature
454:780-3.
49. Wang QY, Zhou C, Johnson KE, Colgrove RC, Coen DM, Knipe DM. 2005. Herpesviral
latency-associated transcript gene promotes assembly of heterochromatin on viral lytic-gene
promoters in latent infection. Proc Natl Acad Sci U S A 102:16055-9.
50. Cliffe AR, Coen DM, Knipe DM. 2013. Kinetics of facultative heterochromatin and polycomb
group protein association with the herpes simplex viral genome during establishment of
latent infection. mBio 4.
51. Cliffe AR, Garber DA, Knipe DM. 2009. Transcription of the herpes simplex virus latency-
associated transcript promotes the formation of facultative heterochromatin on lytic
promoters. J Virol 83:8182-90.
52. Knipe DM, Cliffe A. 2008. Chromatin control of herpes simplex virus lytic and latent
infection. Nat Rev Microbiol 6:211-21.
53. Kwiatkowski DL, Thompson HW, Bloom DC. 2009. The polycomb group protein Bmi1 binds
to the herpes simplex virus 1 latent genome and maintains repressive histone marks during
latency. J Virol 83:8173-81.
54. Dochnal SA, Francois AK, Cliffe AR. 2021. De Novo Polycomb Recruitment: Lessons from
Latent Herpesviruses. Viruses 13.
55. Hu HL, Srinivas KP, Wang S, Chao MV, Lionnet T, Mohr I, Wilson AC, Depledge DP, Huang
TT. 2022. Single-cell transcriptomics identifies Gadd45b as a regulator of herpesvirus-
reactivating neurons. EMBO Rep 23:e53543.
56. Cliffe AR, Arbuckle JH, Vogel JL, Geden MJ, Rothbart SB, Cusack CL, Strahl BD, Kristie TM,
Deshmukh M. 2015. Neuronal Stress Pathway Mediating a Histone Methyl/Phospho Switch Is
Required for Herpes Simplex Virus Reactivation. Cell Host Microbe 18:649-58.
57. Cliffe AR, Wilson AC. 2017. Restarting Lytic Gene Transcription at the Onset of Herpes
Simplex Virus Reactivation. J Virol 91.
58. Cuddy SR, Schinlever AR, Dochnal S, Seegren PV, Suzich J, Kundu P, Downs TK, Farah M,
Desai BN, Boutell C, Cliffe AR. 2020. Neuronal hyperexcitability is a DLK-dependent trigger
of herpes simplex virus reactivation that can be induced by IL-1. Elife 9.
59. Steiner I, Spivack JG, Deshmane SL, Ace CI, Preston CM, Fraser NW. 1990. A herpes simplex
virus type 1 mutant containing a nontransinducing Vmw65 protein establishes latent
infection in vivo in the absence of viral replication and reactivates efficiently from explanted
trigeminal ganglia. J Virol 64:1630-8.
60. Wilson AC, Mohr I. 2012. A cultured affair: HSV latency and reactivation in neurons. Trends
Microbiol 20:604-11.
61. Lachmann R. 2003. Herpes simplex virus latency. Expert Rev Mol Med 5:1-14.
62. Pourchet A, Modrek AS, Placantonakis DG, Mohr I, Wilson AC. 2017. Modeling HSV-1
Latency in Human Embryonic Stem Cell-Derived Neurons. Pathogens 6.
63. Suzich JB, Cliffe AR. 2018. Strength in diversity: Understanding the pathways to herpes
simplex virus reactivation. Virology 522:81-91.
64. Suzich JB, Cuddy SR, Baidas H, Dochnal S, Ke E, Schinlever AR, Babnis A, Boutell C, Cliffe
AR. 2021. PML-NB-dependent type I interferon memory results in a restricted form of HSV
latency. EMBO Rep 22:e52547.

65.   Linderman JA, Kobayashi M, Rayannavar V, Fak JJ, Darnell RB, Chao MV, Wilson AC, Mohr
983   I. 2017. Immune Escape via a Transient Gene Expression Program Enables Productive
984   Replication of a Latent Pathogen. Cell Rep 18:1312-1323.
985   66.   Bertke AS, Swanson SM, Chen J, Imai Y, Kinchington PR, Margolis TP. 2011. A5-positive
986   primary sensory neurons are nonpermissive for productive infection with herpes simplex
987   virus 1 in vitro. J Virol 85:6669-77.
988   67.   Cabrera JR, Charron AJ, Leib DA. 2018. Neuronal Subtype Determines Herpes Simplex Virus
989   1 Latency-Associated-Transcript Promoter Activity during Latency. J Virol 92.
990   68.   Arthur JL, Scarpini CG, Connor V, Lachmann RH, Tolkovsky AM, Efstathiou S. 2001. Herpes
991   simplex virus type 1 promoter activity during latency establishment, maintenance, and
992   reactivation in primary dorsal root neurons in vitro. J Virol 75:3885-95.
993   69.   Camarena V, Kobayashi M, Kim JY, Roehm P, Perez R, Gardner J, Wilson AC, Mohr I, Chao
994   MV. 2010. Nature and duration of growth factor signaling through receptor tyrosine kinases
995   regulates HSV-1 latency in neurons. Cell Host Microbe 8:320-30.
996   70.   Yanez AA, Harrell T, Sriranganathan HJ, Ives AM, Bertke AS. 2017. Neurotrophic Factors
997   NGF, GDNF and NTN Selectively Modulate HSV1 and HSV2 Lytic Infection and Reactivation
998   in Primary Adult Sensory and Autonomic Neurons. Pathogens 6.
999   71.   Wilcox CL, Johnson EM, Jr. 1988. Characterization of nerve growth factor-dependent herpes
1000   simplex virus latency in neurons in vitro. J Virol 62:393-9.
1001   72.   Wilcox CL, Smith RL, Freed CR, Johnson EM, Jr. 1990. Nerve growth factor-dependence of
1002   herpes simplex virus latency in peripheral sympathetic and sensory neurons in vitro. J
1003   Neurosci 10:1268-75.
1004   73.   Thellman NM, Botting C, Madaj Z, Triezenberg SJ. 2017. An Immortalized Human Dorsal
1005   Root Ganglion Cell Line Provides a Novel Context To Study Herpes Simplex Virus 1 Latency
1006   and Reactivation. J Virol 91.
1007   74.   Edwards TG, Bloom DC. 2019. Lund Human Mesencephalic (LUHMES) Neuronal Cell Line
1008   Supports Herpes Simplex Virus 1 Latency In Vitro. J Virol 93.
1009   75.   De Clercq E. 2008. The discovery of antiviral agents: ten different compounds, ten different
1010   stories. Med Res Rev 28:929-53.
1011   76.   De Clercq E, Holy A. 2005. Acyclic nucleoside phosphonates: a key class of antiviral drugs.
1012   Nat Rev Drug Discov 4:928-40.
1013   77.   Morfin F, Thouvenot D. 2003. Herpes simplex virus resistance to antiviral drugs. J Clin Virol
1014   26:29-37.
1015   78.   Forrester A, Farrell H, Wilkinson G, Kaye J, Davis-Poynter N, Minson T. 1992. Construction
1016   and properties of a mutant of herpes simplex virus type 1 with glycoprotein H coding
1017   sequences deleted. J Virol 66:341-8.
1018   79.   Benboudjema L, Mulvey M, Gao Y, Pimplikar SW, Mohr I. 2003. Association of the herpes
1019   simplex virus type 1 Us11 gene product with the cellular kinesin light-chain-related protein
1020   PAT1 results in the redistribution of both polypeptides. J Virol 77:9192-203.
1021   80.   Hilterbrand AT, Heldwein EE. 2019. Go go gadget glycoprotein!: HSV-1 draws on its sizeable
1022   glycoprotein tool kit to customize its diverse entry routes. PLoS Pathog 15:e1007660.
1023   81.   Turner A, Bruun B, Minson T, Browne H. 1998. Glycoproteins gB, gD, and gHgL of herpes
1024   simplex virus type 1 are necessary and sufficient to mediate membrane fusion in a Cos cell
1025   transfection system. J Virol 72:873-5.

1026	82.	Hu HL, Shiflett LA, Kobayashi M, Chao MV, Wilson AC, Mohr I, Huang TT. 2019.
1027		TOP2beta-Dependent Nuclear DNA Damage Shapes Extracellular Growth Factor Responses
1028		via Dynamic AKT Phosphorylation to Control Virus Latency. Mol Cell 74:466-480 e4.
1029	83.	Kobayashi M, Kim JY, Camarena V, Roehm PC, Chao MV, Wilson AC, Mohr I. 2012. A
1030		primary neuron culture system for the study of herpes simplex virus latency and reactivation.
1031		J Vis Exp doi:10.3791/3823.
1032	84.	Chao MV. 2003. Neurotrophins and their receptors: a convergence point for many signalling
1033		pathways. Nat Rev Neurosci 4:299-309.
1034	85.	van Zeijl M, Fairhurst J, Jones TR, Vernon SK, Morin J, LaRocque J, Feld B, O'Hara B, Bloom
1035		JD, Johann SV. 2000. Novel class of thiourea compounds that inhibit herpes simplex virus
1036		type 1 DNA cleavage and encapsidation: resistance maps to the UL6 gene. J Virol 74:9054-61.
1037	86.	Newcomb WW, Brown JC. 2002. Inhibition of herpes simplex virus replication by WAY-
1038		150138: assembly of capsids depleted of the portal and terminase proteins involved in DNA
1039		encapsidation. J Virol 76:10084-8.
1040	87.	Pesola JM, Zhu J, Knipe DM, Coen DM. 2005. Herpes simplex virus 1 immediate-early and
1041		early gene expression during reactivation from latency under conditions that prevent
1042		infectious virus production. J Virol 79:14516-25.
1043	88.	Hollville E, Romero SE, Deshmukh M. 2019. Apoptotic cell death regulation in neurons.
1044		FEBS J 286:3276-3298.
1045	89.	Annis RP, Swahari V, Nakamura A, Xie AX, Hammond SM, Deshmukh M. 2016. Mature
1046		neurons dynamically restrict apoptosis via redundant premitochondrial brakes. FEBS J
1047		283:4569-4582.
1048	90.	Kole AJ, Swahari V, Hammond SM, Deshmukh M. 2011. miR-29b is activated during
1049		neuronal maturation and targets BH3-only genes to restrict apoptosis. Genes Dev 25:125-30.
1050	91.	Smith RL, Pizer LI, Johnson EM, Jr., Wilcox CL. 1992. Activation of second-messenger
1051		pathways reactivates latent herpes simplex virus in neuronal cultures. Virology 188:311-8.
1052	92.	Colgin MA, Smith RL, Wilcox CL. 2001. Inducible cyclic AMP early repressor produces
1053		reactivation of latent herpes simplex virus type 1 in neurons in vitro. J Virol 75:2912-20.
1054	93.	De Regge N, Van Opdenbosch N, Nauwynck HJ, Efstathiou S, Favoreel HW. 2010. Interferon
1055		alpha induces establishment of alphaherpesvirus latency in sensory neurons in vitro. PLoS
1056		One 5.
1057	94.	Danaher RJ, Jacob RJ, Miller CS. 2003. Herpesvirus quiescence in neuronal cells. V: forskolin-
1058		responsiveness of the herpes simplex virus type 1 alpha0 promoter and contribution of the
1059		putative cAMP response element. J Neurovirol 9:489-97.
1060	95.	Moriya A, Yoshiki A, Kita M, Fushiki S, Imanishi J. 1994. Heat shock-induced reactivation of
1061		herpes simplex virus type 1 in latently infected mouse trigeminal ganglion cells in dissociated
1062		culture. Arch Virol 135:419-25.
1063	96.	Halford WP, Gebhardt BM, Carr DJ. 1996. Mechanisms of herpes simplex virus type 1
1064		reactivation. J Virol 70:5051-60.
1065	97.	Halford WP, Schaffer PA. 2001. ICP0 is required for efficient reactivation of herpes simplex
1066		virus type 1 from neuronal latency. J Virol 75:3240-9.
1067	98.	Kushnir AS, Davido DJ, Schaffer PA. 2010. Role of nuclear factor Y in stress-induced
1068		activation of the herpes simplex virus type 1 ICP0 promoter. J Virol 84:188-200.

99.   Sawtell NM, Thompson RL. 1992. Rapid in vivo reactivation of herpes simplex virus in
latently infected murine ganglionic neurons after transient hyperthermia. J Virol 66:2150-6.
100. Perng GC, Osorio N, Jiang X, Geertsema R, Hsiang C, Brown D, BenMohamed L, Wechsler
SL. 2016. Large Amounts of Reactivated Virus in Tears Precedes Recurrent Herpes Stromal
Keratitis in Stressed Rabbits Latently Infected with Herpes Simplex Virus. Curr Eye Res
41:284-91.
101. Shimomura Y, Higaki S, Watanabe K. 2010. Suppression of herpes simplex virus 1
reactivation in a mouse eye model by cyclooxygenase inhibitor, heat shock protein inhibitor,
and adenosine monophosphate. Jpn J Ophthalmol 54:187-90.
102. Hill JM, Lukiw WJ, Gebhardt BM, Higaki S, Loutsch JM, Myles ME, Thompson HW, Kwon
BS, Bazan NG, Kaufman HE. 2001. Gene expression analyzed by microarrays in HSV-1 latent
mouse trigeminal ganglion following heat stress. Virus Genes 23:273-80.
103. Sawtell NM, Thompson RL. 2004. Comparison of herpes simplex virus reactivation in ganglia
in vivo and in explants demonstrates quantitative and qualitative differences. J Virol 78:7784-
94.
104. Thompson RL, Preston CM, Sawtell NM. 2009. De novo synthesis of VP16 coordinates the
exit from HSV latency in vivo. PLoS Pathog 5:e1000352.
105. Patel S, Cohen F, Dean BJ, De La Torre K, Deshmukh G, Estrada AA, Ghosh AS, Gibbons P,
Gustafson A, Huestis MP, Le Pichon CE, Lin H, Liu W, Liu X, Liu Y, Ly CQ, Lyssikatos JP, Ma
C, Scearce-Levie K, Shin YG, Solanoy H, Stark KL, Wang J, Wang B, Zhao X, Lewcock JW,
Siu M. 2015. Discovery of dual leucine zipper kinase (DLK, MAP3K12) inhibitors with
activity in neurodegeneration models. J Med Chem 58:401-18.
106. Heinemann B, Nielsen JM, Hudlebusch HR, Lees MJ, Larsen DV, Boesen T, Labelle M,
Gerlach LO, Birk P, Helin K. 2014. Inhibition of demethylases by GSK-J1/J4. Nature 514:E1-
2.
107. Sawtell NM, Thompson RL. 2016. Herpes simplex virus and the lexicon of latency and
reactivation: a call for defining terms and building an integrated collective framework.
F1000Res 5.
108. Wang L, Brown JL, Cao R, Zhang Y, Kassis JA, Jones RS. 2004. Hierarchical recruitment of
polycomb group silencing complexes. Mol Cell 14:637-46.
109. Chadwick BP, Willard HF. 2003. Barring gene expression after XIST: maintaining facultative
heterochromatin on the inactive X. Semin Cell Dev Biol 14:359-67.
110. Du T, Zhou G, Roizman B. 2011. HSV-1 gene expression from reactivated ganglia is
disordered and concurrent with suppression of latency-associated transcript and miRNAs.
Proc Natl Acad Sci U S A 108:18820-4.
111. Tedeschi A, Bradke F. 2013. The DLK signalling pathway--a double-edged sword in neural
development and regeneration. EMBO Rep 14:605-14.
112. Karney-Grobe S, Russo A, Frey E, Milbrandt J, DiAntonio A. 2018. HSP90 is a chaperone for
DLK and is required for axon injury signaling. Proc Natl Acad Sci U S A 115:E9899-E9908.
113. Summers DW, Frey E, Walker LJ, Milbrandt J, DiAntonio A. 2020. DLK Activation
Synergizes with Mitochondrial Dysfunction to Downregulate Axon Survival Factors and
Promote SARM1-Dependent Axon Degeneration. Mol Neurobiol 57:1146-1158.

114. Hirai S, Kawaguchi A, Suenaga J, Ono M, Cui DF, Ohno S. 2005. Expression of
MUK/DLK/ZPK, an activator of the JNK pathway, in the nervous systems of the developing
mouse embryo. Gene Expr Patterns 5:517-23.
115. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res 27:722-736.
116. Vaser R, Sovic I, Nagarajan N, Sikic M. 2017. Fast and accurate de novo genome assembly
from long uncorrected reads. Genome Res 27:737-746.
117. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and
SAMtools. Bioinformatics 25:2078-9.
118. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26:841-2.
119. Hahne F, Ivanek R. 2016. Visualizing Genomic Data Using Gviz and Bioconductor. Methods
Mol Biol 1418:335-51.
120. Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey
VJ. 2013. Software for computing and annotating genomic ranges. PLoS Comput Biol
9:e1003118.
121. Malin SA, Davis BM, Molliver DC. 2007. Production of dissociated sensory neuron cultures
and considerations for their use in studying neuronal function and plasticity. Nat Protoc
2:152-60.
122. Burman LG, Mauro VP. 2012. Analysis of rRNA processing and translation in mammalian
cells using a synthetic 18S rRNA expression system. Nucleic Acids Res 40:8085-98.
1134