1 Lower limb in sprinters with larger relative mass but not larger
2 normalized moment of inertia: novel insights into human plasticity
3 Running title: Lower limb inertia properties in sprinters
4

Natsuki Sado[1, 2*], Hoshizora Ichinose[3], Yasuo Kawakami[2]
[1] Faculty of Health and Sports Science, University of Tsukuba, Tsukuba, Japan
[2] Faculty of Sports Sciences, Waseda University, Tokorozawa, Japan
[3] Graduate School of Sport Sciences, Waseda University, Tokorozawa, Japan

Corresponding author:

Natsuki Sado
Faculty of Health and Sport Sciences, University of Tsukuba
1-1-1 Tennodai, Tsukuba,
Ibaraki 305-8574 Japan.
Phone: +81-29-853-2611
E-mail: sado.natsuki.gm@u.tsukuba.ac.jp

ORCiDs:

Natsuki Sado: 0000-0002-3958-8193
Yasuo Kawakami: 0000-0003-0588-4039
19

ABSTRACT

Inhomogeneous muscle hypertrophy in response to daily motor executions is a

morphological adaptation in humans. Inhomogeneous morphological adaptation would change

the mass distribution, which in turn may affect the mechanical difficulty in moving from an

inertia perspective; however, the morphological adaptation from the inertia perspective has not

been examined. Here we show no corresponding differences in the normalized mass and

normalized moment of inertia between the sprinters and the untrained non-sprinters. We

analyzed the fat- and water-separated magnetic resonance images from the lower limbs of 11

male sprinters (100 m best time of 10.44–10.83 s) and 12 untrained non-sprinters. The

lower-limb relative mass was significantly larger in sprinters (18.7±0.7% body mass) than in

non-sprinters (17.6±0.6% body mass), while the normalized moment of inertia of the lower

limb around the hip in the anatomical position was similar between them (0.044±0.002 vs.

0.042±0.002 [a. u.]). The thigh relative mass in sprinters (12.9±0.4% body mass) was

significantly larger than that in non-sprinters (11.9±0.4% body mass), whereas the shank and

foot relative masses were similar. We revealed that the adaptation to sprinting in humans does

not relatively increase the mechanical difficulty to swing the lower limb in terms of inertia,

even though the lower-limb mass increases, reflecting muscular hypertrophy. This is suggested

to be one of the significant advantages of human morphology for adaptability to explosive

motor tasks. (224 words)

NEW & NOTEWORTHY

We compared the lower-limb inertia properties between male sprinters and untrained

non-sprinters by using the fat- and water-separated magnetic resonance images. Sprinters had

lower limbs with a larger relative mass but not a larger normalized moment of inertia than

non-sprinters. We suggest one of the practical advantages of human morphological

characteristics for their adaptation to explosive motor tasks, which provides a novel

perspective to the literature for understanding human morphological adaptability. (71 words)

KEYWORDS:

Morphology; Adaptation; Magnetic resonance imaging; Dixon; Hypertrophy

ABBREVIATIONS:

BSIP: body segment inertia parameter.

MRI: magnetic resonance imaging

3D: thee-dimensional

$\hat{I}_{LowerLimb}^{Hip}$:   normalized lower-limb moment of inertia around the hip joint

INTRODUCTION

Human morphological characteristics are mainly the result of adaptation to bipedal locomotion (1, 2). Human morphology enables bipedal walking and endurance running efficiency, resulting in current humans being good endurance runners but poor sprinters (2). Alternately, evolution in humans has left a great plasticity, allowing them to become faster sprinters via training; for example, the Japan National Record of 50 m sprint is 5.75 s while the average time of 50-m sprint in Japanese males is approximately 7.37 ± 0.52 s (3).

Adaptability in humans has been approached from the muscle morphologies in athletes, based on the exertion abilities of the joint torque (4) and power (5) is primarily determined by the muscle size, especially muscle volume (4, 6). Previous MRI studies on sprinters have demonstrated hypertrophy of the trunk and lower-limb muscles (7–9). The hypertrophy was shown to be inhomogeneous with the specific development of the hip flexors and extensors (7–9). The muscle sizes of the hip extensors and flexors are related to the running velocity and/or 100 m sprint time in sub-elite (7, 8, 10) and elite sprinters (9). It has also been reported that the hip extensors' volumes discriminate elite sprinters from the sub-elite (9). The muscularity in sprinters reflects the musculoskeletal demands that increasing running velocity in the high-speed range (>7 m/s) requires like the larger torque/power exertion of hip flexors and extensors (11, 12).

From a mechanical perspective, any motion ($\boldsymbol{a}$:acceleration, $\boldsymbol{\alpha}$:angular acceleration,) is determined by the combination of force-related ($\boldsymbol{f}$: forces, $\boldsymbol{\tau}$: torques) and inertia-related (m: mass, $\boldsymbol{I}$: moment of inertia) factors in line with Newton-Euler equations of motion:

$$m\boldsymbol{a} = \sum \boldsymbol{f}, \tag{1}$$

$$\boldsymbol{I}\boldsymbol{\alpha} = \sum \boldsymbol{\tau}. \tag{2}$$

As humans control their motion via joint torques induced by muscle forces, the relationship between the torque and moment of inertia (Eq. 2) is particularly important for human motor execution. The above muscular characteristics in sprinters (7–10) can be interpreted as an

adaptation to force-related factors, but simultaneously, the morphological changes would

involve changes in the mass distribution and thereby inertia-properties. As lean tissue density

is larger than that of fat tissue (13, 14), the muscular hypertrophy would involve an increase

in the lower-limb mass, leading to a general speculation of the trade-off between greater

torque exertion ability and greater difficulty in moving the lower limb (i.e., moment of inertia).

However, as discussed above, the muscle hypertrophy in sprinters (7–9) is inhomogeneous.

This leads to our hypothesis that the mechanical difficulties in swinging the lower limb and

the moment of inertia around the hip might not correspond to an increase in the lower-limb

mass in sprinters. Anatomists have previously developed scaling coefficients to calculate body

segment inertia parameters (BSIPs) for motion analysis. This is done using the direct

measurements of elderly cadavers (15) and indirect measurements *in vivo* through medical

images or surface scans such as gamma-ray scanning (16) computed tomography (17),

magnetic resonance imaging (MRI) (18, 19) and three-dimensional (3D) laser scanners (20).

However, the effects of adaptation to motor experience on inertial properties have not been

fully examined.

Based on the simple mechanical law shown in Eq. 2), an increase in the moment of

inertia leads to (□) increasing musculoskeletal mechanical loads if the motion is consistent,

and (□) decreasing angular acceleration if the joint torque exertions are consistent.

Understanding the morphological adaptation from an inertial perspective would provide

useful knowledge for human motor performance development from both the injury prevention

and performance improvement perspectives. In this study, we examined the inertia properties

of sprinters by analyzing water- and fat-separated MRIs. Among the inertial properties, we

focused on the normalized lower-limb moment of inertia around the hip joint ($\hat{I}_{LowerLimb}^{Hip}$)

because of the high musculoskeletal demands in the lower limb swing during sprinting (11,

12).

MATERIALS AND METHODS

 Experimental design

We performed *a priori* power analysis for an independent *t*-test with the parameters of

type ☐ error; α=0.05, statistical power: 1-β=0.8, and effect size: Cohen's *d*=1.2 'very large'

(21), which calculated the total sample size (*n*) as 24 (*n*=12 in each group). According to the

power analysis, we recruited a total of 24 healthy males, including 12 well-trained (athletic

career >7 years and 100 m best records: 10.44–10.83 s) sprinters and 12 non-sprinters who did

not undertake resistance training or sports activities for at least 2 years. However, the data of

one of the sprinters could not be analyzed because of an MRI error (fat/water separation error in

the Dixon sequence) around the endpoint of the toe in the foot segment scan, and the main

outcome data (i.e., the lower-limb mass and the moment of inertia around the hip joint) could

not be calculated. Thus, we excluded this participant's data from the analysis. The analyzed

participants were 11 sprinters (age: 20±2 yrs, height: 1.77±0.06 m; body mass: 68.2±4.4 kg,

Mean±SD) and 12 non-sprinters (age: 23±3 yrs, height: 1.69±0.04 m; body mass: 58.6±4.5 kg).

The purpose and experimental protocol of the study were explained to the participants and they

provided written informed consent. The Human Research Ethics Committee at Waseda

University, Japan approved the study protocol (reference number: 2019-174).

We used a 3.0 Tesla MR scanner (SIGNA Premier, GE Healthcare, Milwaukee, USA)

to obtain the MR scans. All lower limb segment scans were performed in the supine position.

The 2-point 3D-dixon axial sequence, called "Lava-Flex" in GE Healthcare, was performed on

the right thigh, shank, and foot segments ('SpacingBetweenSlices': 2 mm, 'Pixel-Spacing':

[0.2930 0.2930] mm, 'RepetitionTime': 4.3 ms, 'EchoTime': 1.7 ms, Width×Height:

1024×1024 pixels). Each scan of the thigh and shank had two sections because of the limitation

in the length scanned in one section. The two sections of each thigh and the shank scans were

taken continuously without changing the position on the MRI machine. In the MRI settings, the

anteroposterior and mediolateral coordinates of the two consecutive scans coincided. To

minimize the effect of magnetic field inhomogeneity on the Dixon sequence, the superoinferior

ranges of the two scans overlapped by > 10 cm.

MRI processing

We analyzed the 16-bit depth MRI images using MATLAB 2019a (MathWorks,

Natick, MA, USA). A flowchart of the segmentation procedure is presented in Fig. 1. To create

a temporal body mask, we first created fat-only and water-only temporal binary images using a

low intensity threshold. These binary images were added together, and the temporal body mask

was defined by filling any internal holes in the added masks based on a binary morphology

operation (22). After creating the temporal body mask, we removed the pixel intensity in the

unwanted areas (other than the range of interest, such as the part of the opposite leg segment

included in the image). After that, we estimated and then homogenized the inhomogeneity field

using a grayscale morphological closing operation with a large circular structure element (22).

The threshold for the foreground and background signal intensity of each of the fat-only and

water-only images was determined using Otsu's method (23), which created water-binary and

fat-binary images. We created a definitive body mask by adding the two masks together and

filling any internal holes with a binary morphology operation (22). We classified each pixel as a

fat tissue, lean (muscle or skin) tissue, or a background using water- and fat-binary images. For

a slice having some voxels involved in both water- and fat-binary images, we further performed

k-means clustering to classify the intensity of the voxels in the sum of the fat-only and

water-only images (i.e., in-phase image) into three categories (fat tissue, lean tissue, or

background), which were applied to the voxels involved in both the water-binary and fat-binary

images. The trabecular bone has MRI intensity characteristics similar to fat tissue because of

the yellow marrow (24), while the cortical bone is not brightened by MRI. These bone tissues

were classified as follows:

Step 1.   Search for a region inside the body mask region where the pixels classified

154   as fat tissue (trabecular bone in fact) are surrounded by pixels classified as

155   background (cortical bone in fact) and define them as trabecular bone and

156   cortical bone.

Step 2.   (When a bone cannot be explored in Step 1 and the trabecular bone is an

independent object in the fat-binary image), select and define the independent

object in the fat-only image as a trabecular bone.

Step 3.   (When a trabecular bone was in contact with the outer fat tissue in the

fat-binary image and could not be defined in Steps 1 and 2) Manually draw a

border with the fat in the fat-only image or in-phase image using a pen tablet

(Artist 15.6 Pro, XP-PEN, Japan) and define the object as a trabecular bone.

Step 4.   (After Steps 2 and 3) Define the pixels that surround the trabecular bone

and are classified as background (cortical bone in fact) as a cortical bone.

Step 5.   Adjust the cortical bone if necessary (for example, if the tissue adjacent to

the cortical bone that is not depressed on MRI is considered to be a cortical

bone).

Data calculation

The 3D coordinates of the anatomical landmarks (Table 1) on the MR coordinate

system were derived via manual digitizing in the 3D mode using the OsiriX software version 13

(Pixmeo, Geneva, Switzerland) by an examiner. The examiner repeated the digitization of all

landmarks from eight participants for whom it had been >1 month following the first

digitizing process to assess the intra-examiner repeatability of manual digitizing.

The ankle joint in the foot and shank scans and the knee joint in the shank and thigh

scans were defined as the midpoints between the malleoli and the lateral and medial articular

cleft of the knee, respectively. The hip joint was defined using a least-squares calculation for a

sphere fitting of 30 points distributed over the surface of the femoral head, similar to Harrington

et al. (25). The ankle and knee boundary planes were defined as the horizontal planes passing

those joint locations (Fig. 2A). The hip joint boundary was defined by a plane 37º (26, 27)

vertical on the medial to the hip joint and a horizontal plane lateral to the hip joint (28) (Fig. 2A).

The lean and fat voxels outside of the border were excluded. The cortical and trabecular bone

voxel in that segment was retained even though it was outside the proximal and distal borders.

Bone voxels within the segment were retained even if they were outside the boundary, whereas

bone voxels in another segment were excluded even if they were inside the boundary.

Based on the literature values (13, 14), we defined the tissue density as:

$$\rho_k = [0.902\ 1.067\ 1.705\ 1.100], \tag{4}$$

187 where $k=1,2,3,4$ shows the tissues of fat, lean, cortical bone, and trabecular bone,

188 respectively. The mass, m_s, volume V_s, density ρ_s, and position coordinates of the center of

189 mass, $\begin{bmatrix} x_{s,\text{CoM}} \\ y_{s,\text{CoM}} \\ z_{s,\text{CoM}} \end{bmatrix}$, of a segment s were calculated as

$$m_s = \sum_{k=1}^{4} \sum_{i=1}^{n_k} [(w^2 h)\rho_k] \tag{5}$$

$$V_s = \sum_{k=1}^{4} [(w^2 h)n_k] \tag{6}$$

$$\rho_s = \frac{m_s}{V_s} \tag{7}$$

$$\begin{bmatrix} x_{s,\text{CoM}} \\ y_{s,\text{CoM}} \\ z_{s,\text{CoM}} \end{bmatrix} = \frac{1}{m_s} \sum_{k=1}^{4} \sum_{i=1}^{n_k} \left[(w^2 h)\rho_k \begin{bmatrix} x_i \\ y_i \\ z_i \end{bmatrix} \right] \tag{8}$$

190 where n_k is the number of voxels classified as the tissue k, w and h were the width

191 ('Pixel-Spacing': 0.2930 mm) and height ('Spacing-Between-Slices': 2.0000 mm) of each

192 voxel (i.e., $w^2 h$ is the volume of each voxel), and $\begin{bmatrix} x_i \\ y_i \\ z_i \end{bmatrix}$ is the position coordinate of the ith

193 voxel, respectively.

194 Inertia tensor of segment s, $\boldsymbol{I}_s^{\text{GCS}}$, was calculated as shown below:

$$\boldsymbol{I}_s^{\text{GCS}} = \begin{bmatrix} \sum_{i=1}^{n} \left[I_{x_i} + m_i(y_i^2 + z_i^2) \right] & -\sum_{i=1}^{n}(m_i x_i y_i) & -\sum_{i=1}^{n}(m_i z_i x_i) \\ -\sum_{i=1}^{n}(m_i x_i y_i) & \sum_{i=1}^{n} \left[I_{y_i} + m_i(z_i^2 + x_i^2) \right] & -\sum_{i=1}^{n}(m_i y_i z_i) \\ -\sum_{i=1}^{n}(m_i z_i x_i) & -\sum_{i=1}^{n}(m_i y_i z_i) & \sum_{i=1}^{n} \left[I_{z_i} + m_i(x_i^2 + y_i^2) \right] \end{bmatrix} \tag{9}$$

195 where m_i is the mass of ith voxel, I_{x_i} I_{y_i} and I_{z_i} were the moments of inertia of each

196 voxel and were calculated as follows.

$$I_{x_i} = I_{y_i} = \frac{m_i(w^2 + h^2)}{12} \text{ and } I_{z_i} = \frac{m_i(2w^2)}{12} \tag{10}$$

197   A right-handed orthogonal segment coordinate system (SCS) was fixed at each of the

198   thigh, shank, and foot segments using the 3D coordinates of landmarks. This definition was

199   consistent with that used by Sado et al. (29, 30). We calculated the inertia tensor in each SCS

200   (I_s^{SCS}) as follows:

$$I_s^{SCS} = R_s{}'I_s^{GCS}R_s \tag{11}$$

where R_s is the rotational transformation matrix from the SCS to the GCS of segment s.

Note that the principal axes of the segments are not consistent with the axes of the SCSs (31);

thus, I_s^{SCS} is not in the form of a diagonal matrix, which is similar to Dumas et al. (31),

considering the differences between the SCS axes and the inertial principal axes.

We hypothetically created the lower-limb data in an anatomical position from the 3D

coordinates in the MRI coordinate system of the thigh, shank, and foot using hierarchically

constructed coordinates with the hip as the origin (Fig. 2B). In this hypothetical creation, the

x-axis directions of their SCSs were consistent with each other. We then calculated

$I_{LowerLimb}^{Hip}$ in the data of this hypothetical anatomical position as

$$I_{Lowerlimb}^{Hip} = \sum_{s=1}^{3} \left\{ I_s^{SCS}{}_{xx} + m_s \left(\left| y_s - y_{Hip} \right|^2 + \left| z_s - z_{Hip} \right|^2 \right) \right\} \tag{12}$$

where $I_s^{SCS}{}_{xx}$ is the moment of inertia of the segment around its CoM, y_s, y_{Hip}, z_s and

z_{Hip} are the anteroposterior or superoinferior coordinates of the CoM of segment s or the

hip joint in a hypothetically created coordinate. In this coordinate system, the origin is set at

the hip joint; thus, y_{Hip}, and z_{Hip} are equal to zero.

The segment and the lower-limb masses were expressed relative to the whole-body

mass. Both the products and the moments of inertia of each segment were expressed relative

to the segment length l_s and the segment mass m_s. For example, the moment of inertia of

segment s around the x-axis is

$$r_{s_{xx}} = \frac{\sqrt{\dfrac{I_{s}^{SCS}{}_{xx}}{m_s}}}{l_s} \tag{13}$$

218 The segment lengths were defined as the Euclidean distances between the proximal and distal

219 joint locations (Fig. 2B). Using the scaling method of gait data established by Hof (32), we

220 normalized $I_{LowerLimb}^{Hip}$ by the lower-limb length ($l_{Lowerlimb}$) and the whole-body mass

221 (m_{body}) as:

$$\hat{I}_{LowerLimb}^{Hip} = \frac{I_{LowerLimb}^{Hip}}{m_{body} l_{Lowerlimb}^{2}} \tag{14}$$

where $l_{Lowerlimb}$ is the height difference between the hip joint center and calcaneus in the

hypothetically created coordinates (Fig.2B).

Statistical analysis

We used MATLAB 2019a for statistical analysis. We used ICCs and CVs to assess

the repeatability of the examiners' manual-digitizing-basis segment lengths. The ICCs can be

interpreted as good (>0.75) or excellent (>0.90) (37). After normality confirmation using the

Jarque-Bera test, we calculated the Pearson's correlation coefficient (*r*) to confirm the effect

of body size on the lower-limb inertia properties.

We tested the significance of the intergroup differences using an independent *t*-test.

The statistical significance was set at 0.05. The effect size of each *t*-test was determined using

Cohen's *d* (33) with classification as small (≤0.49), medium (0.50–0.79), large (0.80–1.19),

very large (1.20–1.99), and huge (≥2.00) (21).

RESULTS

The body size parameters in sprinters were significantly larger than those in

non-sprinters (all *p*<0.01, *d*=1.60, 2.16; Table 2). The absolute values of the lower-limb mass

and moment of inertia were significantly correlated to the body size parameters, while the

relative mass and $\hat{I}_{LowerLimb}^{Hip}$ were not (Fig. 3).

The relative mass of the entire lower limb was larger in sprinters (18.7±0.7% body

mass) than in non-sprinters (17.6±0.6% body mass) ($p<0.01$, $d=1.73$; Fig. 4A). However,

$\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$ in sprinters (0.044±0.002 [a. u.]) and non-sprinters (0.042±0.002 [a. u.]) did not

differ significantly ($p=0.15$, $d=0.62$; Fig. 4B). From individual plots, we can observe that,

although one of the non-sprinters showed a small $\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$ (Fig. 4B), the $\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$ in

non-sprinters and sprinters, except for him, were highly comparable.

The relative thigh mass was significantly larger in sprinters (12.9±0.4%) than in

non-sprinters (11.9±0.5%) ($p<0.01$, $d=2.01$), while those in the shank (4.5±0.3% vs 4.3±0.3%)

and foot (1.4±0.1% vs 1.4±0.1%) were similar between them ($p=0.13$, 0.61, $d=0.67$, 0.22; Fig.

5A). Compared to non-sprinters, all lower-limb segments in sprinters had a significantly

larger lean mass ratio (thigh: 81.7±2.0% vs 72.0±4.8%; shank: 73.0±2.0% vs 66.9±3.1%; foot:

52.3±2.5% vs 48.3±1.9%; all $p<0.01$, $d=1.85$–2.62) and smaller fat mass ratio (thigh:

10.3±2.1% vs 19.2±5.6%; shank: 10.2±2.7% vs 15.3±3.5%; foot: 19.6±1.6% vs 24.0±2.5%; all

$p<0.01$, $d=1.64$–2.09) (Fig. 5B). The mass ratio of the bone tissue in the thigh was smaller in

sprinters (8.1±0.6%) than in non-sprinters (8.9±1.1%) ($p=0.03$, $d=1.00$; Fig. 5C), while those

in the shank and foot were similar between the groups ($p=0.24$, 0.72, $d=0.51$, and 0.15; Fig.

5B). The densities of all lower-limb segments were significantly larger in sprinters than in

non-sprinters (thigh: 1.064±0.004 g/cm^3 vs 1.049±0.012 g/cm^3; shank: 1.084±0.007 g/cm^3

vs 1.073±0.008 g/cm^3; foot: 1.061±0.003 g/cm^3 vs. 1.051±0.008 g/cm^3) (all $p<0.01$,

$d=1.29$–1.60; Fig. 5C). We showed the complete scaling coefficients to calculate the

lower-limb segment inertia parameters for sprinters and non-sprinters in Table 3. The radii of

gyration, the moment of inertia normalized by the segment length, and segment mass around

the segmental mediolateral axis in the thigh (27.7±0.3% vs 27.9±0.2%; $p=0.02$, $d=1.08$) and

shank (27.0±0.3% vs. 27.6±0.3%; $p<0.01$, $d=2.24$) were significantly smaller in sprinters than

in non-sprinters.

For the values relating to the examiner's manual processing (digitizing of the

anatomical landmarks on MRIs), we confirmed the excellent repeatability in segment lengths

with the intraclass correlation coefficients (ICCs) and coefficient of variation (CVs) of 0.993

and 0.01–0.36% for the thigh, 0.994 and 0.00–0.57% for the shank, and 0.963 and

0.17–1.44% for the foot.

DISCUSSION

We found that sprinters have lower limbs with a larger relative mass but not a larger

$\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$ than non-sprinters. Although muscular adaptation, as previously demonstrated

(7–9), increases the force and power exertion capacity, the increase in mass due to

hypertrophy has been speculated to make it difficult for the body to move in terms of inertia.

However, we revealed that the mechanical difficulty in swinging the lower limb in sprinters

does not increase relatively.

A notable difference in the segment mass between sprinters and non-sprinters was

confirmed only in the thigh. Moment of inertia I is mechanically proportional to the square

of the radius of gyration r:

$$I = mr^2. \tag{3}$$

For example, if same mass (e.g., 1 kg) is added to the thigh and shank in the sprinters in this

study ($r_{\text{Hip-to-ThighCoM}}$: 0.19 m; $r_{\text{Hip-to-ShankCoM}}$: 0.60 m), the increase in lower-limb

moment of inertia due to the added mass in the thigh (0.036 kg·m^2) is only 1/10 of that due to

the added mass to the shank (0.360 kg·m^2). Based on simple mechanical laws, a small inertia

can lead to both decreasing musculoskeletal demands and increasing body segment

acceleration. Thus, bottom-heavy, top-light feature of the lower limb in sprinters can be an

optimal solution for their performance, from the biomechanical perspective. All the thigh,

shank, and foot segments in sprinters had greater lean tissue ratios, smaller fat tissue ratios,

and greater densities than those in non-sprinters. The sprinter/non-sprinters ratios of segment

density were very similar in all the three segments, including the thigh (approximately 101%).

The sprinter/non-sprinters ratio of thigh density was smaller than that of the thigh mass

(107.6%). These results indicate that the segment mass is more affected by the segment

volume than the segment density. Furthermore, the bone mass ratio of the sprinters was nearly

10% smaller than that of non-sprinters only in the thigh, which is straightforward to interpret

as the adaptation to sprinting involves an increase in segment volume only in the thigh. Taken

together, we indicated that the lower-limb inertia properties in sprinters, having a larger

relative mass but not a larger $\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$, are the result of the hypertrophic potential being

greater only in the thigh.

The proximal-segment-specific hypertrophic potential in the lower limb in sprinters

is related to the muscle-tendon arrangement. The length ratio of the tendon to the muscle in a

muscle-tendon unit (MTU) in humans has a large variation (0.01–11.25), which is generally

because proximal MTUs have shorter tendons compared to distal ones (34). In a previous

study (35), the Achilles tendon thickness in sprinters and non-sprinters was similar

(sprinters/non-sprinters: 98.6%), whereas the ankle plantar flexor muscle thicknesses were

larger in sprinters than in non-sprinters (113.8–116.8%). A previous meta-analysis (36) also

showed a slight effect (weighted average effect size: 0.24) on the tendon hypertrophy in

response to various types of exercise interventions. As such, the tendon tissue is hardly

hypertrophic. Because of the difference in the hypertrophic potential between the muscle and

tendon, the distal segment with a larger tendon length ratio results in a smaller volumetric

(and mass) plasticity in the distal segment. Thus, it is suggested that the inter-segment

differences in the hypertrophy are not a unique feature of sprinters, but the result of an

adaptation potential in general in humans to explosive physical exercise. Regarding the

significance of a larger tendon length ratio in the distal segment, a simulation study (37)

reported that a larger mechanical energy generation is obtained with a longer tendon when the

mass is small and *vice versa*. This suggests the morphological advantage of the muscle-tendon

arrangement in humans for executing explosive motion from the force/work exertion

perspective. However, to the best of our knowledge, the morphological advantages for the

inertial properties in humans have not been examined. The lower limb inertia properties found

in sprinters, having a larger mass (related to muscle potential) but not a larger $\hat{I}_{\text{LowerLimb}}^{\text{Hip}}$, are

beneficial both in preventing injury and improving performance. As discussed above, the

increase in the lower-limb moment of inertia due to the added mass to the shank is 10-fold of

that due to the added mass to the thigh; a smaller volumetric (and mass) plasticity in the distal

segment is an important factor to keep the lower-limb moment of inertia small. Thus, our

findings further add quantified evidence showing a novel significant advantage of the

muscle-tendon arrangement for the adaptability to explosive (sprinting) motion from an

inertial perspective.

We found differences in the normalized body segment inertia parameters (BSIPs),

such as the radii of gyration, between sprinters and non-sprinters. Researchers have

previously developed scaling coefficients to calculate BSIPs for motion analysis using direct

measurements of elderly cadavers (15) and *in vivo* medical imaging such as gamma-ray

scanning (16), computed tomography (17), or MRI (18, 19); however, these scaling factors do

not adequately accommodate the effects of morphological adaptation on the inertial properties

in athletes. We showed the complete scaling coefficients to estimate the BSIPs of the

sprinter's lower-limb segments, which provides sufficient scaling equations to calculate the

hip, knee, and ankle joint dynamics. Although it is slightly outside the scope of this study, our

data allow future studies to analyze motion in male sprinters more accurately.

Below, we discuss the methodological advantages and limitations of the present

study. First, most of our MRI segmentation processes were computationally and automatically

performed by the widely used standard Otsu's binarization algorithm (23) on

fat/water-separated images, leading to advantages in the methodological simplicity and robust

results. Alternately, the landmark position coordinates were acquired by manual digitizing on

the MRIs, having potential effects on the segment length used for data normalization.

However, we acquired MRI images with a slice thickness of 2 mm, which is thinner than the

previous studies examining inertia properties [for example, Sreenivasa et al. (24): 6 mm–48

mm; Pearsall et al. (28): 10 mm; Cheng et al. (18): 20 mm]. We confirmed excellent

repeatability of the segment lengths (ICC>0.96 and CV<1.5%); thus, the potential effect of

manual digitizing on the segment length would not critically alter our results. Second,

non-sprinters had similar body sizes as the average Japanese 20 year old males shown in the

2019 Japanese Government statistics (38), while the sprinters were larger than the standard

Japanese population, which might affect our conclusion. However, we assessed the

dimensionless inertia parameters calculated according to Hof (32) to eliminate the effect of

body size. Although there were significant correlations between the absolute inertia and body

size parameters, the dimensionless values (the relative mass and the normalized moment of

inertia of the whole lower limb around the hip joint) did not correlate with the body size,

suggesting that these values could reflect the individual differences in body shapes and mass

distributions, thereby not being critically affected by the inter-group differences in the body

size. Third, although we determined the sample size of each group as 12 based on *a priori*

power analysis (see Methods section), we excluded one of the sprinter's data due to an error

in the water/fat separation of his foot scan. However, the main findings regarding intergroup

differences were statistically significant, which suggest that the exclusion did not critically

affect our conclusion.

We examined only male participants. Generally, body composition has large sex

differences (39); therefore, our findings might not be directly applicable to females, which is

an important future theme. It is also unclear whether our findings are applicable to the upper

limbs. It has been suggested that high-speed throwing in the upper limbs is one of the factors

for evolution in current humans (40). It would not be surprising if humans also have

morphological advantages in their upper limbs from an inertial perspective. Furthermore,

unlike the lower limbs, which constantly support the body against gravity, the upper limbs,

being free from such constant gravitational load, would reflect pure trainability in humans; the

morphological adaptation in the upper limbs is an interesting future theme for understanding

human plasticity in motor performance. Nevertheless, our findings regarding

non-correspondence with an increase in lower-limb mass and moment of inertia in male

sprinters open a novel perspective, inertia, for understanding human plasticity.

In conclusion, we demonstrate the lower-limb morphology in sprinters with a larger

relative mass but with a not larger normalized moment of inertia around the hip comparing it

to non-sprinters. This result suggests one of the practical advantages of human morphological

characteristics–a larger tendon length ratio in the distal segment muscles–for their adaptation

to explosive motor tasks, which provides a novel perspective to the literature for

understanding human morphological adaptability.

Acknowledgements

We would like to thank Dr. Gaku Kakehata for his help in participant recruitment.

Author Contributions:

N.S., and Y.K. conceived and designed research; N.S. and H.I. performed

experiments; N.S. analyzed data; N.S. interpreted results of experiments; N.S. prepared figures;

N.S. drafted manuscript; N.S., H.I, and Y.K. edited and revised manuscript; N.S., H.I, and Y.K.

approved final version of manuscript.

Conflict of interest statement:

There are no conflicts of interest to declare.

GRANTS:

Yamaha motor foundation for sports to N.S. and JSPS KAKENHI Grant-in-Aid for

Young Scientists (21K17592) to N.S.

References

1. **Young NM**, **Wagner GP**, **Hallgrímsson B**. Development and the evolvability of

human limbs. *Proc Natl Acad Sci U S A* 107: 3400–5, 2010. doi:

10.1073/pnas.0911856107.

2. **Bramble DM**, **Lieberman DE**. Endurance running and the evolution of Homo. *Nature*

432: 345–352, 2004. doi: 10.1038/nature03052.

3.   **Japan Sports Agency**. The Report of FY 2020 Survey on Physical Strength and

Athletic Performance. 2021.

4.   **Fukunaga T**, **Miyatani M**, **Tachi M**, **Kouzaki M**, **Kawakami Y**, **Kanehisa H**.

Muscle volume is a major determinant of joint torque in humans. *Acta Physiol Scand*

172: 249–255, 2001. doi: 10.1046/j.1365-201x.2001.00867.x.

5.   **O'Brien TD**, **Reeves ND**, **Baltzopoulos V**, **Jones DA**, **Maganaris CN**. Strong

relationships exist between muscle volume, joint power and whole-body external

mechanical power in adults and children. *Exp Physiol* 94: 731–738, 2009. doi:

10.1113/expphysiol.2008.045062.

6.   **Balshaw TG**, **Maden-Wilkinson TM**, **Massey GJ**, **Folland JP**. The Human Muscle

Size and Strength Relationship: Effects of Architecture, Muscle Force, and

Measurement Location. *Med Sci Sports Exerc* 53: 2140–2151, 2021. doi:

10.1249/MSS.0000000000002691.

7.   **Ema R**, **Sakaguchi M**, **Kawakami Y**. Thigh and Psoas Major Muscularity and Its

Relation to Running Mechanics in Sprinters. *Med Sci Sport Exerc* 50: 2085–2091, 2018.

doi: 10.1249/MSS.0000000000001678.

8.   **Tottori N**, **Suga T**, **Miyake Y**, **Tsuchikane R**, **Tanaka T**, **Terada M**, **Otsuka M**,

Nagano A, **Fujita S**, **Isaka T**. Trunk and lower limb muscularity in sprinters: what are

the specific muscles for superior sprint performance? *BMC Res Notes* 14: 10–15, 2021.

doi: 10.1186/s13104-021-05487-x.

9.   **Miller R**, **Balshaw TG**, **Massey GJ**, **Maeo S**, **Lanza MB**, **Johnston M**, **Allen SJ**,

Folland JP. The Muscle Morphology of Elite Sprint Running. *Med Sci Sport Exerc* 53:

804–815, 2020. doi: 10.1249/mss.0000000000002522.

10.   **Takahashi K**, **Kamibayashi K**, **Wakahara T**. Muscle size of individual hip extensors

in sprint runners: Its relation to spatiotemporal variables and sprint velocity during

maximal velocity sprinting. *PLoS One* 16: 1–12, 2021. doi:

10.1371/journal.pone.0249670.

11.   **Schache AG**, **Blanch PD**, **Dorn TW**, **Brown NAT**, **Rosemond D**, **Pandy MG**. Effect

of running speed on lower limb joint kinetics. *Med Sci Sports Exerc* 43: 1260–1271,

2011. doi: 10.1249/MSS.0b013e3182084929.

12.   **Dorn TW**, **Schache AG**, **Pandy MG**. Muscular strategy shift in human running:

dependence of running speed on hip and ankle muscle performance. *J Exp Biol* 215:

1944–56, 2012. doi: 10.1242/jeb.064527.

13.   **Huang HK**, **Wu SC**. The evaluation of mass densities of the human body in vivi from

CT Scans. *Comput Biol Med* 6: 337–343, 1976. doi: 10.1016/0010-4825(76)90070-6.

14.   **Martin PE**, **Mungiole M**, **Maezke MW**, **Longhill JM**. The use of magnetic resonance

imaging for measuring segment inertial propertierties. *J Biomech* 22: 367–376, 1989.

15.   **Clauser CE**, **McConville JT**, **Young JW**. Weight, volume, and center of mass of

segments of the human body. Aerospace Medical Research Laboratory,

Wright–Patterson Air Force Base, Dayton, 1969.

16.   **de Leva P**. Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *J

Biomech* 29: 1223–1230, 1996. doi: 10.1016/0021-9290(95)00178-6.

17.   **Huang HK**, **Suarez FR**. Evaluation of cross-sectional geometry and mass density

distributions of humans and laboratory animals using computerized tomography. *J

Biomech* 16: 821–832, 1983. doi: 10.1016/0021-9290(83)90006-4.

18.   **Cheng CK**, **Chen HH**, **Chen CS**, **Lee CL**, **Chen CY**. Segment inertial properties of

Chinese adults determined from magnetic resonance imaging. *Clin Biomech* 15:

559–566, 2000. doi: 10.1016/S0268-0033(00)00016-4.

19.   **Mungiole M**, **Martin PE**. Estimating segment inertial properties: Comparison of

magnetic resonance imaging with existing methods. *J Biomech* 23: 1039–1046, 1990.

doi: 10.1016/0021-9290(90)90319-X.

20.   **Shan G**, **Bohn C**. Anthropometrical data and coefficients of regression related to

gender and race. *Appl Ergon* 34: 327–337, 2003. doi: 10.1016/S0003-6870(03)00040-1.

21.   **Sawilowsky SS**. New Effect Size Rules of Thumb. *J Mod Appl Stat Methods* 8:

597–599, 2009. doi: 10.22237/jmasm/1257035100.

22. **Klingensmith JD**, **Elliott AL**, **Fernandez-del-Valle M**, **Mitra S**. Automated

segmentation of cardiac adipose tissue in Dixon magnetic resonance images. *J Biomed*

Graph Comput 8: 1, 2017. doi: 10.5430/jbgc.v8n1p1.

23. **Otsu N**. A threshold selection method from gray-level histograms. *IEEE Trans Syst*

Man Cybern 9: 62–66, 1996.

24. **Sreenivasa M**, **Chamorro CJG**, **Gonzalez-Alvarado D**, **Rettig O**, **Wolf SI**.

Patient-specific bone geometry and segment inertia from MRI images for model-based

analysis of pathological gait. *J Biomech* 49: 1918–1925, 2016. doi:

10.1016/j.jbiomech.2016.05.001.

25. **Harrington ME**, **Zavatsky AB**, **Lawson SEM**, **Yuan Z**, **Theologis TN**. Prediction of

the hip joint centre in adults, children, and patients with cerebral palsy based on

magnetic resonance imaging. *J Biomech* 40: 595–602, 2007. doi:

10.1016/j.jbiomech.2006.02.003.

26. **Rossi M**, **Lyttle A**, **El-Sallam A**, **Benjanuvatra N**, **Blanksby B**. Body segment inertial

parameters of elite swimmers using DXA and indirect methods. *J Sport Sci Med* 12:

761–775, 2013.

27. **Pearsall DJ**, **Reid JG**, **Livingston L a**. Segmental inertial parameters of the human

trunk as determined from computed tomography. *Ann Biomed Eng* 24: 198–210, 1996.

doi: 10.1007/BF02667349.

28. **Pearsall DJ**, **Reid JG**, **Ross R**. Inertial properties of the human trunk of males

determined from magnetic resonance imaging. *Ann Biomed Eng* 22: 692–706, 1994.

doi: 10.1007/BF02368294.

29. **Sado N**, **Yoshioka S**, **Fukashiro S**. A non-orthogonal joint coordinate system for the

calculation of anatomically practical joint torque power in three-dimensional hip joint

motion. *Int J Sport Heal Sci* 15: 111–119, 2017. doi: 10.5432/ijshs.201712.

30. **Sado N**, **Yoshioka S**, **Fukashiro S**. Curved approach in high jump induces greater

jumping height without greater joint kinetic exertions than straight approach. *Med Sci*

Sport Exerc 54: 120–128, 2022. doi: 10.1249/mss.0000000000002761.

31.   **Dumas R**, **Chèze L**, **Verriest JP**. Adjustments to McConville et al. and Young et al.

body segment inertial parameters. *J Biomech* 40: 543–553, 2007. doi:

10.1016/j.jbiomech.2006.02.013.

32.   **Hof AL**. Scaling gait data to body size. *Gait Posture* 4: 222–223, 1996. doi:

10.1016/0966-6362(95)01057-2.

33.   **Cohen J**. Statistical power analysis for the behavioral sciences. 2nd ed. Hillsdale:

Lawrence Earlbaum Associates, 1988.

34.   **Hoy MG**, **Zajac FE**, **Gordon ME**. A musculoskeletal model of the human lower

extremity: The effect of muscle, tendon, and moment arm on the moment-angle

relationship of musculotendon actuators at the hip, knee, and ankle. *J Biomech* 23:

157–169, 1990. doi: 10.1016/0021-9290(90)90349-8.

35.   **Kubo K**, **Ikebukuro T**, **Yata H**, **Tomita M**, **Okada M**. Morphological and mechanical

properties of muscle and tendon in highly trained sprinters [Online]. *J Appl Biomech* 27:

336–344, 2011.

http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L36

3098363%5Cnhttp://journals.humankinetics.com/AfcStyle/DocumentDownload.cfm?

DType=DocumentItem&Document=08-Kubo-336-344.pdf%5Cnhttp://sfx.ub.rug.nl:9

003/sfx_local?sid=EMBASE&issn=1.

36.   **Bohm S**, **Mersmann F**, **Arampatzis A**. Human tendon adaptation in response to

mechanical loading: a systematic review and meta-analysis of exercise intervention

studies on healthy adults. *Sport Med - Open* 1, 2015. doi: 10.1186/s40798-015-0009-9.

37.   **Nagano A**, **Komura T**, **Fukashiro S**. Effects of the length ratio between the contractile

element and the series elastic element on an explosive muscular performance. *J*

Electromyogr Kinesiol 14: 197–203, 2004. doi: 10.1016/S1050-6411(03)00085-3.

38.   **Government of Japan**. e-Stat: Portal Site of Official Statistics of Japan [Online]. 2020.

https://www.e-stat.go.jp/en.

39.   **Bredella MA**. Sex Differences in Body Composition BT　- Sex and Gender Factors

Affecting Metabolic Homeostasis, Diabetes and Obesity. In: , edited by Mauvais-Jarvis

F. Springer International Publishing, p. 9–27.

40.   **Roach NT**, **Venkadesan M**, **Rainbow MJ**, **Lieberman DE**. Elastic energy storage in

the shoulder and the evolution of high-speed throwing in Homo. *Nature* 498: 483–6,

2013. doi: 10.1038/nature12267.

FIGURE LEGENDS

        Figure 1   **Flowchart of the segmentation process based on water/fat separation images.**

        Figure 2   **A typical example of the constructed data. A three-dimensional constructed volume data of the thigh, shank and foot (A) and a stick picture created from three-dimensional position coordinates of the entire hypothetically constructed lower limb (B).** Note: The origin of the hypothetical coordinates was at the hip joint center. The lower limb length was defined as the height difference between the hypothetical coordinates of the hip and calcaneus.

        Figure 3   **Relationships between absolute (A, C, E: left column) and normalized (B, D, F: right column) inertia parameters and body size parameters (A–D: body mass, E–F: body height) in sprinters (black) and non-sprinters (grey).** Note: *n*, and *r* were the sample size and Pearson's correlation coefficients in each group, respectively.

        Figure 4   **Whole lower-limb inertia properties. Relative mass (A) and normalized moment of inertia around the hip joint in hypothetically constructed anatomical position data (B).** Note: Bar graphs show means and standard

deviations. Gray plots show individual data. Asterisk (*) indicates a

significant difference between sprinters and non-sprinters.

Figure 5   **Segment mass and its determinants of each lower-limb segment. Relative mass (A), density (B), and composition ratio (C) in each of the thigh, shank, and foot segments.**

Note: Bar graphs show mean and standard deviation. Asterisk (*) indicates a significant difference between sprinters and non-sprinters.

513   **TABLE LEGENDS**

Table 1   **Anatomical landmarks to be acquired by manual digitizing.**

Table 2   **Characteristics of participants.**

Table 3   **Complete scaling coefficients to estimate segment inertia parameters in sprinters and non-sprinters**

A. Segment mass

B. Composition ratio

C. Segment density

Table 1. Anatomical landmarks to be acquired by manual digitizing.

Segments	Landmarks
Thigh	Surface of the femoral head (30 points)
	Lateral articular cleft between the femur and tibia condyle
	Medial articular cleft between the femur and tibia condyle
Shank	Lateral articular cleft between the femur and tibia condyle
	Medial articular cleft between the femur and tibia condyle
	Malleolus lateralis
	Malleolus medialis
Foot	Malleolus lateralis
	Malleolus medialis
	Calcaneus
	1st meta-tarsal heads
	5th meta-tarsal heads
	2nd toe tip

Note: Anatomical landmarks other than the femoral head were digitized closest points on the skin (i.e., points to which reflective markers can be attached when doing motion capture).

Table 2. Characteristics of participants.

Variable	Sprinters	Non-sprinters
n	11	12
Age [years]	20 ± 2	23 ± 3
Height [cm]	176.7 ± 5.7	168.9 ± 3.9
Mass [kg]	68.2 ± 4.4	58.6 ± 4.5
Leg length [cm]	87.6 ± 4.5	82.7 ± 2.9
$d_{\text{HJC-to-KJC}}$ [cm]	44.1 ± 1.9	41.6 ± 1.6
$d_{\text{KJC-to-AJC}}$ [cm]	37.8 ± 2.3	35.4 ± 1.3
$d_{\text{Cal-to-MH1–MH5midpoint}}$ [cm]	14.4 ± 0.5	13.6 ± 0.5
$d_{\text{HJC-to-ThighCoM}}$ [cm]	19.4 ± 0.8	18.3 ± 0.7
$d_{\text{HJC-to-ShankCoM}}$ [cm]	59.5 ± 2.8	56.1 ± 2.1
$d_{\text{HJC-to-FootCoM}}$ [cm]	86.6 ± 4.3	81.6 ± 2.9

Note: Data are shown as means ± standard deviations. n: the number of participants in each group; $d_{\text{a-to-b}}$: distance from a-to-b in the entire hypothetically constructed lower limb; HJC: Hip joint center; KJC: Kip joint center; AJC: Ankle joint center; Cal: Calcaneus; MH1 (5): 1st (5th) Metatarsal Heads; CoM: Center of mass;

Table 3. Complete scaling coefficients to estimate segment inertia parameters in sprinters and non-sprinters.

Group	Segment	Length definition	Origin	Scaling factor for mass	Scaling factors for CoM position			Scaling factors for tensor of inertial					
				m [%]	x [%]	y [%]	z [%]	r_{xx} [%]	r_{yy} [%]	r_{zz} [%]	r_{xy} [%]	r_{yz} [%]	r_{zx} [%]
Sprinters													
	Thigh	HJC to KJC	HJC	12.9	0.6	-2.7	-44.0	27.7	27.4	13.4	1.8 (i)	5.7 (i)	6.1 (i)
	Shank	KJC to AJC	KJC	4.5	4.4	-4.1	-40.7	27.0	26.9	10.0	1.9 (i)	4.1	4.3 (i)
	Foot	CAL to MH$_1$–MH$_5$ midpoint	AJC	1.4	0.1	-31.0	-46.2	40.9	36.9	28.8	13.7	16.9 (i)	4.5
Non-sprinters													
	Thigh	HJC to KJC	HJC	11.9	1.6	-2.5	-43.9	27.9	27.7	13.0	1.8 (i)	5.7(i)	6.2 (i)
	Shank	KJC to AJC	KJC	4.3	4.5	-3.2	-40.9	27.6	27.5	9.9	2.7 (i)	2.6	3.4 (i)
	Foot	Cal to MH$_1$–MH$_5$ midpoint	AJC	1.4	0.6	-32.3	-43.9	42.3	40.3	24.9	10.1	16.4 (i)	6.3

1 Note: HJC: Hip joint center; KJC: Kip joint center; AJC: Ankle joint center; Cal: Calcaneus;

2 MH1 (5): 1st (5th) Metatarsal Heads; CoM: Center of mass;