The role of floral traits in community assembly process at high elevations in Lesser

Himalaya

Mustaqeem Ahmad[1,2,3], Sergey Rosbakh[4], Solveig Franziska Bucher[5,6], Padma Sharma[1],

Sonia Rathee[2], Sanjay Kr. Uniyal[3,] Daizy R. Batish[2] & Harminder P. Singh[1]

[1] Department of Environment Studies, Panjab University, Chandigarh, India

[2] Department of Botany, Panjab University, Chandigarh, India

[3] Department of Environmental Technology, CSIR-IHBT, Palampur, India

[4] Ecology and Conservation Biology, University of Regensburg, Germany

[5] Institute of Ecology and Evolution, Plant Biodiversity Group, Friedrich Schiller University

Jena, Jena, Germany

[6] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Germany

Abstract

1. Ecological theory postulates that plant trait research should consider multiple traits related

to different organs and/or ontogenetic stages as such traits represent different ecological

niche axes. Particularly, floral traits have been suggested to play an important role in

assembling plant communities along environmental gradients as they determine the

reproductive success, one of the key functions in plants. Yet, the predictive power of floral

traits in community assembly research remains largely unverified empirically.

2. We analyzed the predictive power of six floral traits of 139 herbaceous species for

inferring community assembly process in twenty-one sites located along an elevation

gradient in Lesser Himalaya ranging from 2,000 to 4,000 meters above sea level. The floral

trait variability along the gradient was analyzed using community-weighted trait mean

(CWM) values and functional diversities (FD) calculated for each of the study

communities.

3. The CWM values for onset of flowering and flower display area increased significantly

with increasing elevation, whereas specific flower area showed an opposite pattern. In

combination with convergence in onset of flowering and specific area (i.e., lower FD

values in high elevation sites), these patterns suggest that abiotic filtering and plant-

pollinator interactions affected the floral trait composition of the communities studied.

Increasing low-temperature stress towards high-elevation sites selected for late-flowering

species that produce resource-intensive flowers with larger display areas.

4. Low pollinator abundancy and activity in high elevation, could also explain why these

traits were selected in the study communities. Delayed flowering with increasing

elevations might facilitate the phenological overlap of plants and their pollinators, as

pollinator activity at higher elevation peaks in the second half of the vegetation period. The

dominance of species with low specific flower area and larger display area in high

elevation communities were attributed to the increased flower longevity and attraction of

pollinators, respectively, to maximize pollination success under pollinator scarcity.

5. *Synthesis*. Our study provides empirical support of the recent argument that floral traits

contribute considerably to the assembly of plant communities along environmental

gradients. Thus, such traits should be included into community assembly research agenda

as they represent key growth and survival ecological functions.

Keywords: biotic interaction, elevation, filter, flower, gradient, pollinator, temperature

Introduction

Understanding the processes shaping the structure and composition of biological communities

along environmental gradients, i.e., the community assembly mechanisms (Keddy, 1992), is

one of the fundamental aims in ecological research. Disentangling abiotic and biotic factors

selecting for or against species from the regional to local species pools allows us not only to

understand the current community compositions but can also help to answer the question how

these communities will respond to future environmental changes (Götzenberger et al., 2012).

Traditionally, community assembly processes have been scrutinized through the powerful

lenses of trait-based approaches (McGill et al., 2006). These methods are built on the

paradigm that functional traits, that is the physiological, morphological or phenological

characteristics of a plant individual which directly affect its growth, reproduction, and

survival (Violle et al., 2007), determine species' responses to abiotic environment and biotic

interactions (Pillar et al., 2021). Therefore, trait variability within a community reflects the

outcomes of abiotic filtering and biotic interactions which are two key assembly rules (Bello

et al., 2013; Spasojevic & Suding, 2012) across different scales (Kraft et al., 2008, Weiher et

al., 2011).

There is a common agreement that, to maximize our understanding of trait-based community assembly, traits related to different organs should be considered as they relate to different ecological niche axes (Grime et al., 1997; Laughlin, 2014). However, the published studies are biased strongly towards traits connected to vegetative growth, survival and resource acquisition (i.e., vegetative traits) with traits related to sexual reproduction being largely ignored (E-Vojtkó et al., 2020; Jiménez☐Alfaro et al., 2016; Rosbakh et al., 2018), therefore, including these traits into the current research agenda is important to predict complex environmental effects on trait composition within communities (E-Vojtko et al., 2020; Poschlod et al., 2013; Rosbakh et al., 2018).

In this context, floral traits, physiological, morphological and phenological characteristics of flowers and inflorescences and the way they are pollinated (Klotz et al., 2002), are of a particular interest. Several floral traits, such as the number of flowers, flower display area, scent, and petal colour, determine the success of reproduction, one of the main functions in plants (Barrett, 2010; Soltis & Soltis, 2014). Thus, to maximize plant reproductive fitness, these traits display strong intra- and interspecific variability along environmental gradients reflecting adaptations in plants to specific abiotic conditions and/or pollinator availability (E-Vojtkó et al., 2020). For example, soil moisture availability limits plant investment in flowers with smaller floral area in dry habitats (Lambrecht & Dawson, 2007). Further, Burkle and Irwin (2010) reported that low soil nitrogen content has direct effects on survival and adult fecundity, and results in increased length of flowering stalks. These findings suggest that, at the community level, abiotic filtering could select floral traits and thus leads to shifts in community functional trait composition. Another potential driver of community assembly process in flowering plants is plant-plant and plant-pollinator interactions, for which previous research has indicated that some floral traits (e.g., flower morphology, colour, and scent) can be involved in competition and facilitation via pollinators (Sargent & Ackerly, 2008). Pollinators, particularly, bees can differentiate between flower temperatures which influences

their behaviour as they prefer warm flowers (with larger display area) and warm nectar over

unheated ones (Dyer et al., 2006). Norgate et al. (2010) reported that this behaviour becomes

more pronounced with decreasing temperatures. Learning behaviour of pollinators may

indirectly enhance successful pollination via their preference for some colours through their

ability to associate flower temperature with specific colours (van der Kooi et al., 2019).

Additionally, this learning behaviour can also affect flowering communities through

frequency of their visitation as inexperienced pollinators may be attracted to larger display

areas whereas experienced pollinators prefer greater floral rewards (Makino & Sakai, 2007).

Finally, some phenological traits, such as onset and duration of flowering, can be affected

by both, abiotic filtering, and biotic interactions (Bucher & Römermann, 2020; Ulrich et al.,

2020). Temperature has been reported to influence flowering phenology as higher

temperatures lead to an advance in first flowering day (Menzel et al., 2006). Similarly,

temperature also displays strong association with flowering duration, with warmer

temperature leading to extended flowering period (Bucher & Römermann, 2020; Campbell &

Halama, 1993). On the other hand, biotic interactions via pollinators also impact flowering

duration as animal-pollinated flowers have a higher flowering duration as compared with

wind-pollinated plants (Bucher & Römermann, 2020; Rabinowitz et al., 1981).

The aim of this study was to examine the role of six selected floral traits – onset of

flowering, flowering duration, flower number, flower investment, flower display area and

specific flower area – for inferring community assembly process along an elevational gradient

in the Lesser Himalaya. Using *in-situ* measured trait data for 139 herbaceous species

occurring in 21 sites ranging from 2,000 to 4,000 m above sea level (a.s.l.), we analysed

variation in community-weighted trait mean (CWM) values and functional diversities (FD),

two key metrics of community functional structure (Ricotta & Moretti, 2011), to infer the

community assembly processes. Specifically, given the increasing low-temperature stress

towards higher elevations and high frost-sensitivity of flowers (Wagner et al., 2011), we

anticipated a significant shift in plant strategies towards stress-avoidance by delaying the

onset of flowering and consequently shorter flowering duration in high elevation communities

(Ahmad et al., 2021; Bucher & Römermann, 2020; Table 1). In addition to that, the short

growth period with frequent and severe frosts coupled with generally limited soil resources

was expected to select for species which allocate their resources to vegetative growth rather

than to sexual reproduction (Laiolo & Obeso, 2017; Rosbakh & Poschlod, 2021). In terms of

floral traits, we hypothesized that communities at higher elevations are assembled by species

which produce fewer flowers with smaller specific flower area and invest less into flower

biomass and consequently have smaller floral display areas. In other words, the strong abiotic

filtering at high elevations should lead to significant decrease in the CWMs and stronger trait

convergence (low FD values) in all six traits (except for increasing CWMs for onset of

flowering) with increasing elevation.

Alternatively, we hypothesized that elevation-specific plant-pollinator relationships also

affect functional composition of the communities. Considering that higher elevational sites

observe generally sparse vegetation and lower pollinator abundance (Körner, 2021; Trunschke

& Stöcklin, 2017), we expected to observe a pattern of delayed and synchronized flowering

that continues for a longer period at high-elevational communities as a strategy to maximize

pollinator visits and thus to increase the possibility of successful pollination. In other words,

the CWMs for the onset of flowering and flower duration were expected to decrease and

increase, respectively, along the elevational gradient. The increasing role of facilitation with

increasing elevation (i.e., synchronized flowering will attract pollinators and thereby increase

the chances of pollination for all community members; Tachiki et al., 2010) should be

reflected in the FD variation. Particularly, comparatively higher competition for pollinators in

lower elevations was anticipated to result in trait divergence (higher FD at lower than at

higher elevation). With respect to the decreasing pollinator abundance along the elevational

gradient, we further hypothesized an increase in flower number, specific flower area, flower

display area and flower investment in high elevation communities, as it could be

advantageous to enhance pollinator attraction in fragmented alpine habitats combined with a

short flowering season and generally low pollinator abundance (Fabbro & Körner, 2004; Zhu

et al., 2010; Körner, 2021). That is, the CWM and FD values should increase and decrease

with elevation, respectively.

Table 1. Overview of floral traits studied and their hypothesized role in the community

assembly process with increasing elevation.

Traits	Abiotic filtering	Biotic interactions
Onset of flowering	Decreases as a stress-avoidance adaptation. Trait convergence as a result of strong environmental filtering	Delayed and synchronized onset of flowering to maximize pollination success. Highest trait values and trait divergence in lower sites due to high competition for pollinators
Duration of flowering	Shorter as onset of flowering is comparatively later	Longer to increase chance to be pollinated. Trait divergence in lower elevation due to comparatively high competition for pollinators. Higher elevation: trait divergence due to facilitation effects
Flower number	Fewer at higher elevations due to resource limitations and risky sexual reproduction	More at higher elevations (clustering of flowers) for increased pollinator attraction and visitation. Strong trait convergence due to high facilitation effects
Specific flower area	Smaller at higher elevations due to low resource availability	Increases along elevation to ensure successful pollination and fertilization. Trait divergence in lower elevation due to competition for pollinators

	Smaller at higher elevations due to fewer resources and uncertain sexual reproduction	Larger at higher elevation to attract pollinators. Strong trait convergence at higher elevation due to high facilitation effects
Floral display area		
Flower investment	Decreases with increasing elevation because of increasing unfavorability for sexual reproduction	Higher in higher elevation to attract pollinators and ensure successful sexual reproduction. Strong trait convergence due to high facilitation effects. Trait divergence in lower elevation due to plant-plant competition for available pollinators

Materials and methods

Study system

The study was conducted in the Dhauladhar mountain range, Lesser Himalayas (Himachal Pradesh, NW India; Fig. 1). The region is situated at the junction of western, north-western, and trans-Himalayan ranges and is characterized by high levels of floral and faunal diversity due to the combination of diverse geomorphological, geological, and climatic factors (Ahmad, 2021). The study region has a typical alpine relief, with steep mountain peaks reaching up to 6,200 m a.s.l.. The bedrocks of the Dhauladhar ridge are mainly composed of granite and gneiss (Singh & Singh, 1987). The region has a temperate climate with the growth period lasting from April to September in lowlands with average temperatures of 20°C (https://www.adb.org/). There is a decrease in mean annual temperatures with a lapse rate of 6.5° K km^{-1} of elevation (Kattel et al., 2013). The precipitation pattern in Himachal Pradesh is mainly governed by western disturbances and southwest monsoon (Jaswal et al., 2015). Mean annual precipitation in the region is about 2,900 mm a^{-1} (Jaswal et. al., 2014) with a pronounced rainy season from July to September (Singh & Singh, 1987).

The vegetation in the study region between 2,000 and 4,000 m a.s.l. is formed by mixed forests with dominance of *Abies pindrow, Aesculus indica, Cedrus deodara, Pinus*

wallichiana, Picea smithiana, Quercus semecarpifolia and *Rhododendron arboretum.* The

tree line starts at around 3,000 m a.s.l.; above the tree line herbaceous species like *Aconitum*

heterophyllum, Iris kemaonensis, Lindelofia macrostyla, Meconopsis aculeata, Polygonum

amplexicaule, Potentilla argyrophylla and *Rumex nepalensis* dominate the grasslands (Ahmad

et al., 2021). The nival zone with sparse and scattered vegetation starts at about 4,300 m a.s.l..

The lower sites are easily accessible and thus experience high anthropogenic pressure in form

of grazing, tourism, and logging (Ahmad, 2021). In the alpine vegetation belt, land-use is

limited to vertical transhumance of comparatively small flocks of sheep and goat.

In 2015, we established 21 permanent plots, each of 20×20 m in size at every 100 m

increase in elevation ranging from 2,000 to 4,000 m a.s.l. and thus covering the sub-temperate

to alpine vegetation belt. The plant community structure and composition in each plot was

assessed by systematic sampling using the quadrat method (Bellhouse, 2005). Specifically,

within each plot we marked 5 sub-plots of 5×5 m (a total of 105 sub-plots across elevation)

and within these sub-plots, 25 1×1 m quadrats (a total of 525 sub-plots across elevation) were

marked to record presence-absence and abundance of shrubs and herbs in the community,

respectively (Appendix 1). The vegetation in the plots was studied in 2018 during the peak

growing season (February to November, depending on elevation).

Each site was characterized in terms of temperature, soil nutrients and water supply. We

used mean soil temperature at 10-15 cm soil depth measured during the growing season (June

to September) as a proxy for the temperature conditions (Scherer Körner, 2011). The soil

temperatures were measured at a 30-minute interval by data loggers (Geo Precision

Environment Technology, Germany) each installed in the centre of every permanent plot.

To characterise the soil conditions, soil samples were collected from the centre of every

5×5 m sub-plot at each permanent plot, i.e., 5 replicates per plot. The soil samples were air

dried. Subsequently, we measured the concentration of available nitrogen, phosphorous and

potassium (AN, AP, AK, respectively) as well as water saturation (WS; a proxy for soil

moisture content) following the methods provide by Ahmad et al. (2020). Study sites

coordinates, temperature logger code, and mean values of abiotic factors are given in

Appendix 2 Table S1.

Figure 1. The location of the study sites in India (a) and Himachal Pradesh (b), and their

spatial orientation along the elevational gradient (c). See Appendix 2 Table S1 for detailed

site characteristics.

Trait measurements

Based on the vegetation surveys, we selected the 139 most abundant insect-pollinated species

(i.e., species that were present in more than three sites and with an abundance of >3%) from

which flower and phenological traits were measured. We conducted regular phenological

observations from February to November 2018, we collected data for a total of 10,025 visits

to 1 × 1 m quadrats during 401 visits of the 21 permanent plots during the entire study

(Appendix 2; Table S2). We recorded floral abundance of each species (if present) in 25 (1 ×

1 m) sub-plots on each 20 × 20 m plot. We recorded the flowering stages (bud, open flower,

or senescent flower) for every individual of a species in 1 × 1 m quadrats. Within these 1 × 1

m quadrats, the percent (%) flowering of each individual (of a species) was estimated. The

flowering (%) recorded for each individual of a species in 25 (1 × 1 m) quadrats were

averaged to represent flowering (%) of that species' population, for that particular permanent

plot (20 × 20 m). The day of the year (DOY 1 = 1st of January), on which the plant species

reached 20 and 90% of floral abundance was considered as the onset, and end of flowering,

respectively as described in Valencia et al. (2016) and Ahmad et al. (2021). We also estimated

the total flowering duration (TFD) calculated as time span between the end and the onset of

flowering (Crimmins et al., 2013). Additionally, two floral traits namely floral display area

(FDA, mm^2) and specific flower area (SFA, mm^2 mg^{-1}) were measured following the same

protocols as developed for specific leaf area described by Pérez-Harguindeguy et al. (2016).

The FDA is the total area of all flowers or inflorescences of one individual at the peak of

flowering and is an indicator for the investment of a plant into attracting pollinators (higher

values mean higher visibility, pollinator visitation rates and potentially higher fertilization

success) (Fabbro & Körner, 2004; Herrera, 2009; Zhao et al., 2016). SFA is the ratio between

one-sided surface area of fresh flower divided by its dry weight and flower/inflorescence

biomass, a trait that represents resource investment in reproductive parts (Pérez-Harguindeguy

et al., 2016). For the FDA and SFA measurements, 10 to 20 healthy, fully flowering plants

were selected for each species at each plot, dug out completely, wrapped in moist towels and

transported to the laboratory. Fresh flower area was measured with the help of a digital

Vernier calliper (Fabbro & Körner, 2004). Collected plant species were separated into root,

shoot and flowers. These were then dried in oven at 70°C for 48 h followed by weighing

using a balance with a precision of 0.1 mg. Flower/inflorescences number was recorded

manually for each plant species at each plot. Floral investment was calculated as:

$$Floral\ investment = \frac{Flower\ biomass}{Total\ plant\ biomass} \times 100$$

Statistical analysis

The changes in environmental characteristics along the elevational gradient, as well as among

all environmental factors, were analysed using the Pearson correlation coefficient. Polynomial

regressions were applied to test the effect of rainfall and temperature during the sampling

period. A detrended correspondence analysis (DCA) was performed to analyse the

relationship between species occurrence and environmental parameters by using the *vegan*

package in R statistical environment (Oksanen et al., 2015). An environmental fitting test with

999 permutations was performed to analyse correlations of environmental factors with the

DCA axes.

The changes in community functional structure were analysed with the help of community

weighted means (CWMs) and functional diversity (FD). A CWM is the average trait value

weighted by species' relative abundances in a community and thus describes the adaptation

strategy of dominant species to given environmental conditions (de Bello et al., 2021). The

FD is a measure of trait dissimilarity compared to random expectation (i.e., trait convergence

or divergence) and was calculated using Rao's quadratic entropy (Rao, 1982). The deviation

of observed FD from the null expectation, i.e., communities are a random selection of species

from the regional pool was tested with the help of null models. We generated null models by

permuting the columns (species) of the plot-by-species abundance matrix for each plot

(community). Then the standard effect sizes (SES) were calculated to evaluate the deviations

of observed functional diversity values from random expectations. SES were calculated as the

observed FD value minus the mean of the null-FD divided by the standard deviation of the

null-FD. Negative SES indicated that FD was convergent, i.e., lower than expected in the null

expectation. Conversely, positive SES indicated that FD is divergent, i.e., higher than the null

expectations. Further, we assessed the significance of SES by identifying the proportion of

random values that fell below the observed diversity value. The ranks below 0.05 indicated

that FD for a given site was a significant low FD, the ranks above 0.95 indicated significant

high FD.

The variability of the communities (CWM and FD) along the elevational gradient was

estimated with the help of a polynomial regression (i.e., community trait value ~ elevation +

elevation^2), to account for possible non-linear trait-elevation relationship. If the quadratic term

was not significant, the term was removed.

Model assumptions were checked (e.g., homogeneity of variances and normality of

residuals) and met in all cases. The flower display area was log-transformed to improve the

normality of the residuals. All statistical analyses were performed in R version 4.1.0 (R Core

Team 2021).

Results

Variability of environmental conditions along elevational gradient

Soil temperature (r=-0.98, p<0.001) and available phosphorus in soil (r=-0.65, p<0.001)

strongly declined with increasing elevation (Fig. 2). Soil available nitrogen was positively

correlated with soil available potassium (r=0.59, p<0.001) and soil moisture (r=0.66,

p<0.001). Temperature and precipitation peaked from June to September in the study area

(Appendix 2; Fig. S1). The eigenvalue of the first detrended correspondence axis was 0.59

and for the second axis 0.32 (Appendix 2; Fig. S2). Elevation and temperature reflected the

main axis of variation in the multivariate space. Elevation showed negative correlation with

temperature, woody cover, and available phosphorus, whereas available potassium and

available nitrogen showed strong positive correlation with water saturation (Appendix 2; Fig.

S2). High gradient length (DCA axis 1= 4.22) suggests high β-diversity. The sites on the

extreme left of the gradient (S20, S17) share very few similarities with sites on the extreme

right (S5, S6) of DCA axis 1, in terms of environmental factors and vegetation composition

(Appendix 2; Fig. S2).

Figure 2. Correlation matrix for environmental factors measured in the study sites along the

elevational gradient.

Variation of community weighted means along the elevational gradient

CWMs of three out of six traits changed significantly along the elevation gradient (Fig. 3).

Onset of flowering increased strongly with increasing elevations from the third week of May

(DOY 140) at the lower end of the gradient to the second week of July (DOY 190) in the high

elevation areas (Fig. 3A, $R^2 = 0.84$, p<0.001). Specific flower area displayed an opposite

pattern and decreased significantly from 19.6 mm^2 mg^{-1} at 2,100 m a.s.l. (S2) to 9.2 mm^2 mg^{-1}

at 3,800 m a.s.l. (S19 site; Fig. 3D, $R^2 = 0.45$, p<0.001). Floral display area increased with

increasing elevation in a non-linear manner (Fig. 3E, $R^2 = 0.69$, p<0.001) ranging from 8,302

mm^2 at 2,500 m (S6) to the 13,7967 mm^2 at 3,100 m (S10) a.s.l.. In ecological terms, these

patterns suggest that dominant species at high elevations tended to flower later, had larger

display areas, and were composed of flowers with comparatively more resource-intensive

302 flowers (i.e., lower SFA values). The duration of flowering, flower number and flower

303 investment did not show any significant variation with elevation (Fig. 3B, C and F,

304 respectively).

Figure 3. Variation of community weighted means of six traits along the elevational gradient.
Lines indicate significant relationships (p<0.05), the model error is given as a shading.

Variation of functional diversities along the elevational gradient

The FDs of almost all floral traits were non-random across the elevations (Fig. 4). FD of

flowering onset decreased strongly from being divergent at lower elevations to comparatively

strong trait convergence in high elevation areas (Fig. 4A; $R^2 = 0.83$, p<0.001). Additionally,

the FD for specific flower area displayed a significant decrease along the gradient towards

more convergent trait values at higher elevations, yet the amount of variance explained by this

relationship was moderate ($R^2 = 0.32$, p=0.007). The FD of the remaining traits did not show

any significant relationship with the elevation (Fig. 4).

Figure 4. Variation of functional trait diversity (FD) of six traits with elevation. Lines indicate significant relationships according to the linear regressions (p<0.05). Negative and positive standard effect size (SES) indicate a narrower and broader trait range that expected, respectively. Filled circles represent significantly low (rank lower than 0.05) or high (rank above 0.95) FD.

Discussion

Environmental effects on plant regeneration have long been considered as a key to understand plant population dynamics and species distribution patterns (Grubb, 1977; Woodward & Woodward, 1987), yet they have been largely neglected in community ecology research (Jiménez‐Alfaro et al., 2016; Poschlod et al., 2013; Saatkamp et al., 2019). Particularly, floral traits, the main determinants of successful sexual reproduction, are rarely considered in research on community assembly (E-Vojtko et al., 2020). In this study, we close this gap and demonstrate that the so far underused phenological and morphological floral traits are

important for plant community structure, especially in high elevational sites. Specificially, we

reveal that abiotic filters affect phenological traits (e.g. onset of flowering), whereas plant-

pollinator interactions act on flower morphological traits (e.g. floral display area) at the

community level.

As expected, the temperature decreased along the elevational gradient (Fig. 2) and was the

main factor determining the vegetational composition (DCA; Appendix 2 Fig. S2). The low

temperature condition selected for late-flowering species (significantly higher CWM values

for onset of flowering at higher elevations; Fig. 3). The strong filtering effects of low-

temperature could be seen further in the convergent functional diversity of SFA in these sites

(Fig. 4A & D) suggesting exclusion of maladapted early-flowering species. From the

ecological point of view, these patterns can be interpreted as a plant phenological strategy to

avoid frost injury in frost-susceptible flowers (Wagner et al., 2011), which seem to be a

general adaptation of plant reproduction to alpine environments (Körner, 2021). Alternatively,

the delayed and synchronised onset of flowering in high elevation areas (strong trait

convergence suggest a comparatively low number of available temporal niches for flowering)

could be adaptive to facilitate pollination success under conditions with low pollinator

abundancy ('magnet' effect; Koski & Ashman, 2015; Bucher & Römermann, 2020). Further,

delayed flowering with increasing elevation might facilitate plant-pollinator phenological

overlap, as pollinator activity at the higher elevation peaks in the second half of the vegetation

period, when temperatures are high enough (Rawat, 2012). In contrast, the lower elevational

sites seem to harbour a large diversity of plants with different phenological strategies aiming

at reducing plant-plant competition for pollinators (Zhao et al., 2016).

Interestingly, the significantly later onset of flowering did not affect the duration of

flowering in high elevation communities both expressed as CWMs and FDs (Fig. 3B and 4B).

These findings contradict previous observations made in other regions that flowering duration

decreases with increasing elevations, due to decreasing time window for successful

pollination, fertilization, and consequently seed production (Bucher & Römermann, 2020;

Körner, 2021). The most plausible explanation is that the onset of monsoon season in the

study system, which coincides with the peak of flowering, i.e., the middle of the growing

period in the Himalayan alpine vegetation belt (Ram et al., 1988), may relax the low-

temperature filtering effects on plant phenology and thereby promoting longer duration of

flowering. The comparatively long flowering in the resource-limited harsh alpine

environments might be achieved by considerably longer life spans of cost-effective flowers

(Trunschke & Stöcklin, 2017; see below). Cumulatively, these findings contradict our

hypothesis and suggest that flowering duration plays no important role in the community

assembly process in the study system.

Analyzing morphological floral traits, we found that CWMs for specific flower area and

floral display significantly varied along the elevation gradient with high elevation

communities dominated by species with large flower display areas and relatively small SFA

values (Fig. 3D). Most likely, these shifts in CWM can be related to the effects of

environmental filtering and plant-pollinator interactions on sexual regeneration strategies in

alpine environments. As for the former, we assume that low-temperature stress and lower soil

fertility (Fig. 2) as well as stronger winds (Körner, 2021) at higher elevational sites select for

plants that invest more resources per floral display area unit. In alpine regions, flowers adapt

in response to wind induced mechanical perturbations, as heavier flowers have more

mechanical tissue that supports the floral unit, making the structures more flexible and

aerodynamically stable (Cordero et al., 2007; Zhang et al., 2021). The larger flower display

area might be also adaptive in cold alpine climates as it positively correlated with a flowers'

passive heat accumulation (Dietrich and Körner, 2014). As for the plant-pollinator

interactions, the decreased SFA CWM and FD values can be also attributed to the increased

longevity of single flowers at higher elevations to maximize pollination success under

conditions of low pollinator abundancy (Fabbro & Körner 2004; Klatt et al., 2018). Similarly,

a larger display area allows plants from alpine communities to increase pollination probability

by attracting more pollinators (Fabbro & Körner 2004; Kiełtyk, 2021; Zhao et al., 2016).

Regardless of the main driver of SFA variability at the community level, this trait seems to

play an important role in community assembly process as we observed a significantly higher

trait convergence with increasing elevation.

Contrastingly to previous studies in alpine environments (Fabbro & Körner, 2004),

resource investment in reproduction ('flower investment') did not change along the

elevational gradient. Many studies suggested increase in pollinator visitation rates with

increase in flower size and number (Conner & Rush, 1996). However, a reason for no

significant change in flower number with elevation might be that with increasing flower

display size the plant does not need to invest more resources to flower number. In addition,

Samson and Werk (1986) asserted that for understanding the pattern of resource investment in

plants, the species-specific size-dependent effects should be taken into consideration. Obeso

(2002) reported that the responses to plant reproductive investment are extremely variable due

to previous history, individual variations, timing of reproduction, plant size, plant density,

competition, plant architecture, and successional stage.

Thus, the interaction of different life-history traits and plant size effects with reproductive

investment should be studied to understand the resource allocation strategy of plants across

elevation.

Conclusions

The revealed patterns of CMW and FD variations of floral traits have at least three important

points for plant ecological research. First, our study clearly demonstrates that in addition to

widely used vegetation traits, floral traits can also play a significant role in community

assembly process along environmental gradient. Therefore, these finding provide further

support to the recent argument that plant research should consider multiple traits representing

different ecological functions including growth, reproduction, and survival (E-Vojtkó et al.

2020; Laughlin, 2014; Rosbakh et al., 2021).

Second, our findings suggest that the floral traits might be used in applied ecological

research to forecast functional community composition shifts in alpine ecosystems. The

continuous temperature rise observed in the study region (Sabin et al., 2020) may lead to

relaxation of low-temperature stress filtering effects on flowering phenology, specific flower

area and floral display area of species adapted to environmental conditions of lower sites,

resulting in an altered composition of communities in cold sites (i.e., increase in frequency

and abundance of species from lower elevations). The results also revealed a lacuna in our

understanding of reproductive investment patterns of species and its interactions with biotic

and abiotic constraints along elevation, which might provide additional insights into the

factors governing community assembly of flowering plants.

Author contributions

MA, SR, SFB, HPS, DRB and SKU conceive the idea and design the study for this work.

MA, SRA, PS and SKU collected the data. MA, SRA, and PS data entry and did the formal

analyses. SR and MA analysed the data. MA and SR wrote the first version of the manuscript;

all authors contributed to writing of the final version of the manuscript.

Acknowledgements

Authors are grateful to Ministry of Environment, Forest & Climate Change, India, for

providing financial support under National Mission on Himalayan Studies. MA is thankful to

Council of Scientific and Industrial Research (CSIR), India, for providing senior research

fellowship. Authors express gratitude to Sanjay Kumar (Director, CSIR-IHBT) who provided

the mandatory facilities for the research work at CSIR-IHBT. We also acknowledge Christian

Körner and Robert R. Junker for insightful suggestions during the initial stages of project

proposal. We acknowledge Rohit, Om Prakash, Girja Nand, and Ashok Kumar for their

assistance in carrying out the fieldwork.

Data accessibility statement

Should be manuscript be accepted for publication, the data will be published on Zenodo.

References

Ahmad, M., Uniyal, S. K., Batish, D. R., Rathee, S., Sharma, P., & Singh, H. P. (2021).

Flower phenological events and duration pattern is influenced by temperature and

elevation in Dhauladhar mountain range of Lesser Himalaya. *Ecological Indicators*,

129, 107902. https://doi.org/10.1016/j.ecolind.2021.107902

Ahmad, M., Uniyal, S. K., Batish, D. R., Singh, H. P., Jaryan, V., Rathee, S., Sharma, P., &

Kohli, R. K. (2020). Patterns of plant communities along vertical gradient in

Dhauladhar Mountains in Lesser Himalayas in North-Western India. *Science of The*

Total Environment, *716*, 136919. https://doi.org/10.1016/j.scitotenv.2020.136919

Ahmad, M. (2021). Species composition and floral trait diversity along an altitudinal gradient

in the western Himalaya. Graduate thesis, Panjab University, Chandigarh.

Barrett, S. C. H. (2010). Understanding plant reproductive diversity. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *365*, 99–109.

https://doi:10.1098/rstb.2009.0199

Bellhouse, D.R. (2005). Systematic sampling methods. *Encyclopedia of Biostatistics*. John

Wiley & Sons, New York. https://doi.org/10.1002/0470011815.b2a16077

Bello, F. D., Lavorel, S., Lavergne, S., Albert, C. H., Boulangeat, I., Mazel, F., & Thuiller, W.

(2013). Hierarchical effects of environmental filters on the functional structure of plant

communities: a case study in the French Alps. *Ecography*, *36*(3), 393–402.

https://doi.org/10.1111/j.1600-0587.2012.07438.x

Bucher, S. F., & Römermann, C. (2020). Flowering patterns change along elevational

gradients and relate to life-history strategies in 29 herbaceous species. *Alpine Botany*,

130(1), 41–58. https://doi.org/10.1007/s00035-020-00231-w

Burkle, L. A., & Irwin, R. E. (2010). Beyond biomass: measuring the effects of

community☐level nitrogen enrichment on floral traits, pollinator visitation and plant

reproduction. *Journal of Ecology*, *98*(3), 705–717. https://doi.org/10.1111/j.1365-

2745.2010.01648.x

Campbell, D. R., & Halama, K. J. (1993). Resource and pollen limitations to lifetime seed

production in a natural plant population. *Ecology*, *74*(4), 1043–1051.

https://doi.org/10.2307/1940474

Conner, J. K., & Rush, S. (1996). Effects of flower size and number on pollinator visitation to

wild radish, *Raphanus raphanistrum. Oecologia*, *105*(4), 509–516.

https://doi:10.1007/BF00330014

Cordero, R. A., Fetcher, N., & Voltzow, J. (2007). Effects of wind on the allometry of two

species of plants in an elfin cloud forest. *Biotropica*, *39*(2), 177–185.

https://doi.org/10.1111/j.1744-7429.2006.00244.x

Crimmins, T. M., Crimmins, M. A., & Bertelsen, C. D. (2013). Spring and summer patterns in

flowering onset, duration, and constancy across a water☐limited gradient. *American*

Journal of Botany, *100*(6), 1137–1147. https://doi.org/10.3732/ajb.1200633

de Bello, F., Carmona, C. P., Dias, A. T., Götzenberger, L., Moretti, M., & Berg, M. P.

(2021). *Handbook of trait-based ecology: from theory to R tools*. Cambridge University

Press.

Dietrich, L., & Körner, C. (2014). Thermal imaging reveals massive heat accumulation in

flowers across a broad spectrum of alpine taxa. *Alpine Botany*, *124*(1), 27–35.

https://doi.org/10.1007/s00035-014-0123-1

Dyer, A. G., Whitney, H. M., Arnold, S. E., Glover, B. J., & Chittka, L. (2006). Bees

associate warmth with floral colour. *Nature*, *442*(7102), 525–525.

https://doi.org/10.1038/442525a

E-Vojtkó, A., de Bello, F., Durka, W., Kuehn, I., & Goetzenberger, L. (2020). The neglected

importance of floral traits in trait-based plant community assembly. *Journal of*

Vegetation Science, *31*(4), 529–539. https://doi.org/10.1111/jvs.12877

Fabbro, T., & Körner, C. (2004). Altitudinal differences in flower traits and reproductive

allocation. *Flora-Morphology, Distribution, Functional Ecology of Plants*, *199*(1), 70–

81. https://doi.org/10.1078/0367-2530-00128

Götzenberger, L., de Bello, F., Bråthen, K. A., Davison, J., Dubuis, A., Guisan, A., Lepš, J.,

Lindborg, R., Moora, M., Pärtel, M., Pellissier, L., Pottier, J., Vittoz, P., Zobel, K., &

Zobel, M. (2012). Ecological assembly rules in plant communities—approaches,

patterns and prospects. *Biological reviews*, *87*(1), 111–127.

https://doi.org/10.1111/j.1469-185X.2011.00187.x

Grime, J. P., Thompson, K., Hunt, R., Hodgson, J. G., Cornelissen, J. H. C., Rorison, I. H.,

Hendry, G. A. F., Ashenden, T. W., Askew, A. P., Band, S. R., Booth, R. E., Bossard,

C. C., Campbell, B. D., Cooper, J. E. L., Davison, A. W., Gupta, P. L., Hall, W., Hand,

D. W., Hannah, M. A., … Whitehouse, J. (1997). Integrated screening validates primary

axes of specialisation in plants. *Oikos*, *79*, 259–281. https://doi.org/10.2307/3546011

Grubb, P. J. (1977). The maintenance of species□richness in plant communities: the

importance of the regeneration niche. *Biological reviews*, *52*(1), 107–145.

https://doi.org/10.1111/j.1469-185X.1977.tb01347.x

Herrera, J. (2009). Visibility vs. biomass in flowers: exploring corolla allocation in

Mediterranean entomophilous plants. *Annals of Botany*, *103*(7), 1119–1127.

https://doi.org/10.1093/aob/mcp046

Jaswal A.K., Bhan S.C., Karandikar A.S, Gujar, M.K (2015). Seasonal and annual rainfall

trends in Himachal Pradesh during 1951-2005. *Mausam*, *66*(2), 247–264.

Jaswal, A. K., Kumar, N., & Khare, P. (2014). Climate variability in Dharamsala-a hill station

in western Himalayas. *Journal of Indian Geophysical Union*, *18*(3), 336–355.

Jiménez□Alfaro, B., Silveira, F. A., Fidelis, A., Poschlod, P., & Commander, L. E. (2016).

Seed germination traits can contribute better to plant community ecology. *Journal of*

Vegetation Science, *27*(3), 637–645. https://doi.org/10.1111/jvs.12375

Kattel, D. B., Yao, T., Yang, K., Tian, L., Yang, G., & Joswiak, D. (2013). Temperature lapse

rate in complex mountain terrain on the southern slope of the central Himalayas.

Theoretical and applied climatology, *113*(3), 671–682. https://doi.org/10.1007/s00704-

012-0816-6

Keddy, P. A. (1992). Assembly and response rules: two goals for predictive community

ecology. *Journal of vegetation science*, *3*(2), 157–164. https://doi.org/10.2307/3235676

Kiełtyk, P. (2021). Intraspecific morphological variation of *Bellidiastrum michelii*

(Asteraceae) along a 1,155 m elevation gradient in the Tatra Mountains. *PeerJ*, *9*,

e11286. https://doi.org/10.7717/peerj.11286

Klatt, S., Schinkel, C. C., Kirchheimer, B., Dullinger, S., & Hörandl, E. (2018). Effects of

cold treatments on fitness and mode of reproduction in the diploid and polyploid alpine

plant *Ranunculus kuepferi* (Ranunculaceae). *Annals of botany*, *121*(7), 1287–1298.

https://doi: 10.1093/aob/mcy017

Klotz, S., Kühn, I., & Durka, W. (2002). BIOFLOR–Eine Datenbank mit biologisch-

ökologischen Merkmalen zur Flora von Deutschland. – Bundesamt für Naturschutz,

Bonn – Bad Godesberg. https://wiki.ufz.de/biolflor/index.jsp

Körner, C. (2021). Alpine plant life: functional plant ecology of high mountain ecosystems

(3[rd] edistion). Springer Nature. https://doi:10.1007/978-3-642-18970-8

Koski, M. H., & Ashman, T. L. (2015). An altitudinal cline in UV floral pattern corresponds

with a behavioral change of a generalist pollinator assemblage. *Ecology*, *96*(12), 3343-

3353. https://doi.org/10.1890/15-0242.1

Kraft, N. J., Valencia, R., & Ackerly, D. D. (2008). Functional traits and niche-based tree

community assembly in an Amazonian Forest. *Science*, *322*(5901), 580–582.

https://doi:10.1126/science.1160662

Laiolo, P., & Obeso, J. R. (2017). Life-history responses to the altitudinal gradient. In J.

Catalan, Josep. M, Ninot & M. Mercè Aniz (eds.), *High Mountain conservation in a*

changing world (pp. 253–283). Springer Nature. https://doi.org/10.1007/978-3-319-

55982-7_11

Lambrecht, S. C., & Dawson, T. E. (2007). Correlated variation of floral and leaf traits along

a moisture availability gradient. *Oecologia*, *151*(4), 574–583.

https://doi.org/10.1007/s00442-006-0617-7

Laughlin, D. C. (2014). The intrinsic dimensionality of plant traits and its relevance to

community assembly. *Journal of Ecology*, *102*(1), 186–193.

https://doi.org/10.1111/1365-2745.12187

Makino, T. T., & Sakai, S. (2007). Experience changes pollinator responses to floral display

size: from size☐based to reward☐based foraging. *Functional Ecology*, *21*(5), 854–863.

https://doi.org/10.1111/j.1365-2435.2007.01293.x

McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community

ecology from functional traits. *Trends in ecology & evolution*, *21*(4), 178–185.

https://doi.org/10.1016/j.tree.2006.02.002

Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., ... & Zust, A. N. A.

(2006). European phenological response to climate change matches the warming

pattern. *Global change biology*, *12*(10), 1969–1976. https://doi.org/10.1111/j.1365-

2486.2006.01193.x

Norgate, M., Boyd-Gerny, S., Simonov, V., Rosa, M. G., Heard, T. A., & Dyer, A. G. (2010).

Ambient temperature influences Australian native stingless bee (*Trigona carbonaria*)

preference for warm nectar. *PLoS One*, *5*(8), e12000.

https://doi.org/10.1371/journal.pone.0012000

Obeso, J. R. (2002). The costs of reproduction in plants. *New phytologist*, *155*(3), 321–348.

https://doi.org/10.1046/j.1469-8137.2002.00477.x

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.

R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., &

Wagner, H. (2019). vegan: Community Ecology Package. R package version 2.5-6.

https://CRAN.R-project.org/package=vegan

Perez-Harguindeguy, N., Diaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., ... &

Cornelissen, J. H. C. (2016). Corrigendum to: new handbook for standardised

measurement of plant functional traits worldwide. *Australian Journal of botany*, *64*(8),

715–716. https://doi.org/10.1071/BT12225_CO

Pillar, V. D., Sabatini, F. M., Jandt, U., Camiz, S., & Bruelheide, H. (2021). Revealing the

functional traits linked to hidden environmental factors in community assembly.

Journal of Vegetation Science, *32*(1), e12976. https://doi.org/10.1111/jvs.12976

Poschlod, P., Abedi, M., Bartelheimer, M., Drobnik, J., Rosbakh, S. and Saatkamp, A. (2013)

6. Seed ecology and assembly rules in plant communities. In E. Maarel & J. Franklin

(eds.), *Vegetation Ecology* (2nd ed.) (pp. 164–202). John Wiley & Sons Ltd, United

Kingdom. https://doi.org/10.1002/9781118452592.ch6

Rabinowitz, D., Rapp, J. K., Sork, V. L., Rathcke, B. J., Reese, G. A., & Weaver, J. C. (1981).

Phenological properties of wind☐and insect☐pollinated prairie plants. *Ecology*, *62*(1),

49–56. https://doi.org/10.2307/1936667

Ram, J., Singh, S. P., & Singh, J. S. (1988). Community level phenology of grassland above

treeline in central Himalaya, India. *Arctic and Alpine Research*, *20*(3), 325–332.

https://doi:10.1080/00040851.1988.12002680

Rao, C. R. (1982). Diversity and dissimilarity coefficients: a unified approach. *Theoretical*

population biology, *21*(1), 24–43. https://doi.org/10.1016/0040-5809(82)90004-1

Rawat, D. S. (2012). Flowering phenology on altitudinal gradient in the Himalaya. *Current*

Science, *103*(11), 1281–1284.

Ricotta, C., & Moretti, M. (2011). CWM and Rao's quadratic diversity: a unified framework

for functional ecology. *Oecologia*, *167*(1), 181–188. https://doi.org/10.1007/s00442-

011-1965-5

Rosbakh, S., & Poschlod, P. (2021). Plant community persistence strategy is

elevation□specific. *Journal of Vegetation Science*, *32*(3), e13028.

https://doi.org/10.1111/jvs.13028

Rosbakh, S., Hartig, F., Sandanov, D. V., Bukharova, E. V., Miller, T. K., & Primack, R. B.

(2021). Siberian plants shift their phenology in response to climate change. *Global*

Change Biology, *27*(18), 4435–4448. https://doi.org/10.1111/gcb.15744

Rosbakh, S., Pacini, E., Nepi, M., & Poschlod, P. (2018). An unexplored side of regeneration

niche: seed quantity and quality are determined by the effect of temperature on pollen

performance. *Frontiers in Plant Science*, *9*, p1036.

https://doi.org/10.3389/fpls.2018.01036

Saatkamp, A., Cochrane, A., Commander, L., Guja, L. K., Jimenez□Alfaro, B., Larson, J., ...

& Walck, J. L. (2019). A research agenda for seed□trait functional ecology. *New*

Phytologist, *221*(4), 1764–1775. https://doi.org/10.1111/nph.15502

Sabin, T. P., Krishnan, R., Vellore, R., Priya, P., Borgaonkar, H. P., Singh, B. B., & Sagar, A.

(2020). In R. Krishnan, J. Sanjay, C. Gnanaseelan, M. Mujumdar & A. Kulkarni (eds.),

Climate change over the Himalayas. *Assessment of climate change over the Indian region* (pp. 207–222). Springer, Singapore.

Samson, D. A., & Werk, K. S. (1986). Size-dependent effects in the analysis of reproductive effort in plants. *The American Naturalist*, *127*(5), 667–680. https://doi.org/10.1086/284512

Sargent, R. D., & Ackerly, D. D. (2008). Plant–pollinator interactions and the assembly of plant communities. *Trends in Ecology & Evolution*, *23*(3), 123–130. https://doi.org/10.1016/j.tree.2007.11.003

Scherrer, D., & Körner, C. (2011). Topographically controlled thermal habitat differentiation buffers alpine plant diversity against climate warming. *Journal of Biogeography*, *38*(2), 406–416. https://doi.org/10.1111/j.1365-2699.2010.02407.x

Singh, J. S., & Singh, S. P. (1987). Forest vegetation of the Himalaya. *The Botanical Review*, *53*(1), 80-192. https://doi.org/10.1007/BF02858183

Soltis, P., & Soltis, D. (2014). Flower diversity and angiosperm diversification. In J. L. Riechmann & F. Wellmer (Eds.), *Flower development* (pp. 85–102). Humana Press.

Spasojevic, M. J., & Suding, K. N. (2012). Inferring community assembly mechanisms from functional diversity patterns: the importance of multiple assembly processes. *Journal of Ecology*, *100*(3), 652–661. https://doi.org/10.1111/j.1365-2745.2011.01945.x

Tachiki, Y., Iwasa, Y., & Satake, A. (2010). Pollinator coupling can induce synchronized flowering in different plant species. *Journal of Theoretical Biology*, *267*(2), 153–163. https://doi.org/10.1016/j.jtbi.2010.08.023

Trunschke, J., & Stöcklin, J. (2017). Plasticity of flower longevity in alpine plants is increased in populations from high elevation compared to low elevation populations. *Alpine Botany*, *127*(1), 41-51. https://doi.org/10.1007/s00035-016-0176-4

Ulrich, J., Bucher, S. F., Eisenhauer, N., Schmidt, A., Türke, M., Gebler, A., Barry, K., Lange, M. and Römermann, C. (2020). Invertebrate decline leads to shifts in plant

species abundance and phenology. *Frontiers in plant science*, *11*, p1410.

https://doi.org/10.3389/fpls.2020.542125

Valencia, E., Méndez, M., Saavedra, N., & Maestre, F. T. (2016). Plant size and leaf area

influence phenological and reproductive responses to warming in semiarid

Mediterranean species. *Perspectives in Plant Ecology, Evolution and Systematics*, *21*,

31–40. https://doi.org/10.1016/j.ppees.2016.05.003

van der Kooi, C. J., Kevan, P. G., & Koski, M. H. (2019). The thermal ecology of flowers.

Annals of Botany, *124*(3), 343–353. https://doi.org/10.1093/aob/mcz073

Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E.

(2007). Let the concept of trait be functional! *Oikos*, *116*(5), 882–892.

https://doi.org/10.1111/j.0030-1299.2007.15559.x

Wagner, J., Ladinig, U., Steinacher, G., & Larl, I. (2011). From the flower bud to the mature

seed: timing and dynamics of flower and seed development in high-mountain plants. In

C. Lütz (eds.), *Plants in alpine regions* (pp. 135–152). Springer Science & Business

Media. DOI:10.1007/978-3-7091-0136-0_10

Weiher, E., Freund, D., Bunton, T., Stefanski, A., Lee, T., & Bentivenga, S. (2011).

Advances, challenges and a developing synthesis of ecological community assembly

theory. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

366(1576), 2403–2413. https://doi.org/10.1098/rstb.2011.0056

Woodward, F. I., & Woodward, F. I. (1987). Climate and plant distribution. Cambridge

University Press.

Zhang, S., Liu, G., Cui, Q., Huang, Z., Ye, X., & Cornelissen, J. H. (2021). New field wind

manipulation methodology reveals adaptive responses of steppe plants to increased and

reduced wind speed. *Plant Methods*, *17*(1), 1–16. https://doi.org/10.1186/s13007-020-

00705-2

665 Zhao, Y. H., Ren, Z. X., Lázaro, A., Wang, H., Bernhardt, P., Li, H. D., & Li, D. Z. (2016).

666   Floral traits influence pollen vectors' choices in higher elevation communities in the

667   Himalaya-Hengduan Mountains. *BMC Ecology*, *16*(1), 1–8.

668   https://doi.org/10.1186/s12898-016-0080-1

669 Zhu, Y., Siegwolf, R. T., Durka, W., & Körner, C. (2010). Phylogenetically balanced

670   evidence for structural and carbon isotope responses in plants along elevational

671   gradients. *Oecologia*, *162*(4), 853–863. https://doi.org/10.1007/s00442-009-1515-6