Acinetobase: the comprehensive database and repository of *Acinetobacter* strains.

Adam Valcek[1, 2], James Collier[3], Alexander Botzki[3] and Charles Van der Henst[1, 2,#]

[#] Corresponding author: charles.vanderhenst@vub.vib.be

Affiliations

[1] Microbial Resistance and Drug Discovery, VIB-VUB Center for Structural Biology, VIB, Flanders Institute for Biotechnology, Brussels, Belgium.

[2] Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium.

[3] VIB Bioinformatics Core, Ghent, Belgium

Abstract

Background

Acinetobacter baumannii is one of the most problematic nosocomial pathogens that can efficiently thrive within hospital settings, mainly due to resistances towards antibiotics, desiccation, disinfectants, human serum and oxidative stress. Recently, increased resistance against last resort antibiotics earns this bacterium a highest priority concern classified by the CDC and the WHO. An obvious hallmark of this bacterium is the high heterogeneity observed amongst *A. baumannii* isolates, with a limited core genome. This feature complexifies the study of *A. baumannii* bacteria as an entity, subsequently reflected in a diversity of phenotypes of not only antimicrobial and environmental resistance but also virulence.

Main body

High degree of genome plasticity, along with the use of a limited subset of established reference strains, can lead to strain-specific observations, decreasing the global understanding of this pathogenic agent. Phenotypic variability of *A. baumannii* strains is easily observable such as with the microcolony morphologies, *in vitro* and *in vivo* virulence, production of different capsular polysaccharide structures and cellular densities. Some strains encode an extensive amount of virulence factors while other, including the reference strains, lack several key ones. The lack/excess of genes or specific physiological processes might interfere with *in vivo* and *in vitro* experiments, thus providing limited impact on the global understanding of *Acinetobacter* bacteria.

Conclusion

As an answer to the high heterogeneity amongst *A. baumannii* strains, we propose a first comprehensive database that includes the bacterial strains and the associated phenotypic and genetic data. This new repository, freely accessible to the entire community, allow selecting the best bacterial isolate(s) related to any biological question, using an efficient and fast exchange platform.

Keywords

Acinetobase, *Acinetobacter baumannii*, phenotypes, genotypes, database, repository

Background

Acinetobacter baumannii is an opportunistic nosocomial pathogen thriving in hospital environment and endangering especially critically ill patients [1]. This Gram-negative bacterium is part of the most problematic ESKAPE group of human pathogens, against which therapeutic approaches become limited [2]. Increased isolations of antibiotic-resistant strains, especially to last resort antibiotics, earns this bacterium the consideration as "urgent threat to public health" by the Centers for Disease Control and Prevention (CDC) [3] and as "a bacterium for which research and development of new antibiotics is critically needed" by the World Health Organization (WHO) [4]. An obvious hallmark of A. baumannii bacteria is the high heterogeneity observed amongst the isolates [5]. A. baumannii bacteria have a dynamic genome, with an estimated conserved core genome of only 16.5%, while 25% of the genome is unique to each strain, meaning that there is no counterpart in any other A. baumannii genome [5]. Our recent study (Philippe et al., 2022, unpublished) on 47 strains of A. baumannii shows even lower core genome of 2009 genes from the pangenome of 13611 genes, accounting for 14.76%. A. baumannii isolates are impressively variable, at both genetic and phenotypic levels. For example, the capsular polysaccharide, which impacts antibiotic and environmental resistances, host response and virulence, is encoded by at least 137 various capsule locus types, while the outer core of lipooligosaccharide by at least 13 locus types [6, 7]. This genetic variability translates into phenotypic heterogeneity, observable using electron microscopy where the thickness and rigidity of the capsule differs, along with different cellular densities [8]. The established reference strains such as ATCC19606, ATCC17978, DSM30011 or AB5075 are widely used type strains from which significant and validated observations were and are still generated [9–12]. All together, these reference strains greatly contributed to the state of the art of the current and growing A. baumannii field. The reference strains AB5075, ATCC17978 and ATCC19606 are multidrug resistant (MDR) strains of clinical origin [17, 18]

while DSM30011 is a non-MDR environmental strain obtained from plant microbiota [19].

The observations gained about these reference strains shed a strong base of the knowledge

about *A. baumannii*. However, in the other hand, type strains also showed the limitations of

only using these strains when current clinical isolates of *A. baumannii* show extreme genetic

and phenotypic variabilities. In order to fill this modern gap of knowledge, grasp the

complexity of *A. baumannii* and widen the possibilities of the community to use and efficiently

share genotypically and phenotypically characterized strains, we propose the database and

strain repository Acinetobase as a new community tool (Fig. 1). This new data and strains

sharing platform is especially important in the context of novel therapeutic discoveries, for

which conserved targets and antimicrobial mode of actions amongst the highest proportion of

problematic and clinically relevant *A. baumannii* isolates is required.

Figure 1: The view of the "About" page of Acinetobase.

84 Construction and content

This first version of Acinetobase consists of 43 clinical isolates and four sub-cultured reference

strains (AB5075-VUB, ATCC19606-VUB, ATCC17978-VUB and DSM30011-VUB) of *A.*

baumannii. All these strains where whole genome sequenced and the complete chromosomal

sequences of high quality are publicly available via GenBank accession number, where they

were deposited. In addition, the Acinetobase includes a variety of phenotypical characteristics

associated with a strain:

91   i, Quantification of the capsular polysaccharide using gradient density method for each

92   strain [12]

93   ii, Type of macrocolony produced for each strain (designated with letter A to E) with a

94   corresponding photograph

95   iii, Virulence in *Galleria mellonella* infection model for 9 strains [12]

96   iv, Transmission electron microscopy (TEM) photograph depicting the amount and

97   density of the capsule for 11 strains

98 The database also describes basic genotypical characteristics of each strain:

99   i, Multilocus sequence type (ST) according to Pasteur [20] and Oxford [21, 22] typing

100   scheme

101   ii, Capsule locus type (KL) [6, 19]

102   iii, Outer core lipooligosaccharide locus type (OCL) [6, 19]

The above mentioned genotypical and phenotypical traits can be used to search the database

and filter the strains for specific criteria. This way users can efficiently find the strain of desired

characteristics and request it.

The bacterial strains of *A. baumannii* were deposited and are available in lyophilized form as

per order from the Belgian Coordinated Collection of Microorganisms/Lab of Microbiology

(BCCM/LMG) (https://bccm.belspo.be/).

The example of an entry of AB5075-VUB is depicted in Fig. 2.

Figure 2: An entry of AB5075-VUB depicting A, description of the genotype (KL, $ST^{Ox/Pas}$, OCL) with a hyperlink for the complete chromosomal sequence and a reference publication; B, quantification of the capsule; C, macrocolony type; D, TEM micrograph of the capsule and E, virulence in *Galleria mellonella* infection model.

Utility and discussion

Existing databases

The decreasing costs of the whole-genome sequencing allowed genomic databases, filled with high-quality data, to thrive. There are well-established databases such as EnteroBase [20], which does not include data on *Acinetobacter*. In contrast, existing databases which provide whole-genome sequencing data on *A. baumannii*, such as BIGSdb [21], includes metadata and genotype, however, BIGSdb does not include phenotypic data unrelated to the genotype nor the strains themselves. Moreover, centralization of the laboratory providing strains with the genotype and phenotype characteristics will help to decrease the chance of working with subcultured variant which might have altered genotype and phenotype of the parental strain (Valcek et al., unpublished).

Browsing

The users can browse the "Collection" page for the strains (Fig. 3) and pick according to their genomic description. By clicking on the name of the selected strain the full entry with genomic and phenotypic details appears (Fig. 2).

Searching

The users can filter the entries (Fig. 4) according to each genetic and phenotypic-based criterium depicted in the "Collection" page. This feature helps to select specific strains of *A. baumannii* according to criteria demanded by the user, appropriated to their biological question.

Download

To avoid unnecessary redundancy, downloading the complete chromosomal sequences and the whole-genome assemblies is facilitated via GenBank (https://www.ncbi.nlm.nih.gov/). The source information on the strains can be found in the corresponding research articles found in the "Reference" column. By hyperlinking (using Digital Object Identifier - DOI) the corresponding peer-reviewed study, authors of the study provide the protocol on how the strains were obtained, their origin, original figures, and the protocols for assessing the phenotypical traits. The authors of the study will be directly responsible for the information and data the provided and submitted to Acinetobase.

Future perspectives

Acinetobase has a great potential to influence the field of microbiology, especially the *A. baumannii* community, but also other communities which work with similar pathogens (such as *Klebsiella pneumoniae*) and similar biological questions, might be inspired. Further

implementation potential of Acinetobase resides in additional experimental analyses which can

be easily and quickly performed and provide valuable information to researchers whether it is

a correct strain to work with (e.g., if the isolate is capable of natural transformation or if it

forms biofilm). The proposed project is so far aimed to *Acinetobacter baumannii*, yet not

limited to a single species of *Acinetobacter*. With an increasing clinical relevance of other

Acinetobacter species such as *A. pittii*, *A. nosocomialis*, *A. seifertti* and *A. dijkshoorniae* [22],

the project of Acinetobase is open to all *Acinetobacter* species. Acinetobase will be open to

submission of the phenotypic and genotypic data including strains from other laboratories and

research groups. This initiative will, over time, result in a growing database and a platform for

sharing knowledge and *Acinetobacter* spp. strains and might include further molecular tools

such as fluorescently labeled strains or characterized deletion mutant strains.

Figure 3: The "Collection" page depicting the name, Pasteur and Oxford sequence type, KL, OCL and

macrocolony types of each isolate in the collection.

Figure 4: An example of filtering the entries based on the number 3 in the name of the isolate.

Conclusion

In conclusion, Acinetobase is the first database dedicated to *Acinetobacter* genotypes and

associated phenotypes that also serves as a repository of strain. Acinetobase will not only be

the proof-of-concept of high heterogeneity among *A. baumannii* strains but will also benefit

the entire community by helping to select an appropriate strain related to specific conditions,

experiments, and biological questions.

177
178

180 ## Declarations

181 **Ethics approval and consent to participate**

182   Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and analyzed during the current study are linked from the Acinetobase website, http://acinetobase.vib.be/.

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by the Flanders Institute for Biotechnology (VIB).

Authors' contributions

AV and CVDH wrote up the manuscript. CVDH conceptualized the Acinetobase. JC wrote and designed the website, data storage, and update mechanisms. AB assisted with design of the website, is the project sponsor and reviewed the manuscript.

Acknowledgements

We would like to thank to Belgian Coordinated Collections of Microorganisms of Ghent University, Faculty of Sciences, Belgium. Our gratitude belongs to VIB Bioinformatics Core for their support in developing the website, especially to Christof De Bo who helped with graphics. We also would like to thank the National Reference Laboratory for Monitoring of Antimicrobial Resistance in Gram-negative Bacteria, CHU Mont-Godinne, Université

208 Catholique de Louvain (UCL), Yvoir, Belgium for sharing the modern clinical isolates of *A.*

209 *baumannii.*

References

1. Whiteway C, Breine A, Philippe C, van der Henst C. *Acinetobacter baumannii*. Trends in
Microbiology. 2022;30:199–200.
2. Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens.
https://doi.org/101586/eri1312. 2014;11:297–308.
3. CDC. Antibiotic Resistance Threats in the United States 2019. Atlanta, U.S. Department of
Health and Human Services C 2019. Antibiotic Resistance Threats in the United States, 2019.
https://doi.org/10.15620/cdc:82532.
4. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery,
research, and development of new antibiotics: the WHO priority list of antibiotic-resistant
bacteria and tuberculosis. www.thelancet.com/infection. 2018;18.
5. Imperi F, Antunes LCS, Blom J, Villa L, Iacono M, Visca P, et al. The genomics of
Acinetobacter baumannii: Insights into genome plasticity, antimicrobial resistance and
pathogenicity. IUBMB Life. 2011;63:1068–74.
6. Wyres KL, Cahill SM, Holt KE, Hall RM, Kenyon JJ. Identification of *Acinetobacter
baumannii* loci for capsular polysaccharide (KL) and lipooligosaccharide outer core (OCL)
synthesis in genome assemblies using curated reference databases compatible with Kaptive.
Microbial Genomics. 2020;6.
7. Kenyon JJ, Hall RM. Updated Analysis of the Surface Carbohydrate Gene Clusters in a
Diverse Panel of *Acinetobacter baumannii* Isolates. Antimicrobial Agents and Chemotherapy.
2021;66.
8. Whiteway C, Valcek A, Philippe C, Strazisar M, de Pooter T, Mateus I, et al. Scarless
excision of an insertion sequence restores capsule production and virulence in *Acinetobacter
baumannii*. The ISME Journal 2022. 2021;:1–5.
9. Roussin M, Rabarioelina S, Cluzeau L, Cayron J, Lesterlin C, Salcedo SP, et al.
Identification of a Contact-Dependent Growth Inhibition (CDI) System That Reduces Biofilm
Formation and Host Cell Adhesion of *Acinetobacter baumannii* DSM30011 Strain. Frontiers
in Microbiology. 2019;10 OCT:2450.
10. Bravo Z, Orruño M, Parada C, Kaberdin VR, Barcina I, Arana I. The long-term survival of
Acinetobacter baumannii ATCC 19606T under nutrient-deprived conditions does not require
the entry into the viable but non-culturable state. Archives of Microbiology. 2016;198:399–
407.
11. Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, et al. AB5075,
a highly virulent isolate of *Acinetobacter baumannii*, as a model strain for the evaluation of
pathogenesis and antimicrobial treatments. mBio. 2014;5.
12. de Silva PM, Chong P, Fernando DM, Westmacott G, Kumar A. Effect of Incubation
Temperature on Antibiotic Resistance and Virulence Factors of *Acinetobacter baumannii*
ATCC 17978. 2017. https://doi.org/10.1128/AAC.01514-17.

13. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, et al. Resources for
genetic and genomic analysis of emerging pathogen *Acinetobacter baumannii*. Journal of
Bacteriology. 2015;197:2027–35.

14. Harding CM, Hennon SW, Feldman MF. Uncovering the mechanisms of *Acinetobacter*
baumannii virulence. Nature Reviews Microbiology 2017 16:2. 2017;16:91–102.

15. Repizo GD, Viale AM, Borges V, Cameranesi MM, Taib N, Espariz M, et al. The
Environmental *Acinetobacter baumannii* Isolate DSM30011 Reveals Clues into the
Preantibiotic Era Genome Diversity, Virulence Potential, and Niche Range of a Predominant
Nosocomial Pathogen. Genome Biology and Evolution. 2017;9:2292–307.

16. Diancourt L, Passet V, Nemec A, Dijkshoorn L, Brisse S. The Population Structure of
Acinetobacter baumannii: Expanding Multiresistant Clones from an Ancestral Susceptible
Genetic Pool. PLOS ONE. 2010;5:e10034.

17. Bartual SG, Seifert H, Hippler C, Luzon MAD, Wisplinghoff H, Rodríguez-Valera F.
Development of a multilocus sequence typing scheme for characterization of clinical isolates
of *Acinetobacter baumannii*. Journal of Clinical Microbiology. 2005;43:4382–90.

18. Wisplinghoff H, Hippler C, Bartual SG, Haefs C, Stefanik D, Higgins PG, et al. Molecular
epidemiology of clinical *Acinetobacter baumannii* and *Acinetobacter* genomic species 13TU
isolates using a multilocus sequencing typing scheme. Clinical Microbiology and Infection.
2008;14:708–15.

19. Wick RR, Heinz E, Holt KE, Wyres KL, Diekema DJ. Kaptive Web: User-Friendly Capsule
and Lipopolysaccharide Serotype Prediction for *Klebsiella* Genomes. jcm.asm.org 1 Journal of
Clinical Microbiology. 2018;56:197–215.

20. Zhou Z, Alikhan N-F, Mohamed K, Fan Y, Group the AS, Achtman M, et al. The
EnteroBase user's guide, with case studies on *Salmonella* transmissions, *Yersinia pestis*
phylogeny, and *Escherichia* core genomic diversity. Genome Research. 2020;30:138–52.

21. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb
software, the PubMLST.org website and their applications. Wellcome Open Research. 2018;3.

22. Vijayakumar S, Biswas I, Veeraraghavan B. Accurate identification of clinically important
Acinetobacter spp.: an update. Future Sci OA. 2019;5:395–2056.

281