A detailed map of mitochondrial repeat-mediated recombination in Arabidopsis *msh1* mutants using long-read sequencing

Yi Zou[1], Weidong Zhu[1], Daniel B. Sloan[2], and Zhiqiang Wu[1,*]

[1]Guangdong Laboratory for Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of Agriculture, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, Guangdong 518120, China, and

[2]Department of Biology, Colorado State University, Fort Collins, CO 80523

* For Correspondence (e-mail: wuzhiqiang@caas.cn; tel.: +86 135-3040-6763).

Summary

The abundant repeats in plant mitochondrial genomes can cause rapid genome rearrangements and

are also a major obstacle in short-read sequencing studies. Nuclear-encoded proteins such as MSH1

are known to suppress the generation of repeat-associated mitochondrial genome variants, but our

understanding of these mechanisms has been constrained by the limitations of short-read

technologies. Here, we used highly accurate long-read sequencing (PacBio HiFi) to characterize

mitochondrial and plastid genome variants in *Arabidopsis thaliana msh1* mutant individuals. The

HiFi reads successfully quantified parental and crossover recombination products of both large and

small repeats, providing a global view of recombination dynamics. We found that recombination

breakpoints were distributed relatively evenly across the length of repeated sequences and observed

extensive non-crossover recombination (gene conversion) between pairs of imperfect repeats in the

mitochondrial genome of *msh1* mutants. Comparison of long-read assemblies of seven other

Arabidopsis thaliana accessions showed that mitochondrial genome structural divergence between

accessions paralleled the repeat-mediated dynamics observed in *msh1* mutants. The *Arabidopsis*

plastid genome generally lacks small repeats and exhibited a very different pattern of variant

accumulation in *msh1* mutants compared with the mitochondrial genome. Our data illustrate the

power of HiFi technology in studying repeat-mediated recombination in plant mitochondrial

genomes and improved the sequence resolution for recombinational processes suppressed by

MSH1.

Keywords: *Arabidopsis thaliana*, mitochondrial genome, repeat recombination, gene conversion,

long-read

Significance

Plant mitochondrial genomes undergo rapid rearrangements while maintaining protein genes.

Long-read sequencing provides a detailed and quantitative view of mitochondrial repeat-mediated

recombination suppressed by MSH1, advancing our understanding of plant mitochondrial genome

dynamics.

Introduction

Mitochondria and plastids are major energy converters as well as central metabolic compartments

of plant cells. Plant mitochondrial (mt) and plastid (pt) genomes can also be the target of genetic

engineering to alter phenotypes that are responsive to environmental change (Flood et al., 2020).

These two organelle genomes have followed quite different evolutionary trajectories. Compared

with the rather conserved pt genomes (Lee et al., 2016), mt genomes exhibit dramatic variation

among plant species (Wu et al., 2020a). Although coding sequences in angiosperm mt genome have

very low substitution rates, their gene order and intergenic sequences are rarely conserved (Wolfe

et al., 1987; Palmer and Herbon 1988; Christensen, 2021). Angiosperm mt genomes typically

contain numerous non-tandem repeats, and recombination between different copies of repeat

sequences causes a high level of structural instability and variation (Wynn and Christensen, 2019).

These repeats are often grouped based on size ranges with some functional considerations. Large

repeats ($\geq$1000 bp) are usually identical in sequence and recombine at high frequency, leading to

isomerization of the genome (Unseld et al., 1997). The recombination frequencies of intermediate-

sized repeats (50-1000 bp) are lower but sensitive to genetic, environmental and developmental

triggers that make them frequent enough to have important phenotypic effects (Gualberto and

Newton 2017; Mackenzie and Kundariya, 2019). Shorter repeats (<50 bp) typically do not show

recombination activity. These short sequences are often described as "microhomologies", but the

very shortest of them (e.g., <10 bp) are unlikely to reflect true homology, because such similarities

can easily occur at random. Despite their importance and critical roles, repeats in mt genome

assemblies are not always annotated correctly or completely even in model plant species (Wynn

and Christensen, 2019). Moreover, these repeats may produce complex assembly graphs, which

hinder a clear characterization of mt genomes (Masutani et al., 2021). Thus, mechanisms

underlining the repeats and their roles in destabilizing the mt genomes require a more thorough

exploration.

Recombination in plant mt genome is likely involved in repairing DNA damage, such as double-

strand breaks (DSBs), which can disrupt genome integrity and prevent the progression of

replication and transcription activities. To deal with DSBs, two main branches of repair systems

exist, which rely on different sets of genes: homology-directed repair (HDR) and non-homologous

end joining (NHEJ) (Hussmann et al., 2021). HDR requires homologous dsDNA templates for new

strand synthesis. HDR involving both DSB ends will lead to non-crossover repair products, which

is a process known as gene conversion. In contrast, HDR involving only one DSB end and non-

allelic templates is known as break-induced repair (BIR) and leads to crossover products

(Christensen, 2013).

Nuclear genes encode a set of mitochondrial- and plastid-targeted DNA binding proteins, which

are actively involved in processes such as DNA recombination, replication, and repair (RRR;

Gualberto and Newton, 2017; Chevignv et al., 2020; Fuchs et al., 2020). Some RRR genes, such as

RECA, *POLIA/B*, *SSB*, have conserved roles, while others are evolutionary innovations. Unlike

bacterial MutS counterparts, the dual-targeted (mitochondrial and plastid) MSH1 enzyme in plants

combines a mismatch-recognition domain (MutS I) and a predicted DSB-generating GIY-YIG endonuclease domain in a single protein. MSH1 has been hypothesized to participate in novel DNA repair mechanisms in plant organelles (Christensen, 2013; Ayala-Garcia et al., 2018; Fukui et al, 2018; Wu et al., 2020b; Wynn et al., 2020).

Arabidopsis *msh1* (=*chm*) mutants exhibited a series of phenotypes and were originally isolated based on green-white leaf variegation phenotype (Rédei, 1973). Positional cloning identified this locus to encode a plant MutS homolog with dual localization (Abdelnoor et al., 2003). Based on DNA gel blots of dissected white tissues from hemi-complementation plants, leaf variegation was attributed to pt genome rearrangements (Xu et al., 2011). The plastid hemi-complementation plants of *msh1* exhibited curly or wrinkled leaf and slower seed germination, which were thought to be caused by mitochondrial genetic effects (Virdi et al., 2015, 2016). More recently, heritable epigenetic effects in the nucleus were shown to cause diverse growth-related phenotypes of some *msh1* individuals (Virdi et al., 2015; Yang et al., 2020; Kundariya et al., 2020). Disruption of *msh1* results in ectopic recombination of intermediate-sized repeats and sub-stoichiometric shifting of different genomic segments (Abdelnoor et al., 2003; Shedge et al., 2007; Arrieta-Montiel et al., 2009; Davila et al; 2011). Similar results have been observed in moss organelle genomes, implying that these are anciently conserved functions of *MSH1* in land plants (Odahara et al., 2017). It was also reported that lower *MSH1* expression caused mitochondrial rearrangements that produced cytoplasmic male sterile (CMS) lines in tomato and allowed fertility reversion of lines of *Brassica juncea* (Sandhu et al., 2007; Zhao et al., 2016, 2021). These studies demonstrated that MSH1 maintains the structural integrity of the mt genome through suppressing illegitimate recombination at repeats. We recently showed increases in de novo point mutations in organelle genomes of *msh1* mutants by duplex sequencing and droplet digital PCR, suggesting a role of MSH1 in maintaining the lower organellar synonymous substitution rates compared with the nuclear genome (Wu et al., 2020b).

Despite this extensive work on *MSH1* in Arabidopsis, we still lack a complete understanding of its effect on repeat dynamics, in part due to the technical difficulties of traditional DNA gel blots, quantitative PCR, and short-read sequencing. For example, a previous study used early-generation Illumina technology with 35-bp reads to detect variants in the Arabidopsis *msh1* mutants (Davila et al. 2011). This work was successful in detecting repeat-mediated rearrangements, but the short reads available at the time lack resolution for analyzing repeats at a detailed sequence level across their entire lengths. The advent of long-read sequencing technologies has created the opportunity to analyze structural variants in plant mt genome assemblies (Kozik et al., 2019; Yu et al., 2022).

135  In this work, we use PacBio high-fidelity (HiFi) sequencing to comprehensively investigate
136 mitochondrial repeat-mediated recombination in individual Arabidopsis *msh1* plants. These highly
137 accurate long reads are enough to span the entire length of repeats, including the two largest repeats
138 in the Arabidopsis mt genome (6.5 kb and 4.2 kb), enabling unambiguous resolution of structural
139 rearrangements and gene conversion events. We first developed a pipeline to characterize different
140 types of rearrangements based on blastn alignments of HiFi reads to the reference genome. We
141 were able to identify rearrangements mediated by 99 repeat groups ($\geq$ 50 bp), of which 53 groups
142 were not shown recombination-active previously. Notably, we observed frequent gene conversion
143 between a proportion of non-identical repeats in *msh1*, an effect that can be difficult to detect and
144 quantify with earlier short-read sequencing approaches. We found that a dynamic region containing
145 *atp6-2* was consistently lost in mt genomes of 7 other *Arabidopsis thaliana* accessions and loss of
146 *atp6-1* was correlated with phenotypic variation among *msh1* individuals. We also detected novel
147 pt genome variants with HiFi reads. These results advance our understanding of plant mt genome
148 evolution and lay the foundation for engineering mitotype-associated plant phenotypes in the future.
149

150 **Results**

151 **Mapping of HiFi reads identified structural variations in Arabidopsis organelle genomes**

152 We generated F_3 families from a cross between wild type Arabidopsis Columbia-0 (Col-0) and an
153 *msh1* mutant line (*chm1-1*, CS3372) in an earlier work (Wu et al., 2020b). To analyze structural
154 rearrangements in the organelle genomes in homozygous *msh1* mutants, we collected aboveground
155 tissue from 11 individual plants from 3 different F_3 families (Table S1) and prepared separate DNA
156 libraries for each plant for PacBio HiFi sequencing. The libraries generated a total of 34.36 Gb of
157 HiFi reads (12.7-18.1 kb mean read length). These reads were first mapped to the Col-0 reference
158 genome by minimap2 (Li, 2018) to separate nuclear, pt and mt genome-mapped proportions. After
159 excluding one outlier with very low yield, the coverage of the nuclear genome ranged from 8.5$\times$ to
160 46.4$\times$, the coverage of the pt genome ranged from 899.0$\times$ to 3093.8$\times$, and the coverage of the mt
161 genome ranged from 30.9$\times$ to 247.9$\times$ (Table S2). The plastid (4.96%-12.85%) and mitochondrial
162 (0.93%-2.13%) proportions were used for downstream analysis (Figure 1a; Table S3 and S4).

163  The published mt genome is usually represented as a linear sequence of the "master circle".
164 However, various sub-stoichiometric isoforms exist due to frequent repeat-mediated structural
165 rearrangements (Kozik et al., 2019). Potential structural rearrangements could be indicated by reads
166 whose parts were mapped to discontinuous positions in the reference genome (Figure 1b). We
167 developed a pipeline to detect mitochondrial structural rearrangements (available at
168 https://github.com/zouyinstein/hifisr). In this pipeline, mt genome-mapped reads were aligned

against the reference sequence (NC_037304.1; Sloan et al., 2018) via the NCBI blastn software.
Raw blastn results were then processed by custom Python scripts, which can group the reads by
discontinuity, homologous sequence size and mapping coordinates, and calculate read counts for
each group (Table S5). After additional manual checking and curation, we were able to distinguish
zero-rearrangement events and one-, two-, three-, and four-rearrangement events mediated by
repeats (≥50 bp) in a single read. A few low-frequency events were also detected, including
microhomology-mediated end joining (MMEJ; defined as occurring at repeats between 2 and 49
bp in length) and non-homologous end-joining (NHEJ; defined as occurring at breakpoints sharing
0 or only 1 bp of sequence identity), replication slippage associated with tandem repeats, and
insertions (Figure 1c; Table S7). More than 90% of the identified mt genome rearrangements were
crossover recombination (BIR) mediated by non-tandem repeats. In wild type, 7.50% of total reads
accounted for rearrangements mediated by non-tandem repeats, most of which were associated with
the two large repeat pairs that undergo high-frequency recombination. In *msh1* mutants, this ratio
increased to 31.63% of total reads, with an increased frequency of rearrangement events mediated
by intermediate-sized repeats (Figure 1c,d; Table S9). Much lower frequencies of structural
variants were found in the pt genome for both wild-type and *msh1* individuals, but we still identified
rearrangements mediated by the larger inverted repeats (IRa and IRb), MMEJ, NHEJ, replication
slippage associated with tandem repeats, and insertions (Figure 1e,f; Table S6, S8 and S10). In
contrast to the mt genome, the Arabidopsis pt genome contains very few intermediate repeats
(Table S11). Only one read supported rearrangement-mediated by a pair of 123-bp imperfect
repeats in pt genome. And because of the large size (~26 kb), rearrangements mediated by the
IRa/b repeats will be underestimated because few HiFi reads span an entire repeat copy. The
proportions of MMEJ were low in mt and pt genomes, but both were increased in *msh1* mutant
compared with wild-type (Figure 1c,e). Taken together, these results demonstrate the power of
long-read HiFi sequencing for characterizing the increased repeat-mediated structural variants in
the mt genome of *msh1* mutants.
195
Mitochondrial repeat annotations and recombination activity
To further check whether the identified rearrangement events were indeed caused by the repeated
sequence elements of the mt genome, we first visualized the global patterns of the junctions and
relative read counts of each group of rearrangements, including both one-rearrangement events and
two-rearrangement events in a single read (Figure 2a-d; Table S12 and S13). These rearrangements
caused multiple highly dynamic regions across the mt genome, which were most severe at the
positions of large repeats and several repeat-dense regions (Figure 2e). However, we found that

more than half of these rearrangement junctions were not associated with the subset of repeats that

were named and annotated in previous studies (note that annotations of repeats FF, WW, XX, YY,

ZZ among the mentioned repeats A-Z, AA-ZZ are missing in GenBank file NC_037304.1; Arrieta-

Montiel et al., 2009; Davila et al., 2011; Sloan et al., 2018). It was suggested in a recent study that

identifying non-tandem repeats by blastn is incomplete at default parameters and can be

inconsistent at repeat boundaries, especially when mismatches and gaps are abundant (Wynn and

Christensen, 2019). Thus, we decided to perform a blastn search with the lowest word size using

the mitochondrial reference sequence as both query and subject, which generated more candidate

coordinates of repeat boundaries for manual comparison with the junction coordinates of one-

rearrangement HiFi reads. In total, we found 71 groups of two-copy repeats (R2), 23 groups of

three-copy repeats (R3), 5 groups of four-copy repeats (R4), which accounted for most of the read

rearrangement junctions in *msh1* mt genome (Figure S1a; Table S14, S15, S16 and S17). These 99

repeat groups contained all previously reported active repeats except for repeat TT and 53

additional groups of recombinationally active repeats ($\geq$ 50 bp; named HNR1 to HNR53;

annotations of repeats FF, WW, XX, YY, ZZ that are missing in NC_037304.1 were not

considered). The boundaries of some low-identity repeats in our annotation differed from previous

annotations. For example, the direct repeat pair MMJS-1 (767 bp) and MMJS-2 (692 bp) were

previously annotated as separate smaller repeats MM, J, and S. We treated short repeats as part of

the microhomology category (2 to 49 bp), because the blastn software can often generate

overlapping, inaccurate, or incomplete predictions of homology below 50 bp in size. Moreover, the

orientation of a repeat pair determines how they can reorganize the genome. In a hypothetical circle

containing only one repeat pair, crossover causes the interconversion between a large circle and

two smaller circles if they are in direct orientation, while causing the interconversion between two

isomeric circles of the same length if the repeat pair is inverted (Figure S2a). But in real mt genome

with multiple repeat pairs and more complex *in vivo* structures, the orientation of a given repeat

pair can change depending on recombination events between other repeats. Thus, the current

annotations of direct vs. inverted repeat orientations are arbitrary depending on the linear

representation of the reference "master circle" (Table S15, S16 and S17).

The relative recombination frequency of different repeats can be estimated by normalized read

counts of one-rearrangement events. Thus, we further grouped these repeats into high-,

intermediate-, and low-activity categories, with 18 groups showing high activity (read count $\geq$ 50

in *msh1*, normalized to 10000 total mt genome-mapped reads), 17 groups showing intermediate

activity (read count of 10 to 49 in *msh1*) and 64 remaining groups showing low activity (Figure

S1b; Table S18, S19 and S20). Previous studies have suggested a positive correlation between

homology length and recombination frequency of different repeat pairs (Davila et al., 2011). To

investigate this idea, we considered all two-copy repeats with high sequence identity (mismatch +

gap ≤ 2) to exclude the effects of reduced sequence identity or copy number. As indicated by HiFi

read counts, there was indeed a positive relationship between homology length and recombination

frequency (Figure S1c). Some repeats fell far from the regression line, suggesting that additional

factors contribute to recombination frequency, which might explain the different recombination

frequencies for different pairs within the same three-copy repeat group (Table S19). Asymmetrical

recombination or simply preferential accumulation of only one of the crossover products of

intermediate-sized repeats in *msh1* mutant (Arrieta-Montiel et al., 2009, Davila et al., 2011).

Similarly, HiFi reads could also show asymmetrical recombination with conserved polarity but

different intensities among samples and accompanying loss of parental forms for some repeat

groups (Figure S2b; Table S21). Notably, owing to the long read length, HiFi reads also indicated

asymmetrical recombination meditated by the two large repeat pairs, which was more pronounced

in *msh1* mutant than in wild-type (Figure S2b). HiFi reads also identified recombination mediated

by intermediate-sized repeats Q, V, B, F, and a few other repeats at very low frequency in wild-

type (Table S21), which were not detected by less sensitive gel blot experiments previously.

Together, with HiFi reads we confirmed mt genome rearrangements in *msh1* were mostly meditated

by repeated sequence elements of various sizes and activities, and that these rearrangements could

provide additional clues for manual curations of repeat annotations.

Base-level understanding of crossover and non-crossover recombination events in *msh1*

The power of long reads with high accuracy enabled unambiguous mapping across mitochondrial

repeats, which is crucial to investigate the detailed behavior of recombination events. For crossover

recombination events within perfectly matched regions, the actual recombination breakpoints could

not be determined from the product sequence. Among the 18 high-activity and 17 intermediate-

activity repeat groups, 8 groups were 100% identical. For the remaining 27 groups of non-identical

repeats, we could examine the single-nucleotide variants (SNVs) and indels that distinguish the

original repeat copies within one-rearrangement chimeric reads and gain a base-level understanding

of crossover site distribution. For example, repeat group CC showed multiple different

recombination breakpoints distributed along the length of the repeats (Figure 3a). The proportion

of reads with a recombination breakpoint between two SNVs/indels was positively correlated with

the distance between those two variants as a proportion of the total length of the repeat, based on

the analysis of the longest identical fragment in each group of two-copy repeats (Figure 3b; Table

S22).

We also found reads with clusters of SNVs/indels (Table S22) that could be explained by short-
range or partial gene conversion events (Davila et al., 2011). Particularly, in the crossover products
of repeat H-1 and H-3, we observed a highly discontinuous pattern of SNVs indicating multiple
short-range gene conversion events (Figure 3c). A proportion of the one-rearrangement reads
mediated by repeat H-1 and repeat H-3 also contained variants derived from repeat H-2, which
suggested that gene conversion could happen using other homologous dsDNA templates not
directly involved in a crossover recombination event (Figure 3c). To get a detailed view of non-
allelic gene conversion events, we calculated the ratios of gene conversion sites in zero-
rearrangements reads spanning the entire length of 12 high-activity repeat groups and 15
intermediate-activity repeat groups, and indeed observed reads containing SNVs/indels-encoded
by non-allelic repeat homologs in *msh1* mutants, which seldomly occurred in wild-type (Figure 4a;
Table S23, S24, and S25). In *msh1* mutants, repeat groups A, D, G, H (SNVs/indels in E-1 and E-
2 were the same ones in H-2 and H-3), I, M, MMJS, N, R, T, and W showed a high frequency of
non-allelic gene conversion and had higher frequencies of crossover recombination (Figure 4b,c
and S3; Table S21), while the remaining 15 analyzed repeats groups showed a low ratio or no non-
allelic gene conversion (Table S23). Within repeat H-2, SNVs/indels with a higher ratio of gene
conversion were not the ones shared by both H-1 and H-3. Instead, they were the unique variants
located near the H-1 left boundary and the H-3 right boundary (Figure 4b and S4). In repeat
homolog MMJS-2 the unusually high ratios of gene conversion in some individuals were associated
with the near complete loss of parental forms, i.e., very low sequence coverage in this region
(Figure 4c, S2b, and S5). Together, these base-level investigations of HiFi reads indicated that in
addition to suppressing ectopic crossover recombination, *MSH1* also suppresses non-crossover
gene conversion between imperfect repeats of the mt genome. Although less likely, we cannot
exclude the possibility that multiple rounds of crossover recombination might lead to the same
effect.
296

Phenotypic correlations of mt genome variants in *msh1* mutant

It was established by hemi-complementation experiments that the slower germination phenotype
of *msh1* mutants was caused by mitochondrial genetic effects (Virdi et al., 2015). For easy
quantification, we used primary root length of seedlings at 5 days after sowing (DAS) to indirectly
reflect seed germination speed. We observed shorter primary root lengths of *msh1* seedlings at 5
DAS compared with wild-type, which might indicate delayed seed germination (Figure 5a,b; Table
S26). The three *msh1* mutant F_3 families also exhibit quite different patterns of root length
distribution. For example, M3-17 contained both fast- and slow-germinating individuals and M3-

43 contained mostly slow-germinating individuals (Figure 5b). To further check if this slow-
germinating phenotype was heritable, we analyzed root length distribution patterns of F_4 families.
As expected, the progeny of fast-germinating M3-17 F_3 individuals germinated faster than the
progeny of slow-germinating M3-17 F_3 individuals and M3-43 F_3 individuals (Figure S6). The
overall HiFi read coverage of different *msh1* individuals did not vary greatly compared with wild-
type, except for some fluctuations at sporadic stretches (Figure S7; Table S27). Most of these low-
coverage stretches contained no coding sequences, but some overlapped with functional genes,
including tRNA genes and multiple open reading frames (ORFs). For example, the region around
repeat Large1-2 showed the lowest coverage in all *msh1* individuals, which would cause a partial
loss of *atp6-2*, *atp9*, *trnS* and *ORF262* (Figure 5d,e). We reasoned that if any mitochondrial
genomic feature caused the slow-germinating phenotype, it should occur at a substantial ratio and
correlate with these root length distributions of different F_3 families. We found the region between
the boundaries of repeat V-1 and Large1-1 was most likely to meet these criteria because there was
a larger reduction in *atp6-1* copy number in M3-43 individuals, which also had the shortest root
lengths (Figure 5c). Interestingly, it has been reported that *atp6-2* knockouts could be more easily
obtained than *atp6-1* knockouts with mitoTALENs (Arimura et al., 2020). Therefore, we speculate
that the different degrees of sub-stoichiometric shifting at the *atp6-1* gene may contribute to
germination phenotypes in *msh1* mutants, but this hypothesis will need to be tested with more direct
genetic manipulations.

Mt genomes of other accessions showed low-coverage regions similar to *msh1* mutants
We also re-assembled mt genomes for our samples using HiFi reads and metaFlye, which is a fast
long-read assembler for bacterial metagenomes (Kolmogorov et al., 2020). The metaFlye
assemblies of mt genomes were much more complex with fragmented contigs and many unresolved
repeats in *msh1* mutants compared with wild-type, reflecting the highly rearranged and
heteroplasmic mt genome (Figure S8). When assembling wild-type mt genome from published
HiFi datasets (Wang et al., 2021; Naish et al., 2021), the information provided by assembly graphs
was less complete, which suggested that our mapping-based analysis had advantages in detecting
alternative structures and mt genome heteroplasmy (Figure S9; Table S28 and S29).
One recent study used a method combining Flye (not the metagenomic version) and custom
clustering algorithms to study mt genome heteroplasmy using the PacBio continuous long-read
(CLR) datasets of Col-0 and seven other *Arabidopsis thaliana* accessions (Jiao and Schneeberger,
2020; Masutani et al., 2021). We analyzed these same datasets with our mapping-based pipeline.
The resulting metaFlye assembly graphs were completely resolved for Kyoto and Sha. Only large

repeats remained unresolved for other accessions, which suggests low mt genome heteroplasmy in
all accessions, similar to observations in wild-type Col-0 (Figure S10; Table S30 and S31).
Although rearrangements of large repeats could be identified, the blastn alignments of corrected
reads to each pseudo-master circle were less continuous than the more accurate HiFi datasets (Table
S32-S39).
The mt genomes of other accessions showed clear differences in size compared to Col-0 (Table
S40-S42). For example, the size difference between the largest Eri-1 mt genome and Col-0 was
mainly caused by a 1.1 kb insertion (Figure S11). This sequence was present in all other accessions,
suggesting Col-0 specific deletion (Figure S12-S17). The mt genome arrangement of C24 was
closely related to Eri-1, except for a 1.8 kb deletion (Figure S12). These insertions and deletions
were often located within large syntenic regions containing multi-copy genes or no functional genes.
We noticed that the region containing the repeat copy Large1-2 (258638 bp to 265227 bp in Col-0
mt genome) was lost in other accessions except Col-0, which were coincident with the low-
coverage or repeat-mediated highly dynamic regions observed in *msh1* mutants (Figure 6a-c). For
Eri-1 and C24, the region from repeat V-1 to Large1-1 replaced the deleted region near Large1-2,
which formed a pair of 7.1 kb large repeats. But for accessions Cvi-0, An-1, L*er*, Kyoto and Sha,
this deletion caused the loss of one copy of the *trnS* and *atp6-2* genes (Figure 6b, S13-S17). In
particular, the smallest mt genome of Sha accession also had an additional large deletion, which
caused the loss of one copy of *trnK* and *rps19* in repeat A-2 (Figure 6d,e). In an earlier report based
on a 5.8-kb fragment (EMBL: HF543671), loss of *trnK* was experimentally verified and a 400-bp
insertion containing *orf117Sha* was thought to be correlated with the CMS phenotype of Sha
(Gobron et al., 2013). That study also reported the presence of the 1025-bp large repeats near repeat
R-1, which exhibited double coverage and were recombinationally active based on our analysis
(Figure S17; Table S42). Thus, repeat sequences caused fast intraspecies evolution of mt genomes.
Our analysis confirmed previous findings that evolutionary changes of Arabidopsis mt genomes
were constrained by gene functions with changes that are similar rearrangements observed in *msh1*
mutants but a low-heteroplasmy final state (Arrieta-Montiel et al., 2009; Davila et al., 2011).
366
Different patterns of structural variants and mutations in *msh1* pt genome
Loss of MSH1 function in mitochondria and plastids caused similar types of variants, but with quite
different proportions (Figure 1c,e). MMEJ and NHEJ events were detected at very low frequencies
in HiFi reads. They were minor proportions of mt genome rearrangements (Table S7). But with the
higher depth of pt genome-mapped reads, we could observe distinctive patterns of rearrangements
in *msh1* mutants. First, crossover recombination events mediated by short partial palindromic

sequences were over-represented in *msh1* pt genome, compared with the mt genome (Table S43).

A partial palindromic sequence is also a specific pair of inverse repeats or microhomologies at the

same location. For example, we could observe crossover site distribution in chimeric reads caused

by rearrangement-mediated by partial palindromic sequence U-8 in *msh1* (Figure 7a; Table S44).

Second, although most NHEJ events were at low frequencies with a seemingly random distribution,

we could observe one NHEJ event which occurred only in one sample (M3-43-5) at relatively high

frequency (121 one-rearrangement reads versus 1385 zero-rearrangement reads), resulting in a 65-

bp deletion in the *petA* gene (Figure 7b; Table S45). This mutation was predicted to delete a strictly

conserved KQFE motif in the C-terminal stromal extension of cytochrome f, which would

compromise photosynthesis (Choquet et al., 2003). This indicated a different accumulation pattern

compared with the repeatable mitochondrial high-frequency rearrangements among samples, which

might suggest that different yellow sectors of leaves in *msh1* individuals could be caused by

different random MMEJ/NHEJ events of the pt genome. This kind of sample-specific accumulation

pattern of plastid variations was also reproduced when analyzing the SNVs in *msh1*. In total, we

identified 17 SNVs that were enriched in one or a few samples at high ratios (Figure 7c; Table S46).

They were predominantly AT-to-GC transitions, consistent with the mutation spectrum in *msh1*

mutants that was reported previously and suggested to be caused by polymerase errors (Wu et al.,

2020b). Third, in contrast to the indel variants caused by gene conversion between imperfect

repeats observed in the mt genome of *msh1* mutants, the 85 identified indel mutations in the pt

genome were mostly mediated by homopolymers (Tables S47). For example, the 13-bp

homopolymer at position 67164 contained the most variant types detected by GATK, with higher

variant frequencies in *msh1* mutants than the wild-type individual (Figure 7d). Indel variant

frequencies were also much higher in *msh1* individuals compared with wild-type at most other

positions (Figure 7e). Together, these results show that loss of *MSH1* causes different patterns of

structural variants and mutation accumulation in pt and mt genomes.

Discussion

In this work, we used HiFi reads (~15 kb long, > 99% accuracy) to study the behaviors of

mitochondrial repeats using samples from Arabidopsis *msh1* mutant individuals. HiFi reads were

long enough to span the entire lengths of repeats, enabling analyses that were not previously

possible with short reads. We developed a mapping-based analysis, which showed that HiFi reads

performed well in identifying more than one structural variation in a single read and studying

polymorphisms between long non-identical repeats. They were also sufficiently accurate to

distinguish mt genome sequences from those associated with the large nuclear mitochondrial
insertion in Chromosome 2, although that structure has not been fully captured in the reference
nuclear genome assembly, so it may not be possible to fully separate reads from these two different
sources (Stupar et al., 2001). Compared to previous studies, we also identified additional repeats
active in recombination at lower frequencies. However, due to the low number of read counts, HiFi
reads were not cost-effective for studying low-frequency MMEJ and NHEJ variants. This cost
could be reduced by isolating organelles from the cells before DNA extraction for sequencing.
414

MSH1 suppresses gene conversion between non-identical repeats

In meiosis, MutS homologs detect mismatches and recruit the DSB-generating SPO11 complex
to suppress crossover repair between non-identical centromeric repeats (Blackwell et al., 2020;
Naish et al., 2021). SPO11 also repairs double DSB (dDSB) gaps independent of MSH2 at DSB
hotspots (Prieler et al., 2021). It was proposed that DNA mismatches detected by organelle-targeted
MSH1 are converted to DSBs by the GIY-YIG endonuclease domain at its C-terminus and directed
to DSB repair pathways (Christensen, 2014). In this analysis, HiFi reads were unambiguously
mapped across each mitochondrial repeat homolog, leading to the identification of high-frequency
non-crossover recombination or gene conversion in zero-rearrangement reads in *msh1* mutants,
including both SNVs (as in repeats A, H, etc.) and indels (as in repeat MMJS). Gene conversion
was previously observed to explain the discontinuous pattern of SNVs/indels between non-identical
repeats in chimeric reads (Davila et al., 2011), the limitations of short reads have generally
prevented comprehensive analyses across repeats of different lengths and quantitative comparisons
between wild-type and *msh1* mutants.
Numerous earlier studies have focused on the massive increase in repeat-mediated crossover
recombination and mt genome rearrangements in *msh1* mutants (Abdelnoor et al., 2003; Shedge et
al., 2007; Arrieta-Montiel et al., 2009; Davila et al; 2011). More recently, we found a role of MSH1
in suppressing organellar point mutations and indels (Wu et al., 2020b). At first glance, it might
seem contradictory that MSH1 could be involved in suppressing ectopic recombination between
short repeats, while also inducing homologous recombination as part of a mismatch repair pathway
(Christensen, 2014). However, there could very well be a connection between these two processes.
Specifically, MSH1 may provide a surveillance mechanism to check for mismatches in double-
stranded DNA regardless of whether they are introduced by polymerase errors or by strand invasion
between short or non-identical repeats, followed by introduction of DSBs to (re)start repair by
homologous recombination. To suppress recombination at short and intermediate repeats that are
100% identical in sequence, MSH1 may need to detect differences in boundary sequence. During

homologous recombination, the bacterial RecA-ssDNA presynaptic filament checks homology in
~9 bp steps before strand exchange with the dsDNA template (Huang et al., 2020). In such
screening mechanisms, the first mismatches may be detected at the repeat boundaries when the
repeat sequences themselves are identical. Thus, MSH1 may suppress both crossover and non-
crossover forms of ectopic recombination to ensure accurate DSB repair of the genome by detecting
mismatches within repeats or at repeat boundaries.

Conventional recombination activity only accounts for a proportion of DSB repair events
Most previous mitochondrial repeat annotations depended on the penalty parameters of mismatches
and gaps of raw sequence alignment results. We showed that through comparing chimeric HiFi
reads with blastn results, it was possible to identify additional low-frequency repeats, raising
questions about what contributes to the observed differences in recombination activity of each
repeat pair. Increased crossover recombination events of intermediate-sized repeats have been
observed in wild-type plants when DSB frequencies were increased by DNA damaging chemicals
(Wallet et al., 2015). Therefore, repeats near DSB hotspots may show higher ectopic recombination
activity compared to other repeats with similar length and sequence identity, which might explain
cases that fall far from the regression line that defines the relationship between recombination
activity and repeat length (Figure S1c). It may also help to study repair events systematically by
inducing DSBs through mitochondrial-targeting CRISPR-Cas9 or mitoTALENs (Arimura et al.,
2020) and in more RRR gene mutants, such as *why2*, *reca3*, *osb1* (Gualberto and Newton 2017).
DSBs within a repeat or in the flanking sequence would be repaired using homologous DNA
stretches of another intra- or intermolecular copy. After resection of DSB ends, RecA recombinases
bind 3' ssDNA and rapidly search for homology (Wiktor et al., 2021). It was reported that at least
8-nt microhomology was needed for RecA-ssDNA filaments to bind dsDNA templates stably, and
15-nt was sufficient to be resistant to facilitated exchange during homology search (Qi et al., 2015).
These studies suggested that only a small stretch of RecA-ssDNA was needed during each HDR
repair event. Typically, only crossover or BIR products have been used to estimate frequencies of
recombinant conformations (Wallet et al., 2015; Odahara et al., 2017). Such calculations do not
account for the fact that gene conversion events between repeats can also occur at substantial
frequencies. These gene conversion events likely explain why larger repeats tend to be 100%
identical. In wild-type plants, only the two large repeat pairs showed high recombination activities.
The observed difference in *msh1* mutants was the increased proportion of uncontrolled gene
conversion between non-identical repeats and BIR proportion of HDR repair. The lower detected
recombination activity of large repeats in *msh1* could be due to repair of DSBs by other repeats

(Figure 1d). When calculating recombination of some three-copy or four-copy repeats locating in

other longer two-copy repeats, BIR-indicating reads shared by both shorter and longer annotations

were assigned to the larger two-copy repeats (Figure S2b). This indicated that it was more natural

to view recombination events as associated with specific genomic regions per se, but not with repeat

annotations.

Phenotypic contributions of cytoplasmic genetic materials

During seed germination, normal mitochondrial biogenesis and mtDNA copy number are tightly

controlled (Paszkiewicz et al., 2017). This connection between seed germination and mitochondrial

defects was also reported in many other studies (Han et al., 2010; Virdi et al., 2015; Van Aken et

al., 2016; Heidorn-Czarna et al., 2018; Golin et al., 2020). We showed that variation in root lengths

at 5 DAS was heritable among different *msh1* mutant families. Based on sequencing coverage

distributions, the degree of slow seed germination was potentially associated with reduction in

atp6-1 copy number. Generally, coverage analysis indicated similar patterns of fluctuation in

sequence depth across *msh1* individuals, which could be a combined effect of the locations of repeat

pairs and the selection to maintain regions containing functional genes (Figure S7). Interestingly,

the most dynamic region in *msh1* containing Large1-2 was lost in other Arabidopsis accessions.

Despite the very rapid intraspecific structural rearrangements of mt genomes, the various

Arabidopsis accessions were all in a low-heteroplasmy state. During the process of

substoichiometric shifting (Woloszynska, 2009; Tang et al., 2016; Johnston, 2019), the mt genome

may be transiently in a highly rearranged and heteroplasmic state similar to *msh1* mutants,

providing source materials for functional innovation.

In comparison to plant mt genomes, the pt genome is smaller and contains a pair of very long

inverted repeats. The contrasting patterns of variation between mt and pt genomes may result from

differences in repeat spectrum, genome copy number per organelle, and DNA repair mechanisms.

In the pt genome of *msh1* mutants, HiFi-identified rearrangements were mostly MMEJ and NHEJ

events. MMEJ events mediated by partial palindromic sequences were found at a higher frequency

than others and showed repeatability among samples. However, a few other pt genome variants

occurred at high frequency in individual samples (Figure 7b). These individual high-frequency

variants might reflect mutations in different yellow patches, which can often be often in a

homoplasmic state because of random sorting early in leaf primordia (Malinova et al., 2021). Leaf

variegation and albino phenotypes are common in other RRR gene mutants (for example, moc1

single mutant, *why1 why3 pol1b* and *why1 why3 reca1* triple mutants) and transgenic plants with

plastid-targeted restriction endonucleases (Zampini et al., 2015; Kobayashi et al., 2017; Sugimoto

et al., 2020).

In conclusion, *MSH1* acts to control both structural and non-structural variants in plant organellar

genomes. The genomes could be transiently uncontrolled when *MSH1* expression was

downregulated, providing more raw materials for further functional selection (Yang and Mackenzie,

2019). However, the two genomes responded quite differently to the loss of the same gene. In the

future, through understanding the dynamics of these repeats and the detailed functions of RRR

genes in DSB repair processes, we may better harness the cytoplasmic genetic material for crop

improvement with recently developed mt genome editing technologies (Kazama et al., 2019;

Arimura et al., 2020).

Methods

Plant Materials and Growth Conditions

Generation of *msh1* mutant and wild-type F_3 families of *Arabidopsis thaliana* used for HiFi

sequencing was described previously (Wu et al., 2020b). Individuals showing clear long and short

root lengths at 5 days after sowing (DAS) were selected to get corresponding F_4 families. For

estimating the speed of seed germination, seeds were sowed on solid ½ Murashige and Skoog media

with vitamins and 1% sucrose, stratified in darkness (4°C, 2 d), and grown in a light incubator

(22°C, 16 h day and 8 h night) for another three days, e.g., 5 DAS. Photos of seedlings at 5 DAS

were analyzed with ImageJ to measure primary root lengths.

DNA extraction and PacBio HiFi sequencing

Above-ground tissues of 11 individual plants grown in a short-day grown chamber (8 h day and 16

h night) for six weeks were collected to extract DNA with the CTAB method. Total DNA was

purified with a QIAGEN kit (Q13343) and quantified using NanoDrop and Qubit. Purified DNA

(3.3~18.6 μg for each sample) was used for the construction of SMRTbell libraries according to

PacBio's standard protocol. Sequencing was performed using the PacBio Sequel II platform in CCS

mode, which generated a total of 34.36 Gb HiFi reads (12.7~18.1 kb mean read length).

Identification of rearrangements in mt and pt genomes

Raw HiFi reads of each sample were mapped to the Arabidopsis Col-0 reference genome (TAIR10

for nucleus; NC_037304.1 for mitochondria; NC_000932.1 plus a 1-bp expansion at 28673 for

plastid as described in Wu, et al., 2020) with minimap2 (version 2.20, r1061; parameters, -ax map-

hifi; Li, 2018). Reads mapped to mt and pt genomes were separated using bamtools (version 2.5.1) and realigned against each genome with blastn (version 2.12.0). Raw blastn results were further processed by a custom pipeline (available at https://github.com/zouyinstein/hifisr) to group the reads by discontinuity, homologous sequence size and mapping coordinates, and calculate read counts for input samples. Different types of genome rearrangements were manually checked, including zero-, one-, two-, three-, and four-rearrangement events in a single read, microhomology-mediated end joining (MMEJ; 2-49 bp homology sequence size) and non-homologous end-joining (NHEJ; 0-1 bp homology sequence size), replication slippage mediated by tandem repeats, and insertions.

Mitochondrial repeat annotations and recombination activity

Candidate coordinates of repeats were generated by blastn (version 2.12.0+; parameters, -word_size 4) using the mitochondrial reference sequence as both query and subject. The resulting coordinates were manually checked against the junction coordinates of one-rearrangement HiFi reads. In some cases, the boundaries of low-identity repeats were different from previous annotations. Per base reads counts ratios of zero-rearrangement reads versus total reads indicating dynamic regions of mt genome, zero-rearrangement and one-rearrangement read counts indicating parental and recombinant forms of repeat groups were calculated based on blastn outputs of the custom HiFi-SR pipeline.

Identification of gene conversion between imperfect repeats

To analyze patterns of SNVs/indels, zero-rearrangement reads of each sample were re-aligned to the mitochondrial reference genome by PacBio pbmm2 (version 1.3.0; parameters, --sort --preset CCS). The resulting BAM files were used for variant calling by GATK HaplotypeCaller (version 4.2.3.0). Depths of A, T, G, C and read counts of different indel sizes, and the percentage of reference cases of each GATK-called SNVs/indels were calculated from raw BAM files with the Python module pysam (version 0.17.0). SNVs/indels inferred from gene conversion events between some high-activity mitochondrial imperfect repeats were also used for calculations, which were not always detected by GATK.

Calculation of absolute and normalized per base coverage

Total mt genome-mapped reads were re-aligned to mitochondrial reference genomes by PacBio pbmm2 (version 1.3.0; parameters, --sort --preset CCS). Absolute per base coverage across the entire mt genome of each individual and the published datasets was calculated by samtools depth

subcommand (version 1.7) and normalized to total bases of each sample. Annotations are based on

the GenBank file NC_037304.1 (Sloan et al., 2018).

metaFlye assembly and synteny analysis of mt genomes

Raw PacBio CLR reads of each accession were mapped to reference genome (the respective nuclear

assembly in Jiao and Schneeberger, 2020; Col-0 NC_037304.1 for mitochondria and NC_000932.1

plus a 1-bp expansion at 28673 for plastid) with minimap2 (version 2.20, r1061; parameters, -ax

map-pb; Li, 2018). Reads mapped to mt genomes were separated using bamtools (version 2.5.1)

and were re-assembled with metaFlye (Flye version 2.9-b1774; parameters, --meta --pacbio-hifi/--

pacbio-corr --extra-params output_gfa_before_rr=1 --genome-size 370K/2M for mitochondria).

Before assembling, PacBio CLR reads longer than 1000 bp were extracted by Filtlong (version

0.2.1; parameters, --min_length 1000) and were corrected using CONSENT (version 2.2.2;

parameters, --type PB). Assembly graphs before and after repeat resolution were visualized by

Bandage (version 0.8.1). Reads from different *msh1* families are merged. To get linear

representations of pseudo-master circles for mapping-based analysis, we started from the

assemblies (--genome-size 2M) after repeat resolution, discarded low-coverage contigs and

manually resolved the large repeats for accessions Eri-1, C24, Cvi-0, An-1, L*er*, Sha, and re-

resolved Large2 repeats for Kyoto with the Bandage software, followed by rotation or reverse

complementation. Corrected CLR reads were realigned against each pseudo-master circle using the

HiFi-SR pipeline. To analyze the shared synteny between mt genomes, we performed blastn

alignments using the Col-0 mitochondrial reference sequence as query and pseudo-master circle of

the other accession as subject. Raw blastn results were processed by a custom Python script

available in HiFi-SR. Coordinates or large repeats were determined based on blastn alignments

using the pseudo-master circle as both query and subject. Results were generated by the Python

package pycircos (version 1.0.2).

Identification of SNVs/indels pt genome

To analyze patterns of SNVs/indels in pt genome, total reads of each sample were re-aligned to

plastid reference genome by PacBio pbmm2 (version 1.3.0; parameters, --sort --preset CCS). The

resulting BAM files were used for variant calling by GATK HaplotypeCaller (version 4.2.3.0).

Depths of A, T, G, C and read counts of different indel sizes, and the percentage of reference cases

of each GATK-called SNVs/indels were calculated from raw BAM files with the Python module

pysam (version 0.17.0).

Acknowledgments

We thank Xiaojing Zhang from Prof. Li-Zhen Tao's laboratory at South China Agricultural University for assistance with plant growth. This work was supported by grants from National Natural Science Foundation of China (32170238), the Science, Technology and Innovation Commission of Shenzhen Municipality (RCYX20200714114538196), and the U.S. National Institutes of Health (R01 GM118046).

Author contributions

ZW designed the research; YZ performed the experiments; YZ wrote the computer scripts and performed majority of the bioinformatic analysis; YZ, WZ, DS, and ZW discussed the results; YZ, DS and ZW wrote the manuscript; DS and ZW revised the manuscript.

ORCID

Yi Zou: 0000-0003-3117-3898

Weidong Zhu: 0000-0002-9613-354X

Daniel B. Sloan: 0000-0002-3618-0897

Zhiqiang Wu: 0000-0002-4238-7317

Conflict of Interest

The authors have no conflict of interest to declare.

Data Availability

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive (Chen et al, 2021) in National Genomics Data Center (C.-N. Members and Partners, 2022), China National Center for Bioinformation / Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA006060) that are publicly accessible at https://ngdc.cncb.ac.cn/gsa. The Python-based pipeline for the detection of plant mitochondrial structural rearrangements used in this study is available at https://github.com/zouyinstein/hifisr.

Supporting Information

Figure S1 99 repeat groups supported by HiFi reads in the mt genome of *msh1*

Figure S2 Asymmetrical recombination phenomena in mt genome

Figure S3 Imperfect repeats with relatively higher ratios of gene conversion events

Figure S4 Coverages of regions containing repeat group H in wild-type and *msh1* individuals

Figure S5 Coverages of regions containing repeat group MMJS in wild-type and *msh1* individuals

Figure S6 Primary root length of wild-type and *msh1* seedlings at 5 days after sowing

Figure S7 Normalized per base read coverage across mt genome

Figure S8 The metaFlye assemblies of mt genomes are more complex in *msh1* than wild-type

Figure S9 Assembly graphs and mapping-based analysis of published HiFi datasets

Figure S10 The metaFlye assemblies of mt genomes are simple as wild-type in other accessions

Figure S11 Synteny and coverage analysis of Eri-1 mt genome

Figure S12 Synteny and coverage analysis of C24 mt genome

Figure S13 Synteny and coverage analysis of Cvi-0 mt genome

Figure S14 Synteny and coverage analysis of An-1 mt genome

Figure S15 Synteny and coverage analysis of L*er* mt genome

Figure S16 Synteny and coverage analysis of Kyoto mt genome

Figure S17 Synteny and coverage analysis of Sha mt genome

Table S1 DNA content and fresh weight of individual plants used for HiFi sequencing

Table S2 Mapping statistics of HiFi reads of individual plants

Table S3 Statistics of mt genome-mapped reads of individual plants

Table S4 Statistics of pt genome-mapped reads of individual plants

Table S5 Read counts of different types of mitochondrial structural rearrangements of individual
plants

Table S6 Read counts of different types of plastid structural rearrangements of individual plants

Table S7 Read counts of different categories of structural rearrangements in mt genome of wild-
type and *msh1* plants

Table S8 Read counts of different categories of structural rearrangements in pt genome of wild-
type and *msh1* plants

Table S9 Overlapping region length of the two alignments in mitochondrial one-rearrangement
events detected by HiFi reads.

Table S10 Overlapping region length of the two alignments in plastid one-rearrangement events
detected by HiFi reads.

Table S11 Raw blastn outputs using the plastid reference sequence as both query and subject

Table S12 HiFi-SR outputs of one-rearrangement mt genome-mapped reads of individual plants

Table S13 HiFi-SR outputs of two-rearrangement mt genome-mapped reads of individual plants

Table S14 Raw blastn outputs using the mitochondrial reference sequence as both query and
subject

Table S15 Annotations of two-copy repeats based on blastn and one-rearrangement HiFi reads

Table S42 Absolute and normalized per base read coverage across mt genome of each *Arabidopsis*
thaliana accessions from the published PacBio CLR datasets

Table S43 HiFi-SR outputs of one-rearrangement mediated by short partial palindromic sequences
in pt genomes of individual plants

Table S44 HiFi-SR outputs of one-rearrangement mediated by short partial palindromic sequence
U-8 in pt genomes of individual plants

Table S45 HiFi-SR outputs of a NHEJ event in pt genomes of individual plants

Table S46 Ratios of GATK-identified SNVs of individual plants and the published HiFi datasets
in pt genome

Table S47 Ratios of GATK-identified indels of individual plants and the published HiFi datasets
in pt genome

References

1.   Abdelnoor RV, Yule R, Elo A, Christensen AC, Meyer-Gauen G, Mackenzie SA.
Substoichiometric shifting in the plant mitochondrial genome is influenced by a gene
homologous to MutS. **Proc Natl Acad Sci U S A 2003**, 100(10)**:** 5968-5973.

2.   Arimura SI, Ayabe H, Sugaya H, Okuno M, Tamura Y, Tsuruta Y, Watari Y, Yanase S,
Yamauchi T, Itoh T, Toyoda A, Takanashi H, Tsutsumi N. Targeted gene disruption of
ATP synthases 6-1 and *6-2* in the mitochondrial genome of *Arabidopsis thaliana* by
mitoTALENs. **Plant J 2020**, 104(6)**:** 1459-1471.

3.   Arrieta-Montiel MP, Shedge V, Davila J, Christensen AC, Mackenzie SA. Diversity of the
Arabidopsis mitochondrial genome occurs via nuclear-controlled recombination activity.
Genetics 2009, 183(4)**:** 1261-1268.

4.   Ayala-Garcia VM, Baruch-Torres N, Garcia-Medel PL, Brieba LG. Plant organellar DNA
polymerases paralogs exhibit dissimilar nucleotide incorporation fidelity. **FEBS J 2018**,
285(21)**:** 4005-4018.

5.   Barnett DW, Garrison EK, Quinlan AR, Stromberg MP, Marth GT. BamTools: a C++ API
and toolkit for analyzing and managing BAM files. **Bioinformatics 2011**, 27(12)**:** 1691-
1692.

6.   Blackwell AR, Dluzewska J, Szymanska-Lejman M, Desjardins S, Tock AJ, Kbiri N,
Lambing C, Lawrence EJ, Bieluszewski T, Rowan B, Higgins JD, Ziolkowski PA,
Henderson IR. MSH2 shapes the meiotic crossover landscape in relation to interhomolog
polymorphism in Arabidopsis. **EMBO J 2020**, 39(21)**:** e104858.

7.　　Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
BLAST+: architecture and applications. **BMC Bioinformatics 2009**, 10**:** 421.

8.　　Chen T, Chen X, Zhang S, Zhu J, Tang B, Wang A, Dong L, Zhang Z, Yu C, Sun Y, Chi
L, Chen H, Zhai S, Sun Y, Lan L, Zhang X, Xiao J, Bao Y, Wang Y, Zhang Z, Zhao W.
The Genome Sequence Archive Family: Toward Explosive Data Growth and Diverse Data
Types. **Genomics Proteomics Bioinformatics 2021**.

9.　　Chevigny N, Schatz-Daas D, Lotfi F, Gualberto JM. DNA Repair and the Stability of the
Plant Mitochondrial Genome. **Int J Mol Sci 2020**, 21(1).

10.　　Choquet Y, Zito F, Wostrikoff K, Wollman FA. Cytochrome f translation in
Chlamydomonas chloroplast is autoregulated by its carboxyl-terminal domain. **Plant Cell**
2003, 15(6)**:** 1443-1454.

11.　　Christensen AC. Plant mitochondrial genome evolution can be explained by DNA repair
mechanisms. **Genome Biol Evol 2013**, 5(6)**:** 1079-1086.

12.　　Christensen AC. Genes and junk in plant mitochondria-repair mechanisms and selection.
Genome Biol Evol 2014, 6(6)**:** 1448-1453.

13.　　Christensen AC. Plant Mitochondria are a Riddle Wrapped in a Mystery Inside an Enigma.
J Mol Evol 2021, 89(3)**:** 151-156.

14.　　Davila JI, Arrieta-Montiel MP, Wamboldt Y, Cao J, Hagmann J, Shedge V, Xu YZ, Weigel
D, Mackenzie SA. Double-strand break repair processes drive evolution of the
mitochondrial genome in Arabidopsis. **BMC Biol 2011**, 9**:** 64.

15.　　Flood PJ, Theeuwen T, Schneeberger K, Keizer P, Kruijer W, Severing E, Kouklas E,
Hageman JA, Wijfjes R, Calvo-Baltanas V, Becker FFM, Schnabel SK, Willems LAJ,
Ligterink W, van Arkel J, Mumm R, Gualberto JM, Savage L, Kramer DM, Keurentjes
JJB, van Eeuwijk F, Koornneef M, Harbinson J, Aarts MGM, Wijnker E. Reciprocal
cybrids reveal how organellar genomes affect plant phenotypes. **Nat Plants 2020**, 6(1)**:**
13-21.

16.　　Fuchs P, Rugen N, Carrie C, Elsasser M, Finkemeier I, Giese J, Hildebrandt TM, Kuhn K,
Maurino VG, Ruberti C, Schallenberg-Rudinger M, Steinbeck J, Braun HP, Eubel H,
Meyer EH, Muller-Schussele SJ, Schwarzlander M. Single organelle function and
organization as estimated from Arabidopsis mitochondrial proteomics. **Plant J 2020**,
101(2)**:** 420-441.

17.　　Fukui K, Harada A, Wakamatsu T, Minobe A, Ohshita K, Ashiuchi M, Yano T. The GIY-
YIG endonuclease domain of *Arabidopsis* MutS homolog 1 specifically binds to branched
DNA structures. **FEBS Lett 2018**, 592(24)**:** 4066-4077.

18.   Gobron N, Waszczak C, Simon M, Hiard S, Boivin S, Charif D, Ducamp A, Wenes E,
Budar F. A cryptic cytoplasmic male sterility unveils a possible gynodioecious past for
Arabidopsis thaliana. **PLoS One 2013**, 8(4)**:** e62450.

19.   Golin S, Negroni YL, Bennewitz B, Klosgen RB, Mulisch M, La Rocca N, Cantele F,
Vigani G, Lo Schiavo F, Krupinska K, Zottini M. WHIRLY2 plays a key role in
mitochondria morphology, dynamics, and functionality in *Arabidopsis thaliana*. **Plant**
Direct 2020, 4(5)**:** e00229.

20.   Gualberto JM, Newton KJ. Plant Mitochondrial Genomes: Dynamics and Mechanisms of
Mutation. **Annu Rev Plant Biol 2017**, 68**:** 225-252.

21.   Han L, Qin G, Kang D, Chen Z, Gu H, Qu LJ. A nuclear-encoded mitochondrial gene
AtCIB22 is essential for plant development in *Arabidopsis*. **J Genet Genomics 2010**,
37(10)**:** 667-683.

22.   Heidorn-Czarna M, Domanski D, Kwasniak-Owczarek M, Janska H. Targeted Proteomics
Approach Toward Understanding the Role of the Mitochondrial Protease FTSH4 in the
Biogenesis of OXPHOS During Arabidopsis Seed Germination. **Front Plant Sci 2018**, 9**:**
821.

23.   Huang X, Lu Y, Wang S, Sui M, Li J, Ma J, Ma D, Jia Q, Hu S, Xu C, Li M. Mismatch
sensing by nucleofilament deciphers mechanism of RecA-mediated homologous
recombination. **Proc Natl Acad Sci U S A 2020**, 117(34)**:** 20549-20554.

24.   Hussmann D, Starnawska A, Kristensen L, Daugaard I, Thomsen A, Kjeldsen TE, Hansen
CS, Bybjerg-Grauholm J, Johansen KD, Ludvigsen M, Kristensen T, Larsen TS, Moller
MB, Nyvold CG, Hansen LL, Wojdacz TK. *IGHV*-associated methylation signatures more
accurately predict clinical outcomes of chronic lymphocytic leukemia patients than *IGHV*
mutation load. **Haematologica 2021**.

25.   Jiao WB, Schneeberger K. Chromosome-level assemblies of multiple Arabidopsis
genomes reveal hotspots of rearrangements with altered evolutionary dynamics. **Nat**
Commun 2020, 11(1)**:** 989.

26.   Johnston IG. Tension and Resolution: Dynamic, Evolving Populations of Organelle
Genomes within Plant Cells. **Mol Plant 2019**, 12(6)**:** 764-783.

27.   Kazama T, Okuno M, Watari Y, Yanase S, Koizuka C, Tsuruta Y, Sugaya H, Toyoda A,
Itoh T, Tsutsumi N, Toriyama K, Koizuka N, Arimura SI. Curing cytoplasmic male sterility
via TALEN-mediated mitochondrial genome editing. **Nat Plants 2019**, 5(7)**:** 722-730.

28.   Kobayashi Y, Misumi O, Odahara M, Ishibashi K, Hirono M, Hidaka K, Endo M,
Sugiyama H, Iwasaki H, Kuroiwa T, Shikanai T, Nishimura Y. Holliday junction
resolvases mediate chloroplast nucleoid segregation. **Science 2017**, 356(6338)**:** 631-634.

29.   Kolmogorov M, Bickhart DM, Behsaz B, Gurevich A, Rayko M, Shin SB, Kuhn K, Yuan
J, Polevikov E, Smith TPL, Pevzner PA. metaFlye: scalable long-read metagenome
assembly using repeat graphs. **Nat Methods 2020**, 17(11)**:** 1103-1110.

30.   Kozik A, Rowan BA, Lavelle D, Berke L, Schranz ME, Michelmore RW, Christensen AC.
The alternative reality of plant mitochondrial DNA: One ring does not rule them all. **PLoS**
Genet 2019, 15(8)**:** e1008373.

31.   Kundariya H, Yang X, Morton K, Sanchez R, Axtell MJ, Hutton SF, Fromm M, Mackenzie
SA. MSH1-induced heritable enhanced growth vigor through grafting is associated with
the RdDM pathway in plants. **Nat Commun 2020**, 11(1)**:** 5343.

32.   Lee J, Cho CH, Park SI, Choi JW, Song HS, West JA, Bhattacharya D, Yoon HS. Parallel
evolution of highly conserved plastid genome architecture in red seaweeds and seed plants.
BMC Biol 2016, 14**:** 75.

33.   Li H. Minimap2: pairwise alignment for nucleotide sequences. **Bioinformatics 2018**,
34(18)**:** 3094-3100.

34.   Mackenzie SA, Kundariya H. Organellar protein multi-functionality and phenotypic
plasticity in plants. **Philos Trans R Soc Lond B Biol Sci 2020**, 375(1790)**:** 20190182.

35.   Malinova I, Zupok A, Massouh A, Schottler MA, Meyer EH, Yaneva-Roder L, Szymanski
W, Rossner M, Ruf S, Bock R, Greiner S. Correction of frameshift mutations in the *atpB*
gene by translational recoding in chloroplasts of *Oenothera* and tobacco. **Plant Cell 2021**,
33(5)**:** 1682-1705.

36.   Massouh A, Schubert J, Yaneva-Roder L, Ulbricht-Jones ES, Zupok A, Johnson MT,
Wright SI, Pellizzer T, Sobanski J, Bock R, Greiner S. Spontaneous Chloroplast Mutants
Mostly Occur by Replication Slippage and Show a Biased Pattern in the Plastome of
Oenothera. **Plant Cell 2016**, 28(4)**:** 911-929.

37.   Masutani B, Arimura SI, Morishita S. Investigating the mitochondrial genomic landscape
of *Arabidopsis thaliana* by long-read sequencing. **PLoS Comput Biol 2021**, 17(1)**:**
e1008597.

38.   Members C-N, Partners. Database Resources of the National Genomics Data Center, China
National Center for Bioinformation in 2022. **Nucleic Acids Res 2022**, 50(D1)**:** D27-D38.

39.   Morisse P, Marchet C, Limasset A, Lecroq T, Lefebvre A. Scalable long read self-
843   correction and assembly polishing with multiple sequence alignment. **Sci Rep 2021**, 11(1)**:**
844   761.

845   40.   Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmucker A, Mandakova
846   T, Jamge B, Lambing C, Kuo P, Yelina N, Hartwick N, Colt K, Smith LM, Ton J, Kakutani
847   T, Martienssen RA, Schneeberger K, Lysak MA, Berger F, Bousios A, Michael TP, Schatz
848   MC, Henderson IR. The genetic and epigenetic landscape of the *Arabidopsis* centromeres.
849   **Science 2021**, 374(6569)**:** eabi7489.

850   41.   Odahara M, Kishita Y, Sekine Y. MSH1 maintains organelle genome stability and
851   genetically interacts with *RECA* and *RECG* in the moss *Physcomitrella patens*. **Plant J**
852   **2017**, 91(3)**:** 455-465.

853   42.   Palmer JD, Herbon LA. Plant mitochondrial DNA evolves rapidly in structure, but slowly
854   in sequence. **J Mol Evol 1988**, 28(1-2)**:** 87-97.

855   43.   Paszkiewicz G, Gualberto JM, Benamar A, Macherel D, Logan DC. Arabidopsis Seed
856   Mitochondria Are Bioenergetically Active Immediately upon Imbibition and Specialize via
857   Biogenesis in Preparation for Autotrophic Growth. **Plant Cell 2017**, 29(1)**:** 109-128.

858   44.   Prieler S, Chen D, Huang L, Mayrhofer E, Zsoter S, Vesely M, Mbogning J, Klein F. Spo11
859   generates gaps through concerted cuts at sites of topological stress. **Nature 2021**,
860   594(7864)**:** 577-582.

861   45.   Qi Z, Redding S, Lee JY, Gibb B, Kwon Y, Niu H, Gaines WA, Sung P, Greene EC. DNA
862   sequence alignment by microhomology sampling during homologous recombination. **Cell**
863   **2015**, 160(5)**:** 856-869.

864   46.   Redei GP. Extra-chromosomal mutability determined by a nuclear gene locus in
865   Arabidopsis. **Mutat Res 1973**, 18: 149–162.

866   47.   Sandhu AP, Abdelnoor RV, Mackenzie SA. Transgenic induction of mitochondrial
867   rearrangements for cytoplasmic male sterility in crop plants. **Proc Natl Acad Sci U S A**
868   **2007**, 104(6)**:** 1766-1770.

869   48.   Shedge V, Arrieta-Montiel M, Christensen AC, Mackenzie SA. Plant mitochondrial
870   recombination surveillance requires unusual *RecA* and *MutS* homologs. **Plant Cell 2007**,
871   19(4)**:** 1251-1264.

872   49.   Sloan DB, Wu Z, Sharbrough J. Correction of Persistent Errors in Arabidopsis Reference
873   Mitochondrial Genomes. **Plant Cell 2018**, 30(3)**:** 525-527.

874   50.   Stupar RM, Lilly JW, Town CD, Cheng Z, Kaul S, Buell CR, Jiang J. Complex mtDNA
875   constitutes an approximate 620-kb insertion on *Arabidopsis thaliana* chromosome 2:

876   implication of potential sequencing errors caused by large-unit repeats. **Proc Natl Acad**
877   **Sci U S A 2001**, 98(9)**:** 5099-5103.

51.   Sugimoto H, Hirano M, Tanaka H, Tanaka T, Kitagawa-Yogo R, Muramoto N, Mitsukawa N. Plastid-targeted forms of restriction endonucleases enhance the plastid genome rearrangement rate and trigger the reorganization of its genomic architecture. **Plant J 2020**, 102(5)**:** 1042-1057.

52.   Tang H, Zheng X, Li C, Xie X, Chen Y, Chen L, Zhao X, Zheng H, Zhou J, Ye S, Guo J, Liu YG. Multi-step formation, evolution, and functionalization of new cytoplasmic male sterility genes in the plant mitochondrial genomes. **Cell Res 2017**, 27(1)**:** 130-146.

53.   Unseld M, Marienfeld JR, Brandt P, Brennicke A. The mitochondrial genome of *Arabidopsis thaliana* contains 57 genes in 366,924 nucleotides. **Nat Genet 1997**, 15(1)**:** 57-61.

54.   Van Aken O, Ford E, Lister R, Huang S, Millar AH. Retrograde signalling caused by heritable mitochondrial dysfunction is partially mediated by ANAC017 and improves plant performance. **Plant J 2016**, 88(4)**:** 542-558.

55.   Virdi KS, Laurie JD, Xu YZ, Yu J, Shao MR, Sanchez R, Kundariya H, Wang D, Riethoven JJ, Wamboldt Y, Arrieta-Montiel MP, Shedge V, Mackenzie SA. *Arabidopsis* MSH1 mutation alters the epigenome and produces heritable changes in plant growth. **Nat Commun 2015**, 6**:** 6386.

56.   Virdi KS, Wamboldt Y, Kundariya H, Laurie JD, Keren I, Kumar KRS, Block A, Basset G, Luebker S, Elowsky C, Day PM, Roose JL, Bricker TM, Elthon T, Mackenzie SA. MSH1 Is a Plant Organellar DNA Binding and Thylakoid Protein under Precise Spatial Regulation to Alter Development. **Mol Plant 2016**, 9(2)**:** 245-260.

57.   Wallet C, Le Ret M, Bergdoll M, Bichara M, Dietrich A, Gualberto JM. The RECG1 DNA Translocase Is a Key Factor in Recombination Surveillance, Repair, and Segregation of the Mitochondrial DNA in Arabidopsis. **Plant Cell 2015**, 27(10)**:** 2907-2925.

58.   Wang B, Yang X, Jia Y, Xu Y, Jia P, Dang N, Wang S, Xu T, Zhao X, Gao S, Dong Q, Ye K. High-quality *Arabidopsis thaliana* Genome Assembly with Nanopore and HiFi Long Reads. **Genomics Proteomics Bioinformatics 2021**.

59.   Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of *de novo* genome assemblies. **Bioinformatics 2015**, 31(20)**:** 3350-3352.

60.   Wiktor J, Gynna AH, Leroy P, Larsson J, Coceano G, Testa I, Elf J. RecA finds homologous DNA by reduced dimensionality search. **Nature 2021**, 597(7876)**:** 426-429.

61. Wolfe KH, Li WH, Sharp PM. Rates of nucleotide substitution vary greatly among plant
mitochondrial, chloroplast, and nuclear DNAs. **Proc Natl Acad Sci U S A 1987**, 84(24)**:**
9054-9058.

62. Woloszynska M. Heteroplasmy and stoichiometric complexity of plant mitochondrial
genomes--though this be madness, yet there's method in't. **J Exp Bot 2010**, 61(3)**:** 657-671.

63. Wu Z, Liao X, Zhang X, Tembrock LR, Broz A. Genomic architectural variation of plant
mitochondria—A review of multichromosomal structuring. **J Syst Evol 2020a**, 60(1): 160-
168.

64. Wu Z, Waneka G, Broz AK, King CR, Sloan DB. *MSH1* is required for maintenance of
the low mutation rates in plant mitochondrial and plastid genomes. **Proc Natl Acad Sci U**
S A 2020b, 117(28)**:** 16448-16455.

65. Wynn E, Purfeerst E, Christensen A. Mitochondrial DNA Repair in an *Arabidopsis*
thaliana Uracil N-Glycosylase Mutant. **Plants (Basel) 2020**, 9(2).

66. Wynn EL, Christensen AC. Repeats of Unusual Size in Plant Mitochondrial Genomes:
Identification, Incidence and Evolution. **G3 (Bethesda) 2019**, 9(2)**:** 549-559.

67. Xu YZ, Arrieta-Montiel MP, Virdi KS, de Paula WB, Widhalm JR, Basset GJ, Davila JI,
Elthon TE, Elowsky CG, Sato SJ, Clemente TE, Mackenzie SA. MutS HOMOLOG1 is a
nucleoid protein that alters mitochondrial and plastid properties and plant response to high
light. **Plant Cell 2011**, 23(9)**:** 3428-3441.

68. Yang X, Mackenzie SA. Many Facets of Dynamic Plasticity in Plants. **Cold Spring Harb**
Perspect Biol 2019, 11(10).

69. Yang X, Sanchez R, Kundariya H, Maher T, Dopp I, Schwegel R, Virdi K, Axtell MJ,
Mackenzie SA. Segregation of an *MSH1* RNAi transgene produces heritable non-genetic
memory in association with methylome reprogramming. **Nat Commun 2020**, 11(1)**:** 2214.

70. Yu R, Sun C, Zhong Y, Liu Y, Sanchez-Puerta MV, Mower JP, Zhou R. The minicircular
and extremely heteroplasmic mitogenome of the holoparasitic plant *Rhopalocnemis*
phalloides. **Curr Biol 2022**, 32(2)**:** 470-479 e475.

71. Zampini E, Lepage E, Tremblay-Belzile S, Truche S, Brisson N. Organelle DNA
rearrangement mapping reveals U-turn-like inversions as a major source of genomic
instability in *Arabidopsis* and humans. **Genome Res 2015**, 25(5)**:** 645-654.

72. Zhao N, Li Z, Zhang L, Yang X, Mackenzie SA, Hu Z, Zhang M, Yang J. MutS
HOMOLOG1 mediates fertility reversion from cytoplasmic male sterile *Brassica juncea*
in response to environment. **Plant Cell Environ 2021**, 44(1)**:** 234-246.

73.   Zhao N, Xu X, Wamboldt Y, Mackenzie SA, Yang X, Hu Z, Yang J, Zhang M. MutS HOMOLOG1 silencing mediates *ORF220* substoichiometric shifting and causes male sterility in *Brassica juncea*. **J Exp Bot 2016**, 67(1)**:** 435-444.

Figure Legends

Figure 1 Mapping of HiFi reads identified structural variants in Arabidopsis organelle genomes (a) Ratios of HiFi reads mapped to nuclear (nc), plastid (pt), mitochondrial (mt) genomes by minimap2; un denotes the ratio of unmapped reads. (b) Schematics of reads alignment to the reference genome. Reads mapped to two discontinuous positions indicates one-rearrangement events. The overlapping region of two alignments indicated sequence homologies such as non-tandem repeats. (c) Proportions of different types of mt genome rearrangements detected by HiFi reads in wild-type and *msh1* mutant; ref, repeat, repeat_double, repeat_triple, and repeat_quadra denote ratios of zero-, one-, two-, three-, four- rearrangement events; MMEJ, microhomology-mediated end joining (2-49 bp homology sequence size); NHEJ, non-homologous end-joining (0-1 bp homology sequence size); tandem, large-scale tandem repeats-mediated variants; insertion, large-scale nucleotide insertions (mostly are poly-G and poly-C tracts). (d) Distribution of overlapping region length of the two alignments in mitochondrial one-rearrangement events detected by HiFi reads. Reads from different *msh1* individuals are merged. Read counts are normalized to 10000 total reads. (e) Proportions of different types of pt genome rearrangements detected by HiFi reads in wild-type and *msh1* mutant. (f) Distribution of overlapping length of the two alignments in mitochondrial one-rearrangement events detected by HiFi reads. Reads from different *msh1* individuals are merged. Read counts are normalized to 100000 total reads per sample.

Figure 2 Repeat-mediated mt genome rearrangements identified by HiFi reads (a, b) Global patterns of the junctions and relative read counts of mitochondrial one-rearrangement events in (a) wild-type and (b) *msh1* mutant. In wild-type, most recombination events are mediated by large repeats, and intermediate repeats Q, V near Large1-2. In *msh1*, recombination events mediated by intermediate-sized repeats were common. Reads from different *msh1* individuals are merged. There are 5544 one-rearrangement reads out of 24162 total reads in all *msh1* samples. Colors denote repeat sizes. The sizes of dots denote read counts normalized to 10000 total reads per sample. (c, d) Global patterns of the junctions and relative read counts of mitochondrial two-rearrangement events in (c) wild-type and (d) *msh1* mutant. Reads from different *msh1* individuals are merged. There are 1668 two-rearrangement reads out of 24162 total reads in all *msh1* samples.

Colors denote repeat sizes. The thickness of lines linking two dots denotes read counts normalized
to 10000 total reads per sample (line color is the same as the larger dot). (e) Ratios of zero-
rearrangement reads versus total reads indicate multiple highly dynamic regions across the mt
genome in wild-type and *msh1*. Reads from different *msh1* families are merged. Bottom is
annotations of repeats as in NC_037304.1.

Figure 3 Continuous and discontinuous distribution of SNVs/indels in non-identical repeats
(a) Continuous change of SNVs in one-rearrangement HiFi reads indicate possible crossover site
distribution of recombination events between repeat CC-1 and CC-2 in *msh1*. Numbers in brackets
denote the sum of read counts in all samples without normalization. The shaded region denotes the
longest identical fragment in repeat CCs (b) The ratio of read counts between the longest identical
fragment was correlated with the ratio of length versus total length in each group of two-copy
repeats. Blue circles denote data points from non-identical two-copy repeats. Blue line denotes the
linear regression line. The diagonal line denotes equal ratios. (c) Discontinuous patterns of SNVs
in one-rearrangement chimeric reads of repeat H-1 and H-3 indicate partial gene conversion events
in *msh1*. Asterisks denote SNVs introduced by gene conversion from repeat homolog (black) H-1
or H-3, and (red) H-2. Numbers in brackets denote the sum of read counts in all samples without
normalization. Vertical bars in (a) and (c) denote A (green), T (red), G (brown), C (blue) for bases
and I (gray) for indel sites.

Figure 4 Gene conversion events between imperfect repeats are suppressed by *MSH1*
(a) Schematic graph showing mapping of zero-rearrangement reads to two non-identical repeats
with one SNV. Both long and short reference reads are correctly mapped based on the SNV in the
repeated sequence. Long gene-conversion (GC) reads could be correctly mapped based on flanking
sequences, enabling the identification of gene conversion events. Short GC reads failed to identify
gene conversion events because of incorrect mapping to the other repeat homolog. (b,c) Proportions
of gene conversion events at predicted sites in repeat group (b) H and (c) MMJS, which illustrated
unsuppressed gene conversion events in samples of *msh1* compared to wild-type. The stacked bar
height in each box indicated the ratio of a given base for A (green), T (red), G (brown), C (blue),
and I (gray) for indel sites. Reference bases are indicated below. Results were calculated from zero-
rearrangement reads. SNVs and indels are arranged from small to large coordinates in the reference.
Coordinates of the first and last SNVs/INDs (black triangles) are shown. H-2 and H-3 are in the
reverse orientation compared to H-1, and MMJS-2 is in the reverse orientation compared to MMJS-

1. Asterisks in (b) denote predicted gene conversion sites corresponding to two other repeat

homologs.

1011

Figure 5 Phenotypic correlations of mt genome variants in *msh1*

(a) Photos showing wild-type and *msh1* seedlings of different families at 5 days after sowing (DAS).

(b) Distributions of primary root length frequency of F_3 families at 5 DAS indicate delayed seed

germination ($120 > n \geq 100$). (c-e) Normalized per-base read coverage of mt genome regions: (c)

49201-51200, (d) 263901-265900, (e) 270501-271500. Blue arrows denote coding regions of *atp6-*

1, *atp6-2* and *atp9*. Purple arrows denote tRNA and ORF genes. Orange arrows denote different

mitochondrial repeats. Annotations are based on the GenBank accession NC_037304.1. Purple and

orange lines denote normalized per-base read coverage of M3-17 and M3-43 mutant individuals.

Green lines denote normalized per-base read coverage of W3-5-2, and the published datasets WT-

XJTU and WT-CEN. Red horizontal lines in c to e denote 100% coverage. The white bar in (a)

represents 10 mm.

1023

Figure 6 Loss of mt genome regions in other *Arabidopsis thaliana* accessions

(A-E) Normalized per-base read coverage of mt genome regions: (A) 49201-51200, (B) 263901-

265900, (C) 270501-271500, (D) 19201-21200, (E) 345801-347800. Blue arrows denote coding

regions of *atp6-1*, *atp6-2*, *atp9*, *rps3*, and *rps19*. Purple arrows denote tRNA and ORF genes.

Orange arrows denote different mitochondrial repeats. Annotations are based on the GenBank file

NC_037304.1. Green lines denote normalized per-base read coverage of published Col-0 CLR

dataset. Purple lines denote normalized per-base read coverage of published Eri-1 and C24 datasets.

Orange lines denote normalized per-base read coverage of published Cvi-0, An-1, L*er*, Kyoto and

Sha datasets. Red horizontal lines in C-E denote 100% coverage.

1033

Figure 7 Mutations and structural variants of in pt genome detected by HiFi reads

(A) A MMEJ event mediated by a partial palindromic sequence (U-8) in *msh1*. A partial

palindromic sequence is also a specific pair of inverse repeats or microhomologies at the same

location. Numbers in brackets denote the sum of read counts showing crossover site distribution in

chimeric reads in all samples without normalization. The ratio of one-rearrangement reads versus

zero-rearrangement reads across U-8 is 160/14786 (1.1% of total reads) in *msh1* and 0/1966 in

wild-type. (B) A NHEJ event occurring only in the M3-43-5 individual was predicted to disrupt the

petA gene. The ratio of one-rearrangement reads versus zero-rearrangement reads across this

deletion is 121/1385 (8.0% of total reads). (C) Proportions of AT-to-GC transitions in 17 GATK-

called SNVs in wild-type and *msh1* individuals, which were enriched in one or a few samples at

high ratios. The stacked bar height in each box indicated the ratio of a given base for A (green), T

(red), G (brown), and C (blue). Reference bases are indicated below. (D) Proportions of an IND

mutation caused by a 13-bp homopolymer started near position 67164 with wild-type showing the

larger proportion of reference genotype. Multiple indel sizes (ranging from 6-bp deletion to 18-bp

insertion) were calculated from raw BAM-alignment files by the pysam package. (E) Colors

indicate ratios of variant genotypes of individual samples minus the mean ratios of variant

genotypes of three wild-type samples of the 85 GATK-called indel mutations. The proportions of

variant genotype at most positions were much higher in *msh1* individuals compared with wild-type,

which supported the previously proposed role of *MSH1* in maintaining low mutation rates to some

extent.

1054

1055

Figure 1 Mapping of HiFi reads identified structural variants in *Arabidopsis* organelle genomes

1056

1057

Figure 2 Repeat-mediated mt genome rearrangements identified by HiFi reads

1058

1059

1060

1061

Figure 3 Continuous and discontinuous distribution of SNVs/indels in non-identical repeats

1062

1063

1064

1065

Figure 4 Gene conversion events between imperfect repeats are suppressed by *MSH1*

1066

1067

1068

1069

Figure 5 Phenotypic correlations of mt genome variants in *msh1*

1070

1071

1072

1073

Figure 6 Loss of mt genome regions in other *Arabidopsis thaliana* accessions

1074

1075

1076

1077

Figure 7 Mutations and structural variants of in pt genome detected by HiFi reads

1078

1079