Title

Loading-Induced Bone Formation is Mediated by *Wnt1* Induction in Osteoblast-Lineage Cells

Authors

Lisa Y. Lawson [a,b*], Nicole Migotsky [a,b,c], Christopher J. Chermside-Scabbo [a,b,d], John T. Shuster

[a,b], Kyu Sang Joeng[e], Roberto Civitelli [b,f], Brendan Lee [g], Matthew J. Silva [a,b,c]

[a] Department of Orthopaedic Surgery, Washington University School of Medicine, Saint Louis,

MO, United States

[b] Musculoskeletal Research Center, Washington University School of Medicine, Saint Louis,

MO, United States

[c] Department of Biomedical Engineering, Washington University, Saint Louis, MO, United

States

[d] Medical Scientist Training Program, Washington University School of Medicine, St. Louis,

MO, USA

[e] McKay Orthopaedic Research Laboratory, Department of Orthopaedic Surgery, Perelman

School of Medicine, University of Pennsylvania, Philadelphia, PA, United States

[f] Division of Bone and Mineral Diseases, Department of Medicine, Washington University, Saint

Louis, MO, United States

[g] Department of Molecular and Human Genetics, Baylor College of Medicine, Waco, TX, United

States

[*] Corresponding author: lylawson@wustl.edu

April 15, 2022

Abstract

Mechanical loading on the skeleton stimulates bone formation. Although the exact mechanism underlying this process remains unknown, a growing body of evidence indicates that the Wnt signaling pathway is necessary for the skeletal response to loading. Recently, we showed that Wnts produced by osteoblast lineage cells mediate the osteo-anabolic response to tibial loading in adult mice. Here, we report that *Wnt1* specifically plays a crucial role in mediating the mechano-adaptive response to loading. Independent of loading, short-term loss of *Wnt1* in the Osx-lineage resulted in decreased cortical bone area in the tibias of 5-month old mice. In females, strain-matched loading enhanced periosteal bone formation in Wnt1F/F controls, but not in Wnt1F/F; OsxCreERT2 knockouts. In males, strain-matched loading increased periosteal bone formation in both control and knockout mice; however, the periosteal relative bone formation rate was 65% lower in *Wnt1* knockouts versus controls. Together, these findings show that *Wnt1* supports adult bone homeostasis and mediates the bone anabolic response to mechanical loading.

Keywords

Osteoblasts, Osteocytes, Bone, Osteogenesis, Mechanobiology, Wnt Signaling Pathway

Introduction

Mechanical loading on the skeleton stimulates bone formation. Although the mechanisms underlying this process are not fully defined, a growing body of evidence indicates that the Wnt signaling pathway plays a role in loading-induced bone formation. For example, ulnar loading suppresses expression of Wnt antagonist Sclerostin, and downregulation of this antagonist is in fact necessary for the bone anabolic response to loading[1-2]. Furthermore, mice lacking Wnt co-receptor Lrp5 exhibit a diminished bone anabolic response to loading, whereas mice with high-bone mass (HBM) mutations in Lrp5 have enhanced responses to loading[3-5].

Wnt signaling initiates when one of 19 Wnt ligands ('Wnts') binds to Frizzled and Lrp5/6 co-receptors[6]. Recently, we reported that blocking Wnts secretion in the Osx-expressing cells of skeletally mature mice reduced the osteo-anabolic response to loading[7]. OsxCreERT2; Wls[F/F]; mice were tamoxifen-treated to inactivate Wntless – the Wnt-specific transport receptor responsible for shuttling Wnt ligands to the cell surface for exocytosis[8] – thereby blocking the secretion of all Wnts from osteoblasts and osteocytes. We showed that the periosteal response to tibial loading was 65% lower in Wls knockouts relative to WlsF/F controls, indicating that Wnts produced by osteoblast-lineage cells are important mediators of loading-induced bone formation[7].

Gene expression studies have aimed to identify the key factors that mediate loading-induced bone formation. In our lab and others, RNASeq analysis of cortical bone samples showed that *Wnt1* and *Wnt7b* were differentially expressed in loaded vs non-loaded bones, suggesting that these Wnt ligands in particular may be necessary for the skeletal response to loading[9-11]. Furthermore, age-related declines in the skeletal response to loading are associated with dampened upregulation of *Wnt1* and *Wnt7b* in the bone, while *in vivo* overexpression of

Wnt1 and *Wnt7b* in the osteo-lineage causes increased bone mass[12-14]. Based on these studies,

we hypothesized that loading-induced upregulation of *Wnt1* and *Wnt7b* in osteoblasts is

necessary for the bone anabolic response to loading. In this communication, we report that

blocking *Wnt1* upregulation in the osteoblast-lineage via *Wnt1* deletion in Osx-expressing cells

blunts the bone anabolic response to skeletal loading in mice.

Methods

Mice

Maintenance and genotyping. All animal studies were approved by the Washington University

IACUC. Mice were derived from breeders originally gifted by Drs. Henry Kronenberg

(OsxCreERT2[15]), Brendan Lee (Wnt1 floxed[13]), or obtained from Jackson Labs (Wnt7b[C3]

floxed; B6; 129X1-Wnt7b[tm2Amc]/J; stock number 008467). Experimental mice were generated by

breeding Wnt1F/F dams to OsxCreERT2; Wnt1F/F sires (for _Wnt1_ single conditional mutants),

or Wnt1F/F; Wnt7bF/F dams to OsxCreERT2; Wnt1F/F; Wnt7F/F sires (for _Wnt1/7b_ double

conditional mutants). Mice were weaned at 21 days and housed up to five per cage with _ad_

libitum access to normal chow and water. Mice were genotyped by Transnetyx (Transnetyx,

USA) using tail snip DNA with probes for wild-type _Wnt1_ (_Wnt1-1 WT_), conditional _Wnt1_

(_L1L2-Bact-P MD_), wild-type _Wnt7b_ (_Wnt7b-2 WT_), conditional Wnt7b (_Wnt7b-2 FL_), and

OsxCreERT2 (_Cre_). Mice were euthanized by CO_2 asphyxiation.

Tamoxifen induction. Gene deletion was induced in 20 to 22 week old mice. Tamoxifen was

dissolved in corn oil to a final concentration of 10mg/ml, and delivered by oral gavage for 5

consecutive days at a dose of 50mg/kg/day. The first day of dosing was "Day 1" in the

experimental timeline. Tamoxifen-treated Wnt1F/F and Wnt1F/F; Wnt7bF/F (cre-negative) mice

served as genotype controls for _Wnt1_ and _Wnt1/7b_ knockout experiments, respectively.

Knockdown validation

Bone-specific deletion was confirmed by DNA recombination PCR and knockdown efficiency

was evaluated by RT-qPCR (see Gene Expression below for RT-qPCR methods). Validation

experiments were performed 3 weeks after the first tamoxifen dose, on Day 22. Mice used for

validation experiments were not subjected to tibial loading.

Recombination PCR. In preparation for DNA recombination analysis, skeletal and extra-skeletal

tissues were crushed manually with mortar and pestle, then digested overnight in Proteinase K.

Genomic DNA was isolated using a commercially available kit from Qiagen (DNeasy). 5'-

CTGCCCAGCTGGGTTTCTACTACG-3' (FOR) and 5'-

ACCAGCTGCAGACTCTTGGAATCCG-3' (REV) were used to amplify wild-type Wnt1 DNA

from the intact/conditional allele (800bp). 5'-AGTGAGCTAGTACGGGGTCC-3' (FOR) and

5'-AGGACCATGAACTGATGGCG-3' (REV) were used to amplify the modified/recombined

Wnt1 locus, which produces a *Wnt1* null allele (368bp). 5'-TGACAGAGGATGGGGAGAAG-

3' (FOR) and 5'-GGTCTTTCCAAGGGTGGTCT-3' (REV) were used to amplify wild-type

Wn7b DNA from the intact locus. 5'-GAGGAAGTCAGGCAGGTGTC-3' (FOR) and 5'-

TATCCCACCGATACGCAAAC-3' (REV) were used to amplify the modified/mutant Wnt7b

allele as described previously[16]. PCR reactions were run separately then combined and resolved

by electrophoresis on a 2.5% agarose gel.

Knockdown efficiency qPCR. Tibias were dissected and processed for knockdown efficiency

testing as described under *Gene Expression* (RT-qPCR). PrimeTime assay Mm.PT.58.30187381

(Integrated DNA Technologies, USA) was used to assay *Wnt1* expression in the cortical bone.

117 These primers amplify a region spanning exons 2 and 3, which is deleted in knockout tissues.

118 Deletion of exons 2-3 produces a *Wnt1* null allele; thus, qPCR data reflects the abundance of

119 wild-type *Wnt1* mRNA in the bone. *Wnt7b* was assayed using primers spanning exons 2-3:

120 FOR: 3'-CGGGCAAGAACTCCGAGTAG-5'; REV: 3'-GCGACGAGAAAAGTCGATGC-5'.

121 Exon 3 is deleted in the Wnt7b^{C3} conditional mouse.

122
Tibial MicroCT

Sequential in vivo microCT scans of the tibia were used to evaluate the effects of short-term

Wnt1 and *Wnt1/7b* deletion on cortical bone morphometry. Tibias were scanned at the mid-

diaphysis 24-48 hours before the start of tamoxifen induction (Day 0), followed by a second scan

approximately 3 weeks later on Day 22. Under anesthesia (1-3% isoflurane), the left tibia was

pulled straight and a 2.1 mm region centered 5 mm proximal to the distal TFJ was scanned on a

vivaCT40 with settings optimized for bone (70 kVp, 115 μA, 300 ms integration, 0.4 sigma, 1

support, 1000 projections, 10.5 μm/voxel) (Scanco Medical, Switzerland). Post-scans were

completed on mice in situ after sacrifice. For all scans a 1.05 mm region in the center of the scan

was contoured in the Scanco software and evaluated for cortical bone area, total area, cortical

thickness, polar moment of inertia, and tissue mineral density.

Tibial loading

Starting on Day 22 after tamoxifen induction, mice were subjected to an *in vivo* tibial loading

regimen to stimulate periosteal lamellar bone formation as previously described[17-18]. Briefly, the

right tibias of isoflurane-anesthetized mice were positioned vertically between two fixtures and

an ElectroPuls E1000 instrument (Instron, USA) was used to apply cyclic axial compression (60

cycles/day, 4Hz) on the right tibia for 5 consecutive days. Contralateral left tibias served as non-

loaded controls.

Strain gage analysis was performed on mice that were sacrificed 3 weeks after tamoxifen

induction (Day 22). The right tibia was exposed and cleaned of soft tissue before a single-

element strain gage was glued to the antero-medial surface 5mm proximal to the distal

tibiofibular junction (TFJ). The tibia was axially loaded on a material testing machine (Dynamite

8841) at a peak-to-peak force ranging from -2N to -8N for 12 cycles at 4Hz. LabView data

acquisition software was used to collect data, which was analyzed by linear regression to

calculate the force-strain equation for each sex-genotype group.

Strain-matched and force-matched loading. Force-strain equations for each group were used to

define the loading force necessary to engender strain-matched loading at a peak target strain of

approximately -3000 $\mu\varepsilon$. In males, this required a peak force of -10.5 N for both genotypes. In

females, strain-matched loading required -9.2 N and -5.4 N in control and knockout mice,

respectively. These forces were found to induce a robust lamellar bone formation response in

Wnt1F/F males and females (Fig S3). We loaded an additional set of female *Wnt1* knockout

mice to -9.2 N (-5600 $\mu\varepsilon$) to provide a force-matched comparison with control females, and

determine if increasing strain magnitude induced a greater loading response.

Dynamic histomorphometry

In preparation for bone formation analysis by dynamic histomorphometry, mice were loaded for

5 days (Days 22-26), then dosed with two fluorochromes to label sites of active mineralization.

Calcein green (10 mg/kg; Sigma-Aldrich, St. Louis, MO, USA) and alizarin red (30 mg/kg;

Sigma-Aldrich) were given by IP injection on Days 26 and 31, respectively, followed by

sacrifice on Day 33 (see timeline, Fig 4A). Bilateral tibias (loaded and non-loaded controls) were

de-hydrated through ethanol and embedded in MMA plastic as described previously[19].

Transverse sections from the mid-diaphysis (5mm proximal to the TFJ) were used to capture

20µm z-stack images on a Leica confocal microscope (Leica DMi8) for analysis. Bioquant

analysis software was used to calculate periosteal (Ps) and endocortical (Ec) MS/BS, MAR, and

BFR/BS[20]. Relative bone formation rate (rBFR/BS) was calculated as [loaded – non-loaded].

Relative (r) values were used as a net measure of the effect of loading within each animal.

Gene Expression

Gene expression was analyzed by in situ hybridization and RT-qPCR 3 weeks after tamoxifen

induction, 4-hr after the 5th bout of loading.

RNAScope in situ hybridization. Tibias from loaded Wnt1F/F mice were fixed for 24 hours in

10% neutral buffered formalin (NBF), de-calcified in 14% EDTA, and submitted to the

Musculoskeletal Histology and Morphometry Core (Washington University in St Louis, USA)

for paraffin embedding and sectioning. Bones were sectioned to a thickness of 5µm through the

transverse plane approximately 5mm distal to the tibial plateau. Hybridization was performed

using the HybEZ II Hybridization System (ACDBio) in conjunction with the RNAScope 2.5HD

Assay-BROWN kit. Probes for murine *Wnt1*, *Bmp2*, *Bglap, Axin2, Opg, and Id1* were purchased

from ACDBio (Catalog No. 401091, 406661, 441211, 488961, and 322335). Sections were

counterstained with Hematoxylin and imaged on a Hamamatsu NanoZoomer slide scanning

system at the Alafi Neuroimaging Laboratory (Washington University in St. Louis, USA).

Expression was quantified by scoring cells within the ROI (site of peak compressive strain) as

negative, low, medium, or high. A two-sided Chi-square test was used for statistical analysis.

RT-qPCR. Bilateral tibias were collected and snap-frozen in liquid nitrogen as described

previously[7, 12]. Briefly, tibias were exposed and soft tissues were removed from the bone surface

using forceps, leaving the periosteum intact. Tibias were cut 2mm below the tibial plateau and

1mm below the tibio-fibrular junction to remove the epiphysis and distal tibia. The remainder,

consisting largely of compact cortical bone, was spun at 13,000rpm for 30s to remove bone

marrow. Bones were pulverized with a Mikro Dismembrator, lysed in Trizol reagent, and phase

separated with chloroform reagent. Total RNA was purified using RNeasy Total RNA kit

(Qiagen), and cDNA was prepared from samples with RIN > 6.0 using the iScript cDNA

Synthesis Kit (BioRad). Gene expression was analyzed with SYBR-based reagents on a StepOne

Plus Machine (Applied Biosystems), and reported as a relative expression value (2^{-dCT}) relative

to reference gene *Tbp*.

Statistical Analysis

The main effects of genotype (control vs Wnt1Δ or Wnt1/7bΔ) and loading (non-loaded vs

loaded) and their interaction were analyzed by 2-factor ANOVA, and *post hoc* Sidak's multiple

comparisons test was used for pairwise comparisons (Prism 7.0, GraphPad Software, Inc., La

Jolla, CA, USA). In this analysis the significance of the interaction ("Int") term is crucial

because it indicates whether the effect of loading differed between genotype groups. One-factor

ANOVA was used for outcomes where loading was not a factor (eg knockdown efficiency,

rBFR/BS). Significance was defined at $p<0.05$, with trends noted at $0.05<p<0.10$. Individual data

points and the mean $\pm$ standard deviation are plotted.

Results

Tibial loading induces Wnt1 expression in osteocytes

Tibial loading induces *Wnt1* expression in bone[9, 11-12]. To identify the source of Wnt1 in response to loading, *in situ* hybridization was used to evaluate *Wnt1* RNA expression in the loaded and non-loaded tibias of Wnt1F/F control mice after 5 days of loading (Fig 1A). In loaded tibias *Wnt1* expression was detected in 39.3% ($\pm$ 14.8%) of the osteocytes at the site of peak compressive strain, while in non-loaded control tibias, *Wnt1* expression was detected in only 3.8% ($\pm$ 2.8%) of the osteocytes in the same region of interest ($p<0.0001$, chi-square test) (Fig 1B). In contrast, little expression was observed in the bone lining cells of either loaded or non-loaded bones after 5 days of loading (Fig S1).

Figure 1. Tibial loading increased _Wnt1_ expression in the bone. (A) Tibial _Wnt1_ expression was analyzed by RNAScope _in situ_ hybridization after 5 days of loading. Results representative of n=4. (B) Osteocytes were scored for _Wnt1_ expression at the site of peak compressive strain (lower panels).

An inducible Cre/LoxP approach was used to delete Wnt1 in the Osx-lineage cells of 5-month

old mice

To investigate the role of _Wnt1_ in the skeletal response to loading, a tamoxifen-inducible

Cre/LoxP approach was used to conditionally inactivate _Wnt1_ in the osteoblast-lineage cells of 5-

month old mice. OsxCreERT2; Wnt1F/F mice were dosed with tamoxifen by oral gavage for 5

consecutive days to induce exon 2-3 deletion in the osteoblast lineage. Tamoxifen-treated

Wnt1F/F mice (Cre-negative) served as genotype controls. Gene deletion was confirmed on Day

22 by DNA recombination PCR and knockdown efficiency testing by RT-qPCR (Fig 2A). DNA

recombination PCR indicated that deletion was specific to bone; no deletion occurred in the

extra-skeletal tissues surveyed (Fig 2B). PCR analysis also confirmed that bone-specific DNA

deletion occurred in 5/5 of the knockout mice tested, but not in any (0/4) of the controls (Fig 2C).

Knockdown efficiency testing by qPCR indicated that *Wnt1* mRNA expression was 78% lower

in cortical bones of knockouts relative to controls (Fig 2D).

Figure 2. *Wnt1* **was conditionally deleted in the Osx-expressing cells of adult mice**. (A) Five-month old Wnt1F/F (control) and OsxCreERT2; Wnt1F/F (knockout) mice were dosed with tamoxifen to induce *Wnt1* deletion in the osteoblast lineage. (B-C) DNA Recombination PCR indicated that Wnt1 deletion was specific to the tibias of OsxCreERT2-positive mice. (D) Tibial *Wnt1* RNA expression was 78% lower in *Wnt1* knockouts compared to controls. Individual data points and the mean ± std deviation are shown. n=3-6/group.

Bone area is decreased after short-term Wnt1 deletion

To evaluate the effects of short-term *Wnt1* deletion on cortical bone morphometry, tibias

were serially scanned by *in vivo* microCT 3 weeks apart (Fig 3A). At baseline, there were no

differences in cortical morphology between control and *Wnt1* knockout mice, consistent with no

phenotype prior to induction of gene deletion. But, 22 days after initial tamoxifen dosing, in both

males and females the cortical area and cortical thickness were lower in *Wnt1* knockout mice

than in controls, and in females pMOI was also lower in knockouts (Fig. 3B, 3D, 3F). These

changes are consistent with a reduction in cortical area in *Wnt1* knockouts from day 0 to 22,

whereas cortical area did not change in controls. Similarly, total area was reduced significantly

from day 0 to 22 in female knockouts, and marginally in male knockouts (p=0.123), whereas it

did not change in control mice. In slight contrast, cortical thickness increased from day 0 to 22 in

control mice of both sexes, while in knockout mice it was marginally reduced (p=0.055, males)

or unchanged (p>0.05, females). Therefore, *Wnt1* deletion for only 3 weeks led to reduced

cortical bone mass, due either to loss of bone or reduced accrual of bone.

Figure 3. Cortical bone area decreased after short-term *Wnt1* deletion. The tibias of naïve male and female mice were serially scanned to determine the effects of short-term (3 week) *Wnt1* deletion. Tibias were scanned before tamoxifen induction on Day 0, and again on Day 22. Individual data points and the mean ± std deviation are shown. n=3-6/group.

Loading-induced periosteal bone formation is impaired in Wnt1 knockouts

Next, to evaluate the role of *Wnt1* in the loading response, *in vivo* tibial loading was used

to induce bone formation. Three weeks following 5 days of tamoxifen treatment, mice were

subjected to daily compressive loading on the right tibia for 5 consecutive days and dosed with

calcein and alizarin to label newly mineralizing bone (Fig 4A). Based on *a priori* strain analysis,

the majority of mice were subjected to strain-matched loading to a peak target strain of

approximately -3000 $\mu\epsilon$. Preliminary analysis indicated that this stimulus elicited a robust bone

formation response in Wnt1F/F controls (Fig S3).

In male mice, tibial loading enhanced indices of periosteal (Ps) bone formation in both

control and *Wnt1* knockout groups, but the response to loading was diminished in knockouts.

Two-way ANOVA indicated a significant effect of loading (main effect) for periosteal bone

formation indices, and a significant (or trending) interaction between loading and genotype,

indicating that the effect of load was different in control versus *Wnt1* knockout mice (Fig. 4B-D).

In controls, Ps.MS/BS, Ps.MAR, and Ps.BFR/BS were 4, 2.5, and 9.6-fold higher, respectively,

in loaded vs non-loaded limbs ($p<0.01$) (Fig 4B-4D). By comparison, in knockouts Ps.MS/BS,

Ps.MAR, and Ps.BFR/BS were only 1.9 ($p=0.004$), 1.4 ($p=0.35$), and 2.8-fold ($p=0.16$) higher,

respectively, in loaded vs. non-loaded limbs. We also calculated the relative bone formation rate

(rBFR/BS), defined as BFR/BS$_{Loaded}$ minus BFR/BS$_{Non-Loaded}$, as a measure of loading-induced

bone formation. Periosteal rBFR/BS was 65% lower in knockouts relative to controls ($p=0.037$;

Fig 4E), which further illustrates that the periosteal response to loading was impaired by the loss

of *Wnt1*.

Figure 4. Loading-induced periosteal bone formation was blunted in *Wnt1* knockout males. (A) After 5 days of loading, mice received calcein and alizarin injections in preparation for bone formation analysis by dynamic histomorphometry. (B-D) Periosteal bone formation indices in males subjected to strain-matched loading (-10.5N). Loading increased periosteal bone formation in both groups, albeit to a lesser degree in *Wnt1* knockouts. (E) Relative bone formation rate (rBFR/BS), used as a net index of loading-induced bone formation, was 65% less in *Wnt1* knockouts relative to controls.

In females, we observed similar negative effects of *Wnt1* deletion on the periosteal

response to loading (Fig 5). First, comparing mice loaded to a similar peak strain (~3000 με;

"strain matched"), loading stimulated increases in periosteal bone formation in control but not

knockout (Wnt1Δ.) mice. In controls, Ps.MS/BS, Ps.MAR and Ps.BFR/BS were 2.8, 3.5 and 8.3-

fold higher, respectively, in the loaded vs non-loaded bones ($p \leq 0.0001$) (Fig 5A-5C). By

comparison, in knockout females loading-associated changes in Ps.MS/BS, Ps.MAR, and

Ps.BFR/BS were minimal and not significant. Moreover, periosteal relative bone formation rate

(rBFR/BS) was 94% lower in knockouts relative to controls after strain-matched loading

(p<0.0001; Fig 5D).

To achieve a strain-matched stimulus of -3000με, the force applied to female *Wnt1*

knockout mice was 40% less than the force applied to control mice (-5.4N vs. -9.2N; Fig S2),

which is consistent with difference in bone morphology (lower Ct.Ar and pMOI in knockouts;

Fig 3B, F). Because this level of stimulus did not induce a significant loading response in female

knockouts (Wnt1Δ^{SM} group), we asked if a higher magnitude stimulus might induce a response.

Thus, we subjected a second cohort of knockout females (Wnt1Δ^{FM}) to -9.2N to achieve a "force

match" (compared to control); this engendered an estimated -5600με peak strain (Fig S2), a level

that typically induces a woven bone response[12]. Knockout females subjected to force-matched

loading did display a significant periosteal loading response. Ps.MS/BS, Ps.MAR, and

Ps.BFR/BS were 1.8, 2.4 and 4-fold higher, respectively, in loaded vs non-loaded limbs (Fig 5A-

5C). Notably, these values were nominally less than the respective fold-increases in control mice

(2.8, 3.5 and 8.3), and Ps.MAR and Ps.BFR/BS were significantly less in loaded limbs of force-

matched knockout females compared to controls. Moreover, periosteal rBFR/BS was 56% lower

in knockouts relative to controls after force-matched loading (*p*<0.01; Fig 5D). Therefore,

despite being loaded to 85% higher peak strain the control mice, *Wnt1* knockout females still had

a blunted periosteal response to loading.

Figure 5. Loading-induced periosteal bone formation was blunted in *Wnt1* knockout females. (A-C) Periosteal bone formation indices in females. Wnt1F/F females were loaded to -9.2N to engender a peak target strain of -3040με. *Wnt1* knockouts were subjected to strain-matched loading (SM) or force-matched loading (FM) by applying either -5.4N or -9.2N, respectively.

Finally, in contrast to the robust loading response elicited on the periosteal surface of

control mice, tibial loading did not induce a strong endocortical response, which is consistent

with the lower values of strain engendered on this surface. In males, indices of endocortical bone

formation were not significantly greater in loaded compared to non-loaded limbs, for both

control and knockout mice (Fig S4). In female control mice, loading did stimulate a modest

increase in Ec.MAR ($p<0.05$, loaded vs. non-loaded) and Ec.BFR/BS ($p=0.052$), but not in

strain-matched *Wnt1* knockout mice (Fig S5). Force-matched loading did induce a significant

increase in Ec.MAR and Ec.BFR in knockout females, with values in loaded limbs not different

from the loaded limbs of control mice.

Induction of osteogenic genes was diminished in the bones of Wnt1 knockouts

Gene expression analysis was performed 4-hr after the 5th bout of loading. Congruent

with previous reports [11-12], we found that loading potently induced *Wnt1* expression in the bone

(Fig 6A). RT-qPCR analysis indicated that *Wnt1* expression was 6.6-fold higher in the loaded vs

non-loaded tibias of Wnt1F/F controls ($p<0.0001$). In contrast, loading had no effect on the

abundance of *Wnt1* mRNA in the tibias of *Wnt1* knockouts ($p=0.999$) (Fig 6A), consistent with

bone-targeted knockout.

Expression of genes associated with osteo-induction, osteoblast maturation, and matrix

synthesis were also analyzed. RNAScope *in situ* hybridization showed that loading increased

Bmp2 expression in osteocytes and *Bglap* expression on the bone surface (Fig S6). These

findings were supported by qPCR, which showed that *Bmp2* and *Bglap* were 1.8 and 3.7-fold

higher, respectively, in the loaded vs non-loaded bones of Wnt1F/F controls ($p<0.001$) (Fig 6B,

6D). In contrast, loading did not significantly increase *Bmp2* or *Bglap* expression in knockouts.

Similarly, loading increased expression of the matrix gene *Col1a1* in the tibias of control mice

(5.5-fold higher in loaded vs non-loaded, *p*<0.001), while in knockouts *Col1a1* did not increase

with loading (Fig 6C).

Figure 6. Loading-induced upregulation of bone formation genes was impaired in *Wnt1* knockouts. (A) *Wnt1* induction was observed in Wnt1F/F controls, but not in *Wnt1* knockouts. (B-D) Loading-induced *Bmp2*, *Col1a1*, and *Bglap* upregulation was blunted in *Wnt1* knockouts relative to controls. n=4-5 per sex per genotype in each group for a total of n=9/genotype. Two-factor ANOVA was used to evaluate the effect of tibial loading ("Load") and genotype ("Gen"), and their interaction ("Int"), on gene expression.

Finally, we assayed several reported Wnt-target genes. Neither loading nor genotype had

a significant effect on the expression of *Axin2* or *Nkd2* in the bone (Fig S7). In contrast, gene

expression analysis by two-factor ANOVA revealed that loading had a modest but significant

effect on the expression of *Opg*. In controls, *Opg* was 1.4-fold higher in loaded vs non-loaded

limbs (*p*=0.077), while in knockouts *Opg* was only 1.2-fold higher (*p*=0.482) (Fig S7).

In sum, these results indicate that *Wnt1* induction in Osx-lineage cells is required for the

bone anabolic effects of tibial loading.

The skeletal phenotype caused by Wnt1 deletion was recapitulated in Wnt1/7b knockout mice

Because *Wnt7b* is osteo-anabolic[14] and is induced in the bone by loading[12], we asked

whether the phenotypes observed in *Wnt1* knockouts would be exacerbated by also targeting

Wnt7b. We confirmed recombination of *Wnt1* and *Wnt7b* genes in tibias of *Wnt1/7b* knockout

mice (tamoxifen-treated OsxCreERT2; Wnt1F/F; Wnt7bF/F) (Fig.S8A). We also confirmed

knockdown of *Wnt1* (-63% in the loaded tibias of Wnt1/7bΔ mice vs control, *p*=0.083) and

Wnt7b (-87% in the loaded tibias of *Wnt1/7b* knockouts vs control, *p*=0.194) at the mRNA level

(Fig S9A).

The cortical bone phenotype observed in *Wnt1* knockouts was recapitulated in *Wnt1/7b*

knockouts. In brief, relative to Wnt1F/F; Wnt7bF/F controls, the bone morphology of *Wnt1/7b*

knockouts was not different on Day 0 (prior to tamoxifen induction), but on Day 22 cortical bone

area, cortical thickness, and pMOI were less in *Wnt1/7b* knockouts. In general, we observed

similar magnitude differences in *Wnt1/7b* knockout mice compared to its control, as observed in

Wnt1 knockouts compared to its control. For example, cortical area was 10-15% less in *Wnt1*

knockouts vs. control (Fig. 3B), and 15-18% less in *Wnt1/7b* knockouts vs. control (Fig. S8C).

Finally, when *Wnt1/7b* knockout mice and their controls were subjected to strain-matched

tibial loading (approx. -3000 με), *Wnt1/7b* knockouts had impaired periosteal bone formation,

similar to *Wnt1* single knockouts. In controls, loading significantly enhanced Ps.MS/BS and

Ps.MAR, resulting in an overall increase in Ps.BFR/BS of 11 and 5.6-fold in the loaded bones of

male and female mice, respectively, compared to non-loaded controls (*p*<0.0001; Figure S9B).

By comparison, loading did not significantly increase bone formation indices in *Wnt1/7b*

knockouts. Consequently, the relative bone formation rate (rBFR/BS) in *Wnt1/7b* knockout mice

was 90% lower in males and 86% lower in females compared to their respective controls. These

data further support the finding that *Wnt1* is a crucial mediator of loading-induced bone

formation.

Discussion

Mechanical loading on the skeleton stimulates bone formation. To investigate the role of *Wnt1* in the skeletal response to loading, we tamoxifen-dosed 5-month old OsxCreERT2; Wnt1F/F mice to inactivate *Wnt1* in the osteoblast lineage, then subjected mice to *in vivo* tibial loading 21 days later. Over 3 weeks, bone area decreased 10-15% in *Wnt1* knockouts, concomitant with a 78% reduction in tibial *Wnt1* expression (Figs 2D, 3B). In Wnt1F/F control mice, loading potently increased *Wnt1* expression in the bone and *in situ* hybridization showed that this induction occurred in osteocytes (Figs 1A, 6A). Loading-induced *Wnt1* upregulation was associated with a 8.4 to 9.6-fold increase in periosteal bone formation rate (Ps.BFR/BS) in control mice (Figs 4D, 5C). In knockouts, Osx-targeted *Wnt1* deletion blocked *Wnt1* upregulation (Fig 6A), leading to a 65 to 94% lower periosteal bone formation response (Figs 4E, 5D). In sum, these findings show that *Wnt1* in the Osx-lineage mediates the bone anabolic effects of skeletal loading in adult mice.

Loading increases Wnt1 expression in osteocytes

Whole-bone transcriptional profiling studies have shown that mechanical loading increases the expression of Wnt family genes – including *Wnt1* and *Wnt7b* – in the bone[9-10]. Subsequently, Harris and Silva used laser capture microdissection to isolate intracortical bone samples from loaded and non-loaded bones of C57Bl/6 mice, and showed that *Wnt1* in these osteocyte-enriched samples was upregulated by loading[11]. Congruent with these results, our gene expression analyses showed that *Wnt1* was 6.6-fold higher in the loaded vs non-loaded bones of Wnt1[F/F] mice after 5 days of loading (Fig 6A). Additionally, analysis by RNAscope® *in situ* hybridization showed that *Wnt1* induction occurred primarily in osteocytes embedded in the

intra-cortical matrix rather than in cells lining the bone surface (i.e., osteoblasts) (Figs 1A and

S1). Analysis of *in situ* hybridization further revealed that loading increased not only the total

percentage of *Wnt1*-positive cells in the bone, but also the magnitude of *Wnt1* expression within

those cells (Fig 1B). Together, these data show that mechanical loading stimulates *Wnt1*

expression in the bone, particularly in osteocytes.

Loading-induced bone formation is reduced in Wnt1 knockouts

OsxCreERT2 was used to inactivate *Wnt1* in the Osx-lineage cells of 5-month old mice.

Previously, we used a similar tamoxifen dosing strategy for Cre reporter analysis in 5-month old

OsxCreERT2; Rosa-Ai9 mice, which showed robust TdTomato reporter expression on the bone

surface (osteoblasts) and within the cortical bone (osteocytes) within days of tamoxifen

treatment, indicating that OsxCreERT2 targets both osteoblasts and osteocytes in the bones of

adult mice[7]. OsxCreERT2-mediated *Wnt1* deletion completely neutralized the effect of loading

on *Wnt1* expression in the bone (Fig 6A). While tibial *Wnt1* expression increased 6.6-fold in

loaded bones of Wnt1F/F controls, *Wnt1* expression remained unchanged in the loaded vs non-

loaded limbs of *Wnt1* knockouts. Additionally, tibial loading had a robust inductive effect on the

expression of *Col1a1*, *Bmp2*, and *Bglap* in the bones of control but not knockout mice, indicating

that loading-induced *Wnt1* expression was required for the regulation of these genes (Figs 6B-C).

In combination with our *in situ* hybridization findings, these results suggest that loading

stimulates osteoblast differentiation/activity on the bone surface (*Bglap*, Fig S6C) by inducing

Wnt1 ligand expression in osteocytes (Figs 1A, S6A-B).

At the tissue level, *Wnt1* inactivation in the Osx-lineage reduced the periosteal response

to loading. In males, periosteal bone formation rate was higher in the loaded vs non-loaded bones

of all mice, independent of genotype. However, the relative bone formation rate – which reflects

the overall magnitude of the loading response – showed that the periosteal response to loading

was 65% lower in knockouts (Fig 4E). In females, strain-matched loading increased all indices of

periosteal bone formation in control but not knockout mice (Fig 5A-C). Compared to Wnt1F/F

controls, the periosteal response to loading (i.e., Ps.rBFR/BS) was 94% lower in strain-matched

Wnt1 knockouts (Fig 5D). Moreover, analysis of a force-matched cohort (-9.2 N, *Wnt1* knockout

females) showed that although a loading response could be elicited with greater strain magnitude

(-5600ue vs -3000ue), the periosteal loading response was still 39% less in knockouts vs controls

(Fig 5D). Together, these results demonstrate that *Wnt1* is required for loading-induced

periosteal bone formation.

Bone area decreased in Wnt1 knockouts after 21 days deletion

Independent of loading, short-term *Wnt1* deletion (3 weeks) in the Osx-lineage was

associated with decreased bone area and cortical thickness (Fig 2B, 2D). These findings are

congruent with a report by Wang, et al. which showed that deleting *Wnt1* in the Osx-lineage

impaired periosteal bone growth in young mice. Using a Dox-repressible system, Wang et al.

showed that inactivating *Wnt1* in the Osx-lineage of 4-week old mice produced measurable

deficits in the skeleton 4 and 8 weeks later. Analysis of distal tibias from 12 week old mice

revealed that these deficits were caused by a reduction in periosteal bone formation, while

changes to endocortical bone formation were not significant[21].

Similar findings were reported by Joeng, et al. and Luther, et al. Mice lacking *Wnt1* in

late osteoblasts (Dmp1Cre; Wnt1F/F) had skeletal deficits – including reduced Ct.Th, Tb.N, and

Tb.Th – 2 months after birth[13], while mice lacking *Wnt1* in osteo-progenitors (Runx2Cre;

Wnt1F/F) exhibited BV/TV and Ct.Th deficits in the femur at 24 weeks, leading to increased risk

of spontaneous fractures[22]. Hypomorphic *Wnt1* variants have also been liked to low bone mass

and spontaneous fractures in mice[23]. Our study extends these findings by showing that short-term

Wnt1 deletion in adults leads to bone loss. Thus, osteoblast/osteocyte *Wnt1* is required for adult

bone homeostasis.

The skeletal response to bone anabolic loading declines with age[24-26]. For example, tibial

loading enhances periosteal bone formation in adult mice of different ages; however, the

magnitude of the osteogenic response to loading is significantly lower in aged 22-month old vs.

young adult 5-month old mice[12]. To understand the basis for these differences, Chermside-

Scabbo, et al. used RNASeq to characterize the transcriptional responses to loading in 5 and 22-

month old C57Bl/6 mice after 1, 3, and 5 days of loading[10]. In both age groups, *Wnt1* emerged as

a top 10 upregulated gene on days 1 and 3. The fold-change difference for *Wnt1* in 5-month old

mice was 5.8 and 6.7 on days 1 and 3, respectively, while in 22-month old mice the fold-change

difference for *Wnt1* was 1.4 and 2.3. Thus, loading increased *Wnt1* in both age groups, albeit to a

lesser degree in aged mice – concomitant with a diminished osteogenic response to loading.

Taken together, these findings suggest that diminished induction of *Wnt1* by loading may

contribute to reduced loading-induced bone formation in the bones of aged mice.

The bone anabolic function of *Wnt1* in the Osx-lineage is further supported by our

findings in *Wnt1/7b* double conditional knockout mice, which exhibited cortical bone deficits 21

days post-deletion (Fig S8C). Moreover, like *Wnt1* knockouts, *Wnt1/7b* knockouts did not

exhibit a loading-induced increase in *Wnt1* expression and had a blunted periosteal response to

loading (Fig S9B). In general, the magnitude of the deficits in *Wnt1/7b* knockouts were

comparable to those in *Wnt1* knockouts. Importantly, while we did see evidence of *Wnt7b* gene

recombination in *Wnt1/7b* double knockout mice (Fig. S8A), we did not confirm significant

knockdown of *Wnt7b* at the mRNA level (Fig. S9A). Thus, we are not able to infer whether

Wnt7b has a functional role in loading-induced bone formation. Additional studies, perhaps using

a different *Wnt7b* floxed mouse, are needed to address this question. Nonetheless, the consistent

deficits in loading-induced bone formation in *Wnt1* and *Wnt1/7b* knockout mice show that in two

different mouse lines, *Wnt1* in the Osx-lineage was essential for the response to loading.

In conclusion, our findings support a growing body of evidence that *Wnt1* is essential for

bone homeostasis in the adult skeleton. In particular, we showed that mechanical loading

induced *Wnt1* expression in osteocytes, and that loading-induced *Wnt1* expression in the Osx-

lineage was required for the periosteal bone formation response to tibial loading.

 Supplemental Figures

Figure S1. *Wnt1* expression in a loaded tibia after 5 days of loading. *Wnt1* was observed in osteocytes on both the tensile and compressive sides of the bone, including osteocytes near the bone surface (red arrowheads). *Wnt1* expression was not observed in the periosteal layer (white arrowheads). Results representative of n=4.

Figure S2. Strain gage analysis was used to experimentally determine the force-strain relationships in male (A) and female (B) mice 3 weeks after tamoxifen induction. In males, strain-matched loading was achieved by loading controls and knockouts to -10.5N, which engendered a peak target strain of -2800µℇ in both genotypes. In females, strain-matched (SM) loading was achieved by loading controls to -9.2N (-3040µℇ) and knockouts to -5.4N (-3200µℇ). Female knockouts in the force-matched (FM) group were subjected to a peak target strain of -5600µℇ.

Figure S3 (A-B) Wnt1F/F mice were loaded to a peak target strain of approximately -3000µℇ (-10.5N males; -9.2N females) to stimulate lamellar bone formation. (C) A Wnt1F/F female loaded to -5600µℇ (-15N) had a woven bone formation response to loading. Scale bars represent 500um.

Figure S4. Endocortical bone formation indices in males. In both groups, loading had a negligible effect on endocortical indices.

Figure S5. Endocortical bone formation indices in females.

Figure S6. Supplemental RNAScope. After 5 days of loading, bilateral tibias from Wnt1F/F mice were harvested for analysis by RNAScope *in situ* hybridization. (A) Loading increased osteo-inductive *Bmp2* expression in the bone, particularly in osteocytes located at the site of peak compressive strain. (B) Loading also increased the expression of Bmp target gene *Id1* on the bone surface. (C) Robust *Bglap* expression was observed on the surface of loaded bones. Images representative of n=3.

Figure S7. Wnt target genes. Gene expression analysis by RNAScope (A-B) and RT-qPCR (C-E) showed that *Axin2* and *Nkd2* were not different in the loaded vs non-loaded bones of control or knockout mice. Loading increased *Opg* in the tibias of Wnt1F/F controls but not *Wnt1* knockouts. RNAScope images representative of n=3.

Figure S8.   **The skeletal deficits associated with *Wnt1* deletion were re-capitulated in *Wnt1/7b* knockouts**. OsxCreERT2; Wnt1F/F; Wnt7bF/F (*Wnt1/7b* knockout) mice were tamoxifen-treated at 5-months old to induce *Wnt1* and *Wnt7b* deletion in the Osx-lineage. (A) DNA recombination PCR was performed on Day 22 to confirm bone-specific deletion of *Wnt1* (exon 2-3) and *Wnt7b* (exon 1). (B-G) Serial microCT scans were used to evaluate the effect of *Wnt1/7b* deletion (Wnt1/7bΔ) over a 3 week period. Tamoxifen-treated cre-negative Wnt1F/F; Wnt7bF/F mice served as control. n=6-9/group.

Figure S9. **Loading-induced bone formation was reduced in *Wnt1/7b* knockouts**. (A) *Wnt1* and *Wnt7b* expression was assayed 4 hours after the 5th bout of loading. (B) Loading-induced periosteal bone formation was reduced in Wnt1/7b knockout male (top) and female (bottom) mice, relative to sex-matched controls.

Data Availability Statement

The data that support the findings of this study are openly available in Biorxiv at

https://doi.org/10.1101/2022.02.28.482178.

Conflict of Interest Statement

LYL, NM, CCS, JTS, KJ, and BL have no financial or non-financial competing interests to

disclose. MJS is on the editorial board at *Bone*, *Journal of Orthopaedic Research*, and *Calcified*

Tissue International, and served on the board of directors at the Orthopaedic Research Society

(2018-2022). RC serves as Editor-in-Chief for *Journal of Bone and Mineral Research* and has

received research support from Ultragenyx.

Author Contributions

Authors' roles: Study design: LYL, NM, CCS, and MJS. Study conduct and data collection:

LYL, NM, CCS, and JTS. Data analysis and interpretation: LYL, NM, CCS, KJ, RC, BL, and

MJS. Drafting manuscript: LYL and MJS.

Acknowledgments

This work was supported by NIH grants R01 AR047867, T32 AR060719 and the Washington

University Musculoskeletal Research Center (P30 AR074992). We thank Crystal Idleburg and

Samantha Coleman for histology support.

References

1. Robling AG, Niziolek PJ, Baldridge LA, Condon KW, Allen MR, Alam I, Mantila SM,

Gluhak-Heinrich J, Bellido TM, Harris SE, Turner CH. Mechanical stimulation of bone in

vivo reduces osteocyte expression of Sost/sclerostin. *J Biol Chem*. 2008 Feb 29;283(9):5866-

75. doi: 10.1074/jbc.M705092200. Epub 2007 Dec 17. PMID: 18089564.

2. Tu X, Rhee Y, Condon KW, et al. Sost downregulation and local Wnt signaling are required

for the osteogenic response to mechanical loading. *Bone*. 2012;50(1):209-217.

doi:10.1016/j.bone.2011.10.025

3. Sawakami K, Robling AG, Ai M, Pitner ND, Liu D, Warden SJ, Li J, Maye P, Rowe DW,

Duncan RL, Warman ML, Turner CH. The Wnt co-receptor LRP5 is essential for skeletal

mechanotransduction but not for the anabolic bone response to parathyroid hormone

treatment. *J Biol Chem*. 2006 Aug 18;281(33):23698-711. doi: 10.1074/jbc.M601000200.

Epub 2006 Jun 20. PMID: 16790443.

4. Saxon LK, Jackson BF, Sugiyama T, Lanyon LE, Price JS. Analysis of multiple bone

responses to graded strains above functional levels, and to disuse, in mice in vivo show that

the human Lrp5 G171V High Bone Mass mutation increases the osteogenic response to

loading but that lack of Lrp5 activity reduces it. *Bone*. 2011 Aug;49(2):184-93. doi:

10.1016/j.bone.2011.03.683. Epub 2011 Mar 16. PMID: 21419885; PMCID: PMC3121951.

5. Niziolek PJ, Farmer TL, Cui Y, Turner CH, Warman ML, Robling AG. High-bone-mass-

producing mutations in the Wnt signaling pathway result in distinct skeletal phenotypes.

Bone. 2011 Nov;49(5):1010-9. doi: 10.1016/j.bone.2011.07.034. Epub 2011 Aug 9. PMID:

21855668; PMCID: PMC3412139.

6. Clevers H, Nusse R. Wnt/β-catenin signaling and disease. *Cell*. 2012 Jun 8;149(6):1192-205.

doi: 10.1016/j.cell.2012.05.012. PMID: 22682243.

7. Lawson LY, Brodt MD, Migotsky N, Chermside-Scabbo CJ, Palaniappan R, Silva MJ.

Osteoblast-Specific Wnt Secretion Is Required for Skeletal Homeostasis and Loading-

Induced Bone Formation in Adult Mice. *J Bone Miner Res*. 2021 Sep 20. doi:

10.1002/jbmr.4445. Epub ahead of print. PMID: 34542191.

8. Bänziger C, Soldini D, Schütt C, Zipperlen P, Hausmann G, Basler K. Wntless, a conserved

membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell*. 2006

May 5;125(3):509-22. doi: 10.1016/j.cell.2006.02.049. PMID: 16678095.

9. Kelly NH, Schimenti JC, Ross FP, van der Meulen MC. Transcriptional profiling of cortical

versus cancellous bone from mechanically-loaded murine tibiae reveals differential gene

expression. *Bone*. 2016 May;86:22-9. doi: 10.1016/j.bone.2016.02.007. Epub 2016 Feb 12.

PMID: 26876048; PMCID: PMC4833881.

10. Chermside-Scabbo CJ, Harris TL, Brodt MD, Braenne I, Zhang B, Farber CR, Silva MJ. Old

Mice Have Less Transcriptional Activation But Similar Periosteal Cell Proliferation

Compared to Young-Adult Mice in Response to in vivo Mechanical Loading. *J Bone Miner

Res*. 2020 Sep;35(9):1751-1764. doi: 10.1002/jbmr.4031. Epub 2020 Jun 1. PMID:

32311160; PMCID: PMC7486279.

11. Harris TL, Silva MJ. Gene expression of intracortical bone demonstrates loading-induced

increases in Wnt1 and Ngf and inhibition of bone remodeling processes. *Bone*. 2021

Sep;150:116019. doi: 10.1016/j.bone.2021.116019. Epub 2021 May 21. PMID: 34023542;

PMCID: PMC8408835.

12. Holguin N, Brodt MD, Silva MJ. Activation of Wnt Signaling by Mechanical Loading Is

Impaired in the Bone of Old Mice. *J Bone Miner Res*. 2016 Dec;31(12):2215-2226. doi:

10.1002/jbmr.2900. Epub 2016 Sep 7. PMID: 27357062; PMCID: PMC5397287.

13. Joeng KS, Lee YC, Lim J, Chen Y, Jiang MM, Munivez E, Ambrose C, Lee BH. Osteocyte-

specific WNT1 regulates osteoblast function during bone homeostasis. *J Clin Invest*. 2017

Jun 30;127(7):2678-2688. doi: 10.1172/JCI92617. Epub 2017 Jun 19. PMID: 28628032;

PMCID: PMC5490765.

14. Song D, He G, Song F, Wang Z, Liu X, Liao L, Ni J, Silva MJ, Long F. Inducible expression

of Wnt7b promotes bone formation in aged mice and enhances fracture healing. *Bone Res*.

2020 Feb 3;8:4. doi: 10.1038/s41413-019-0081-8. PMID: 32047703; PMCID: PMC6997361.

15. Maes C, Kobayashi T, Kronenberg HM. A novel transgenic mouse model to study the

osteoblast lineage in vivo. *Ann N Y Acad Sci*. 2007 Nov;1116:149-64. doi:

10.1196/annals.1402.060. PMID: 18083926.

16. Rajagopal J, Carroll TJ, Guseh JS, Bores SA, Blank LJ, Anderson WJ, Yu J, Zhou Q,

McMahon AP, Melton DA. Wnt7b stimulates embryonic lung growth by coordinately

increasing the replication of epithelium and mesenchyme. *Development*. 2008

May;135(9):1625-34. doi: 10.1242/dev.015495. Epub 2008 Mar 26. PMID: 18367557;

PMCID: PMC2556060.

17. Sun D, Brodt MD, Zannit HM, Holguin N, Silva MJ. Evaluation of loading parameters for

murine axial tibial loading: Stimulating cortical bone formation while reducing loading

duration. J Orthop Res. 2018 Feb;36(2):682-691. doi: 10.1002/jor.23727. Epub 2017 Oct 9.

PMID: 28888055; PMCID: PMC5839947.

18. Main RP, Shefelbine SJ, Meakin LB, Silva MJ, van der Meulen MCH, Willie BM. Murine

Axial Compression Tibial Loading Model to Study Bone Mechanobiology: Implementing the

Model and Reporting Results. J Orthop Res. 2020 Feb;38(2):233-252. doi:

10.1002/jor.24466. Epub 2019 Oct 23. PMID: 31508836.

19. Erben RG. Embedding of bone samples in methylmethacrylate: an improved method suitable

for bone histomorphometry, histochemistry, and immunohistochemistry. J Histochem

Cytochem. 1997 Feb;45(2):307-13. doi: 10.1177/002215549704500215. PMID: 9016319.

20. Dempster DW, Compston JE, Drezner MK, Glorieux FH, Kanis JA, Malluche H, Meunier

PJ, Ott SM, Recker RR, Parfitt AM. Standardized nomenclature, symbols, and units for bone

histomorphometry: a 2012 update of the report of the ASBMR Histomorphometry

Nomenclature Committee. *J Bone Miner Res*. 2013 Jan;28(1):2-17. doi: 10.1002/jbmr.1805.

PMID: 23197339; PMCID: PMC3672237.

21. Wang F, Rummukainen P, Heino TJ, Kiviranta R. Osteoblastic Wnt1 regulates periosteal

bone formation in adult mice. *Bone*. 2021 Feb;143:115754. doi:

10.1016/j.bone.2020.115754. Epub 2020 Nov 13. PMID: 33189914.

22. Luther J, Yorgan TA, Rolvien T, Ulsamer L, Koehne T, Liao N, Keller D, Vollersen N,

Teufel S, Neven M, Peters S, Schweizer M, Trumpp A, Rosigkeit S, Bockamp E, Mundlos S,

Kornak U, Oheim R, Amling M, Schinke T, David JP. Wnt1 is an Lrp5-independent bone-

anabolic Wnt ligand. *Sci Transl Med*. 2018 Nov 7;10(466):eaau7137. doi:

10.1126/scitranslmed.aau7137. PMID: 30404864.

23. Joeng KS, Lee YC, Jiang MM, Bertin TK, Chen Y, Abraham AM, Ding H, Bi X, Ambrose

CG, Lee BH. The swaying mouse as a model of osteogenesis imperfecta caused by WNT1

mutations. *Hum Mol Genet.* 2014 Aug 1;23(15):4035-42. doi: 10.1093/hmg/ddu117. Epub

2014 Mar 14. PMID: 24634143; PMCID: PMC4082367.

24. Turner CH, Takano Y, Owan I. Aging changes mechanical loading thresholds for bone

formation in rats. J Bone Miner Res. 1995 Oct;10(10):1544-9. doi:

10.1002/jbmr.5650101016. PMID: 8686511.

25. Birkhold AI, Razi H, Duda GN, Weinkamer R, Checa S, Willie BM. The influence of age on

adaptive bone formation and bone resorption. *Biomaterials*. 2014 Nov;35(34):9290-301. doi:

10.1016/j.biomaterials.2014.07.051. Epub 2014 Aug 13. PMID: 25128376.

26. Meakin LB, Galea GL, Sugiyama T, Lanyon LE, Price JS. Age-related impairment of bones'

adaptive response to loading in mice is associated with sex-related deficiencies in osteoblasts

but no change in osteocytes. *J Bone Miner Res*. 2014;29(8):1859-1871.

doi:10.1002/jbmr.2222

Figure captions

Figure 1. **Tibial loading increased *Wnt1* expression in the bone**. (A) Tibial *Wnt1* expression was analyzed by RNAScope *in situ* hybridization after 5 days of loading. Results representative of n=4. (B) Osteocytes were scored for *Wnt1* expression at the site of peak compressive strain (lower panels).

Figure 2. *Wnt1* **was conditionally deleted in the Osx-expressing cells of adult mice**. (A) Five-month old Wnt1F/F (control) and OsxCreERT2; Wnt1F/F (knockout) mice were dosed with tamoxifen to induce *Wnt1* deletion in the osteoblast lineage. (B-C) DNA Recombination PCR indicated that Wnt1 deletion was specific to the tibias of OsxCreERT2-positive mice. (D) Tibial *Wnt1* RNA expression was 78% lower in *Wnt1* knockouts compared to controls. Individual data points and the mean ± std deviation are shown. n=3-6/group.

Figure 3. **Cortical bone area decreased after short-term *Wnt1* deletion**. The tibias of naïve male and female mice were serially scanned to determine the effects of short-term (3 week) *Wnt1* deletion. Tibias were scanned before tamoxifen induction on Day 0, and again on Day 22. Individual data points and the mean ± std deviation are shown. n=3-6/group.

Figure 4. **Loading-induced periosteal bone formation was blunted in *Wnt1* knockout males**. (A) After 5 days of loading, mice received calcein and alizarin injections in preparation for bone formation analysis by dynamic histomorphometry. (B-D) Periosteal bone formation indices in males subjected to strain-matched loading (-10.5N). Loading increased periosteal bone formation in both groups, albeit to a lesser degree in *Wnt1* knockouts. (E) Relative bone formation rate (rBFR/BS), used as a net index of loading-induced bone formation, was 65% less in *Wnt1* knockouts relative to controls.

Figure 5. **Loading-induced periosteal bone formation was blunted in *Wnt1* knockout**

females. (A-C) Periosteal bone formation indices in females. Wnt1F/F females were loaded to -

9.2N to engender a peak target strain of -3040µℇ. *Wnt1* knockouts were subjected to strain-

matched loading (SM) or force-matched loading (FM) by applying either -5.4N or -9.2N,

respectively.

Figure 6. **Loading-induced upregulation of bone formation genes was impaired in *Wnt1***

knockouts. (A) *Wnt1* induction was observed in Wnt1F/F controls, but not in *Wnt1* knockouts.

(B-D) Loading-induced *Bmp2*, *Col1a1*, and *Bglap* upregulation was blunted in *Wnt1* knockouts

relative to controls. n=4-5 per sex per genotype in each group for a total of n=9/genotype. Two-

factor ANOVA was used to evaluate the effect of tibial loading ("Load)" and genotype ("Gen"),

and their interaction ("Int"), on gene expression.

Figure S1. *Wnt1* expression in a loaded tibia after 5 days of loading. *Wnt1* was observed in

osteocytes on both the tensile and compressive sides of the bone, including osteocytes near the

bone surface (red arrowheads). *Wnt1* expression was not observed in the periosteal layer (white

arrowheads). Results representative of n=4.

Figure S2. Strain gage analysis was used to experimentally determine the force-strain

relationships in male (A) and female (B) mice 3 weeks after tamoxifen induction. In males,

strain-matched loading was achieved by loading controls and knockouts to -10.5N, which

engendered a peak target strain of -2800µℇ in both genotypes. In females, strain-matched (SM)

loading was achieved by loading controls to -9.2N (-3040µℇ) and knockouts to -5.4N (-3200µℇ).

Female knockouts in the force-matched (FM) group were subjected to a peak target strain of -

5600µℇ.

Figure S3 (A-B) Wnt1F/F mice were loaded to a peak target strain of approximately -3000µℇ (-

10.5N males; -9.2N females) to stimulate lamellar bone formation. (C) A Wnt1F/F female loaded

to -5600µℇ (-15N) had a woven bone formation response to loading. Scale bars represent 500um.

Figure S4. Endocortical bone formation indices in males. In both groups, loading had a

negligible effect on endocortical indices.

Figure S5. Endocortical bone formation indices in females.

Figure S6. Supplemental RNAScope. After 5 days of loading, bilateral tibias from Wnt1F/F

mice were harvested for analysis by RNAScope *in situ* hybridization. (A) Loading increased

osteo-inductive *Bmp2* expression in the bone, particularly in osteocytes located at the site of peak

compressive strain. (B) Loading also increased the expression of Bmp target gene *Id1* on the

bone surface. (C) Robust *Bglap* expression was observed on the surface of loaded bones. Images

representative of n=3.

Figure S7. Wnt target genes. Gene expression analysis by RNAScope (A-B) and RT-qPCR (C-

E) showed that *Axin2* and *Nkd2* were not different in the loaded vs non-loaded bones of control

or knockout mice. Loading increased *Opg* in the tibias of Wnt1F/F controls but not *Wnt1*

knockouts. RNAScope images representative of n=3.

Figure S8. The skeletal deficits associated with *Wnt1* deletion were re-capitulated in

***Wnt1/7b* knockouts**. OsxCreERT2; Wnt1F/F; Wnt7bF/F (*Wnt1/7b* knockout) mice were

tamoxifen-treated at 5-months old to induce *Wnt1* and *Wnt7b* deletion in the Osx-lineage. (A)

DNA recombination PCR was performed on Day 22 to confirm bone-specific deletion of *Wnt1*

(exon 2-3) and *Wnt7b* (exon 1). (B-G) Serial microCT scans were used to evaluate the effect of

Wnt1/7b deletion (Wnt1/7bΔ) over a 3 week period. Tamoxifen-treated cre-negative Wnt1F/F;

Wnt7bF/F mice served as control. n=6-9/group.

<u>Figure S9.</u> Loading-induced bone formation was reduced in *Wnt1/7b* knockouts. (A) *Wnt1*

and *Wnt7b* expression was assayed 4 hours after the 5th bout of loading. (B) Loading-induced

periosteal bone formation was reduced in Wnt1/7b knockout male (top) and female (bottom)

mice, relative to sex-matched controls.