Arctic evergreening - encroaching allelopathy in a management blind spot

Tuomi M.W.[1][*][†], Utsi T.Aa.[2], Yoccoz N.[1], Armstrong C.W.[3], Gonzalez V.[4], Hagen S.B.[4], Jonsdottir

I.S.[5], Pham N.M.[3], Pugnaire F.I.[6], Shea K.[7], Wardle D.A.[8], Zielosko S.[1], Bråthen K.A.[1][*][†]

Affiliations

[1] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Tromsø, Norway

[2] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Alta, Norway

[3] The Norwegian College of Fishery Science, UiT – Arctic University of Norway, Tromsø, Norway

[4] Department of Ecosystems in the Barents region, Norwegian institute of Bioeconomy Research-NIBIO, Svanvik,

Norway

[5] Institute of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland

[6] Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas (EEZA-CSIC), Almería,

Spain

[7] Department of Biology and IGDP in Ecology, The Pennsylvania State University, University Park, Pennsylvania,

16802 USA

[8] Asian School of the Environment, Nanyang Technological University, Singapore

* corresponding authors

maria.w.tuomi@uit.no, kari.brathen@uit.no

†These authors contributed equally to this work

*Abstract*

Management blind spots, i.e., influential processes not included in management models, may

undermine management objectives and reduce system resilience. Here we address one such

blind spot in the management of *Rangifer* in Norway. The current model is based on

regulation of pastures and population vital rates through reindeer density, but lacks validation

of the underlying assumption of forage stability, which is a critical blind spot. We report

pasture changes over 18 growing seasons, under variable reindeer densities and a changing

climate. We show a strong increase of the native evergreen, allelopathic crowberry,

contrasted by smaller increases of deciduous dwarf-shrubs and stagnation in herbaceous

species. We found no evidence of reindeer density influencing these vegetation changes. Our

results unveil a trajectory of Arctic evergreening and increasing allelopathy, and a shift

towards lower biodiversity and ecosystem functioning. It exemplifies an ongoing change in a

management blind spot that greatly impacts social-ecological systems.

Sustainable management of ecosystems and social-ecological systems (SES) is contingent on

models that are representative of the managed system and its uncertainties (Milner-Gulland

and Shea 2017; Regan, Colyvan, and Burgman 2002; Latombe et al. 2019). Management blind

spots, i.e. *influential variables or processes not included in a management model*, may

severely undermine management objectives (Madden and McQuinn 2014), lead to misguided

management efforts, and reduce the capacity of the system to adapt and maintain system

resilience against undesirable states (Dudney et al. 2018). Particularly challenging are slow

and poorly detectable changes in external or internal drivers of the system (Schneider, Leifeld,

and Malang 2013) that, if overlooked through being a management blind spot, may develop

unnoticed and potentially push the system towards a less favorable state. Hence, sustainable

SES-management under climate change is contingent on assessment and revision of

management models to identify potential critical blind spots.

In the Arctic, climate change brings novel threats (Post et al. 2009; Mallory and Boyce

2017), and opportunities (Mallory and Boyce 2017; Henden et al. 2020) for caribou and

reindeer (*Rangifer* sp.) populations and associated indigenous pastoral systems (Horstkotte

et al. 2017). The anticipated warming-induced vegetation trends in summer pastures include

increased productivity, biomass and leaf area, which contribute to Arctic greening (Myers-

Smith et al. 2020), and shifts towards more resource acquisitive species (Bjorkman et al. 2018;

Elina Kaarlejärvi, Eskelinen, and Olofsson 2013). Such changes are expected to manifest

through biomass increases of plants with taller growth and higher nitrogen (N)

concentrations, such as willows, graminoids and forbs (Bråthen, Gonzalez, and Yoccoz 2018;

Bjorkman et al. 2018; Myers-Smith et al. 2019). Increased productivity of these palatable

species should translate into higher population growth rates of ungulates through positive

bottom-up effects (Henden et al. 2020; Tveraa et al. 2013). By foraging, ungulates can even

counteract shrubification (Christie et al. 2015), keeping willows in a "browse-trap" (Bråthen

et al. 2017), thereby preventing the most nutritious forbs and graminoids from being

overgrown (Bret-Harte et al. 2001; Mod and Luoto 2016).

However, although rapid changes in climate in the Arctic (IPCC 2021) are altering

primary productivity, biodiversity and ecosystem functions (Myers-Smith et al. 2020; Barry,

Berteaux, and Bültmann 2013; Post et al. 2019; Taylor et al. 2020), recent observations

indicate that ongoing Arctic greening may conceal qualitatively and functionally contrasting

trends. The satellite-derived greening signal in North-America is negatively related to caribou

population growth rates, and this has been attributed to expansion of deciduous shrubs with

high levels of anti-browsing defenses (Fauchald et al. 2017). Contradicting satellite-derived

inference, field-derived observations show increments in especially evergreen dwarf-shrubs

across the circumpolar Arctic (Wilson and Nilsson 2009; Elmendorf et al. 2012; Maliniemi et

al. 2018; Vowles et al. 2017). These plants often have high phenolics and low N content, and

can contribute to lowering ecosystem productivity (Vowles and Björk 2019; Tuomi et al.

2018). These observations suggest a decline of pasture quality, an "evergreening" trend that

is functionally distinct from that of greening. In Arctic pastoral SES, already under pressure

from a multitude of anthropogenic pressures (Horstkotte et al. 2017; Hausner et al. 2020),

ecosystem bottom-up effects that erode pasture quality could pose a significant management

risk, and undermine system resilience if left unchecked.

In this context, of particular interest is the niche-constructing, native, allelopathic

evergreen dwarf-shrub, black crowberry (*Empetrum nigrum*) (Bråthen, Gonzalez, and Yoccoz

2018; Wardle et al. 1998) (Table S1). In Fennoscandia, crowberry is already abundant and

commonly dominant, and as a boreal-Arctic species it appears to thrive under climate change

and tolerates various environmental stressors (Tybirk et al. 2000; Aerts 2010; Preece and

Phoenix 2014). Owing to its low foliar nutrient concentrations, high levels of allelochemical

polyphenolic compounds in its leaves and a dense, clonal growth, crowberry is highly

unpalatable, and can significantly retard ecosystem processes such as litter decomposition,

soil nutrient fluxes and seedling establishment (Bråthen, Fodstad, and Gallet 2010; González

et al. 2021; Tybirk et al. 2000; Wardle et al. 1998; Nilsson and Wardle 2005). It can also

suppress biodiversity and herbaceous plant growth in summer pastures (Bråthen, Gonzalez,

and Yoccoz 2018) (Table S1). While herbivores can suppress growth of palatable deciduous

shrubs (Bråthen et al. 2017; Olofsson et al. 2009), their effect on crowberry growth is less

clear (Maliniemi et al. 2018; Vowles et al. 2017), as reindeer avoid crowberry-dominated

areas for most of the growing season (Iversen et al. 2014). This indicates that ecosystem

evergreening may severely undermine a management system based on an assumption of

density-dependent population regulation derived from stable-state conditions (Krebs 2002).

In the present paper, our focus is on tundra pastures used for reindeer husbandry,

which are core to the indigenous Sámi livelihood and culture (James 2020). *Rangifer* (reindeer

**Figure 1.** Pastures as a blind spot of a reindeer management system. **The current management of reindeer in Norway applies reindeer density as the regulatory mechanism of sustainable pastures** (*Reindeer Husbandry Act*). The central norm for assessing sustainability of the summer pastures is the autumn slaughter weight of animals, as a proxy for body condition, and no measures pertaining to the pastures themselves. **a) Empirical density-dependence between reindeer densities and weights within and across herding districts**. Over the time-period of 2000-2019 and among the management districts of the current study, weights declined with increasing reindeer density both within districts representing different ecological contexts and across all districts, here exemplified with calf weights (*Ministry of Agriculture*). **b) Available forage quality and quantity in summer pastures in northern Fennoscandia in 2003.** Pasture forage consists of plant groups of varying species richness (Bråthen et al 2007), quality (nitrogen content (Murguzur et al 2019)) and abundance (biomass (Bråthen et al 2007)). Forbs, the most nutritious and species-rich group, make up the least biomass in pastures. Conversely, the least nutritious and evergreen dwarf-shrub crowberry makes up the most biomass. **c) Exposing pastures as a blind spot in the current reindeer management model.** The link between reindeer densities and weights is indirect, assumed to be mediated through pasture carrying capacity in an equilibrium system, i.e. forage (*Reindeer Husbandry Act; White Paper 32 (2016-2017)*). The decomposed model shows the assumed negative top-down relationship between animal density and forage in an equilibrium system (left panel), and an empirical positive bottom-up relationship between forage and weights (right panel, see Tveraa et al. 2013). However, negative density-dependence in animal populations can operate independent of pasture development (Krebs 2002). Framed graphs in panels b) and c) indicate patterns and relationships that are empirically addressed here.

and caribou) grazing systems span the circumpolar area (Mallory and Boyce 2017) and, akin

all extensive pastoral or grazing systems, rely on high-quality plant resources in pastures as

forage for the animals (Sinclair and Krebs 2002). Changes in pastures are therefore focal to

understanding mismanagement and adaptation risks in pastoral systems, as exemplified in

our study system: the Norwegian reindeer herding SES. The current Norwegian reindeer

management model (Fig 1a), set by the reindeer herding law  of 1978 and 2007 (*Reindeer*

*Husbandry Act*, lovdata.no/dokument/NL/lov/2007-06-15-40), operates based on an

empirically supported negative density-dependent relationship between reindeer numbers

and body mass (Henden et al. 2020; Stien et al. 2021), and an implicit assumption of a stable-
state population dynamics, suggesting that population vital rates can be predictably managed
via animal density (Krebs 2002) (Fig 1a, Supplementary Figure S1). Whenever animal weights
fall below a threshold, husbandry sustainability is, according to this model, attained via
lowering animal densities. A key implicit mediator of this relationship are summer pastures
and their quality, assumed to be in an equilibrium state (Krebs 2002), but this is a blind spot
in the management model (Fig 1c). In 2003, crowberry made up the majority of plant biomass
across the northern Norwegian tundra, resulting in pastures that are abundant in forage of
low nitrogen content (Fig 1b). Monitoring the change in crowberry abundance over time
relative to that of the more nutritious shrubs, forbs and graminoids, is vital to understanding
the development of these tundra ecosystems and their pastoral SES.

Specifically, we ask 1) to what extent crowberry and other forage plant groups are
changing their abundance in tundra pastures, 2) whether the densities of reindeer, which are
the primary regulating factor in reindeer management, are linked to these changes, and 3)
what are the implications of the observed changes in forage quantity for sustainability of the
reindeer management system. To answer these questions, we present results from a unique,
large-scale resurvey of 292 plant communities within 56 landscape areas sampled across 20
reindeer summer pastures, or management districts, in northern Norway over 18 growing
seasons from 2003 to 2020 (Figure S2). Finally, we use these empirical findings to support a
revision to the reindeer
management model.

Located at the transition
from subarctic to the low-Arctic,
Northern Fennoscandia is a
bellwether for changes in tundra
vegetation (Vuorinen et al. 2017;
Elmendorf et al. 2012) and
ecosystem functioning (Parker et al.
2021). The region has experienced
lengthening of the growing season as
spring and autumn have become
warmer over the past century

**Figure 2.** Growing degree days (GDD), as annual sums of daily temperatures above 5°C during May-October 1957-2019, averaged across the studied reindeer summer pastures (Pedersen et al. 2020/met.no). Solid thick line and shaded areas represent mean ± 95% confidence intervals from segmented linear regression with a breakpoint at year 1977. Parameter estimates are provided at the top of the figure.

(Kivinen et al. 2017). With an approximate increase of 100 growing degree days in the studied

summer pastures since the late 70's (Fig. 2), the pastures have been subject to a growing

season prolongation, which is a key driver behind increased shrub growth in the Arctic (Post

et al. 2019; Weijers et al. 2018; Myers-Smith et al. 2019). We used identical georeferenced

plant communities and sampling methods within each community in both 2003 and 2020.

Each plant community was represented by a 50m long transect with 11 equidistant, triangular

measurement plots with 40cm sides (Bråthen et al. 2007). We used Bayesian generalized

linear mixed models (Bürkner 2017) to test if 1) community-level standing biomass of forbs,

graminoids, deciduous woody plants (shrubs and dwarf-shrubs) and crowberry, and the area

covered by these groups, changed over the study period, and if 2) the standing biomass of

these forage groups was associated with reindeer density. For the latter, we linked the plant

biomass estimated for each of the survey-years with their averaged reindeer densities from

the previous two decades (*Ministry of Agriculture,*

https://www.landbruksdirektoratet.no/nb/nyhetsrom/rapporter/ressursregnskapet-for-

reindriftsnaeringen, accessed via reinbase.no). We report results as posterior mean effect

sizes (pmes) and their 95% credible intervals (CI), unless stated otherwise.

160                                                    Results

161                               *Vegetation changes in tundra pastures*

Over the 18 growing season interval of our surveys, crowberry standing biomass

increased by 60% (Figure 3, Table S2), and by 14% in cover (Supplementary Table S3). The

increase in crowberry is in sharp contrast with a complete stagnation in the standing biomass

and cover of the more nutrient rich and faster growing forbs and graminoids over this period

(Figure 3a, Table 1, S2 and S3). Crowberry's increase is nearly an order of magnitude larger

than the increase of deciduous dwarf-shrubs and shrubs (Figure 3a, Table 1, S3). The

frequency of communities for which more than 25% of total vascular biomass was crowberry

rose significantly over this period [pmes 0.57; 95%CI 0.10-1.05], from an already high 0.76 to

0.83, suggesting that an increasing number of communities is subject to allelopathic effects

from crowberry.

In contrast, we found little evidence of reindeer densities correlating with abundances

of any of the plant forage groups either in 2003 or 2020 across the different management

districts (Figure 3b), with small effect sizes and 95% credible intervals for all parameter

estimates overlapping zero (Table 1).

While crowberry biomass did not vary

with reindeer density, it varied

substantially among landscape areas

within management districts (Fig 3b,

Table S3), suggesting strong ecological

context-dependence in its abundance.

Deciduous shrubs and graminoids varied

m arkedly between districts and in

landscape areas, while variation in forb

biomass was largest between districts

(Fig 3b, Tables S3, S4).

**Table 1.** Parameter estimates for plant biomass change from 2003 to 2020 and plant biomass variation with animal density in 2003 and 2020. For full population-level estimates (fixed effects) and estimates of group-level standard deviation (random effects), see Supplementary Table S2. Error terms Q2.5 and Q97.5 represent the 95% credible interval, within which the true estimate lies with a 95% probability. Bold font indicates strong support for the effect, and italic indicates relatively strong support for the effect, given the data.

| response | parameter | estimate | Q2.5 | Q97.5 |
|---|---|---|---|---|
| Forbs | Density 2003 | -0.089 | -0.190 | 0.013 |
| | Biomass 2020 | -0.306 | -0.846 | 0.240 |
| | Density 2020 | 0.032 | -0.043 | 0.107 |
| Graminoids | Density 2003 | -0.0004 | -0.067 | 0.066 |
| | Biomass 2020 | -0.115 | -0.514 | 0.289 |
| | Density 2020 | 0.009 | -0.047 | 0.067 |
| Deciduous woody | Density 2003 | -0.015 | -0.067 | 0.037 |
| | *Biomass 2020* | *0.283* | *-0.001* | *0.575* |
| | Density 2020 | 0.004 | -0.039 | 0.045 |
| Crowberry | Density 2003 | 0.026 | -0.019 | 0.072 |
| | **Biomass 2020** | **0.601** | **0.301** | **0.905** |
| | Density 2020 | -0.019 | -0.062 | 0.025 |

**Figure 3.** Changes in the abundance of plant groups of pasture forage in the management blind spot. **a)** Modelled posterior predictions of biomass of forbs, graminoids, deciduous woody plants and crowberry, separately for each plant group in 2003 (grey) and in 2020 (greens). Means and 95% confidence intervals from observed data for each year are provided as point intervals below each biomass density plot. **b)** Modelled association with confidence bands (model-derived 95% credible intervals of the posterior mean estimate) between plant groups of pasture forage (forbs, graminoids, deciduous shrubs and dwarf-shrubs and crowberry), and reindeer densities across management districts, separately for 2003 (grey) and 2020 (greens). Observed associations within each district and year (points) are linked per district to display the trajectory of change in forage group biomass with changing reindeer densities over time.

Discussion

We show that tundra pastures over the last two decades have been subject to a strong

evergreening forcing, attributable to the increased abundance of the native, allelopathic

evergreen dwarf-shrub crowberry (Fig. 4a), and a weaker greening forcing. The process of

pasture evergreening (Maliniemi et al. 2018; Vuorinen et al. 2017; Vowles et al. 2017) far

outpaces that of the deciduous shrubs – species typically linked to the process of greening

(Post et al. 2019; Weijers et al. 2018; Myers-Smith et al. 2019). Moreover, we show this

process juxtaposed with no increase of the already scarce yet most species-rich and

productive plant groups, the forbs and graminoids, likely resulting in an important biodiversity

loss through reduced species evenness.

**Figure 4. a)** Mechanisms of crowberry encroachment and increased allelopathy over time. Growth of established plants results in infilling and increased biomass, leading to stronger allelopathic litter effects. Dispersal via seeds and clonal reproduction are means of lateral encroachment. Already an abundant and long-lived species, slow growth may still cause substantial increments in abundance. **b)** Conceptual model of the pasture state transition based on bottom-up effects. Increasing crowberry abundance may push the system towards higher allelopathy and induce legacy effects (green arrow) that over time will reduce new plant establishment (Gonzalez et al 2021) at levels as low as 25% crowberry out of total community standing biomass (Bråthen et al 2018). The average proportion of crowberry in community standing biomass in Fennoscandia was above 25% in 2003 (upper panel) and even higher in 2020. A new management norm of *pastures in a state of low allelopathy* (red arrow) could push the system back. **c)** Conceptual model of a revision to the management model based on results from the current study. Weak or no relationship between the abundance of pasture forage and reindeer densities decouples the regulatory mechanism and the norm in the current management model, suggesting primacy of bottom-up regulatory effects in reindeer summer pastures, prompting the inclusion of a new management focus: pasture forage.

200     Across management districts, representing a range of different ecological contexts, reindeer densities were not related to abundance patterns of any forage group (Fig 3b). Our results therefore point at evergreening by crowberry encroachment as a bottom-up forcing on the pastures (Fig 4) (cf. le Roux et al. 2013), an ever stronger process in the management blind spot of the reindeer herding SES with potentially adverse long-term consequences for the diversity and productivity of this critical indigenous Arctic pastoral system. Furthermore, our results suggest crowberry encroachment delimits areas available for pastures (Skarin et al. 2020; Iversen et al. 2014), adding to ongoing loss of land with productive pastures due to human disturbance (Hausner et al. 2020) and other stressors (Vaas et al. 2021; Horstkotte et al. 2017). Such loss may even amplify other stressors, further eroding resilience of the system.

211     *Evergreening: tundra pastures in transition*

212     As a strong niche-constructing species, crowberry can modify the environment once established (González et al. 2021; Bråthen, Gonzalez, and Yoccoz 2018; Wardle et al. 1998) and through legacy effects and reinforcing feed-backs, push the system towards a state of strong allelopathy (Supplementary Table S4, Fig 4a-b). Consequently, we suspect that the legacy effects at the community- and ecosystem-level from crowberry allelopathy may represent an ongoing shift with potential context-specific thresholds (Hughes et al. 2013) (Fig 4bc). Diminishing diversity may happen gradually through reduced seedling recruitment (González et al. 2015; 2021), and over time induce an extinction debt on the plant communities (Kuussaari et al. 2009, Gilbert & Levine 2013). As a consequence of such reinforcing feedbacks, along with the potential longevity, dense growth and poorly palatable leaves of crowberry (Aerts 2010), crowberry-dominance of communities is likely to be a highly resilient state (E.a Kaarlejärvi, Hoset, and Olofsson 2015; González et al. 2021; Wardle et al. 1998; Nilsson and Wardle 2005), and once established, may require strong external disturbances to reverse (Fig 4a-b).

226     While we show that crowberry encroachment occurs in landscapes across northern Fennoscandia, we also find large spatial variability in this encroachment, indicating there are landscape areas and ecological contexts in which the shift seems weaker or absent (Fig. 3b, Supplementary Table S3). Tundra vegetation changes are repeatedly found to be spatially heterogeneous across scales (Björkman et al 2020, van Wijk et al. 2004) and can be linked to e.g., variation in microclimate or herbivory. For instance, cyclic small rodent outbreaks across

the resurveyed region can decimate dwarf-shrubs and especially crowberry in patches across

landscapes (Tuomi et al. 2018; Hoset et al. 2017) (Fig. S3) - a temporary reduction even

detectable from space (Olofsson, Tømmervik, and Callaghan 2012). However, strong localized

rodent impact has not limited the overall high-magnitude encroachment of crowberry

documented here.

While neither small nor large herbivores may halt the trend of allelopathic

encroachment across tundra pastures, they are likely adversely affected by the observed,

extremely low abundance of forbs and diminishing prevalence of communities with little

crowberry (Oksanen et al. 2020; Iversen et al. 2014). Apart from reindeer, other endotherm

herbivores, such as small rodents, ptarmigans, domesticated sheep and musk-ox also rely on

N-rich forage (Schmidt et al. 2018), and seek pastures with high-quality food (Iversen et al.

2014; Jenkins et al. 2020; Skarin et al. 2020). Furthermore, crowberry being a wind-pollinated

plant (Bell and Tallis 1973), its encroachment over forbs may over time also affect insect

pollinators. In summary, crowberry encroachment of the magnitude documented here raises

significant concerns of cascading effects onto the tundra ecosystem and human beneficiaries

relying on them – including pastoralists, sheep farmers and game hunters.

*Encroaching allelopathy in a management blind spot*

A trajectory towards stronger allelopathy (Fig 4b) – resulting in lowered pasture process rates,

forage availability and carrying capacity – induces a risk of mismanagement and loss of

resilience of the SES (Valéry, Fritz, and Lefeuvre 2013; Carey et al. 2012; Shackleton,

Shackleton, and Kull 2019). Therefore our results call for explicit inclusion of norms relating

to pasture plant diversity and productivity, especially that of forbs (Bråthen, Pugnaire, and

Bardgett 2021), in the management model (Fig 4c), in line with intentions for sustainable

reindeer husbandry (lovdata.no) and the urgency of translating the IPBES Global Assessment

to policy (Ruckelshaus et al. 2020).

Presently, however, crowberry encroachment, and its ecological and socio-economic

impacts, remain largely unobserved and subject to high epistemic uncertainty (cf. Shackleton,

Shackleton, and Kull 2019; van der Wal et al. 2015) that can impede development of

management norms and objectives (Vaas et al. 2021; Davis et al. 2019). Validation and

monitoring of Arctic vegetation change has relied heavily on remote sensing indices (Myers-

Smith et al. 2020; Henden et al. 2020), which may not necessarily capture changes in

functional composition (Wang et al. 2021), or indeed, distinguish greening and evergreening as functionally different processes (Vowles and Björk 2019). Adaptive and long-term ecosystem and pasture monitoring programmes are key to mitigating uncertainty about climate-driven vegetation change (Post et al. 2009; Myers-Smith et al. 2019; Lindenmayer and Likens 2009). Such programmes have recently been implemented, for example, in Iceland (Marteinsdóttir et al. 2021), and in the Varanger peninsula and Svalbard, in Norway (Ims and Yoccoz 2017).

Management interventions or norms that promote diversity, productivity and spatial heterogeneity of pastoral landscapes have centennial or even millennial roots in extensive management practices of European coastal heathlands (Webb 2008; Vandvik et al. 2005) and in European agri-environmental policy focused on preventing loss of open semi-natural grasslands (Critchley, Burke, and Stevens 2004). Especially fire has the capacity to ameliorate soil conditions against crowberry's allelopathic effects (Bråthen, Fodstad, and Gallet 2010; Keech, Carcaillet, and Nilsson 2005), suggesting management through burning as a way to control ecosystem state shifts similar to that of coastal heathlands (Vandvik et al. 2005) or boreal *Pinus*-dominated forests (Wardle et al. 1998; Nilsson and Wardle 2005). Here, the challenge lies in the already scarce and diminishing herbaceous resource, the forbs (Bråthen, Pugnaire, and Bardgett 2021), for adaptive monitoring and as a basis for management, emphasising the need to see SES and their management in light of constantly changing biodiversity.

*System-wide consequences of native encroachment*

Encroachment puts an increasing pressure on sustainable land-use planning and prioritization, in particular to preserve remaining forb-rich pastures and enhance productive and diverse tundra landscapes. Evergreening through crowberry encroachment exemplifies emergence of super-dominance among native species, a phenomenon linked with anthropogenic pressures or novel climates across biomes (Pivello et al. 2018; Shackelford et al. 2013; Valéry et al. 2009). We show how the lack of monitoring or management norms for a native species with niche constructing potential can cause an unnoticed development of biotic, bottom-up forcing of a pastoral management system. This also demonstrates how a native species can cause ecosystem effects similar to non-native invasive species, and can

occur in the blind spot of any management system focusing on potential harm from invasive species or changed top-down forcing, only.

Ignoring the capacity of native species to be internal drivers of change is a critical blind spot that threatens the biodiversity and sustainability of ecosystems. We recommend that norms and objectives constantly informed by ecosystem-based monitoring will enhance sustainable management of diversity, and complex social-ecological systems. For the Arctic pastoral system studied here, the management of tundra ecosystems and SES post-2020 should align targets of reindeer and biodiversity management, and ensure resilience through explicit norms related to its resource base; the biodiversity of forage.

*Materials and Methods*

**Study area and resurvey design.** We conducted a vegetation survey of 292 remote, georeferenced vegetation communities in summer pastures across Northern Norway (lat N69° 25.806'- N70° 58.471', lon E20° 47.186'- E27° 31.099') during peak growing season in 2003 and again in 2020. The study area includes strong climatic gradients from west to east and from coast to inland, altitudinal variation from 60 to 600 m asl, as well as variation in bedrock types. Sampling included the most common vegetation types in the region: heaths, mires, snow beds, meadows, and windblown ridges. The survey design (Supplementary Figure S2) incorporates summer pasture areas of 20 reindeer districts, where adjacent districts of similar climatic conditions were organized into 10 district pairs of contrasting reindeer densities, based on average reindeer density from 1980 to 2003. Within each district, a 2x2 km vegetated grid was assigned out of which a random subset was chosen as landscape areas for sampling. Within each landscape area, a random set of 25 of the 100, 200 x 200m squares that constitute the vegetated grid were assigned for plant community sampling. Squares were then sampled by the transect method (each transect representing a plant community), whereby a 50m transect was placed from the midpoint towards a random GPS-position along a circle with a 50m radius. We used the point-frequency method in 11 plots every 5m along the transect to sample each community (Supplementary Figure S2). Each plot was measured by placing a triangular frame with sides of 40cm and one pin in each corner, counting all intercepts with the vegetation. The *a priori* stratification of the original 2003 design, inclusion rules of transects and sampling method are described in more detail in Bråthen et al. (2007; 2018). In 2020, we resurveyed a subset of 292 (of the original 1450) communities in 56 (of if the original 151) landscape areas. The re-sampling retained the

geographic extent as well as most of climatic and abiotic variability. *A priori*, we also set a higher

priority to resample communities where crowberry occurrence was lower in 2003, i.e., in transects

where crowberry did not occur along the entire transect, to allow for detection of changes in

crowberry extent. This was accomplished by limiting sampling to transects where crowberry

*occurrence* was in lower 70% quantiles of the entire 2003 dataset. We were only partially able to follow

this prioritization due to limited accessibility to sites, and hence also transects with higher than 70%

crowberry occurrence in 2003 were included in the sample. Based on transect descriptions from 2003

(e.g., "transect was moved 10m backwards due to a lake") we accurately relocated all resampled

communities that were measured in 2003.

**Environmental data.** We estimated climatic trends between 1957 and 2019 in the studied summer

pastures for growing degree days (GDD), using data described in Pedersen (2021). We applied

segmented linear regression models using the R-package segmented (Muggeo 2017) to explore trends

and breakpoints in the mean estimates for the regional climate.

We retrieved data on reindeer numbers from the onset of the reindeer herding year (cf. Stien et al.

2021) for each studied reindeer herding district (Ministry of Agriculture, accessed via reinbase.no),

which we then divided by summer pasture area ($km^2$) to obtain reindeer density for each district

(individuals/$km^2$). We used the average density from 1980 to 2003 as a predictor for the 2003

vegetation data, and average density from 2003 to 2019 as a predictor for the 2020 vegetation data.

**Vegetation sampling.** We used the point-intercept method (Bråthen and Hagberg 2004) with a

triangular 3-pin frame to obtain a measure of vascular plant abundance. We counted all hits of all

vascular species in 11 0.08$m^2$ plots spaced along each 50m transect. Prior to further analysis, we

converted point-frequency hits per species per plot to biomass estimates (g/$m^2$) using established

calibration equations (Tuomi et al. 2018; Ravolainen et al. 2010). For analyzing biomass data, we first

pooled species-specific biomass estimates based on functional grouping to forbs, graminoids,

deciduous woody shrubs and dwarf-shrubs (deciduous woody), and crowberry. Other groups not

included in the analysis (Supplementary table S2) were other evergreen woody dwarf-shrubs, non-

woody evergreen plants and vascular cryptogams. We then averaged biomass of all 11 plots along

each transect to reach a community-averaged estimate of g/$m^2$ for forbs (non-zero sample size $N_{year}$:

$N_{2003}$ = 136, $N_{2020}$ = 136), graminoids ($N_{2003}$ = 257, $N_{2020}$ = 252), deciduous woody ($N_{2003}$ = 282, $N_{2020}$ =

279), and crowberry ($N_{2003}$ = 261, $N_{2020}$ = 269). We also calculated and analyzed the cover of crowberry

within each community, using plot-level presence-absence data and summarized numbers of plots

with crowberry present in each transect (results in Supplementary table S4).

**Statistical analyses.** To test our first two hypothesis, we used Bayesian linear multilevel models fitted

with the package *brms* (Bürkner 2017) in the R statistical environment (version 4.0.4 / 15.02.2021).

We modeled biomass of each plant group and crowberry cover separately, with year (2003 vs. 2020)

and reindeer density and their interaction as population-level predictors. Landscape areas nested

within reindeer districts were set as the group-level predictors of the intercept. We fitted gamma-

hurdle models for biomass data and zero-inflated beta models for crowberry extent data. We used

package default weakly informative priors for population and group-level predictors and family

parameters (Bürkner 2017), and evaluated convergence and independence of HMC chains based on

the ^R statistics (< 1.1) and effective sample size (> 1000) (Muth, Oravecz, and Gabry 2018). We

confirmed model fit visually through posterior predictive checks as well as comparing model-

simulated data to observed data (Muth, Oravecz, and Gabry 2018). Data visualization was done with

packages ggplot2 (Wickham 2009) and ggdist (Kay 2021).

Data availability

The datasets collected and analyzed in this study as well as R scripts for analysis are available in UiT

repository at ---.

Acknowledgements

The work was funded by the Norwegian Research Council (FRIPRO project MONEC, code 302749 to

K.A.B.). We thank Karoline Helene Aares, Hanna Böhner, Lea Lipphardt, Hans Ivar Hortmann, Kinga

Skalska, Katrine Skamfer Hoset and Sindre Natvik for conducting field work, and the Norwegian Coast

Guard, especially the crew of KV Farm, for their hospitality and invaluable logistic help during the field

work campaign. We thank Audun Stien for discussions on the calf weight-density relationship. We also

wish to thank MONEC partners for discussions on implications of crowberry encroachment for their

reindeer and sheep herding SES. M.T. thanks the Mikkeli University Consortium's (MUC) for providing

working facilities.

Author contributions

K.A.B. conceived the idea, M.T., K.A.B. and N.Y. planned the re-sampling design. K.A.B. designed

conceptual figures, based on discussions with all authors. M.T., K.A.B., T.Aa.U., N.Y. and S.Z. collected

the data. M.T., K.A.B. and N.Y. planned data analysis. M.T. analyzed the data and extracted the results.

M.T. and K.A.B. led data interpretation, with all authors contributing to interpreting the results. M.T.

wrote the manuscript with frequent advice and support from K.A.B. All authors contributed

substantially to editing the manuscript.

Additional information

**Extended data: Extended discussion**

**Supplementary Material:**

**Supplementary Table S1.** Literature synthesis of key characteristics of crowberry (*Empetum nigrum*

ssp. *nigrum / hermaphroditum*).

**Supplementary Table S2.** Observed biomass and species richness of all functional groups.

**Supplementary Table S3.** Parameter estimates for plant biomass change from 2003 to 2020 and plant

biomass variation with animal density in 2003 and 2020.

**Supplementary Table S4.** Parameter estimates for plant extent change from 2003 to 2020 and plant

biomass variation with animal density in 2003 and 2020.

**Supplementary Figure S1.** Map of the study area and illustration of the sampling design.

**Supplementary Figure S2.** Small rodent outbreak impacts on dwarf-shrub vegetation during 2020 re-
sampling.

References

Aerts, Rien. 2010. "Nitrogen-Dependent Recovery of Subarctic Tundra Vegetation after
Simulation of Extreme Winter Warming Damage to Empetrum Hermaphroditum."
*Global Change Biology* 16 (3): 1071–81. https://doi.org/10.1111/j.1365-
2486.2009.01999.x.
Barry, Tom, Dominique Berteaux, and Helga Bültmann, eds. 2013. *Arctic Biodiversity*
*Assessment: Status and Trends in Arcitc Biodiversity*. Akureyri, Iceland: The
Conservation of Arctic Flora and Fauna.
Bell, J. N. B., and J. H. Tallis. 1973. "Empetrum Nigrum L." *The Journal of Ecology* 61 (1): 289.
https://doi.org/10.2307/2258934.
Bjorkman, Anne D., Isla H. Myers-Smith, Sarah C. Elmendorf, Signe Normand, Nadja Rüger,
Pieter S. A. Beck, Anne Blach-Overgaard, et al. 2018. "Plant Functional Trait Change
across a Warming Tundra Biome." *Nature* 562 (7725): 57–62.
https://doi.org/10.1038/s41586-018-0563-7.
Bråthen, Kari Anne, Cathrine Helene Fodstad, and Christiane Gallet. 2010. "Ecosystem
Disturbance Reduces the Allelopathic Effects of Empetrum Hermaphroditum Humus
on Tundra Plants." *Journal of Vegetation Science* 21 (4): 786–95.
https://doi.org/10.1111/j.1654-1103.2010.01188.x.
Bråthen, Kari Anne, Victoria T. Gonzalez, and Nigel G. Yoccoz. 2018. "Gatekeepers to the
Effects of Climate Warming? Niche Construction Restricts Plant Community Changes
along a Temperature Gradient." *Perspectives in Plant Ecology, Evolution and*
*Systematics*, Special issue on Alpine and arctic plant communities : a worldwide
perspective, 30 (February): 71–81. https://doi.org/10.1016/j.ppees.2017.06.005.
Bråthen, Kari Anne, and Oskar Hagberg. 2004. "More Efficient Estimation of Plant Biomass."
*Journal of Vegetation Science* 15 (5): 653–660.
Bråthen, Kari Anne, Rolf A. Ims, Nigel G. Yoccoz, Per Fauchald, Torkild Tveraa, and Vera H.
Hausner. 2007. "Induced Shift in Ecosystem Productivity? Extensive Scale Effects of
Abundant Large Herbivores." *Ecosystems* 10 (5): 773–89.
https://doi.org/10.1007/s10021-007-9058-3.
Bråthen, Kari Anne, Francisco I Pugnaire, and Richard D Bardgett. 2021. "The Paradox of
Forbs in Grasslands and the Legacy of the Mammoth Steppe." *Frontiers in Ecology*
*and the Environment* 19 (10): 584–92. https://doi.org/10.1002/fee.2405.
Bråthen, Kari Anne, Virve Tuulia Ravolainen, Audun Stien, Torkild Tveraa, and Rolf A. Ims.
2017. "Rangifer Management Controls a Climate-Sensitive Tundra State Transition."
*Ecological Applications* 27 (8): 2416–27. https://doi.org/10.1002/eap.1618.
Bret-Harte, M Syndonia, GAIus R Shaver, Jennifer P Zoerner, Jill F Johnstone, Joanna L
Wagner, Andreas S Chavez, Ralph F Gunkelman Iv, Suzanne C Lippert, and James A
Laundre'. 2001. "Developmental Plasticity Allows Betula Nana to Dominate Tundra
Subjected to an Altered Environment," 16.
Bürkner, Paul-Christian. 2017. "Brms: An R Package for Bayesian Multilevel Models Using
Stan." *Journal of Statistical Software* 80 (1). https://doi.org/10.18637/jss.v080.i01.
Carey, Michael P., Beth L. Sanderson, Katie A. Barnas, and Julian D. Olden. 2012. "Native
Invaders – Challenges for Science, Management, Policy, and Society." *Frontiers in*
*Ecology and the Environment* 10 (7): 373–81. https://doi.org/10.1890/110060.
Christie, Katie S., John P. Bryant, Laura Gough, Virve T. Ravolainen, Roger W. Ruess, and Ken
D. Tape. 2015. "The Role of Vertebrate Herbivores in Regulating Shrub Expansion in

the Arctic: A Synthesis." *BioScience* 65 (12): 1123–33.
https://doi.org/10.1093/biosci/biv137.

Critchley, C. N. R, M. J. W Burke, and D. P Stevens. 2004. "Conservation of Lowland Semi-
Natural Grasslands in the UK: A Review of Botanical Monitoring Results from Agri-
Environment Schemes." *Biological Conservation* 115 (2): 263–78.
https://doi.org/10.1016/S0006-3207(03)00146-0.

Davis, Katrina J., Iadine Chadès, Jonathan R. Rhodes, and Michael Bode. 2019. "General
Rules for Environmental Management to Prioritise Social Ecological Systems
Research Based on a Value of Information Approach." *Journal of Applied Ecology* 56
(8): 2079–90. https://doi.org/10.1111/1365-2664.13425.

Dudney, Joan, Richard J. Hobbs, Robert Heilmayr, John J. Battles, and Katharine N. Suding.
2018. "Navigating Novelty and Risk in Resilience Management." *Trends in Ecology &*
*Evolution* 33 (11): 863–73. https://doi.org/10.1016/j.tree.2018.08.012.

Elmendorf, Sarah C., Gregory H. R. Henry, Robert D. Hollister, Robert G. Björk, Noémie
Boulanger-Lapointe, Elisabeth J. Cooper, Johannes H. C. Cornelissen, et al. 2012.
"Plot-Scale Evidence of Tundra Vegetation Change and Links to Recent Summer
Warming." *Nature Climate Change* 2 (6): 453–57.
https://doi.org/10.1038/nclimate1465.

Fauchald, Per, Taejin Park, Hans Tømmervik, Ranga Myneni, and Vera Helene Hausner.
2017. "Arctic Greening from Warming Promotes Declines in Caribou Populations."
*Science Advances* 3 (4): e1601365. https://doi.org/10.1126/sciadv.1601365.

González, Victoria T., Olavi Junttila, Bente Lindgård, Rigmor Reiersen, Kajetan Trost, and Kari
Anne Bråthen. 2015. "Batatasin-III and the Allelopathic Capacity of Empetrum
Nigrum." *Nordic Journal of Botany* 33 (2): 225–31.
https://doi.org/10.1111/njb.00559.

González, Victoria T., Bente Lindgård, Rigmor Reiersen, Snorre B. Hagen, and Kari Anne
Bråthen. 2021. "Niche Construction Mediates Climate Effects on Recovery of Tundra
Heathlands after Extreme Event." Edited by Craig Eliot Coleman. *PLOS ONE* 16 (2):
e0245929. https://doi.org/10.1371/journal.pone.0245929.

Hausner, Vera H., Sigrid Engen, Camilla Brattland, and Per Fauchald. 2020. "Sámi Knowledge
and Ecosystem-Based Adaptation Strategies for Managing Pastures under Threat
from Multiple Land Uses." *Journal of Applied Ecology* 57 (9): 1656–65.
https://doi.org/10.1111/1365-2664.13559.

Henden, Ja, T Tveraa, A Stien, Jp Mellard, F Marolla, Ra Ims, and Ng Yoccoz. 2020. "Direct
and Indirect Effects of Environmental Drivers on Reindeer Reproduction." *Climate*
*Research*. https://doi.org/10.3354/cr01630.

Horstkotte, T., T. Aa Utsi, Å Larsson-Blind, P. Burgess, B. Johansen, J. Käyhkö, L. Oksanen,
and B. C. Forbes. 2017. "Human–Animal Agency in Reindeer Management: Sámi
Herders' Perspectives on Vegetation Dynamics under Climate Change." *Ecosphere* 8
(9): e01931. https://doi.org/10.1002/ecs2.1931.

Hoset, Katrine S., Lise Ruffino, Maria Tuomi, Tarja Oksanen, Lauri Oksanen, Aurelia
Mäkynen, Bernt Johansen, and Torunn Moe. 2017. "Changes in the Spatial
Configuration and Strength of Trophic Control Across a Productivity Gradient During
a Massive Rodent Outbreak." *Ecosystems*, March, 1–15.
https://doi.org/10.1007/s10021-017-0124-1.

Hughes, Terry P., Cristina Linares, Vasilis Dakos, Ingrid A. van de Leemput, and Egbert H. van
Nes. 2013. "Living Dangerously on Borrowed Time during Slow, Unrecognized

Regime Shifts." *Trends in Ecology & Evolution* 28 (3): 149–55.
https://doi.org/10.1016/j.tree.2012.08.022.

Ims, Rolf A, and Nigel G Yoccoz. 2017. "Ecosystem-Based Monitoring in the Age of Rapid
Climate Change and New Technologies." *Current Opinion in Environmental*
*Sustainability* 29 (December): 170–76. https://doi.org/10.1016/j.cosust.2018.01.003.

IPCC, ed. 2021. *IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of*
*Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on*
*Climate Change*. Cambridge University Press.

Iversen, Marianne, Per Fauchald, Knut Langeland, Rolf A. Ims, Nigel G. Yoccoz, and Kari Anne
Bråthen. 2014. "Phenology and Cover of Plant Growth Forms Predict Herbivore
Habitat Selection in a High Latitude Ecosystem." Edited by John F. Valentine. *PLoS*
*ONE* 9 (6): e100780. https://doi.org/10.1371/journal.pone.0100780.

James, Simon P. 2020. "Legal Rights and Nature's Contributions to People: Is There a
Connection?" *Biological Conservation* 241 (January): 108325.
https://doi.org/10.1016/j.biocon.2019.108325.

Jenkins, Deborah A., Nicolas Lecomte, Geoffrey Andrews, Glenn Yannic, and James A.
Schaefer. 2020. "Biotic Interactions Govern the Distribution of Coexisting Ungulates
in the Arctic Archipelago – A Case for Conservation Planning." *Global Ecology and*
*Conservation* 24 (December): e01239. https://doi.org/10.1016/j.gecco.2020.e01239.

Kaarlejärvi, E.a, K.S.b Hoset, and J.a Olofsson. 2015. "Mammalian Herbivores Confer
Resilience of Arctic Shrub-Dominated Ecosystems to Changing Climate." *Global*
*Change Biology* 21 (9): 3379–88. https://doi.org/10.1111/gcb.12970.

Kaarlejärvi, Elina, Anu Eskelinen, and Johan Olofsson. 2013. "Herbivory Prevents Positive
Responses of Lowland Plants to Warmer and More Fertile Conditions at High
Altitudes." *Functional Ecology* 27 (5): 1244–53. https://doi.org/10.1111/1365-
2435.12113.

Kay, Matthew. 2021. *Ggdist: Visualizations of Distributions and Uncertainty*.
https://doi.org/10.5281/zenodo.3879620.

Keech, Olivier, Christopher Carcaillet, and Marie-Charlotte Nilsson. 2005. "Adsorption of
Allelopathic Compounds by Wood-Derived Charcoal: The Role of Wood Porosity."
*Plant and Soil* 272 (1): 291–300. https://doi.org/10.1007/s11104-004-5485-5.

Kivinen, Sonja, Sirpa Rasmus, Kirsti Jylhä, and Mikko Laapas. 2017. "Long-Term Climate
Trends and Extreme Events in Northern Fennoscandia (1914–2013)." *Climate* 5 (1):
16. https://doi.org/10.3390/cli5010016.

Krebs, Charles J. 2002. "Two Complementary Paradigms for Analysing Population
Dynamics." *Philosophical Transactions of the Royal Society of London. Series B:*
*Biological Sciences* 357 (1425): 1211–19. https://doi.org/10.1098/rstb.2002.1122.

Latombe, G., S. Canavan, H. Hirsch, C. Hui, S. Kumschick, M. M. Nsikani, L. J. Potgieter, et al.
2019. "A Four-Component Classification of Uncertainties in Biological Invasions:
Implications for Management." *Ecosphere* 10 (4): e02669.
https://doi.org/10.1002/ecs2.2669.

Lindenmayer, David B., and Gene E. Likens. 2009. "Adaptive Monitoring: A New Paradigm
for Long-Term Research and Monitoring." *Trends in Ecology & Evolution* 24 (9): 482–
86. https://doi.org/10.1016/j.tree.2009.03.005.

Madden, Francine, and Brian McQuinn. 2014. "Conservation's Blind Spot: The Case for
Conflict Transformation in Wildlife Conservation." *Biological Conservation* 178
(October): 97–106. https://doi.org/10.1016/j.biocon.2014.07.015.

Maliniemi, Tuija, Jutta Kapfer, Patrick Saccone, Anu Skog, and Risto Virtanen. 2018. "Long-Term Vegetation Changes of Treeless Heath Communities in Northern Fennoscandia: Links to Climate Change Trends and Reindeer Grazing." *Journal of Vegetation Science* 29 (3): 469–79. https://doi.org/10.1111/jvs.12630.

Mallory, Conor D., and Mark S. Boyce. 2017. "Observed and Predicted Effects of Climate Change on Arctic Caribou and Reindeer." *Environmental Reviews*, June. https://doi.org/10.1139/er-2017-0032.

Marteinsdóttir, B, E F Þórarinsdóttir, G Halldórsson, J H Stefánsson, J Þórsson, K Svavarsdóttir, R Finnsdóttir, and S Jónsdóttir. 2021. "GróLind – Sustainable Land Use Based on Ecological Knowledge." *International Grassland Congress Proceedings*, 6.

Milner-Gulland, E. J., and Katriona Shea. 2017. "Embracing Uncertainty in Applied Ecology." *Journal of Applied Ecology* 54 (6): 2063–68. https://doi.org/10.1111/1365-2664.12887.

Mod, Heidi K., and Miska Luoto. 2016. "Arctic Shrubification Mediates the Impacts of Warming Climate on Changes to Tundra Vegetation." *Environmental Research Letters* 11 (12): 124028. https://doi.org/10.1088/1748-9326/11/12/124028.

Muggeo, Vito M.R. 2017. "Interval Estimation for the Breakpoint in Segmented Regression: A Smoothed Score-Based Approach." *Australian & New Zealand Journal of Statistics* 59: 311–22.

Muth, Chelsea, Zita Oravecz, and Jonah Gabry. 2018. "User-Friendly Bayesian Regression Modeling: A Tutorial with Rstanarm and Shinystan." *The Quantitative Methods for Psychology* 14 (2): 99–119. https://doi.org/10.20982/tqmp.14.2.p099.

Myers-Smith, Isla H., Meagan M. Grabowski, Haydn J. D. Thomas, Sandra Angers-Blondin, Gergana N. Daskalova, Anne D. Bjorkman, Andrew M. Cunliffe, et al. 2019. "Eighteen Years of Ecological Monitoring Reveals Multiple Lines of Evidence for Tundra Vegetation Change." *Ecological Monographs* 89 (2): e01351. https://doi.org/10.1002/ecm.1351.

Myers-Smith, Isla H., Jeffrey T. Kerby, Gareth K. Phoenix, Jarle W. Bjerke, Howard E. Epstein, Jakob J. Assmann, Christian John, et al. 2020. "Complexity Revealed in the Greening of the Arctic." *Nature Climate Change* 10 (2): 106–17. https://doi.org/10.1038/s41558-019-0688-1.

Nilsson, Marie-Charlotte, and David A. Wardle. 2005. "Understory Vegetation as a Forest Ecosystem Driver: Evidence from the Northern Swedish Boreal Forest." *Frontiers in Ecology and the Environment* 3 (8): 421–28. https://doi.org/10.1890/1540-9295(2005)003[0421:UVAAFE]2.0.CO;2.

Oksanen, Tarja, Lauri Oksanen, Katariina E. M. Vuorinen, Christopher Wolf, Aurelia Mäkynen, Johan Olofsson, William J. Ripple, Risto Virtanen, and Tove Aa. Utsi. 2020. "The Impact of Thermal Seasonality on Terrestrial Endotherm Food Web Dynamics: A Revision of the Exploitation Ecosystem Hypothesis." *Ecography* 43 (12): 1859–77. https://doi.org/10.1111/ecog.05076.

Olofsson, Johan, Lauri Oksanen, Terry Callaghan, Philip E. Hulme, Tarja Oksanen, and Otso Suominen. 2009. "Herbivores Inhibit Climate-Driven Shrub Expansion on the Tundra." *Global Change Biology* 15 (11): 2681–93. https://doi.org/10.1111/j.1365-2486.2009.01935.x.

Olofsson, Johan, Hans Tømmervik, and Terry V. Callaghan. 2012. "Vole and Lemming Activity Observed from Space." *Nature Climate Change* 2 (12): 880–83. https://doi.org/10.1038/nclimate1537.

Parker, Thomas C., Alana M. Thurston, Katrine Raundrup, Jens-Arne Subke, Philip A. Wookey, and Iain P. Hartley. 2021. "Shrub Expansion in the Arctic May Induce Large-scale Carbon Losses Due to Changes in Plant-soil Interactions." *Plant and Soil* 463 (1): 643–51. https://doi.org/10.1007/s11104-021-04919-8.

Pedersen, Åshild Ønvik. 2021. *Norwegian Arctic Tundra: A Panel-Based Assessment of Ecosystem Condition*. Norsk Polarinstitutt. https://brage.npolar.no/npolar-xmlui/handle/11250/2754696.

Pivello, Vânia Regina, Marcus Vinicius Vieira, Maria Tereza Grombone-Guaratini, and Dalva Maria Silva Matos. 2018. "Thinking about Super-Dominant Populations of Native Species – Examples from Brazil." *Perspectives in Ecology and Conservation* 16 (2): 74–82. https://doi.org/10.1016/j.pecon.2018.04.001.

Post, Eric, Richard B. Alley, Torben R. Christensen, Marc Macias-Fauria, Bruce C. Forbes, Michael N. Gooseff, Amy Iler, et al. 2019. "The Polar Regions in a 2°C Warmer World." *Science Advances* 5 (12): eaaw9883. https://doi.org/10.1126/sciadv.aaw9883.

Post, Eric, Mads C. Forchhammer, M. Syndonia Bret-Harte, Terry V. Callaghan, Torben R. Christensen, Bo Elberling, Anthony D. Fox, et al. 2009. "Ecological Dynamics Across the Arctic Associated with Recent Climate Change." *Science* 325 (5946): 1355–58. https://doi.org/10.1126/science.1173113.

Preece, C., and G. K. Phoenix. 2014. "Impact of Early and Late Winter Icing Events on Sub-Arctic Dwarf Shrubs." *Plant Biology* 16 (1): 125–32. https://doi.org/10.1111/plb.12015.

Ravolainen, V. T., N. G. Yoccoz, K. A. Bråthen, R. A. Ims, M. Iversen, and V. T. González. 2010. "Additive Partitioning of Diversity Reveals No Scale-Dependent Impacts of Large Ungulates on the Structure of Tundra Plant Communities." *Ecosystems* 13 (1): 157–70. https://doi.org/10.1007/s10021-009-9308-7.

Regan, Helen M., Mark Colyvan, and Mark A. Burgman. 2002. "A Taxonomy and Treatment of Uncertainty for Ecology and Conservation Biology." *Ecological Applications* 12 (2): 618–28. https://doi.org/10.1890/1051-0761(2002)012[0618:ATATOU]2.0.CO;2.

Roux, Peter C. le, Jonathan Lenoir, Loïc Pellissier, Mary S. Wisz, and Miska Luoto. 2013. "Horizontal, but Not Vertical, Biotic Interactions Affect Fine-Scale Plant Distribution Patterns in a Low-Energy System." *Ecology* 94 (3): 671–682.

Ruckelshaus, Mary H., Stephen T. Jackson, Harold A. Mooney, Katharine L. Jacobs, Karim-Aly S. Kassam, Mary T. K. Arroyo, András Báldi, et al. 2020. "The IPBES Global Assessment: Pathways to Action." *Trends in Ecology & Evolution* 35 (5): 407–14. https://doi.org/10.1016/j.tree.2020.01.009.

Schmidt, Niels M., Jesper B. Mosbacher, Eero J. Vesterinen, Tomas Roslin, and Anders Michelsen. 2018. "Limited Dietary Overlap amongst Resident Arctic Herbivores in Winter: Complementary Insights from Complementary Methods." *Oecologia* 187 (3): 689–99. https://doi.org/10.1007/s00442-018-4147-x.

Schneider, Volker, Philip Leifeld, and Thomas Malang. 2013. "Coping with Creeping Catastrophes: National Political Systems and the Challenge of Slow-Moving Policy Problems." In *Long-Term Governance for Social-Ecological Change*, edited by Bernd Siebenhüner, Marlen Arnold, Klaus Eisenack, Klaus Jacob, and Michael Pregernig, 221–38. Environmental Politics / Routledge Research in Environmental Politics, v. 21. New York: Routledge.

Shackelford, Nancy, Michael Renton, Michael P. Perring, and Richard J. Hobbs. 2013. "Modeling Disturbance-Based Native Invasive Species Control and Its Implications for Management." *Ecological Applications* 23 (6): 1331–44. https://doi.org/10.1890/12-1881.1.

Shackleton, Ross T., Charlie M. Shackleton, and Christian A. Kull. 2019. "The Role of Invasive Alien Species in Shaping Local Livelihoods and Human Well-Being: A Review." *Journal of Environmental Management*, The human and social dimensions of invasion science and management, 229 (January): 145–57. https://doi.org/10.1016/j.jenvman.2018.05.007.

Sinclair, A. R. E., and Charles J. Krebs. 2002. "Complex Numerical Responses to Top–down and Bottom–up Processes in Vertebrate Populations." Edited by R. M. Sibly, J. Hone, and T. H. Clutton–Brock. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 357 (1425): 1221–31. https://doi.org/10.1098/rstb.2002.1123.

Skarin, Anna, Mariana Verdonen, Timo Kumpula, Marc Macias-Fauria, Moudud Alam, Jeffrey Kerby, and Bruce C Forbes. 2020. "Reindeer Use of Low Arctic Tundra Correlates with Landscape Structure." *Environmental Research Letters* 15 (11): 115012. https://doi.org/10.1088/1748-9326/abbf15.

Stien, Audun, Torkild Tveraa, Rolf Anker Ims, Jennifer Stien, and Nigel Gilles Yoccoz. 2021. "Unfounded Claims about Productivity beyond Density for Reindeer Pastoralism Systems." *Pastoralism* 11 (1): 20. https://doi.org/10.1186/s13570-021-00209-1.

Taylor, Jason J., James P. Lawler, Mora Aronsson, Tom Barry, Anne D. Bjorkman, Tom Christensen, Stephen J. Coulson, et al. 2020. "Arctic Terrestrial Biodiversity Status and Trends: A Synopsis of Science Supporting the CBMP State of Arctic Terrestrial Biodiversity Report." *Ambio* 49 (3): 833–47. https://doi.org/10.1007/s13280-019-01303-w.

Tuomi, Maria, Sari Stark, Katrine S. Hoset, Maria Väisänen, Lauri Oksanen, Francisco J. A. Murguzur, Hanna Tuomisto, Jonas Dahlgren, and Kari Anne Bråthen. 2018. "Herbivore Effects on Ecosystem Process Rates in a Low-Productive System." *Ecosystems* 22 (4): 827–43. https://doi.org/10.1007/s10021-018-0307-4.

Tveraa, Torkild, Audun Stien, Bård-J. Bårdsen, and Per Fauchald. 2013. "Population Densities, Vegetation Green-Up, and Plant Productivity: Impacts on Reproductive Success and Juvenile Body Mass in Reindeer." Edited by Gil Bohrer. *PLoS ONE* 8 (2): e56450. https://doi.org/10.1371/journal.pone.0056450.

Tybirk, Knud, Marie-Charlotte Nilsson, Anders Michelsen, Hanne Lakkenborg Kristensen, Anna Shevtsova, Morten Tune Strandberg, Marianne Johansson, et al. 2000. "Nordic Empetrum Dominated Ecosystems: Function and Susceptibility to Environmental Changes." *AMBIO: A Journal of the Human Environment* 29 (2): 90–97. https://doi.org/10.1579/0044-7447-29.2.90.

Vaas, J., P. P. J. Driessen, M. Giezen, F. van Laerhoven, and M. J. Wassen. 2021. "Moving from Latent to Manifest Problem: Trajectories Across Scientific and Public Salience of Invasive Alien Species." *Environmental Management* 67 (5): 901–19. https://doi.org/10.1007/s00267-020-01404-3.

Valéry, Loïc, Hervé Fritz, and Jean-Claude Lefeuvre. 2013. "Another Call for the End of Invasion Biology." *Oikos* 122 (8): 1143–46. https://doi.org/10.1111/j.1600-0706.2013.00445.x.

Valéry, Loïc, Hervé Fritz, Jean-Claude Lefeuvre, and Daniel Simberloff. 2009. "Invasive
Species Can Also Be Native…." *Trends in Ecology & Evolution* 24 (11): 585.
https://doi.org/10.1016/j.tree.2009.07.003.
Vandvik, Vigdis, Einar Heegaard, Inger Elisabeth Måren, and Per Arild Aarrestad. 2005.
"Managing Heterogeneity: The Importance of Grazing and Environmental Variation
on Post-Fire Succession in Heathlands." *Journal of Applied Ecology* 42 (1): 139–49.
https://doi.org/10.1111/j.1365-2664.2005.00982.x.
Vowles, Tage, and Robert G. Björk. 2019. "Implications of Evergreen Shrub Expansion in the
Arctic." *Journal of Ecology* 107 (2): 650–55. https://doi.org/10.1111/1365-
2745.13081.
Vowles, Tage, Bengt Gunnarsson, Ulf Molau, Thomas Hickler, Leif Klemedtsson, and Robert
G. Björk. 2017. "Expansion of Deciduous Tall Shrubs but Not Evergreen Dwarf Shrubs
Inhibited by Reindeer in Scandes Mountain Range." *Journal of Ecology* 105 (6): 1547–
61. https://doi.org/10.1111/1365-2745.12753.
Vuorinen, Katariina E. M., Lauri Oksanen, Tarja Oksanen, Anni Pyykönen, Johan Olofsson,
and Risto Virtanen. 2017. "Open Tundra Persist, but Arctic Features Decline—
Vegetation Changes in the Warming Fennoscandian Tundra." *Global Change Biology*
23 (9): 3794–3807. https://doi.org/10.1111/gcb.13710.
Wal, René van der, Anke Fischer, Sebastian Selge, and Brendon M. H. Larson. 2015. "Neither
the Public nor Experts Judge Species Primarily on Their Origins." *Environmental
Conservation* 42 (4): 349–55. https://doi.org/10.1017/S0376892915000053.
Wang, Hao, Huiying Liu, Ni Huang, Jian Bi, Xuanlong Ma, Zhiyuan Ma, Zijian Shangguan, et al.
2021. "Satellite-Derived NDVI Underestimates the Advancement of Alpine
Vegetation Growth over the Past Three Decades." *Ecology* 102 (12): e03518.
https://doi.org/10.1002/ecy.3518.
Wardle, David A., Marie-Charlotte Nilsson, Christiane Gallet, and Olle Zackrisson. 1998. "An
Ecosystem-Level Perspective of Allelopathy." *Biological Reviews* 73 (3): 305–319.
Webb, N.R. 2008. "The Traditional Management of European Heathlands." *Journal of
Applied Ecology* 35 (6): 987–90. https://doi.org/10.1111/j.1365-
2664.1998.tb00020.x.
Weijers, Stef, Roland Pape, Jörg Löffler, and Isla H. Myers-Smith. 2018. "Contrasting Shrub
Species Respond to Early Summer Temperatures Leading to Correspondence of
Shrub Growth Patterns." *Environmental Research Letters* 13 (3): 034005.
https://doi.org/10.1088/1748-9326/aaa5b8.
Wickham, Hadley. 2009. *Ggplot2: Elegant Graphics for Data Analysis*. Springer Science &
Business Media.
Wilson, Scott D., and Christer Nilsson. 2009. "Arctic Alpine Vegetation Change over 20
Years." *Global Change Biology* 15 (7): 1676–84. https://doi.org/10.1111/j.1365-
2486.2009.01896.x.
729
730