Arctic evergreening - encroaching allelopathy in a management blind spot

Tuomi M.W.[1*†], Utsi T.Aa.[2], Yoccoz N.[1], Armstrong C.W.[3], Gonzalez V.[4], Hagen S.B.[4], Jónsdóttir

I.S.[5], Pugnaire F.I.[6], Shea K.[7], Wardle D.A.[8], Zielosko S.[1], Bråthen K.A.[1*†]

Affiliations

[1] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Tromsø, Norway

[2] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Alta, Norway

[3] The Norwegian College of Fishery Science, UiT – Arctic University of Norway, Tromsø, Norway

[4] Department of Ecosystems in the Barents region, Norwegian institute of Bioeconomy Research-NIBIO, Svanvik,

Norway

[5] Institute of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland

[6] Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas (EEZA-CSIC), Almería,

Spain

[7] Department of Biology and IGDP in Ecology, The Pennsylvania State University, University Park, Pennsylvania,

16802 USA

[8] Asian School of the Environment, Nanyang Technological University, Singapore

* corresponding authors

maria.w.tuomi@uit.no, kari.brathen@uit.no

†These authors contributed equally to this work

24   *Abstract*

Management blind spots, i.e., influential processes not included in management models, may

undermine management objectives and reduce system resilience. Here we address one such

blind spot in the management of *Rangifer* in Norway. The current model is based on

regulation of pastures and population vital rates through reindeer density, but lacks validation

of the underlying assumption of forage stability, which is a critical blind spot. We report

pasture changes over 18 growing seasons, under variable reindeer densities and a changing

climate. We show a strong increase of the native evergreen, allelopathic crowberry,

contrasted by smaller increases of deciduous dwarf-shrubs and stagnation in herbaceous

species. We found no evidence of reindeer density influencing these vegetation changes. Our

results unveil a trajectory of Arctic evergreening and increasing allelopathy, and a shift

towards lower biodiversity and ecosystem functioning. It exemplifies an ongoing change in a

management blind spot that greatly impacts social-ecological systems.

                                        Introduction

Sustainable management of ecosystems and social-ecological systems (SES) is contingent on

models that are representative of the managed system and its uncertainties[1–3]. Management

blind spots, i.e. *influential variables or processes not included in a management model*, may

severely undermine management objectives[4], lead to misguided management efforts, and

reduce the capacity of the system to adapt and maintain system resilience against undesirable

states[5]. Particularly challenging are slow and poorly detectable changes in external or internal

drivers of the system[6] that, if overlooked through being a management blind spot, may

develop unnoticed and potentially push the system towards a less favorable state. Hence,

sustainable SES-management under climate change is contingent on assessment and revision

of management models to identify potential critical blind spots.

In the Arctic, climate change brings novel threats[7,8], and opportunities[8,9] for caribou

and reindeer (*Rangifer* sp.) populations and associated indigenous pastoral systems[10]. The

anticipated warming-induced vegetation trends in summer pastures include increased

productivity, biomass and leaf area, which contribute to Arctic greening[11], and shifts towards

more resource acquisitive species[12,13]. Such changes are expected to manifest through

biomass increases of plants with taller growth and higher nitrogen (N) concentrations, such

as willows, graminoids and forbs[12,14,15]. Increased productivity of these palatable species

should translate into higher population growth rates of ungulates through positive bottom-

up effects[9,16]. By foraging, ungulates can even counteract shrubification[17], keeping willows in

a "browse-trap"[18], thereby preventing the most nutritious forbs and graminoids from being

overgrown[19,20].

However, although rapid changes in climate in the Arctic[21] are altering primary

productivity, biodiversity and ecosystem functions[11,22–24], recent observations indicate that

ongoing Arctic greening may conceal qualitatively and functionally contrasting trends. The

satellite-derived greening signal in North-America is negatively related to caribou population

growth rates, and this has been attributed to expansion of deciduous shrubs with high levels

of anti-browsing defenses[25]. Contradicting satellite-derived inference, field-derived

observations show increments in especially evergreen dwarf-shrubs across the circumpolar

Arctic[26–29]. These plants often have high phenolics and low N content, and can contribute to

lowering ecosystem productivity[30,31]. These observations suggest a decline of pasture quality,

an "evergreening" trend that is functionally distinct from that of greening. In Arctic pastoral

SES, already under pressure from a multitude of anthropogenic pressures[10,32], ecosystem

bottom-up effects that erode pasture quality could pose a significant management risk, and

undermine system resilience if left unchecked.

In this context, of particular interest is the niche-constructing, native, allelopathic

evergreen dwarf-shrub, black crowberry (*Empetrum nigrum*)[14,33] (Table S1). In Fennoscandia,

crowberry is already abundant and commonly dominant, and as a boreal-Arctic species it

appears to thrive under climate change and tolerates various environmental stressors[34–36].

Owing to its low foliar nutrient concentrations, high levels of allelochemical polyphenolic

compounds in its leaves and a dense, clonal growth, crowberry is highly unpalatable, and can

significantly retard ecosystem processes such as litter decomposition, soil nutrient fluxes and

seedling establishment[33,34,37–39]. It can also suppress biodiversity and herbaceous plant

growth in summer pastures[14] (Table S1). While herbivores can suppress growth of palatable

deciduous shrubs[18,40], their effect on crowberry growth is less clear[28,29], as reindeer avoid

crowberry-dominated areas for most of the growing season[41]. This indicates that ecosystem

evergreening may severely undermine a management system based on an assumption of

density-dependent population regulation derived from stable-state conditions[42].

In the present paper, our focus is on tundra pastures used for reindeer husbandry,

which are core to the indigenous Sámi livelihood and culture[43]. *Rangifer* (reindeer and

caribou) grazing systems span the circumpolar area[8] and, akin all extensive pastoral or grazing

systems, rely on high-quality plant resources in pastures as forage for the animals[44]. Changes

in pastures are therefore focal to understanding mismanagement and adaptation risks in

pastoral systems, as exemplified in our study system: the Norwegian reindeer herding SES.

The current Norwegian reindeer management model (Fig 1a), set by the reindeer herding law

of 1978 and 2007 (*Reindeer Husbandry Act*, lovdata.no/dokument/NL/lov/2007-06-15-40),

operates based on an empirically supported negative density-dependent relationship

between reindeer numbers and body mass[9,45], and an implicit assumption of a stable-state

population dynamics, suggesting that population vital rates can be predictably managed via

animal density[42] (Fig 1a, Fig S1). Whenever animal weights fall below a threshold, husbandry

sustainability is, according to this model, attained via lowering animal densities. A key implicit

mediator of this relationship are summer pastures and their quality, assumed to be in an

equilibrium state[42], but this is a blind spot in the management model (Fig 1c). In 2003,

crowberry made up the majority of plant biomass across the northern Norwegian tundra,

resulting in pastures that are abundant in forage of low nitrogen content (Fig 1b). Monitoring

the change in crowberry abundance over time relative to that of the more nutritious shrubs,

forbs and graminoids, is vital to understanding the development of these tundra ecosystems

and their pastoral SES.

Figure 1. Pastures as a blind spot of a reindeer management system. **The current management of reindeer in Norway applies reindeer density as the regulatory mechanism of sustainable pastures** (*Reindeer Husbandry Act*). The central norm for assessing sustainability of the summer pastures is the autumn slaughter weight of animals, as a proxy for body condition, and no measures pertaining to the pastures themselves. **a) Empirical density-dependence between reindeer densities and weights within and across herding districts.** Over the time-period of 2000-2019 and among the management districts of the current study, weights declined with increasing reindeer density both within districts representing different ecological contexts and across all districts, here exemplified with calf weights (*Ministry of Agriculture*). **b) Available forage quality and quantity in summer pastures in northern Fennoscandia in 2003.** Pasture forage consists of plant groups of varying species richness (Bråthen et al 2007), quality (nitrogen content (Murguzur et al 2019)) and abundance (biomass (Bråthen et al 2007)). Forbs, the most nutritious and species-rich group, make up the least biomass in pastures. Conversely, the least nutritious and evergreen dwarf-shrub crowberry makes up the most biomass. **c) Exposing pastures as a blind spot in the current reindeer management model.** The link between reindeer densities and weights is indirect, assumed to be mediated through pasture carrying capacity in an equilibrium system, i.e. forage (*Reindeer Husbandry Act*; *White Paper 32 (2016-2017)*). The decomposed model shows the assumed negative top-down relationship between animal density and forage in an equilibrium system (left panel), and an empirical positive bottom-up relationship between forage and weights (right panel, see Tveraa et al. 2013). However, negative density-dependence in animal populations can operate independent of pasture development (Krebs 2002). Framed graphs in panels b) and c) indicate patterns and relationships that are empirically addressed here.

Specifically, we ask 1) to what extent crowberry and other forage plant groups are

changing their abundance in tundra pastures, 2) whether the densities of reindeer, which are

the primary regulating factor in reindeer management, are linked to these changes, and 3)

what are the implications of the observed changes in forage quantity for sustainability of the

reindeer management system. To answer these questions, we present results from a unique,

large-scale resurvey of 292 plant communities within 56 landscape areas sampled across 20

reindeer summer pastures, or management districts, in northern Norway over 18 growing

seasons from 2003 to 2020 (Fig S2). Finally, we use these empirical findings to support a

revision to the reindeer management model.

Located at the transition from subarctic to the low-Arctic, Northern Fennoscandia is a

bellwether for changes in tundra vegetation[27,46] and ecosystem functioning[47]. The region has

experienced lengthening of the growing season as spring and autumn have become warmer

over the past century[48]. With an approximate increase of 100 growing degree days in the

studied summer pastures since the late 70's (Fig 2), the pastures have been subject to a

growing season prolongation, which is a key driver behind increased shrub growth in the

Arctic[15,23,49]. We used identical georeferenced plant communities and sampling methods

within each community in both 2003 and 2020. Each plant community was represented by a

50m long transect with 11 equidistant, triangular measurement plots with 40cm sides[50]. We

used Bayesian generalized linear mixed models[51] to test if 1) community-level standing

biomass of forbs, graminoids,

deciduous woody plants (shrubs and

dwarf-shrubs) and crowberry, and

the area covered by these groups,

changed over the study period, and if

2) the standing biomass of these

forage groups was associated with

reindeer density. For the latter, we

linked the plant biomass estimated

for each of the survey-years with

their averaged reindeer densities

from the previous two decades

(*Ministry of Agriculture,*

Figure 2. Growing degree days (GDD), as annual sums of daily temperatures above 5°C during May-October 1957-2019, averaged across the studied reindeer summer pastures (Pedersen et al. 2020/met.no). Solid thick line and shaded areas represent mean ± 95% confidence intervals from segmented linear regression with a breakpoint at year 1977. Parameter estimates are provided at the top of the figure.

142   https://www.landbruksdirektoratet.no/nb/nyhetsrom/rapporter/ressursregnskapet-for-

143   reindriftsnaeringen, accessed via reinbase.no). We report results as posterior mean effect

144   sizes (pmes) and their 95% credible intervals (CI), unless stated otherwise.

146   Results

147   *Vegetation changes in tundra pastures*

Over the 18 growing season interval of our surveys, crowberry standing biomass

increased by 60% (Fig 3, Table S2), and by 14% in cover (Table S3). The increase in crowberry

is in sharp contrast with a complete stagnation in the standing biomass and cover of the more

nutrient rich and faster growing forbs and graminoids over this period (Fig 3a, Table 1, Tables

S2 and S3). Crowberry's increase is

nearly an order of magnitude larger

than the increase of deciduous dwarf-

shrubs and shrubs (Fig 3a, Table 1, Table

S3). The frequency of communities for

which more than 25% of total vascular

biomass was crowberry rose significantly

over this period [pmes 0.57; 95%CI 0.10-

1.05], from an already high 0.76 to 0.83,

suggesting that an increasing number of

communities is subject to allelopathic

effects from crowberry.

Table 1. Parameter estimates for plant biomass change from 2003 to 2020 and plant biomass variation with animal density in 2003 and 2020. For full population-level estimates (fixed effects) and estimates of group-level standard deviation (random effects), see Supplementary Table S2. Error terms Q2.5 and Q97.5 represent the 95% credible interval, within which the true estimate lies with a 95% probability. Bold font indicates strong support for the effect, and italic indicates relatively strong support for the effect, given the data.

response	parameter	estimate	Q2.5	Q97.5
Forbs	Density 2003	-0.089	-0.190	0.013
	Biomass 2020	-0.306	-0.846	0.240
	Density 2020	0.032	-0.043	0.107
Graminoids	Density 2003	-0.0004	-0.067	0.066
	Biomass 2020	-0.115	-0.514	0.289
	Density 2020	0.009	-0.047	0.067
Deciduous woody	Density 2003	-0.015	-0.067	0.037
	Biomass 2020	*0.283*	*-0.001*	*0.575*
	Density 2020	0.004	-0.039	0.045
Crowberry	Density 2003	0.026	-0.019	0.072
	Biomass 2020	**0.601**	**0.301**	**0.905**
	Density 2020	-0.019	-0.062	0.025

In contrast, we found little

evidence of reindeer densities correlating with abundances of any of the plant forage groups

either in 2003 or 2020 across the different management districts (Fig 3b), with small effect

sizes and 95% credible intervals for all parameter estimates overlapping zero (Table 1). While

crowberry biomass did not vary with reindeer density, it varied substantially among landscape

areas within management districts (Fig 3b, Table S3), suggesting strong ecological context-

dependence in its abundance. Deciduous shrubs and graminoids varied markedly between

districts and in landscape areas, while variation in forb biomass was largest between districts

(Fig 3b, Tables S3, S4).

Figure 3. Changes in the abundance of plant groups of pasture forage in the management blind spot. **a)** Modelled posterior predictions of biomass of forbs, graminoids, deciduous woody plants and crowberry, separately for each plant group in 2003 (grey) and in 2020 (greens). Means and 95% confidence intervals from observed data for each year are provided as point intervals below each biomass density plot. **b)** Modelled association with confidence bands (model-derived 95% credible intervals of the posterior mean estimate) between plant groups of pasture forage (forbs, graminoids, deciduous shrubs and dwarf-shrubs and crowberry), and reindeer densities across management districts, separately for 2003 (grey) and 2020 (greens). Observed associations within each district and year (points) are linked per district to display the trajectory of change in forage group biomass with changing reindeer densities over time.

175   Discussion

We show that tundra pastures over the last two decades have been subject to a strong

evergreening forcing, attributable to the increased abundance of the native, allelopathic

evergreen dwarf-shrub crowberry (Fig 4a), and a weaker greening forcing. The process of

pasture evergreening[28,29,46] far outpaces that of the deciduous shrubs – species typically

linked to the process of greening[15,23,49]. Moreover, we show this process juxtaposed with no

increase of the already scarce yet most species-rich and productive plant groups, the forbs

and graminoids, likely resulting in an important biodiversity loss through reduced species

evenness.

Across management districts, representing a range of different ecological contexts,

reindeer densities were not related to abundance patterns of any forage group (Fig 3b). Our

results therefore point at evergreening by crowberry encroachment as a bottom-up forcing

on the pastures (Fig 4)[cf. 52], an ever stronger process in the management blind spot of the

reindeer herding SES with potentially adverse long-term consequences for the diversity and

productivity of this critical indigenous Arctic pastoral system. Furthermore, our results

suggest crowberry encroachment delimits areas available for pastures[41,53], adding to ongoing

loss of land with productive pastures due to human disturbance[32] and other stressors[10,54].

Such loss may even amplify other stressors, further eroding resilience of the system.

Figure 4. a) Mechanisms of crowberry encroachment and increased allelopathy over time. Growth of established plants results in infilling and increased biomass, leading to stronger allelopathic litter effects. Dispersal via seeds and clonal reproduction are means of lateral encroachment. Already an abundant and long-lived species, slow growth may still cause substantial increments in abundance. **b)** Conceptual model of the pasture state transition based on bottom-up effects. Increasing crowberry abundance may push the system towards higher allelopathy and induce legacy effects (green arrow) that over time will reduce new plant establishment (Gonzalez et al 2021) at levels as low as 25% crowberry out of total community standing biomass (Bråthen et al 2018). The average proportion of crowberry in community standing biomass in Fennoscandia was above 25% in 2003 (upper panel) and even higher in 2020. A new management norm of *pastures in a state of low allelopathy* (red arrow) could push the system back. **c)** Conceptual model of a revision to the management model based on results from the current study. Weak or no relationship between the abundance of pasture forage and reindeer densities decouples the regulatory mechanism and the norm in the current management model, suggesting primacy of bottom-up regulatory effects in reindeer summer pastures, prompting the inclusion of a new management focus: pasture forage.

194   *Evergreening: tundra pastures in transition*

As a strong niche-constructing species, crowberry can modify the environment once

established[14,33,38] and through legacy effects and reinforcing feed-backs, push the system

towards a state of strong allelopathy (Fig 4a-b, Table S1). Consequently, we suspect that the

legacy effects at the community- and ecosystem-level from crowberry allelopathy may

represent an ongoing shift with potential context-specific thresholds [55] (Fig 4b-c). Diminishing

diversity may happen gradually through reduced seedling recruitment [38,56], and over time

induce an extinction debt on the plant communities[57,58]. As a consequence of such reinforcing

feedbacks, along with the potential longevity, dense growth and poorly palatable leaves of

crowberry[35], crowberry-dominance of communities is likely to be a highly resilient
state[33,38,39,59], and once established, may require strong external disturbances to reverse (Fig
4a-b).

While we show that crowberry encroachment occurs in landscapes across northern
Fennoscandia, we also find large spatial variability in this encroachment, indicating there are
landscape areas and ecological contexts in which the shift seems weaker or absent (Fig 3b,
Table S3). Tundra vegetation changes are repeatedly found to be spatially heterogeneous
across scales[12,60,61] and can be linked to e.g., variation in microclimate or herbivory. For
instance, cyclic small rodent outbreaks across the resurveyed region can decimate dwarf-
shrubs and especially crowberry in patches across landscapes[31,62] (Fig S3) - a temporary
reduction even detectable from space[63]. However, strong localized rodent impact has not
limited the overall high-magnitude encroachment of crowberry documented here.

While neither small nor large herbivores may halt the trend of allelopathic
encroachment across tundra pastures, they are likely adversely affected by the observed,
extremely low abundance of forbs and diminishing prevalence of communities with little
crowberry[41,64]. Apart from reindeer, other endotherm herbivores, such as small rodents,
ptarmigans, domesticated sheep and musk-ox also rely on N-rich forage[65], and seek pastures
with high-quality food[41,53,66]. Furthermore, crowberry being a wind-pollinated plant[67], its
encroachment over forbs may over time also affect insect pollinators. In summary, crowberry
encroachment of the magnitude documented here raises significant concerns of cascading
effects onto the tundra ecosystem and human beneficiaries relying on them – including
pastoralists, sheep farmers and game hunters.

Encroaching allelopathy in a management blind spot
A trajectory towards stronger allelopathy (Fig 4b) – resulting in lowered pasture process rates,
forage availability and carrying capacity – induces a risk of mismanagement and loss of
resilience of the SES[68–70]. Therefore our results call for explicit inclusion of norms relating to
pasture plant diversity and productivity, especially that of forbs[71], in the management model
(Fig 4c), in line with intentions for sustainable reindeer husbandry (lovdata.no) and the
urgency of translating the IPBES Global Assessment to policy[72].

Presently, however, crowberry encroachment, and its ecological and socio-economic
impacts, remain largely unobserved and subject to high epistemic uncertainty[cf. 70,73] that can

impede development of management norms and objectives[54,74]. Validation and monitoring

of Arctic vegetation change has relied heavily on remote sensing indices[9,11], which may not

necessarily capture changes in functional composition[75], or indeed, distinguish greening and

evergreening as functionally different processes[30]. Adaptive and long-term ecosystem and

pasture monitoring programmes are key to mitigating uncertainty about climate-driven

vegetation change[7,15,76]. Such programmes have recently been implemented, for example, in

Iceland[77], and in the Varanger peninsula and Svalbard, in Norway[78].

Management interventions or norms that promote diversity, productivity and spatial

heterogeneity of pastoral landscapes have centennial or even millennial roots in extensive

management practices of European coastal heathlands[79,80] and in European agri-

environmental policy focused on preventing loss of open semi-natural grasslands[81]. Especially

fire has the capacity to ameliorate soil conditions against crowberry's allelopathic effects[37,82],

suggesting management through burning as a way to control ecosystem state shifts similar to

that of coastal heathlands[80] or boreal *Pinus*-dominated forests[33,39]. Here, the challenge lies in

the already scarce and diminishing herbaceous resource, the forbs[71], for adaptive monitoring

and as a basis for management, emphasising the need to see SES and their management in

light of constantly changing biodiversity.

253   *System-wide consequences of native encroachment*

Encroachment puts an increasing pressure on sustainable land-use planning and

prioritization, in particular to preserve remaining forb-rich pastures and enhance productive

and diverse tundra landscapes. Evergreening through crowberry encroachment exemplifies

emergence of super-dominance among native species, a phenomenon linked with

anthropogenic pressures or novel climates across biomes[83–85]. We show how the lack of

monitoring or management norms for a native species with niche constructing potential can

cause an unnoticed development of biotic, bottom-up forcing of a pastoral management

system. This also demonstrates how a native species can cause ecosystem effects similar to

non-native invasive species, and can occur in the blind spot of any management system

focusing on potential harm from invasive species or changed top-down forcing, only.

Ignoring the capacity of native species to be internal drivers of change is a critical blind

spot that threatens the biodiversity and sustainability of ecosystems. We recommend that

norms and objectives constantly informed by ecosystem-based monitoring will enhance

sustainable management of diversity, and complex social-ecological systems. For the Arctic

pastoral system studied here, the management of tundra ecosystems and SES post-2020

should align targets of reindeer and biodiversity management, and ensure resilience through

explicit norms related to its resource base; the biodiversity of forage.

Materials and Methods

Study area and resurvey design. We conducted a vegetation survey of 292 remote, georeferenced

vegetation communities in summer pastures across Northern Norway (lat N69° 25.806'- N70° 58.471',

lon E20° 47.186'- E27° 31.099') during peak growing season in 2003 and again in 2020. The study area

includes strong climatic gradients from west to east and from coast to inland, altitudinal variation from

60 to 600 m asl, as well as variation in bedrock types. Sampling included the most common vegetation

types in the region: heaths, mires, snow beds, meadows, and windblown ridges. The survey design

(Figure S2) incorporates summer pasture areas of 20 reindeer districts, where adjacent districts of

similar climatic conditions were organized into 10 district pairs of contrasting reindeer densities, based

on average reindeer density from 1980 to 2003. Within each district, a 2x2 km vegetated grid was

assigned out of which a random subset was chosen as landscape areas for sampling. Within each

landscape area, a random set of 25 of the 100, 200 x 200m squares that constitute the vegetated grid

were assigned for plant community sampling. Squares were then sampled by the transect method

(each transect representing a plant community), whereby a 50m transect was placed from the

midpoint towards a random GPS-position along a circle with a 50m radius. We used the point-

frequency method in 11 plots every 5m along the transect to sample each community (Figure S2).

Each plot was measured by placing a triangular frame with sides of 40cm and one pin in each corner,

counting all intercepts with the vegetation. The *a priori* stratification of the original 2003 design,

inclusion rules of transects and sampling method are described in more detail in Bråthen et al. [14,50]. In

2020, we resurveyed a subset of 292 (of the original 1450) communities in 56 (of if the original 151)

landscape areas. The re-sampling retained the geographic extent as well as most of climatic and abiotic

variability. *A priori*, we also set a higher priority to resample communities where crowberry occurrence

was lower in 2003, i.e., in transects where crowberry did not occur along the entire transect, to allow

for detection of changes in crowberry extent. This was accomplished by limiting sampling to transects

where crowberry *occurrence* was in lower 70% quantiles of the entire 2003 dataset. We were only

partially able to follow this prioritization due to limited accessibility to sites, and hence also transects

with higher than 70% crowberry occurrence in 2003 were included in the sample. Based on transect

descriptions from 2003 (e.g., "transect was moved 10m backwards due to a lake") we accurately

relocated all resampled communities that were measured in 2003.

Environmental data. We estimated climatic trends between 1957 and 2019 in the studied summer

pastures for growing degree days (GDD), using data described in Pedersen [86]. We applied segmented

linear regression models using the R-package segmented [87] to explore trends and breakpoints in the

mean estimates for the regional climate.

We retrieved data on reindeer numbers from the onset of the reindeer herding year [cf. 45] for each

studied reindeer herding district (Ministry of Agriculture, accessed via reinbase.no), which we then

divided by summer pasture area (km^2) to obtain reindeer density for each district (individuals/km^2).

We used the average density from 1980 to 2003 as a predictor for the 2003 vegetation data, and

average density from 2003 to 2019 as a predictor for the 2020 vegetation data.

Vegetation sampling. We used the point-intercept method [88] with a triangular 3-pin frame to obtain

a measure of vascular plant abundance. We counted all hits of all vascular species in 11 $0.08m^2$ plots

spaced along each 50m transect. Prior to further analysis, we converted point-frequency hits per

species per plot to biomass estimates (g/m^2) using established calibration equations [31,89]. For analyzing

biomass data, we first pooled species-specific biomass estimates based on functional grouping to

forbs, graminoids, deciduous woody shrubs and dwarf-shrubs (deciduous woody), and crowberry.

Other groups not included in the analysis (Table S2) were other evergreen woody dwarf-shrubs, non-

woody evergreen plants and vascular cryptogams. We then averaged biomass of all 11 plots along

each transect to reach a community-averaged estimate of g/m^2 for forbs (non-zero sample size N_{year}:

N_{2003} = 136, N_{2020} = 136), graminoids (N_{2003} = 257, N_{2020} = 252), deciduous woody (N_{2003} = 282, N_{2020} =

279), and crowberry (N_{2003} = 261, N_{2020} = 269). We also calculated and analyzed the cover of crowberry

within each community, using plot-level presence-absence data and summarized numbers of plots

with crowberry present in each transect (results in Table S4).

Statistical analyses. To test our first two hypothesis, we used Bayesian linear multilevel models fitted

with the package *brms*[51] in the R statistical environment[90] (version 4.0.4/15.02.2021 and later). We

modeled biomass of each plant group and crowberry cover separately, with year (2003 vs. 2020) and

reindeer density and their interaction as population-level predictors. Landscape areas nested within

reindeer districts were set as the group-level predictors of the intercept. We fitted gamma-hurdle

models for biomass data and zero-inflated beta models for crowberry extent data. We used package

default weakly informative priors for population and group-level predictors and family parameters[51],

and evaluated convergence and independence of HMC chains based on the ^R statistics (< 1.1) and

effective sample size (> 1000)[91]. We confirmed model fit visually through posterior predictive checks

as well as comparing model-simulated data to observed data[91]. Data visualization was done with packages ggplot2[92] and ggdist[93].

Data availability

The datasets collected and analyzed in this study as well as reproducible R scripts for analysis are made publicly available (CC-BY license) via UiT repository (https://dataverse.no/) at DOI: ---.

Acknowledgements

The work was funded by the Norwegian Research Council (FRIPRO project MONEC, code 302749 to K.A.B.). We thank Nhat Minh Pham for comments on previous drafts of the manuscript, and Karoline Helene Aares, Hanna Böhner, Lea Lipphardt, Hans Ivar Hortmann, Kinga Skalska, Katrine Skamfer Hoset and Sindre Natvik for conducting field work, and the Norwegian Coast Guard, especially the crew of KV Farm, for their hospitality and invaluable logistic help during the field work campaign. We thank Audun Stien for discussions on the calf weight-density relationship. We also wish to thank MONEC partners for discussions on implications of crowberry encroachment for their reindeer and sheep herding SES. M.T. thanks the Mikkeli University Consortium (MUC) for providing working facilities.

Author contributions

K.A.B. conceived the idea, M.T., K.A.B. and N.Y. planned the re-sampling design. K.A.B. designed conceptual figures, based on discussions with all authors. M.T., K.A.B., T.Aa.U., N.Y. and S.Z. collected the data. M.T., K.A.B. and N.Y. planned data analysis. M.T. analyzed the data and extracted the results. M.T. and K.A.B. led data interpretation, with all authors contributing to interpreting the results. M.T. wrote the manuscript with frequent advice and support from K.A.B. All authors contributed substantially to editing the manuscript.

Competing interests

The authors declare no competing interests.

Materials & Correspondence

Maria W. Tuomi (maria.w.tuomi@uit.no)

Kari Anne Bråthen (kari.brathen@uit.no)

Additional information

Supplementary Material:

Supplementary Table S1. Literature synthesis of key characteristics of crowberry (*Empetum nigrum*

ssp. *nigrum / hermaphroditum*).

Supplementary Table S2. Observed biomass and species richness of all functional groups.

Supplementary Table S3. Parameter estimates for plant biomass change from 2003 to 2020 and plant

biomass variation with animal density in 2003 and 2020.

Supplementary Table S4. Parameter estimates for plant extent change from 2003 to 2020 and plant

biomass variation with animal density in 2003 and 2020.

Supplementary Figure S1. Map of the study area and illustration of the sampling design.

Supplementary Figure S2. Small rodent outbreak impacts on dwarf-shrub vegetation during 2020 re-

sampling.

Extended discussion.

References

1. Milner-Gulland, E. J. & Shea, K. Embracing uncertainty in applied ecology. *J. Appl. Ecol.* **54**, 2063–2068 (2017).

2. Regan, H. M., Colyvan, M. & Burgman, M. A. A Taxonomy and Treatment of Uncertainty for Ecology and Conservation Biology. *Ecol. Appl.* **12**, 618–628 (2002).

3. Latombe, G. *et al.* A four-component classification of uncertainties in biological invasions: implications for management. *Ecosphere* **10**, e02669 (2019).

4. Madden, F. & McQuinn, B. Conservation's blind spot: The case for conflict transformation in wildlife conservation. *Biol. Conserv.* **178**, 97–106 (2014).

5. Dudney, J., Hobbs, R. J., Heilmayr, R., Battles, J. J. & Suding, K. N. Navigating Novelty and Risk in Resilience Management. *Trends Ecol. Evol.* **33**, 863–873 (2018).

6. Schneider, V., Leifeld, P. & Malang, T. Coping with Creeping Catastrophes: National Political Systems and the Challenge of Slow-Moving Policy Problems. in *Long-term governance for social-ecological change* (eds. Siebenhüner, B., Arnold, M., Eisenack, K., Jacob, K. & Pregernig, M.) 221–238 (Routledge, 2013).

7. Post, E. *et al.* Ecological Dynamics Across the Arctic Associated with Recent Climate Change. *Science* **325**, 1355–1358 (2009).

8. Mallory, C. D. & Boyce, M. S. Observed and predicted effects of climate change on Arctic caribou and reindeer. *Environ. Rev.* (2017) doi:10.1139/er-2017-0032.

9. Henden, J. *et al.* Direct and indirect effects of environmental drivers on reindeer reproduction. *Clim. Res.* (2020) doi:10.3354/cr01630.

10. Horstkotte, T. *et al.* Human–animal agency in reindeer management: Sámi herders' perspectives on vegetation dynamics under climate change. *Ecosphere* **8**, e01931 (2017).

11. Myers-Smith, I. H. *et al.* Complexity revealed in the greening of the Arctic. *Nat. Clim. Change* **10**, 106–117 (2020).

12. Bjorkman, A. D. *et al.* Plant functional trait change across a warming tundra biome. *Nature* **562**, 57–62 (2018).

13. Kaarlejärvi, E., Eskelinen, A. & Olofsson, J. Herbivory prevents positive responses of lowland plants to warmer and more fertile conditions at high altitudes. *Funct. Ecol.* **27**, 1244–1253 (2013).

14. Bråthen, K. A., Gonzalez, V. T. & Yoccoz, N. G. Gatekeepers to the effects of climate warming? Niche construction restricts plant community changes along a temperature gradient. *Perspect. Plant Ecol. Evol. Syst.* **30**, 71–81 (2018).

15. Myers-Smith, I. H. *et al.* Eighteen years of ecological monitoring reveals multiple lines of evidence for tundra vegetation change. *Ecol. Monogr.* **89**, e01351 (2019).

16. Tveraa, T., Stien, A., Bårdsen, B.-J. & Fauchald, P. Population Densities, Vegetation Green-Up, and Plant Productivity: Impacts on Reproductive Success and Juvenile Body Mass in Reindeer. *PLoS ONE* **8**, e56450 (2013).

17. Christie, K. S. *et al.* The Role of Vertebrate Herbivores in Regulating Shrub Expansion in the Arctic: A Synthesis. *BioScience* **65**, 1123–1133 (2015).

18. Bråthen, K. A., Ravolainen, V. T., Stien, A., Tveraa, T. & Ims, R. A. Rangifer management controls a climate-sensitive tundra state transition. *Ecol. Appl.* **27**, 2416–2427 (2017).

19. Bret-Harte, M. S. *et al.* Developmental Plasticity Allows Betula nana to Dominate Tundra Subjected to an Altered Environment. 16 (2001).

20. Mod, H. K. & Luoto, M. Arctic shrubification mediates the impacts of warming climate on changes to tundra vegetation. *Environ. Res. Lett.* **11**, 124028 (2016).

21. *IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. (Cambridge University Press, 2021).

22. *Arctic Biodiversity Assessment: status and trends in Arcitc biodiversity*. (The Conservation of
Arctic Flora and Fauna, 2013).

23. Post, E. *et al.* The polar regions in a 2°C warmer world. *Sci. Adv.* **5**, eaaw9883 (2019).

24. Taylor, J. J. *et al.* Arctic terrestrial biodiversity status and trends: A synopsis of science
supporting the CBMP State of Arctic Terrestrial Biodiversity Report. *Ambio* **49**, 833–847 (2020).

25. Fauchald, P., Park, T., Tømmervik, H., Myneni, R. & Hausner, V. H. Arctic greening from warming
promotes declines in caribou populations. *Sci. Adv.* **3**, e1601365 (2017).

26. Wilson, S. D. & Nilsson, C. Arctic alpine vegetation change over 20 years. *Glob. Change Biol.* **15**,
1676–1684 (2009).

27. Elmendorf, S. C. *et al.* Plot-scale evidence of tundra vegetation change and links to recent
summer warming. *Nat. Clim. Change* **2**, 453–457 (2012).

28. Maliniemi, T., Kapfer, J., Saccone, P., Skog, A. & Virtanen, R. Long-term vegetation changes of
treeless heath communities in northern Fennoscandia: Links to climate change trends and
reindeer grazing. *J. Veg. Sci.* **29**, 469–479 (2018).

29. Vowles, T. *et al.* Expansion of deciduous tall shrubs but not evergreen dwarf shrubs inhibited by
reindeer in Scandes mountain range. *J. Ecol.* **105**, 1547–1561 (2017).

30. Vowles, T. & Björk, R. G. Implications of evergreen shrub expansion in the Arctic. *J. Ecol.* **107**,
650–655 (2019).

31. Tuomi, M. *et al.* Herbivore Effects on Ecosystem Process Rates in a Low-Productive System.
Ecosystems **22**, 827–843 (2018).

32. Hausner, V. H., Engen, S., Brattland, C. & Fauchald, P. Sámi knowledge and ecosystem-based
adaptation strategies for managing pastures under threat from multiple land uses. *J. Appl. Ecol.*
57, 1656–1665 (2020).

33. Wardle, D. A., Nilsson, M.-C., Gallet, C. & Zackrisson, O. An ecosystem-level perspective of
allelopathy. *Biol. Rev.* **73**, 305–319 (1998).

34. Tybirk, K. *et al.* Nordic Empetrum Dominated Ecosystems: Function and Susceptibility to
Environmental Changes. *AMBIO J. Hum. Environ.* **29**, 90–97 (2000).

35. Aerts, R. Nitrogen-dependent recovery of subarctic tundra vegetation after simulation of
extreme winter warming damage to Empetrum hermaphroditum. *Glob. Change Biol.* **16**, 1071–
1081 (2010).

36. Preece, C. & Phoenix, G. K. Impact of early and late winter icing events on sub-arctic dwarf
shrubs. *Plant Biol.* **16**, 125–132 (2014).

37. Bråthen, K. A., Fodstad, C. H. & Gallet, C. Ecosystem disturbance reduces the allelopathic effects
of Empetrum hermaphroditum humus on tundra plants. *J. Veg. Sci.* **21**, 786–795 (2010).

38. González, V. T., Lindgård, B., Reiersen, R., Hagen, S. B. & Bråthen, K. A. Niche construction
mediates climate effects on recovery of tundra heathlands after extreme event. *PLOS ONE* **16**,
e0245929 (2021).

39. Nilsson, M.-C. & Wardle, D. A. Understory vegetation as a forest ecosystem driver: evidence
from the northern Swedish boreal forest. *Front. Ecol. Environ.* **3**, 421–428 (2005).

40. Olofsson, J. *et al.* Herbivores inhibit climate-driven shrub expansion on the tundra. *Glob. Change*
Biol. **15**, 2681–2693 (2009).

41. Iversen, M. *et al.* Phenology and Cover of Plant Growth Forms Predict Herbivore Habitat
Selection in a High Latitude Ecosystem. *PLoS ONE* **9**, e100780 (2014).

42. Krebs, C. J. Two complementary paradigms for analysing population dynamics. *Philos. Trans. R.*
Soc. Lond. B. Biol. Sci. **357**, 1211–1219 (2002).

43. James, S. P. Legal rights and nature's contributions to people: Is there a connection? *Biol.*
Conserv. **241**, 108325 (2020).

44. Sinclair, A. R. E. & Krebs, C. J. Complex numerical responses to top–down and bottom–up
processes in vertebrate populations. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **357**, 1221–1231
(2002).

45. Stien, A., Tveraa, T., Ims, R. A., Stien, J. & Yoccoz, N. G. Unfounded claims about productivity
beyond density for reindeer pastoralism systems. *Pastoralism* **11**, 20 (2021).
46. Vuorinen, K. E. M. *et al.* Open tundra persist, but arctic features decline—Vegetation changes in
the warming Fennoscandian tundra. *Glob. Change Biol.* **23**, 3794–3807 (2017).
47. Parker, T. C. *et al.* Shrub expansion in the Arctic may induce large-scale carbon losses due to
changes in plant-soil interactions. *Plant Soil* **463**, 643–651 (2021).
48. Kivinen, S., Rasmus, S., Jylhä, K. & Laapas, M. Long-Term Climate Trends and Extreme Events in
Northern Fennoscandia (1914–2013). *Climate* **5**, 16 (2017).
49. Weijers, S., Pape, R., Löffler, J. & Myers-Smith, I. H. Contrasting shrub species respond to early
summer temperatures leading to correspondence of shrub growth patterns. *Environ. Res. Lett.*
13, 034005 (2018).
50. Bråthen, K. A. *et al.* Induced Shift in Ecosystem Productivity? Extensive Scale Effects of Abundant
Large Herbivores. *Ecosystems* **10**, 773–789 (2007).
51. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **80**,
(2017).
52. le Roux, P. C., Lenoir, J., Pellissier, L., Wisz, M. S. & Luoto, M. Horizontal, but not vertical, biotic
interactions affect fine-scale plant distribution patterns in a low-energy system. *Ecology* **94**,
671–682 (2013).
53. Skarin, A. *et al.* Reindeer use of low Arctic tundra correlates with landscape structure. *Environ.*
Res. Lett. **15**, 115012 (2020).
54. Vaas, J., Driessen, P. P. J., Giezen, M., van Laerhoven, F. & Wassen, M. J. Moving from Latent to
Manifest Problem: Trajectories Across Scientific and Public Salience of Invasive Alien Species.
Environ. Manage. **67**, 901–919 (2021).
55. Hughes, T. P., Linares, C., Dakos, V., van de Leemput, I. A. & van Nes, E. H. Living dangerously on
borrowed time during slow, unrecognized regime shifts. *Trends Ecol. Evol.* **28**, 149–155 (2013).
56. González, V. T. *et al.* Batatasin-III and the allelopathic capacity of Empetrum nigrum. *Nord. J. Bot.*
33, 225–231 (2015).
57. Kuussaari, M. *et al.* Extinction debt: a challenge for biodiversity conservation. *Trends Ecol. Evol.*
24, 564–571 (2009).
58. Gilbert, B. & Levine, J. M. Plant invasions and extinction debts. *Proc. Natl. Acad. Sci.* **110**, 1744–
1749 (2013).
59. Kaarlejärvi, E. a, Hoset, K. S. b & Olofsson, J. a. Mammalian herbivores confer resilience of Arctic
shrub-dominated ecosystems to changing climate. *Glob. Change Biol.* **21**, 3379–3388 (2015).
60. Van Wijk, M. T. *et al.* Long-term ecosystem level experiments at Toolik Lake, Alaska, and at
Abisko, Northern Sweden: generalizations and differences in ecosystem and plant type
responses to global change. *Glob. Change Biol.* **10**, 105–123 (2004).
61. Graae, B. J. *et al.* Stay or go – how topographic complexity influences alpine plant population
and community responses to climate change. *Perspect. Plant Ecol. Evol. Syst.* **30**, 41–50 (2018).
62. Hoset, K. S. *et al.* Changes in the Spatial Configuration and Strength of Trophic Control Across a
Productivity Gradient During a Massive Rodent Outbreak. *Ecosystems* 1–15 (2017)
doi:10.1007/s10021-017-0124-1.
63. Olofsson, J., Tømmervik, H. & Callaghan, T. V. Vole and lemming activity observed from space.
Nat. Clim. Change **2**, 880–883 (2012).
64. Oksanen, T. *et al.* The impact of thermal seasonality on terrestrial endotherm food web
dynamics: a revision of the Exploitation Ecosystem Hypothesis. *Ecography* **43**, 1859–1877
(2020).
65. Schmidt, N. M., Mosbacher, J. B., Vesterinen, E. J., Roslin, T. & Michelsen, A. Limited dietary
overlap amongst resident Arctic herbivores in winter: complementary insights from
complementary methods. *Oecologia* **187**, 689–699 (2018).

66. Jenkins, D. A., Lecomte, N., Andrews, G., Yannic, G. & Schaefer, J. A. Biotic interactions govern
the distribution of coexisting ungulates in the Arctic Archipelago – A case for conservation
planning. *Glob. Ecol. Conserv.* **24**, e01239 (2020).
67. Bell, J. N. B. & Tallis, J. H. Empetrum Nigrum L. *J. Ecol.* **61**, 289 (1973).
68. Valéry, L., Fritz, H. & Lefeuvre, J.-C. Another call for the end of invasion biology. *Oikos* **122**,
1143–1146 (2013).
69. Carey, M. P., Sanderson, B. L., Barnas, K. A. & Olden, J. D. Native invaders – challenges for
science, management, policy, and society. *Front. Ecol. Environ.* **10**, 373–381 (2012).
70. Shackleton, R. T., Shackleton, C. M. & Kull, C. A. The role of invasive alien species in shaping local
livelihoods and human well-being: A review. *J. Environ. Manage.* **229**, 145–157 (2019).
71. Bråthen, K. A., Pugnaire, F. I. & Bardgett, R. D. The paradox of forbs in grasslands and the legacy
of the mammoth steppe. *Front. Ecol. Environ.* **19**, 584–592 (2021).
72. Ruckelshaus, M. H. *et al.* The IPBES Global Assessment: Pathways to Action. *Trends Ecol. Evol.*
35, 407–414 (2020).
73. van der Wal, R., Fischer, A., Selge, S. & Larson, B. M. H. Neither the public nor experts judge
species primarily on their origins. *Environ. Conserv.* **42**, 349–355 (2015).
74. Davis, K. J., Chadès, I., Rhodes, J. R. & Bode, M. General rules for environmental management to
prioritise social ecological systems research based on a value of information approach. *J. Appl.*
Ecol. **56**, 2079–2090 (2019).
75. Wang, H. *et al.* Satellite-derived NDVI underestimates the advancement of alpine vegetation
growth over the past three decades. *Ecology* **102**, e03518 (2021).
76. Lindenmayer, D. B. & Likens, G. E. Adaptive monitoring: a new paradigm for long-term research
and monitoring. *Trends Ecol. Evol.* **24**, 482–486 (2009).
77. Marteinsdóttir, B. *et al.* GróLind – Sustainable Land Use Based on Ecological Knowledge. *Int.*
Grassl. Congr. Proc. 6 (2021).
78. Ims, R. A. & Yoccoz, N. G. Ecosystem-based monitoring in the age of rapid climate change and
new technologies. *Curr. Opin. Environ. Sustain.* **29**, 170–176 (2017).
79. Webb, N. R. The traditional management of European heathlands. *J. Appl. Ecol.* **35**, 987–990
(2008).
80. Vandvik, V., Heegaard, E., Måren, I. E. & Aarrestad, P. A. Managing heterogeneity: the
importance of grazing and environmental variation on post-fire succession in heathlands. *J.*
Appl. Ecol. **42**, 139–149 (2005).
81. Critchley, C. N. R., Burke, M. J. W. & Stevens, D. P. Conservation of lowland semi-natural
grasslands in the UK: a review of botanical monitoring results from agri-environment schemes.
Biol. Conserv. **115**, 263–278 (2004).
82. Keech, O., Carcaillet, C. & Nilsson, M.-C. Adsorption of allelopathic compounds by wood-derived
charcoal: the role of wood porosity. *Plant Soil* **272**, 291–300 (2005).
83. Pivello, V. R., Vieira, M. V., Grombone-Guaratini, M. T. & Matos, D. M. S. Thinking about super-
dominant populations of native species – Examples from Brazil. *Perspect. Ecol. Conserv.* **16**, 74–
82 (2018).
84. Shackelford, N., Renton, M., Perring, M. P. & Hobbs, R. J. Modeling disturbance-based native
invasive species control and its implications for management. *Ecol. Appl.* **23**, 1331–1344 (2013).
85. Valéry, L., Fritz, H., Lefeuvre, J.-C. & Simberloff, D. Invasive species can also be native…. *Trends*
Ecol. Evol. **24**, 585 (2009).
86. Pedersen, Å. Ø. *Norwegian Arctic tundra: a panel-based assessment of ecosystem condition*.
(Norsk Polarinstitutt, 2021).
87. Muggeo, V. M. R. Interval estimation for the breakpoint in segmented regression: a smoothed
score-based approach. *Aust. N. Z. J. Stat.* **59**, 311–322 (2017).
88. Bråthen, K. A. & Hagberg, O. More efficient estimation of plant biomass. *J. Veg. Sci.* **15**, 653–660
(2004).

89. Ravolainen, V. T. *et al.* Additive Partitioning of Diversity Reveals No Scale-dependent Impacts of
Large Ungulates on the Structure of Tundra Plant Communities. *Ecosystems* **13**, 157–170 (2010).
90. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for
Statistical Computing, 2021).
91. Muth, C., Oravecz, Z. & Gabry, J. User-friendly Bayesian regression modeling: A tutorial with
rstanarm and shinystan. *Quant. Methods Psychol.* **14**, 99–119 (2018).
92. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer Science & Business Media,
2009).
93. Kay, M. *ggdist: Visualizations of Distributions and Uncertainty*. (2021).
doi:10.5281/zenodo.3879620.
588
589