Greening conceals evergreening: contrasting trends for a socio-ecological system in Arctic

Europe

Tuomi M.W.[1*†], Utsi T.Aa.[2], Yoccoz N.G.[1], Armstrong C.W.[3], Gonzalez V.[4], Hagen S.B.[4],

Jónsdóttir I.S.[5], Pugnaire F.I.[6], Shea K.[7], Wardle D.A.[8,9], Zielosko S.[1], Bråthen K.A.[1*†]

Affiliations

[1] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Tromsø, Norway

[2] Department of Arctic and Marine Biology, UiT – Arctic University of Norway, Alta, Norway

[3] The Norwegian College of Fishery Science, UiT – Arctic University of Norway, Tromsø, Norway

[4] Department of Ecosystems in the Barents region, Norwegian institute of Bioeconomy Research-NIBIO,

Svanvik, Norway

[5] Institute of Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland

[6] Estación Experimental de Zonas Áridas, Consejo Superior de Investigaciones Científicas (EEZA-CSIC), Almería,

Spain

[7] Department of Biology and IGDP in Ecology, The Pennsylvania State University, University Park, Pennsylvania,

16802 USA

[8] Asian School of the Environment, Nanyang Technological University, Singapore

[9] Department of Ecology and Environmental Science, Umeå University, SE-901 87 Umeå, Sweden

ORCID:

Maria W. Tuomi 0000-0002-7154-5177, Utsi T.Aa. 0000-0003-0776-2369, Yoccoz N. 0000-0003-2192-1039,

Armstrong C.W. 0000-0002-1837-9165, Gonzalez V. 0000-0002-3997-1803, Hagen S.B. 0000-0001-8289-7752,

Jónsdóttir I.S. 0000-0003-3804-7077, Pugnaire F.I. 0000-0002-1227-6827, Shea K. 0000-0002-7607-8248,

Wardle D.A. 0000-0002-0476-7335, Zielosko S. 0000-0002-6259-0873, Bråthen K.A. 0000-0003-0942-1074

* corresponding authors

maria.w.tuomi@uit.no, kari.brathen@uit.no

†These authors contributed equally to this work

34 **Open Research Statement**

35 The datasets collected and analyzed in this study as well as reproducible R scripts for analysis are upon

36 acceptance made publicly available (CC-BY license) via UiT repository (https://dataverse.no/) at DOI: ---.

Abstract

1. Ongoing Arctic greening can increase productivity and reindeer pasture quality in the tundra. However, greening may also entail proliferation of unpalatable species, with distinct consequences for pastoral socio-ecological systems (SES).

2. We show extensive greening across 20 reindeer districts in northern Norway between 2003 and 2020. The allelopathic, evergreen dwarf-shrub crowberry biomass increased by 60%, contrasted by smaller increases of deciduous dwarf-shrubs and stagnating forb and graminoid biomass.

3. We found evidence, although uncertain, of a negative relationship between biomass and reindeer densities, but only among forbs, the least abundant plant group.

4. Our results challenge the management decision-making, which aims at sustainable pasture management, but which assumes stationary density-dependent relationships. Changes in temporal vs. spatial relationships should be included as management criteria, to avoid mismanaging a SES in transition.

5. Large-scale shift towards increased allelopathy may undermine the resource base of a key Arctic herbivore and pastoral SES.

Keywords

Arctic greening, *Empetrum nigrum*, deciduous shrubs, forbs, graminoids, pastoralism, *Rangifer*

 Introduction

Rapid changes in the Arctic climate (IPCC 2021) are altering primary productivity,

biodiversity and ecosystem functions (I. H. Myers-Smith et al. 2020; Barry, Berteaux, and

Bültmann 2013; Post et al. 2019; Taylor et al. 2020), with effects cascading to and

interacting with herbivore populations and coupled social-ecological systems (SES).

Prevalent warming-induced vegetation trends include increased productivity, biomass and

leaf area, identified as ecological greening of the Arctic (I. H. Myers-Smith et al. 2020), and

shifts towards more resource acquisitive plant species (Bjorkman et al. 2018; Elina

Kaarlejärvi, Eskelinen, and Olofsson 2013). In practice, such changes are expected to

manifest as increased abundance of plants with taller stature and higher nitrogen (N)

concentrations, especially willows and other deciduous shrubs, but also graminoids and

forbs (Bråthen, Gonzalez, and Yoccoz 2018; Bjorkman et al. 2018; I. H. Myers-Smith et al.

2019). In areas where shrubs are already present or dominant, their growth can occur via

increased cover, i.e. infilling of existing patches, or accumulation of biomass, i.e. vertical

growth (I. H. a b Myers-Smith et al. 2011). Recent observations, however, indicate that

Arctic vegetation changes conceal functionally contrasting trends. Field observations show

increases especially in evergreen dwarf-shrubs across the circumpolar Arctic (Wilson and

Nilsson 2009; Elmendorf et al. 2012; Maliniemi et al. 2018; Vowles et al. 2017; Vuorinen et

al. 2017; Stewart et al. 2018). Evergreen plants often have high phenolic and low N content,

and can contribute to lowering ecosystem productivity (Vowles and Björk 2019; Tuomi et al.

2018). An "evergreening" trend may therefore be functionally distinct from that of greening

by deciduous plants, as it may suggest an ongoing decline in process rates, herbivore forage

quality and biodiversity (Bråthen, Gonzalez, and Yoccoz 2018), despite increasing biomass in

vegetation.

Proliferation of poorly palatable vs. palatable plants will likely have distinct

consequences for herbivore populations and associated socio-ecological systems. In a

rapidly changing Arctic, reindeer herding systems are readily subject to multiple climate and

anthropogenic stressors that affect food availability (e.g. rain-on-snow events that lead to

impenetrable ice layers that prevent foraging) and access to pasture (e.g. tourism and

infrastructure) (Horstkotte et al. 2017; Hausner et al. 2020). Until now, however, changes in

Arctic vegetation composition have been little considered among the novel threats to

caribou and reindeer populations and indigenous pastoral systems (Post et al. 2009; Mallory

and Boyce 2017; Horstkotte et al. 2017).

Rangifer (reindeer and caribou) are the most numerous Arctic ungulate, with grazing

systems that span the circumpolar area (Mallory and Boyce 2017), and that are core to the

indigenous Sámi livelihood and culture (James 2020). Akin to all extensive pastoral or

grazing systems, *Rangifer* rely on high-quality plant resources (Sinclair and Krebs 2002; K. L.

Parker, Barboza, and Gillingham 2009). Increasing plant productivity and abundance of N-

rich and palatable herbaceous and deciduous species are expected to translate into higher

survival and population growth rates of reindeer through positive bottom-up effects (K. L.

Parker, Barboza, and Gillingham 2009; Tveraa et al. 2013). In contrast, poorly palatable

species may reduce pasture productivity and quality to the extent that reindeer avoid areas

where their dominance is high (Iversen et al. 2014). In a recent example, the satellite-

derived greening signal in North-America was negatively related to caribou population

growth rates, which was attributed to expansion of deciduous shrubs with high levels of

anti-browsing defenses (Fauchald et al. 2017). The balance between the proliferation of

poorly palatable and palatable plant species in *Rangifer* pastures across space and time,

however, is poorly known.

Arctic terrestrial ecosystems often host strong herbivore-plant interactions, with co-

occurring bottom-up and top-down dynamics (L. Oksanen and Oksanen 2000; Hoset et al.

2017; Tveraa et al. 2007). Arctic reindeer and caribou have highly context-dependent

impacts on vegetation (Bernes et al. 2015), although they theoretically can strongly impact

plant biomass along gradients of productivity through top-down effects (L. Oksanen and

Oksanen 2000). In line with top-down effects, *Rangifer* have been found to modulate

warming-induced vegetation changes. For instance, reindeer can counteract shrubification

(Christie et al. 2015), e.g. keeping nutritious willows in a "browse-trap" (Bråthen et al.

2017), and thereby prevent the even more nutritious forbs and graminoids from being

overgrown (Bret-Harte et al. 2001; Mod and Luoto 2016). Suppression of palatable species

may, however, not be representative of how herbivores affect less palatable plants. For

instance, the ability of *Rangifer* to modulate warming-induced changes in dominant,

nutrient-poor evergreen dwarf-shrubs is less clear (Maliniemi et al. 2018; Vowles et al.

2017).

Central to many Arctic ecosystems and their change is the evergreen dwarf-shrub 124   crowberry (*Empetrum nigrum*), a niche-constructing, allelopathic species (Bråthen, 125   Gonzalez, and Yoccoz 2018; Wardle et al. 1998) (Supplementary Content Table). In 126   Fennoscandia, crowberry is already abundant and commonly dominant. As a boreal-Arctic 127   species it appears to thrive under warming climate in tundra and it tolerates various 128   environmental stressors (Tybirk et al. 2000; Aerts 2010; Preece and Phoenix 2014; González 129   et al. 2019). Owing to its low foliar nutrient concentrations, high levels of allelopathic 130   polyphenolic compounds in its leaves and a dense, clonal growth form, crowberry is highly 131   unpalatable, and can significantly retard ecosystem processes such as litter decomposition, 132   soil nutrient fluxes and seedling establishment (Bråthen, Fodstad, and Gallet 2010; González 133   et al. 2021; Tybirk et al. 2000; Wardle et al. 1998; Nilsson and Wardle 2005). Its dominance 134   is related to suppressed biodiversity and herbaceous plant growth in summer pastures 135   (Bråthen, Gonzalez, and Yoccoz 2018) (Supplementary Content Table), and reindeer have 136   been found to avoid crowberry-dominated areas for most of the growing season (Iversen et 137   al. 2014). Nevertheless, higher reindeer densities may be associated with spatially higher 138   abundance of crowberry (Bråthen, Gonzalez, and Yoccoz 2018; Bråthen et al. 2007).

139   In the present work, we analyze vegetation across 20 reindeer districts in Norway to 140   assess both temporal changes and spatial variation in reindeer summer pastures, and to 141   what extent the associated Norwegian reindeer management (*Reindeer Husbandry Act* 142   2007) is capturing these changes (Fig. 1). The current reindeer management decision-143   making process aims at sustainable management of pastures (*Reindeer Husbandry Act* 2007; 144   Ministry of Agriculture 2017, 32). However, the indicators used in the decision process do 145   not measure the summer pastures and their quality directly, as the indicators are assumed 146   to reflect the state of the system. Instead, the decision-making is based on an empirically 147   supported negative density-dependent relationship between reindeer numbers and body 148   mass as the focal indicator (Henden et al. 2020; Stien et al. 2021) (Fig. 1c, Supplementary 149   Figure 2), and this relationship is assumed to be unaffected by long-term changes (Krebs 150   2002). Hence, the pastures themselves are not directly included in the decision-making 151   process, as management indicators such as reindeer number and weights are easily 152   accessible and they are linked to the economic valuation of the industry. The consequence is 153   that long-term changes in productivity – a key determinant of sustainability of pastures and 154   the husbandry – are managed through mandatory reductions of reindeer densities, easing

the assumed top-down regulation of the pastures (Fig. 1b) whenever animal slaughter
weights fall below a threshold. However, negative density-dependence in animal
populations is not due only to changes in resources (Krebs 2002), and pasture productivity
and composition may develop independently of animal densities, leading to bottom-up
effects (Fig. 1b).

In contrast to the current management decision making, which emphasizes short-
term and homogenous spatial and temporal effects, a community ecological perspective
sees the pastures themselves as not uniform nor stable (Fig. 1b, d). There is a large variation
in nutrient content between the plant functional groups in pastures (Fig. 1d), and the plant
groups may respond differently to herbivory (Fig. 1e) and to other temporal and spatial
factors, such as climate (temporal confounding) and bedrock nutrient content (spatial
confounding). Therefore, plant responses to reindeer densities are likely to vary between
plant functional groups in both space and/or time. Spatial and temporal patterns of plant
groups and community compositions are therefore important determinants of changes in
pasture quality (Fig. 1d), and hence of reindeer nutrition and growth (Fig. 1e). For instance,
pastures abundant with the most nutritious plant groups could support high calf weights,
whereas weights may saturate at lower levels if pastures consist mainly of crowberry (Fig
1d, e). In 2003, crowberry made up the majority of plant biomass across the northern
Norwegian tundra, already resulting in pastures that were abundant in poor-quality forage
of low nitrogen content (Fig. 1d).

We apply a unique, large-scale resurvey of 292 plant communities within 56
landscape areas sampled in 2003 and 2020 across 20 reindeer districts in northern Norway
(Supplementary Figure 1) to assess variation in vegetation biomass relative to reindeer
densities. Specifically, we ask 1) how preferred forage plants (forbs, graminoids and
deciduous dwarf-shrubs) and less palatable evergreen dwarf-shrubs – with focus on
crowberry – vary with reindeer density across space and in time, and 2) what are the
implications of these changes for reindeer management decision-making. Given the ongoing
warming trend in the region (Fig. 2), we hypothesize that all species and functional groups
increase in abundance, including crowberry and other evergreen dwarf-shrubs. However, if
the existing decision making process is adequate for achieving a sustainable management of
pastures and the husbandry, observed changes in vegetation should be linked to reindeer
density, or its changes. We thus ask if palatable plant groups have a more positive

abundance response in management districts with low-to-intermediate or decreasing

reindeer density (cf. Fig. 1e), and if evergreen dwarf shrubs have a more positive abundance

response with high or increasing reindeer density. If the current management decision

making is insufficient or incomplete, we expect opposite patterns (for dwarf-shrubs, Fig. 1e),

or no association to reindeer density.

193   Methods

Located at the transition from subarctic to the low-Arctic, the Northern

Fennoscandian study system is a bellwether for changes in tundra vegetation (Vuorinen et

al. 2017; Elmendorf et al. 2012) and ecosystem functioning (T. C. Parker et al. 2021). The

region has experienced lengthening of the growing season as spring and autumn have

become warmer over the past century (Kivinen et al. 2017). With an approximate increase

of 100 growing degree days in the studied summer pastures since the late 1970's (Fig. 2),

the pastures have been subject to a growing season prolongation, which is a key driver

behind increased shrub growth in the Arctic (Post et al. 2019; Weijers et al. 2018; I. H.

Myers-Smith et al. 2019). We revisited a hierarchical sampling design from 2003, with plant

communities nested in blocks and reindeer districts (Bråthen et al. 2007) (Supplementary

Figure 1). We used identical georeferenced plant communities and sampling methods within

each community in both 2003 and 2020. Each plant community was represented by a 50m

long transect with 11 equidistant, triangular measurement plots with 40cm sides (Bråthen et

al. 2007) (Supplementary Figure 1). We analyzed plant data as biomass (g/m^2) and cover

(number of plots within transect with observed presence).

We decomposed the reindeer density to its spatial, temporal, and residual

components (Oedekoven et al. 2017), and standardized all three predictors to a mean of 0

and variance of 1 for better effect comparability and model convergence. The temporal

component represents the change in average reindeer densities between the two time

points, the spatial component the difference in the time-averaged district reindeer densities

and the residuals the district-specific change in reindeer densities that is not captured in the

average spatial and temporal densities. Models for forbs and graminoids included a

quadratic term for the spatial component, while deciduous dwarf-shrubs and crowberry

included a linear term only (cf. Fig 1e), and all models included block and district as group-

level intercepts. We fitted Bayesian generalized linear mixed models (Bürkner 2017) with

weakly informative default priors, and checked model convergence and independence of
HMC chains based on the $\hat{R}$ statistics (< 1.002) and effective sample size (> 2000) (Muth,
Oravecz, and Gabry 2018). Lack of spatial autocorrelation in group-level effects and model
residuals was assessed visually. We used averaged reindeer densities from the previous 23
years for 2003 (1980-2003, same data as in Bråthen et al. 2007) and 17 years for 2020 to
avoid temporal overlap in the data (2003-2019, Ministry of Agriculture, n.d., accessed via
reinbase.no).
226
227   Results
Across the entire study area, crowberry standing biomass increased by 60% (Fig. 3a,
Supplementary Result Table 1A), and cover by 14% (Supplementary Result Table 1B and 2)
from 2003 to 2020. The increase of deciduous dwarf-shrub biomass was nearly an order of
magnitude smaller than that of crowberry's (Fig. 3a, Table 1), and we found no change in
the cover of deciduous dwarf-shrubs. The average reindeer densities declined from 6.47 to
6.24 animals/km^2 over the same time period. The change in crowberry and deciduous
dwarf-shrubs and shrubs was spatially consistent, as they increased in biomass in 90% and
85% of the districts, respectively. The increase in the woody shrubs' biomass, crowberry in
particular, was in sharp contrast with no temporal effects on the biomass and cover of forbs
and graminoids (Fig. 3a, Table 1, Supplementary Result Table 2). The frequency of pasture
communities in which more than 25% of total vascular biomass was crowberry rose over
this period from an already high 0.76 to 0.83.
While the overall temporal change in reindeer densities did not translate to change
in forbs, we found that the spatial pattern in district-level reindeer densities had a negative
relationship with forb biomass (Table 1). This effect had, however, a relatively high
uncertainty (Table 1), and it tended to become weaker in 2020 (Fig. 3c). The shape of the
relationship was also different from the expectation, with lowest estimated biomass at
intermediate reindeer densities and a non-significant quadratic term (Table 1). There was no
evidence that the spatial differences in reindeer densities explained variation in any of the
other plant groups.
In the studied reindeer districts over the study period, densities declined in 50%,
increased in 45% and stayed stable in 5% of reindeer districts (calculated with a change
threshold of 0.1 reindeer/km^2). None of these district-specific changes in density explained

variation in any of the plant groups (Table 1, Supplementary Figure 3). In our dataset, the

spatial scale for important group-level effects differed between plant groups. Crowberry

biomass varied substantially among landscape areas within reindeer districts, but little at

the scale of districts (Fig. 3b, Supplementary Result Table 3). Deciduous shrubs and

graminoids varied markedly between both districts and among landscape areas. In contrast,

variation in forb biomass was largest between – and not within – districts (Fig. 3b,

Supplementary Result Table 3).

259   Discussion

In line with our first expectation, we found substantial increases in woody vegetation

biomass. The difference in magnitude between the evergreen and deciduous woody plant

proliferation, however, was surprising, with pasture evergreening far outpacing greening by

deciduous shrubs. We found no evidence that variation in woody species linked with spatial

density patterns nor with district-specific changes in reindeer densities. Therefore, it is

plausible that the observed temporal changes are not attributable to the reindeer density

change *per se*, but are mainly confounded by other temporal effects, such as changes in

climate (Fig 2), corroborating the hypothesized evergreening and greening trends.

Proliferation of crowberry appears to mainly occur through accumulation of biomass

(vertical growth), but also through infilling, whereby ever larger surface areas are potentially

impacted by crowberry.

In contrast to the expectation of temporal change, the least abundant, yet most

species-rich and productive plant groups, the forbs and the graminoids, show no change

over 18 years. This result was also surprising, given the overall decrease in reindeer

densities and an ongoing warming trend, which both could be expected to favor herbaceous

growth forms. Our results indicate that the spatial negative relationship of forbs and

reindeer density may not be informative to predict changes over time. We note that the

abundance of forbs and graminoids is extremely low, their rarity making assessment of

patterns difficult at a large spatial scale and increasing uncertainty of the estimates.

Nevertheless, the persisting rarity of the most productive plant groups combined with rapid

proliferation of the least palatable evergreen dwarf-shrubs across landscapes is likely of high

importance for pasture management and the pastoral SES.

Our results point at evergreening and crowberry proliferation as a potential major
bottom-up forcing on the pastures (Fig. 3a, c) (cf. le Roux et al. 2013), effectively decoupled
from district-level variation in reindeer density. Sustainable management of tundra pastures
and the pastoral SES will be contingent on models that are representative of the managed
system and its spatial and temporal uncertainties (Milner-Gulland and Shea 2017; Regan,
Colyvan, and Burgman 2002; Latombe et al. 2019). An insufficient decision making process
that misses influential variables or processes may severely undermine management
objectives (Madden and McQuinn 2014), and lead to non-resilient management efforts,
reducing the capacity of the system to adapt and maintain system resilience against
undesirable states (Dudney et al. 2018). A pastoral management practice that does not
monitor changes in the plant resource *per se*, such as the Norwegian reindeer management
decision-making process, would likely remain functional or even thrive under increasing
productivity and greening of palatable plants. However, managers would be ill equipped to
detect and manage bottom-up effects that slowly reduce pasture quality. In the following,
we argue that if left unchecked, allelopathic evergreening may have severe, adverse, long-
term consequences for the diversity, productivity and resilience of reindeer pastures and
the pastoral SES.
299
300   *Evergreening: allelopathy and the mechanisms of transition*
As a niche-constructing species, crowberry can modify the environment once
established (González et al. 2021; Bråthen, Gonzalez, and Yoccoz 2018; Wardle et al. 1998),
as allelochemicals in crowberry's leaves and accumulating litter can push the system
towards a state of strong allelopathy (Fig. 4). The growth-inhibiting effects of crowberry
litter can remain after the plant itself is gone (Pilsbacher et al. 2021; Dorrepaal, Cornelissen,
and Aerts 2007; Aerts 2010), giving rise to legacy effects. Diminishing diversity due to
allelopathy happens gradually through reduced seedling recruitment (González et al. 2015;
2021), and over longer time scales induce an extinction debt on the local plant communities
(Kuussaari et al. 2009; Gilbert and Levine 2013), as local seedbanks disappear (Fig. 4a).
These legacy effects, along with the potential longevity, recovery potential, dense growth
and poorly palatable leaves of crowberry (Aerts 2010), suggests crowberry dominance of
communities is likely to be a highly resilient state (E.a Kaarlejärvi, Hoset, and Olofsson 2015;
González et al. 2021; Wardle et al. 1998; Nilsson and Wardle 2005). Once established, the

state may require strong external disturbances to reverse (Fig. 4b, left panel). Consequently,
we hypothesize that the long-term effects from crowberry allelopathy at the community-
and ecosystem-level may represent an ongoing shift with potential context-specific
thresholds (Hughes et al. 2013) (Fig. 4b, left panel).

While we show that crowberry proliferation occurs in landscapes across northern
Fennoscandia, we also find spatial variability in its increase especially at relatively local
landscape scales, indicating there are ecological contexts in which the shift is stronger,
weaker, or absent. Tundra vegetation changes are repeatedly found to be spatially
heterogeneous across scales (Bjorkman et al. 2018; Van Wijk et al. 2004; Graae et al. 2018)
and can be linked to e.g., variation in microclimate or herbivory. For instance, cyclic small
rodent outbreaks across the resurveyed region can decimate dwarf-shrubs and especially
crowberry in patches across landscapes (Tuomi et al. 2018; Hoset et al. 2017)
(Supplementary Figure 4) - a temporary reduction even detectable from space (Olofsson,
Tømmervik, and Callaghan 2012). However, strong localized small rodent grazing has not
limited the overall high-magnitude, long-term encroachment of crowberry documented
here.

While neither small nor large herbivores may halt the overall trend of crowberry
encroachment across tundra pastures, they are also likely adversely affected by the
observed low abundance of forbs and diminishing prevalence of communities with little
crowberry (T. Oksanen et al. 2020; Iversen et al. 2014). Apart from reindeer, other
endotherm herbivores, such as small rodents, ptarmigans, domesticated sheep and musk-ox
also rely on N-rich forage (Schmidt et al. 2018), and seek pastures with high-quality food
(Iversen et al. 2014; Jenkins et al. 2020; Skarin et al. 2020). Furthermore, crowberry being a
wind-pollinated plant (Bell and Tallis 1973), its encroachment may over time also affect
insect pollinators. In summary, crowberry encroachment of the magnitude documented
here – and predicted e.g. for Arctic Greenland (Stewart et al. 2018) – raises significant
concerns of cascading effects onto the tundra biota, ecosystem and human beneficiaries
relying on them, including pastoralists, sheep farmers and game hunters.

343   *Way forward: monitor and manage*
In the context of the reindeer management system, an evergreening trajectory
towards higher evergreen dwarf-shrub (particularly crowberry) abundance with potentially

lowered pasture process rates, forage availability and carrying capacity, threaten a risk of
mismanagement and loss of resilience of the pastoral SES (Valéry, Fritz, and Lefeuvre 2013;
Carey et al. 2012; Shackleton, Shackleton, and Kull 2019) (Fig. 4b, right panel). Our results
suggest that revision of the current reindeer management decision-making process should
add an adaptive component sensitive to long-term, decadal changes to the current
approach focusing on relatively short-term fluctuations. Inclusion of measures relating to
pasture plant diversity and productivity, in line with intentions for sustainable reindeer
husbandry (*Reindeer Husbandry Act* 2007) and for translating the IPBES Global Assessment
to policy (Ruckelshaus et al. 2020), will be crucial. In this regard, particular attention should
be given to the most species rich and nutritious, yet scarce growth form, the forbs (Bråthen,
Pugnaire, and Bardgett 2021).

Presently, however, crowberry proliferation, and its ecological and socio-economic
impacts, remain poorly understood; this can impede development of novel management
norms and objectives (Vaas et al. 2021; Davis et al. 2019). First, general acceptance and
evidence-based baselines are often difficult to establish for slow and poorly detectable
changes (Schneider, Leifeld, and Malang 2013), such as the creeping infilling and biomass
accumulation of a slow-growing dwarf-shrub. The trajectory of evergreening is likely long,
but large-scale empirical evidence goes back barely half-a century (Maliniemi et al. 2018).
Second, contemporary validation and monitoring of Arctic vegetation change has relied
heavily on remote sensing indices (I. H. Myers-Smith et al. 2020). Remote sensing may not
necessarily capture changes in functional composition (Wang et al. 2021), distinguish
greening and evergreening as functionally different processes (Vowles and Björk 2019), or
document the species diversity of pastures. Adaptive and long-term ecosystem and pasture
monitoring programmes are key to mitigating uncertainty about climate-driven vegetation
change (Post et al. 2009; I. H. Myers-Smith et al. 2019; Lindenmayer and Likens 2009). Such
programmes have recently been implemented, for example, in Iceland (Marteinsdóttir et al.
2021), and in the Varanger peninsula and Svalbard, in Norway (Ims and Yoccoz 2017).

Our results suggest that the current regulator of the management system (reindeer
numbers) may not function in an efficient manner to support resilient and sustainable
pastures. Additional management strategies, interventions and indicators that directly
address spatial and temporal variation in diversity, productivity and heterogeneity of
pastoral landscapes are needed. Such strategies have centennial or even millennial roots in

extensive management practices of European coastal heathlands (Webb 2008; Vandvik et al.

2005) and in European agri-environmental policy that has focused on preventing loss of

open semi-natural grasslands (Critchley, Burke, and Stevens 2004). Fire has the capacity to

ameliorate soil conditions against crowberry's allelopathic effects (Bråthen, Fodstad, and

Gallet 2010; Keech, Carcaillet, and Nilsson 2005), suggesting management through burning

as one potential way to locally control ecosystem state shifts similar to those in coastal

heathlands (Vandvik et al. 2005) or boreal *Pinus*-dominated forests (Wardle et al. 1998;

Nilsson and Wardle 2005). Here, the challenge lies in the already scarce and diminishing

herbaceous resource, the forbs (Bråthen, Pugnaire, and Bardgett 2021). Scarcity begets

scarcity through increasing seed limitation under encroaching crowberry dominance and

allelopathic effects (González et al. 2021), and promotion of productive vegetation would

likely require re-building seed banks alongside soil amelioration. Spatial rarity also poses

challenges for monitoring efforts. Despite such challenges, we believe a resilient

management approach to tundra reindeer pastures and pastoral SES is both attainable and

urgent.

Conclusions

Evergreening puts an increasing pressure on sustainable land-use planning and

prioritization to preserve remaining forb and graminoid-rich pastures (Webber et al. 2022)

and the biodiversity of tundra landscapes against rapid homogenization (Stewart et al.

2018). Evergreening through crowberry encroachment exemplifies emergence of super-

dominance among native species, a phenomenon linked with anthropogenic pressures or

novel climates across biomes (Pivello et al. 2018; Shackelford et al. 2013; Valéry et al. 2009),

with ecological consequences not unlike those of non-native invasive species.

Overall, our results on crowberry encroachment suggest an ongoing decline in the

quality of pasture land (Skarin et al. 2020; Iversen et al. 2014), adding to ongoing loss of land

with productive pastures due to human disturbance (Hausner et al. 2020) and other

stressors (Vaas et al. 2021; Horstkotte et al. 2017). Loss of pasture quality may additionally

amplify other stressors, further eroding the resilience of an increasingly vulnerable Arctic

system.

Ignoring the capacity of native species to be drivers of change is a critical blind spot

that threatens the biodiversity and sustainability of ecosystems. For the Arctic pastoral

system studied here, the management of tundra ecosystems and pastoral SES post-2020
should align targets of reindeer and biodiversity management and ensure resilience through
ecosystem-based monitoring of the biodiversity of pastures. Only with such comprehensive
approaches will sustainable management of diversity in complex social-ecological systems
be possible in changing climates.
415
416
417   *Methods*
418
Study area and resurvey design. We conducted a vegetation survey of 292 remote, georeferenced
vegetation communities in summer pastures across Northern Norway during peak growing season in
2003 and again in 2020. The study area (lat N69° 25.806'- N70° 58.471', lon E20° 47.186'- E27°
31.099') includes strong climatic gradients from west to east and from coast to inland, altitudinal
variation from 60 to 600 m asl, as well as variation in bedrock types. Sampling included the most
common vegetation types in the region: heaths, mires, snow beds, meadows, and windblown ridges.
The survey design (Supplementary Figure 1) incorporates summer pasture areas of 20 reindeer
districts, where adjacent districts of similar climatic conditions were organized into 10 district pairs
of contrasting reindeer densities, based on average reindeer density from 1980 to 2003. Within each
district, a 2x2 km vegetated grid was assigned out of which a random subset was chosen as
landscape areas for sampling. Within each landscape area, a random set of 25 of the 100, 200 x
200m squares that constitute the vegetated grid were assigned for plant community sampling.
Squares were then sampled by the transect method (each transect representing a plant community),
whereby a 50m transect was placed from the midpoint towards a random GPS-position along a circle
with a 50m radius. We used the point-frequency method in 11 plots every 5m along the transect to
sample each community (Supplementary Figure 1). Each plot was measured by placing a triangular
frame with sides of 40cm and one pin in each corner, counting all intercepts with the vegetation
(Bråthen and Hagberg 2004). The a priori stratification of the original 2003 design, inclusion rules of
transects and sampling method are described in more detail in Bråthen et al. (2007, 2018). In 2020,

we resurveyed a subset of 292 (of the original 1450) communities in 56 (of if the original 151)

landscape areas. The re-sampling retained the geographic extent as well as most of climatic and

abiotic variability. Based on transect descriptions from 2003 (e.g., "transect was moved 10m

backwards due to a lake") we deemed the relocation of communities accurate.

Environmental data. We estimated climatic trends between 1957 and 2019 in the studied summer

pastures for growing degree days (GDD), using data described in Pedersen (2021). We applied

segmented linear regression models using the R-package segmented (Muggeo 2017) to explore

trends and breakpoints in the mean estimates for the regional climate.

We retrieved data on reindeer numbers from the onset of the reindeer herding year (cf. Stien et al.

2021) for each studied reindeer herding district (Ministry of Agriculture, n.d.), which we then divided

by summer pasture area (km^2) to obtain reindeer density for each district (individuals/km^2). We used

the average density from 1980 to 2003 as a predictor for the 2003 vegetation data (following

Bråthen et al. 2007), and average density from 2003 to 2019 as a predictor for the 2020 vegetation

data, to not overlap with the previous interval. Results were not sensitive to the length of the

intervals (2003-2019 vs. 2000-2019).

Vegetation sampling. We used the point-intercept method (Bråthen and Hagberg 2004) with a

triangular 3-pin frame to obtain a measure of vascular plant abundance. We counted all hits of all

vascular species in 11 $0.08m^2$ plots spaced along each 50m transect. Prior to further analysis, we

converted point-frequency hits per species per plot to biomass estimates (g/m^2) using established

calibration equations (Tuomi et al. 2018; Ravolainen et al. 2010). For analyzing biomass data, we first

pooled species-specific biomass estimates based on functional grouping to forbs, graminoids,

deciduous woody shrubs and dwarf-shrubs (deciduous woody), and crowberry. Other groups not

included in the analysis (Supplementary Result Table 1A) were other evergreen woody dwarf-shrubs,

non-woody evergreen plants and vascular cryptogams. Other evergreen woody dwarf-shrubs were

not included in statistical analyses as responded very similarly to crowberry but have comparably

low biomass. We then averaged biomass of all 11 plots along each transect to reach a community-

averaged estimate of g/m^2 for forbs (non-zero sample size N_{year}: N_{2003} = 136, N_{2020} = 136), graminoids

(N_{2003} = 257, N_{2020} = 252), deciduous woody (N_{2003} = 282, N_{2020} = 279), and crowberry (N_{2003} = 261,

N_{2020} = 269). We also calculated the cover of the functional groups within each community, using

plot-level presence-absence data and summarized numbers of plots with each functional group

present in each transect.

Statistical analyses. To test our hypothesis, we fitted Bayesian linear multilevel gamma-hurdle

models with the package *brms* (Bürkner 2017) in the R statistical environment (R Core Team 2021)

(version 4.0.4/15.02.2021 and later). We decomposed the reindeer density to its spatial, temporal,

and residual components (Oedekoven et al. 2017), and standardized all three predictors to a mean

of 0 and variance of 1 for better effect comparability and model convergence. The temporal

component represents the change in average reindeer densities over two time points, the spatial

component the difference associated with the district reindeer densities averaged over time and the

residuals the district- and year specific variation in reindeer densities from the spatial and temporal

averages. To fully test the expectations in Fig. 1e, models for forbs and graminoids included a

quadratic term for the spatial component, while deciduous dwarf-shrubs and crowberry were

included linear term only, and all models included block and district as group-level intercepts. We

fitted the Bayesian generalized linear mixed models with weakly informative default priors (Bürkner

2017) and checked model convergence and independence of HMC chains based on the ^R statistics

(< 1.002) and effective sample size (> 2000) (Muth, Oravecz, and Gabry 2018). In addition, we fitted

negative-binomial hurdle models for plant functional group cover in the same way as described

above for biomass data. We used package default weakly informative priors for population and

group-level predictors and family parameters (Bürkner 2017). We confirmed model fit visually

through posterior predictive checks as well as comparing model-simulated data to observed data

(Muth, Oravecz, and Gabry 2018). Lack of spatial autocorrelation in group-level effects and model

residuals was assessed visually. Data visualization was done with packages ggplot2 (Wickham 2009)

ggdist (Kay 2021) and ggpmisc (Aphalo 2023).

Author contributions

K.A.B. conceived the idea, M.T., K.A.B. and N.Y. planned the re-sampling design. K.A.B. designed
conceptual figures with support from M.T., and based on discussions with all authors. M.T., K.A.B.,
T.Aa.U., N.Y. and S.Z. collected the data. M.T., K.A.B. and N.Y. planned data analysis. M.T. analyzed
the data and extracted the results. M.T. and K.A.B. led data interpretation, with all authors
contributing to interpreting the results. M.T. wrote the manuscript with and support from K.A.B. All
authors contributed substantially to editing the manuscript.

Acknowledgements

The work was funded by the Norwegian Research Council (FRIPRO project MONEC, code 302749 to
K.A.B.). We thank Nhat Minh Pham for comments on previous drafts of the manuscript, and Karoline
Helene Aares, Hanna Böhner, Lea Lipphardt, Hans Ivar Hortmann, Kinga Skalska, Katrine Skamfer
Hoset and Sindre Natvik for conducting field work, and the Norwegian Coast Guard, especially the
crew of KV Farm, for their hospitality and invaluable logistic help during the field work campaign. We
thank Audun Stien for discussions on the calf weight-density relationship. We also wish to thank
MONEC partners for discussions on implications of crowberry encroachment for their reindeer and
sheep herding SES. M.T. thanks the Mikkeli University Consortium (MUC) for providing working
facilities.

Competing interests

We disclose a familial link of one co-author, an associate professor, to a reindeer herder. Apart from
this, the authors declare no competing interests.

References

Aerts, Rien. 2010. "Nitrogen-Dependent Recovery of Subarctic Tundra Vegetation after
520   Simulation of Extreme Winter Warming Damage to Empetrum Hermaphroditum."
521   *Global Change Biology* 16 (3): 1071–81. https://doi.org/10.1111/j.1365-
522   2486.2009.01999.x.
523   Aphalo, Pedro J. 2023. "Ggpmisc: Miscellaneous Extensions to 'Ggplot2.'"
524   Barry, Tom, Dominique Berteaux, and Helga Bültmann, eds. 2013. *Arctic Biodiversity
525   Assessment: Status and Trends in Arcitc Biodiversity*. Akureyri, Iceland: The
526   Conservation of Arctic Flora and Fauna.
527   Bell, J. N. B., and J. H. Tallis. 1973. "Empetrum Nigrum L." *The Journal of Ecology* 61 (1): 289.
528   https://doi.org/10.2307/2258934.
529   Bjorkman, Anne D., Isla H. Myers-Smith, Sarah C. Elmendorf, Signe Normand, Nadja Rüger,
530   Pieter S. A. Beck, Anne Blach-Overgaard, et al. 2018. "Plant Functional Trait Change
531   across a Warming Tundra Biome." *Nature* 562 (7725): 57–62.
532   https://doi.org/10.1038/s41586-018-0563-7.
533   Bråthen, Kari Anne, Cathrine Helene Fodstad, and Christiane Gallet. 2010. "Ecosystem
534   Disturbance Reduces the Allelopathic Effects of Empetrum Hermaphroditum Humus
535   on Tundra Plants." *Journal of Vegetation Science* 21 (4): 786–95.
536   https://doi.org/10.1111/j.1654-1103.2010.01188.x.
537   Bråthen, Kari Anne, Victoria T. Gonzalez, and Nigel G. Yoccoz. 2018. "Gatekeepers to the
538   Effects of Climate Warming? Niche Construction Restricts Plant Community Changes
539   along a Temperature Gradient." *Perspectives in Plant Ecology, Evolution and
540   Systematics*, Special issue on Alpine and arctic plant communities: a worldwide
541   perspective, 30 (February): 71–81. https://doi.org/10.1016/j.ppees.2017.06.005.
542   Bråthen, Kari Anne, and Oskar Hagberg. 2004. "More Efficient Estimation of Plant Biomass."
543   *Journal of Vegetation Science* 15 (5): 653–60.
544   Bråthen, Kari Anne, Rolf A. Ims, Nigel G. Yoccoz, Per Fauchald, Torkild Tveraa, and Vera H.
545   Hausner. 2007. "Induced Shift in Ecosystem Productivity? Extensive Scale Effects of
546   Abundant Large Herbivores." *Ecosystems* 10 (5): 773–89.
547   https://doi.org/10.1007/s10021-007-9058-3.
548   Bråthen, Kari Anne, Francisco I Pugnaire, and Richard D Bardgett. 2021. "The Paradox of
549   Forbs in Grasslands and the Legacy of the Mammoth Steppe." *Frontiers in Ecology
550   and the Environment* 19 (10): 584–92. https://doi.org/10.1002/fee.2405.
551   Bråthen, Kari Anne, Virve Tuulia Ravolainen, Audun Stien, Torkild Tveraa, and Rolf A. Ims.
552   2017. "Rangifer Management Controls a Climate-Sensitive Tundra State Transition."
553   *Ecological Applications* 27 (8): 2416–27. https://doi.org/10.1002/eap.1618.
554   Bret-Harte, M Syndonia, GAIus R Shaver, Jennifer P Zoerner, Jill F Johnstone, Joanna L
555   Wagner, Andreas S Chavez, Ralph F Gunkelman Iv, Suzanne C Lippert, and James A
556   Laundre'. 2001. "Developmental Plasticity Allows Betula Nana to Dominate Tundra
557   Subjected to an Altered Environment," 16.
558   Bürkner, Paul-Christian. 2017. "Brms: An R Package for Bayesian Multilevel Models Using
559   Stan." *Journal of Statistical Software* 80 (1). https://doi.org/10.18637/jss.v080.i01.
560   Carey, Michael P., Beth L. Sanderson, Katie A. Barnas, and Julian D. Olden. 2012. "Native
561   Invaders – Challenges for Science, Management, Policy, and Society." *Frontiers in
562   Ecology and the Environment* 10 (7): 373–81. https://doi.org/10.1890/110060.

Christie, Katie S., John P. Bryant, Laura Gough, Virve T. Ravolainen, Roger W. Ruess, and Ken
D. Tape. 2015. "The Role of Vertebrate Herbivores in Regulating Shrub Expansion in
the Arctic: A Synthesis." *BioScience* 65 (12): 1123–33.
https://doi.org/10.1093/biosci/biv137.
Critchley, C. N. R, M. J. W Burke, and D. P Stevens. 2004. "Conservation of Lowland Semi-
Natural Grasslands in the UK: A Review of Botanical Monitoring Results from Agri-
Environment Schemes." *Biological Conservation* 115 (2): 263–78.
https://doi.org/10.1016/S0006-3207(03)00146-0.
Davis, Katrina J., Iadine Chadès, Jonathan R. Rhodes, and Michael Bode. 2019. "General
Rules for Environmental Management to Prioritise Social Ecological Systems
Research Based on a Value of Information Approach." *Journal of Applied Ecology* 56
(8): 2079–90. https://doi.org/10.1111/1365-2664.13425.
Dorrepaal, Ellen, Johannes H. C. Cornelissen, and Rien Aerts. 2007. "Changing Leaf Litter
Feedbacks on Plant Production across Contrasting Sub-Arctic Peatland Species and
Growth Forms." *Oecologia* 151 (2): 251–61. https://doi.org/10.1007/s00442-006-
0580-3.
Dudney, Joan, Richard J. Hobbs, Robert Heilmayr, John J. Battles, and Katharine N. Suding.
2018. "Navigating Novelty and Risk in Resilience Management." *Trends in Ecology &*
Evolution 33 (11): 863–73. https://doi.org/10.1016/j.tree.2018.08.012.
Elmendorf, Sarah C., Gregory H. R. Henry, Robert D. Hollister, Robert G. Björk, Noémie
Boulanger-Lapointe, Elisabeth J. Cooper, Johannes H. C. Cornelissen, et al. 2012.
"Plot-Scale Evidence of Tundra Vegetation Change and Links to Recent Summer
Warming." *Nature Climate Change* 2 (6): 453–57.
https://doi.org/10.1038/nclimate1465.
Fauchald, Per, Taejin Park, Hans Tømmervik, Ranga Myneni, and Vera Helene Hausner.
2017. "Arctic Greening from Warming Promotes Declines in Caribou Populations."
Science Advances 3 (4): e1601365. https://doi.org/10.1126/sciadv.1601365.
Gilbert, Benjamin, and Jonathan M. Levine. 2013. "Plant Invasions and Extinction Debts."
Proceedings of the National Academy of Sciences 110 (5): 1744–49.
https://doi.org/10.1073/pnas.1212375110.
González, Victoria T., Olavi Junttila, Bente Lindgård, Rigmor Reiersen, Kajetan Trost, and Kari
Anne Bråthen. 2015. "Batatasin-III and the Allelopathic Capacity of Empetrum
Nigrum." *Nordic Journal of Botany* 33 (2): 225–31.
https://doi.org/10.1111/njb.00559.
González, Victoria T., Bente Lindgård, Rigmor Reiersen, Snorre B. Hagen, and Kari Anne
Bråthen. 2021. "Niche Construction Mediates Climate Effects on Recovery of Tundra
Heathlands after Extreme Event." Edited by Craig Eliot Coleman. *PLOS ONE* 16 (2):
e0245929. https://doi.org/10.1371/journal.pone.0245929.
González, Victoria T., Mikel Moriana-Armendariz, Snorre B. Hagen, Bente Lindgård, Rigmor
Reiersen, and Kari Anne Bråthen. 2019. "High Resistance to Climatic Variability in a
Dominant Tundra Shrub Species." *PeerJ* 7 (June): e6967.
https://doi.org/10.7717/peerj.6967.
Graae, Bente J., Vigdis Vandvik, W. Scott Armbruster, Wolf L. Eiserhardt, Jens-Christian
Svenning, Kristoffer Hylander, Johan Ehrlén, et al. 2018. "Stay or Go – How
Topographic Complexity Influences Alpine Plant Population and Community
Responses to Climate Change." *Perspectives in Plant Ecology, Evolution and*

Systematics, Special issue on Alpine and arctic plant communities⸫: a worldwide
perspective, 30 (February): 41–50. https://doi.org/10.1016/j.ppees.2017.09.008.
Hausner, Vera H., Sigrid Engen, Camilla Brattland, and Per Fauchald. 2020. "Sámi Knowledge
and Ecosystem-Based Adaptation Strategies for Managing Pastures under Threat
from Multiple Land Uses." *Journal of Applied Ecology* 57 (9): 1656–65.
https://doi.org/10.1111/1365-2664.13559.
Henden, Ja, T Tveraa, A Stien, Jp Mellard, F Marolla, Ra Ims, and Ng Yoccoz. 2020. "Direct
and Indirect Effects of Environmental Drivers on Reindeer Reproduction." *Climate*
Research. https://doi.org/10.3354/cr01630.
Horstkotte, T., T. Aa Utsi, Å Larsson-Blind, P. Burgess, B. Johansen, J. Käyhkö, L. Oksanen,
and B. C. Forbes. 2017. "Human–Animal Agency in Reindeer Management: Sámi
Herders' Perspectives on Vegetation Dynamics under Climate Change." *Ecosphere* 8
(9): e01931. https://doi.org/10.1002/ecs2.1931.
Hoset, Katrine S., Lise Ruffino, Maria Tuomi, Tarja Oksanen, Lauri Oksanen, Aurelia
Mäkynen, Bernt Johansen, and Torunn Moe. 2017. "Changes in the Spatial
Configuration and Strength of Trophic Control Across a Productivity Gradient During
a Massive Rodent Outbreak." *Ecosystems*, March, 1–15.
https://doi.org/10.1007/s10021-017-0124-1.
Hughes, Terry P., Cristina Linares, Vasilis Dakos, Ingrid A. van de Leemput, and Egbert H. van
Nes. 2013. "Living Dangerously on Borrowed Time during Slow, Unrecognized
Regime Shifts." *Trends in Ecology & Evolution* 28 (3): 149–55.
https://doi.org/10.1016/j.tree.2012.08.022.
Ims, Rolf A, and Nigel G Yoccoz. 2017. "Ecosystem-Based Monitoring in the Age of Rapid
Climate Change and New Technologies." *Current Opinion in Environmental*
Sustainability 29 (December): 170–76. https://doi.org/10.1016/j.cosust.2018.01.003.
IPCC, ed. 2021. *IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of*
Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on
Climate Change. Cambridge University Press.
Iversen, Marianne, Per Fauchald, Knut Langeland, Rolf A. Ims, Nigel G. Yoccoz, and Kari Anne
Bråthen. 2014. "Phenology and Cover of Plant Growth Forms Predict Herbivore
Habitat Selection in a High Latitude Ecosystem." Edited by John F. Valentine. *PLoS*
ONE 9 (6): e100780. https://doi.org/10.1371/journal.pone.0100780.
James, Simon P. 2020. "Legal Rights and Nature's Contributions to People: Is There a
Connection?" *Biological Conservation* 241 (January): 108325.
https://doi.org/10.1016/j.biocon.2019.108325.
Jenkins, Deborah A., Nicolas Lecomte, Geoffrey Andrews, Glenn Yannic, and James A.
Schaefer. 2020. "Biotic Interactions Govern the Distribution of Coexisting Ungulates
in the Arctic Archipelago – A Case for Conservation Planning." *Global Ecology and*
Conservation 24 (December): e01239. https://doi.org/10.1016/j.gecco.2020.e01239.
Kaarlejärvi, E.a, K.S.b Hoset, and J.a Olofsson. 2015. "Mammalian Herbivores Confer
Resilience of Arctic Shrub-Dominated Ecosystems to Changing Climate." *Global*
Change Biology 21 (9): 3379–88. https://doi.org/10.1111/gcb.12970.
Kaarlejärvi, Elina, Anu Eskelinen, and Johan Olofsson. 2013. "Herbivory Prevents Positive
Responses of Lowland Plants to Warmer and More Fertile Conditions at High
Altitudes." *Functional Ecology* 27 (5): 1244–53. https://doi.org/10.1111/1365-
2435.12113.

Kay, Matthew. 2021. *Ggdist: Visualizations of Distributions and Uncertainty*.
https://doi.org/10.5281/zenodo.3879620.
Keech, Olivier, Christopher Carcaillet, and Marie-Charlotte Nilsson. 2005. "Adsorption of
Allelopathic Compounds by Wood-Derived Charcoal: The Role of Wood Porosity."
Plant and Soil 272 (1): 291–300. https://doi.org/10.1007/s11104-004-5485-5.
Kivinen, Sonja, Sirpa Rasmus, Kirsti Jylhä, and Mikko Laapas. 2017. "Long-Term Climate
Trends and Extreme Events in Northern Fennoscandia (1914–2013)." *Climate* 5 (1):
16. https://doi.org/10.3390/cli5010016.
Krebs, Charles J. 2002. "Two Complementary Paradigms for Analysing Population
Dynamics." *Philosophical Transactions of the Royal Society of London. Series B:
Biological Sciences* 357 (1425): 1211–19. https://doi.org/10.1098/rstb.2002.1122.
Kuussaari, Mikko, Riccardo Bommarco, Risto K. Heikkinen, Aveliina Helm, Jochen Krauss,
Regina Lindborg, Erik Öckinger, et al. 2009. "Extinction Debt: A Challenge for
Biodiversity Conservation." *Trends in Ecology & Evolution* 24 (10): 564–71.
https://doi.org/10.1016/j.tree.2009.04.011.
Latombe, G., S. Canavan, H. Hirsch, C. Hui, S. Kumschick, M. M. Nsikani, L. J. Potgieter, et al.
2019. "A Four-Component Classification of Uncertainties in Biological Invasions:
Implications for Management." *Ecosphere* 10 (4): e02669.
https://doi.org/10.1002/ecs2.2669.
Lindenmayer, David B., and Gene E. Likens. 2009. "Adaptive Monitoring: A New Paradigm
for Long-Term Research and Monitoring." *Trends in Ecology & Evolution* 24 (9): 482–
86. https://doi.org/10.1016/j.tree.2009.03.005.
Madden, Francine, and Brian McQuinn. 2014. "Conservation's Blind Spot: The Case for
Conflict Transformation in Wildlife Conservation." *Biological Conservation* 178
(October): 97–106. https://doi.org/10.1016/j.biocon.2014.07.015.
Maliniemi, Tuija, Jutta Kapfer, Patrick Saccone, Anu Skog, and Risto Virtanen. 2018. "Long-
Term Vegetation Changes of Treeless Heath Communities in Northern Fennoscandia:
Links to Climate Change Trends and Reindeer Grazing." *Journal of Vegetation Science*
29 (3): 469–79. https://doi.org/10.1111/jvs.12630.
Mallory, Conor D., and Mark S. Boyce. 2017. "Observed and Predicted Effects of Climate
Change on Arctic Caribou and Reindeer." *Environmental Reviews*, June.
https://doi.org/10.1139/er-2017-0032.
Marteinsdóttir, B, E F Þórarinsdóttir, G Halldórsson, J H Stefánsson, J Þórsson, K
Svavarsdóttir, R Finnsdóttir, and S Jónsdóttir. 2021. "GróLind – Sustainable Land Use
Based on Ecological Knowledge." *International Grassland Congress Proceedings*, 6.
Milner-Gulland, E. J., and Katriona Shea. 2017. "Embracing Uncertainty in Applied Ecology."
Journal of Applied Ecology 54 (6): 2063–68. https://doi.org/10.1111/1365-
2664.12887.
Ministry of Agriculture. 2017. "White Paper 32 (2016–2017). Reindrift— Lang Tradisjon –
Unike Muligheter." 32. https://www.regjeringen.no/no/dokumenter/meld.-st.-32-
20162017/id2547907/?ch=1.
———. n.d. "Ressursregnskapet for Reindriftsnæringen."
https://www.landbruksdirektoratet.no/nb/nyhetsrom/rapporter/ressursregnskapet-
for-reindriftsnaeringen.
Mod, Heidi K., and Miska Luoto. 2016. "Arctic Shrubification Mediates the Impacts of
Warming Climate on Changes to Tundra Vegetation." *Environmental Research Letters*
11 (12): 124028. https://doi.org/10.1088/1748-9326/11/12/124028.

Muggeo, Vito M.R. 2017. "Interval Estimation for the Breakpoint in Segmented Regression:
A Smoothed Score-Based Approach." *Australian & New Zealand Journal of Statistics*
59: 311–22.
Muth, Chelsea, Zita Oravecz, and Jonah Gabry. 2018. "User-Friendly Bayesian Regression
Modeling: A Tutorial with Rstanarm and Shinystan." *The Quantitative Methods for*
Psychology 14 (2): 99–119. https://doi.org/10.20982/tqmp.14.2.p099.
Myers-Smith, I.H.a b, B.C.c Forbes, M.d Wilmking, M.d Hallinger, T.e Lantz, D.f Blok, K.D.g
Tape, et al. 2011. "Shrub Expansion in Tundra Ecosystems: Dynamics, Impacts and
Research Priorities." *Environmental Research Letters* 6 (4).
https://doi.org/10.1088/1748-9326/6/4/045509.
Myers-Smith, Isla H., Meagan M. Grabowski, Haydn J. D. Thomas, Sandra Angers-Blondin,
Gergana N. Daskalova, Anne D. Bjorkman, Andrew M. Cunliffe, et al. 2019. "Eighteen
Years of Ecological Monitoring Reveals Multiple Lines of Evidence for Tundra
Vegetation Change." *Ecological Monographs* 89 (2): e01351.
https://doi.org/10.1002/ecm.1351.
Myers-Smith, Isla H., Jeffrey T. Kerby, Gareth K. Phoenix, Jarle W. Bjerke, Howard E. Epstein,
Jakob J. Assmann, Christian John, et al. 2020. "Complexity Revealed in the Greening
of the Arctic." *Nature Climate Change* 10 (2): 106–17.
https://doi.org/10.1038/s41558-019-0688-1.
Nilsson, Marie-Charlotte, and David A. Wardle. 2005. "Understory Vegetation as a Forest
Ecosystem Driver: Evidence from the Northern Swedish Boreal Forest." *Frontiers in*
Ecology and the Environment 3 (8): 421–28. https://doi.org/10.1890/1540-
9295(2005)003[0421:UVAAFE]2.0.CO;2.
Oksanen, Lauri, and Tarja Oksanen. 2000. "The Logic and Realism of the Hypothesis of
Exploitation Ecosystems." *The American Naturalist* 155 (6): 703–23.
Oksanen, Tarja, Lauri Oksanen, Katariina E. M. Vuorinen, Christopher Wolf, Aurelia
Mäkynen, Johan Olofsson, William J. Ripple, Risto Virtanen, and Tove Aa. Utsi. 2020.
"The Impact of Thermal Seasonality on Terrestrial Endotherm Food Web Dynamics: A
Revision of the Exploitation Ecosystem Hypothesis." *Ecography* 43 (12): 1859–77.
https://doi.org/10.1111/ecog.05076.
Olofsson, Johan, Hans Tømmervik, and Terry V. Callaghan. 2012. "Vole and Lemming Activity
Observed from Space." *Nature Climate Change* 2 (12): 880–83.
https://doi.org/10.1038/nclimate1537.
Parker, Katherine L., Perry S. Barboza, and Michael P. Gillingham. 2009. "Nutrition Integrates
Environmental Responses of Ungulates." *Functional Ecology* 23 (1): 57–69.
https://doi.org/10.1111/j.1365-2435.2009.01528.x.
Parker, Thomas C., Alana M. Thurston, Katrine Raundrup, Jens-Arne Subke, Philip A.
Wookey, and Iain P. Hartley. 2021. "Shrub Expansion in the Arctic May Induce
Large-scale Carbon Losses Due to Changes in Plant-soil Interactions." *Plant and Soil*
463 (1): 643–51. https://doi.org/10.1007/s11104-021-04919-8.
Pedersen, Åshild Ønvik. 2021. *Norwegian Arctic Tundra: A Panel-Based Assessment of*
Ecosystem Condition. Norsk Polarinstitutt. https://brage.npolar.no/npolar-
xmlui/handle/11250/2754696.
Pilsbacher, Anna Katharina, Bente Lindgård, Rigmor Reiersen, Victoria T. González, and Kari
Anne Bråthen. 2021. "Interfering with Neighbouring Communities: Allelopathy
Astray in the Tundra Delays Seedling Development." *Functional Ecology* 35 (1): 266–
76. https://doi.org/10.1111/1365-2435.13694.

Pivello, Vânia Regina, Marcus Vinicius Vieira, Maria Tereza Grombone-Guaratini, and Dalva
Maria Silva Matos. 2018. "Thinking about Super-Dominant Populations of Native
Species – Examples from Brazil." *Perspectives in Ecology and Conservation* 16 (2): 74–
82. https://doi.org/10.1016/j.pecon.2018.04.001.
Post, Eric, Richard B. Alley, Torben R. Christensen, Marc Macias-Fauria, Bruce C. Forbes,
Michael N. Gooseff, Amy Iler, et al. 2019. "The Polar Regions in a 2°C Warmer
World." *Science Advances* 5 (12): eaaw9883.
https://doi.org/10.1126/sciadv.aaw9883.
Post, Eric, Mads C. Forchhammer, M. Syndonia Bret-Harte, Terry V. Callaghan, Torben R.
Christensen, Bo Elberling, Anthony D. Fox, et al. 2009. "Ecological Dynamics Across
the Arctic Associated with Recent Climate Change." *Science* 325 (5946): 1355–58.
https://doi.org/10.1126/science.1173113.
Preece, C., and G. K. Phoenix. 2014. "Impact of Early and Late Winter Icing Events on Sub-
Arctic Dwarf Shrubs." *Plant Biology* 16 (1): 125–32.
https://doi.org/10.1111/plb.12015.
R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing. https://www.R-project.org/.
Ravolainen, V. T., N. G. Yoccoz, K. A. Bråthen, R. A. Ims, M. Iversen, and V. T. González. 2010.
"Additive Partitioning of Diversity Reveals No Scale-Dependent Impacts of Large
Ungulates on the Structure of Tundra Plant Communities." *Ecosystems* 13 (1): 157–
70. https://doi.org/10.1007/s10021-009-9308-7.
Regan, Helen M., Mark Colyvan, and Mark A. Burgman. 2002. "A Taxonomy and Treatment
of Uncertainty for Ecology and Conservation Biology." *Ecological Applications* 12 (2):
618–28. https://doi.org/10.1890/1051-0761(2002)012[0618:ATATOU]2.0.CO;2.
Reindeer Husbandry Act. 2007. lovdata.no/dokument/NL/lov/2007-06-15-40.
Roux, Peter C. le, Jonathan Lenoir, Loïc Pellissier, Mary S. Wisz, and Miska Luoto. 2013.
"Horizontal, but Not Vertical, Biotic Interactions Affect Fine-Scale Plant Distribution
Patterns in a Low-Energy System." *Ecology* 94 (3): 671–82.
Ruckelshaus, Mary H., Stephen T. Jackson, Harold A. Mooney, Katharine L. Jacobs, Karim-Aly
S. Kassam, Mary T. K. Arroyo, András Báldi, et al. 2020. "The IPBES Global
Assessment: Pathways to Action." *Trends in Ecology & Evolution* 35 (5): 407–14.
https://doi.org/10.1016/j.tree.2020.01.009.
Schmidt, Niels M., Jesper B. Mosbacher, Eero J. Vesterinen, Tomas Roslin, and Anders
Michelsen. 2018. "Limited Dietary Overlap amongst Resident Arctic Herbivores in
Winter: Complementary Insights from Complementary Methods." *Oecologia* 187 (3):
689–99. https://doi.org/10.1007/s00442-018-4147-x.
Schneider, Volker, Philip Leifeld, and Thomas Malang. 2013. "Coping with Creeping
Catastrophes: National Political Systems and the Challenge of Slow-Moving Policy
Problems." In *Long-Term Governance for Social-Ecological Change*, edited by Bernd
Siebenhüner, Marlen Arnold, Klaus Eisenack, Klaus Jacob, and Michael Pregernig,
221–38. Environmental Politics / Routledge Research in Environmental Politics, v. 21.
New York: Routledge.
Shackelford, Nancy, Michael Renton, Michael P. Perring, and Richard J. Hobbs. 2013.
"Modeling Disturbance-Based Native Invasive Species Control and Its Implications for
Management." *Ecological Applications* 23 (6): 1331–44. https://doi.org/10.1890/12-
1881.1.

Shackleton, Ross T., Charlie M. Shackleton, and Christian A. Kull. 2019. "The Role of Invasive
796   Alien Species in Shaping Local Livelihoods and Human Well-Being: A Review." *Journal*
797   *of Environmental Management*, The human and social dimensions of invasion
798   science and management, 229 (January): 145–57.
799   https://doi.org/10.1016/j.jenvman.2018.05.007.
800   Sinclair, A. R. E., and Charles J. Krebs. 2002. "Complex Numerical Responses to Top–down
801   and Bottom–up Processes in Vertebrate Populations." Edited by R. M. Sibly, J. Hone,
802   and T. H. Clutton–Brock. *Philosophical Transactions of the Royal Society of London.*
803   *Series B: Biological Sciences* 357 (1425): 1221–31.
804   https://doi.org/10.1098/rstb.2002.1123.
805   Skarin, Anna, Mariana Verdonen, Timo Kumpula, Marc Macias-Fauria, Moudud Alam, Jeffrey
806   Kerby, and Bruce C Forbes. 2020. "Reindeer Use of Low Arctic Tundra Correlates with
807   Landscape Structure." *Environmental Research Letters* 15 (11): 115012.
808   https://doi.org/10.1088/1748-9326/abbf15.
809   Stewart, Lærke, Caroline E. Simonsen, Jens-Christian Svenning, Niels Martin Schmidt, and
810   Loïc Pellissier. 2018. "Forecasted Homogenization of High Arctic Vegetation
811   Communities under Climate Change." *Journal of Biogeography* 45 (11): 2576–87.
812   https://doi.org/10.1111/jbi.13434.
813   Stien, Audun, Torkild Tveraa, Rolf Anker Ims, Jennifer Stien, and Nigel Gilles Yoccoz. 2021.
814   "Unfounded Claims about Productivity beyond Density for Reindeer Pastoralism
815   Systems." *Pastoralism* 11 (1): 20. https://doi.org/10.1186/s13570-021-00209-1.
816   Taylor, Jason J., James P. Lawler, Mora Aronsson, Tom Barry, Anne D. Bjorkman, Tom
817   Christensen, Stephen J. Coulson, et al. 2020. "Arctic Terrestrial Biodiversity Status
818   and Trends: A Synopsis of Science Supporting the CBMP State of Arctic Terrestrial
819   Biodiversity Report." *Ambio* 49 (3): 833–47. https://doi.org/10.1007/s13280-019-
820   01303-w.
821   Tuomi, Maria, Sari Stark, Katrine S. Hoset, Maria Väisänen, Lauri Oksanen, Francisco J. A.
822   Murguzur, Hanna Tuomisto, Jonas Dahlgren, and Kari Anne Bråthen. 2018.
823   "Herbivore Effects on Ecosystem Process Rates in a Low-Productive System."
824   *Ecosystems* 22 (4): 827–43. https://doi.org/10.1007/s10021-018-0307-4.
825   Tveraa, Torkild, Per Fauchald, Nigel Gilles Yoccoz, Rolf Anker Ims, Ronny Aanes, and Kjell
826   Arild Høgda. 2007. "What Regulate and Limit Reindeer Populations in Norway?"
827   *Oikos* 116 (4): 706–15.
828   Tveraa, Torkild, Audun Stien, Bård-J. Bårdsen, and Per Fauchald. 2013. "Population
829   Densities, Vegetation Green-Up, and Plant Productivity: Impacts on Reproductive
830   Success and Juvenile Body Mass in Reindeer." *PLOS ONE* 8 (2): e56450.
831   https://doi.org/10.1371/journal.pone.0056450.
832   Tybirk, Knud, Marie-Charlotte Nilsson, Anders Michelsen, Hanne Lakkenborg Kristensen,
833   Anna Shevtsova, Morten Tune Strandberg, Marianne Johansson, et al. 2000. "Nordic
834   Empetrum Dominated Ecosystems: Function and Susceptibility to Environmental
835   Changes." *AMBIO: A Journal of the Human Environment* 29 (2): 90–97.
836   https://doi.org/10.1579/0044-7447-29.2.90.
837   Vaas, J., P. P. J. Driessen, M. Giezen, F. van Laerhoven, and M. J. Wassen. 2021. "Moving
838   from Latent to Manifest Problem: Trajectories Across Scientific and Public Salience of
839   Invasive Alien Species." *Environmental Management* 67 (5): 901–19.
840   https://doi.org/10.1007/s00267-020-01404-3.

841   Valéry, Loïc, Hervé Fritz, and Jean-Claude Lefeuvre. 2013. "Another Call for the End of
842   Invasion Biology." *Oikos* 122 (8): 1143–46. https://doi.org/10.1111/j.1600-
843   0706.2013.00445.x.
844   Valéry, Loïc, Hervé Fritz, Jean-Claude Lefeuvre, and Daniel Simberloff. 2009. "Invasive
845   Species Can Also Be Native…." *Trends in Ecology & Evolution* 24 (11): 585.
846   https://doi.org/10.1016/j.tree.2009.07.003.
847   Van Wijk, M. T., K. E. Clemmensen, G. R. Shaver, M. Williams, T. V. Callaghan, F. S. Chapin III,
848   J. H. C. Cornelissen, et al. 2004. "Long-Term Ecosystem Level Experiments at Toolik
849   Lake, Alaska, and at Abisko, Northern Sweden: Generalizations and Differences in
850   Ecosystem and Plant Type Responses to Global Change." *Global Change Biology* 10
851   (1): 105–23. https://doi.org/10.1111/j.1365-2486.2003.00719.x.
852   Vandvik, Vigdis, Einar Heegaard, Inger Elisabeth Måren, and Per Arild Aarrestad. 2005.
853   "Managing Heterogeneity: The Importance of Grazing and Environmental Variation
854   on Post-Fire Succession in Heathlands." *Journal of Applied Ecology* 42 (1): 139–49.
855   https://doi.org/10.1111/j.1365-2664.2005.00982.x.
856   Vowles, Tage, and Robert G. Björk. 2019. "Implications of Evergreen Shrub Expansion in the
857   Arctic." *Journal of Ecology* 107 (2): 650–55. https://doi.org/10.1111/1365-
858   2745.13081.
859   Vowles, Tage, Bengt Gunnarsson, Ulf Molau, Thomas Hickler, Leif Klemedtsson, and Robert
860   G. Björk. 2017. "Expansion of Deciduous Tall Shrubs but Not Evergreen Dwarf Shrubs
861   Inhibited by Reindeer in Scandes Mountain Range." *Journal of Ecology* 105 (6): 1547–
862   61. https://doi.org/10.1111/1365-2745.12753.
863   Vuorinen, Katariina E. M., Lauri Oksanen, Tarja Oksanen, Anni Pyykönen, Johan Olofsson,
864   and Risto Virtanen. 2017. "Open Tundra Persist, but Arctic Features Decline—
865   Vegetation Changes in the Warming Fennoscandian Tundra." *Global Change Biology*
866   23 (9): 3794–3807. https://doi.org/10.1111/gcb.13710.
867   Wang, Hao, Huiying Liu, Ni Huang, Jian Bi, Xuanlong Ma, Zhiyuan Ma, Zijian Shangguan, et al.
868   2021. "Satellite-Derived NDVI Underestimates the Advancement of Alpine
869   Vegetation Growth over the Past Three Decades." *Ecology* 102 (12): e03518.
870   https://doi.org/10.1002/ecy.3518.
871   Wardle, David A., Marie-Charlotte Nilsson, Christiane Gallet, and Olle Zackrisson. 1998. "An
872   Ecosystem-Level Perspective of Allelopathy." *Biological Reviews* 73 (3): 305–19.
873   Webb, N.R. 2008. "The Traditional Management of European Heathlands." *Journal of
874   Applied Ecology* 35 (6): 987–90. https://doi.org/10.1111/j.1365-
875   2664.1998.tb00020.x.
876   Webber, Quinn M.R., Kristy M. Ferraro, Jack G. Hendrix, and Eric Vander Wal. 2022. "What
877   Do Caribou Eat? A Review of the Literature on Caribou Diet." *Canadian Journal of
878   Zoology* 100 (3): 197–207. https://doi.org/10.1139/cjz-2021-0162.
879   Weijers, Stef, Roland Pape, Jörg Löffler, and Isla H. Myers-Smith. 2018. "Contrasting Shrub
880   Species Respond to Early Summer Temperatures Leading to Correspondence of
881   Shrub Growth Patterns." *Environmental Research Letters* 13 (3): 034005.
882   https://doi.org/10.1088/1748-9326/aaa5b8.
883   Wickham, Hadley. 2009. *Ggplot2: Elegant Graphics for Data Analysis*. Springer Science &
884   Business Media.
885   Wilson, Scott D., and Christer Nilsson. 2009. "Arctic Alpine Vegetation Change over 20
886   Years." *Global Change Biology* 15 (7): 1676–84. https://doi.org/10.1111/j.1365-
887   2486.2009.01896.x.

888
889

Tables

Table 1. Model parameter estimates and credible intervals, with plant functional group biomass as response and decomposed reindeer density as predictors. Note that the temporal coefficient links with decrease in reindeer density over time, and hence a negative value indicates an increase over time. For estimates of group-level standard deviation (random effects), see Supplementary Results Table 3. Error terms Q2.5 and Q97.5 represent the 95% credible interval. Bold font indicates strong support for the effect, and italic indicates relatively strong support for the effect, given the data.

Response	Parameter	Estimate	Q2.5	Q97.5
Biomass ~ spatial + spatial^2 + temporal + residual				
Forbs	Intercept	1.777	1.394	2.168
	Spatial	*-0.424*	*-0.831*	*-0.006*
	Spatial^2	0.205	-0.071	0.475
	Temporal	0.041	-0.093	0.174
	Residual	-0.069	-0.161	0.020
Graminoids	Intercept	3.132	2.799	3.471
	Spatial	0.012	-0.378	0.393
	Spatial^2	0.037	-0.213	0.280
	Temporal	0.041	-0.072	0.152
	Residual	-0.013	-0.088	0.062
Biomass ~ spatial + temporal + residual				
Deciduous woody	Intercept	4.491	4.251	4.791
	Spatial	0.028	-0.224	0.289
	Temporal	**-0.220**	**-0.302**	**-0.137**
	Residual	-0.028	-0.082	0.027
Crowberry	Intercept	5.229	5.079	5.391
	Spatial	0.081	-0.092	0.256
	Temporal	**-0.340**	**-0.420**	**-0.261**
	Residual	0.001	-0.052	0.054

Figures and legends

Figure 1. Role of pastures in the reindeer management system. The current management decision-

making process of reindeer in Norway applies reindeer density as the regulatory mechanism of

sustainable pastures (*Reindeer Husbandry Act* 2007). Autumn slaughter weight of animals, as a proxy

for body condition, is used as the indicator for assessing sustainability of the summer pastures (state

of the system), and no measures pertaining to the pastures themselves. **a)** The study area includes

20 management districts, spanning two latitudinal and six longitudinal degrees and representing

different ecological contexts. Within each district we analysed transects for plant quantity, quality

and diversity in 2003 and in 2020. **b)** Reindeer interact with their pasture through top-down and

bottom-up effects. Approximate pasture functional group composition is visualized based on 2003

data. **c)** The management decision process applies animal weight as an indicator of pasture

condition, here expressed as density-dependence of calf body mass in 2000-2019 within and across

the management districts of the current study (Ministry of Agriculture, n.d.). **d)** In 2003 across the

studied summer pastures, the average pasture consisted of plant functional groups of varying

species richness (Bråthen et al 2007), nitrogen content (Murguzur et al 2019) and biomass (Bråthen

et al 2007). Forbs, the most nutritious and species-rich group, made up the least biomass in the

pastures. Conversely, the evergreen dwarf-shrubs, the least nutritious and species-poor group, made

up the most biomass and with crowberry (red arrows) making up most of this biomass but not of the

nitrogen pool. **e)** The role of plant biomass in linking reindeer densities and weights is indirect

(*Reindeer Husbandry Act* 2007; Ministry of Agriculture 2017). The decision-making assumes a

negative top-down relationship between animal density and plant biomass (left panel), and a

positive bottom-up relationship between plant biomass and weights (right panel) (see Tveraa et al.

2013). Here, we inspect these assumptions further for plant functional groups. Left panel: Forbs and

grasses, as more grazing tolerant groups, are expected to have a unimodal response to animal

density (Wardle and Bardgett 2004). Deciduous shrubs are expected to show patterns of a browsing

trap at densities above 5 animals/km^2 (Bråthen et al 2017), whereas evergreen shrubs are expected

to be little affected from browsing (Iversen et al. 2014) but may decline under trampling effects

(Tybirk et al. 2000). Right panel: Per biomass unit (right panel), we assume the most nutritious forbs

to have the highest contribution to animal weights followed by graminoids, deciduous and

evergreen shrubs in declining order of nutrient content respectively. Framed graphs in panels D and

E are patterns and relationships that are empirically addressed here.

Figure 2. Growing degree days (GDD), as annual sums of daily temperatures above 5°C during May-

October 1957-2019, averaged across the studied reindeer summer pastures (Pedersen et al.

2020/met.no). Solid thick line and shaded areas represent mean ± 95% confidence intervals from

segmented linear regression with a breakpoint at year 1977. Parameter estimates are provided at

the top of the figure.

Figure 3. Modeled temporal and spatial variation in plant group biomass (forbs, graminoids,
deciduous shrubs and dwarf-shrubs and crowberry). **A)** Estimates of posterior mean distributions of
biomass of forbs, graminoids, deciduous woody plants and crowberry, separately for each plant
group in 2003 (grey) and in 2020 (greens). Means and 95% confidence intervals from observed data
for each year are provided as point intervals below each biomass density plot (note full overlap for
forbs and graminoids). **B)** Association between plant group biomass (mean and model-derived 95%
and 80% credible intervals of the estimated posterior mean distribution) and the district-averaged
reindeer density. Each point represents a reindeer district. **C)** Model-derived estimates of the
relationship of plant biomass and the (recomposed) reindeer density. Each short solid black line
connects averaged model estimates in 2003 and 2020 in each district, and a regression line is fitted
with log(density) across all districts. In panels B and C, the the coloured regression lines and adjusted
R^2 across all districts are estimated with the package *ggpmisc* functions stat_poly_line and
stat_poly_eq (Aphalo 2023).

Figure 4. Mechanisms and consequences of crowberry proliferation. A) Mechanisms of crowberry

proliferation and increased allelopathy over time. Already an abundant and long-lived species, slow

growth may still cause substantial increments in abundance. Growth of established plants results in

infilling and increased biomass, leading to increasing allelopathic litter effects in the soil. In addition,

dispersal via seeds and clonal reproduction are means of lateral encroachment, increasing the areal

extent of the plant. **B)** Potential transition in pasture state and its implication for the state-indicator

relationship. Left panel: Increasing crowberry abundance may push the system towards a state of

strong allelopathy. The state may be highly resilient, with litter and seed bank -mediated legacy

effects that over time will reduce new plant establishment (Gonzalez et al 2021) at levels as low as

25% crowberry out of total community standing biomass (Bråthen et al 2018). Right panel:

Conceptual model of effects of evergreening and crowberry proliferation on reindeer weights.

Relationship between pasture biomass and calf weights is dependent on plant composition, and

unlikely to be invariable over time. Changing bottom-up effects and transition to increasing

allelopathy in pastures may have adverse, yet slow impacts on reindeer condition.

997