Multiscale structural anisotropy in pulvinus motor organs of the sensitive plant *Mimosa pudica*

David A. Sleboda[1,2,*], Anja Geitmann[2], and Reza Sharif-Naeini[1]

[1]Department of Physiology, McGill University

[2]Department of Plant Science, McGill University

*Corresponding Author

Present Address: David A. Sleboda, Department of Ecology and Evolutionary Biology, University of California Irvine

Abstract

Pulvini are joint-like motor organs that power active leaf movement in many plants.

Multiple structural specializations spanning subcellular, cellular, and tissue scales of pulvinus

organization have been described; however, the impacts of multiscale mechanics on pulvinus

physiology remain poorly understood. To investigate the influence of multiscale morphology on

turgor-induced deformation, we visualized *Mimosa pudica* pulvinus morphology at multiple

hierarchical scales of organization and used osmotic perturbations to experimentally swell

pulvini in incremental states of dissection. We observed directional cellulose microfibril

reinforcement, oblong, spindle-shaped primary pit fields, and flattened, disk-like cell geometries

in the parenchyma of *M. pudica*. Consistent with these observations, isolated parenchyma

tissues displayed highly anisotropic swelling behaviors indicating a high degree of mechanical

anisotropy. Swelling behaviors at higher scales of pulvinus organization were also influenced by

the presence of the pulvinus epidermis, which displayed oblong epidermal cells oriented

transverse to the pulvinus long axis. Our findings indicate that structural specializations spread

across multiple hierarchical scales of organization guide hydraulic deformation of pulvini,

suggesting that multiscale mechanics are crucial to the translation of cell-level turgor variations

into organ-scale pulvinus motion *in vivo*.

Introduction

Pulvini are fleshy swellings located at the bases of petioles, leaves, and leaflets in many

plants. They are sites of growth-independent motion and facilitate active, reversible bending of

leafy plant appendages. Anatomically, pulvini are roughly cylindrical in shape and consist of a

flexible core of vasculature surrounded by a thick cortex of parenchyma cells. During active

deformation, bulk migration of water occurs between extensor and flexor parenchyma cell

groups located on opposing sides of the central vasculature core (Tamiya et al., 1988), resulting

in a length differential between the two groups that induces long-axis pulvinus bending.

Pulvinus-driven leaf folding can occur gradually over the course of hours under the influence of

a circadian rhythm, or rapidly as a thigmonastic response to touch, shaking, or wounding of a

plant. Active pulvinus bending is associated with efflux of K^+ and Cl^- ions from parenchyma cells

(Kumon and Suda, 1984), offloading of sugar from vascular elements (Fromm and Eschrich,

1988a), and consumption of ATP (Fromm and Eschrich, 1988b), indicating that pulvinus motion

is associated with active transport of ions and other osmolites to which water is attracted

(Scorza and Dornelas, 2011).

While multiple theoretical models of pulvinus physiology provide insight into the

biochemical mechanisms that power active movement of fluid between parenchyma cell groups

(e.g. Kagawa and Saito, 2000; Morillon et al., 2001; Kwan et al., 2013; Wang and Li, 2020),

relatively little is known regarding the mechanisms by which volumetric changes at the cellular

level are translated into physiologically useful motion at the organ scale. Multiple morphological

features have been identified which have the potential to bridge this gap. In pulvini of the scarlet

runner bean *Phaseolus coccineus L.*, observations with polarized light microscopy suggest that

the walls of parenchyma cells are reinforced by cellulose microfibrils oriented primarily

perpendicular to the long axis of the pulvinus, confining turgor-induced cell deformation to the

long axis of the organ where its effect on bending is greatest (Mayer et al., 1985). In pulvini of

Phaseolus coccineus L. and of the common bean *Phaseolus vulgaris*, parenchyma cells are

flattened in the direction parallel to the long axis of the pulvinus, displaying approximately disc-

shaped morphologies that are also conducive to deformation preferentially along the long axis of

the organ (Mayer et al., 1985; Koller and Zamski, 2002). In pulvini of *Albizia julibrissin*,

Phaseolus vulgaris, and the sensitive plant *Mimosa pudica,* wrinkles in the pulvinus surface run

preferentially transverse to the organ long axis and have been identified as structural features

that facilitate longitudinal deformation of the pulvinus epidermis without sacrificing

circumferential stiffness (Satter et al., 1970; Koller and Zamski, 2002; Song et al., 2014; Mano

and Hasebe, 2021). These structural features have the potential to physically guide hydraulic

deformation of pulvini, ensuring that volumetric changes at the cellular level are ultimately

translated into long-axis tissue deformations that induce macroscopic organ bending. Despite

the functional importance of this role, a multiscale study surveying structural anisotropy across

scales of pulvinus organization within a single plant species has not been carried out, and the

distinct contributions of cell wall ultrastructure, cell geometry, and epidermis morphology to the

management of hydraulic deformation remain unclear.

Here we disentangle the influences of cell wall ultrastructure, cell geometry, and

epidermis morphology on turgor-induced deformation in the pulvini of *Mimosa pudica*, a touch-

sensitive plant which displays both rapid thigmonastic and slow circadian leaf movements.

Using a combination of scanning electron and confocal microscopy, we visualized *M. pudica*

pulvinus morphology at multiple hierarchical scales of organization and characterized the

degree of structural anisotropy present at each. Additionally, we used osmotic bathing solutions

to experimentally swell live, isolated *M. pudica* pulvinus tissues in various states of dissection.

This approach allowed us to characterize the degree of anisotropic tissue deformation that

results from purely isotropic loads of intracellular fluid pressure. Differences in swelling

behaviors across scales of organization are discussed in relation to the anisotropic structural

features observed.

Materials and Methods

All pulvinus organs and tissue samples were harvested fresh from *Mimosa pudica* plants

grown under laboratory conditions. Plants were grown from seed in a growth chamber with a 12-

hour light-dark cycle at 21-23°C and 60% humidity. Plants were watered approximately once

every 3 days and fertilized monthly.

Epidermis morphology

The surface morphology of the pulvinus epidermis was visualized using a Hitachi TM-

1000 scanning electron microscope (SEM). Primary, secondary, and tertiary pulvini located at

the bases of petioles, leaves, and leaflets, respectively, were removed from plants and

dehydrated in a graded series of ethanol. Isolated pulvini were then critical point dried in liquid

CO_2 and mounted in lateral view on aluminum stubs with carbon double-sided tape. Mounted

organs were sputter coated with a 4 nm layer of gold and palladium and imaged at an excitation

voltage of 15 kV. Aspect ratios and orientations of epidermal cells were measured manually

from scanning electron micrographs of primary pulvini using ImageJ (Schindelin et al., 2012).

The positions of epidermal cells along the lengths of pulvini were defined relative to their

distance from the pulvinus base, measured along a path traversing the central axis of pulvinus

and petiole (see yellow line depicted in fig. 2a). Total path length examined was determined by

doubling the distance from the base of the pulvinus to the center of the knuckle-like

protuberance immediately distal to the pulvinus (indicated as length 0.50 in fig. 2a), such that

roughly equal lengths of pulvinus and petiole were examined. To facilitate comparison of pulvini

of varying size, distances of epidermal cells from the pulvinus base were expressed as portions

of the total path length examined for each organ. Epidermal cell orientation was measured

relative to the orientation of the central path.

Cell wall ultrastructure

The ultrastructures of individual parenchymal cell walls were visualized using an

environmental scanning electron microscope (FEI Quanta 450 FE-ESEM, FEI Co.) at the McGill

University Facility for Electron Microscopy Research. In order to observe the inner structures of

individual cell walls, primary pulvini were decellularized using an adapted protocol for the

generation of acellular, cellulosic tissue engineering scaffolds from plant material (Adamski et

al., 2018). Fresh pulvini were fixed overnight in 4% formaldehyde in 0.1 M phosphate buffer, pH

7.2 (Kwan et al., 2013) and then bisected longitudinally using a fine razor blade under a

stereomicroscope. Fixed, bisected pulvini were immersed in an aqueous solution of 3% NaOH,

5% bleach for approximately 20 hours at room temperature until they appeared uniformly white

and translucent under transmitted light illumination, indicating maceration of cellular material.

Pulvini were then gently rinsed in distilled water for 24 hours, dehydrated in 100% ethanol,

critical point dried in liquid CO_2, mounted on aluminum stubs with carbon double-sided tape, and

sputter coated with a 4 nm layer of gold and palladium. Exposed internal faces of the bisected

pulvini were imaged at an accelerating voltage of 10 kV, providing lateral views of the internal

structure of the decellularized organs.

The spatial orientations of cellulose microfibrils composing parenchyma cell walls were

measured from high-magnification (~50,000-65,000x) scanning electron micrographs. Cellulose

microfibril orientation was quantified using the Directionality plugin in ImageJ (Liu, 1991).

Analysis via either the Fourier components or local gradient modes of the plugin yielded similar

results, and results from the local gradient method are reported here. Orientations of identifiable

structures in each cell wall micrograph were expressed as a histogram with portion of structures

observed on the Y axis and angular orientation in degrees on the X axis. The peak of a

gaussian curve fit to the collective histogram compiled from all micrographs provided a

quantitative metric of the predominant orientation of cellulose microfibrils across samples, and

the standard deviation of the gaussian curve provided a metric of histogram width, i.e., the

dispersion of microfibril orientations about the predominant angle.

Aspect ratios and spatial orientations of primary pit fields, which physically connect

adjacent parenchyma cells, were also measured from scanning electron micrographs manually

using ImageJ. Pit fields were identified as matted areas of cell wall perforated by numerous

plasmodesmata. The spatial orientations of both pit fields and cellulose microfibrils were

measured relative to the pulvinus long axis, as indicated by the orientations of nearby

vasculature running longitudinally through the organ core.

Multiscale osmotic swelling

To explore differences in osmotic swelling behaviors across scales of pulvinus

organization, pulvini in incremental states of dissection were imaged in aqueous mannitol baths

of varying tonicity. Mannitol is a biologically inert molecule and has been used previously to

induce osmotic deformation of pulvinus tissues (Mayer et al., 1985). Bathing in relatively dilute,

hypotonic mannitol solutions produces an osmotic gradient that favors movement of water into

cells, resulting in tissue expansion, while bathing in relatively concentrated, hypertonic mannitol

solutions favors movement of water out of cells, resulting in tissue shrinkage.

Three preparations of pulvinus organ were subjected to graded osmotic manipulation:

cylindrical sections of pulvini, cylindrical sections skinned of their epidermis, and isolated blocks

of parenchyma tissue. Cylindrical sections of pulvinus organ were produced by removing the

proximal and distal ends of isolated pulvini using a fine razor blade under a stereomicroscope,

resulting in sushi roll-like sections of pulvinus organ 1000-1500 μm long and 1000-1400 μm

wide. Sections comprised a core of vasculature surrounded concentrically by a thick

parenchyma cortex and thin enveloping epidermis. Removal of the epidermis from a subset of

sections was accomplished via manual skinning with a fine razor under a stereomicroscope and

produced cylindrical sections containing only the vasculature core and surrounding parenchyma

cortex. Isolated parenchyma blocks were taken from the relatively large extensor (i.e. ventral or

abaxial) cortexes of primary pulvini and were free of both vascular and epidermis components.

Flexor (i.e. dorsal or adaxial) cortexes of *M. pudica* pulvini were found to be too small to reliably

yield parenchyma samples via manual dissection under a stereomicroscope, and were not

examined in the current study.

Dissected pulvinus preparations were transferred through a graded series of osmotic

mannitol baths ranging in concentration from 0 to 878 milliosmoles (mOsm). Solution

concentrations were calculated and prepared via serial dilution of measured quantities of

mannitol dissolved in distilled water. Following an acclimation time of at least 30 minutes in each

osmotic bath, samples were imaged in both lateral and ventral views using a digital camera

(Zeiss AxioCam MRc, Carl Zeiss AG) mounted on a stereomicroscope (Zeiss SteREO

Discovery V8, Carl Zeiss AG). Widths, lengths, and heights of dissected pulvinus preparations

were measured at their maxima from stereomicrographs using ImageJ. Measurements of

dissected pulvinus preparations were normalized to dimensions in the most hypertonic osmotic

bath employed (878 mOsm). Statistically significant changes in tissue size were identified via

two-tailed, one sample t-tests comparing normalized dimensions in the most hypertonic (878

mOsm) and most hypotonic (distilled water) bathing solutions.

Parenchyma cell geometry

The geometries of live parenchyma cells were visualized using a laser scanning confocal

microscope (LSM710, Carl Zeiss AG). Thin sections of fresh parenchyma tissue were collected

from the relatively large extensor (i.e. ventral or abaxial) cortexes of primary pulvini using a pair

of razor blades spaced by ~400 µm of double-sided tape. Thin sections were prepared for

confocal microscopy by staining briefly (7-9 minutes) with 0.5 mg/ml propidium iodide (PI) in

osmotically isotonic bathing solution. PI binds to cell wall pectin and is a common stain for live

imaging of plant anatomical features (Bidhendi et al., 2020). Isotonic bathing tonicity was

determined by bathing parenchyma tissue in a graded series of aqueous mannitol solutions and

measuring changes in tissue size. Osmotic baths more concentrated than 384 mOsm of

mannitol were found to cause shrinkage of freshly dissected parenchyma tissues, while osmotic

baths less concentrated than 384 mOsm caused expansion. After fluorescent staining,

parenchyma sections were rinsed briefly of unbound PI and then excited with a 543 nm Helium-

Neon laser at 5% power. Fluorescence emission in a range of 554-711 nm was captured from

parenchyma cells deep to the sectioned tissue surface. Cells with a fluorescing nucleus

indicating penetration of PI into the cell protoplasm were considered damaged by sectioning and

excluded from the study. Optical sections 2.4-6.4 µm thick were captured at the midlines of cells

of interest, located by changing focus along the z-axis until the largest apparent cell dimensions

were presented. Slice thickness was changed by manipulating the size of the confocal pinhole.

To examine the influence of increased intracellular fluid volume on the geometry of

individual parenchyma cells, parenchyma tissues visualized via confocal microscopy were

initially imaged in osmotically isotonic bathing solution and subsequently re-imaged in

osmotically hypotonic baths of distilled water. Isotonic and hypotonic baths were exchanged

without moving parenchyma tissue samples from the confocal stage, allowing visualization of

individual parenchyma cells of interest before and after osmotic swelling. An acclimation time of

a least 5 minutes in hypotonic solution was given prior to re-imaging of individual cells. The

lengths, heights, widths, lateral perimeters, and frontal perimeters of individual parenchyma

cells were measured from confocal micrographs in ImageJ and normalized to dimensions in

osmotically isotonic baths. Changes between isotonic and hypotonic solutions were expressed

as percentages, and significant changes in cell dimensions between the two bath conditions

were identified as those significantly different from zero in a two-tailed one sample t-test. Lateral

aspect ratios of parenchyma cells were calculated by dividing cell length (measured in the

direction parallel to the long axis of the pulvinus) by cell height (measured along the dorsal-

ventral axis of the organ). Frontal aspect ratios of cells were calculated by dividing cell width by

cell height. Non-circular aspect ratios were defined as those that differed significantly from 1.0 in

a two-tailed one sample t-test.

Anatomical descriptors

All anatomical descriptors employed in the current study are defined relative to organ-

scale pulvinus anatomy. Lengths are measured along the long axis of the pulvinus, defined by

the orientation of the vasculature tissue running longitudinally through the organ center. Heights

are measured along the dorsal-ventral axis of the pulvinus, perpendicular to the neutral axis of

active pulvinus bending. Widths are measured laterally across the cross-section of pulvini.

These terms are employed across all hierarchical scales of organization examined. For

example, the length of an individual parenchyma cell refers to size measured along the long

axis of the pulvinus it composes.

<u>**Results**</u>

Pulvini of *Mimosa pudica* displayed structural anisotropy at all scales of organization

examined. Epidermal cells bounding the surfaces of pulvini were oblong in shape with long axes

oriented perpendicular to the organ long axis (Fig. 1). Wrinkles in the pulvinus surface were

oriented predominantly transverse to the pulvinus long axis, i.e., parallel to the long axes of

epidermal cells. Surface wrinkles were confined exclusively to pulvinus organs, and were not

apparent on neighboring surfaces of stems, petioles, or leaflets.

Figure 1: Pulvinus locations and external morphologies. [A] Line drawing depicting the locations of primary (B), secondary (C), and tertiary (D) pulvini of *Mimosa pudica*. [B,C,D] Scanning electron micrographs of pulvinus organs identified in A. Inset micrographs show epidermis morphology at higher magnifications. In the immediate regions of pulvini, epidermal cells are oblong and oriented perpendicular to the organ long axis. Wrinkles in the epidermis run parallel to the long axes of epidermal cells and are absent outside of the immediate regions of pulvini.

The shapes of epidermal cells of primary pulvini transitioned from oblong and

perpendicularly oriented in the region of the pulvinus to oblong and parallelly oriented in the

region of the petiole (Fig. 2). Epidermal cells in the intermediate region between pulvinus and

petiole displayed relatively circular aspect ratios (Fig. 2b) and did not display an obvious

orientation relative to the organ long axis (Fig. 2c). The average epidermal cell aspect ratio in

the region of the pulvinus was 3.70 ± 1.15 (mean ± SD) oriented at an average angle of 88.2° ±

6.85° (mean ± SD) relative to the organ long axis.

230
231
Figure 2: Spatial variation in epidermal cell morphology. [A] Stitched scanning electron
micrographs showing a primary pulvinus organ in lateral view. Yellow line depicts a path traced
along the central axis of pulvinus and petiole. Inset micrographs show epidermis morphology at
higher magnification. Labels on inset micrographs indicate the approximate location at which
each was taken, expressed as a portion of the total path length examined. [B] Epidermal cell
aspect ratio varies across the lengths of pulvini and petioles. Dashed blue line denotes a
circular aspect ratio of 1. [C] Epidermal cell orientations expressed relative to the orientation of
the central path. Dashed blue lines fall at 90° and 0° denoting perpendicular and parallel cell

orientations, respectively. Data points collected from the same pulvini are depicted in matching
shades of gray in both B and C.
242

243   High magnification scanning electron microscopy allowed visualization of individual

244   cellulose microfibrils at the inner surfaces of parenchymal cell walls (Fig. 3). Cellulose

245   microfibrils were aligned primarily dorso-ventrally in the plane of the wall with a predominant

246   microfibril orientation of 82.5° ± 29.1° (mean ± SD) relative to the orientation of nearby

247   vasculature.

Figure 3: Orientations of cellulose microfibrils reinforcing parenchymal cell walls. [A] Scanning electron micrograph showing a lateral view of a longitudinally bisected primary pulvinus. Protoplasts have been removed chemically to reveal architecture at the inner face of the cell wall. Labels indicate dorsal, i.e. adaxial, epidermis (d.e.), dorsal parenchyma (d.p.), vasculature (vasc.), ventral, i.e. abaxial, parenchyma (v.p.), and ventral epidermis (v.e.). Organ is oriented with petiole (pet.; not visible) left of the frame. [B] Magnified view of the highlighted region in A showing ventral parenchyma bounded by vasculature and ventral epidermis. [C] Magnified view of the highlighted region in B showing the inner face of one parenchymal cell wall. [D] High

magnification view of the highlighted region in C showing cellulose microfibrils running primarily
dorso-ventrally in the plane of the cell wall. Fiber angles are described relative to the long axis
of nearby vasculature (defined as 0°) with 90° indicating a perpendicular orientation. [E]
Histograms showing distributions of fiber orientations in 9 cells from 4 pulvini. Each gray trace
shows the distribution of fiber angles in a high magnification scanning electron micrograph
similar to D. A gaussian curve (solid red trace) fit to all histograms indicates a predominant fiber
orientation of 82.5 degrees (solid vertical line) with a standard deviation of ± 29.1 degrees
(dotted vertical lines).

Consistent with the necessity for rapid fluid exchange during active movement, primary

pit fields connecting adjacent parenchyma cells were prevalent in pulvini of *M. pudica*. Pit fields

displayed a mix of circular, oval, and spindle-shaped morphologies (Fig. 4). In many

parenchyma cells, pit fields were predominantly oval or spindle-shaped with long axes oriented

perpendicular to the organ axis. For example, pit fields in the five parenchyma cells depicted in

figure 4b displayed aspect ratios ranging from 1.2 to 4 with an average aspect ratio of 2.22 ±

0.78 (mean ± SD, n=30 pit fields) and an average long axis orientation of 93.0 ± 25.0 degrees

(mean ± SD) relative to the organ axis. Parenchyma cells with predominantly oval or spindle-

shaped pit fields were often directly adjacent to cells with predominantly circular pit fields,

indicating that parenchyma cells of similar size, shape, and position can possess pit fields of

varying morphology.

Figure 4: Primary pit fields visible in the walls of decellularized parenchyma cells. [A] Scanning
electron micrograph showing a lateral view of a longitudinally bisected primary pulvinus. The
organ long axis is oriented horizontally with the central vascular core visible at the top of the
panel. Pit fields are visible as darkened regions of cell wall and display a mix of circular, oval,
and spindle-shaped morphologies. Yellow asterisks highlight cells displaying predominantly
oblong pit fields oriented perpendicular to the long axis of the pulvinus. White asterisks highlight
cells with more circular pit field morphologies. [B] Magnified view of the highlighted region in A
showing parenchyma cells with predominantly spindle-shaped pit fields oriented perpendicular
to the organ long axis. [C] A single parenchyma cell displaying a mix of circular and oval shaped
pit fields. [D] A single spindle-shaped pit field composed of mat-like cellulose microfibrils
perforated by numerous plasmodesmata channels.

Osmotic swelling of dissected pulvinus organs and tissue samples yielded anisotropic

deformation across all scales of pulvinus organization studied (Fig. 5). Across a graded series of

osmotic baths ranging from 0 to 878 mOsm, cylindrical sections of pulvinus organ with an intact

epidermis (Fig. 5c) displayed a 15.3 ± 2.75% (mean ± SD, $p < 0.01$) change in length without

significant changes in either width (1.74 ± 1.83%, $p = 0.11$) or height (1.02 ± 2.42%, $p = 0.10$).

Cylindrical pulvinus sections skinned of their epidermis (Fig. 5d) varied in length 17.7 ± 4.26%

($p < 0.01$), increased marginally in width (2.91 ± 1.98%, $p < 0.01$), and displayed increased

variation in height (7.80 ± 8.55%, $p = 0.05$). The increased variation in the height of cylindrical

organ sections skinned of their epidermis was attributable to a tendency for elongating

parenchyma tissues to bow dorsally and ventrally away from the central vasculature of the

pulvinus organ. This bowing resulted in skinned pulvinus sections adopting spindle-shaped

lateral profiles reminiscent of rugby or American footballs when swollen osmotically (Fig. 5f).

Isolated blocks of parenchyma tissue dissected free of epidermis and vasculature (Fig. 5e)

changed in length 26.2 ± 3.15% ($p < 0.01$) without statistically significant changes in either width

(-1.39 ± 2.40%, $p = 0.26$) or height (3.76 ± 5.75%, $p = 0.22$).

Figure 5: Osmotic perturbations reveal anisotropic swelling behaviors across scales of organization. [A] Lateral view of an intact *Mimosa pudica* primary pulvinus with flanking leaves removed. Dashed white lines illustrate cross-sectional cuts used to isolate a cylindrical section of pulvinus. Orange arrowheads in A and C indicate proximal, ventral corner of sections. [B] Isolated cylindrical section of pulvinus with epidermis partially removed to illustrate skinning technique. [C, D, E] Effect of graded osmotic bathing solutions on the length (L, black circles), cross-sectional width (w, blue squares), and height (h, red diamonds) of excised pulvinus sections. Inset line drawings depict dimensions and typical morphologies in lateral and cross-sectional views. Graph markers are means ± SD. Data is normalized to tissue dimensions in the most hypertonic bathing solution (878 mOsm). Asterisks denote values significantly different

from 1.0 in the most hypotonic bathing solution (0 mOsm). [C] Isolated pulvinus sections expand longitudinally in increasingly hypotonic bathing solutions with negligible change in width or height. [D] Sections manually skinned of their epidermis expand primarily in length but display a small increase in width and increased variability in height. [E] Isolated blocks of parenchyma tissue expand longitudinally with negligible change in width or height. Sample sizes (n) for each condition are noted beside inset illustrations. [F] Representative images illustrating shape changes of pulvinus preparations. Upper and lower images of each pair display sample shape in hypertonic and hypotonic baths, respectively. Scale bars 500 μm.

The geometries of live parenchyma cells visualized by confocal microscopy depended

on the tonicity of the solution in which they were bathed (Fig. 6). In isotonic bathing solution, the

average ratio of cell width to cell height was 0.95 ± 0.15 (mean ± SD), which does not differ

significantly from a circular cross-sectional aspect ratio of 1.0 (p=0.54). The average ratio of cell

length to cell height was 0.85 ± 0.27 (mean ± SD), which differs significantly from a circular

lateral aspect ratio of 1.0 (p=0.01), confirming that live parenchyma cells bathed in isotonic

solution are approximately disk-shaped; circular in cross section but flattened longitudinally in

the direction parallel to the long axis of the pulvinus. Cells reimaged in osmotically hypotonic

distilled water baths did not change significantly in width (-0.56 ± 1.88%, mean ± SD, p = 0.13),

displayed a small but significant change in height (1.78 ± 2.99%, p < 0.01), and displayed a

relatively large increase in length of 14.5 ± 5.07% (p < 0.01) occurring along the long axis of the

pulvinus (Fig. 6b). Following transfer to hypotonic bathing solution, cells displayed an average

ratio of cell length to cell height of 0.96 ± 0.30, which was not significantly different from circular

(p=0.54), indicating that longitudinally flattened, disk-shaped cells expanded towards more

perfectly circular lateral profiles under increased turgor.

Measurements of frontal and lateral cell perimeters revealed anisotropy in the amount of

cell wall distension that occurred during osmotic expansion. Lateral cell perimeter, measured in

the plane parallel to the organ long axis, increased 6.87 ± 3.10% (p < 0.01) following

acclimation in hypotonic bathing solution. In contrast, frontal cell perimeter measured in cross

section perpendicular to the organ long axis changed only marginally (0.96 ± 1.41%, p < 0.01).

This difference in cell wall distension under isotropic turgor rise indicates that the parenchyma

cell wall is materially anisotropic, displaying greater extensibility in the longitudinal direction than

in the circumferential.

Figure 6: Nonuniform osmotic swelling of individual parenchyma cells. [A] Confocal micrographs
of parenchyma tissue imaged in osmotically isotonic (384 mOsm) and hypotonic (distilled water)
baths. Representative micrographs show cell morphology as seen in frontal and lateral views.
All scale bars 20 μm. Inset line drawings illustrate anatomical dimensions and highlight changes
in cell shape between osmotic baths. [B] Effect of osmotic swelling on frontal and lateral
dimensions of 53 cells from 8 pulvini. Gray dots represent percent changes in cell dimensions
following transfer from isotonic to hypotonic bath. Black markers represent means ± standard
errors. Asterisks denote percent changes significantly different from zero (p<0.05).

Discussion

The hydraulic nature of pulvinus-driven movement has been recognized since the 1800s

and is discussed in Darwin's *The Power of Movement in Plants* (Darwin and Darwin, 1880).

However, an understanding of the functional morphologies that transform isotropic cellular

turgor pressures into anisotropic, directed movement at the organ scale has remained

fragmented. Here we demonstrate that in pulvini of *Mimosa pudica,* multiple anisotropic

structural features are present across subcellular, cellular, and tissue levels of organization.

These include perpendicularly oriented cellulose microfibrils and primary pit fields in

parenchymal cell walls, longitudinally flattened parenchyma cell geometries, and perpendicularly

oriented epidermal cells associated with transverse surface wrinkles. We observed highly

anisotropic osmotic swelling of parenchyma cells and tissues; however, swelling behaviors at

higher levels of organization were also influenced by the presence of the enveloping epidermis,

suggesting that hydraulic deformation of pulvini is influenced by structural anisotropy at multiple

hierarchical scales.

Cell wall anisotropy

Consistent with polarized light investigations of pulvinus morphology in the scarlet runner

bean *Phaseolus coccineus L* (Mayer et al., 1985), scanning electron micrographs of

decellularized *M. pudica* pulvini revealed cellulose microfibrils oriented predominantly

perpendicular to the long axes of pulvini. Cellulose microfibrils are crystalline structures

composed of closely packed chains of parallelly-oriented cellulose molecules (Brett, 2000) and

display high tensile strengths and longitudinal stiffnesses of approximately 1 GPa and 130 GPa

respectively (Gibson, 2012). Cellulose microfibrils function as the primary mechanical

component of the plant cell wall, ultimately providing the tensile strength needed to balance

large intracellular turgor pressures ranging from 0.4-0.8 MPa in typical hydrated plant tissues

(Taiz et al., 2015) to over 1 MPa in some specialized plant cells (Franks, 2003).

Anisotropic patterns of cellulose reinforcement are not unique to pulvini. Cellulose

orientation is actively controlled by the microtubule-guided motion of cellulose synthase

complexes that extrude cellulose microfibrils into the extracellular space (Brett, 2000), and

anisotropic cellulose deposition is the primary means by which mechanical anisotropy is

produced and maintained in plant cell walls. The interplay of intracellular turgor pressure and

cell wall anisotropy ultimately governs the direction of irreversible turgor induced shape changes

(Baskin, 2005) and underlies the development of complex non-spherical morphologies

(Bidhendi and Geitmann, 2016) in growing plant cells. Interactions between cell wall anisotropy

and turgor pressure have also been shown to underlie reversible deformation of certain mature

plant cell types. The hydraulic guard cells flanking stomatal pores, for example, are reinforced

by cellulose microfibrils oriented primarily perpendicular to the guard cell long axes (Galatis and

Apostolakos, 2004). The perpendicular orientation of microfibrils inhibits radial expansion of

guard cells while permitting elongation during hydraulic expansion, ultimately facilitating the

microscopic outward bowing of guard cells that facilitates repeated opening and closing of

stomatal pores (Aylor et al., 1973). Parenchyma cells observed in the current study displayed a

high degree of directional anisotropy in the extensibilities of their cell walls, with osmotic

expansion causing wall distension approximately 7 times greater in the longitudinal direction

than in the circumferential. Analogous to the effect of cellulose anisotropy in stomatal guard

cells, this finding suggests that cellulose microfibrils selectively restrict circumferential

expansion of parenchymal cell walls in the pulvini of *M. pudica*, resulting in predominantly

longitudinal cell wall deformation in response to turgor change.

In addition to unidirectionally oriented cellulose microfibrils, frequent primary pit fields

also have the potential to contribute to mechanical anisotropy in the parenchyma cell wall.

Primary pit fields are typically circular, sometimes oval regions of the primary cell wall perforated

by numerous plasmodesmata. In addition to being perforated, the cell wall is significantly thinner

in pit fields compared to the surrounding wall, commonly by a factor of 5 or more (e.g. Kobiyama

and Crandall-Stotler, 1999). The presence of numerous primary pit fields is consistent with the

importance of rapid water exchange between cells in the *M. pudica* pulvinus; however, the

presence of many oblong and spindle-shaped pit fields oriented perpendicular to the organ axis

suggests their shape and orientation may also serve a mechanical function. Assuming the

reduced thickness of the pit field is associated with a lower elastic modulus, oblong pit fields

oriented perpendicular to the pulvinus axis may serve as expandable seams that facilitate

distension of the cell wall preferentially in the longitudinal direction. Direct measurements of pit

field deformation or elastic modulus are needed to verify this hypothesis, and exploration of the

influence of pit field material properties and shape on cell wall mechanics may prove a fruitful

topic for future research.

Geometric deformation of parenchyma cells

While perpendicularly oriented cellulose microfibrils and pit fields are conducive to long-

axis deformation of parenchyma cells, our results indicate that direct stretching of the cell wall

cannot account for all of the longitudinal cell deformation observed in the current study.

Following transfer from isotonic to hypotonic bathing solution, the average length of parenchyma

cells increased by 14.5 ± 5.07% (mean ± SD); however, the lateral perimeters of these same

parenchyma cells increased by only 6.87 ± 3.10%, indicating that direct stretching of the cell

wall accounts for less than half of longitudinal cell deformation on average, even under the

assumption that all cell wall distension contributes exclusively to elongation. Geometric

deformation of cells from longitudinally flattened lateral profiles towards more longitudinally

expanded lateral profiles likely accounts for the remaining observed cell deformation. Like the

corrugated bellows of an accordion or concertina, the longitudinally flattened morphologies of

parenchyma cells are conducive to longitudinal expansion and contraction via geometric

changes in cell shape. We observed both direct distension of the cell wall and geometric shape

changes of cells under increased turgor, indicating that both of these mechanisms contributed to

anisotropic parenchyma cell deformation during experimentally imposed swelling. Further

studies of parenchyma cell deformation at physiologically realistic intracellular turgor pressures

are needed to determine the relative contributions of these two mechanisms to anisotropic

deformation *in vivo*.

Here we focus exclusively on parenchyma cells from the relatively large extensor (i.e.

ventral or abaxial) cortex of the primary pulvinus of *M. pudica*. Comparison with morphology and

swelling behavior of the flexor cortex was not carried out in the current study but is a worthwhile

topic for future research. We expect that parenchyma cells of the flexor cortex would display

similar structural anisotropy and directional swelling behavior to that of the extensor cortex;

however, subtle differences in morphology across cortexes likely influence their behaviors. In

pulvini of the scarlet runner bean *Phaseolus coccineus L.,* for example, both extensor and flexor

cortexes are composed of longitudinally flattened, disk-like parenchyma cells. Cells of the

extensor compartment are notably larger in size, and isolated blocks of extensor parenchyma

tissue display relatively larger percent changes in length than flexor tissues when osmotically

swollen in distilled water (Mayer et al. 1985). Such differences may be relevant to pulvinus

mechanics *in vivo*, and further investigation may provide insight into the granularity of

morphological specialization displayed by parenchyma cells that are subject to differing

functional demands.

Mechanical contribution of the pulvinus epidermis

Most plant stems and petioles are ensheathed in a skin of elongate epidermal cells

oriented parallel to the long axis of the organ (Kutschera, 2008). This parallel orientation of

epidermal cells is a natural consequence of turgor-driven morphogenesis, in which elongated

plant organs form developmentally through anisotropic expansion and elongation of the cells

that compose them. Over most of their lengths, the petioles of *M. pudica* display a typical

epidermis morphology of elongate cells oriented parallel to the organ long axis. Surprisingly,

however, we observed epidermal cells oriented perpendicular to the growth axis in the

immediate regions of pulvini. Perpendicular cells correlated spatially with the presence of

transverse wrinkles oriented perpendicular to the long axis of the pulvinus. Transverse wrinkles

in the epidermis of *Mimosa pudica* have been described previously and have been shown to

expand and contract dynamically during active movement of live plants (Song et al., 2014).

They are thought to provide extra surface area that increases the longitudinal extensibility of the

epidermis without compromising its circumferential stiffness (Mano and Hasebe, 2021).

We suggest that perpendicularly oriented epidermal cells are a morphological

specialization that facilitates transverse wrinkling of the epidermis by reducing its effective

flexural stiffness in the longitudinal direction. Epidermal cell orientation has been shown to have

a direct effect on the material stiffness of plant epidermis, inducing anisotropic tensile properties

and relatively lower tensile stiffness in the direction perpendicular to the orientation of epidermal

cells (Bidhendi et al. 2020). Importantly, a perpendicular orientation of epidermal cells also

allows transverse wrinkles to form between epidermal cells rather than across their lengths,

minimizing bending strains imposed on individual epidermal cells. Direct mechanical tests

exploring the impact of epidermal cell orientation on flexural stiffness are needed; however,

facilitation of transverse wrinkle formation provides a potential explanation for the observation of

perpendicularly oriented epidermal cells confined to the immediate regions of pulvini.

In cylindrical sections of pulvinus organ skinned of an enveloping epidermis, we

observed outward bowing of parenchyma tissue away from the central axis of cylindrical

pulvinus sections during osmotic swelling. We suspect that this parenchymal bowing results

from a difference in the longitudinal extensibility of parenchyma tissues and the vascular tissue

they surround. Isolated parenchyma tissues dissected away from the mechanical context of the

pulvinus organ display a strong bias for purely longitudinal expansion; however, when coupled

to a relatively stiff vasculature core in the context of an intact organ, pure longitudinal expansion

of the parenchyma along a straight line is resisted, and outward curvature of longitudinally

expanding parenchyma tissues results. We did not observe the phenomenon of outward

parenchyma bowing in pulvinus preparations with an intact epidermis, suggesting that the

circumferential stiffness of the epidermis resists outward bowing of parenchyma tissues. While

anisotropic cell wall reinforcement and cell geometry are well suited to ensure longitudinal

expansion at the scale of individual parenchyma cells, our findings suggest that the transversely

wrinkled epidermis serves as an encompassing sleeve that restricts radial deformations that

emerge at higher scales of tissue organization, ensuring that hydraulic expansion of

parenchyma tissue *en masse* remains aligned along the long axis of the pulvinus where its

effect on organ bending is greatest.

Osmotic swelling as a model for *in vivo* physiology

Simple osmotic swelling and shrinkage of pulvinus tissues does not replicate the rapid

and complex mechanisms of intercellular fluid movement employed by *Mimosa pudica* pulvini *in*

vivo, the details of which are not fully understood. Rather, osmotic perturbations employed in

the current study provide a means of introducing purely isotropic, nondirectional intracellular

loads into parenchyma cells in the form of hydrostatic fluid pressures, the effects of which

provide insight into multiscale tissue mechanics. While the precise physiological mechanisms

powering active fluid movement and turgor change *in vivo* is an important topic for further

investigation, here we show that hydraulic deformation of *M. pudica* pulvini is a product not only

of biochemical specialization, but also of a suite of downstream structural features that span

multiple hierarchical scales of organization. These features are ideally situated to translate

isotropic, non-directional cellular turgor pressures into the physiologically relevant motor task of

long-axis organ bending, suggesting that they are important contributors to the mechanism of

pulvinus-driven motion *in vivo*.

Acknowledgements

511 We thank Youssef Chebli for help in microscope operation and helpful discussions about data

512 collection, and David Liu and S. Kelly Sears at the Facility for Electron Microscopy Research of

513 McGill University for help in microscope operation and data collection.

Funding

516 Funded by a Human Frontier Science Program long-term postdoctoral fellowship (LT - 000531/

517 2020).

Competing Interests

520 We have no competing interests to declare.

Data availability

523 Tabulated data from all figures are available as supplemental files.

References Cited

Adamski, M., Fontana, G., Gershlak, J. R., Gaudette, G. R., Le, H. D. and Murphy, W. L. (2018).

Two methods for decellularization of plant tissues for tissue engineering applications. *J.*

Vis. Exp., 135, e57586. 10.3791/57586.

Aylor, D. E., Parlange, J. Y. and Krikorian, A. D. (1973). Stomatal mechanics. *Am. J. Bot., 60*(2),

163-171. 10.1002/j.1537-2197.1973.tb10213.x.

Baskin, T. I. (2005). Anisotropic expansion of the plant cell wall. *Annu. Rev. Cell Dev. Biol., 21*,

203-222. 10.1146/annurev.cellbio.20.082503.103053.

Bidhendi, A. J., Chebli, Y. and Geitmann, A. (2020). Fluorescence visualization of cellulose and

pectin in the primary plant cell wall. *J. Microsc., 278*(3), 164-181. 10.1111/jmi.12895.

Bidhendi, A. J. and Geitmann, A. (2016). Relating the mechanics of the primary plant cell wall to

morphogenesis. *J. Exp. Bot., 67*(2), 449-461. 10.1093/jxb/erv535.

Bidhendi, A. J., Li, H. and Geitmann, A. (2020). Modeling the nonlinear elastic behavior of plant

epidermis. *Botany, 98*(1), 49-64. 10.1139/cjb-2019-0058.

Brett, C. T. (2000). Cellulose microfibrils in plants: biosynthesis, deposition, and integration into

the cell wall. *Int. Rev. Cytol.,* 199, 161-199. 10.1016/S0074-7696(00)99004-1.

Darwin, C. and Darwin, F. (1880). *The power of movement in plants*. Appleton, New York, NY.

Franks, P. J. (2003). Use of the pressure probe in studies of stomatal function. *J. Exp. Bot.,*

54(387), 1495-1504. 10.1093/jxb/erg162.

Fromm, J. and Eschrich, W. (1988a). Transport processes in stimulated and non-stimulated

leaves of *Mimosa pudica*. *Trees, 2*(1), 7-17. 10.1007/BF00196974.

Fromm, J. and Eschrich, W. (1988b). Transport processes in stimulated and non-stimulated

leaves of *Mimosa pudica*. *Trees, 2*(1), 18-24. 10.1007/BF00196975.

Galatis, B. and Apostolakos, P. (2004). The role of the cytoskeleton in the morphogenesis and

function of stomatal complexes. *New Phytol., 161*(3), 613-639. 10.1046/j.1469-

8137.2003.00986.x.

Gibson, L. J. (2012). The hierarchical structure and mechanics of plant materials. *J. R. Soc.*

Interface, 9(76), 2749-2766. 10.1098/rsif.2012.0341.

Kagawa, H. and Saito, E. (2000). A model on the main pulvinus movement of *Mimosa pudica*.

JSME Int. J. C-Mech Sy., 43(4), 923-928. 10.1299/jsmec.43.923.

Kobiyama, Y. and Crandall-Stotler, B. (1999) Studies of specialized pitted parenchyma cells of

the liverwort *Conocephalum* Hill and their phylogenetic implications. *Int. J. Plant Sci.*,

160(2), 351-370. 10.1086/314126.

Koller, D. and Zamski, E. (2002). The phototropic pulvinus of bean *Phaseolus vulgaris L.*-

Functional features. *Plant Biol., 4*(5), 584-594. 10.1055/s-2002-35442.

Kumon, K. and Suda, S. (1984). Ionic Fluxes from Pulvinar Cells during the Rapid Movement of

Mimosa pudica L. Plant. Cell. Physiol., 25(6), 975-979.

10.1093/oxfordjournals.pcp.a076813.

Kutschera, U. (2008). The growing outer epidermal wall: design and physiological role of a

composite structure. *Ann. Bot., 101*(5), 615-621. 10.1093/aob/mcn015.

Kwan, K., Ye, Z., Chye, M. and Ngan, A. (2013). A mathematical model on water redistribution

mechanism of the seismonastic movement of *Mimosa pudica*. *Biophys. J., 105*(1), 266-

275. 10.1016/j.bpj.2013.06.001.

Liu, Z. Q. (1991). Scale space approach to directional analysis of images. *Appl. Opt., 30*(11),

1369-1373. 10.1364/AO.30.001369.

Mano, H. and Hasebe, M. (2021). Rapid movements in plants. *J. Plant. Res., 134*(1), 3-17.

10.1007/s10265-020-01243-7.

Mayer, W.-E., Flach, D., Raju, M., Starrach, N. and Wiech, E. (1985). Mechanics of circadian

pulvini movements in *Phaseolus coccineus L. Planta, 163*(3), 381-390.

10.1007/BF00395147.

Morillon, R., Liénard, D., Chrispeels, M. J. and Lassalles, J.-P. (2001). Rapid movements of

plants organs require solute-water cotransporters or contractile proteins. *Plant Physiol.,*

127(3), 720-723. 10.1104/pp.010732.

Satter, R. L., Sabnis, D. D. and Galston, A. W. (1970). Phytochrome controlled nyctinasty in

Albizzia julibrissin. I. Anatomy and fine structure of the pulvinule. *Am. J. Bot., 57*(4), 374-

381. 10.1002/j.1537-2197.1970.tb09826.x.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V.,

Eliceiri, K., Tomancak, P. and Cardona, A. (2012). Fiji: an open-source platform for

biological-image analysis. *Nat. Methods, 9*(7), 676-682. 10.1038/nmeth.2019.

Scorza, L. C. T. and Dornelas, M. C. (2011). Plants on the move: towards common mechanisms

governing mechanically-induced plant movements. *Plant Signal. Behav., 6*(12), 1979-

1986. 10.4161/psb.6.12.18192.

Song, K., Yeom, E. and Lee, S. J. (2014). Real-time imaging of pulvinus bending *in Mimosa*

pudica. *Sci. Rep., 4*, 6466. 10.1038/srep06466.

Taiz, L., Zeiger, E., Møller, I. M. and Murphy, A. (2015). *Plant physiology and development*:

Sinauer Associates, Sunderland, CT.

Tamiya, T., Miyazaki, T., Ishikawa, H., Iriguchi, N., Maki, T., Matsumoto, J. J. and Tsuchiya, T.

(1988). Movement of water in conjunction with plant movement visualized by NMR

imaging. *J. Biochem., 104*(1), 5-8. 10.1093/oxfordjournals.jbchem.a122421.

Wang, Y. and Li, H. (2020). Bio-chemo-electro-mechanical modelling of the rapid movement of

Mimosa pudica. *Bioelectrochemistry, 134*, 107533. 10.1016/j.bioelechem.2020.107533.