**Mucin-degrading gut bacteria promote anti-parasitic immunity**

**Short title:** *Nematodes and gut microbial mucin foraging*

Mathis Wolter[a,b], Marie Boudaud[a#], Erica T. Grant[a,b], Amy Parrish[a,b], Alessandro De Sciscio[a],

Seona Thompson[c,d], Jean-Jacques Gerardy[e,f], Michel Mittelbronn[e,f,g,h,i,j], David J. Thornton[c,d],

Andrew J. Macpherson[k,l,m], Richard K. Grencis[c,d] and Mahesh S. Desai[a,n,*]

[a]*Department of Infection and Immunity, Luxembourg Institute of Health, Esch-sur-Alzette,*

*Luxembourg*

[b]*Faculty of Science, Technology and Medicine, University of Luxembourg, Esch-sur-Alzette,*

*Luxembourg*

[c]*Lydia Becker Institute for Immunology and Inflammation. Faculty of Biology, Medicine and*

*Health, University of Manchester, Manchester, United Kingdom*

[d]*Wellcome Centre for Cell Matrix Research, Faculty of Biology, Medicine and Health,*

*University of Manchester, Manchester, United Kingdom*

[e]*National Center of Pathology, Laboratoire National de Santé, Dudelange, Luxembourg*

[f]*Luxembourg Center of Neuropathology, Dudelange, Luxembourg*

[g]*Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette,*

*Luxembourg*

[h]*Department of Cancer Research, Luxembourg Institute of Health, Luxembourg, Luxembourg*

[i]*Faculty of Science, Technology and Medicine, University of Luxembourg, Esch-sur-Alzette*

[j]*Department of Life Science and Medicine, University of Luxembourg, Esch-sur-Alzette,*

*Luxembourg*

[k]*Department of Visceral Surgery and Medicine, Bern University Hospital, University of Bern,*

*Bern, Switzerland*

[l]*Maurice Müller Laboratories, Department for Biomedical Research, University of Bern,*

*Bern, Switzerland*

[m]*CMPG, Institute of Ecology and Evolution, University of Bern, Bern, Switzerland*

[n]*Odense Research Center for Anaphylaxis, Department of Dermatology and Allergy Center,*

*Odense University Hospital, University of Southern Denmark, Odense, Denmark*

[#]These authors contributed equally

[*]Corresponding author: Mahesh S. Desai, mahesh.desai@lih.lu

**Grant support**

This work was supported by the following grants in the laboratory of M.S.D.: Luxembourg National Research Fund (FNR) CORE grants (C15/BM/10318186 and C18/BM/12585940) and BRIDGES grant (22/17426243) to M.S.D.; FNR PRIDE (17/11823097) and the Fondation du Pélican de Mie et Pierre Hippert- Faber, under the aegis of the Fondation de Luxembourg grants to E.T.G.; FNR AFR individual PhD fellowship to A.P. (11602973); and European Commission Horizon 2020 Marie Skłodowska-Curie Actions individual fellowship to M.B. (897408). MM thanks the Luxembourg National Research Fond (FNR) for the support (FNR PEARL P16/BM/11192868 grant). The Wellcome Trust supports work of R.K.G and S.T. (Z10661/Z/18/Z; 203128A/Z/16/Z) and D.J.T (203128A/Z/16/Z).

**Correspondence**

Mahesh S. Desai, PhD

Luxembourg Institute of Health

Department of Infection and Immunity

29, rue Henri Koch, L-4354 Esch-sur-Alzette

Luxembourg

Tel: +352 26970-389

Fax: +352 26970-390

Email: mahesh.desai@lih.lu

**Disclosures**

M.S.D. works as a consultant and an advisory board member at Theralution GmbH, Germany.

**Author Contributions**

M.S.D. supervised the study and obtained research funding; M.W., M. B., D.J.T., R.K.G. and M.S.D. conceived and designed the study; M.W., M. B., E.T.G., A.P., A.D.S., and S.T. performed the experiments; M.W., E.T.G., R.K.G., A.P. and M.B. analyzed the data; J.-J. G. and M.M. prepared slides for histology; A.J.M. supplied germ-free mice; M.W., M. B., and M.S.D. primarily wrote the manuscript. All authors reviewed and approved the manuscript.

**Data Transparency Statement**

All data, study materials and analytical methods will be made available to other researchers

upon request.

**Abstract**

**BACKGROUND & AIMS**

Host-secreted gastrointestinal mucus plays a key role in the expulsion of intestinal nematode

parasites. A balance between mucin secretion by the host and the gut microbial mucin foraging

is essential to maintain the intestinal homeostasis, yet little is known about how changes in the

mucin–microbiome interactions affect worm infections. Here, we aimed to examine how mucin

foraging activity by the microbiome changes the course of parasitic worm infections by

modulating the host immune responses.

**METHODS**

We utilized a gnotobiotic mouse model containing a synthetic human gut microbiota that

allows for: 1) a complete removal of the mucin-degrading bacteria from the community; and

2) diet-driven manipulation of the microbiota toward mucin foraging. We infected mice with a

murine nematode, *Trichuris muris*, which resembles human infection with *Trichuris trichiura*.

We examined the temporal dynamics of worm infection including worm burden and the host

immune responses, and coupled these readouts to the microbial changes and mucin foraging

activity.

**RESULTS**

The absence of mucin-degrading bacteria in the microbiota enhances susceptibility to parasitic

infection—evidenced by higher worm number—by promoting stronger Th1 immune

responses. Dietary fiber deprivation increases the microbial mucin-foraging activity, which

coincides with a shift in host immune responses from susceptible (chronic, Th1 type) to

resistant (acute, Th2 type), thereby promoting worm clearance.

**CONCLUSIONS**

Our results provide mechanistic insights into how the colonic mucin-degrading bacteria

promote anti-parasitic immunity through modulation of the host immune responses. Our study

documents a clinically-relevant, novel link in the microbiome–parasite–host immune axis that

is useful prerequisite knowledge in managing parasitic infections.

**Keywords**

Mucin; gut microbiome; anti-parasitic immunity; nematode; *Trichuris muris*

## Introduction

Intestinal parasitic worms (helminths) make an invaluable model to study the host immunomodulatory mechanisms due to their ability to subvert the host immune responses[1]. In this context, the murine parasite *Trichuris muris*—a relevant biological model for the closely related human parasite *Trichuris trichiura*—has been extensively investigated[2]. *T. muris* is an especially attractive experimental model as either a susceptible or resistant immune response can be investigated[2,3]. Susceptibility in mice, which is more reflective of the naturally-occurring chronic infections[4], is generally characterized by a Th1 response resulting in long-term worm infections and chronic colitis. On the other hand, resistance is characterized by an acute Th2 response leading to worm clearance and tissue repair[2].

Since they occupy the same host environment, it is no surprise that enteric parasites such as helminths strongly interact with the gut microbiome[5,6]. Although these interactions may lead to the suppression of either the gut microbiome or the helminth[5–7], in case of *T. muris*, bacterial–helminth crosstalk is essential for worm hatching, as the parasite eggs fail to hatch and establish in germ-free environments[8–10]. Once the worm has settled in the intestinal epithelium, the bacterial–helminth dynamics appear mainly unidirectional, with helminths altering the microbiome composition through various mechanisms, including through helminth excretory-secretory products or host anti-microbial peptides, released in response to the parasite[11–15]. Furthermore, the helminths alter the nutrients available to the microbiome, indirectly influencing bacteria selection[16]. The effect of the infection on host immunology has an additional impact on the microbiome, as the worms or their metabolic byproducts can alter toll-like receptor (TLR) expression profiles[17,18] or responsiveness of host immune cells to TLR ligands[19] and interfere with local IgA-production[17]. While the gastrointestinal (GI) mucins are essential for worm clearance[20–22] and the impacts of the helminth infection on the microbiome during worm clearance are documented at the broader level of microbial phylogeny[23–28], questions remain regarding how the mucus–microbiome interactions are affected by and affect worm infection and the underlying host immune responses.

Worm expulsion during an acute response is largely driven by goblet-cell secreted gastrointestinal mucin that is tightly linked to the Th2-type immune response[22]. Along these lines, mice lacking the *Muc2* gene, which encodes the major glycoprotein of the GI mucus,

show a delayed clearance of *T. muris*[29] and *Muc5ac* deficient mice completely fail to clear the helminth[22].

We have previously shown that a fiber-deprived microbiota leads to a disrupted balance towards excessive gut microbial mucin foraging activity[30–32]. As the balance between mucin secretion and its consumption by the gut microbes is critical for intestinal homeostasis[33,34], here, we hypothesized that excessive mucin foraging by the microbiota alters the course of the parasite infection by modulating the immune response. To test this hypothesis, we employed a tractable 14-member synthetic human microbiota (14SM) characterized for the capability of individual members to degrade various glycans including mucin oligosaccharides[31,35]. In a longitudinal experimental setup, using ex-germ-free mice with a mucin-foraging or a non-mucin-foraging synthetic microbiota, we studied how mucin–gut microbiome interactions shape anti-parasitic immunity. Our findings suggest a mechanism of how mucin-degrading bacteria promote anti-parasitic immunity, which sheds a new light on the microbiome–parasite–host immune interactions.

## Methods

### *Ethical statement*

All animal experiments were performed according to the "Règlement Grand-Ducal du 11 janvier 2013 relatif à la protection des animaux utilisés à des fins scientifiques" based on Directive 2010/63/EU on the protection of animals used for scientific purposes, and approved by the Animal Experimentation Ethics Committee of the University of Luxembourg and by the Luxembourgish Ministry of Agriculture, Viticulture, and Rural Development (national authorization no. LUPA2020/02). The mice were housed in accordance with the recommendations stated by the Federation of European Laboratory Animal Science Associations (FELASA).

### *Experimental design and dietary treatment*

8–23 weeks old C57BL/6 mice were housed in ISOcages with up to five animals per cage. Sterile food and water were provided *ad libitum* and light cycles consisted of 12 hours. Before the administration of 14SM, the germ-free status of the mice was confirmed using aerobic and anaerobic culturing of stools samples. Mice were intragastrically gavaged with 0.2 ml of the 14SM or 10SM gavage mix as described previously[35]. Before and for 6–9 days following the

gavage, mice were maintained on a standard mouse chow, which we refer to as Fiber-rich (FR)

diet. Afterwards, half of the mice were assigned to a Fiber-free (FF) diet and the rest remained

on the FR diet. Twenty days after the diet switch, the mice were infected by oral gavage with

a high dose of approximately 450 eggs of *T. muris* strain E and observed for up to 30 days post-

infection (dpi). Mice were euthanized by cervical dislocation to determine worm counts and

other final readouts. A piece (~0.5 cm) of the cecum was fixed in methacarn in order to

determine goblet cell numbers. The remaining part of the cecum and 2 cm of the attached colon

were frozen at –20 °C to determine *T. muris* worm burden. A piece (~0.5 cm) of the colon was

emptied from content, longitudinally cut-opened and weighted before an incubation of 24 h at

37 °C, 5 % $CO_2$, in culture medium (RPMI with L-glutamine, 5% fetal bovine serum, 100U

penicillin-streptomycin) for analysis of cytokine secretion. The remaining colonic tissue and

mesenteric lymph nodes were stored in RNAprotect Tissue Reagent (Qiagen, Hilden,

Germany; catalog no. 76106, according to the manufacturer's protocol) or processed for

immune cell isolation and FACS analysis. Serum of the euthanized mice was collected and

stored at –20 °C.

***Parasitological techniques***

*T. muris* maintenance and infection was performed as previously described[36]. Mice were orally

infected with ~450 eggs and worm burdens were assessed by counting the number of worms

present in the cecum and 2 cm of the attached colon on days 20, 25 and 30 post infection, as

described previously[36]. The researcher, S.T., who counted the worms, was blinded for both the

individual time points and dietary groups.

**ELISA**

Lipocalin-2 ELISA was performed on stool samples using the lipocalin-2/NGAL DuoSet

ELISA R&D system (Bio-Techne, Minneapolis, MN, USA; catalog no. DY1857) according to

the manufacturer's instructions and samples were prepared as previously described[31]. MCPT1

concentration of serum samples was determined using the MCPT-1 (mMCP-1) Mouse

Uncoated ELISA Kit from Invitrogen (Waltham, MA, United States, catalog. no. 88-7503-88)

according to manufacturer's instructions. Serum ELISA for total IgG1 and IgE were performed

by coating plates using 0.5 μg/μl rat α mouse IgE purified UNLB (Imtec Diagnostics, 1130-

01) or rat α-mouse IgG1 purified UNLB (Imtec Diagnostics, Ardmore, OK, United States;

catalog no. 1144-01) capture antibody diluted in 0.05 M carbonate/bicarbonate buffer (pH 9.6).

Washing steps were performed using 1% Tween, 154 mM NaCl, 10 mM Trizma Base.

Blocking of the plates was performed using 1% w/v of BSA in a TBS buffer (15 mM Trizma-
acetate, 136 mM NaCl & 2 mM KCl). Mouse IgG1 Isotype Control UNLB (Imtec Diagnostics,
Ardmore, OK, United States; catalog no 0102-01) or Mouse IgE Isotype Control (UNLB),
Southern Biotech (Imtec Diagnostics, Ardmore, OK, United States; catalog no 0114-01) served
as controls. Serum samples were two-fold diluted using the aforementioned TBS buffer with
0.1% w/v Tween-20 and 1% BSA. The secondary antibody was Goat anti-mouse IgG1-AP,
Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog no 1071-04) or Goat
anti-mouse IgE-AP, Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog
no 1110-04), respectively. The substrate solution consisting of 0.5 mg/ml phosphate substrate
(Sigma-Aldrich, St. Louis, MO, United States; catalog no. S0642-200 TAB) in 1 mM AMP
and 0.1 mM $MgCl_2 \cdot 6H_2O$ and plates were read at 405 nm using a SpectraMax ABS PLUS
spectrophotometer (Molecular Devices, San Jose, CA, United States). A detailed protocol can
be found in the supplementary methods section. Parasite specific IgG1 and IgG2c ELISA was
performed as previously described[37].
215

216 ***LEGENDplex bead-based immunoassay***
217 The LEGENDplex assay (BioLegend, San Diego, CA, USA) was performed on serum samples
218 or colon tissue supernatant according to manufacturers' instruction.
219

220 ***RNA extraction and RT-qPCR from mesenteric lymph nodes, colonic tissue***
221 A detailed protocol can be found in the supplemental methods section.
222

***RNA-Seq***
RNA extracted from samples at 30 dpi were utilized for performing RNA-Seq. Illumina
Stranded Total RNA Prep with Ribo-Zero Plus was used to prepare the RNA sequencing library
according to the reference guide's instructions. Sequencing was performed using NovaSeq
6000 SP Reagent Kit v1.5 (Illumina, San Diego, CA, USA) on an Illumina NovaSeq 6000
system.
229

***Flow cytometry analysis of lamina propria immune cells***
Colonic immune cells were isolated using a Lamina Propria Dissociation Kit (Miltenyi Biotec,
Bergisch Gladbach, Germany) on the GentleMACS Dissociator (Miltenyi Biotec), following
manufacturer's instructions. Cells were washed, counted, and stained as previously described[38].
Samples were acquired on a NovoCyte Quanteon flow cytometer (ACEA Biosciences Inc., San

Diego, CA, USA). Raw fcs files were analyzed in FlowJo version 10.8.1 as previously
described[38].

***Statistical analyses***

Statistical analysis was performed using Prism 9.2.0. (GraphPad Software, Inc., San Diego,

CA, USA). Error bars in all figures represent SEM. Unless specified in the figure legend, a

two-way analysis of variance (ANOVA) was performed in Prism and *p* values were adjusted

with the original FDR method of Benjamini and Hochberg. P values lower than 0.1 are reported

in the figures.

**Results**

***Dietary fibers tune the worm infection dynamics***

We colonized 22-weeks-old germ-free, C57BL/6 mice with a 14-member synthetic human

microbiota (14SM), fed them a fiber-rich (FR) or a fiber-free (FF) diet, and then infected them

with a high dose of *T. muris* consisting of approximately 450 eggs from the same egg

suspension **(Fig. 1A)**. Throughout the experiment, none of the mice exhibited any obvious

physiological abnormalities. While FF-fed mice were overall heavier compared to the FR-fed

mice, the weights of both groups remained largely stable over time **(Fig. S1A)**. In line with our

published work[31,39], the FF diet allowed proliferation of two dominant mucin-degrading

bacteria *Akkermansia muciniphila* and *Bacteroides caccae* and reduction of two prominent

fiber-degrading bacteria *Eubacterium rectale* and *Bacteroides ovatus* **(Fig. 1B)**. Overall,

colonization with the synthetic human microbiota permitted establishment of a low-level *T.*

*muris* infection. Interestingly, although the worm burdens were statistically similar between

FR and FF groups on 20 and 25 dpi, subsequent infection dynamics significantly differed

between both groups, with FF-fed mice clearing the pathogen by 30 dpi and FR-fed mice

remaining chronically infected **(Fig. 1C)**.

In order to investigate whether the two distinct diets and/or the skewed gut microbiome

composition toward mucin polymer degradation caused an altered baseline inflammation prior

to infection with *T. muris*, we determined fecal lipocalin-2 (LCN-2) levels, which serves as a

neutrophilic marker for low-grade inflammation[40]. There were no significant differences

between FR and FF groups at day 0, however, after infection, the levels of LCN-2 were

significantly higher in the FR group **(Fig. 1D)**.

### *Diet-driven* T. muris *chronic infection or clearance are supported by Th1 or Th2 immune responses*

Chronic *T. muris* infection has been associated with Th1 inflammation while clearance of the worm is usually attributed to a strong Th2 response[2]. To better understand the change in host immunological responses that drive differential *T. muris* infection dynamics, we analyzed the host colonic gene expression at 30 dpi **(Fig. 2A, B, Fig. S1B, C and Table S1)**. We found 147 and 184 genes upregulated in the FR- and FF-fed mice, respectively **(Fig. 2A and Tables S2, S3)**. Ingenuity Pathway Analysis (IPA) revealed that both the interferon signaling and Th1 pathways were upregulated in the FR-fed mice **(Fig. 2B)**, which was consistent with the failure to successfully clear the parasite among FR-fed mice **(Fig. 1C)**. In accordance with the upregulation of the interferon signaling and Th1 pathways, related disease pathways, such as osteoarthritis, systemic lupus erythematosus and even neuroinflammation, were also upregulated as their gene expression profiles overlap with those of a Th1 response **(Fig. 2B)**. Similarly, the Th1 response is potentially linked to the observed increase in antimicrobial responses that results in the activation of cytotoxic T cells, lymphopoiesis, increased apoptosis of macrophages and a reduced replication of viruses **(Fig. S1B)**. In contrast to this, the increased MSP-RON signaling in macrophage pathway hints at a Th2 response under FF conditions **(Fig. 2B),** as an activation of the MSP-RON pathway causes M2 macrophage polarization[41], which is associated with a Th2-type immune response against *T. muris*[42,43]. Furthermore, the upregulation of the white adipose tissue browning pathway is also in line with this observation, as M2 macrophage differentiation has been linked to adaptive thermogenesis[44].

In order to further support conclusions from our RNA-Seq data, we used RT-qPCR to investigate transcript levels of cytokines in mesenteric lymph nodes (MLNs) and colon **(Fig. 2C, D, Table S4)**. In accordance with the global transcriptome analysis **(Fig. 2A, B)**, the FR-fed mice exhibited a Th1-type response in the MLN, characterized by a peak of *Ifng*, *Tnfa* and *Il1b* transcripts at 25 dpi and a continuously increasing expression of the Th1 transcription factor *Tbet* **(Fig. 2C and Fig. S1D)**. This Th1-type response was also observed in the colon with an increasing expression of *Ifng* over time and a peak of *Tnfa* expression at 25 dpi **(Fig 2D)**, as well as higher titers of IFNɣ, TNFα and IL-6 in the serum of FR-fed mice at 30 dpi **(Fig. 2E)**. Despite an initial induction of the anti-parasitic Th2-type response in the MLN of FR-fed mice at 15 dpi, *Il13* and *Il4* transcript levels decreased by 20 dpi, consistent with the

simultaneous increased expression of the counteracting Th1-type cytokines **(Fig. 2C)**. By
contrast, transcript levels of the Th2-type cytokines IL-4, IL-5 and IL-13 and the Th2
transcription factor GATA-3 were upregulated over time in either the MLN or colons of FF-
fed mice **(Fig. 2D and S1D)**. Further supporting the efficient anti-parasitic response in FF-fed
mice, the transcript levels of *Muc2*, the major component of the intestinal mucus, was strongly
upregulated in FF-fed mice by 25 and 30 dpi, while remaining low in FR-fed mice **(Fig. 2D)**.
308
Assessment of serum concentrations of total IgE and IgG1, both of which are reflective of the
protective Th2-type responses against helminths, resulted in no significant changes, albeit an
increased trend was observed for both antibodies at 30 dpi  **(Fig. S1E)**. We then assessed
parasite-specific IgG1 and IgG2, characteristic of Th2 and Th1 responses, respectively[45].
Chronic *T. muris* infections are usually characterized by significantly elevated serum levels of
parasite-specific IgG2a/c and a low increase in parasite-specific IgG1[45]. In contrast, protection
against *T. muris* is characterized by high parasite-specific IgG1 levels, but a lack of IgG2a/c[45].
In accordance with the observed worm burdens **(Fig. 1C)**, at 30 dpi, we saw an increase in
specific IgG1 in the serum of mice fed the FF diet, while we saw an increase in specific IgG2a/c
in mice fed the FR diet **(Fig. 2F)**. Furthermore, serum MCPT1—a proxy for IgE-induced mast
cell degranulation[46]—increased at 30 dpi in the FF-fed group **(Fig. 2G)**. These data are in line
with an increased tendency of total IgE in the FF-fed group at 30 dpi, and further support the
elevated anti-helminth type 2 immunity under fiber deprivation.
322
Overall, our data support that worm infection in mice hosting the 14SM and being fed fibers,
induce a Th1-type immune response that is driven by an overall increase in IFNγ. On the other
hand, in absence of dietary fibers, a similar infectious dose of *T.muris*, induces a Th2-type
immune response, resulting in the successful clearance of the worm. We have previously shown
that fiber deprivation leads to excessive gut microbial mucin foraging activity[30–32]. Since mucus
secretion is essential to the anti-parasitic immunity, and as the gut microbiome is intimately
connected to changes in the diet and the host immune responses, we hypothesized that mucin-
degrading bacteria promote anti-parasitic immunity.
331
***Mucin-degrading bacteria and absence of fiber promote anti-parasitic immunity***
To study the causal mechanism of mucin-degrading bacteria in modulating immunity to
helminth infection, we next infected 8-to-12-weeks-old mice, both male and female, colonized
with a 10-member synthetic microbiota (10SM) consisting of the 14SM minus the 4 previously

characterized mucin-degrading bacteria[31]. In this experiment, 14SM-colonized mice had lower

worm burdens compared to the previous experiment (maximum 19 worms versus 53,

respectively) (**Fig. 1C and 3A**), and no difference was detected between males and females

(data not shown). However, in line with the previous experiment (**Fig. 1C**), the mice fed the

FR diet had a higher worm burden than mice on the FF diet which completely cleared the

worms by 30 dpi (**Fig. 3A**). The 10SM FR-fed mice harbored worms at all three time points,

whereas FF-fed mice presented with few or no worms at any time point (**Fig. 3A**). Nevertheless,

all mice developed parasite-specific IgG, confirming that they were properly infected (**Fig.**

**3B**). Interestingly, mice fed the FR diet and devoid of mucin-degrading bacteria (10SM FR)

had a higher worm burden by 30 dpi compared to 14SM-colonized FR mice (**Fig. 3A**), while

they were similar between both groups on 20 and 25 dpi (**Fig. S2A**). Additionally, among both

FF groups at 30 dpi, 10SM-colonized mice had a higher titer of serum specific IgG1 (**Fig. 3B).**

On the other hand, among both FR groups at 30 dpi, 10SM-colonized mice had higher amounts

of fecal lipocalin-2 (**Fig. S2B**). These data suggest that mucin-degrading bacteria limit worm

infection by altering anti-parasitic immunity.

To better assess the anti-parasitic immune response, we analyzed the immune profiles of

mesenteric lymph nodes (MLN) and colons at 20, 25 and 30 dpi. Regardless of bacterial

colonization, flow cytometry analysis of immune cell populations in the MLN and colons

revealed a higher proportion of Th1 cells in FR-fed mice, in contrast to more Th2 (both MLN

and colon) and Treg (colon only) cells in the FF-fed mice (**Fig. 3C, D and S2C**). This diet

effect was further supported by transcript and protein expression profiles of Th1-related

cytokines (IFNγ, TNFα and IL-1β) in the colons of FR-fed mice and more Th2-related

cytokines (IL4, IL-5, IL-9 and IL-13) and MCPT1 in the colons of FF-fed mice (**Fig. 3E and**

**S3**).

Among FR-fed mice, 14SM-colonized mice had lower expression levels of Th1-related

cytokine transcripts *Ifng*, *Tnfa* and *Il1b* than 10SM-colonized mice at 30 dpi (**Fig. 3E**).

Although these trends were not always consistent at the protein level (**Fig. S3B**), they are

perfectly in line with the higher worm burden in 10SM FR mice (**Fig. 3A**), corroborating that

the presence of mucin-degrading bacteria further modulates intensity of the anti-parasitic

immune response. Among FF-fed mice, 14SM colonization tended to increase the proportion

of Th2 cells in the MLN at 25 dpi but then reduced it at 30 dpi in both the MLN and colon (**Fig.**

**3C, D**). Supporting an earlier T cell response in 14SM-colonized mice, the colonic transcript

levels of the growth factor IL-2 and the Type 2 promoting cytokines IL-25 and IL-33 were
already higher at 20 dpi in FF-fed, 14SM-colonized mice compared to FF-fed, 10SM-colonized
mice (**Fig. S3A**).

Intriguingly, FF-fed, 10SM-colonized mice had increased proportions of Tregs, ILC1 and ILC3
in the colon at 25 dpi (**Fig. 3D and S2C**), which may impair a timely Th2-driven anti-parasitic
response[47]. Further confirming an impaired anti-parasitic response in FF-fed, 10SM-colonized
mice, the increased proportion of colonic Th2 and ILC2 at 30 dpi was poorly reflected by IL-
5 and IL-13 cytokine production (**Fig. 3D, S2C and S3B**). Consistently, *Muc2* transcript was
strongly increased in FF-fed, 14SM-colonized mice but not in 10SM-colonized mice (**Fig. 3E**).
Additionally, at 25 dpi, 14SM-colonized FF-fed colons also produced more IL-10 and Th17-
related cytokines (IL17-A, IL-17F and IL-22) compared to all other groups (**Fig. S3B**),
suggesting a more efficient barrier healing and protective process in the presence of mucin-
degrading bacteria and in the absence of dietary fibers. These results strongly support that
mucin-degrading bacteria and absence of fiber act together to promote an anti-parasitic immune
response.

***Mucin-degrading bacteria promote mucus production under fiber deprivation***
As a result of a strong type-2 immune response, mucus production is a key mechanism of host
defense during parasite infection and can be regulated at the transcriptional, translational and
post-translational levels. From 15 to 30 dpi, both FR- and FF-fed 14SM-colonized mice had
moderately increased numbers of goblet cells per crypt, but no difference was seen between
dietary groups (**Fig. 4A**). Transcriptomic analysis of colon tissues at 30 dpi revealed that the
expression of numerous genes involved in mucus production and glycosylation pathways were
upregulated in FF-fed mice compared to FR-fed mice (**Fig. 4B**). Moreover, targeted qPCR
revealed that, in addition to *Muc2* (**Fig. 2C and 3E**), *Muc5ac* expression was higher in 14SM-
colonized FF-fed mice at 30 dpi, compared to FR-fed mice or FF-fed, 10SM-colonized mice
(**Fig. 4C**). Similarly, the combination of both mucin-degrading bacteria and fiber deprivation
induced the expression of *C1galt1*, *B3gnt6*, *Fut2*, *Galnt3* and *Gal3st2*, which encode key
enzymes involved in mucin glycosylation (**Fig. 4C, Table S4**). These results suggest that
mucin-degrading bacteria support production of mucus under fiber deprivation, and possibly
ensure its proper glycosylation so that this new mucus can be secreted in order to expulse the
worms.

*Increased mucin–microbiome interactions promote worm clearance*

To better understand the role of specific microbiota members in the course of the parasite

infection, we first analyzed the composition of the fecal microbiota (**Fig. S4**). Surprisingly and

despite the reproducible phenotypes showed above, the 14SM microbiota composition at the

time of infection was different between the first experiment (**Figs. 1, 2**) and the second

experiment (**Fig. 3**) (**Fig. S4, 5**, **Supplementary results**). The analysis of the relative

abundance over time for individual bacteria revealed different profiles between groups with

little reproducibility between independent experiments, except for *A. muciniphila* and *B.*

*caccae* (**Fig. 5A, B**, **S4B and S6, 7**). Despite different initial relative abundances between the

two experiments, these 2 mucin degraders expanded significantly and reproducibly from 14 to

20 dpi (**Fig. 5A, B**), a timepoint when mice usually start to expel their worms.

Consistent with the evolving mucus production and microbiota composition, the enzymatic

activities targeting either mucin glycans only (fucosidase and *N*-acetyl-glucosaminidase) or

both mucin and dietary glycans (glucosidase) evolved over time during *T. muris* infection (**Fig.**

**5C, Tables S5–7**). The activities of mucin-targeting enzymes were higher in 14SM-colonized

and FF-fed mice (**Fig. 5C**), and correlated negatively with the worm burden (**Fig. 5D**), further

supporting a role for mucin-degrading bacteria in facilitating worm expulsion. Of note,

fucosidase activity in 14SM-colonized FF-fed mice (**Fig. 5C**) followed the expression profile

of the fucosyltransferase transcript *Fut2* (**Fig. 4C**), decreasing at 25 dpi to increase again by 30

dpi, supporting a microbial enzyme activity that may be driven by the glycosylation profile of

the mucins. Finally, among the four mucin-degrading strains, the relative abundance of *B.*

*caccae* and *A. muciniphila* correlated positively with *N*-acetyl-glucosaminidase activity (**Fig.**

**5E and Table S6**), while the relative abundance of *B. thetaiotaomicron* and *B. intestinihominis*

correlated negatively with *N*-acetyl-glucosaminidase and fucosidase activities, respectively

(**Fig. 5F**, **Tables S5, 6**).

**Discussion**

Helminth infection models have been profoundly utilized to elucidate the underlying host

responses for the past several decades, yet open questions remain about the mechanisms of how

the intricate interactions between the gut microbiome and GI mucus affect worm infection

dynamics including the host immune responses. Here, we shed insight into the role of bacterial

mucin foraging in altering the host immune responses during the infection with *T. muris*. Using

a functionally characterized synthetic human microbiota that allows either an ablation of
microbial mucin foraging or increase in microbial mucin foraging through reduced fiber
consumption, our study supports a model in which the mucin-degrading bacteria promote anti-
parasitic immunity, that can be further enhanced by reduced dietary fiber **(Fig. 6)**. Our results
identify mucin-degrading bacteria as an innovative link in the gut microbiome–parasite axis,
suggesting a causal mechanism through which the anti-parasitic immunity is a modulated.

It is well known that bacteria are important for *T. muris* worm hatching[8,9], although the specific
bacterial members involved are not well defined and it is possible that the selected bacteria in
our synthetic human microbiota might not be ideal for inducing egg hatching for mouse
*Trichuris*, resulting in a lower level of infection of *T. muris* that provides a chronic low-level
infection characteristic of infections under natural conditions. Thus, our tractable synthetic
microbiota, that facilitates functional interpretations, is a good model to understand the
mechanisms linking the worm infection and mucin–microbiome interactions, which would be
difficult to decipher in such great detail using a complex, native microbiota. Although the fiber-
deficient mice, lacking the mucin-degrading bacteria, contained none to very few worms, these
mice still mounted a robust anti-parasitic immune response **(Fig. 3B)**. Differences in the
parasitic load observed between the two experiments **(Figs. 1, 2 vs. Fig. 3)** could be due to
several factors such as the age of the mice used for these experiments (22 weeks vs. 8–12
weeks) or in some other unknown factors, yet both experiments provided similar key outcomes.
We noticed inconsistency between some readouts e.g., immune cell profiles (MLNs and
colons) and serum cytokines at protein level, which is not unexpected for such detailed,
complex longitudinal data sets. Nevertheless, overall, the data from our gnotobiotic mouse
models allow us to adequately support our main conclusions.

Our study shows an IFNγ-dominated Th1 response, along with a chronic *T. muris* infection,
when mice were fed a normal chow (which we called a fiber-rich (FR) diet), in contrast to a
Th2 response and worm clearance under fiber deprivation **(Figs. 1–3 and 6)**. Two previous
studies[27,48] observed a similar Th1 response when specific pathogen free mice—containing
their native, complex microbiotas—were fed diets containing purified 10% inulin or dried
chicory leaves. Nevertheless, these studies did not provide an explanation for the underlying
mechanisms. Using our 14SM model, we offer an explanation that the mucin–microbiome
interactions driven by dietary changes alter the host immune responses and the worm infection
dynamics **(Fig. 6)**. Our study lays groundwork for investigating future molecular mechanisms

of how mucolytic bacteria drive specific alterations in mucus glycosylation and how such

alterations impact host immune response to worm infection.

*T. muris* has been shown to broadly alter the host's gut microbiome composition[23–28]. Here,

repeating the experiment in several batches of animals revealed different, most likely age-

dependent, profiles of microbiota composition over the course of the infection (**Figs. S4–7**).

Nevertheless, the infectious phenotype was reproducible as well as a decline of *A. muciniphila*

at 15 dpi followed by an expansion of both *A. muciniphila* and *B. caccae* at 20 dpi in fiber-

deprived mice (**Fig. 5A, B**). The role of mucin-degrading bacteria such as *A. muciniphila* and

*B. caccae* in our model is evident because colonization with a synthetic microbiota without

mucin-degrading bacteria prevented the fiber deprivation-induced Th2 immunity (**Fig. 3**).

In contrast to the observations in *Muc2$^{-/-}$* mice[29], in our model, increased microbial mucin

foraging under fiber-deprived conditions did not delay the parasite clearance, but instead

promoted the clearance. These results are not unexpected as genetic ablation of Muc2 largely

destabilizes the mucus structure, whereas the increased microbial mucin foraging seems to

either resist transition to a Th1 response and aids in a Th2 response. We could not detect an

increase in goblet cell numbers/mucin production associated with this (**Fig. 4A**). Nevertheless,

we observed increased transcription of genes involved in mucin glycosylation (**Fig. 4B, C**),

which has been shown previously to alter susceptibility to *T. muris*[49]. Considering that mucin

glycosylations are built-up by the host glycosyltransferases and targeted by the microbial

enzymes[50], our results suggest that fiber deprivation promotes changes in mucin glycosylation

that facilitates mucus degradation by the fiber-deprived microbiota and aids in clearance of *T.*

*muris*.

Given the high morbidity associated with the parasitic infections in humans[51], our study has

generated important prerequisite knowledge for clinical settings: 1) for treating parasitic

infections – in the absence of co-infections, periodic fiber deprivation or fasting to boost

microbial mucin foraging could be potentially employed in the clinic to manage worm

infections; and 2) for harnessing the immunomodulatory properties of helminths for their

potential therapeutic use in inflammatory disorders[52] or allograft rejection[53]. Further research

is required about how the model proposed in our study (**Fig. 6)** could be translated into the

clinic.

**References**

1.  Douglas B, Oyesola O, Cooper MM, et al. Immune System Investigation Using
Parasitic Helminths. Annu Rev Immunol 2021;39:639–665.

2.  Klementowicz JE, Travis MA, Grencis RK. Trichuris muris: A model of
gastrointestinal parasite infection. Semin Immunopathol 2012;34:815–828.

3.  Yousefi Y, Haq S, Banskota S, et al. Trichuris muris model: Role in understanding
intestinal immune response, inflammation and host defense. Pathogens 2021;10.

4.  Colombo SAP, Grencis RK. Immunity to Soil-Transmitted Helminths: Evidence From
the Field and Laboratory Models. Front Immunol 2020;11.

5.  Zaiss MM, Harris NL. Interactions between the intestinal microbiome and helminth
parasites. Parasite Immunol 2016;38:5–11.

6.  Brosschot TP, Reynolds LA. The impact of a helminth-modified microbiome on host
immunity review-article. Mucosal Immunol 2018;11:1039–1046.

7.  Jaenike J, Unckless R, Cockburn SN, et al. Adaptation via symbiosis: Recent spread of
a drosophila defensive symbiont. Science (80- ) 2010;329:212–215.

8.  Hayes KS, Bancroft AJ, Goldrick M, et al. Exploitation of the intestinal microflora by
the parasitic nematode trichuris muris. Science (80- ) 2010;328:1391–1394.

9.  White EC, Houlden A, Bancroft AJ, et al. Manipulation of host and parasite
microbiotas: Survival strategies during chronic nematode infection. Sci Adv 2018;4.

10. Robertson A, Sall J, Venzon M, et al. Bacterial contact induces polar plug
disintegration to mediate whipworm egg hatching. PLOS Pathog 2023;19:e1011647.

11. Hewitson JP, Harcus Y, Murray J, et al. Proteomic analysis of secretory products from
the model gastrointestinal nematode Heligmosomoides polygyrus reveals dominance
of Venom Allergen-Like (VAL) proteins. J Proteomics 2011;74:1573–1594.

12. Abner SR, Parthasarathy G, Hill DE, et al. Trichuris suis: Detection of antibacterial
activity in excretory-secretory products from adults. Exp Parasitol 2001;99:26–36.

13. Su L, Su CW, Qi Y, et al. Coinfection with an intestinal helminth impairs host innate
immunity against Salmonella enterica serovar typhimurium and exacerbates intestinal
inflammation in mice. Infect Immun 2014;82:3855–3866.

14. Fricke WF, Song Y, Wang AJ, et al. Type 2 immunity-dependent reduction of
segmented filamentous bacteria in mice infected with the helminthic parasite
Nippostrongylus brasiliensis. Microbiome 2015;3:40.

15. Forman RA, DeSchoolmeester ML, Hurst RJM, et al. The Goblet Cell Is the Cellular

Source of the Anti-Microbial Angiogenin 4 in the Large Intestine Post Trichuris muris
Infection. PLoS One 2012;7.

16.  Shea-Donohue T, Sullivan C, Finkelman FD, et al. The Role of IL-4 in
Heligmosomoides polygyrus -Induced Alterations in Murine Intestinal Epithelial Cell
Function. J Immunol 2001;167:2234–2239.

17.  Harris JB, Podolsky MJ, Bhuiyan TR, et al. Immunologic responses to Vibrio cholerae
in patients co-infected with intestinal parasites in Bangladesh. PLoS Negl Trop Dis
2009;3.

18.  Kosik-Bogacka DI, Wojtkowiak-Giera A, Kolasa A, et al. Hymenolepis diminuta:
Analysis of the expression of Toll-like receptor genes and protein (TLR3 and TLR9) in
the small and large intestines of rats. Exp Parasitol 2014;145:61–67.

19.  Zakeri A, Borji H, Haghparast A. Interaction Between Helminths and Toll-Like
Receptors: Possibilities and Potentials for Asthma Therapy. Int Rev Immunol
2016;35:219–248.

20.  Marillier RG, Michels C, Smith EM, et al. IL-4/IL-13 independent goblet cell
hyperplasia in experimental helminth infections. BMC Immunol 2008;9.

21.  Hasnain SZ, McGuckin MA, Grencis RK, et al. Serine Protease(s) Secreted by the
Nematode Trichuris muris Degrade the Mucus Barrier. PLoS Negl Trop Dis 2012;6.

22.  Hasnain SZ, Evans CM, Roy M, et al. Muc5ac: A critical component mediating the
rejection of enteric nematodes. J Exp Med 2011;208:893–900.

23.  Li RW, Wu S, Li W, et al. Alterations in the porcine colon microbiota induced by the
gastrointestinal nematode Trichuris suis. Infect Immun 2012;80:2150–2157.

24.  Broadhurst MJ, Ardeshir A, Kanwar B, et al. Therapeutic Helminth Infection of
Macaques with Idiopathic Chronic Diarrhea Alters the Inflammatory Signature and
Mucosal Microbiota of the Colon. PLoS Pathog 2012;8.

25.  Holm JB, Sorobetea D, Kiilerich P, et al. Chronic Trichuris muris infection decreases
diversity of the intestinal microbiota and concomitantly increases the abundance of
lactobacilli. PLoS One 2015;10.

26.  Houlden A, Hayes KS, Bancroft AJ, et al. Chronic Trichuris muris infection in
C57BL/6 mice causes significant changes in host microbiota and metabolome: Effects
reversed by pathogen clearance. PLoS One 2015;10.

27.  Myhill LJ, Stolzenbach S, Mejer H, et al. Fermentable Dietary Fiber Promotes
Helminth Infection and Exacerbates Host Inflammatory Responses. J Immunol
2020;204:3042–3055.

28.    Rosa BA, Snowden C, Martin J, et al. Whipworm-Associated Intestinal Microbiome
572           Members Consistent Across Both Human and Mouse Hosts. Front Cell Infect
573           Microbiol 2021;11.

574    29.    Hasnain SZ, Wang H, Ghia JE, et al. Mucin Gene Deficiency in Mice Impairs Host
575           Resistance to an Enteric Parasitic Infection. Gastroenterology 2010;138:1763-1771.e5.

576    30.    Neumann M, Steimle A, Grant ET, et al. Deprivation of dietary fiber in specific-
577           pathogen-free mice promotes susceptibility to the intestinal mucosal pathogen
578           Citrobacter rodentium. Gut Microbes 2021;13.

579    31.    Desai MS, Seekatz AM, Koropatkin NM, et al. A Dietary Fiber-Deprived Gut
580           Microbiota Degrades the Colonic Mucus Barrier and Enhances Pathogen
581           Susceptibility. Cell 2016;167:1339-1353.e21.

582    32.    Parrish A, Boudaud M, Grant ET, et al. Akkermansia muciniphila exacerbates food
583           allergy in fibre-deprived mice. Nat Microbiol 2023 810 2023;8:1863–1879.

584    33.    Martens EC, Neumann M, Desai MS. Interactions of commensal and pathogenic
585           microorganisms with the intestinal mucosal barrier. Nat Rev Microbiol 2018;16:457–
586           470.

587    34.    Parrish A, Boudaud M, Kuehn A, et al. Intestinal mucus barrier: a missing piece of the
588           puzzle in food allergy. Trends Mol Med 2022;28:36–50.

589    35.    Steimle A, Sciscio A De, Neumann M, et al. Constructing a gnotobiotic mouse model
590           with a synthetic human gut microbiome to study host–microbe cross talk. STAR
591           Protoc 2021;2:100607.

592    36.    Wakelin D. Acquired immunity to Trichuris muris in the albino laboratory mouse.
593           Parasitology 1967;57:515–524.

594    37.    Hayes KS, Cliffe LJ, Bancroft AJ, et al. Chronic Trichuris muris infection causes
595           neoplastic change in the intestine and exacerbates tumour formation in APC min/+
596           mice. PLoS Negl Trop Dis 2017;11.

597    38.    Grant ET, Boudaud M, Muller A, et al. Maternal diet and gut microbiome composition
598           modulate early-life immune development. EMBO Mol Med 2023;15.

599    39.    Wolter M, Steimle A, Parrish A, et al. Dietary Modulation Alters Susceptibility to
600           Listeria monocytogenes and Salmonella Typhimurium with or without a Gut
601           Microbiota. mSystems 2021;6:e00717-21.

602    40.    Chassaing B, Srinivasan G, Delgado MA, et al. Fecal Lipocalin 2, a Sensitive and
603           Broadly Dynamic Non-Invasive Biomarker for Intestinal Inflammation Bereswill S,
604           ed. PLoS One 2012;7:e44328.

41.  Chaudhuri A. Regulation of macrophage polarization by RON receptor tyrosine kinase
signaling. Front Immunol 2014;5.

42.  Forman R, Logunova L, Smith H, et al. Trichuris muris infection drives cell-intrinsic
IL4R alpha independent colonic RELMα+ macrophages. PLoS Pathog 2021;17.

43.  Sorobetea D, Svensson-Frej M, Grencis R. Immunity to gastrointestinal nematode
infections. Mucosal Immunol 2018;11:304–315.

44.  Hallowell RW, Collins SL, Craig JM, et al. MTORC2 signalling regulates M2
macrophage differentiation in response to helminth infection and adaptive
thermogenesis. Nat Commun 2017;8.

45.  Else KJ, Entwistle GM, Grencis RK. Correlations between worm burden and markers
of Th1 and Th2 cell subset induction in an inbred strain of mouse infected with
Trichuris muris. Parasite Immunol 1993;15:595–600.

46.  Kasakura K, Nagata K, Miura R, et al. Cooperative Regulation of the Mucosal Mast
Cell–Specific Protease Genes Mcpt1 and Mcpt2 by GATA and Smad Transcription
Factors. J Immunol 2020;204:1641–1649.

47.  Hepworth MR, Monticelli LA, Fung TC, et al. Innate lymphoid cells regulate CD4+ T-
cell responses to intestinal commensal bacteria. Nat 2013 4987452 2013;498:113–117.

48.  Valente AH, Jensen KMR, Myhill LJ, et al. Dietary non-starch polysaccharides impair
immunity to enteric nematode infection. BMC Biol 2023;21:1–16.

49.  Hasnain SZ, Dawson PA, Lourie R, et al. Immune-driven alterations in mucin
sulphation is an important mediator of Trichuris muris helminth expulsion. PLoS
Pathog 2017;13.

50.  Arike L, Holmén-Larsson J, Hansson GC. Intestinal Muc2 mucin O-glycosylation is
affected by microbiota and regulated by differential expression of glycosyltranferases.
Glycobiology 2017;27:318–328.

51.  World Health Organization. Global Health Estimates 2020: Disease burden by Cause,
Age, Sex, by Country and by Region, 2000-2019. 2020.

52.  Helmby H. Human helminth therapy to treat inflammatory disorders- where do we
stand? BMC Immunol 2015;16.

53.  Kiss M, Burns H, Donnelly S, et al. Effectiveness of Helminth Therapy in the
Prevention of Allograft Rejection: A Systematic Review of Allogeneic
Transplantation. Front Immunol 2020;11.

**Figure legends**

**Figure 1. Diet drives changes in mucin-degrading bacteria expansion and alters worm infection.**

(A) Experimental timeline.

(B) Relative bacterial abundance at the time of infection (day 0), determined by qPCR on fecal DNA. Red latin crosses denote known mucin-degrading bacteria.

(C) Worm burdens assessed at 20, 25 and 30 dpi. Each dot represents a mouse (n=4–5). Multiple Mann-Whitney test, *p* values adjusted with the FDR method of Benjamini and Hochberg.

(D) Fecal Lipocalin-2 (LCN-2) levels assessed by ELISA before (day 0) and after infection with *T. muris* (13, 15, 20, 23, 30 dpi) (n=4–5). *p* values indicate comparisons between FR and FF groups.

**Figure 2. Dietary fibers skew the anti-parasitic immune response towards Th1-type responses.**

(A, B) Analysis of data generated by RNA-seq of RNA extracted from colonic tissue at 30 dpi (n=4). (A) Volcano plot of the distribution of all differentially expressed genes. (B) Barplot showing significantly up- or downregulated host pathways ((Fiber Rich (FR)/Fiber Free (FF)) determined by Ingenuity Pathway Analysis (IPA). The higher the opacity of the bar, the more up- or downregulated the pathway.

(C, D) Relative expression of indicated transcripts in (C) mesenteric lymph nodes and (D) colonic tissue over time after *T. muris* infection (n=3–5); *p* values indicate comparisons between FR and FF groups.

(E) Serum cytokine concentrations at 30 dpi (n=4–5). Unpaired, two-tailed t-test.

(F) Parasite-specific IgG1 and IgG2a/c concentrations in mouse serum (n=4–5).

(G) Serum mast cell protease 1 (MCPT1) concentrations (n=2–5); *p* values indicate comparisons between FR and FF groups.

**Figure 3. Mucin-degrading bacteria promote the anti-parasitic immunity.**

(A) Worm burdens assessed at 20, 25 and 30 dpi in 14SM- and 10SM-colonized mice fed a fiber-rich (FR) or a fiber-free (FF) diet (n=5–6).

(B) Parasite-specific IgG1 and IgG2a/c concentrations in mouse serum (n=5–6). *P* values indicate comparisons between FR and FF groups among 10SM- or 14SM-colonized mice.

(C, D) Proportion of indicated immune cell populations in (C) the mesenteric lymph nodes
and (D) colonic tissue (n=3–6).

(E) Relative expression of indicated transcripts in colonic tissue (n=3–6).

**Figure 4. Mucin-degrading bacteria promote mucus production under fiber deprivation.**
(A) Goblet cell counts per crypt of *T. muris* infected mice determined by histological
assessment of methacarn-fixed cecum sections (n=2–5).

(B) Relative gene expression (FF/FR) at 30 dpi of the 5 most significant activated pathways in
the colonic tissue under FF conditions (n=4 per group). Pathways are based on GO terms for
biological processes and REVIGO was used to reduce redundant terms.

(C) Relative expression of indicated transcripts in colonic tissue (n=2–6).

**Figure 5. Increased mucin–microbiome interactions promote worm clearance.**
(A, B) Relative abundance of *B. caccae* (A) and *A. muciniphila* (B) over the course of *T. muris*
infection in 14SM-colonized mice (n=2–6). Significance labels indicate comparisons between
FR versus FF groups (black), or between indicated timepoints among FF-fed mice of Fig. 1–2
(yellow), FR-fed mice of Fig. 3 (green) and FF-fed mice of Fig. 3 (red).

(C) Glycan-degrading enzyme activities of the gut microbiome determined by stool-based p-
nitrophenyl glycoside-based enzyme assays. α-fucosidase and β-N-acetyl-glucosaminidase are
key mucin-degrading enzymes, while β-glucosidase serves as a control for general glycan-
degrading activity (n=3–6). Significance labels indicate comparisons between [&]FR versus FF
groups among 14SM-colonized mice from Figs. 1, 2, *FR versus FF groups among 14SM-
colonized mice from Fig. 3, [$]FR versus FF groups 10SM-colonized mice from Fig.3, and
[#]10SM versus 14SM groups among FF-fed mice. [&/*/#]$p<0.05$; [&&/$$/##]$p<0.01$; ***$p<0.001$;
[&&&&/****/$$$$/####]$p<0.0001$.

(D, E) Spearman correlations between indicated bacteria relative abundance and (D) N-acetyl-
glucosaminidase activity (n = 41–70) or (E) fucosidase activity (n = 40).

(F) Spearman correlations between worm burdens and enzymatic activities (n=31).

**Figure 6. Model for the role of mucin-degrading bacteria in facilitating anti-parasitic**
**immunity and *T. muris* clearance.**
(1) Fiber deprivation promotes microbial mucus foraging and thinning, which may facilitate
hatching of the worm to the epithelium and its sensing by the host. (2) This sensing induces a
strong type-2 response, leading to mucus production and preventing the worm to stay and

develop. (3) Under fiber deprivation the worms reaching the epithelium promote an IFNg
response, which inhibits the anti-parasitic response and allows the worm to settle and grow.

# Figure 1. Diet drives changes in mucin-degrading bacteria and alters worm infection.

**A**

**Colonization**
◇ *Trichuris muris*
□ Germ-free (GF)
■ synthetic human microbiota (14SM or 10SM)

**Diet**
● Fiber rich (FR)
● Fiber free (FF)

**B**

Relative abundance (%)

- *Akkermansia muciniphila†*
- *Collinsella aerofaciens*
- *Desulfovibrio piger*
- *Escherichia coli*
- *Faecalibacterium prausnitzii*
- *Roseburia intestinalis*
- *Marvinbryantia formatexigens*
- *Clostridium symbiosum*
- *Eubacterium rectale*
- *Barnesiella intestinihominis†*
- *Bacteroides caccae†*
- *Bacteroides uniformis*
- *Bacteroides ovatus*
- *Bacteroides thetaiotaomicron†*

**C** Worm burden

0.047619

Worm count / Days post-infection

**D** Fecal Lipocalin-2

0.0410

ng / g of feces / Days post-infection

Fiber-rich / Fiber-free

# Figure 2. Dietary fiber skew the anti-parasitc immune response towards Th1-type responses.

# Figure3. Mucin-degrading bacteria promote the anti-parasitic immunity.

# Figure 4. Mucin-degrading bacteria mediate mucus production under fiber deprivation.

# Figure 5. Increased mucin–microbiome interactions promote worm clearance.

Fig. 6

Dietary
fiber

IFNγ
TNFα

Th1

IL-4
IL-5
IL-13

ILC2

Th2

Mast

IL-25
IL-33

Fiber-rich response

Fiber-free response

# Fig. S3