Increased gut microbial mucin foraging promotes clearance of a parasitic worm

Short title: *Nematodes and gut microbial mucin foraging*

Mathis Wolter[a,b], Erica T. Grant[a,b], Amy Parrish[a,b], Alessandro De Sciscio[a], Seona

Thompson[c,d], Marie Boudaud[a], Jean-Jacques Gerardy[e,f], Michel Mittelbronn[e,f,g,h,i,j], David J.

Thornton[c,d], Andrew J. Macpherson[k,l,m], Richard K. Grencis[c,d] and Mahesh S. Desai[a,n,*]

[a]*Department of Infection and Immunity, Luxembourg Institute of Health, Esch-sur-Alzette,*

Luxembourg

[b]*Faculty of Science, Technology and Medicine, University of Luxembourg, Esch-sur-Alzette,*

Luxembourg

[c]*Lydia Becker Institute for Immunology and Inflammation. Faculty of Biology, Medicine and*

Health, University of Manchester, Manchester, United Kingdom

[d]*Wellcome Centre for Cell Matrix Research, Faculty of Biology, Medicine and Health,*

University of Manchester, Manchester, United Kingdom

[e]*National Center of Pathology, Laboratoire National de Santé, Dudelange, Luxembourg*

[f]*Luxembourg Center of Neuropathology, Dudelange, Luxembourg*

[g]*Luxembourg Centre for Systems Biomedecine, University of Luxembourg, Esch-sur-Alzette,*

Luxembourg

[h]*Department of Cancer Research, Luxembourg Institute of Health, Luxembourg, Luxembourg*

[i]*Faculty of Science, Technology and Medecine, University of Luxembourg, Esch-sur-Alzette*

[j]*Department of Life Science and Medicine, University of Luxembourg, Esch-sur-Alzette,*

Luxembourg

[k]*Department of Visceral Surgery and Medicine, Bern University Hospital, University of Bern,*

Bern, Switzerland

[l]*Maurice Müller Laboratories, Department for Biomedical Research, University of Bern,*

Bern, Switzerland

[m]*CMPG, Institute of Ecology and Evolution, University of Bern, Bern, Switzerland*

Odense Research Center for Anaphylaxis, Department of Dermatology and Allergy Center,

Odense University Hospital, University of Southern Denmark, Odense, Denmark

*Corresponding author: Mahesh S. Desai, mahesh.desai@lih.lu

Grant support

This work was supported by the following grants in the laboratory of M.S.D.: Luxembourg

National Research Fund (FNR) CORE grants (C15/BM/10318186 and C18/BM/12585940) to

M.S.D.; FNR PRIDE (17/11823097) and the Fondation du Pélican de Mie et Pierre Hippert-

Faber, under the aegis of the Fondation de Luxembourg grants to E.T.G.; FNR AFR individual

PhD fellowship to A.P. (11602973); and European Commission Horizon 2020 Marie

Skłodowska-Curie Actions individual fellowship to M.B. (897408). MM thanks the

Luxembourg National Research Fond (FNR) for the support (FNR PEARL P16/BM/11192868

grant). The Wellcome Trust supports work of R.K.G and S.T. (Z10661/Z/18/Z;

203128A/Z/16/Z) and D.J.T (203128A/Z/16/Z).

Correspondence

Mahesh S. Desai, PhD

Luxembourg Institute of Health

Department of Infection and Immunity

29, rue Henri Koch, L-4354 Esch-sur-Alzette

Luxembourg

Tel: +352 26970-389

Fax: +352 26970-390

Email: mahesh.desai@lih.lu

Disclosures

M.S.D. works as a consultant and an advisory board member at Theralution GmbH, Germany.

Author Contributions

M.S.D. supervised the study and obtained research funding; M.W., D.J.T., R.K.G. and M.S.D.

conceived and designed the study; M.W., E.T.G., A.P., A.D.S., and S.T. performed the

experiments; M.W., E.T.G., R.K.G., A.P. and M.B. analyzed the data; J.-J. G. and M.M.

prepared slides for histology; A.J.M. supplied germ-free mice; M.W. and M.S.D. primarily

wrote the manuscript with input from E.T.G and A.P. All authors reviewed and approved the
manuscript.
65
Data Transparency Statement
All data, study materials and analytical methods will be made available to other researchers
upon request.

Abstract

BACKGROUND & AIMS

Host-secreted gastrointestinal mucus plays a key role in the expulsion of intestinal nematode

parasites. A balance between mucin secretion by the host and the gut microbial mucin foraging

is essential to maintain the intestinal homeostasis, yet little is known about how changes in the

mucin–microbiome interactions affect worm infection. Here, we aimed to examine how

increased mucin foraging activity by the microbiome changes the course of the parasitic

infection by modulating the host immune responses.

METHODS

We employed a gnotobiotic mouse model containing a 14-member synthetic human gut

microbiota that facilitates functional interpretations, including diet-driven manipulation of the

microbiota toward mucin foraging. We infected the mice with a robust murine nematode,

Trichuris muris, that closely resembles human infection with *Trichuris trichiura*. We

investigated the temporal dynamics of worm infection including worm burden and the host

immune responses, and connected these readouts to the microbial changes and metabolic

activity toward mucin foraging.

RESULTS

The microbial mucin foraging was further increased during worm infection only in mice with

pre-enhanced mucin degrading capacity. The elevated mucin foraging coincides with a shift in

host immune responses from susceptible (chronic, Th1 type) to resistant (acute, Th2 type),

which promotes worm clearance. The relative abundances of mucin-generalist bacteria

drammatically increased during worm clearance, but not during worm retention.

CONCLUSIONS

These results point to a mechanism whereby skewing the metabolic activity of the microbiome

toward mucin glycoproteins promotes resistance to worm infection. Our study documents a

clinically-relevant, novel link in the microbiome–parasite–host immune axis, which is

important prerequisite knowledge in treating parasitic infections.

Keywords

Mucin; gut microbiome; nematode; *Trichuris muris*

Introduction

Chronic infections by intestinal parasitic worms (helminths) are common in animals and humans, especially in developing countries, where the infections are associated with high morbidity[1]. Due to their ability to subvert the host immune response, helminths make an invaluable model to study the host immunomodulatory mechanisms[2]. In this context, the murine parasite *Trichuris muris*—a relevant biological model for the closely related human parasite *Trichuris trichiura*—has been extensively investigated[3]. *T. muris* is an especially attractive model as either a susceptible (chronic, Th1 type) or resistant (acute, Th2 type) immune response is triggered, depending on the infectious dose[3,4]. Low-dose infections in mice, which are more reflective of the naturally-occurring chronic infections[5], are generally characterized by a Th1 response resulting in long-term worm infections and chronic colitis. On the other hand, high-dose infections, which are employed as a laboratory model, are generally characterized by an acute Th2 response leading to worm clearance and tissue repair[3].

Since they occupy the same host environment, it is no surprise that enteric parasites such as helminths strongly interact with the gut microbiome[6,7]. Although these interactions may lead to the suppression of either the gut microbiome or the helminth[6–8], in case of *T. muris*, bacterial–helminth crosstalk is essential for worm hatching, as the parasite eggs fail to hatch and establish in germ-free environments[9,10]. Once the worm has settled in the intestinal epithelium, the bacterial–helminth dynamics appear mainly unidirectional, with helminths altering the microbiome composition through various mechanisms, including through helminth excretory-secretory products or host anti-microbial peptides, released in response to the parasite[11–15]. Furthermore, the helminths alter the nutrients available to the microbiome, indirectly influencing which bacteria proliferate[16]. The effect of the infection on host immunology has an additional impact on the microbiome, as the worms or their metabolic byproducts can alter toll-like receptor (TLR) expression profiles[17,18] or responsiveness of host immune cells to TLR ligands[19] and interfere with local IgA-production[17]. While the gastrointestinal (GI) mucins are essential for worm clearance[20–22] and the impacts of the helminth infection on the microbiome during worm clearance are documented at the broader level of microbial phylogeny[23–28], questions remain regarding how the mucus–microbiome interactions affect worm infection and the underlying host immune responses.

Worm expulsion during an acute response is largely driven by goblet-cell secreted gastrointestinal mucin that is tightly linked to the Th2-type immune response[22]. Along these lines, mice lacking the *Muc2* gene, which encodes the major glycoprotein of the GI mucus,

show a delayed clearance of *T. muris*[29] and *Muc5ac* deficient mice completely fail to clear the

helminth[22]. In germ-free mice, the mucus layer expands upon exposure to gut bacteria, and the

colonic mucin secretion by goblet cells is regulated by the gut microbial activity[30,31]. The

balance between mucin secretion and its consumption by the gut microbes is critical for

intestinal homeostasis[32,33]. We have previously shown that a disrupted balance, for example,

excessive mucin foraging by a fiber-deprived microbiota leads to an increase in the gut

microbial mucin foraging activity[31,34].

Here, we hypothesized that excessive mucin foraging by the microbiota alters the

course of the parasite infection by modulating the immune response. To test this hypothesis,

we employed a tractable 14-member synthetic human microbiota (14SM) characterized for the

capability of individual members to degrade various glycans including mucin

oligosaccharides[31,35]. In a longitudinal experimental setup, using C57BL/6 ex-germ-free mice

with a homeostatic mucin-foraging 14SM, we induced a chronic *T. muris* infection.

Simultaneously, using fiber-deprived diet as a precursor, we skewed 14SM toward excessive

mucin foraging. We assessed the worm burden and immune outcome of infection. Moreover,

our 14SM model combined with a longitudinal infection setup enabled us to understand

specific microbial functional changes along the course of the infection in both homeostatic

14SM and 14SM skewed toward mucin foraging. Our findings suggest a mechanism through

which the increased gut microbial mucin foraging promotes worm clearance by enhancing a

Th2 immune response, which sheds a new light on the microbiome–parasite–host immune

interactions.

Methods

Ethical statement

All animal experiments were performed according to the "Règlement Grand-Ducal du 11

janvier 2013 relatif à la protection des animaux utilisés à des fins scientifiques" based on

Directive 2010/63/EU on the protection of animals used for scientific purposes, and approved

by the Animal Experimentation Ethics Committee of the University of Luxembourg and by the

Luxembourgish Ministry of Agriculture, Viticulture, and Rural Development (national

authorization no. LUPA2020/02). The mice were housed in accordance with the

recommendations stated by the Federation of European Laboratory Animal Science

Associations (FELASA).

Experimental design and dietary treatment

18–23 weeks old C57BL/6 male mice (n=4–5 per group) were housed in ISOcages with up to

five animals per cage. Sterile food and water were provided *ad libitum* and light cycles

consisted of 12 hours. Before the administration of 14SM, the germ-free status of the mice was

confirmed using aerobic and anaerobic culturing of stools samples. Mice were intragastrically

gavaged with 0.2 ml of the 14SM gavage mix as described previously[35]. Before and for 6 days

following the gavage, mice were maintained on a standard mouse chow, which we refer to as

Fiber-rich (FR) diet. Afterwards, half of the mice (n=24) were assigned to a Fiber-free (FF)

diet and the rest remained on the FR diet (n=23). Twenty days after the diet switch, the mice

were infected by oral gavage with a high dose of approximately 450 eggs of *T. muris* strain E.

Mice were observed for up to 30 days post-infection (DPI). On 13, 15, 20, 25 and 30 DPI, four

to five mice per group were euthanized by cervical dislocation to determine worm counts and

other final readouts. A small piece (~0.5 cm) of the cecum was fixed in methacarn fixative in

order to determine goblet cell numbers. The remaining part of the cecum and 2 cm of the

attached colon were frozen at −80 °C to determine *T. muris* worm burden. The remaining

colonic tissue and mesenteric lymph nodes were stored in RNAprotect Tissue Reagent (Qiagen,

Hilden, Germany; catalog no. 76106, according to the manufacturer's protocol). Serum of the

euthanized mice was collected and stored at −80 °C.

Animal diets

The Fiber-rich diet was a standard autoclaved rodent chow (LabDiet, St. Louis, MO, USA;

catalog no. 5013), while the Fiber-free diet was a custom diet manufactured and irradiated by

SAFE Diets (Augy, France). The FF diet was manufactured as the previously described

TD.140343 diet[31], which is a modified version of the Harlan.TD08810 diet from Envigo

(Indianapolis, IN, USA).

14SM cultivation and quantification

All bacterial strains of the 14SM were cultured as previously described[35]. The colonization of

the 14SM was confirmed using strain-specific qPCR primers and relative abundances of the

individual bacteria were determined using the same qPCR protocol as described previously[35].

Samples from different dates were run on the same plates in order to avoid plate-specific

confounders.

Parasitological techniques

T. muris maintenance and infection was performed as previously described[36]. Mice were orally infected with ~450 eggs and worm burdens were assessed by counting the number of worms present in the cecum and 2 cm of the attached colon as described previously[36]. Worm burdens on 13 and 15 DPI were excluded from the analysis as determining the burden during the early larval stages, where low worm numbers establish, is technically challenging and is unreliable. The researcher, S.T., who counted the worms, was blinded for both the individual time points and dietary groups.

Goblet cell counting

Goblet cell counting was performed on a small piece (~0.5 cm) of methacarn-fixed and Alcian blue-stained cecal tissue. To preserve the mucus within the intestinal samples, a modified direct pretreatment in 99% ethanol and toluene was performed, omitting the usual previous ascending ethanol steps before embedding the samples in paraffin. All samples were cut using a microtome at 3 μm thickness before performing automated HE (Tissue-Tek Prisma Plus and Film, Sakura, Alphen aan den Rijn, Netherlands) and Alcian blue (Artisan Link Pro Special Staining System, Dako, Glostrup, Denmark) staining according to the manufacturers' instructions. The numbers of goblet cells per crypt were counted and averaged for each individual mouse before the mean per experimental group was determined. The researcher, S.T., who counted the goblet cells, was blinded for both the individual time points and dietary groups.

ELISA

Lipocalin-2 ELISA was performed on stool samples using the lipocalin-2/NGAL DuoSet ELISA R&D system (Bio-Techne, Minneapolis, MN, USA; catalog no. DY1857) according to the manufacturer's instructions and samples were prepared as previously described[31]. MCPT1 concentration of serum samples was determined using the MCPT-1 (mMCP-1) Mouse Uncoated ELISA Kit from Invitrogen (Waltham, MA, United States, catalog. no. 88-7503-88) according to manufacturer's instructions. Serum ELISA for total IgG1 and IgE were performed by coating plates using 0.5 μg/μl rat α mouse IgE purified UNLB (Imtec Diagnostics, 1130-01) or rat α-mouse IgG1 purified UNLB (Imtec Diagnostics, Ardmore, OK, United States; catalog no. 1144-01) capture antibody diluted in 0.05 M carbonate/bicarbonate buffer (pH 9.6). Washing steps were performed using 1% Tween, 154 mM NaCl, 10 mM Trizma Base. Blocking of the plates was performed using 1% w/v of BSA in a TBS buffer (15 mM Trizma-

acetate, 136 mM NaCl & 2 mM KCl). Mouse IgG1 Isotype Control UNLB (Imtec Diagnostics,

Ardmore, OK, United States; catalog no 0102-01) or Mouse IgE Isotype Control (UNLB),

Southern Biotech (Imtec Diagnostics, Ardmore, OK, United States; catalog no 0114-01) served

as controls. Serum samples were two-fold diluted using the aforementioned TBS buffer with

0.1% w/v Tween-20 and 1% BSA. The secondary antibody was Goat anti-mouse IgG1-AP,

Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog no 1071-04) or Goat

anti-mouse IgE-AP, Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog

no 1110-04), respectively. The substrate solution consisting of 0.5 mg/ml phosphate substrate

(Sigma-Aldrich, St. Louis, MO, United States; catalog no. S0642-200 TAB) in 1 mM AMP

and 0.1 mM $MgCl_2 \cdot 6H_2O$ and plates were read at 405 nm using a SpectraMax ABS PLUS

spectrophotometer (Molecular Devices, San Jose, CA, United States). A detailed protocol can

be found in the supplementary methods section. Parasite specific IgG1 and IgG2c ELISA was

performed as previously described[37].

LEGENDplex bead-based immunoassay

The LEGENDplex assay (BioLegend, San Diego, CA, USA) was performed on serum samples

according to manufacturers' instruction.

Detection of bacterial glycan-degrading enzyme activities

The enzymatic activities of α-fucosidase, β-*N*-acetyl-glucosaminidase, and β-glucosidase in

fecal samples were determined using *p*-nitrophenyl glycoside-based enzyme assays as

described previously[38].

RNA extraction from mesenteric lymph nodes and colonic tissue and RT-qPCR

A detailed protocol can be found in the supplemental methods section.

RNA-Seq

RNA extracted from samples at 30 DPI were utilized for performing RNA-Seq. Illumina

Stranded Total RNA Prep with Ribo-Zero Plus was used to prepare the RNA sequencing library

according to the reference guide's instructions. Sequencing was performed using NovaSeq

6000 SP Reagent Kit v1.5 (Illumina, San Diego, CA, USA) on an Illumina NovaSeq 6000

system.

RNA-Seq analysis

Following adapter removal with Cutadapt[39], reads were mapped and gene counts were
generated with STAR 2.7.9a[40]. Transcripts not appearing at least once on average across all
samples were removed from subsequent analysis. Samples were normalized using the median
of ratios method[41] supported by DESeq2[42] and one outlier was removed based on PC score
using the prcomp function of the base R stats 4.0.2 package. Differential expression analysis
was performed using DESeq2 1.30.1 with default parameters and p-value adjustment using the
Benjamini-Hochberg method. Significant genes were identified based on adjusted p-value <
0.05. The expressed log-ratios for significant genes were imported into Ingenuity Pathway
Analysis (Qiagen, Hilden, Germany) and analyzed considering all available reference datasets
for mice. Pathway analysis was also performed using the enrichGO function of clusterProfiler
3.18.1[43] in R to identify upregulated pathways using Gene Ontology (GO) terms for biological
processes. REVIGO[44] was employed to reduce redundancy in GO terms before visualisation.
281
Statistical analyses
Statistical analysis was performed using Prism 9.2.0. (GraphPad Software, Inc., San Diego,
CA, USA), Statistical significances are represented by asterisks as follows: *, $p < 0.05$; **,
$p < 0.01$; ***, $p < 0.001$; and ****, $p < 0.0001$. Unless otherwise specified in the figure legend,
for normally distributed values, unpaired two-tailed t tests were used, while for non-normally
distributed values, a Mann–Whitney test was used. For multiple comparisons, a two-way
analysis of variance (ANOVA) with a Šidák post hoc test was performed in Prism. The specific
test used for each experiment is detailed in the figure legends.

Results

Increased mucin-degrading bacteria coincides with altered worm infection dynamics

We colonized germ-free, C57BL/6 mice with a 14-member synthetic human microbiota (14SM). Successful colonization of each bacterial strain was checked by qPCR primers specific to each of the 14 bacterial strains as described previously[35]. Mice were maintained for six days on the Fiber-rich (FR) diet, which is a standard mouse chow, before half of the mice (n=24) were switched to a Fiber-free (FF) diet **(Fig. 1A)**. Twenty days after the diet switch, mice in both dietary groups were gavaged with a high dose of *T. muris* consisting of approximately 450 eggs from the same egg suspension **(Fig. 1A)**. Mice were monitored for up to 30 days post-infection (DPI) and, on 13, 15, 20, 25 and 30 DPI, 4–5 mice per group were sacrificed and cecum, colon and mesenteric lymph nodes (MLNs) were frozen in order to collect various experimental readouts **(Fig. 1A)**. In line with our published work[31,45], the FF diet allowed for proliferation of two dominant mucin-degrading bacteria *Akkermansia muciniphila* and *Bacteroides caccae* and reduction of two prominent fiber-degrading bacteria *Eubacterium rectale* and *Bacteroides ovatus* **(Fig. 1B)**. Throughout the experiment, none of the mice exhibited any obvious physiological abnormalities. While FF-fed mice were overall heavier compared to the FR-fed mice, the weights of both groups remained largely stable over time **(Fig. 1C)**.

Frozen cecal tissues, including the ascending colon containing their luminal contents, were defrosted and the worms were counted in a blinded fashion. Following the establishment of low level infections, during the early larval stages, the *T. muris* larvae are too small to allow reliable enumeration which is why the worm burdens were only assessed starting day 20 post infection. Infection of *T. muris* with a high egg number (~450) in conventional or specific-pathogen-free (SPF) mice, results in an acute infection, and only lower number of eggs (<40) result in a chronic infection[3]. In our synthetic human gut microbiota under standard dietary conditions (that is fed the FR diet), infection with a high number of eggs caused a low level infection resulting in a chronic infection. The reasons for this could be rooted in an initially skewed immune response in our gnotobiotic model or in the fact that our study uses a synthetic human gut microbiota, as opposed to the complete, native murine microbiota that may be more suitable for egg hatching of the naturally occurring murine whipworm or improving the ability of the larvae to establish in the gut. Considering that this a new model, it is particularly an important point because previous studies have shown that the gut bacteria are essential for effective egg hatching, as the eggs do not hatch in the absence of gut bacteria *in vitro* or in

germe-free mice[9,10]. Our results showed that, although the worm burdens were statistically
insignificant between FR and FF groups on 20 and 25 DPI, subsequent infection dynamics
statistically differed between both groups, with FF-fed mice clearing the pathogen by 30 DPI
and FR-fed mice remaining chronically infected **(Fig. 1D)**.

In order to investigate whether the two distinct diets and/or the skewed gut microbiome
composition toward mucin polymer degradation caused an altered baseline inflammation prior
to infection with *T. muris*, we determined fecal lipocalin-2 (LCN-2) levels, which serves as a
marker for low-grade inflammation[46]. There were no significant differences between FR and
FF groups at day 0 **(Fig. 1E)**, which is in line with our previous results in C57BL/6 mice[45].
However, after the infection, the levels of LCN-2 were significantly higher in the FR group
(Fig. 1E). Since chronic *T. muris* infections are characterized by a Th1 response leading to
colitis, the elevated LCN-2 in the FR group compared to the FF group is indicative of an early
induction of a Th1 response in the FR group, because LCN-2 has been shown to be induced by
the Th1 cytokines IFNγ and TNF-α[47]. Altogether, these results are supportive of the notion that
the worm retention in FR-fed 14SM group is driven by a Th1 immune response and similarly
the worm clearance in FF-fed 14SM group might be driven by a Th2 response.

Global transcriptomic analyses support chronic and acute outcomes of worm infection
In order to better understand the change in host immunological responses that drive differential
T. muris infection dynamics, we analyzed the host colonic gene expression at 30 DPI **(Fig. 2,**
Fig. S1 and Table S1). We found numerous significant differences in the host transcriptome
of FR- and FF-fed mice, as evident in the volcano plot showing the differentially expressed
genes, with 147 and 184 genes upregulated in the FR- and FF-fed groups, respectively **(Fig**
2A). Ingenuity Pathway Analysis (IPA) revealed that both the interferon signaling and Th1
pathways were upregulated in the FR-fed mice that had a homeostatic 14SM before infection
(Fig. 2B). These two pathways are characteristic of chronic *T. muris* infections[3], which was
consistent with the failure to successfully clear the parasite among FR-fed mice **(Fig. 2B)**. In
accordance with upregulation of the interferon signaling and Th1 pathways, related disease
pathways—that is, diseases often associated with an increased Th1 response such as
osteoarthritis, systemic lupus erythematosus and even neuroinflammation—were also
upregulated as their gene expression profiles overlap with those of a Th1 response **(Fig. 2B)**.

A closer look at the Ingenuity-generated summary of the transcriptomic data using IPA,
showed that indeed most of the genes upregulated under FR conditions during the chronic *T.*

muris infection are strongly linked to interferon signaling, which results in a Th1 response **(Fig.**
2C). This Th1 response is potentially linked to the observed increase in antimicrobial responses
that results in the activation of cytotoxic T cells, lymphopoiesis, increased apoptosis of
macrophages and a reduced replication of viruses **(Fig. 2C)**. In contrast to this, the host
transcriptomic response on the FF diet was not clearly characterized by an increased Th2-type
response typical for an acute *T. muris* infection **(Fig. 2B)**. However, the moderate type 2
response might be a result of the chosen sample time point as the infection is mostly cleared
by 30 DPI **(Fig. 1D)**. In line with this, the increased MSP-RON signaling in macrophage
pathway hints at an increased Th2 response under FF conditions **(Fig. 2B),** as an activation of
the MSP-RON pathway causes M2 macrophage polarization[48], which is associated with a Th2-
type immune response against *T. muris*[49,50]. Furthermore, the upregulation of the white adipose
tissue browning pathway is also in line with this observation, as M2 macrophage differentiation
has been linked to adaptive thermogenesis[51]. The observed results are consistent with a major
shift toward the Th1 pathway and a chronic infection in the FR-fed group, whereas the acute
response observed in the FF-fed group seems most likely to be based on the Th2 immunity or
a failure to generate the Th1 immunity.

Maintenance or moderate increase of Th2 immunity permits worm expulsion
In order to further support conclusions from our RNA-Seq data **(Fig. 2)**, we used RT-qPCR to
investigate transcription of cytokines IFNγ, IL-13 and IL-4 in RNA extracted from MLNs and
colon at different time points **(Fig. 3A)**. These three cytokines are indicative of whether *T.*
muris infection leads to a Th1- (IFNγ) or a Th2-type response (IL-4 and IL-13)[50,52]. In
accordance with the global transcriptome analysis **(Fig. 2)**, chronic infection in the FR-fed
group was dominated by a Th1-type response, as exhibited by over 6-fold upregulation of IFNγ
in the colon by 30 DPI **(Fig. 3A)**. Cytokine expression in the MLNs was even more
pronounced, with IFNγ having an over 30-fold increase in the FR group at day 20 and dropping
to a 6-fold increase at 30 DPI. IL-13 was only significantly upregulated in the MLNs of FF-fed
mice at 20 DPI and IL-4 showed a significant increase in the MLNs of the FR group at D15
before being upregulated under FF conditions for the subsequent days **(Fig. 3A)**. We performed
similar analysis of the cytokine transcripts of TNF-α, IL-5, IL-22 and IL-17a at 13 and 30 DPI,
but only TNF-α showed significant changes, with upregulation in the colon of the FR group
both at 13 and 30 DPI **(Fig. S2A)**. No diet-independent shifts in cytokine production were
found by analyzing the absolute cytokine data **(Fig. S3)**.

To lend support to our transcript data, we used the LEGENDplex bead-based
immunoassay to determine the cytokine protein concentration in the serum samples of mice at
30 DPI. Only one of the detected Th1 cytokines, TNF-α, was significantly upregulated under
FR conditions **(Fig. 3B)** and another Th1 cytokine, IFNγ, showed an upward trend with
p=0.0587. Some of the detected cytokines showed a trend toward increased production under
the FR conditions, hinting at a generally more active immune system during the chronic
infection **(Fig. 3B and Fig. S2B)**. The expression of transcription factors GATA3 and Tbet
(Fig. S2C) further supports these trends by showing significantly increased levels of Tbet
expression in the colon of the FR group at 30 DPI and a significant increase in GATA3 in the
MLN of the FF group at 20 DPI **(Fig. S2C)**.

Assessment of serum concentrations of total IgE and IgG1, both of which are reflective
of the protective responses against helminths, such as *T. muris*, resulted in no significant
changes, albeit a slightly increased trend was observed for both antibodies at 30 DPI but not at
25 DPI **(Fig. 3C)**. We then assessed parasite specific IgG1and IgG2, characteristic of Th2 and
Th1 responses, respectively[53]. Chronic *T. muris* infections are usually characterized by
significantly elevated serum levels of parasite-specific IgG2a/c and a low increase in parasite
specific IgG1[53]. In contrast, protection against *T. muris* is characterized by high parasite
specific IgG1 levels, but a lack of IgG2a/c[53]. Importantly, although there was no significant
increase in total IgG1, *T. muris*-specific IgG1 and IgG2a/c showed clear significant differences
at 30 DPI, but not at 25 DPI **(Fig. 3D)**. In accordance with the observed worm burdens **(Fig.**
1D), we saw an increase in specific IgG1 in mice fed the FF diet, while we see an increase in
specific IgG2a/c in mice fed the FR diet **(Fig. 3D)**. Despite a similar upward trend for both
parasite-specific antibodies at 25 DPI in the two respective dietary groups, the differences were
not significant **(Fig. 3D)**. Furthermore, we quantified MCPT1—a proxy for mast cell
degranulation[54]—in sera at all time points (13, 15, 20, 25 and 30 DPI) and observed
significantly increased MCPT1 concentration at 30 DPI in the FF-fed group that cleared the
parasite **(Fig. 3E)**. These data are in line with an increased tendency of total IgE in the FF-fed
group at 30 DPI. Moreover, these data support the known association of mucosal mast cells
and protection against helminths[55,56] supporting the elevation of the type 2 immunity.

Overall, our transcript **(Fig. 2 and Fig. 3A)** and protein expression data **(Fig. 3B)**
support that, mostly during the later stages of infection, the worm infection in mice containing
a homeostatic 14SM induce a Th1-type immune response that is driven by an overall increase
in IFNγ. On the other hand, absence of a similar increase in IFNγ, after the worm infection, in

mice with increased microbial mucin foraging, induces a Th2-type immune response, resulting

in the successful clearance of *T. muris*. This heightened Th2 response seems to peak earlier

compared to the Th1 response of the FR group and likely levels off by 30 DPI; thus, the 30

DPI transcriptomics data **(Fig. 2)** might not have fully captured the Th2 response. Since the gut

microbiome is intimately connected to changes in the diet and the host immune responses, and

since specific changes in the microbial taxa could contribute to the observed immune

responses, we ventured to determine which bacteria change in their abundance during the

course of the worm infection.

Mucin generalists increase during worm clearance, but not during worm retention

Although previous studies have reported noticeable changes in the gut microbial taxa in mice

with conventional gut microbiota after worm infection, these changes are described at broader

level of microbial taxonomy[23–28]. It also remains to be studied which precise microbial species

respond most to worm infection particularly during a longitudinal course of the worm infection

both during chronic and acute phases of infection. Better understanding such changes would

aid in further exploration of the microbiota–worm–host immune system interactions. In this

regard, our tractable, fully-characterized 14SM model[31] is well-suited to support functional

interpretations of how the underlying microbial metabolism changes as the worms start to

moult and are either expelled (acute infection) or remain persistent (chronic infection). Thus,

we next investigated the fecal gut microbial compositions using specific qPCR primers for all

five time points post infection (13, 15, 20, 25 and 30 DPI) in both dietary groups **(Fig. 4).** To

facilitate comparison, we combined these data with the microbial compositions prior to worm

infection in Fig. 1B (shown in Fig. 4 as day 0).

Throughout the experiment, the gut microbiome composition in the mice that had been

FR-fed for 20 days prior to infection remained distinct from the composition of the FF-fed mice

as seen in the principal coordinates analysis (PCoA) plots **(Fig. 4A)**. While the 14SM

composition has been shown to be stable over time[31], here we show that infection with *T. muris*

led to substantial changes in the gut microbiome composition within each dietary group **(Fig.**

4A). These changes seemed to be more dynamic under FF conditions than under FR conditions,

which is likely explained by the nature of the chronic infection under FR conditions and an

acute infection in the FF conditions, but might also be reflective of the worm burden of

individual mice **(Fig. 4A)**. Looking at the microbiome composition over the course of the

infection, while diet impacts the initial colonization of the gut microbiome, the infection with

T. muris led to the bloom of some bacterial species and a decline of others **(Fig. 4B)**.

Interestingly, some of these changes seem to revert to initial levels upon clearance of *T. muris*

in the mice fed the FF diet **(Fig. 4A and 4B)**.

Most of the individual bacteria changed in relative abundance either in response to the

early larval stages (L2 (~day 8/9)–L3 (~day 17)) of *T. muris* or in response to the later larval

stages (L3–L4 (~day 23)) **(Fig. 4C)**. These infection-induced abundance changes seem to be

reversible for most bacteria **(Fig. 4C)**. *Marvinbryantia formatexigens* appears to be an

exception as it does not significantly change in relative abundance due to either diet or infection

(Fig. 4C). Considering that we observed that nearly all bacteria change in abundance during

the early larval stages of *T. muris*, it is likely that the growing worms competitively take over

certain nutrients. While the overall bacterial abundances are modulated by the diet

composition, some bacteria such as *Collinsella aerofaciens* or *Escherichia coli* seemed

unaffected by diet. It has been reported that tissue damage induced by *T. muris* might induce

the hypoxia response gene HIF-1α[57]. The resulting increased oxygen availability might explain

the diet-independent bloom of the facultative anaerobic *E. coli*, just after the infection **(Fig.**

4C). Furthermore, other highly oxygen sensitive bacteria such as *Roseburia intestinalis* and

Faecalibacterium prausnitzii dropped in abundance around the same time, while the more

oxygen-resistant *Bacteroides* spp. remained largely unaffected **(Fig. 4C)**[58].

Among the bacteria with mucin-degrading capacities, we observed that the abundance

of *B. caccae* and *Bacteroides thetaiotaomicron* increased in the FF-fed group during the course

of the infection, but not in the FR-fed group **(Fig. 4C)**. The normalized average fold-change of

the abundance of these two mucin-generalist bacteria (*B. caccae* and *B. thetaiotaomicron*)[31]

significantly increased by 15 DPI and kept increasing until doubling their abundance at 20 DPI

in the FF-fed group but not in the FR fed group **(Fig 4D)**. In contrast, the abundance of mucin

specialist, *A. muciniphila* slightly decreased in the FF-fed group **(Fig 4C)**. Since being a mucin-

generalist and being capable of removing sulfate linkages on mucins[59] as well as secreting M-

60 like proteases that degrade the mucin protein backbone[31,32], *B. caccae* and *B.*

thetaiotaomicron are more aggressive mucin foragers than *A. muciniphila*, especially in the

absence of dietary fibers. These data suggest an overall increased mucin-foraging activity in

the FF-fed group as the worms start to grow and moult. This is further supported by the halted

expansion of the mucin-specialist *Barnesiella intestinihominis* at 15 DPI in the FF group **(Fig**

4C). A sharp significant increase in the abundance of a sulfate-reducing bacterium

Desulfovibrio piger on 20 DPI in the FF-fed group, but not in the FR-fed group, further supports

the increase in mucin-foraging, as this bacterium increasingly utilizes the released sulfate from

microbial mucin foraging as an electron acceptor[60].

Intriguingly, even in the absence of fiber, a classical fiber-degrader *E. rectale* bloomed

in the FF-fed group around 25 DPI, after being almost non-existent before this and finally

became almost non-existent again on 30 DPI **(Fig. 4C)**. This coincides with the bloom of the

mucin generalists **(Fig. 4C)**. A similar trend was observed for *E. rectale* in the FR-fed group

(Fig. 4C), with the abundance of *E. rectale* doubling in the FR-fed group as compared to its

abundance at 0 DPI. We have previously shown that *E. rectale* is adept at growing on two

mucosal monosaccharides, galactose and *N*-acetylglucosamine, but it fails to grow on mucin

oligosaccharides[31]. A potential explanation for the dramatic increase in *E. rectale* is that,

despite an absence of dietary fibers in the FF-fed group, simple mucin sugars that can be readily

metabolized, such as galactose and *N*-acetylglucosamine, become readily available through the

elevated mucin degradation in the FF-fed group at around 20 DPI by the two mucin generalists

B. caccae and *B. thetaiotaomicron*. Considering that the expansion of *E. rectale* occurs under

both diets, another supportive explanation is that the *T. muris*-secreted serine proteases, which

can depolymerize mucin[21], may make these structurally complex glycoproteins accessible to

bacteria that are otherwise unable to utilize them, such as *E. rectale*. The loss of exclusivity of

this niche offers a potential explanation for the halted expansion of the mucin specialists, *A.*

muciniphila and *B. intestinihominis* **(Fig. 4D)**. Collectively, these results strongly connect

increased mucin-degrading bacterial activity to worm clearance in the FF-fed group. As mucin

plays a key role in the clearance of *T. muris* infection, we next rationalized that enhanced

mucolytic activity during infection leads to increased mucus turnover, meaning an increased

cycle of mucin degradation and secretion, which could facilitate clearance of *T. muris* in the

FF-fed mice.

Enhanced mucin–microbiome interactions promote worm clearance

Since we found that the worm clearance is strongly correlated to increased abundance of

mucin-degrading generalist bacteria, we next set out to study how this altered microbial

composition has a functional impact on the mucin-degrading activity. To assess this, we

measured the enzyme activity of the key mucin-degrading bacterial enzymes β-*N*-

acetylglucosaminidase (NAG) and α-fucosidase (FUC) before and at days 20 and 25 post-

infection **(Fig. 5A)**. As a contrast to these key mucin-degrading enzymes, we similarly

determined the activity of β-glucosidase (GLU), a microbial enzyme specific for plant

polysaccharides **(Fig. 5A)**. At day 0, NAG activity was higher under FF conditions; however,

after the infection, the activity of this enzyme significantly rose by 25 DPI in mice resistant to

persistent worm infection, but not in the mice where worm infection became chronic. A similar

pattern was observed for FUC activity. In contrast, GLUC activity showed a significant

increase in the resistant mice only at 20 DPI **(Fig. 5A)**. Comparing NAG and FUC activity

(Fig. 5A) to the abundance of mucin-degrading bacteria **(Fig. 4C and D)**, we observed a delay

between the increased abundance of mucin-generalist bacteria under FF conditions at 20 DPI

and the increased activity of mucin-degrading enzymes at 25 DPI, possibly due to a slight delay

in extracellular secretion of the enzymes into the feces. A potential explanation is the tight

regulation of mucin-degrading enzymes, as observed for *B. thetaiotaomicron*[61]. The

depolymerized mucin released by the serine proteases of *T. muris* might initially suppress the

expression of the NAG and FUC until fiber-degrading bacteria such as *E. rectale* expand

sufficiently to fully scavenge those glycans[61]. Looking at the strong increase in the relative

abundance of some fiber-degrading bacteria, such as *E. rectale,* around 20 DPI, it is also

possible that the absolute abundance of the mucin-degrading bacteria under FF conditions

keeps increasing past 20 DPI, but that this continuous expansion is not observable in the

relative abundance due to an even stronger expansion of other bacteria (**Fig 4C and D)**. In this

case, the NAG and FUC enzyme activity might be reflective of the absolute, but not relative

abundance of the mucin-degrading bacteria.

Since we found increased abundance of mucin-generalist bacteria and subsequently

increased mucin-degrading enzyme activity during the course of worm clearance, but not

during the course of worm retention, we questioned whether this is a result of underlying

qualitative changes in mucin production in goblet cells or whether a compensatory response to

increased mucin-foraging leads to qualitative changes in mucin secretion. To better understand

the changes in mucin secretion, we performed blinded counting of goblet cells in small (~0.5

cm) pieces of cecal tissue that were preserved in methacarn fixative. The goblet cell counts

revealed no significant changes in numbers throughout the experiment **(Fig. 5B)**. Furthermore,

the expression of neither the gene *Muc2* nor the gene *Muc5ac*, which play an important role in

worm clearance[62], were upregulated **(Tables S1–S3)**. However, since these transcripts do not

necessarily reflect protein production, involvement of these mucins and/or other mechanisms

(such as epithelial cell turnover), that contribute to host protection, cannot be ruled out.

Among the genes involved in the five most upregulated pathways in resistant mice, we

identified several genes involved in glycoprotein synthesis and glycosylation **(Fig. 5C)**. This

is especially interesting as many of the upregulated genes are involved in the glycosylation of
the colonic mucins[32,63,64]. For example, the expression of the glycosyltransferases *Fut2*,
St3gal6, *C1galt1*, *Galnt3* and *Galnt4* or the sulfotransferase *Gal3st2* were upregulated in the
resistant mice[32,63,64] **(Fig. 5C)**. Altered mucin glycosylation, in particular sulfation, has been
linked to modified *T. muris* susceptibility[64]. Thus, altered glycosylation might be responsible
for the improved clearance of *T. muris* in resistant mice. This is particularly intriguing
considering that the microbiome has a major influence on the glycosylation of the Muc2
glycoprotein, the major constituent of the colonic mucus layer[63]. Furthermore, enhanced
microbial mucin foraging could potentially increase interaction of the gut microbiome with the
host mast cells, which could explain the observed increase in MCPT1 **(Fig. 3E)**. Altogether,
our results suggest that the increased mucolytic activity of the gut microbiome under fiber
deprivation **(Fig. 5A)** without the compensatory mucin production **(Fig. 5B)**, but with possible
changes in mucin glycosylation **(Fig. 5C)**, aids in clearance of *T. muris*.
567
Discussion

Helminth infection models have been profoundly utilized to elucidate the underlying host
responses for the past several decades, yet open questions remain about the mechanisms of how
the intricate interactions between the gut microbiome and GI mucus affect worm infection
dynamics including the host immune responses. Here, we shed insight into the role of bacterial
mucin foraging in altering the host immune responses during the infection with *T. muris*. Using
a diet reduced in fiber to increase bacterial mucin foraging in a gnotobiotic mouse model
containing a 14-member synthetic human gut microbiota, our study supports a model in which
the increased microbial mucolytic activity promotes Th2-driven worm clearance **(Fig. 6)**. Our
results document an intricate link in the gut microbiome–parasite axis, suggesting a potential
mechanism through which increased microbial mucin foraging could alter the host immune
response to the worm.

Our study demonstrates that an IFNγ-dominated Th1 response, which has been shown
to drive chronic *T. muris* infections, is characterized by a balanced fiber vs. mucin degradation,
while successful clearance of the worm characterized by a type 2 response is potentially the
result of increased microbial mucin foraging during worm infection **(Figs. 2, 3 and 6)**. A recent
study by Myhill et al.[27] observed a similar IFNγ-driven shift from a Th2 toward a Th1 response
when feeding SPF mice a high-fiber diet containing 10% inulin, however, this study did not
provide an explanation for the underlying mechanisms. Using our 14SM model, a standard

mouse chow containing 3.8% dietary fiber and a diet deprived of fiber, we offer an explanation

that the mucin–microbiome interactions driven by dietary changes alter the host immune

responses and the worm infection dynamics **(Fig. 6)**. Despite the well-known specificity of

parasites for their hosts, it is clear that *T. muris* is able to establish and develop in the intestine

colonized by the synthetic, human 14SM for at least 30 days. It is well known that bacteria are

important for *T. muris* worm hatching[9,10], although the specific bacterial members involved are

not well known and it is possible that the selected bacteria in our synthetic human microbiota

might not be ideal for inducing egg hatching, resulting in a lower level of infection than would

normally be seen in SPF mice with a natural microflora. The system used here, however, does

provide a chronic low-level infection characteristic of infections under natural conditions.

T. muris has been shown to broadly alter the host's gut microbiome composition[23–28],

but our longitudinal characterization of the gut microbiome provides explicit details about how

the microbial shifts are characterized by distinct phases and specific microbial species based

on their functional roles **(Fig. 6)**. In resistant mice, the microbial functional changes during the

worm infection and clearance are largely centered around microbial mucin foraging. During

the early larval stages (L2–L3) around 13 to 15 DPI, we observed a major disruption of the gut

microbiome composition, especially in the FF-fed group **(Fig. 4)**, which was characterized by

a drastic expansion of *E. coli* on both diets as well as a decline of *A. muciniphila* in resistant

mice. A similar expansion of *Enterobacteriaceae* (*E. coli*) and decline of *Akkermansia* (*A.*

muciniphila) was observed following a high-dose infection of SPF mice fed an inulin-

supplemented diet[27]. The second major shift in microbial abundances occurs around 20 to 25

DPI which coincides with the moulting of the worm from the L3 to the L4 larval stage **(Fig.**

4). During this transition, the worm moves to occupy cells nearer to the top of the crypt,

protruding into the lumen, thus most likely increasing its interaction with the gut microbiome[3].

In resistant mice, this shift is characterized by an increased abundance of the mucin generalists

B. caccae and *B. thetaiotaomicron*, resulting in an overall increased mucolytic activity of the

gut microbiome **(Figs. 4, 5A and 6)**. It is likely that the resulting secretion of M60-like

proteases by mucin generalists and the *T. muris*-secreted serine proteases depolymerize the

mucin backbone[21,31,32] generating new niches for other bacteria. This is reflected by the

increased abundance of *D. piger*, which forages the released sulfate and the increased

abundance of a strict fiber-degrader *E. rectale,* which can degrade the released mucin

monosaccharides.

In contrast to the observations in *Muc2*[-/-] mice[29], in our model, increased microbial
mucin foraging under fiber-deprived conditions did not delay the parasite clearance, but instead
promoted the clearance. These results are not unexpected as genetic ablation of Muc2 largely
destabilizes the mucus structure, whereas the increased microbial mucin foraging seems to
either resist transition to a Th1 response and/or aids in a Th2 response. We could not detect an
increase in goblet cell numbers/mucin production associated with this **(Fig. 5**B**)**. Nevertheless,
we observed increased transcription of genes involved in mucin glycosylation **(Fig. 5C)**, which
has been shown previously to alter susceptibility to *T. muris*[64]. Considering that the microbiome
can directly influence mucin glycosylation[63], our results suggest that the fiber-deprived diet-
induced changes in the microbiome might lead to an alteration in the glycosylation of the GI
mucins, which aids in clearance of *T. muris*. One caveat being that the gene transcription was
not assessed longitudinally, thus limiting interpretations considering the infection dynamics.
However, even if indirectly assessed, these data support the hypothesis that excessive mucin
degradation leads to closer interaction of the gut microbiome and the host mast cells, leading
to their activation or accumulation, as evidenced by an increase in MCPT1 concomitant with
worm clearance **(Figs. 3E and 6)**. Mast cells have been linked to *T. muris* clearance[56] and based
on the work with another nematode—*Trichinella spiralis*—this might be related to a mast cell-
dependent increase in epithelial permeability[65]. Therefore, an increase in MCPT1 offers an
additional explanation as to why excessive mucin foraging renders mice resistant to *T. muris*
infection. Finally, mast cell activation suppresses Th1 responses by decreasing IFNγ expression
and increasing Th2 cytokine expression[55], which offers an additional explanation how
excessive foraging of mucin could alter the host immune response. However, it is also likely
that the strong IFNγ expression on the FR diet simply impairs the type 2 and later mast cell
response.

Given the high morbidity associated with the parasitic infections in humans[1], our study
has generated important prerequisite knowledge for clinical settings: 1) for treating parasitic
infections – in the absence of co-infections, periodic fiber deprivation or fasting to boost
microbial mucin foraging could be potentially employed in the clinic to manage worm
infections; and 2) for harnessing the immunomodulatory properties of helminths for their
potential therapeutic use in inflammatory disorders[66] or allograft rejection[67]. Further research
is required about how the model proposed in our study **(Fig. 6)** could be translated into the
clinic.

References

1. World Health Organization. Global Health Estimates 2020: Disease burden by Cause, Age, Sex, by Country and by Region, 2000-2019. 2020.

2. Douglas B, Oyesola O, Cooper MM, et al. Immune System Investigation Using Parasitic Helminths. Annu Rev Immunol 2021;39:639–665.

3. Klementowicz JE, Travis MA, Grencis RK. Trichuris muris: A model of gastrointestinal parasite infection. Semin Immunopathol 2012;34:815–828.

4. Yousefi Y, Haq S, Banskota S, et al. Trichuris muris model: Role in understanding intestinal immune response, inflammation and host defense. Pathogens 2021;10.

5. Colombo SAP, Grencis RK. Immunity to Soil-Transmitted Helminths: Evidence From the Field and Laboratory Models. Front Immunol 2020;11.

6. Zaiss MM, Harris NL. Interactions between the intestinal microbiome and helminth parasites. Parasite Immunol 2016;38:5–11.

7. Brosschot TP, Reynolds LA. The impact of a helminth-modified microbiome on host immunity review-article. Mucosal Immunol 2018;11:1039–1046.

8. Jaenike J, Unckless R, Cockburn SN, et al. Adaptation via symbiosis: Recent spread of a drosophila defensive symbiont. Science (80-) 2010;329:212–215.

9. Hayes KS, Bancroft AJ, Goldrick M, et al. Exploitation of the intestinal microflora by the parasitic nematode trichuris muris. Science (80-) 2010;328:1391–1394.

10. White EC, Houlden A, Bancroft AJ, et al. Manipulation of host and parasite microbiotas: Survival strategies during chronic nematode infection. Sci Adv 2018;4.

11. Hewitson JP, Harcus Y, Murray J, et al. Proteomic analysis of secretory products from the model gastrointestinal nematode Heligmosomoides polygyrus reveals dominance of Venom Allergen-Like (VAL) proteins. J Proteomics 2011;74:1573–1594.

12. Abner SR, Parthasarathy G, Hill DE, et al. Trichuris suis: Detection of antibacterial activity in excretory-secretory products from adults. Exp Parasitol 2001;99:26–36.

13. Su L, Su CW, Qi Y, et al. Coinfection with an intestinal helminth impairs host innate immunity against Salmonella enterica serovar typhimurium and exacerbates intestinal inflammation in mice. Infect Immun 2014;82:3855–3866.

14.   Fricke WF, Song Y, Wang AJ, et al. Type 2 immunity-dependent reduction of
segmented filamentous bacteria in mice infected with the helminthic parasite
Nippostrongylus brasiliensis. Microbiome 2015;3:40.

15.   Forman RA, DeSchoolmeester ML, Hurst RJM, et al. The Goblet Cell Is the Cellular
Source of the Anti-Microbial Angiogenin 4 in the Large Intestine Post Trichuris muris
Infection. PLoS One 2012;7.

16.   Shea-Donohue T, Sullivan C, Finkelman FD, et al. The Role of IL-4 in
Heligmosomoides polygyrus -Induced Alterations in Murine Intestinal Epithelial Cell
Function. J Immunol 2001;167:2234–2239.

17.   Harris JB, Podolsky MJ, Bhuiyan TR, et al. Immunologic responses to Vibrio cholerae
in patients co-infected with intestinal parasites in Bangladesh. PLoS Negl Trop Dis
2009;3.

18.   Kosik-Bogacka DI, Wojtkowiak-Giera A, Kolasa A, et al. Hymenolepis diminuta:
Analysis of the expression of Toll-like receptor genes and protein (TLR3 and TLR9) in
the small and large intestines of rats. Exp Parasitol 2014;145:61–67.

19.   Zakeri A, Borji H, Haghparast A. Interaction Between Helminths and Toll-Like
Receptors: Possibilities and Potentials for Asthma Therapy. Int Rev Immunol
2016;35:219–248.

20.   Marillier RG, Michels C, Smith EM, et al. IL-4/IL-13 independent goblet cell
hyperplasia in experimental helminth infections. BMC Immunol 2008;9.

21.   Hasnain SZ, McGuckin MA, Grencis RK, et al. Serine Protease(s) Secreted by the
Nematode Trichuris muris Degrade the Mucus Barrier. PLoS Negl Trop Dis 2012;6.

22.   Hasnain SZ, Evans CM, Roy M, et al. Muc5ac: A critical component mediating the
rejection of enteric nematodes. J Exp Med 2011;208:893–900.

23.   Li RW, Wu S, Li W, et al. Alterations in the porcine colon microbiota induced by the
gastrointestinal nematode Trichuris suis. Infect Immun 2012;80:2150–2157.

24.   Broadhurst MJ, Ardeshir A, Kanwar B, et al. Therapeutic Helminth Infection of
Macaques with Idiopathic Chronic Diarrhea Alters the Inflammatory Signature and
Mucosal Microbiota of the Colon. PLoS Pathog 2012;8.

25.   Holm JB, Sorobetea D, Kiilerich P, et al. Chronic Trichuris muris infection decreases
710   diversity of the intestinal microbiota and concomitantly increases the abundance of
711   lactobacilli. PLoS One 2015;10.

712   26.   Houlden A, Hayes KS, Bancroft AJ, et al. Chronic Trichuris muris infection in
713   C57BL/6 mice causes significant changes in host microbiota and metabolome: Effects
714   reversed by pathogen clearance. PLoS One 2015;10.

715   27.   Myhill LJ, Stolzenbach S, Mejer H, et al. Fermentable Dietary Fiber Promotes
716   Helminth Infection and Exacerbates Host Inflammatory Responses. J Immunol
717   2020;204:3042–3055.

718   28.   Rosa BA, Snowden C, Martin J, et al. Whipworm-Associated Intestinal Microbiome
719   Members Consistent Across Both Human and Mouse Hosts. Front Cell Infect
720   Microbiol 2021;11.

721   29.   Hasnain SZ, Wang H, Ghia JE, et al. Mucin Gene Deficiency in Mice Impairs Host
722   Resistance to an Enteric Parasitic Infection. Gastroenterology 2010;138:1763-1771.e5.

723   30.   Johansson MEV, Jakobsson HE, Holmén-Larsson J, et al. Normalization of host
724   intestinal mucus layers requires long-term microbial colonization. Cell Host Microbe
725   2015;18:582–592.

726   31.   Desai MS, Seekatz AM, Koropatkin NM, et al. A Dietary Fiber-Deprived Gut
727   Microbiota Degrades the Colonic Mucus Barrier and Enhances Pathogen
728   Susceptibility. Cell 2016;167:1339-1353.e21.

729   32.   Martens EC, Neumann M, Desai MS. Interactions of commensal and pathogenic
730   microorganisms with the intestinal mucosal barrier. Nat Rev Microbiol 2018;16:457–
731   470.

732   33.   Parrish A, Boudaud M, Kuehn A, et al. Intestinal mucus barrier: a missing piece of the
733   puzzle in food allergy. Trends Mol Med 2022;28:36–50.

734   34.   Neumann M, Steimle A, Grant ET, et al. Deprivation of dietary fiber in specific-
735   pathogen-free mice promotes susceptibility to the intestinal mucosal pathogen
736   Citrobacter rodentium. Gut Microbes 2021;13.

737   35.   Steimle A, Sciscio A De, Neumann M, et al. Constructing a gnotobiotic mouse model
738   with a synthetic human gut microbiome to study host–microbe cross talk. STAR

739   Protoc 2021;2:100607.

740 36. Wakelin D. Acquired immunity to Trichuris muris in the albino laboratory mouse.
741   Parasitology 1967;57:515–524.

742 37. Hayes KS, Cliffe LJ, Bancroft AJ, et al. Chronic Trichuris muris infection causes
743   neoplastic change in the intestine and exacerbates tumour formation in APC min/+
744   mice. PLoS Negl Trop Dis 2017;11.

745 38. Steimle A, Grant ET, Desai MS. Quantitative assay to detect bacterial glycan-
746   degrading enzyme activities in mouse and human fecal samples. STAR Protoc
747   2021;2:100326.

748 39. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
749   reads. EMBnet.journal 2011;17:10.

750 40. Dobin A, Davis CA, Schlesinger F, et al. STAR: Ultrafast universal RNA-seq aligner.
751   Bioinformatics 2013;29:15–21.

752 41. Anders S, Huber W. Differential expression analysis for sequence count data. Genome
753   Biol 2010;11.

754 42. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
755   RNA-seq data with DESeq2. Genome Biol 2014;15.

756 43. Yu G, Wang LG, Han Y, et al. ClusterProfiler: An R package for comparing biological
757   themes among gene clusters. Omi A J Integr Biol 2012;16:284–287.

758 44. Supek F, Bošnjak M, Škunca N, et al. Revigo summarizes and visualizes long lists of
759   gene ontology terms. PLoS One 2011;6.

760 45. Wolter M, Steimle A, Parrish A, et al. Dietary Modulation Alters Susceptibility to
761   Listeria monocytogenes and Salmonella Typhimurium with or without a Gut
762   Microbiota. mSystems 2021;6:e00717-21.

763 46. Chassaing B, Srinivasan G, Delgado MA, et al. Fecal Lipocalin 2, a Sensitive and
764   Broadly Dynamic Non-Invasive Biomarker for Intestinal Inflammation Bereswill S,
765   ed. PLoS One 2012;7:e44328.

766 47. Zhao P, Elks CM, Stephens JM. The induction of lipocalin-2 protein expression in
767   vivo and in vitro. J Biol Chem 2014;289:5960–5969.

768   48.   Chaudhuri A. Regulation of macrophage polarization by RON receptor tyrosine kinase
769   signaling. Front Immunol 2014;5.

770   49.   Forman R, Logunova L, Smith H, et al. Trichuris muris infection drives cell-intrinsic
771   IL4R alpha independent colonic RELMα+ macrophages. PLoS Pathog 2021;17.

772   50.   Sorobetea D, Svensson-Frej M, Grencis R. Immunity to gastrointestinal nematode
773   infections. Mucosal Immunol 2018;11:304–315.

774   51.   Hallowell RW, Collins SL, Craig JM, et al. MTORC2 signalling regulates M2
775   macrophage differentiation in response to helminth infection and adaptive
776   thermogenesis. Nat Commun 2017;8.

777   52.   Deschoolmeester ML, Else KJ. Cytokine and chemokine responses underlying acute
778   and chronic Trichuris muris infection. Int Rev Immunol 2002;21:439–467.

779   53.   Else KJ, Entwistle GM, Grencis RK. Correlations between worm burden and markers
780   of Th1 and Th2 cell subset induction in an inbred strain of mouse infected with
781   Trichuris muris. Parasite Immunol 1993;15:595–600.

782   54.   Kasakura K, Nagata K, Miura R, et al. Cooperative Regulation of the Mucosal Mast
783   Cell–Specific Protease Genes Mcpt1 and Mcpt2 by GATA and Smad Transcription
784   Factors. J Immunol 2020;204:1641–1649.

785   55.   Mazzoni A, Siraganian RP, Leifer CA, et al. Dendritic Cell Modulation by Mast Cells
786   Controls the Th1/Th2 Balance in Responding T Cells. J Immunol 2006;177:3577–
787   3581.

788   56.   Hayes KS, Grencis RK. Trichuris muris and comorbidities-within a mouse model
789   context. Parasitology 2021;148:1774–1782.

790   57.   Hayes KS, Hager R, Grencis RK. Sex-dependent genetic effects on immune responses
791   to a parasitic nematode. BMC Genomics 2014;15.

792   58.   Browne HP, Neville BA, Forster SC, et al. Transmission of the gut microbiota:
793   Spreading of health. Nat Rev Microbiol 2017;15:531–543.

794   59.   Luis AS, Jin C, Pereira GV, et al. A single sulfatase is required to access colonic
795   mucin by a gut bacterium. Nature 2021;598:332–337.

60.   Rey FE, Gonzalez MD, Cheng J, et al. Metabolic niche of a prominent sulfate-

reducing human gut bacterium. Proc Natl Acad Sci U S A 2013;110:13582–13587.

61.   Rogers TE, Pudlo NA, Koropatkin NM, et al. Dynamic responses of Bacteroides
thetaiotaomicron during growth on glycan mixtures. Mol Microbiol 2013;88:876–890.

62.   Larsson JMH, Thomsson KA, Rodríguez-Piñeiro AM, et al. Studies of mucus in
mouse stomach, small intestine, and colon. III. Gastrointestinal Muc5ac and Muc2
mucin O-glycan patterns reveal a regiospecific distribution. Am J Physiol -
Gastrointest Liver Physiol 2013;305.

63.   Arike L, Holmén-Larsson J, Hansson GC. Intestinal Muc2 mucin O-glycosylation is
affected by microbiota and regulated by differential expression of glycosyltranferases.
Glycobiology 2017;27:318–328.

64.   Hasnain SZ, Dawson PA, Lourie R, et al. Immune-driven alterations in mucin
sulphation is an important mediator of Trichuris muris helminth expulsion. PLoS
Pathog 2017;13.

65.   McDermott JR, Bartram RE, Knight PA, et al. Mast cells disrupt epithelial barrier
function during enteric nematode infection. Proc Natl Acad Sci U S A 2003.

66.   Helmby H. Human helminth therapy to treat inflammatory disorders- where do we
stand? BMC Immunol 2015;16.

67.   Kiss M, Burns H, Donnelly S, et al. Effectiveness of Helminth Therapy in the
Prevention of Allograft Rejection: A Systematic Review of Allogeneic
Transplantation. Front Immunol 2020;11.

 Figure legends

Figure 1. Diet drives changes in mucin-degrading bacteria and alters worm infection. (A)
Experimental timeline. Age-matched 18–23 weeks old germ-free (GF) C57BL/6 mice were
gavaged with the synthetic 14-member gut microbiota (14SM) on two consecutive days. These
colonized mice were maintained for six days on the fiber-rich (FR) diet, after which nearly half
of the mice were switched to a fiber-free (FF) diet. The mice were maintained for 20 days on
their respective diets, after which they were gavaged with *T. muris* eggs. The infection was
observed for up to 30 days and on day 13, 15, 20, 25 and 30 post infection 4–5 mice were
euthanized for final readouts. (B) Relative bacterial abundance at day 0, just before the
infection, determined by qPCR on DNA extracted from fecal pellets. Red latin crosses denote
known mucin-degrading bacteria. (C) Weekly assessed weights of *T. muris* infected mice. Error
bars represent SEM and significance labels comparisons between FR and FF group as
determined by an unpaired, two-tailed t-test. (D) Worm burdens assessed on the mice
euthanized at day 13, 15, 20, 25 and 30 post infection. Error bars represent SEM; unpaired,
two-tailed t-test. (E) Fecal Lipocalin-2 (LCN-2) levels assessed by ELISA before (day 0) and
after infection with *T. muris* (13, 15, 20, 23, 30 DPI). Error bars represent SEM; unpaired t-
test. ns, non-significant; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

Figure 2. Host transcriptomics supports chronic and acute outcome of worm infection.
Analysis of data generated by RNA-seq of RNA extracted from colonic tissue at day 30 post
infection (n = 4 per group). (A) Volcano plot of the distribution of all differentially expressed
genes. (B) Barplot showing significantly up- or downregulated host pathways (Fiber Rich
(FR)/Fiber Free (FF)) determined by Ingenuity Pathway Analysis (IPA). The higher the opacity
of the bar, the more up- or downregulated the pathway. (C) Graphical summary generated by
IPA showing the major upregulated genes and pathways as well as their interactions.

Figure 3. Maintenance or moderate increase of Th2 immunity leads to worm expulsion.
(A) Relative cytokine expression of INFγ, IL-04 and IL-13 (Fiber-free versus Fiber-rich fed
mice) determined by qPCR using RNA extracted from mesenteric lymph nodes and colonic
tissue (colon) at 30 days post infection (n = 3-5). Error bars represent 95% confidence intervals;
unpaired, two-tailed t-test. (B) Serum cytokine concentrations determined by LEGENDplex
bead-based immunoassay at days 30 post infection. Error bars represent SEM; unpaired, two-
tailed t-test. (C) Total IgE and total IgG1 concentrations determined by ELISA on mouse

serum. Error bars represent SEM; two-way ANOVA using a Šidák multiple comparison test.

(D) Parasite-specific IgG1 and IgG2c concentrations determined by ELISA on mouse serum.

Error bars represent SEM; two-way ANOVA using a Šidák multiple comparison test. (E)

Serum mast cell protease 1 (MCPT1) concentrations determined by ELISA. Error bars

represent SEM. Black significance labels denote comparisons between FR and FF groups. Red

asterisk denotes significant comparison to day 0 within the FF group. Error bars represent SEM;

Two-way ANOVA using a Šidák multiple comparison test; ns, non-significant; *p<0.05;

**p<0.01; ***p<0.001; ****p<0.0001.

Figure 4. Mucin-generalist bacteria increase during worm clearance. (A) PCoA plot of the

microbiome composition of all mice split by diet (left panel) or by day and diet (middle and

right panels). PCs were selected based on Eigenvalues. (B) Microbial composition of the gut

microbiota before the infection (day 0) and at day 13, 15, 20, 25 and 30 post infection (denoted

by dotted lines). Relative bacterial abundances were determined by qPCR of DNA extracted

from fecal pellets using primers specific to each bacterial strain. Red crosses denote known

mucin-degrading bacteria. (C) Relative abundances of individual bacterial strains. Black

significance labels denote comparisons between the Fiber-rich (FR) and Fiber-free (FF) groups

on a specific day. Green asterisks denote significant comparisons to day 0 within the FR group.

Red asterisks similarly denote significant comparisons to day 0 within the FF group. Error bars

represent SEM; two-way ANOVA using a Šidák multiple comparison test. (D) Fold-change of

mucin generalists (*Bacteroides caccae* and *Bacteroides thetaiotaomicron*) and mucin

specialists *(Akkermansia muciniphila* and *Barnesiella intestinihominis*) normalized by their

average day 0 relative abundance. Black significance labels denote comparisons between the

FR and FF group on a specific day. Green asterisks denote significant comparisons to day 0

within the FR group. Red asterisks similarly denote significant comparisons to day 0 within

the FF group. Error bars represent SEM; two-way ANOVA using a Šidák multiple comparison

test. ns, non-significant; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

Figure 5. Increased mucin–microbiome interactions promote worm clearance. (A)

Glycan-degrading enzyme activities of the gut microbiome determined by stool-based p-

nitrophenyl glycoside-based enzyme assays. β-N-acetyl-glucosaminidase (NAG) and α-

fucosidase (FUC) are key mucin-degrading enzymes, while β-glucosidase (GLUC) serves as a

control for general glycan-degrading activity. Error bars represent SEM; Mann-Whitney test

between Fiber-rich (FR) and Fiber-free (FF) groups. (B) Goblet cell counts per crypt of *T.*

muris infected mice determined by histological assessment of methacarn-fixed cecum sections.

Error bars represent SEM; two-way ANOVA using a Šidák multiple comparison test. (C)

Relative gene expression (FF/FR) at 30 days post infection of the 5 most significant activated

pathways in the colonic tissue under FF conditions (n = 4 per group). Pathways are based on

GO terms for biological processes and REVIGO was used to reduce redundant terms. ns, non-

significant; *p<0.05; **p<0.01. †Adaptive immune response is based on somatic

recombination of immune receptors built from immunoglobulin superfamily domains.

Figure 6. Model for clearance of *T. muris* following excessive microbial mucin foraging.

Under homeostatic conditions there is a balance between bacterial fiber- and mucin-

degradation. Deprivation of dietary fiber leads to an increased bacterial mucin-degrading

activity of the gut microbiome, which is further increased after infection with *T. muris*. This

increase is a result of the expansion of mucin-generalist bacteria around 20 DPI, which in turn

opens up a growth niche for other bacteria such as *D. piger* and *E. rectale*, which can utilize

the sulfate or simple mucin glycoproteins, respectively, resulting from the depolymerization of

the complex host mucins by the mucin-generalist bacteria. In contrast to mucin-generalists,

mucin-specialists drastically drop in abundance following worm hatching in Fiber-free (FF)-

diet fed mice. In Fiber-rich (FR)-diet fed mice, the host develops a Th1 response characterized

by an increase of Th1-type cytokines, particularly IFNγ, and the production of parasite specific

IgG2 resulting in a chronic *T. muris* infection. In contrast, the excessive mucin foraging in FF-

fed mice leads to a moderate Th2 response characterized by the secretion of Th2 cytokines,

expression of parasite-specific IgG1 and increased degranulation of mast cells ultimately

leading to the clearance of *T. muris* around 30 DPI.

Graphical abstract

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary figure legends

Figure S1. Host gene expression on the FR diet is dominated by genes involved in Th1 response.
Relative gene expression (FR/FF) at 30 days post infection of the 5 most significant activated pathways in the colonic tissue under FF conditions (n = 4 per group). Pathways are based on GO terms for biological processes and REVIGO was used to reduce redundant terms.

Figure S2. Cytokine transcript and protein abundances on different days post worm infection.
(A) Relative cytokine expression of TNF-α, IL-5, IL-22 and IL-17a (fiber-free over fiber-rich fed mice) determined by qPCR of cDNA from RNA extracted from mesenteric lymph nodes (MLN) and colonic tissue (colon) at 13 and 30 DPI. Error bars represent 95% confidence intervals; unpaired, two-tailed t-test. (B) Serum cytokine concentrations of IL-22 and IL-5 at 30 DPI determined by LEGENDplex bead-based immunoassay. Error bars represent SEM; unpaired, two-tailed t-test. (C) Relative transcription factor expression of Tbet and GATA3 (Fiber-free over Fiber-rich fed mice) determined by qPCR using RNA extracted from mesenteric lymph nodes (MLN) and colonic tissue (colon) at 13–30 DPI. In the colon, GATA3 for all DPI and Tbet for 13 DPI were below the detection threshold. Error bars represent 95% confidence intervals; unpaired, two-tailed t-test; ns, non-significant; *p<0.05; **p<0.01.

Figure S3. Absolute abundances of cytokine transcripts on different days post worm infection.
(A) Absolute cytokine and transcription factor expression as determined by qPCR using RNA extracted from mesenteric lymph nodes (MLN) and colonic tissue (colon) at 13, 15, 20, 25 and 30 DPI. Error bars represent SEM; unpaired, two-tailed t-test. (B) Absolute cytokine expression as determined by qPCR of using RNA extracted from mesenteric lymph nodes (MLN) and colonic tissue (colon) at 13 and 30 DPI. Error bars represent SEM; unpaired, two-tailed t-test. In the colon, GATA3 for all DPI and Tbet for 13 DPI were below the detection threshold. Error bars represent 95% confidence intervals; unpaired, two-tailed t-test; ns, non-significant; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

Supplementary table legends

Table S1. Relative gene expression (FR/FF) based differential expression analysis using DESeq2.

Table S2. Significantly upregulated pathways in FR diet based on gene oncology (GO) terms for biological processes.

Table S3. Significantly upregulated pathways in FF diet based on gene oncology (GO) terms for biological processes.

Table S4. RT-qPCR primers for genes encoding various cytokines, transcription factors and a reference (GAPDH).

Supplemental methods

Serum ELISA for total IgG1 and IgE

Serum ELISA for total IgG1 and IgE were performed as follows. Plate were coated overnight at room temperature using 0.5 μg/μl rat α mouse IgE purified UNLB (Imtec Diagnostics, 1130-01) or rat α-mouse IgG1 purified UNLB (Imtec Diagnostics, Ardmore, OK, United States; catalog no. 1144-01) capture antibody, respectively, diluted in 0.05 M carbonate/bicarbonate buffer (pH 9.6). Wells were washed three times using a washing buffer consisting of 1% Tween, 154 mM NaCl, 10 mM Trizma Base. 1% w/v of BSA in a TBS buffer (15 mM Trizma-acetate, 136 mM NaCl & 2 mM KCl) was used to block plates by incubating them for 2 hours at room temperature. Afterwards, plates were washed three times using the washing buffer. As controls, Mouse IgG1 Isotype Control UNLB (Imtec Diagnostics, Ardmore, OK, United States; catalog no 0102-01) or Mouse IgE Isotype Control (UNLB), Southern Biotech (Imtec Diagnostics, Ardmore, OK, United States; catalog no 0114-01) was used to generate two-fold serial dilutions. Serum samples were two-fold diluted from 1/40 to 1/1280 using the aforementioned TBS buffer with 0.1% w/v Tween-20 and 1% BSA. Plates were loaded the standards and samples were incubated for 90 minutes at room temperature. Another three wash steps were performed before loading a 1/500 dilution of the 2nd antibody, Goat anti-mouse IgG1-AP, Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog no 1071-04) or Goat anti-mouse IgE-AP, Southern Biotech (Imtec diagnostics, Ardmore, OK, United States; catalog no 1110-04), respectively. The final three wash steps were performed before loading the substrate solution consisting of 0.5 mg/ml phosphate substrate (Sigma-Aldrich, St. Louis, MO, United States; catalog no. S0642-200 TAB) in 1 mM AMP and 0.1 mM MgCl$_2$•6H$_2$O. Plates were incubated for 1 hour at 37°C before they were read at 405 nm using a SpectraMax ABS PLUS spectrophotometer (Molecular Devices, San Jose, CA, United States).

RNA extraction from mesenteric lymph nodes and colonic tissue

Serum ELISA for total IgG1 and IgE were performed as follows. Plate were coated overnight at room temperature using 20 μl/well of 0.5 μg/μl rat α mouse IgE purified UNLB (Imtec Diagnostics, 1130-01) or rat α-mouse IgG1 purified UNLB (Imtec Diagnostics, Ardmore, OK, United States; catalog

no. 1144-01) capture antibody, respectively, diluted in 0.05 M carbonate/bicarbonate buffer at pH

9.6. Wells were washed three times using 100 μl/well of a washing buffer consisting of 1% Tween,

154 mM NaCl, 10 mM Trizma Base. Then plates were blocked by incubating them for 2 hours at

room temperature with 75 μl of blocking buffer consisting of 1% w/v of BSA in a TBS buffer (15

mM Trizma-acetate, 136 mM NaCl & 2 mM KCl). Afterwards, plates were washed three times using

the washing buffer. As controls, Mouse IgG1 Isotype Control UNLB (Imtec Diagnostics, Ardmore,

OK, United States; catalog no 0102-01) was used to generate a two-fold serial dilution from 1/500 to

1/512000 and Mouse IgE Isotype Control (UNLB), Southern Biotech (Imtec Diagnostics, Ardmore,

OK, United States; catalog no 0114-01) was used to generate a two-fold serial dilution from 1/1000

to 1/1024000. Serum samples were two-fold diluted from 1/40 to 1/1280. The dilution buffer

consisted of 0.1% w/v Tween-20 and 1% BSA in the aforementioned TBS buffer. Plates were loaded

with 20 μl/well of the standards and diluted samples and incubated for 90 minutes at room

temperature. Another three wash steps were performed before loading 20 μl/well of a 1/500 dilution

of the 2nd antibody, Goat anti-mouse IgG1-AP, Southern Biotech (Imtec diagnostics, Ardmore, OK,

United States; catalog no 1071-04) or Goat anti-mouse IgE-AP, Southern Biotech (Imtec diagnostics,

Ardmore, OK, United States; catalog no 1110-04), respectively. The final three wash steps were

performed before loading 40 μl/well of the substrate solution, which consists of 0.5 mg/ml phosphate

substrate Sigma-Aldrich, St. Louis, MO, United States; catalog no. S0642-200 TAB) in a 1 mM AMP

and 0.1 mM $MgCl_2 \cdot 6H_2O$. Plates were incubated for 1 hour at 37°C before they were read at 405 nm

using a SpectraMax ABS PLUS spectrophotometer (Molecular Devices, San Jose, CA, United

States).

RT-qPCR

Targeted analysis of cytokine and transcription factor expression was determined by RT-qPCR of

RNA extracted from mesenteric lymph nodes (MLN) and colonic tissue (colon). The cDNA library

was prepared by combining 500 μM dNTP Set (100 mM) Solution (Invitrogen, Waltham, USA), 2.5

μM Random Primers (Invitrogen), 100 ng/μl of RNA sample and 1ul of ddH2O per sample, followed

by heating to 65°C for 5 min and incubation on ice for at least 1 min. Next, 1× SSIV Buffer, 100 mM

DTT, 40 U RNaseOUT (Invitrogen, Waltham, USA), Recombinant Ribonuclease Inhibitor

(Invitrogen, Waltham, USA), and 200 U Superscript IV Reverse Transcriptase (Invitrogen, Waltham,

USA) was added to each sample and incubated at 23°C for 10 min. Samples were incubated at 50°C

for 10 min, then heat-inactivated at 80°C for 10 min. A master mix consisting of 1× Buffer, 2.5 mM

$MgCl_2$, 400uM dNTP, 1× SYBR Green I Nucleic Acid Gel Stain, 10,000× concentrate in DMSO

(Invitrogen, Waltham, USA), and 0.5 U Platinum *Taq* DNA Polymerase (Life Technologies,

Carlsbad, USA), was added to 1 µl each cDNA sample along with 0.2 µM each of forward and reverse

primers (the specific primers are listed in **Table S4)**. The PCR cycle consisted of pre-denaturation at

94°C for 5 min, followed by 40 cycles of 20 sec denaturation at 94°C, 50 sec annealing at 60°C, and

45 sec extension at 72°C. Samples were held at 72°C for 5 min post-extension and then a melting

curve was generated by heating from 65°C to 95°C with 0.3°C interval increases over 15 sec.

Figure S1

Figure S2

Figure S3