In vivo chromatic and spatial tuning of foveolar retinal ganglion cells in *Macaca fascicularis*

Tyler Godat*[1,2], Nicolas P. Cottaris[3], Sara Patterson[1], Kendall Kohout[1,2], Keith Parkins[1], Qiang

Yang[1], Jennifer M. Strazzeri[1,4], Juliette E. McGregor[1], David H. Brainard[3], William H. Merigan[1,4] &

David R. Williams*[1,2,4]

[1] Center for Visual Science, University of Rochester, Rochester, NY, United States of America.

[2] Institute of Optics, University of Rochester, Rochester, NY, United States of America.

[3] Department of Psychology, University of Pennsylvania, Philadelphia, PA, United States of

America.

[4] David & Ilene Flaum Eye Institute, University of Rochester Medical Center, Rochester, NY,

United States of America.

* Corresponding authors

E-mail: tgodat@ur.rochester.edu (TG), david.williams@rochester.edu (DRW)

Abstract

The primate fovea is specialized for high acuity chromatic vision, with the highest density of cone

photoreceptors and a disproportionately large representation in visual cortex. The unique visual

properties conferred by the fovea are conveyed to the brain by retinal ganglion cells, the somas

of which lie at the margin of the foveal pit. Microelectrode recordings of these centermost retinal

ganglion cells have been challenging due to the fragility of the fovea in the excised retina. Here

we overcome this challenge by combining high resolution fluorescence adaptive optics

ophthalmoscopy with calcium imaging to optically record functional responses of foveal retinal

ganglion cells in the living eye. We use this approach to study the chromatic responses and spatial

transfer functions of retinal ganglion cells using spatially uniform fields modulated in different

directions in color space and monochromatic drifting gratings. We repeated these recordings in

over 350 cells across three *Macaca fascicularis* primates over a time frame of weeks to months.

We find that the majority of the L vs. M cone opponent cells serving the most central foveolar

cones have spatial transfer functions that peak at high spatial frequencies (25-40 c/deg), reflecting

strong surround inhibition that sacrifices sensitivity at low frequencies but preserves the

transmission of fine detail in the retinal image. In addition, we fit to the drifting grating data a

detailed model of how ganglion cell responses draw on the cone mosaic to derive receptive field

properties of L vs. M cone opponent cells at the very center of the foveola. The fits are consistent

with the hypothesis that midget ganglion cells are specialized to preserve information at the

resolution of the cone mosaic. By characterizing the functional properties of retinal ganglion cells

in vivo through adaptive optics, we characterize the response characteristics of these cells *in situ*.

Introduction

In primates, the fovea is a specialized region of the retina characterized by a high density of cones

at its center and a corresponding radial displacement of inner retinal cells and retinal vasculature.

These specializations, along with a vastly expanded representation in cortex [1-3], contribute to

the superiority of foveal vision relative to peripheral vision in spatial resolution and color

discrimination [1, 4-21]. To date, in excess of 20 classes of retinal ganglion cells (RGCs) have been

identified in macaque retina [1, 4, 6, 7, 13, 17-20, 22-52], reflecting a complex array of retinal

computations that precede additional processing in the brain. Despite these advances, a complete

understanding of these independent circuits and their role in vision and behavior remains elusive.

Even the precise role and scope of function of the most numerous and commonly studied RGCs,

the midget ganglion cells, remains controversial, and there has been relatively little study at the

very center of the foveola.

Electrophysiological responses from primate RGCs have been studied *in vivo* with a

microelectrode that penetrates the sclera [53, 54] and *in vitro* using a single microelectrode [6,

27] or microelectrode arrays [55, 56]. However, many recordings in primate have been made in

peripheral retina, often 30 to 70 degrees from the fovea because of the difficulty penetrating the

internal limiting membrane near the fovea. Moreover, the survival of an excised fovea is time

limited, and the delicate structure of the foveal pit is often strained when flattened, sometimes

tearing crucial circuitry. While the anatomical basis for foveal circuitry has been studied

extensively [7, 32, 34, 38, 57, 58] and there has been some *ex vivo* physiology of foveal RGCs, most

individual cell classes present in the primate fovea have not been directly studied, especially at

the most central locations. Retinal processing can also be inferred by recording from

retinorecipient neurons in the LGN, but the foveal representation of the LGN is located within a

relatively thin cell layer making such recordings challenging [59]. Moreover, it has been difficult

to identify the exact foveal center *in vivo* and determining the precise retinal eccentricity of single

cell recordings in LGN is difficult [5, 9, 60-62]. Furthermore, RGCs project to many different targets

other than LGN, so studying these cells at the retinal level gives the best chance at providing a

more complete sampling.

In this study, we use functional adaptive optics cellular imaging in the living eye (FACILE) [12, 16,

63, 64], a technique that utilizes an adaptive optics light scanning ophthalmoscope (AOSLO) to

image fluorescence elicited from the activity of RGCs that express genetically encoded calcium

indicators, such as those described by Chen et al [65]. Simultaneously, the cone mosaic is imaged

with infrared wavelengths to map the cone density profile at the fovea. The cone mosaic images

have a relatively high signal to noise ratio (SNR) and are also used to track residual eye motion in

the fluorescence channel which has a relatively low SNR. Knowing the cone density profile for an

individual subject allows for a more precise modeling of receptive field structure and more precise

motion correction. Using this technique, we can both image and stimulate the centermost

foveolar cones with spatial or chromatic stimuli and optically record the activity of the ganglion

cells that are served by those cones, characterizing the physiology of the innermost foveal primate

RGCs *in vivo*. Indeed, we can return repeatedly to the same ganglion cells for weeks to months,

allowing repeated functional measures of the same cells. With this approach, we can use data

from an individual animal that includes the optics, the cone density profile, and the RGC responses

to create a detailed model of the first stages of visual processing.

Materials and Methods

Animal Care. The macaques (*Macaca fascicularis)* were housed in pairs in an AAALAC accredited

facility. They had free access to food and water, providing a complete and nutritious diet. To

supplement their ordinary lab chow, animals were given various treats daily such as nuts, raisins,

and a large variety of fresh fruits and vegetables. An animal behaviorist provided novel

enrichment items once per week, including items such as grass and treat filled bags, grapevines,

and forage boxes. Daily enrichment included several pieces of manipulata: mirrors, puzzle feeders

rotated among the animals, daily movies and/or music, as well as rotating access to a large free-

ranging space with swings and elevated perches during any long periods between imaging

sessions. All macaques were cared for by the Department of Comparative Medicine veterinary

staff, including four full-time veterinarians, five veterinary technicians, and an animal care staff

who monitored animal health and checked for signs of discomfort at least twice daily. This study

was carried out in strict accordance with the Association for Research in Vision and

Ophthalmoscopy (ARVO) Statement for the Use of Animals and the recommendations in the

Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol

was approved by the University Committee on Animal Resources of the University of Rochester

(PHS assurance number: D16-00188(A3292-01)).

Immune Suppression. Two macaques (M2, M3) received subcutaneous Cyclosporine A prior to

intravitreal injection and imaging. Blood trough levels were collected weekly to titrate the dose

into a therapeutic range of 150-200 ng ml^{-1} and then maintained at that level. M2 began immune

suppression in March 2018 with 6 mg kg^{-1}, then stepped down a month later to 4 mg kg^{-1} which

was maintained until October 2019 when suppression was stopped; we believe that the cessation

of immune suppression was responsible for widespread apparent death of foveal ganglion cells

some months later in this animal (data not shown in this manuscript). M3 started immune

suppression in May 2019 at 6 mg kg^{-1}, then stepped down to 3.4 mg kg^{-1} which continues at time

of submission. M1 did not receive any immune suppression.

AAV mediated gene delivery to retina. Intravitreal injections were carried out in each animal as

previously described [66]. In M1, *7m8-SNCG-GCaMP6f* [67] was injected into the right eye; *AAV2-*

CAG-GCaMP6s was injected into the left eye of M2 and into the right eye of M3. All vectors were

synthesized by the University of Pennsylvania Vector Core. Before the injections, the eyes were

sterilized with 50% diluted Betadine, and the vector was injected into the middle of the vitreous

at a location approximately 4 mm behind the limbus using a tuberculin syringe and 30-gauge

needle. Following injection, each eye was imaged with a conventional scanning light

ophthalmoscope (Heidelberg Spectralis) using the 488 nm autofluorescence modality to

determine onset of GCaMP expression and to periodically monitor image quality and eye health.

Anesthesia and animal preparation. Anesthesia and animal preparation followed established lab

protocols as previously reported [e.g. 16, 68] and were performed by a veterinary technician

licensed by the State of New York (USA). All monkeys were fasted overnight prior to anesthesia

induction the morning of an imaging session. The animal was placed prone onto a custom

stereotaxic cart where it remained for the duration of the imaging session. A Bair Hugger warming

system was placed over the animal to maintain body temperature. Monitoring devices including

rectal temperature probe, blood pressure cuff, electrocardiogram leads, capnograph, and a pulse

oximeter, were used to track vital signs. Temperature, heart rate and rhythm, respirations and

end tidal CO_2, blood pressure, SPO$_2$, and reflexes were monitored consistently and recorded every

fifteen minutes. A full description of all medications used in anesthesia induction, pupil dilation,

intubation, and animal recovery can be found in McGregor *et. al.* [68].

Adaptive optics imaging. Data were collected using an AOSLO system previously described in Gray

et al. [63]. An 847 nm diode laser source (QPhotonics) was used as a beacon for a Shack-Hartmann

Wavefront Sensor (SHWS) to measure optical aberrations in each animal's eye in real time, and a

Deformable Mirror (ALPAO) was used to correct those aberrations in a closed loop. A 796 nm

superluminescent diode source (Superlum) was focused on the photoreceptor layer of the retina

to collect reflectance images at an approximately 2 Airy disk confocal pinhole (20 µm), with the

images used for both placement of stimuli and later registration (motion correction) of

fluorescence images. A 561 nm laser (Toptica) and three LEDs (Thorlabs, center wavelengths 420

nm, 530 nm, 660 nm) were used for visual stimulation of foveal cones. A 488 nm laser source

(Qioptiq) was focused on the ganglion cell layer to excite GCaMP fluorescence, which was

detected through a 520/35 nm emission filter at an approximately 2 Airy disk confocal pinhole (20

µm) for animals M2 and M3 and an approximately 7.5 Airy disk confocal pinhole (75 µm) for

animal M1, each of which was chosen to balance the tradeoff between signal strength and axial

resolution. It is likely that the lack of immune suppression in M1 resulted in the much poorer

expression of GCaMP and consequently worse SNR in that animal, which thus motivated using the

much larger confocal pinhole for imaging M1. The 488 nm excitation light was presented only

during forward scans and filled only the portion of the imaging window where ganglion cells were

present to avoid exposing foveal photoreceptors and biasing stimulation. The 488 nm excitation

light intensities on the retina were 3.4 mW cm^{-2} in M1, 3.2 mW cm^{-2} in M2, and 1.7 mW cm^{-2} in

M3, as presented through a dilated pupil size of approximately 6.7 mm for each animal. Retinal

subtenses were 780 x 570 µm in M1, 765 x 560 µm in M2, and 740 x 540 µm or 505 x 380 µm in

M3.

Light safety. The total retinal exposure to all light sources was calculated for each proposed retinal

location before each imaging session after measuring the optical power of each source at the

pupil plane (nominally 488 nm at 7-15 µW, 561 nm at 5 µW, 796 nm at 250 µW, 847 nm at 30 µW,

and combined 5 µW from the LEDs at 420 nm, 530 nm, and 660 nm). For all animals, the total

exposure was kept below the maximum permissible exposure for human retina according to the

2014 American National Standards Institute [69-71]. Most exposures were also kept below a

reduced ANSI limit that was further scaled by the squared ratio of the numerical aperture of the

human eye to the primate eye (~0.78) [72], though imaging sessions in M3 at the smaller FOV

exceeded this reduced limit. Imaging sessions were at least five days apart, so accumulated

exposure was not taken account into the calculations.

Methods to maximize optical quality and signal across experiments. Our method affords the

advantage of being able to record from hundreds of cells simultaneously, but there is a tradeoff.

The best axial resolution that can be achieved with the confocal AOSLO required for our

experiments is 25-30 microns, approximately the diameter of 2-3 RGC somas [27], which means

that in areas where the RGC layer is thick, such as the fovea [26, 46], there may be some optical

crosstalk from multiple cells contributing to the recorded fluorescence signal. Therefore, as part

of the validation of our method and to be certain in describing the physiology of individual cells,

we operate in two distinct modalities. The first modality is a wide imaging field of view (FOV) of

3.70 x 2.70 deg where we can image hundreds of RGCs across a breadth of soma locations (in M1,

M2 and M3), but not be certain that each cell is free of crosstalk. The second modality (only used

in M3) is a smaller imaging FOV of 2.54 x 1.92 deg that is centered and focused on the innermost

foveal slope, where the RGCs are in a monolayer—in this modality we can only image tens of cells

at a time, but we are certain that the signal is not contaminated by optical crosstalk. We present

wide field data from all three animals to show cell populations and general physiology, while we

present data from the smaller field only from animal M3 to draw conclusions about the physiology

of individual cells—the very innermost foveolar RGCs driven by the centermost cones. To ensure

the desired imaging FOVs were used in each experiment, the AOSLO system was calibrated using

Ronchi gratings of known spacing (80 lp/mm) to translate scanner control voltages into FOV

measurements in degrees of visual angle.

To correct for longitudinal chromatic aberration between the cone reflectance source (796 nm),

visual stimulus laser (561 nm), and GCaMP excitation laser (488 nm), axial source and detector

positions of the visible channels were optimized by initial calculation using chromatic dispersion

data in humans [73] scaled for the macaque eye and then further optimized empirically *in vivo*

using algorithms such as simplex [74] to iterate over lateral pinhole positions and find the location

with maximum detector mean pixel value. For the data in M3 at the inner edge of the foveal slope,

an additional experiment optimized the responses of 37 RGCs to 28 c/deg drifting gratings at 6 Hz

to find the best axial source position of the 488 nm excitation laser, to account for RGCs on the

foveal slope differing in axial position. Transverse chromatic aberration was not critical to stimulus

placement in these experiments, but lateral positions of confocal detection pinholes were also

optimized using similar algorithms [75].

Perhaps as a side effect of the *in vivo* viewing eye being held open for long periods of time during

imaging sessions, optical quality tends to degrade over the course of the experiment, slowly at

first, but more quickly after the first hour of imaging. To mitigate this, all grating stimuli were

presented at the beginning of a session when optical quality was the highest, and only for the first

30 minutes, to minimize degradation of high spatial frequency responses.

Lastly, the isoplanatic patch size of our system is approximately 1-1.5 deg, which is smaller than

both of the imaging fields (3.70 x 2.70 deg and 2.54 x 1.92 deg) used in this study. This degrades

quality across the entire image and produces field aberrations towards the edges of the images

where some ganglion cells were imaged or cones were stimulated. This anisoplanatism, coupled

with the fact that the wavefront correction is at a different wavelength than stimulation, means

that there could be defocus (see Modeling section) or higher order residual aberrations that

preferentially blur high spatial frequency gratings. Care was taken to mitigate or optimize these

sources of residual blur, but they cannot completely be eliminated with our current instrument.

Therefore, the spatial frequency responses we measured represent a lower bound for the cell

responses *in situ* when all sources of residual blur are removed.

Chromatic Stimuli. To drive the chromatic responses of RGCs connected to foveal photoreceptors,

we presented a 1.3 deg diameter spatially uniform, circular LED stimulus (Thorlabs, center

wavelengths 420nm, 530nm, 660nm) through a Maxwellian view system [76]. The stimulus was

sinusoidally modulated in time. Mean intensities on the retina and temporal frequencies were 6.8

mW cm^{-2} at 0.2 Hz in M1, 7.0 mW cm^{-2} at 0.2 Hz in M2, and 7.6 mW cm^{-2} at 0.15 Hz in M3. The

slow temporal frequencies were chosen to accommodate the slow temporal response of the

GCaMP6 calcium indicator [65] and were adjusted by 0.05 Hz in M3 to avoid potential confound

with the respiration rate in that animal. The LED primary wavelengths were chosen to maximize

excitation of the L, M, and S cone photopigments [77] while minimizing significant confound by

macular pigment absorption [78, 79], the spatial distribution of which was not measured in this

study. Each stimulus presentation was 90 s long, following a 30 s adaptation period in which the

mean luminance white point and excitation laser were on. The cone fundamentals were

constructed using the Govardovskii standard template [80, 81] and the wavelengths of peak

absorption for macaque primates [82]. The transmissivity of the eye (including cornea, aqueous,

lens, and vitreous) was modeled using measured data from the rhesus macaque [83]. To generate

the luminous efficiency function, an equal weighting of the L and M cone fundamentals was used,

as the ratio of L and M cones in macaques is close to unity on average [84]. The LEDs were

calibrated using a spectrometer (Ocean Optics) validated by a NIST traceable blackbody source.

Five silent substitution [85] stimuli were presented at the foveal center while GCaMP responses

were recorded. The Psychophysics Toolbox [86] for MATLAB was used to calculate the appropriate

power modulations needed: a nominally isoluminant stimulus targeting both L and M cones in

counterphase (L modulation 15%, M modulation 17%), an L cone isolating stimulus (24%

modulation), an M cone isolating stimulus (33% modulation), an S cone isolating stimulus (92%

modulation), and an achromatic luminance stimulus (100% modulation, all cone classes). There

was also a control stimulus where the adaptation period was followed by continued presentation

of the mean luminance white point. Within one imaging session, each of the stimuli used in that

session was presented three times and responses were averaged within the experiment. In M1,

only one experiment was performed. In M2 and M3, two experiments across the entire width of

soma locations (large imaging FOV) were performed and responses across sessions were

averaged. In M3, three separate experiments at the innermost foveal RGCs (small imaging FOV)

were performed and averaged.

Grating Stimuli. To probe the spatial configuration of foveal RGC receptive fields, we presented

monochromatic (561 nm) horizontally-oriented sinusoidal drifting gratings at 6 Hz and 100%

contrast. The grating stimuli were presented within the scanned imaging FOV and occupied a

square subregion that was either 1.9 deg (large imaging FOV) or 1.3 deg (small imaging FOV) in

extent. Mean light intensities on the retina were 1.72 mW cm^{-2} in M2 (large imaging FOV), 1.8 mW

cm^{-2} in M3 (large imaging FOV), and 3.2 mW cm^{-2} in M3 (small imaging FOV) imaging only the

innermost RGCs closest to the center of the foveola. No data from drifting gratings is shown for

animal M1, as the signal was too weak and noisy to produce reliable data from these stimuli. The

temporal frequency of these gratings was too fast to allow GCaMP6 to track modulation, so

cellular responses were characterized by quantifying the increase in steady fluorescence during

stimulation, as opposed to using fluorescence modulation as was done for the slower chromatic

stimuli. Grating stimuli were 45 s in duration and were preceded by 15 s adaptation to both the

mean luminance of the 561 nm stimulus laser and the 488 nm imaging laser.

In the small imaging FOV used with M3, the increased resolution meant that the spatial frequency

range was 4 c/deg to 49 c/deg. A control and 14 gratings were presented twice each within a

session and responses were averaged. Gratings were presented in decreasing order of spatial

frequency, to minimize reduction in high spatial frequency response related to decreasing optical

quality over the course of the experiment. There were three sessions at the small FOV in M3, and

the variability and averages are shown in the results. We note that as gratings were only

presented in one orientation, the responses of some cells might be modified by orientation

selective effects [87].

In M2 and M3 at the large imaging FOV, a control (spatially uniform mean luminance) and 14

gratings varying in spatial frequency from 2 c/deg to 34 c/deg were presented. Each grating was

presented twice within a session and responses were averaged for one session in M2 and across

two sessions in M3. These data are not presented in the results because they did not include high

enough spatial frequency responses that permit the modeling used in this work (see Supplement

S2 Fig.).

Data analysis. To remove blurring and other effects of residual eye movements present in the raw

recordings, each video frame of the fluorescence recording was co-registered using corresponding

frames from a high signal-to-noise reflectance video of the photoreceptor layer at the same

retinal location. During imaging, a reference frame was taken in the reflectance channel to allow

for real-time stabilization; this reference frame was also used as the reference for the frame-to-

frame cross-correlation method image registration [88] of all videos taken at that location. One

registered video, typically one of the videos corresponding to the peak spatial frequency response

(usually 10 – 30 cycles/deg), was temporally summed to create a high SNR fluorescence image of

the ganglion cell layer, and individual ganglion cells were segmented using the open-source

software GIMP. Typically, all identifiable cells in the focal plane were segmented, though cells

were excluded if the boundaries between two or more adjacent cells were unclear. The

segmentation mask from GIMP was exported to MATLAB, where it was applied to all registered

videos at that location to isolate data from individual RGCs. For every frame, the mean of the

signal within each cell mask was computed to produce an average signal time course for each cell.

When the same stimuli were repeated across multiple sessions, the signal time courses for each

cell were averaged across all sessions.

For the chromatic data, the temporal speed of the flicker (0.15-0.20 Hz) was well within the

temporal resolution of the GCaMP6, thus cells responding to the stimulus exhibited a quasi-

sinusoidal modulation in fluorescence level where the amplitude and phase depended on the

receptive field properties of the cell and the frequency of the response depended on the stimulus

frequency. Each cell's time course was filtered using a Hann windowing function before being

Fourier transformed into the frequency domain. The Fourier amplitudes were normalized by

subtracting the mean noise at higher frequencies (0.32 Hz - 1.08 Hz) and then dividing by the

standard deviation of the noise to produce a response metric equivalent to the sensitivity index

d'. The metric was computed for each stimulus for every cell, so that the cells could be plotted in

two different configurations from the literature [e.g. 60]. In animals M1 and M2, where an L-M

nominally isoluminant stimulus, an S-isolating stimulus, and an achromatic/luminance stimulus

were used, no further analysis was needed, and the data was simply plotted along S vs. L-M axes.

In animal M3, since single cone isolating stimuli were used for L, M, and S cones, it was possible

to reproduce cone weighting plots such as those in Derrington *et. al.* [60]. As in that paper, we

calculated the cone weights by dividing the response metric by the cone modulation from the

stimuli and normalized them all such that the sum of all cone weights added to unity.

For the spatial frequency data, the temporal speed of the drifting gratings (6 Hz) exceeded the

temporal resolution of the GCaMP6s indicator [65], thus cells responding to the grating exhibited

an overall increase in fluorescence that plateaued after approximately 10-20 s. The background

level for each cell was calculated as the mean of the first 10 s of fluorescence during the

adaptation period, while the signal was calculated as the mean of the last 10 s of fluorescence

recorded during the presentation of the stimulus. The $\Delta F/F$ metric commonly used in calcium

imaging was calculated for each spatial frequency as the background response at that frequency

subtracted from the signal at that frequency, all divided by the signal at 17 c/deg, a spatial

frequency producing a robust response across the majority of the cells. This function of $\Delta F/F$ with

respect to spatial frequency is what we refer to as a spatial transfer function (STF) for each RGC

measured. An error metric was also calculated for each $\Delta F/F$ measurement by using the partial

differential error propagation formula using variances [89] and the measured variances in the

background response, the signal response, and the signal at 17 c/deg.

Modeling of photoreceptor mosaic and RGC response characteristics. To connect the measured

spatial transfer functions (STFs) to the receptive field (RF) organization of the underlying RGCs,

we employed a computational model that simulates optical, spectral, spatial, and temporal

components of the AOSLO stimulation apparatus, as well as the animal's optics and cone mosaic

structure. The model computed cone mosaic responses to simulations of the stimuli used to

measure the STF and derived an STF model fit for the RGC under study. The model assumed that

cone signals are pooled linearly and instantaneously (ignoring temporal pooling dynamics) by

center and surround mechanisms of an RGC according to a difference of Gaussians (DoG) [90]

spatial profile. The parameters of the DoG (and therefore the cone pooling weights) were

estimated by minimizing the error between the model STF and the measured *in vivo* fluorescence-

based STF. A model scenario where the center mechanism's Gaussian weighting was replaced by

center weighting on just a single cone was also considered. The modeling pipeline, depicted

schematically in Figure 6, was implemented within the Imaging Systems Engineering Tools for

Biology (ISETBio) software framework [91, 92].

The drifting monochromatic sinusoidal gratings (Fig 6a) used to measure RGC STFs via the AOSLO

apparatus (Fig 6b) were modeled as temporal sequences of ISETBio spatial-spectral radiance

scenes, where each scene models one frame of the displayed stimulus. The spectral

characteristics, spatial extent, and the temporal properties of the AOSLO display subsystem were

taken into account in generating the ISETBio scenes. The spectral profile of the monochromatic

beam was modeled as Gaussian shaped with a peak at 561 nm and a full-width-half-max (FWHM)

of 5 nm with a mean irradiance on the retina of 1.29 mW cm^{-2}. The power estimate included

absorption by the lens pigment. The visual stimulus as imaged on the retina had a pixel size of

1.03 µm (0.005 deg) with spatial extent 140 x 140 µm (0.7 x 0.7 deg). All sinusoidal gratings were

modeled with a nominal contrast of 1.0, drifting at 6 Hz with a refresh rate of 25.3 Hz for a duration

of 666 ms (4 cycles).

The animal's optics during AOSLO-based stimulation were modeled as a diffraction-limited optical

system (Fig. 6c) with a small amount of residual defocus blur (Fig. 6d). The diffraction limit was

obtained using the 6.7 mm pupil diameter employed in the experiment, assuming a conversion

between degrees of visual angle and retinal extent of 199 µm/deg. Residual blur in the AOSLO,

which might occur due to a slight defocus of the stimulus with respect to the plane of cone inner

segments in the retina, was modeled by adjusting the Zernike defocus coefficient used to compute

the optical point spread function. Residual blur could also be generated by any off-axis aberrations

in the system, imperfect AO correction, or other sources, so the Zernike defocus coefficient was

used as a proxy for all sources of residual blur. The amount of residual blur was not known *a priori*

and was estimated as part of model fitting as described below. The temporal sequence of ISETBio

spatial-spectral radiance scenes that model the AOSLO stimulus were passed via the simulated

optical system to generate a corresponding sequence of spatial-spectral retinal irradiance images,

which were processed by the cone mosaic model as described next.

An ISETBio model of the animal's cone mosaic (Fig. 6e) was generated from cone density maps

measured during AOSLO imaging, using an iterative algorithm described previously [91]. The

spatial extent of the modeled cone mosaic was 1.3 x 1.3 deg, with a maximal cone density of

270,200 cones mm^{-2} with relative L:M:S cone densities of 0.48:0.48:0.04 (quasi-regular S cone

packing, random L and M cone packing), and without a tritanopic (S-cone free) area at the fovea.

Cones were modeled with Gaussian entrance apertures with a characteristic radius equal to

$0.204 * \sqrt{2} * D$, where D is the inner segment diameter measured during AOSLO imaging [10]. In

this cone mosaic model, cone outer segment lengths and macular pigment all varied with

eccentricity [91], and the cone quantal efficiencies were based on the Stockman-Sharpe (2000)

normalized absorbance measurements [93, 94]. The modeled macular pigment density is taken

as   the   human   values   provided   by   Stockman   *et. al.*   (1999)   [93]

(http://www.cvrl.org/database/text/maclens/macss.htm), and the modeled lens pigment density

is for young human subjects (< 20 years) as reported by Pokorny *et. al.* [95].

To compute a cone's excitation, the spatial-spectral irradiance impinging on the retina was first

spectrally weighted by the product of macular pigment transmittance and by each cone's spectral

quantal efficiency, subsequently integrated over wavelength, and spatially integrated over the

cone's Gaussian aperture. This excitation response was integrated over the temporal duration of

each stimulus frame (39.5 ms), to estimate a spatial map of the expected excitation events count

$E^j(\omega,t)$, for the *j*-th cone in the mosaic, at time *t*, in response to a drifting grating of spatial

frequency ω (Fig 6f). In these simulations, we did not include Poisson noise in $E^j(\omega,t)$, nor did we

introduce positional jitter between the cone mosaic and the retinal stimulus due to fixational eye

movements. In the measured AOSLO data, the animals were anesthetized with eye muscles

paralyzed, and digital tracking was employed for stimuli, so retinal motion relative to stimuli was

minimized.

Assuming that cones are adapted to the mean background irradiance, the spatiotemporal

excitation response $E^j(\omega,t)$ was converted to a spatiotemporal modulation response, $R^j(\omega,t)$, (Fig

6g) by first subtracting the excitation of the cone to the background stimulus E_0^j, and then dividing

by it separately for each cone j, i.e.:

400
$$R^j(\omega, t) = \frac{E^j(\omega, t) - E_0^j}{E_0^j}$$

This operation captures in broad strokes an important effect of the photocurrent generation

process which converts cone absorption events in the inner segment into ionic currents flowing

through the cone outer segment, and which in effect down-regulates the stimulus-induced cone

excitation rate with respect to the background cone excitation rate.

Model ganglion cell responses $RGC(\omega,t)$, were computed from the cone contrast responses by

weighting responses with corresponding center and surround cone weights (Fig 6h), followed by

spatial pooling within the antagonistic center and surround mechanisms (Fig. 6i) as follows:

409
$$RGC(\omega, t) = RGC_c(\omega, t) - RGC_s(\omega, t)$$

410
$$= \sum_j W_c^j * R^j(\omega, t) - \sum_j W_s^j * R^j(\omega, t)$$

where W_c^j and W_s^j are the weights with which the center and surround mechanisms respectively

pool the responses $R^j(\omega,t)$. We did not model temporal filtering or delay between center and

surround responses. Although real RGC responses may be affected by both temporal filtering and

a center-surround delay, measurements in this study only recorded the amplitude of responses

at one temporal frequency with no response phase information, so we could not meaningfully

estimate temporal RGC parameters.

To make computation of cone weights more tractable, we assumed that the spatial distribution

of cone weights to the center and surround mechanisms had concentric, radially-symmetric

Gaussian profiles (multi-cone RF center scenario), or that the center drew on a single cone with a

Gaussian surround (single-cone RF center scenario):

422
$$W_c^j = \begin{cases} k_c * \exp\left[-\left(\dfrac{d_j}{r_c}\right)^2\right] & (multi - cone\ RF\ center\ scenario) \\ k_c & (single - cone\ RF\ center\ scenario) \end{cases}$$

423
$$W_s^j = k_s * \exp\left[-\left(\dfrac{d_j}{r_s}\right)^2\right]$$

where d_j is the distance between cone-j and the spatial position of the center mechanism of the

model RGC (which is taken as the geometric centroid of the locations of the cones driving the

center mechanism). The parameters k_c, k_s, r_c, and r_s, which represent the center and surround

peak sensitivities and characteristic radii, respectively, of the DoG RF model (Fig. 6 l,m) were

determined by minimizing the root mean squared error (RMSE) between the model-predicted

STF, $STF^m(\omega)$, and the measured STF, $STF^{\Delta F/F}(\omega)$, (Fig. 6k) accumulated over all spatial frequencies

ω:

431
$$RMSE = \sqrt{\frac{1}{N} * \sum_{\omega=\omega_1}^{\omega_n} \frac{\beta(\omega)}{\epsilon(\omega)} * [STF^m(\omega) - STF^{\Delta F/F}(\omega)]^2}$$

where $\epsilon(\omega)$ is the standard error of the mean of the $STF^{\Delta F/F}(\omega)$ measurement, and $\beta(\omega)$ is a high-

spatial frequency boost factor which linearly increases with spatial frequency from 0.1 to 1.0 over

the spatial frequency range 4.7 to 49 c/deg. This boost factor was introduced to emphasize high

spatial frequency measurements, the stimulus regime which is maximally informative about the

properties of the center mechanism.

In some cells, and for certain recording sessions, the measured $STF^{\Delta F/F}(\omega)$ dropped below zero for

the lowest spatial frequencies. When this occurred, it was accompanied by an apparent overall

downward shift in STF$^{\Delta F/F}(\omega)$ across all frequencies, so we assumed that in such cases the

background fluorescence was overestimated. To compensate for this, we fit the model to

STF$^{\Delta F/F}(\omega)$ − min(STF$^{\Delta F/F}(\omega)$) instead of STF$^{\Delta F/F}(\omega)$ in cases where STF$^{\Delta F/F}(\omega)$ dropped below zero

for any spatial frequency.

To reduce the chance of the minimization algorithm getting stuck at a local minimum of the error

function, we employed a multi-start minimizer which was run 512 times, keeping the results from

the starting point with the minimum RMSE.

The model-predicted STF, STF$^m(\omega)$, was computed by fitting a sinusoidal function to $RGC^m(\omega, t)$,

$$A(\omega) * \sin[2\pi f t - \theta]$$

where f is set to the temporal frequency of the drifting gratings, and θ, A(ω) are free parameters.

At each spatial frequency, the amplitude of the fitted sinusoid, A(ω), was taken as STF$^m(\omega)$ (Fig.

6j).

To interpret the measured STFs, we considered four different model scenarios: single cone RF

centers with and without residual defocus in the AOSLO apparatus, and multiple cone RF centers

with Gaussian weighting, with and without AOSLO residual defocus. For each scenario, the model

was fit to the STF data of an examined RGC at multiple positions within the model cone mosaic

and the final model was selected as the one with the minimum RMSE over the examined positions.

The multi-position model fitting/selection was performed to take into account local

inhomogeneities of the cone mosaic as exact RF center of each recorded RGC is only known to

reside approximately within the central 40 microns. Details of the model fitting differences

between each scenario, model training, and comparisons between the modeled cone mosaic and

the measured mosaic from M3 are provided in the Supplement (see **S4 Fig, S5 Fig, S6 Fig, S7 Fig**).

All figures in the main body of the paper assume the single cone center with fixed residual 0.067

D defocus scenario, unless otherwise noted.

Results

***In vivo* functional imaging of foveal RGCs.** In this study, optical stimulation of foveal cones elicits

the activity of retinal ganglion cells in the living macaque eye. Intravitreal injection of an adeno-

associated virus (7m8 or AAV2) with a promoter (ubiquitous CAG or RGC-specific) produced

expression of the calcium indicator GCaMP6f [65] in animal M1 and GCaMP6s [65] in animals M2

and M3. As previously reported, injections in all three animals produced expression in a ring of

RGCs surrounding and radiating out from the foveal avascular zone where the internal limiting

membrane is thin [66, 68, 96] (Fig. 1a). Stimuli were presented to the centermost foveal cones

while a 488 nm imaging laser stimulated fluorescence from GCaMP expressing RGCs in a

rectangular area placed either nasally or temporally relative to the cones. Fig. 2a-b show the

fluorescence response of a single RGC to a simple luminance flicker stimulus in time as well as the

resulting Fourier transform, which shows the cell responding primarily at the stimulus frequency.

Figure 2c shows that, under our stimulus paradigm, the imaging light did not stimulate foveal

cones and confound the recorded functional responses after a 561 nm adapting light was

presented to the foveal cones. In each animal, hundreds of RGCs were recorded simultaneously

using this paradigm, and we used the precision of the AOSLO system to return to the same cells

over multiple sessions spanning weeks to months.

Fig 1. Stimulus presentation and recording paradigm. In **(a)** the general stimulation paradigm is

shown. Background is an image of the retina of M3 using the blue reflectance imaging modality

of a Heidelberg Spectralis instrument. In false color green are GCaMP-expressing cells from M3

imaged using the blue autofluorescence modality of the same Spectralis instrument. When using

the AOSLO instrument, videos are captured within a rectangular field of view (white dashed line)

while stimuli (white circle) such as cone-isolating flicker or drifting gratings (which were

rectangular in shape and covered approximately the same area as the white circle) are presented

to and centered on the centermost foveal cone photoreceptors. A 488 nm imaging laser in the

AOSLO (blue rectangle) excites GCaMP-mediated fluorescent responses of ganglion cells

surrounding the foveal slope. **(b)** shows the recorded cone densities at the fovea of M3 and **(c)**

shows the ganglion cells segmented (green) at the innermost edge of the foveal slope in M3.

Fig 2. Data collection from ganglion cells *in vivo*. (a) shows the smoothed time course of a single

RGC fluorescence response to a 0.15 Hz luminance flicker stimulus. **(b)** shows the amplitude of

the Fourier transform of the response time course, recovering the response peak at the stimulus

frequency of 0.15 Hz (red dashed line). **(c)** shows the effect of the imaging light on cone-mediated

response to stimuli in 62 foveal RGCs. The 488 nm imaging laser which excites GCaMP

fluorescence can scatter to foveal cones and confound cone stimulation. If the imaging light is

turned on without adaptation to any light, there is a quick rise and slow falloff in fluorescence due

to cone responses feeding RGCs. However, if the 561nm stimulus laser (used for drifting gratings)

or LED white point (used for chromatic stimuli) is presented at a mean luminance of 2-2.5 μW

during a short 30s adaptation period prior to imaging onset, then there is no corresponding effect

of the imaging light confounding the fluorescence signal. In all recordings, an adaptation light

preceded each recording to prevent this transient effect of scattered light from the imaging laser.

Reliability and precision of foveal RGC recordings. Across multiple sessions, RGC somas were

recorded from over a range of soma locations between 1.2 and 3.5 degrees from the center of

the foveal avascular zone (FAZ). When averaging data across multiple experiments, only cells that

were visible in all experiments were used. In M3, a follow-up experiment examining only cone

density (Fig. 1b) found the center of the FAZ to be roughly 41 microns inferior and 51 microns

nasal from the peak cone density which agrees with the range of data reported from humans [14].

Based on detailed receptive field mapping performed by McGregor *et. al.* [16], the cones driving

the receptive fields of the cells examined in all three animals are likely to be located at

eccentricities less than 36 arcmin from the foveal center, or a retinal radius of 120 microns

calculated using the primate model eye [97] with the axial lengths of each animal (M1: 17.51 mm,

M2: 17.20 mm, M3: 16.56 mm). According to the same mapping, the receptive fields of the

innermost cells we recorded from in M3 (Fig. 1c) are likely to be driven by cones found at

eccentricities less than 6 arcmin, or a retinal radius of 20 microns calculated using the same

primate model eye and the axial length of M3 (16.56 mm). All visual stimuli delivered to foveal

cones were large enough to cover both the FAZ center and the location of maximum cone density

and were some 10-60% larger than the cone area servicing even the furthest cells we recorded

from, ensuring the best possible chance of recording responses from the maximum number of

RGCs.

Functional measurements of RGCs made in the past often involved flashed stimuli and marking

approximate estimates of soma location on fundus images of the retina [e.g. 9, 98], a method that

leads to uncertainty about precise locations. In contrast, our imaging method captures

photoreceptors, vasculature, and RGC somas in the same field of view, allowing more accurate

measurements of receptive field eccentricity and soma position. One shortcoming of *in vivo*

functional imaging compared to traditional electrophysiology is that the excitation and imaging

sources can interfere with stimulation or fluorescence detection. However, unlike in the

periphery, foveal RGCs are displaced laterally from their photoreceptor inputs [1, 11, 15, 99].

Thus, our paradigm avoids this pitfall, and we can reliably image foveal RGCs without the 488 nm

excitation light stimulating foveal cones (Fig. 2c) or the photoreceptor stimuli interfering with the

RGC GCaMP fluorescence imaging.

Chromatic tuning of foveal RGCs. For functional classification of RGCs with receptive fields at the

foveal center, chromatic tuning was measured across multiple imaging sessions. Amplitude and

phase responses (Fig. 3a-d) to spatially uniform 0.15 Hz or 0.20 Hz flicker modulations that were

directed along a combination of L, M, L-M, S, and achromatic/luminance directions in color space

enabled reliable identification of chromatic tuning of RGCs in all three animals. For each cell, the

raw fluorescence time course was measured in response to each visual stimulus, and the Fourier

transform was used to find the cell's response amplitude at the stimulus frequency. The signal-

to-noise (SNR) response metric was calculated as the response Fourier amplitude (signal) minus

the mean amplitudes of higher spatial frequencies (noise), all divided by the standard deviation

of the noise. This analysis revealed several functional groups including L-M and M-L opponent

cells, S-ON and S-OFF cells, ON and OFF luminance cells, and cells with mixed L vs. M ± S responses.

In M3, single cone isolating stimuli were presented to the fovea and the individual cone weights

feeding RGCs were calculated as was done previously in LGN [60] (Fig. 4a-b). This analysis

estimates the relative contribution of each cone class to the response of a cell and revealed

further diversity of functional groups, including cells with responses to only L or M cones, as well

as various S cone connections to cells with either L or M cone input that were not L vs. M

opponent.

Fig 3. Chromatic response properties of RGCs from three different animals. Black dots show

luminance only responding cells, red dots show L-M and M-L cone opponent cells, blue dots show

S responding cells, while magenta dots show cells that responded to both L-M and S stimuli.

Dotted lines show the significance cutoff of SNR = 2 calculated based on cell responses to the

control stimulus (see Methods). Cell response is captured by our signal-to-noise metric described

in the Methods. Response phases were also calculated but are not directly shown; however,

positive SNRs correspond to responses that were in phase with the stimulus (ON response), while

negative SNRs correspond to responses that were approximately 180 deg out of phase with the

stimulus (OFF response). The luminance axis is perpendicular to the plane shown and is not

included for readability and because not all cells responded to the luminance stimulus. Luminance

only cells shown in black thus include both ON and OFF cells. In **(a)**, responses from 15 cells in M1

are shown after presentation of spatially uniform luminance, L-M isoluminant and S-isolating

stimuli at 0.2 Hz. In **(b)**, responses from 126 cells in M2 are shown after presentation of spatially

uniform luminance, L-M isoluminant and S-isolating stimuli at 0.2 Hz. In **(c)**, responses from 83

cells (a subset of the cells in Fig 4a) in M3 are shown after presentation of spatially uniform

luminance, L-isolating, M-isolating, and S-isolating stimuli at 0.15 Hz. In **(d)**, responses from 22

cells (a subset of the cells in Fig 4b) serving the very central fovea of M3 are shown after

presentation of spatially uniform luminance, L-isolating, M-isolating, and S-isolating stimuli at 0.15

Hz. These cells are closer to the foveal center than any of the cells shown in (c). In **(c-d)**, all stimuli

were single-cone isolating, so the L-M axis is calculated from the single cone L-isolating and M-

isolating responses and their relative phase.

Fig 4. Cone weights of RGCs from M3. The horizontal axis shows L cone weight, the vertical axis

the M cone weight, and the distance from the origin shows the S cone weight where further away

from the origin corresponds to less S-cone weighting. In **(a)**, 100 RGCs (a superset of the cells in

Fig 3c) were recorded across the entire width of the GCaMP-expressing ring in M3. There were 3

cells that only showed S cone responses, represented by overlapping circles at the origin. There

were also 4 L+M and 11 -L-M luminance only cells that are represented by asterisks in the upper

right and lower left. Because these cells only responded to the luminance stimulus but not the L-

isolating or M-isolating stimulus, the response sign is known but not the exact L or M cone

weighting. In **(b)**, 34 cells (a superset of the cells in Fig 3d) at the innermost edge of the foveal

slope in M3, served by the centermost foveal cones, were recorded from. No consistent S only

responses were observed, and there were 5 L+M and 2 -L-M luminance only cells.

Identification of foveolar midget RGCs using chromatic tuning. Across all three animals, cells with

L-M or M-L chromatic opponent responses (and no response to S-cone stimuli) were identified as

putative midget RGCs. There were a large number of these cells as can be seen in Figures 3,4: the

proportions of these putative midget cells were 44% (95% CI true proportion of 44±17%) of

responsive RGCs closest to the fovea (1.2-1.8 deg soma locations) in M3 (small imaging FOV), 33%

(95% CI true proportion of 33±9%) of responsive RGCs over the entire measured range in M3

(large imaging FOV), 72% (95% CI true proportion of 72±8%) of responsive RGCs in M2 (large

imaging FOV), and 47% (95% CI true proportion of 47±25%) of responsive RGCs in M1. It is

expected that midget RGCs comprise some 70-90% of foveal cells [20, 51, 99], and the confidence

intervals in M1 and M2 contain values in that range. In M3, single cone isolating stimuli were

used as detailed in the chromatic stimuli section—it is expected that these stimuli produce lower

responses in the classical cone opponent model of midget RGCs by not driving the center and

surround in counterphase as would the isoluminant L-M stimulus used in M1 and M2. There were

many cells in M3 (an additional 35% of the closest foveal RGCs, and an additional 32% over the

entire measured range) with only an L cone response or only an M cone response and no S cone

response. It is possible that some of these cells are cone opponent, but that the single cone

isolating stimuli produced too small of a response from the antagonistic surround in those cases.

Based on this uncertainty, we cannot say for sure whether the observed number of putative

midget RGCs in M3 is outside the range of expected values.

Soma size is a useful anatomical measure for distinguishing more peripheral ganglion cell types

and was examined here as a potential correlate with our putative midget RGCs. RGC soma sizes

were measured by two different observers see (Supplement **S1 Fig.**) using different methods, but

as others have reported, soma sizes did not exhibit bimodalities near the fovea [46]. Soma size

also did not distinguish between functional groups identified by chromatic tuning (Wilcoxon

ranksum test, $p > 0.1$ for comparisons between L-M, S only, luminance only, and L-M/S cells).

Spatial tuning of foveolar putative midget RGCS. Spatially patterned drifting gratings were used

to probe the spatial frequency response of RGCs and to augment measurements of chromatic

tuning (Fig. 5). In M3, at the small field of view focused on the innermost RGCs, a control and

drifting gratings of fourteen different spatial frequencies (561 nm, 6 Hz, 4-49 c/deg) were used to

stimulate foveal cones while the 488 nm imaging light elicited fluorescence from the same GCaMP

expressing RGCs shown in Figure 3d and Figure 4b. At this temporal frequency, GCaMP could not

track modulation, and responding cells exhibited a sustained increase in fluorescence in response

to photoreceptor stimulation. A ΔF/F metric was calculated for each cell as the change in

fluorescence under stimulation divided by the mean fluorescence in response to a grating of 17

c/deg (a spatial frequency with a robust response across the majority of the cells). Most of these

putative midget RGCs at the foveal center in M3 (Fig. 5), which were all identified according to

their L-M or M-L cone opponency, showed two salient features when their spatial frequency

response functions were measured: a consistent and very strong low frequency cut, and a peak

in the higher spatial frequencies (25-40 c/deg).

Fig 5. Spatial frequency responses of putative midget cells. Fifteen cells in M3 serving the central

foveal cones were identified as L-M or M-L opponent (Fig. 3d). Each cell is numbered L1-11 or M1-

4 based on whether the suspected center cone was an L or M cone. The green points (squares),

blue points (triangles), and red points (circles) are measurements from three different

experiments each one week apart. The solid orange line in each panel is the average response for

that particular cell across all three experiments. The vertical axis is a response metric where the

mean baseline fluorescence is subtracted from the fluorescence under stimulation and divided by

the fluorescence under stimulation at the peak spatial frequency (see Methods). The error bars

show the standard error at each data point, which represents the error in obtaining the signal and

background needed to calculate the metric. Additional spatial frequency responses of cells from

the large imaging FOV (**S2 Fig**) and non-midget RGCs (**S3 Fig**) can be found in the supplement.

Modeling of foveal midget RGCs. Using the ISETBio-based approach described in the Methods

and depicted schematically in Figure 6, we derived estimates of the spatial RF organization from

the measured STF data under four different model scenarios involving combinations of residual

instrument defocus (zero or some residual defocus) and RF center composition (single or multiple

cone inputs). Model cone mosaics matched well in both density and cone size with the measured

data in M3 along the horizontal and vertical meridians (see supplement **S4 Fig**). Modeling results

from one RGC are depicted in Figure 7. Figure 7a shows results from the modeling scenario that

assumes a single cone in the RF center and zero residual defocus. Note that this scenario predicts

a diffuse surround and that the derived STF does not agree well with the measured STF, resulting

in the largest RMSE across the four examined model scenarios. The STFs derived by the remaining

three modeling scenarios match the measured STF approximately equally well even though the

resulting cone pooling weights differ. The single cone RF center with 0.067 D residual defocus (Fig.

7b) results in a more focal surround than would be true if there were no residual defocus. The

multi-cone center RF with zero residual defocus (Fig. 7c) results in a diffuse RF center and a

surround that is marginally more diffuse than the center, which is not consistent with known

anatomy of foveal midget RFs. Finally, the multi-cone center 0.067 D residual defocus scenario

(Fig. 7d) results in an RF center with heavy input from one cone and very weak input from nearby

cones, and a focal surround. This scenario essentially collapses to the single cone center/0.067 D

residual defocus scenario. In Fig. 7e-f, the single and multi-cone RF center scenario performances

are depicted as a function of assumed residual defocus. Note that for the particular cell shown,

the minimal error in the single cone center scenario was at 0.077 D of residual defocus, and that

the multi-cone scenario never performs better than the best single cone scenario at all residual

defocus values modeled. Across all cells examined, the best performance for the single cone RF

center scenario was obtained for values of residual defocus between 0 D and 0.082 D with a mean

value of 0.067 D. The comparison shown for one cell in Fig. 7e-f is provided for all cells modeled

in Supplement **S6 Fig**. In addition, a cross-validation analysis (Supplement **S5 Fig**) similarly

suggests that across all cells, the data do not reject the single cone RF center/0.067 D model

relative to the multi-cone center models. Therefore, for the remainder of this paper, we use the

single cone RF center/0.067 D residual defocus scenario as the vehicle for interpreting the

measured STFs.

Fig 6. Schematic overview of the ISETBio modeling approach. To interpret the AOSLO-based STF

measurements (Fig. 5) in terms of the spatial pooling of cone signals to the measured RGCs we

model the spatial, spectral and temporal aspects of the stimulus, the optics, the cone mosaic and

the cone pooling within the center and surrounds mechanisms of an RGC using the ISETBio

toolbox. **(a)** A frame of the drifting monochromatic grating stimulus which is projected through

the AOSLO system to stimulate and measure the GCaMP fluorescence – based spatial transfer

function (STF) of RGCs, $STF^{\Delta F/F}(\omega)$ **(b).** The drifting grating is modeled as a temporal sequence of

spatial spectral radiance scenes. These scenes model the AOSLO display including a diffraction-

limited optical system with a 6.7 mm pupil **(c)** combined with residual defocus in the eye's wave

aberrations **(d).** The resulting sequence of retinal irradiance maps are spectrally integrated and

spatially averaged within the apertures of cones in a model of the animal's cone mosaic **(e)**,

resulting in a spatiotemporal sequence of cone excitations **(f).** Cone excitations are transformed

to cone modulations by first subtracting and subsequently dividing the mean cone excitation level

(g). Modulation signals from different cones are scaled by the spatial weights with which the RF

center and the RF surround mechanism pool cone signals **(h).** The weighted cone signals are

subsequently summed **(i)** to compute the temporal response of the model RGC. A sinusoid is fitted

to the RGC temporal response **(j)** and the amplitude of the sinusoid is taken as the STF of the

model RGC at the stimulus spatial frequency, $STF^m(\omega)$. The RMSE between $STF^m(\omega)$ and $STF^{\Delta F/F}(\omega)$

(k) is minimized using a multi-start solver which optimizes the center and surround peak

sensitivity and radius parameters **(l)** of a difference of Gaussians (DoG) RF model **(m).** The model

yields the best-fitting cone pooling weights for the RGC.

Figure 7. Model interpretation of measured STF for a single RGC. a-b show single cone center

receptive field models with zero and 0.067 D residual defocus respectively. **c-d** show multi cone

center receptive field models with zero and 0.067 D residual defocus respectively. In a – d, the

first and second columns depict cone pooling weights for the center and surround mechanism,

derived by fitting the model STF to the measured STF. The third column depicts the integrated

(along the y-axis) center and surround sensitivity profiles, and the fourth column depicts the

model-predicted STF (orange lines) together with the measured STF (green disks). **e-f** show single

and multi-cone model residual errors for this particular cell with varying levels of residual defocus.

The best residual defocus for individual cells in the single cone center receptive field modeled

varies slightly, so 0.067 D was chosen as the value optimizing models across all cells measured.

Figure 8 depicts the results of the single-cone RF center/0.067 D scenario for four cells from the

center of the fovea of M3. These cells were chosen to showcase the model's fitting to slightly

different STF shapes and are not necessarily special in any way. Full results of the model fitting to

all 15 cells, including fits for all four scenarios examined, can be found in Supplement **S7 Fig**.

Figure 8. ISETBio model receptive field estimates for four additional RGCs. Each row shows the

model fits to a different cell's STF using the single cone center/0.067D residual defocus model.

The ISETBio model expresses the RFs directly in terms of the weights with which they pool signals

from cones in the mosaic, and as such is appropriate for comparison with anatomical

measurements and with *in vitro* physiological data where the optics of the eye are not in the

stimulus light path. In order to compare our data to other *in vivo* data collected in macaque [e.g.

9], where the eye's optics form a blurred image of the visual stimulus, we predicted *in vivo* STFs

using the cone pooling weights obtained from the single cone RF center/0.067 D residual defocus

scenario. To do so, we swapped the AOSLO optics with the animal's (M3) own wavefront

aberration optics as measured *in vivo* and assumed a pupil diameter of 2.5 mm. These

physiological optics STFs were computed for 100% contrast achromatic sinusoidal gratings, which

matched in their spatial and temporal properties the monochromatic gratings used to measure

the STFs in the AOSLO. Results of this analysis are displayed in Figure 9, with three example cells

shown in Figure 9a. Here, gray disks and black lines depict the AOSLO STF data and corresponding

fit of the model for the AOSLO measurement conditions (single cone RF center/0.067 D residual

defocus). Red disks depict the STFs computed using the model combined with M3's optics, and

red lines depict a simple DoG model fit to the computed STFs, as is typically done in analysis of *in*

vivo data. Note that the STFs predicted under physiological optics differ substantially from the

STFs obtained under near diffraction-limited optics: their magnitude is lower, and their shape is

less bandpass. They have weaker attenuation at low frequencies and peak at lower spatial

frequencies. This is due to the differential effect of the optical modulation transfer function (MTF)

to the STF of the center and surround mechanism, as illustrated in Figure 9b. The left panel depicts

the STFs of the center (pink) and surround (blue) mechanism of a model RGC under diffraction-

limited optics, and the gray disks represent the corresponding overall STF of the cell. The middle

panels depict MTFs for three hypothetical physiological optical systems with increasing amounts

of Gaussian blur. The physiological STFs that would be obtained under the three MTF conditions

are depicted in the right panels of Figure 9B. Note that as the physiological optics become more

blurred, the mapped STF becomes less bandpass and its peak shifts towards lower spatial

frequencies. This occurs because the difference between center STF and surround STF decreases

as the optics become more blurred. Also note the overall decrease in the magnitude of the STF

predicted under physiological optics conditions.

Figure 9. Effect of physiological optics on RGC STFs. a. Gray disks depict STF data measured using

the AOSLO for three RGCs, and the black lines show the STFs from the ISETBio model fit to these

cells. The cone weights from the model fits are used to predict the STF that would be measured

under the animal's own physiological optics (as characterized by wavefront-aberration

measurements taken during the experiment) with a 2.5 mm pupil. These model-predicted STFs

are depicted by the red disks. A simple DoG model is then fitted to the predicted STF data (red

line) for comparison to measurements obtained in traditional *in vivo* neurophysiological

experiments (e.g. Croner & Kaplan [9]). **b. Demonstration of the effect of physiological optics on**

the STF. The left panel depicts the STFs of the RF center and the RF surround of a model RGC as

they would be measured using diffraction-limited optics (pink and blue, respectively), whereas

the composite STF is depicted by the gray disks. The middle panels depict the MTFs of three

hypothetical physiological optical systems with progressively larger Gaussian point spread

functions. The corresponding STFs that would be measured under these physiological optical

systems are depicted in the right panels. Note the difference in scale between left y-axis (for the

center and surround STFs) and right y-axis (for composite STFs).

To compare our data with published results on midget RGC STFs obtained *in vivo* from outside the

central fovea, we fit the DoG model to the synthesized STFs obtained under the animal's optics

and we contrasted the DoG parameters derived using the procedure illustrated in Figure 9a to

those reported by Croner & Kaplan [9]. Figure 10 shows the characteristic radii of centers and

surrounds of data from M3, as predicted with physiological optics, along with data from Croner &

Kaplan. Cone characteristic radii as measured in M3 (blue circles) along with those obtained in

Macaca nemestrina from Packer *et al.* [100] (blue dashed line) are also shown. The midget RGC

center and surround characteristic radii from M3 lie along a reasonable extrapolation from the

Croner & Kaplan data which overall follows the general slope of the decreasing cone characteristic

radii. This supports the hypothesis that the midget pathway is specialized to preserve information

at the level of the cone photoreceptor mosaic, and that midget RGCs at the foveola transmit high

acuity chromatic information from retinal images.

Figure 11a depicts population data for R_c/R_s, the ratio of center to surround characteristic radii,

which typically shows a smaller center size compared to the surround in the Gaussian model and

can vary across RGC types. The data of Croner & Kaplan [9], which were obtained from a wide

range of eccentricities, are depicted in gray, whereas data from our foveal RGCs are shown in blue

(AOSLO optics) and yellow (M3 as predicted for physiological optics). Note that our population of

foveal RGCs has a higher mean R_c/R_s ratio (0.32 under AOSLO, 0.45 under M3 optics) than the

population of Croner & Kaplan (0.15), which is not surprising since foveal RGCs are expected to

have smaller surrounds than peripheral cells [6, 8, 20].

Figure 11b depicts population data for K_s/K_c, the ratio of surround to center peak sensitivity, a

measure of the surround compared to the center strength weighting, where typically it has been

found that the surrounds are much weaker than the centers when expressed in terms of peak

sensitivity [9]. Again, there is a difference between our foveal cells with mean K_s/K_c values of 0.28

under AOSLO optics and 0.11 under M3's optics, whereas the Croner & Kaplan data have a mean

of 0.0094. To the extent that the midget RGC data from M3 and the Croner & Kaplan study [9] are

comparable, this suggests that the surround strength expressed as a peak sensitivity ratio is

stronger at the foveola than at more eccentric locations, supported by the general trend of the

K_s/K_c ratio as a function of eccentricity in both the Croner & Kaplan and M3 data.

Figure 11c provides the surround/center integrated sensitivity ratio, which is essentially the ratio

of the total sensitivities (in a volumetric sense) of surrounds and centers. This does not vary with

eccentricity in the Croner & Kaplan data [9], as the variation in R_c/R_s and the variation of K_s/K_c

counteract each other such that the integrated sensitivity ratio is relatively constant across midget

cells. Consistent with this, the integrated sensitivity ratio for M3 predicted for physiological optics

conditions is similar to that found in Croner & Kaplan's data (means of 0.71 and 0.54 respectively).

Note also the large effect that the optics have on this value, which rises to 3.37 under the

measured AOSLO conditions. Overall, this population analysis shows that our foveal RGCs have a

large degree of surround antagonism when the receptive field is expressed directly in terms of

how the center and surround draw on the cones in the mosaic, but that this antagonism is greatly

attenuated under physiological optics viewing conditions, resulting in center/surround structure

in our data that is consistent with what has been observed in previous midget RGC studies across

a wide range of eccentricities.

Figure 10. Comparison to extra-foveal *in vivo* midget RGC data. Data from M3 model fits are

compared with extra-foveal data from Croner & Kaplan [9]. Gray circles and squares respectively

show center and surround characteristic radii from Croner & Kaplan as a function of eccentricity,

while pink circles and green squares show characteristic radii of centers and surrounds from M3

predicted using the ISETBio model with physiological optics accounted for. Light blue dashed line

shows cone characteristic radii data from Packer *et. al* [99]., while blue dots represent cone

characteristic radii measured in foveal cones of M3.

Figure 11. Relationship between our findings in foveal midget RGCs and findings in midget RGCs

from earlier studies. Data from Croner & Kaplan [9] is shown in gray, data from M3 using the

ISETBio model and physiological optics is shown in yellow, and data from M3 using the AOSLO

optics is shown in blue. **a.** Analysis of Rc/Rs ratio. Left panel: relationship between the Rc/Rs ratio

and retinal eccentricity. Right panel: distribution of Rc/Rs. The mean Rc/Rs ratio in the Croner &

Kaplan data is 0.15, vs. 0.45 in our foveal RGCs under physiological optics, and 0.32 in our foveal

RGCs under AOSLO optics. **b.** Analysis of Ks/Kc ratio. The mean Ks/Kc ratio in the Croner & Kaplan

data is 9.4e-3, vs. 1.1e-1 in our foveal RGCs under physiological optics, and 0.28 in our foveal RGCs

under AOSLO optics. **c.** Analysis of integrated surround-to-center ratio, (Ks/Kc) x (Rs/Rc)^2. The

mean integrated surround-to-center ratio in the Croner & Kaplan data is 0.54, vs. 0.71 in our

foveal RGCs under physiological optics, and 3.37 in our foveal RGCs under AOSLO optics.

Discussion

In this study, we repeatedly imaged the same RGCs serving the foveal center *in vivo*, in three

animals, returning to the same retinal locations each time. Spatially uniform chromatic visual

stimuli enabled the identification of L-M and M-L opponent cells which were further interrogated

in one animal with drifting gratings of various spatial frequencies. Detailed modeling revealed

how the receptive fields of these RGCs draw on the cones of the retinal mosaic. The L vs. M

opponency and high spatial frequency responses of these cells identifies them as candidate

midget RGCs. Our modelling is consistent with the hypothesis that these cells have single cone

center inputs and are specialized to preserve the resolution of the cone photoreceptor mosaic in

the signals they send to the visual cortex.

Measuring the precise location of cells from the foveal center. In our data, the foveal avascular

zone (FAZ) was used as the 'center' measurement of the fovea, as the imaging paradigm precluded

using the location of maximum photoreceptor density for all sessions. The preferred retinal locus

of fixation (PRL) is also of interest but could not be determined using our method, since the

animals are anesthetized during the experiments and cannot freely fixate. The center of the

ganglion cell ring surrounding the fovea is also sometimes used to define the center of the fovea,

but this measure is highly correlated with the center of the FAZ [11, 101]. A more sophisticated

measure of the 'physiological center' can be identified using white noise stimuli as in McGregor

et. al. [16], but those stimuli were not used in this study. Others have characterized the precise

nature of the relationship between these various measures [14], and as noted in the results, our

measurements of the difference between the center of the FAZ and the location of maximum

cone density in M3 are in agreement with that literature. In any case, the difference in

identification of the 'foveal center' is not large enough under our method to affect the conclusion

that, using the mapping of McGregor *et. al.* [16], the receptive fields we examined are located

near the very center of the foveola.

S-cone mediated color vision in the fovea. As shown in Figures 3,4, cells with either S cone input

without L vs. M cone opponency (pure) or S cone input on top of L vs. M opponency (mixed) were

observed in each animal. 40% of responsive cells exhibited pure S responses and 7% mixed in M1,

9% pure and 14% mixed in M2, 16% pure and 3% mixed in the near fovea of M3 (large imaging

FOV), while no consistent S cone responses were recorded from the centermost foveolar cells

(small imaging FOV) in M3. Other than in M1 where the sample size was only 15 cells, these

numbers are consistent with the expected S cone density of approximately 10% across primate

fovea [102] and the known reduction of S cone density at the very center of the primate fovea [8,

15]. Others have reported subtypes of midget RGCs with S cone input or even that midget RGCs

may indiscriminately contact S cones along with L and M cones in the surround [4, 17, 48, 103,

104]. Our current data set has no repeatable S cone responses for RGCs at the innermost foveal

edge in M3, where RGCs are optically well-isolated, so we cannot rule out that the mixed cells we

measure are partially or fully the product of optical crosstalk resulting from the 25-30µm axial

resolution of the AOSLO instrument.

Putative parasol cells at the fovea. As shown in Figures 3,4, cells with no chromatic responses but

only a L+M or -L-M luminance response were recorded from each of the three animals: 7% of

responsive cells in M1, 4% in M2, 15% in the entirety of M3 and 20% of the centermost foveal

cells in M3, consistent with expected proportions of parasol ganglion cells, which are known to

mediate achromatic vision [20]. Simple modeling based on hexagonal packing of cones and

random wiring of 6 cones to midget surrounds suggests that approximately 9% of midget RGCs

could be dominated by a single cone type and appear achromatic (Supplement **S8 Fig.**). It is

probable then that some of these achromatic cells are midget RGCs but that most of them are

candidate parasol cells, though at this time we cannot make that distinction for an individual cell.

The spatial frequency response properties of these cells were even more varied than the

chromatic opponent cells (Supplement **S3 Fig.**), suggesting that they might be responsible for a

wide range of achromatic or even temporal vision properties that complement the midget system.

As noted in the results, we have not found significant differences between soma sizes of putative

foveal parasol and midget RGCs, so additional anatomical information such as histological studies

may be required in the future to confirm functional designations.

The role of midget RGCs in foveal vision. Among the putative midget RGCs from M3 (Fig. 5,7-9),

there was some variability among the responses to spatial frequencies in that a subset of cells

possessed lower spatial frequency response peaks (0-20 c/deg) compared to the peak spatial

frequencies of the majority of the cells (20-40 c/deg), though both of these groups of cells were

well fit by the single-cone RF center/0.067 D residual defocus model scenario, suggesting that the

change in their peak spatial frequency is related to the surround size and/or ratio of center and

surround strengths, as opposed to the spatial extent of the center. Previous studies examining

parvocellular neurons in the lateral geniculate nucleus (LGN) serving the central five degrees of

retina found an average of 4.57±2.75 c/deg as the peak spatial frequency response [62], and other

studies in the retina found parvocellular cells in the central five degrees to have a peak spatial

frequency response around 3 c/deg [9]. However, these previous studies relied on presenting

stimuli through the animals' optics, whereas the adaptive optics method we use approaches near

diffraction-limited stimulus presentation with some possible residual defocus. Indeed, as shown

in Figures 9,10, when the animal's optics are added back into a simulation of our data from M3,

we recover the classically measured bandpass shape and lower spatial frequency peaks that are

expected of parvocellular neurons (midget RGCs). We thus measure putative midget RGCs not

only closer to the foveal center and with greater precision than previously achievable, but also

with diminished optical blur in the *in vivo* retina compared to previous studies, with the exception

of studies that used interferometric stimuli to bypass the optics altogether [e.g. 10]. We thus posit

that the parameters of spatial DoG models fit to data collected through the optics do not directly

represent the *in situ* characteristics of RGCs, and that future *in vivo* measurements and models

need to take into account the effects of diffractive and optical blurring on the measured spatial

transfer functions.

In all 15 L-M and M-L cone opponent RGCs serving the central foveola, we could not reject the

hypothesis that the receptive fields were best described by a single cone center with some

residual instrumental defocus (0.067 D). There were some cells where the single cone

center/0.067 D scenario was clearly the best fit, but for many cells several of the scenarios

performed similarly in terms of RMSE—full details for all scenarios can be found in Supplement

(**S7 Fig**). We posit that since the fovea is specialized for high acuity vision, it must be capable of

extracting high spatial frequency information from natural scenes which are known to follow a

$1/f^2$ law [105] and are dominated by low frequency information. Our modeling that adds the

optics of M3 back into the predictions (Fig. 9) suggests that natural ocular aberrations reduce the

peak spatial frequency obtained under AOSLO conditions by a factor of three or more, which

means that high spatial frequency information is greatly reduced by ocular transmission. Thus, to

extract meaningful high spatial frequency information from natural scenes through the eye's

native optics, detectors that preserve information at the resolution of the cone photoreceptor

mosaic are required. It seems highly probable that the putative midget RGCs shown in Figures 5,8

are responsible for this resolution, as they have responses consistent with single cone receptive

field centers and most have spatial transfer functions that peak well into the 20-40 c/deg range

when measured with respect to a retinal image under AOSLO conditions. McMahon *et al.* [10]

measured the spatial frequency responses of parvocellular LGN cell centers near the foveola using

interference fringes, which are not subject to blur by diffraction. They found that cells still had

detectable responses to spatial frequencies above 100 c/deg, with some responses that were

complex and inconsistent with a simple difference of Gaussians model. Because the AO-corrected

stimuli used in the present study were subject to blur by diffraction, we were unable to explore

responses at the high spatial frequencies where such effects were observed.

Putative midget RGCs with the lowest spatial frequency peaks comprised only 2/15 cells (cells L2

and M3) compared to ones with higher spatial frequency peaks (13/15 cells). The two cells with

lowest spatial frequency peaks had stronger surround weighting compared to the centers (higher

K_s/K_c ratio) coupled with surrounds that were almost spatially coexistent with the centers. These

two cells are inconsistent with previous data finding the center responses of midget RGCs to be

some 10x-1000x stronger than those of the surround and finding centers to be much smaller in

spatial extent than surrounds [9]. The cells with higher spatial frequency peaks had center and

surround strength ratios closer to previous data, though all 15 cells had higher surround/center

integrated sensitivity ratios than previous results such as those reported by Croner & Kaplan [9].

These differences, however, are explainable by the blurring effect of the optics as presented in

Figures 9,10,11. Further study of low-frequency L-M and M-L cone opponent cells will confirm if

they exhibit any other physiological differences or if they belong to one or more of several midget

subtypes as others have postulated to exist [48, 103]. There is also evidence that parasol ganglion

cells can respond weakly to L vs. M modulation [106], but they are not expected to have single

cone centers from anatomy [6, 20, 27, 107], so it is unlikely that any of the 15 cells studied here

are parasols.

Modeling foveal RGCs. In our modeling, we could not reject the hypothesis that the 15 cells

studied from the center of the fovea in M3 had single cone centers with some residual optical

blur, beyond diffraction, present in the measurements. We think it likely that there was some

residual optical blur, and certainly cannot reject this idea. However, we note that model scenarios

that did not restrict to a single-cone center also describe the data, and that if no residual blur is

assumed then the single-cone center model can be rejected for some cells. There are two reasons

it is difficult to differentiate between single and multi-cone scenarios from our data. First, due to

limitations imposed by the pixel density of the raster scan, our measurements did not extend to

high spatial frequencies (>50 c/deg) where the different scenarios would make more clearly

different predictions. Second, there was inter-session variability in the measured STFs (Fig. 5),

which decreases the power of the data. In future experiments, we plan to (i) extend the range of

spatial frequencies examined and (ii) test multiple grating orientations. Despite these limitations,

as noted in the Supplement (**S7 Fig**), the multi-cone RF scenario with residual defocus often

produces a RF strikingly similar to the single-cone RF scenario with residual defocus, with only a

minimal amount of cone coupling, and the classical understanding of midget RGCs that postulates

single-cone centers is consistent with our data.

A notable observation of the STFs in our population of foveal RGCs (Fig. 8) is the large amount of

low-spatial frequency attenuation. Such observations are not typical of midget ganglion cells

reported in other *in vivo* studies [9]. The question then arises as to whether these most central

foveal cells have distinctively different RF organization from the less-foveal cells reported in other

studies, or whether the underlying RF organization is similar, but the measured data differ

between the two types of studies simply because of differences in the optical systems involved.

When the animal's optics are added back in (Figures 9,10,11), the STFs have a clear shift of peak

sensitivity towards lower spatial frequencies, approaching those of previous *in vivo* studies but

not exactly replicating them, suggesting that the optics account for most of the observed

differences with the possibility of modest foveal specialization or instrument differences

accounting for the rest.

The modeling used in this study has the capability to add a subject's optical PSF (here through the

measured wavefront aberrations from M3) to predict responses to stimuli presented through the

natural (non-AO-corrected) optics. The robust nature of the entire simulation allows for any cone

density profile or PSF to be substituted in the model so that the simulation can be generalized to

population data or be tailored to individual subjects, as it has been here for M3.

Acknowledgments

The authors would like to thank Amber Walker for performing animal anesthesia and monitoring

during imaging, Deniz Dalkara for providing the viral vector used in M1, Max Snodderly for

assistance with estimating the effect of macular pigment, Charles Granger for optical alignment

assistance, Daniel Guarino and Martin Gira for maintaining the stereotaxic cart and other

electronic components, as well as Aby Joseph, Jennifer Hunter, and Jesse Schallek for data analysis

suggestions. We thank the vector core at the Perelman School of Medicine, University of

Pennsylvania and the Genetically-Encoded Neuronal Indicator and Effector (GENIE) Project and

the Janelia Research Campus of the Howard Hughes Medical Institute, specifically Vivek

Jayaraman, Ph.D., Douglas S. Kim, Ph.D., Loren L. Looger, Ph.D., and Karel Svoboda, Ph.D.

1002

References

1.   Schein SJ. Anatomy of macaque fovea and spatial densities of neurons in foveal
representation. J Comp Neurol. 1988;269(4):479-505. Epub 1988/03/22. doi:
10.1002/cne.902690403. PubMed PMID: 3372725.
2.   Wassle H, Grunert U, Rohrenbeck J, Boycott BB. Cortical magnification factor and the
ganglion cell density of the primate retina. Nature. 1989;341(6243):643-6. Epub 1989/10/19.
doi: 10.1038/341643a0. PubMed PMID: 2797190.

3.   Wassle H, Grunert U, Rohrenbeck J, Boycott BB. Retinal ganglion cell density and cortical
magnification factor in the primate. Vision Res. 1990;30(11):1897-911. Epub 1990/01/01.
PubMed PMID: 2288097.
4.   De Monasterio FM, Gouras P. Functional properties of ganglion cells of the rhesus
monkey retina. J Physiol. 1975;251(1):167-95. Epub 1975/09/01. doi:
10.1113/jphysiol.1975.sp011086. PubMed PMID: 810576; PubMed Central PMCID:
PMCPMC1348381.
5.   Derrington AM, Lennie P. Spatial and temporal contrast sensitivities of neurones in
lateral geniculate nucleus of macaque. J Physiol. 1984;357:219-40. Epub 1984/12/01. doi:
10.1113/jphysiol.1984.sp015498. PubMed PMID: 6512690; PubMed Central PMCID:
PMCPMC1193256.
6.   Watanabe M, Rodieck RW. Parasol and midget ganglion cells of the primate retina. J
Comp Neurol. 1989;289(3):434-54. Epub 1989/11/15. doi: 10.1002/cne.902890308. PubMed
PMID: 2808778.
7.   Kolb H, Dekorver L. Midget ganglion cells of the parafovea of the human retina: a study
by electron microscopy and serial section reconstructions. J Comp Neurol. 1991;303(4):617-36.
Epub 1991/01/22. doi: 10.1002/cne.903030408. PubMed PMID: 1707423.
8.   Hendrickson A. A morphological comparison of foveal development in man and monkey.
Eye (Lond). 1992;6 (Pt 2):136-44. Epub 1992/01/01. doi: 10.1038/eye.1992.29. PubMed PMID:
1624035.
9.   Croner LJ, Kaplan E. Receptive fields of P and M ganglion cells across the primate retina.
Vision Res. 1995;35(1):7-24. Epub 1995/01/01. doi: 10.1016/0042-6989(94)e0066-t. PubMed
PMID: 7839612.
10.   McMahon MJ, Lankheet MJ, Lennie P, Williams DR. Fine structure of parvocellular
receptive fields in the primate fovea revealed by laser interferometry. J Neurosci.
2000;20(5):2043-53. Epub 2000/02/24. PubMed PMID: 10684905.
11.   Hendrickson A. Organization of the Adult Primate Fovea. In: Penfold PL, Provis JM,
editors. Macular Degeneration: Springer Berlin Heidelberg; 2005.
12.   Yin L, Masella B, Dalkara D, Zhang J, Flannery JG, Schaffer DV, et al. Imaging light
responses of foveal ganglion cells in the living macaque eye. J Neurosci. 2014;34(19):6596-605.
Epub 2014/05/09. doi: 10.1523/jneurosci.4438-13.2014. PubMed PMID: 24806684; PubMed
Central PMCID: PMCPMC4012315.
13.   Sinha R, Hoon M, Baudin J, Okawa H, Wong ROL, Rieke F. Cellular and Circuit
Mechanisms Shaping the Perceptual Properties of the Primate Fovea. Cell. 2017;168(3):413-
26.e12. Epub 2017/01/28. doi: 10.1016/j.cell.2017.01.005. PubMed PMID: 28129540; PubMed
Central PMCID: PMCPMC5298833.
14.   Wilk MA, Dubis AM, Cooper RF, Summerfelt P, Dubra A, Carroll J. Assessing the spatial
relationship between fixation and foveal specializations. Vision Research. 2017;132:53-61. doi:
https://doi.org/10.1016/j.visres.2016.05.001.
15.   Bringmann A, Syrbe S, Gorner K, Kacza J, Francke M, Wiedemann P, et al. The primate
fovea: Structure, function and development. Prog Retin Eye Res. 2018;66:49-84. Epub
2018/04/03. doi: 10.1016/j.preteyeres.2018.03.006. PubMed PMID: 29609042.
16.   McGregor JE, Yin L, Yang Q, Godat T, Huynh KT, Zhang J, et al. Functional architecture of
the foveola revealed in the living primate. PLoS One. 2018;13(11):e0207102. Epub 2018/11/30.
doi: 10.1371/journal.pone.0207102. PubMed PMID: 30485298; PubMed Central PMCID:
PMCPMC6261564 following competing interests: DRW and WHM receive funding from the
National Eye Institute. DRW receives additional funding from The Arnold and Mabel Beckman

Foundation and WHM receives funding from Research to Prevent Blindness as detailed in the
Financial Disclosure statement. DRW has patents with the University of Rochester for adaptive
optics imaging of the retina: US patent #6,199,986 "Rapid, automatic measurement of the eye's
wave aberration". US patent #6,264,328 "Wavefront sensor with off-axis illumination" and US
patent 6,338,559 "Apparatus and method for improving vision and retinal imaging". QY has
patents with the University of Rochester, Canon Inc. and the University of Montana, for image
stabilization algorithms: US patent #9,226,656: "Real-time optical and digital image stabilization
for adaptive optics scanning ophthalmoscopy", US patent # 9,406,133: "System and method for
real-time image registration", US patent #: 9,485,383, "Imaging based correction of distortion
from a scanner" and US patent #: 9,454,084, "Light source modulation for a scanning
microscope". Additionally QY has undertaken consultancy work for Oculus VR and Boston
Micromachine Corporation. This does not alter our adherence to PLOS ONE policies on sharing
data and materials.

17.   Thoreson WB, Dacey DM. Diverse Cell Types, Circuits, and Mechanisms for Color Vision
in the Vertebrate Retina. Physiol Rev. 2019;99(3):1527-73. Epub 2019/05/30. doi:
10.1152/physrev.00027.2018. PubMed PMID: 31140374; PubMed Central PMCID:
PMCPMC6689740.
18.   Dacey DM. The mosaic of midget ganglion cells in the human retina. J Neurosci.
1993;13(12):5334-55. Epub 1993/12/01. PubMed PMID: 8254378.
19.   Dacey DM. Primate retina: cell types, circuits and color opponency. Prog Retin Eye Res.
1999;18(6):737-63. Epub 1999/10/26. doi: 10.1016/s1350-9462(98)00013-5. PubMed PMID:
10530750.
20.   Dacey DM. Origins of Perception: Retinal Ganglion Cell Diversity and the Creation of
Parallel Visual Pathways. In: Gazzaniga MS, editor. The Cognitive Neurosciences: The MIT Press;
2004. p. 281-302.
21.   Hansen T, Pracejus L, Gegenfurtner KR. Color perception in the intermediate periphery
of the visual field. J Vis. 2009;9(4):26.1-12. Epub 2009/09/18. doi: 10.1167/9.4.26. PubMed
PMID: 19757935.
22.   Leventhal AG, Rodieck RW, Dreher B. Retinal ganglion cell classes in the Old World
monkey: morphology and central projections. Science. 1981;213(4512):1139-42. Epub
1981/09/04. doi: 10.1126/science.7268423. PubMed PMID: 7268423.
23.   Perry VH, Cowey A. Retinal ganglion cells that project to the superior colliculus and
pretectum in the macaque monkey. Neuroscience. 1984;12(4):1125-37. Epub 1984/08/01. doi:
10.1016/0306-4522(84)90007-1. PubMed PMID: 6483194.
24.   Perry VH, Oehler R, Cowey A. Retinal ganglion cells that project to the dorsal lateral
geniculate nucleus in the macaque monkey. Neuroscience. 1984;12(4):1101-23. Epub
1984/08/01. doi: 10.1016/0306-4522(84)90006-x. PubMed PMID: 6483193.
25.   Kaplan E, Shapley RM. The primate retina contains two types of ganglion cells, with high
and low contrast sensitivity. Proceedings of the National Academy of Sciences of the United
States of America. 1986;83(8):2755-7. Epub 1986/04/01. doi: 10.1073/pnas.83.8.2755. PubMed
PMID: 3458235; PubMed Central PMCID: PMCPMC323379.
26.   Perry VH, Silveira LCL. Functional lamination in the ganglion cell layer of the macaque's
retina. Neuroscience. 1988;25(1):217-23. doi: https://doi.org/10.1016/0306-4522(88)90020-6.
27.   Dacey DM, Petersen MR. Dendritic field size and morphology of midget and parasol
ganglion cells of the human retina. Proceedings of the National Academy of Sciences of the
United States of America. 1992;89(20):9666-70. doi: 10.1073/pnas.89.20.9666. PubMed PMID:
1409680.

28.   Grunert U, Greferath U, Boycott BB, Wassle H. Parasol (P alpha) ganglion-cells of the
primate fovea: immunocytochemical staining with antibodies against GABAA-receptors. Vision
Res. 1993;33(1):1-14. Epub 1993/01/01. PubMed PMID: 8383899.
29.   Dacey DM. Morphology of a small-field bistratified ganglion cell type in the macaque
and human retina. Vis Neurosci. 1993;10(6):1081-98. Epub 1993/11/01. doi:
10.1017/s0952523800010191. PubMed PMID: 8257665.
30.   Dacey DM, Lee BB. The 'blue-on' opponent pathway in primate retina originates from a
distinct bistratified ganglion cell type. Nature. 1994;367(6465):731-5. Epub 1994/02/24. doi:
10.1038/367731a0. PubMed PMID: 8107868.
31.   Graham DM, Wong KY. Melanopsin-expressing, Intrinsically Photosensitive Retinal
Ganglion Cells (ipRGCs). In: Kolb H, Fernandez E, Nelson R, editors. Webvision: The Organization
of the Retina and Visual System. Salt Lake City (UT)1995.
32.   Dacey DM. Circuitry for color coding in the primate retina. Proceedings of the National
Academy of Sciences of the United States of America. 1996;93(2):582-8. Epub 1996/01/23. doi:
10.1073/pnas.93.2.582. PubMed PMID: 8570599; PubMed Central PMCID: PMCPMC40095.
33.   Lee BB. Receptive field structure in the primate retina. Vision Research. 1996;36(5):631-
44. doi: https://doi.org/10.1016/0042-6989(95)00167-0.
34.   Calkins DJ, Tsukamoto Y, Sterling P. Microcircuitry and mosaic of a blue-yellow ganglion
cell in the primate retina. J Neurosci. 1998;18(9):3373-85. Epub 1998/05/09. PubMed PMID:
9547245; PubMed Central PMCID: PMCPMC6792640.
35.   Chichilnisky EJ, Baylor DA. Receptive-field microstructure of blue-yellow ganglion cells in
primate retina. Nat Neurosci. 1999;2(10):889-93. Epub 1999/09/24. doi: 10.1038/13189.
PubMed PMID: 10491609.
36.   Telkes I, Distler C, Hoffmann KP. Retinal ganglion cells projecting to the nucleus of the
optic tract and the dorsal terminal nucleus of the accessory optic system in macaque monkeys.
Eur J Neurosci. 2000;12(7):2367-75. Epub 2000/08/18. doi: 10.1046/j.1460-9568.2000.00133.x.
PubMed PMID: 10947815.
37.   Dacey DM, Liao HW, Peterson BB, Robinson FR, Smith VC, Pokorny J, et al. Melanopsin-
expressing ganglion cells in primate retina signal colour and irradiance and project to the LGN.
Nature. 2005;433(7027):749-54. Epub 2005/02/18. doi: 10.1038/nature03387. PubMed PMID:
15716953.
38.   Calkins DJ, Sterling P. Microcircuitry for two types of achromatic ganglion cell in primate
fovea. J Neurosci. 2007;27(10):2646-53. Epub 2007/03/09. doi: 10.1523/JNEUROSCI.4739-
06.2007. PubMed PMID: 17344402; PubMed Central PMCID: PMCPMC6672494.
39.   Petrusca D, Grivich MI, Sher A, Field GD, Gauthier JL, Greschner M, et al. Identification
and characterization of a Y-like primate retinal ganglion cell type. J Neurosci.
2007;27(41):11019-27. Epub 2007/10/12. doi: 10.1523/jneurosci.2836-07.2007. PubMed PMID:
17928443.
40.   Field GD, Sher A, Gauthier JL, Greschner M, Shlens J, Litke AM, et al. Spatial Properties
and Functional Organization of Small Bistratified Ganglion Cells in Primate Retina. The Journal of
Neuroscience. 2007;27(48):13261-72. doi: 10.1523/jneurosci.3437-07.2007.
41.   Crook JD, Peterson BB, Packer OS, Robinson FR, Gamlin PD, Troy JB, et al. The smooth
monostratified ganglion cell: evidence for spatial diversity in the Y-cell pathway to the lateral
geniculate nucleus and superior colliculus in the macaque monkey. J Neurosci.
2008;28(48):12654-71. Epub 2008/11/28. doi: 10.1523/jneurosci.2986-08.2008. PubMed PMID:
19036959; PubMed Central PMCID: PMCPMC2778055.

42.   Percival KA, Martin PR, Grunert U. Organisation of koniocellular-projecting ganglion cells
and diffuse bipolar cells in the primate fovea. Eur J Neurosci. 2013;37(7):1072-89. Epub
2013/01/15. doi: 10.1111/ejn.12117. PubMed PMID: 23311464.
43.   Crook JD, Packer OS, Troy JB, Dacey DM. Synaptic Mechanisms of Color and Luminance
Coding: Rediscovering the X-Y-cell dichotomy in primate retinal ganglion cels. . In: Werner JS,
Chalupa LM, editors. The New Visual Neurosciences: The MIT Press; 2013. p. 123-44.
44.   Puller C, Manookin MB, Neitz J, Rieke F, Neitz M. Broad thorny ganglion cells: a
candidate for visual pursuit error signaling in the primate retina. J Neurosci. 2015;35(13):5397-
408. Epub 2015/04/04. doi: 10.1523/jneurosci.4369-14.2015. PubMed PMID: 25834063;
PubMed Central PMCID: PMCPMC4381007.
45.   Liao HW, Ren X, Peterson BB, Marshak DW, Yau KW, Gamlin PD, et al. Melanopsin-
expressing ganglion cells on macaque and human retinas form two morphologically distinct
populations. J Comp Neurol. 2016;524(14):2845-72. Epub 2016/03/15. doi: 10.1002/cne.23995.
PubMed PMID: 26972791; PubMed Central PMCID: PMCPMC4970949.
46.   Liu Z, Kurokawa K, Zhang F, Lee JJ, Miller DT. Imaging and quantifying ganglion cells and
other transparent neurons in the living human retina. Proceedings of the National Academy of
Sciences of the United States of America. 2017;114(48):12803-8. Epub 2017/11/16. doi:
10.1073/pnas.1711734114. PubMed PMID: 29138314; PubMed Central PMCID:
PMCPMC5715765.
47.   Peng YR, Shekhar K, Yan W, Herrmann D, Sappington A, Bryman GS, et al. Molecular
Classification and Comparative Taxonomics of Foveal and Peripheral Cells in Primate Retina. Cell.
2019;176(5):1222-37 e22. Epub 2019/02/05. doi: 10.1016/j.cell.2019.01.004. PubMed PMID:
30712875; PubMed Central PMCID: PMCPMC6424338.
48.   Wool LE, Packer OS, Zaidi Q, Dacey DM. Connectomic Identification and Three-
Dimensional Color Tuning of S-OFF Midget Ganglion Cells in the Primate Retina. J Neurosci.
2019;39(40):7893-909. Epub 2019/08/14. doi: 10.1523/JNEUROSCI.0778-19.2019. PubMed
PMID: 31405926; PubMed Central PMCID: PMCPMC6774400.
49.   Rhoades CE, Shah NP, Manookin MB, Brackbill N, Kling A, Goetz G, et al. Unusual
Physiological Properties of Smooth Monostratified Ganglion Cell Types in Primate Retina.
Neuron. 2019;103(4):658-72 e6. Epub 2019/06/23. doi: 10.1016/j.neuron.2019.05.036. PubMed
PMID: 31227309; PubMed Central PMCID: PMCPMC6817368.
50.   Dacey DM, Kim YJ, Packer OS, Detwiler PB. ON-OFF direction selective ganglion cells in
macaque monkey retina are tracer-coupled to an ON-OFF direction selective amacrine cell type.
Investigative Ophthalmology & Visual Science. 2019;60(9):5280-.
51.   Kim US, Mahroo OA, Mollon JD, Yu-Wai-Man P. Retinal Ganglion Cells—Diversity of Cell
Types and Clinical Relevance. Frontiers in Neurology. 2021;12(635). doi:
10.3389/fneur.2021.661938.
52.   Grünert U, Martin PR. Cell types and cell circuits in human and non-human primate
retina. Progress in Retinal and Eye Research. 2020;78:100844. doi:
https://doi.org/10.1016/j.preteyeres.2020.100844.
53.   Lee BB, Martin PR, Valberg A. Sensitivity of macaque retinal ganglion cells to chromatic
and luminance flicker. The Journal of physiology. 1989;414:223-43. doi:
10.1113/jphysiol.1989.sp017685. PubMed PMID: 2607430.
54.   Lee BB, Pokorny J, Smith VC, Martin PR, Valberg A. Luminance and chromatic
modulation sensitivity of macaque ganglion cells and human observers. J Opt Soc Am A.
1990;7(12):2223-36. Epub 1990/12/01. doi: 10.1364/josaa.7.002223. PubMed PMID: 2090801.

55.   Soto F, Hsiang J-C, Rajagopal R, Piggott K, Harocopos GJ, Couch SM, et al. Efficient
Coding by Midget and Parasol Ganglion Cells in the Human Retina. Neuron. 2020;107(4):656-
66.e5. doi: https://doi.org/10.1016/j.neuron.2020.05.030.
56.   Brackbill N, Rhoades C, Kling A, Shah NP, Sher A, Litke AM, et al. Reconstruction of
natural images from responses of primate retinal ganglion cells. eLife. 2020;9:e58516. doi:
10.7554/eLife.58516. PubMed PMID: 33146609.
57.   Calkins DJ, Schein SJ, Tsukamoto Y, Sterling P. M and L cones in macaque fovea connect
to midget ganglion cells by different numbers of excitatory synapses. Nature.
1994;371(6492):70-2. doi: 10.1038/371070a0.
58.   Crook JD, Dacey DM. Circuitry and receptive field structure underlying 'double duty'
performance by midget ganglion cells. Journal of Vision. 2010;10(15):5-. doi: 10.1167/10.15.5.
59.   Kaas JH, Guillery RW, Allman JM. Some Principles of Organization in the Dorsal Lateral
Geniculate Nucleus; pp. 253–267. Brain, Behavior and Evolution. 1972;6(1-6):253-67. doi:
10.1159/000123713.
60.   Derrington AM, Krauskopf J, Lennie P. Chromatic mechanisms in lateral geniculate
nucleus of macaque. J Physiol. 1984;357:241-65. Epub 1984/12/01. doi:
10.1113/jphysiol.1984.sp015499. PubMed PMID: 6512691; PubMed Central PMCID:
PMCPMC1193257.
61.   Solomon SG, Peirce JW, Dhruv NT, Lennie P. Profound contrast adaptation early in the
visual pathway. Neuron. 2004;42(1):155-62. Epub 2004/04/07. doi: 10.1016/s0896-
6273(04)00178-3. PubMed PMID: 15066272.
62.   Levitt JB, Schumer RA, Sherman SM, Spear PD, Movshon JA. Visual response properties
of neurons in the LGN of normally reared and visually deprived macaque monkeys. J
Neurophysiol. 2001;85(5):2111-29. Epub 2001/05/16. doi: 10.1152/jn.2001.85.5.2111. PubMed
PMID: 11353027.
63.   Gray DC, Merigan W, Wolfing JI, Gee BP, Porter J, Dubra A, et al. In vivo fluorescence
imaging of primate retinal ganglion cells and retinal pigment epithelial cells. Optics express.
2006;14(16):7144-58. Epub 2006/08/07. doi: 10.1364/oe.14.007144. PubMed PMID: 19529085.
64.   Yin L, Geng Y, Osakada F, Sharma R, Cetin AH, Callaway EM, et al. Imaging light
responses of retinal ganglion cells in the living mouse eye. J Neurophysiol. 2013;109(9):2415-21.
Epub 2013/02/15. doi: 10.1152/jn.01043.2012. PubMed PMID: 23407356; PubMed Central
PMCID: PMCPMC3652215.
65.   Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, et al. Ultrasensitive
fluorescent proteins for imaging neuronal activity. Nature. 2013;499(7458):295-300. doi:
10.1038/nature12354. PubMed PMID: 23868258.
66.   Yin L, Greenberg K, Hunter JJ, Dalkara D, Kolstad KD, Masella BD, et al. Intravitreal
injection of AAV2 transduces macaque inner retina. Invest Ophthalmol Vis Sci. 2011;52(5):2775-
83. Epub 2011/02/12. doi: 10.1167/iovs.10-6250. PubMed PMID: 21310920; PubMed Central
PMCID: PMCPMC3088562.
67.   Chaffiol A, Caplette R, Jaillard C, Brazhnikova E, Desrosiers M, Dubus E, et al. A New
Promoter Allows Optogenetic Vision Restoration with Enhanced Sensitivity in Macaque Retina.
Molecular therapy : the journal of the American Society of Gene Therapy. 2017;25(11):2546-60.
Epub 2017/08/16. doi: 10.1016/j.ymthe.2017.07.011. PubMed PMID: 28807567; PubMed
Central PMCID: PMCPMC5675708.
68.   McGregor JE, Godat T, Dhakal K, Parkins K, Strazzeri J, Bateman B, et al. Optogenetic
restoration of retinal ganglion cell activity in the living primate. Nature Communications. 2020.

69.   Delori FC, Webb RH, Sliney DH. Maximum permissible exposures for ocular safety (ANSI
2000), with emphasis on ophthalmic devices. J Opt Soc Am A. 2007;24(5):1250-65. doi:
10.1364/JOSAA.24.001250.
70.   Schulmeister K. The upcoming new editions of IEC 60825-1 and ANSI Z136.1 – Examples
on impact for classification and exposure limits. International Laser Safety Conference.
2013;2013(1):330-7. doi: 10.2351/1.5056809.
71.   Rockwell B, Thomas R, Zimmerman S. Updates to the ANSI Z136.1 standard.
International Laser Safety Conference. 2015;2015(1):75-7. doi: 10.2351/1.5056858.
72.   Geng Y, Dubra A, Yin L, Merigan WH, Sharma R, Libby RT, et al. Adaptive optics retinal
imaging in the living mouse eye. Biomedical optics express. 2012;3(4):715-34. Epub 2012/05/11.
doi: 10.1364/boe.3.000715. PubMed PMID: 22574260; PubMed Central PMCID:
PMCPMC3345801.
73.   Atchison DA, Smith G. Chromatic dispersions of the ocular media of human eyes. J Opt
Soc Am A. 2005;22(1):29-37. doi: 10.1364/JOSAA.22.000029.
74.   Nelder JA, Mead R. A Simplex Method for Function Minimization. The Computer Journal.
1965;7(4):308-13. doi: 10.1093/comjnl/7.4.308.
75.   Rossi EA, Granger CE, Sharma R, Yang Q, Saito K, Schwarz C, et al. Imaging individual
neurons in the retinal ganglion cell layer of the living eye. Proceedings of the National Academy
of Sciences of the United States of America. 2017;114(3):586-91. Epub 2017/01/05. doi:
10.1073/pnas.1613445114. PubMed PMID: 28049835; PubMed Central PMCID:
PMCPMC5255596.
76.   Baron WS. Maxwellian view stimulator for electrophysiological or psychophysical work.
Applied optics. 1973;12(11):2560-2. Epub 1973/11/01. doi: 10.1364/AO.12.002560. PubMed
PMID: 20125830.
77.   Schnapf JL, Kraft TW, Nunn BJ, Baylor DA. Spectral sensitivity of primate photoreceptors.
Vis Neurosci. 1988;1(3):255-61. Epub 1988/01/01. PubMed PMID: 3154798.
78.   Snodderly DM, Auran JD, Delori FC. The macular pigment. II. Spatial distribution in
primate retinas. Invest Ophthalmol Vis Sci. 1984;25(6):674-85. Epub 1984/06/01. PubMed PMID:
6724837.
79.   Snodderly DM, Brown PK, Delori FC, Auran JD. The macular pigment. I. Absorbance
spectra, localization, and discrimination from other yellow pigments in primate retinas. Invest
Ophthalmol Vis Sci. 1984;25(6):660-73. Epub 1984/06/01. PubMed PMID: 6724836.
80.   Lamb TD. Photoreceptor spectral sensitivities: Common shape in the long-wavelength
region. Vision Research. 1995;35(22):3083-91. doi: https://doi.org/10.1016/0042-
6989(95)00114-F.
81.   Govardovskii VI, Fyhrquist N, Reuter TOM, Kuzmin DG, Donner K. In search of the visual
pigment template. Visual Neuroscience. 2000;17(4):509-28. Epub 2000/07/01. doi:
10.1017/S0952523800174036.
82.   Bowmaker JK. Cone Visual Pigments in Monkeys and Humans. Advances in
Photoreception: Proceedings of a Symposium on Frontiers of Visual Science: National Academies
Press; 1990.
83.   Boettner EA. Spectral Transmission of the Eye. The University of Michigan, Final Report,
Air Force Contract F41(609)-2966. 1967.
84.   Dobkins KR, Thiele A, Albright TD. Comparison of red-green equiluminance points in
humans and macaques: evidence for different L:M cone ratios between species. Journal of the
Optical Society of America A, Optics, image science, and vision. 2000;17(3):545-56. Epub
2000/03/09. PubMed PMID: 10708036.

85.   Estevez O, Spekreijse H. The "silent substitution" method in visual research. Vision Res.
1982;22(6):681-91. Epub 1982/01/01. doi: 10.1016/0042-6989(82)90104-3. PubMed PMID:
7112962.
86.   Brainard DH. The Psychophysics Toolbox. Spat Vis. 1997;10(4):433-6. Epub 1997/01/01.
PubMed PMID: 9176952.
87.   Passaglia CL, Troy JB, Rüttiger L, Lee BB. Orientation sensitivity of ganglion cells in
primate retina. Vision Research. 2002;42(6):683-94. doi: https://doi.org/10.1016/S0042-
6989(01)00312-1.
88.   Yang Q, Zhang J, Nozato K, Saito K, Williams DR, Roorda A, et al. Closed-loop optical
stabilization and digital image registration in adaptive optics scanning light ophthalmoscopy.
Biomedical optics express. 2014;5(9):3174-91. Epub 2014/11/18. doi: 10.1364/boe.5.003174.
PubMed PMID: 25401030; PubMed Central PMCID: PMCPMC4230869.
89.   Ku HH. Notes on the use of propagation of error formulas. Journal of Research of the
National Bureau of Standards. 1966;70(4):263-73.
90.   Enroth-Cugell C, Robson JG. The contrast sensitivity of retinal ganglion cells of the cat. J
Physiol. 1966;187(3):517-52. Epub 1966/12/01. doi: 10.1113/jphysiol.1966.sp008107. PubMed
PMID: 16783910; PubMed Central PMCID: PMCPMC1395960.
91.   Cottaris NP, Jiang H, Ding X, Wandell BA, Brainard DH. A computational-observer model
of spatial contrast sensitivity: Effects of wave-front-based optics, cone-mosaic structure, and
inference engine. Journal of Vision. 2019;19(4):8-. doi: 10.1167/19.4.8.
92.   Cottaris NP, Wandell BA, Rieke F, Brainard DH. A computational observer model of
spatial contrast sensitivity: Effects of photocurrent encoding, fixational eye movements, and
inference engine. Journal of Vision. 2020;20(7):17-. doi: 10.1167/jov.20.7.17.
93.   Stockman A, Sharpe LT, Fach C. The spectral sensitivity of the human short-wavelength
sensitive cones derived from thresholds and color matches. Vision Res. 1999;39(17):2901-27.
Epub 1999/09/24. doi: 10.1016/s0042-6989(98)00225-9. PubMed PMID: 10492818.
94.   Stockman A, Sharpe LT. The spectral sensitivities of the middle- and long-wavelength-
sensitive cones derived from measurements in observers of known genotype. Vision Research.
2000;40(13):1711-37. doi: https://doi.org/10.1016/S0042-6989(00)00021-3.
95.   Pokorny J, Smith VC, Lutze M. Aging of the human lens. Applied optics. 1987;26(8):1437-
40. doi: 10.1364/AO.26.001437.
96.   Dalkara D, Kolstad KD, Caporale N, Visel M, Klimczak RR, Schaffer DV, et al. Inner limiting
membrane barriers to AAV-mediated retinal transduction from the vitreous. Molecular therapy :
the journal of the American Society of Gene Therapy. 2009;17(12):2096-102. Epub 2009/08/13.
doi: 10.1038/mt.2009.181. PubMed PMID: 19672248; PubMed Central PMCID:
PMCPMC2814392.
97.   Helga Kolb EF, Ralph N. Webvision : the organization of the retina and visual system:
[Bethesda, Md.] : National Library of Medicine : [National Center for Biotechnology
Information], 2007.; 2007.
98.   Croner LJ, Purpura K, Kaplan E. Response variability in retinal ganglion cells of primates.
Proceedings of the National Academy of Sciences. 1993;90(17):8128-30. doi:
10.1073/pnas.90.17.8128.
99.   Yan W, Peng Y-R, van Zyl T, Regev A, Shekhar K, Juric D, et al. Cell Atlas of The Human
Fovea and Peripheral Retina. Scientific reports. 2020;10(1):9802. doi: 10.1038/s41598-020-
66092-9.

100.   Packer O, Hendrickson AE, Curcio CA. Photoreceptor topography of the retina in the
adult pigtail macaque (Macaca nemestrina). Journal of Comparative Neurology.
1989;288(1):165-83. doi: https://doi.org/10.1002/cne.902880113.
101.   Provis JM, Hendrickson AE. The Foveal Avascular Region of Developing Human Retina.
Archives of Ophthalmology. 2008;126(4):507-11. doi: 10.1001/archopht.126.4.507.
102.   Bumsted K, Hendrickson A. Distribution and development of short-wavelength cones
differ between Macaca monkey and human fovea. J Comp Neurol. 1999;403(4):502-16. Epub
1999/01/15. PubMed PMID: 9888315.
103.   Patterson SS, Neitz M, Neitz J. Reconciling Color Vision Models With Midget Ganglion
Cell Receptive Fields. Front Neurosci. 2019;13:865. Epub 2019/09/03. doi:
10.3389/fnins.2019.00865. PubMed PMID: 31474825; PubMed Central PMCID:
PMCPMC6707431.
104.   Tailby C, Solomon SG, Lennie P. Functional Asymmetries in Visual Pathways Carrying S-
Cone Signals in Macaque. The Journal of Neuroscience. 2008;28(15):4078-87. doi:
10.1523/jneurosci.5338-07.2008.
105.   van der Schaaf A, van Hateren JH. Modelling the Power Spectra of Natural Images:
Statistics and Information. Vision Research. 1996;36(17):2759-70. doi:
https://doi.org/10.1016/0042-6989(96)00002-8.
106.   Lee BB, Sun H. The chromatic input to cells of the magnocellular pathway of primates.
Journal of vision. 2009;9(2):1-18. doi: 10.1167/9.2.15. PubMed PMID: 19271925.
107.   Diller L, Packer OS, Verweij J, McMahon MJ, Williams DR, Dacey DM. L and M cone
contributions to the midget and parasol ganglion cell receptive fields of macaque monkey
retina. J Neurosci. 2004;24(5):1079-88. Epub 2004/02/06. doi: 10.1523/JNEUROSCI.3828-
03.2004. PubMed PMID: 14762126; PubMed Central PMCID: PMCPMC6793593.

1358

1359

 Supporting Information

S1 Fig. Soma sizes of RGCs for animals M2 and M3 across chromatic functional groups. Observer

1 used the open source software GIMP to segment RGCs in fluorescence images from M2 (large

FOV), M3 (large FOV) and M3 (small FOV). The ellipse tool was used to segment the rough

boundary of individual RGCs. Observer 2 used the open source software ImageJ to segment the

same RGCs in fluorescence images from M2 and M3 using a hand tracing tool to trace the

observable edges of each cell's fluorescence. Under both methods, the area for each cell was

computed in terms of pixels^2 and then converted to μm^2 using the following formula: $\frac{\mu m}{pixel} =$

$291.2 \frac{\mu m}{deg} * \frac{axial\ length}{24.2\ mm} * \frac{FOV\ width}{496\ pixels}$, where 291.2 μm/deg is the human model eye visual angle to

retinal extent conversion, 24.2 mm is the human model eye axial length, 496 pixels is the width

of the imaging PMT used in the AOSLO system, axial length is the animal's axial length in mm, and

FOV width is the width in degrees of the FOV used. The axial length of animal M2 is 17.2 mm, and

the FOV width used was 3.64 deg. The axial length of animal M3 is 16.56 mm, and the FOV widths

used were 3.69 deg (large FOV) and 2.54 deg (small FOV). Cells were compared across functional

groups identified as L-M/M-L chromatic opponent (L-M), S only responding (S), Luminance only

or achromatic (LUM), and mixed L-M/S responses (L-M/S). In **(a)**, the soma areas in μm^2 for the

four functional groups in M2 are listed as the mean and standard deviation of each group across

both observers. In **(b)**, the same comparisons are made for the four functional groups in M3 at

the large FOV. In **(c)**, comparisons for the two functional groups found in M3 at the inner edge of

the foveal slope (small FOV) are made across both observers. In, **(d)**, summary tables show the p-

scores from a Mann-Whitney U test (MATLAB function ranksum(x,y)) comparing the distributions

of soma areas across functional groups as measured by both observers separately. All p-values

were greater than 0.1, except for Observer 2's comparison of the L-M group to the S only group

in M2 which had p = 0.0118. Based on these results, we cannot reject the hypothesis that the

distributions of soma sizes for these functional groups are roughly the same across the two

animals measured at the range of eccentricities at which we imaged cell somas.

1386

S2 Fig. Spatial response of RGCs from M3 in the large FOV condition. Some example responses

of L-M/M-L chromatic opponent RGCs from M3 at the large FOV, where each plot is the response

of a different cell to drifting gratings (6 Hz) of spatial frequencies from 2 – 34 c/deg. These cells

were chosen to showcase different cell responses and are not necessarily special or

representative of the population. Purple and Orange curves show two separate experiments that

were averaged together (Gray curve). The Blue curves are simple difference of Gaussians fits (not

the full ISETBio modeling, for reasons explained below) to each average. Each plot title includes

the cell's unique label "cN", and five parameters—the center strength K_c, the center radius r_c, the

surround strength K_s, the surround radius r_s, and a goodness of fit value for the difference of

Gaussians. As can be easily seen, many of the cells do not exhibit high spatial frequency falloff at

the range of spatial frequencies measured, and so the difference of Gaussians either fits poorly

or produces fit parameters that are highly irregular or suspect such as center sizes much smaller

than the size of a single cone. For this reason, these data were not incorporated into the modelling

in the main body, as higher spatial frequency responses were needed to properly fit many of these

cell responses. There were 48 such L-M/M-L chromatic opponent cells in the large FOV in M3 out

of 145 measured cells. In M2, there was only one experiment measuring response to spatial

frequency and it suffered from the same lack of high spatial frequency measurements as the data

shown here. Full data from both animals can be shared upon request.

1405

S3 Fig. Spatial response of achromatic RGCs from M3 at the center of the foveola. Each graph

represents a single cell identified as responding only to achromatic stimuli from the center of the

foveola of M3. The measured responses were to drifting gratings (6 Hz) of spatial frequencies

from 4 – 49 c/deg. Purple, Orange, and Green lines represent the cell's response in three different

experiments. The Gray line is the average of the three experiments for each cell, and the Blue line

is a simple difference of Gaussians fit to the average (not the full ISETBio model, see S2 Fig). Each

plot title includes the cell's unique label "cN", and five parameters—the center strength K_c, the

center radius r_c, the surround strength K_s, the surround radius r_s, and a goodness of fit value for

the difference of Gaussians. Compared to the 15 L-M and M-L cone opponent cells from this same

location, these 7 achromatic cells were had noisier responses across experiments and more varied

response characteristics. Though most of these cells are likely to be parasol RGCs, some could be

achromatic midget RGCs or other rarer achromatic RGC types. Due to this uncertainty, these data

were not included in the modelling for putative midget RGCs presented in the main manuscript.

There were also achromatic cells identified in M2 and in M3 at the large FOV condition, but those

data suffer from decreased resolution and a decreased range of spatial frequencies presented to

the cells. Full data from all animals can be shared upon request.

1422

S4 Fig. Cone mosaic modelling fits compared to data from M3. Images showing the performance

of the ISETBio cone mosaic generator compared to the actual data from M3. At top, the model

trichromatic photoreceptor mosaic generated by ISETBio (left) and the corresponding model cone

density (right) (compare to Fig. 1b). At bottom left, a comparison of the median diameter of cones

are compared between the model and the M3 data. At bottom middle and bottom right, the cone

diameters of the model and measured data from M3 are compared across the horizontal and

vertical meridians respectively.

1430

S5 Fig. Model cross-validation. RMS errors for the 4 modeling scenarios we considered are depicted for 12 cells. Yellow and blue bars indicate insample and out-of-sample performance, respectively. During in-sample performance assessment the model is trained and evaluated using data from the same recording session. During out-of-sample performance assessment, the model is trained in one session and evaluated using data from another session. The data do not have enough power to reveal a model with best generalizing (out-of-sample) performance. In a few cells, the 1-cone/0.00D residual defocus model can be ruled out as its performance is significantly worse than the remaining 3. A two sample t-test with unequal variance was used to test against the hypothesis that there is a significant difference in the mean RMS fit errors between 2 modeling scenarios.

S6 Fig. Optimization of residual defocus for different RGCs. For each cell the optimal residual defocus value for the single cone center and multi-cone center model scenarios was calculated to gauge variability from the chosen 0.067 D reported in the main text. For each cell, the RMSE is shown for various residual defocus values for both model scarios. Cells are labeled L1-11 or M1-4 according to whether we believed they were likely to contain an L cone or M cone at their center.

S7 Fig. All four model scenario fits for the 15 L-M and M-L cone opponent RGCs from the center of the fovea of M3. Four model scenarios were considered for all 15 cells: single cone centers with 0 D residual defocus, single cone centers with 0.067 D residual defocus, multi-cone centers with 0 D residual defocus, and multi-cone centers with 0.067 D residual defocus. As can be seen from the fits, the single cone center with 0.067 D defocus performs well for all 15 cells and produces center-surround structures that are consistent with what has been measured from

1454   physiology. Cells are labeled L1-11 or M1-4 according to whether we believed they were likely to

1455   contain an L cone or M cone at their center.

1456

S8 Fig. Simple center-surround model estimating percentage of achromatic midget RGCs. A

simple model cone mosaic using hexagonal packing was created in MATLAB for the centermost

2400 cones (radius of 25 cones from the foveal center). Each cone was randomly assigned to be

an L cone (48%), M cone (48%), or S cone (4%). For this simple model, a midget RGC was assumed

to connect to one cone at its receptive field center and 6 cones at its surround (immediately

adjacent to the center cone). For each model midget RGC, the proportion of cones that were the

same type as the center cone was calculated. RGCs with all 6 surround cones of the same type as

the center were classified as 'true achromatic' (orange histogram), and RGCs with at least 5

surround cones of the same type as the center were classified as 'thresholded achromatic' (blue

histogram). The entire model simulation was then repeated 10,000 times, with a different

randomly assigned cone mosaic each time. The histograms above show the number of simulations

for which each percentage of midget RGCs that were either true achromatic or thresholded

achromatic occurred. Based on this simple model, truly achromatic midget RGCs might make up

approximately 1.5% of total midget RGCs at the foveal center, while thresholded achromatic

midget RGCs might make up approximately 9% of total midget RGCs at the foveal center.

Figure 1

Spatial frequency (cyc/deg)

Click here to access/download
Supporting Information
S1_Fig.pdf

Supporting Information 2

Click here to access/download
Supporting Information
S2_Fig.pdf

Click here to access/download
Supporting Information
S3_Fig.pdf

Click here to access/download
Supporting Information
S4_Fig.pdf

Click here to access/download
Supporting Information
S5_Fig.pdf

Supporting Information 6

Click here to access/download
Supporting Information
S6_Fig.pdf

Click here to access/download
Supporting Information
S7_Fig.pdf

Click here to access/download
Supporting Information
S8_Fig.pdf