1

Drastic differential CRISPR-Cas9 induced mutagenesis influenced by DNA methylation and chromatin features

Trevor Weiss[1,3,5,6], Peter A. Crisp[1,2,3], Krishan M. Rai[1], Meredith Song[4], Nathan M. Springer[1,3], Feng Zhang[1,3,5,6]*.

[1]Department of Plant and Microbial Biology, University of Minnesota, Saint Paul, MN 55108
[2]School of Agriculture and Food Sciences, The University of Queensland, Brisbane QLD 4072, Australia
[3]Center for Precision Plant Genomics, University of Minnesota, Saint Paul, MN 55108
[4]Department of Genetics, Cell Biology and Development, University of Minnesota, Minneapolis, MN 55108
[5]Microbial and Plant Genomics Institute, University of Minnesota, Minneapolis, MN, 55108
[6]Center for Genome Engineering, University of Minnesota, Minneapolis, MN, 55108

*Corresponding author: Feng Zhang
Room 318, Cargill Building
1500 Gortner Ave
University of Minnesota
St. Paul, 55108
Email: zhangumn@umn.edu

Short running title: Epigenetic features impact CRISPR-Cas9 efficacy

Abstract

CRISPR-Cas9 mediated genome editing has been widely adopted for basic and

applied biological research in eukaryotic systems. While many studies

considered DNA sequences of CRISPR target sites as the primary determinant

for CRISPR mutagenesis efficiency and mutation profiles, increasing evidence

revealed the significant role of chromatin context. Nonetheless, most of the prior

studies were limited by the lack of sufficient epigenetic resources and/or by only

transiently expressing CRISPR-Cas9 in a short time window. In this study, we

leveraged the wealth of high-resolution epigenomic resources in *Arabidopsis*

thaliana to address the impact of chromatin features on CRISPR-Cas9

mutagenesis using stable transgenic plants. Our results indicated that DNA

methylation and chromatin features could lead to significant variations in

mutagenesis efficiency by up to 250 folds. Low mutagenesis efficiencies were

mostly associated with repressive heterochromatic features. This repressive

effect appeared to persist through cell divisions but could be alleviated through

substantial reduction of DNA methylation at CRISPR target sites. Moreover,

specific chromatin features, such as H3K4me1, H3.3, and H3.1, appear to be

associated with significant variations in CRISPR-Cas9 mutation profiles reflected

by the 1 bp insertion rates. Our findings provided strong evidence that specific

chromatin features could have significant and lasting impacts on both CRISPR-

Cas9 mutagenesis efficiency and DNA double strand break repair outcomes.

Introduction

CRISPR-Cas based genome editing technologies have greatly advanced

both basic and applied biological research. Among them, CRISPR-Cas9, the

class II bacterial CRISPR system, has been the first and the most widely adopted

in eukaryotes (Jinek et al. 2012). The key steps in CRISPR-Cas9 mediated

genome editing involve searching, binding and then cleaving a 20-nucleotide

target site directed by a guide RNA (gRNA). The resulting cleavage product with

double-strand breaks (DSBs) can then be repaired by either error-prone DNA

repair pathways, such as classical non-homologous end-joining (NHEJ) or

microhomology-mediated end-joining (MMEJ), or by a template DNA-dependent

pathway, *i.e.* homology directed repair (HDR) (Chen et al. 2019). Thus, specific

mutations, including insertions, deletions or point mutations, can be introduced

by employing distinct DNA repair machineries (Chen et al. 2019).

Previous studies indicated that the CRISPR targeted sequence is the primary

determinant for mutagenesis efficiency and mutation profile (Allen et al. 2018;

Lazzarotto et al. 2020). Several tools have been developed to predict the

efficiency and mutation outcomes solely based on CRISPR targeted sequences

(Allen et al. 2018; Concordet and Haeussler 2018; Xiang et al. 2021). However,

the predictability of these tools, primarily based on the large dataset from human

cells, often varies and appears to translate poorly to other species, such as

plants (Naim et al. 2020). This observation suggested that non-sequence

features could influence CRISPR-Cas9 mutagenesis.

Increasing evidence revealed negative correlations between CRISPR-

Cas9 mutagenesis rates and heterochromatic signatures or low chromatin

accessibility in multiple systems, such as yeast, zebrafish, mouse, human and

rice (Daer et al. 2017; G. Liu et al. 2019; Uusi-Mäkelä et al. 2018; Wu et al. 2014;

Yarrington et al. 2018). However, most of these studies were conducted at

various genomic locations, making it difficult to separate the effect of chromatin

context from those of DNA sequences. Recently two studies investigated the

impact of chromatin features by using over 1,000 copies of integrated reporter

sequences to fix the sequence variables (Gisler et al. 2019; Schep et al. 2021).

Their findings confirmed the previous observations that heterochromatin has a

negative impact on CRISPR-Cas9 mutagenesis efficacy. Notably, specific

chromatin features were also identified to impact both efficiency and mutation

outcomes (Gisler et al. 2019; Schep et al. 2021). Nevertheless, these studies

were limited by two factors: 1) the bias of the integration sites in certain genomic

locations, and 2) the ambiguity from whether the newly integrated sequences can

quickly and faithfully adopt the local chromatin context. Furthermore, most of

these previous studies were conducted in cell lines within a short time window

(usually less than 72 hours) using transiently expressed CRISPR-Cas9. It is still

unclear whether the heterochromatic features have a long-lasting effect on

CRISPR-Cas9 mutagenesis, or only delay it (Kallimasioti-Pazi et al. 2018).

In this study, we leveraged the high resolution epigenomic resources in

the model plant species, *Arabidopsis thaliana*, to address the impact of chromatin

features on both CRISPR-Cas9 mutagenesis efficiency and mutation outcomes.

To fix the sequence variable, the Arabidopsis genome was scanned to identify

identical CRISPR target sites located in various chromatin contexts. By using

stable CRISPR-Cas9 transgenic plants targeting multiple chromosomal regions,

we systematically characterized mutagenesis efficiency and mutation outcomes

with 23 distinct DNA methylation and chromatin features using a Next Generation

Sequencing (NGS) approach. Our findings provided insight into the influences of

chromatin features on CRISPR-Cas9 mutagenesis and DNA repair outcomes.

This could help develop better technologies for more efficient and precise

genome editing.

Results

Identification of identical CRISPR-Cas9 sites in diverse chromatin contexts

To identify identical CRISPR target sites in various chromatin contexts, the

Arabidopsis *Col*-0 reference genome was scanned for 20 bp CRISPR-Cas9

recognition sequences with 3 bp NGG (the PAM sequence) at the 3' end. Out of

7,376,476 distinct target sites, 19,161 were identified as repeating 7-25 times

across the genome (Figure 1A). A series of filters were then applied to remove

the sites with one of the following features: simple repeats, GC content outside

the range of 40-60%, matching sequences in mitochondria or chloroplast

genomes, or containing no overlapping restriction enzyme site for subsequent

mutation genotyping. The remaining 7,971 sequences, representing 92,117 total

genomic sites, were assessed for three key chromatin features using 100 bp

windows: chromatin accessibility indicated by ATAC-seq scores (Lu et al. 2016),

DNA methylation patterns categorized as DNA methylation domain (RdDM,

heterochromatin, CG-only, unmethylated and intermediate) (Crisp et al. 2017;

Springer and Schmitz 2017), and nine chromatin states (Supplemental Dataset

1) (Sequeira-Mendes et al. 2014). Seven multicopy CRISPR sites (MCsites) were

identified with individual sequences in each family having highly diverse

chromatin contexts, including both open and closed chromatin, at least 3 different

DNA methylation domains, and at least 2 distinct chromatin states (Figure 1B-C;

Supplemental Dataset 2).

136
137 **Differential CRISPR-Cas9 efficiencies were associated with distinct**
138 **chromatin features**
139
Next, we evaluated CRISPR-Cas9 efficacy for each of the seven MCsite

families. T-DNA constructs, containing a CRISPR-Cas9 expression cassette, a

firefly luciferase reporter and a BAR selection marker gene, were made to target

each MCsite (Supplemental Figure S1A). In each construct, the CRISPR single

guide RNA (sgRNA) cassette consisted of 2 guide RNAs (gRNAs), one targeting

the MCsite and the other targeting a single-copy endogenous gene, Cheletase2

(CHLl2), as the CRISPR mutagenesis control (Figure 2A) (Mao et al. 2013). It is

worth noting that we intentionally chose the CaMV 35S promoter to drive

expression of the Cas9 and gRNA sequences because this promoter has much

lower activity in Arabidopsis embryos than leaves (Wang et al. 2015; Yan et al.

2015). By reducing the mutagenesis potential in early embryo development

stages, we were able to capture more independent mutation events in somatic

cells during leaf development.

After transformation of each T-DNA construct, the resulting CRISPR-Cas9

transgenic plants (T1) were first assessed for mutagenesis efficiency at the

CHLI2 control site using the Cleaved Amplified Polymorphic Sequences (CAPS)

method. Individual plants with detectable mutagenesis at the control site were

then analyzed at each target site. Using the CAPS assay we were able to identify

two MCsites, MCsite4 and MCsite5, that produced notable mutagenesis for at

least one of the CRISPR target sites (Supplemental Figure S1B). MCsite4 and 5,

totaling 15 individual sites, were found spanning all 5 chromosomes

(Supplemental Figure S1C). Among the 15 sites, four seemed to overlap with

Arabidopsis genes, and only 1 site, MCsite5.8, was detected in transcribed RNA

sequences (Supplemental Dataset 2).

CRISPR-Cas9 mutagenesis efficiency was then assessed at the individual

sequences of MCsite4 and 5 by quantifying indel mutations using the NGS

assay. Consistent with the observation from the CAPS assay, substantial

variations were observed across individual target sites (Supplemental Figure

S2A). To control for the variation in CRISPR-Cas9 expression levels between

plants, the mutation rate at each targeted site was normalized to the CHLI2

control. As a result, the normalized frequencies at MCsite4 and MCsite5 sites

ranged from an average 0.61-152.28% and 9.17%-69.17%, respectively (Figure

2B-C). Pairwise comparisons indicated mutagenesis frequencies of MCsite4 sites

can be categorized into three distinct groups (p value < 0.001): the high editing

group (group H: MCsite4.8, 152.28%), the moderate editing frequency group

(group M: MCsite4.4 with 17.04% and 4.7 with 16.20%), and the low editing

frequency group (group L: MCsite4.1, 4.2, 4.3, 4.5, and 4.9, ranging from 0.61%

to 1.71%) (Figure 2B). Comparison between the highest and lowest MCsite4

edited sites revealed a 249.64-fold difference. Similarly, mutagenesis frequencies

at the MCsite5 sequences can also be grouped into the high, moderate and low

editing groups (*p* value < 0.01) with 7.54-fold differences between the highest

and lowest edited sites (Figure 2C). We then characterized the local sequence

context surrounding each target site (25 bp from each side). High sequence

similarities were found in these extended sequences for both MCsite4 and 5

families (Supplemental Figure S2B and C). Phylogenetic analyses within each

target site family found no evident correlations between sequence similarity and

the mutagenesis frequency groups (Supplemental Figure S2B and C). Thus, the

local sequence context could not explain differential mutagenesis frequencies

observed from individual target sites.

The initial association analysis of indel frequencies with DNA methylation

domains and chromatin accessibility indicated unmethylated and accessible sites

generally had higher mutagenesis levels than the methylated and inaccessible

sites (Figure 2B-C). Notably, the strongly negative correlations between

mutagenesis frequency and DNA methylation levels at both MCsite4 and 5 sites

could also be observed when the cytosine methylation status was examined at

the single nucleotide level of individual CRISPR target sites (Supplemental

Figure S3). To systematically investigate the correlation between mutagenesis

efficiency and chromatin features, we further characterized individual MCsites

using all 23 chromatin features, including distinct histone modifications and

histone variants (Supplemental Dataset 3) (Y. Liu et al. 2018). When hierarchical

cluster analysis was performed, all lowly edited sites (group L) clustered with the

heterochromatic features, such as H2A.W, H3K9me1, H3K9me2, H3K27me1,

and hyper DNA methylation, for both MCsite4 and 5 (Figure 2D-E). On the

contrary, the highly and moderately edited groups (group H and M) appeared to

be associated with accessible, active chromatin features such as histone

acetylation, H3K36me3, and H3K4 methylation (Figure 2D-E) (Roudier et al.

2011). To examine the impact of individual features, we performed correlation

analyses by plotting mutagenesis efficiency at all 15 target sites with each

chromatin and DNA methylation feature (Supplemental Figure S4). Consistent

with the hierarchical cluster analysis, strong positive correlations were observed

between mutagenesis frequency and euchromatin-related features such as

H3K56ac ($R = 0.85$, $p = 1.8e\text{-}13$), H3K9ac ($R = 0.82$, $p = 8.6e\text{-}12$), H3K36ac ($R =$

0.75, $p = 4.2e\text{-}9$), H3K27ac ($R = 0.71$, $p = 4.1e\text{-}8$), H3K36me3 ($R = 0.71$, $p =$

$3.5e\text{-}8$), H3K4me3 ($R = 0.65$, $p = 1.6\text{--}6$), and accessibility ($R = 0.51$, $p =$

0.00038) measured with ATAC-Seq data (Supplemental Figure S4). On the

contrary, strong negative correlations were found between mutagenesis

frequencies and the heterochromatin-related features H2A.W ($R = \text{-}0.57$, $p =$

$4.1e\text{-}5$), H3K9me1 ($R = \text{-}0.55$, $p = 0.00011$), H3K9me2 ($R = \text{-}0.55$, $p = 1e\text{-}4$) and

cytosine DNA methylation ($R = \text{-}0.45$, $p = 0.0022$) (Supplemental Figure S4).

219
220 **Improving CRISPR-Cas9 mutagenesis efficiency through combined**
221 **reduction of DNA methylation**
222
Because of the strong negative association observed between the lowly

edited sites and heterochromatic features, we hypothesized that altering these

chromatin states could improve mutagenesis efficiency at the refractory sites. In

this study, we chose to perturb DNA methylation at the lowly edited sites

because high levels of DNA methylation are often correlated with

heterochromatic features (Crisp et al. 2020; Johnson et al. 2007; Zemach and

Grafi 2003). We first sought to test the possible impact of CHG methylation on

mutagenesis efficiencies because CHG methylation is often associated with

H3K9me2 and heterochromatin (Springer and Schmitz 2017). To this end, the

chromomethylase3 (*cmt3-11t*) mutant was chosen due to the well-documented

genome-wide reduction in CHG methylation (Stroud et al. 2013). Analysis of the

single-base resolution DNA methylation profiles indicated that a substantial

reduction in CHG methylation was confirmed at individual MCsite4 and 5

CRISPR targets in this mutant (Supplemental Figure S5). The CRISPR-Cas9 T-

DNA constructs targeting MCsite4 and 5 sequences were transformed into the

cmt3-11t mutant using the same procedure described above. As a result,

transgenic *cmt3-11t* plants (T1) were identified with detectable mutagenesis

activities at both MCsite4 and 5 target sites (Supplemental Figure S6). After

normalizing to the CHLI2 control target site, we observed a nearly identical

pattern in mutagenesis efficiency between WT and *cmt3-11t* plants for both

MCsite4 and MCsite5 (Figure 3A-B). These results suggested that a reduction of

CHG methylation alone is not sufficient to improve CRISPR-Cas9 mutagenesis at

the refractory sites.

Although additional genetic mutant lines with perturbed DNA methylation,

such as *met1*, were available (Stroud et al. 2013), we have not been able to

obtain transgenic CRISPR-Cas9 T-DNA events in these mutant lines. Therefore,

we chose a chemical approach to reduce genome-wide DNA methylation in all

5mC contexts using 5-azacytidine (Griffin, Niederhuth, and Schmitz 2016).

Analysis of the DNA methylation for 5-azacytidine treated plants revealed

substantial reductions in DNA methylation at the highly methylated sequences at

both MCsite4 and 5 (Supplemental Figure S7A) (Griffin, Niederhuth, and Schmitz

2016). We focused on MCsite4 in the subsequent 5-azacytidine experiment due

to the lower mutagenesis frequencies at the lowly edited sites. The T2 plants with

the CRISPR-Cas9 T-DNA targeting MCsite4 in the wildtype background were

obtained from a self-pollinated T1 plant used above. After 2-weeks of growth with

or without 100 µM 5-azacytidine, the plants were subjected to mutagenesis

analyses using the NGS assay (Supplemental Figure S7B). No significant

differences were observed for the normalized mutagenesis frequency at each

MCsite4 site between the untreated and treated samples (Figure 3C). Thus,

partial reductions in all DNA methylation contexts using 5-azacytidine did not

reveal changes in mutagenesis frequencies.

Lastly, we tested the impact of combined reductions in DNA methylation

using both the *cmt3-11t* mutant and 5-azacytidine treatments. T2 plants with the

MCsite4-targeting T-DNA were obtained from a self-pollinated T1 plant in the

cmt3-11t mutant background, grown for 2 weeks with or without 100 µM 5-

azacytidine, and subjected to NGS analysis using the same procedure

(Supplemental Figure S7C). After normalizing to the CHLI2 control, we observed

two distinct patterns for mutagenesis efficiency across individual MCsite4

CRISPR targets. At the unmethylated and accessible sites (groups H and M), no

significant differences in mutagenesis frequency were found between the 5-

azacytidine treated mutants and the control group (Figure 3D). On the contrary,

significant increases in mutagenesis frequency were observed at the

inaccessible and hypermethylated sites (group L) when 5-azacytidine treatment

was combined with the *cmt3-11t* mutant, *i.e.,* MCsite4.9 (4.01-fold), MCsite4.3

(2.12-fold), MCsite4.1 (4.73-fold), MCsite4.5 (4.83-fold), and MCsite4.2 (3.87-

fold) (Figure 3D). Together, these data indicated that strong reduction in DNA

methylation in multiple contexts could result in significant improvement of

CRISPR-Cas9 mutagenesis efficiency at the hypermethylated refractory sites.

281
282   **Differential CRISPR-Cas9 mutational profiles are associated with distinct**
283   **chromatin features**
284
CRISPR-Cas9 induced mutations are typically composed of either small

deletions or 1 bp insertions (Allen et al. 2018). Small deletions are mainly derived

from the NHEJ or MMEJ pathway through exonuclease-mediated end processing

and ligation, while the 1 bp insertions were mostly generated from blunt-end or 1

bp staggered cleavage by Cas9 followed by DNA polymerase-mediated end

filling (Gisler et al. 2019). To investigate the potential impact of chromatin context

on CRISPR-Cas9 mutation outcome, we examined the insertion and deletion

profiles for both MCsite4 and 5. As expected, most mutations contained 1 bp

insertions and small deletions (< 10 bp) for both MCsite4 and 5 (Figure 4A).

MCsite4 was preferentially repaired as 1 bp insertions, while MCsite5 showed a

strong bias towards small deletion outcomes (Figure 4A). While the major

mutation types at individual sites within each family seemed to be highly similar

(Supplemental Figure S8A-D), further analysis revealed significant variations for

the insertion rate between individual sites. The rate of insertion outcomes ranged

from 56.25-81.73% and 7.57-30.27% in MCsite4 and MCsite5 sites, respectively

(Figure 4B-C). We then conducted a correlation analysis using the 23 epigenetic

features with the insertion rates at all 15 sites. Three histone H3 related features,

H3K4me1 (R = -0.64, p = 0.01), H3.3 (R = -0.83, p = 3e-4), and H3.1 (R = -0.91,

p = 2.8e-6), were identified with significant negative correlations with 1 bp

insertional mutations (Figure. 4D; Supplemental Figure S9). Thus, these findings

strongly suggested that chromatin features could not only affect CRISPR-Cas9

mutagenesis efficiency but also influence mutation outcomes.

Discussion

In recent studies, chromatin contexts have been demonstrated to have

significant impacts on CRISPR-Cas9 mediated genome editing (Daer et al. 2017;

G. Liu et al. 2019; Wu et al. 2014; Yarrington et al. 2018). Most of these findings

indicated that heterochromatic features at the CRISPR target regions could

impede CRISPR-Cas9 mutagenesis efficiency in multiple systems, such as

yeast, rice, mouse, and human cell lines (Daer et al. 2017; G. Liu et al. 2019; Wu

et al. 2014; Yarrington et al. 2018). However, it remained unclear whether

heterochromatic features could only temporarily delay CRISPR-Cas9

mutagenesis (Kallimasioti-Pazi et al. 2018). In this study, we systematically

characterized the impact of 23 distinct DNA methylation and chromatin features

on CRISPR-Cas9 mutagenesis efficiency and mutation outcomes by

investigating CRISPR-Cas9 transgenic Arabidopsis plants. Consistent with the

previous studies, our results demonstrated that inaccessible and heterochromatic

features were strongly associated with low mutagenesis efficiency. Such

repressive effects can be long-lasting through plant development leading up to a

250-fold difference between identical CRISPR target sites. On the other hand,

mutagenesis efficiency at the target sequences in accessible chromatin regions

could also have significant variations ranging from 1.53-fold to 10-fold at MCsite5

and MCsite4, respectively (Figure 2B-C). This observation suggested that

specific chromatin features other than just open or closed chromatin should be

considered to account for CRISPR-Cas9 mutagenesis efficacy. Close

examination of individual chromatin features identified several euchromatic

marks, such as H3K9ac, H3K56ac, H3K36ac, H3K27ac, H3K4 methylation and

H3K36m3, that were positively correlated with mutagenesis efficiency. Further

investigation with a larger data set will be needed to dissect their impacts on

CRISPR-Cas9 mutagenesis in greater detail.

In this study, we observed strongly negative correlations between

CRISPR-Cas9 mutagenesis efficiency and repressive chromatin features, such

as hyper DNA methylation, low DNA accessibility, H3K9 methylations. To test the

hypothesis that modulating some of these features could improve mutagenesis

frequency at the lowly edited sites, we sought to reduce DNA methylation by

using both genetic and chemical approaches. Our results indicated that partial

reduction of DNA methylation using either a mutant affecting CHG methylation or

5-azacytidine treatment alone was not sufficient to improve mutagenesis

efficiency at any of the tested sites. When combining the CHG deficiency mutant

with 5-azacytidine chemical treatment, 2.1 to 4.8-fold improvements in

mutagenesis efficiency were found at the lowly edited sites but not at the highly

and moderately edited sites. Combined reduction of DNA methylation in multiple

contexts has been demonstrated to increase chromatin accessibility and even

alter the higher order 3D chromatin organization in Arabidopsis (Zhong et al.

2021). Thus, mutagenesis efficiency improvement at lowly edited sites observed

here could have resulted from increasing chromatin accessibility and/or changing

3D chromatin organization due to the significant reduction of DNA methylation at

multiple contexts.

In addition to the impacts on mutagenesis efficiency, chromatin features

have been suggested to influence CRISPR-Cas9 mutation outcomes. For

example, a recent study with more than 1,000 copies of identical insertion sites

indicated the 1 bp insertions were found more prevalent in euchromatin than in

heterochromatin, likely through recruiting different DNA repair machinery (Schep

et al. 2021). However, conflicting results were also reported showing little impact

of chromatin features on mutation outcomes (Gisler et al. 2019; Kallimasioti-Pazi

et al. 2018). In this study, we observed significant variations for the 1 bp insertion

rate in different chromatin contexts. Of the 23 chromatin features analyzed, three

distinct histone H3 features, H3K4me1, H3.3, and H3.1, exhibited significantly

strong negative correlations with the 1 bp insertion rate. Interestingly, H3K4me1

was also identified by Schep et al. as a marker to correlate with distinct mutation

outcomes, while H3.1 and H3.3 were not included in their study (Schep et al.

2021). This suggested that the balance between 1 bp insertions and small

deletions could be influenced by specific chromatin markers. Distinct chromatin

features have been reported to recruit different DNA repair machinery and result

in different repair outcomes in mammalian systems (Fnu et al. 2011; Jacquet et

al. 2016; Luijsterburg et al. 2016). Further investigation is needed to address the

potential roles of specific chromatin markers in determining DNA repair

outcomes.

We proposed a model to account for the impacts of chromatin features on

CRISPR-Cas9 mutagenesis efficiency and mutation outcomes (Figure 5). In the

first step, chromatin features are the key determinants for CRISPR-Cas9

recognition and binding efficiency (Figure 5). In general, heterochromatic

features, such as H3K9 methylations, H3K27me1, H2A.W, and hyper DNA

methylation, could significantly reduce chromatin accessibility and thus reduce

the recognition and binding efficiency of CRISPR-Cas9. Such repressive effects

could persist through cell division and development. After CRISPR-Cas9 locates

the target site and binds the genomic DNA, the Cas9 nuclease can introduce

double stranded DNA breaks either with a 1-bp 5' overhang (staggered cut), or

blunt ends (Gisler et al. 2019). The cleaved product can be repaired to yield three

outcomes: wild-type sequence by perfect ligation, 1 bp insertions, or small

deletions (less than 10 bp) (Figure 5). It has been proposed that the staggered

cut primarily leads to 1 bp insertion via template-dependent repair, facilitated by a

DNA polymerase, while the blunt cut mainly results in small deletions through

end resection (Schmid-Burgk et al. 2020; Gisler et al. 2019). The blunt ends

could occasionally be repaired by DNA polymerase, likely the members from

DNA polymerase family X, without DSB end resection, resulting in template-

independent 1 bp insertions (Gisler et al. 2019). In this study, we observed both

templated 1 bp insertion and template-independent insertions, as exemplified in

MCsite5 and MCsite4, respectively (Supplemental Figure S8A-B). Nevertheless,

the balance between 1 bp insertion and small deletion products is primarily

dependent on the repair choices between short-range DSB end resection and

DNA polymerase end filling (Figure 5) (Schmid-Burgk et al. 2020; Lemos et al.

2018). While sequence features are a key determinant for mutation profiles, our

data indicated that chromatin features, such as H3K4me1, H3.3, and H3.1, could

also significantly impact the balance between 1 bp insertion and small deletion

outcomes. These specific chromatin features may exert their influences on the

balance between the staggered and blunt cut, or through modulating the balance

between DNA polymerase end filling and short-range DSB end resection during

NHEJ. In fact, previous studies have demonstrated that distinct chromatin

features, such as the H3.3 variant, could impact DSB end resection (Luijsterburg

et al. 2016).

Recently much effort has been made to develop computational tools to

predict CRISPR-Cas9 editing efficiency and/or repair outcomes. To our

knowledge, these tools primarily relied on sequence features (Allen et al. 2018;

Concordet and Haeussler 2018; Xiang et al. 2021). Yet our results indicated that

non-sequence features should also be taken into consideration for the prediction

of both CRISPR-Cas9 efficiency and mutation outcomes. One implication from

this study is that heterochromatin related features, such as low accessibility,

H3K9me1, H3K9me2, H3K27me1, H2A.W and hyper DNA methylation, should

be avoided in order to design gRNAs with high efficiency. When it comes to

predicting CRISPR-Cas9 mutation profiles, chromatin features such as

H3K4me1, H3.3, and H3.1 should also be considered. Future studies using

genetic mutants involved in histone medications will be interesting to further

dissect the impact of distinct chromatin features on CRISPR-Cas9 mutagenesis.

A better understanding of the interplay between chromatin dynamics and

CRISPR-Cas9 will enable the development of more precise and efficient genome

engineering technologies.

Methods and Materials

Identification of multicopy CRISPR sites

To identify gRNAs that matched to multiple places in the genome, the

Arabidopsis Col-0 reference genome (TAIR10) was parsed to identify every NGG

PAM site using *seqkit locate* (Shen et al. 2016) with the search motif

NNNNNNNNNNNNNNNNNNNNNGG. The number of occurrences of each non-

redundant sequence in the genome was summarized using *csvtk*. The resulting

distinct target sites were then filtered in R (v4.1.2) to retain gRNAs that had

between 7 to 25 distinct matches. Sites were then eliminated if they had simple

sequence motifs consisting of 5 As, Ts, Gs or Cs in a row; CG content outside

40%-60% or if the sequence was found in the chloroplast or mitochondria

genome; to give a final list of 9,902 candidate gRNAs. The gRNA identification

script is available at

https://github.com/pedrocrisp/Weiss_et_al_gRNA_chromatin. The potential

gRNAs were also screened to identify those with restriction enzyme recognition

motifs (from a list commercially available enzymes from NEB) that overlapped

position 17-18 of the gRNA (between position 3 and 4 bp from the PAM) such

that an indel mutation in this position would disrupt restriction enzyme recognition

and cleave for efficient screening of edited transgenic lines. A final list of 7,971

gRNAs were then annotated with chromatin state information by overlapping the

coordinates of the gRNA target sites with chromatin annotation files using

bedtools (Quinlan and Hall 2010). Chromatin data included nine histone states

(Sequeira-Mendes et al. 2014); chromatin accessibility (Lu et al. 2016); DNA

methylation (Crisp et al. 2017; Springer and Schmitz 2017). DNA methylation

data was converted to methylation domains for each 100 bp non-overlapping

window of the TAIR10 genome using the method detailed in (Crisp et al. 2020).

Chromatin accessibility at the CRISPR target site was called either Open or

Closed based on the presence of an overlapping ATAC-Seq peak or lack thereof,

respectively, using the accessibility profiles in (Lu et al. 2016). Gene and

transposable element annotations were downloaded from Araport v11.

T-DNA vector construction

The CRISPR-Cas9 constructs were created using the Golden Gate assembly

method. The gRNA sequences were first assembled into the pMOD_B2301

backbone containing the gRNA targeting the MCsite and the CHLl2 positive

control (Čermák et al. 2017). T-DNA constructs were assembled using

pMOD_A0101 (Cas9), pMOD_B2301 containing the gRNA array, pMZ105

(luciferase reporter), and pTRANS230d (T-DNA backbone) (Čermák et al. 2017).

All of the T-DNA constructs described in this study are available at Addgene. The

modular components used to build the T-DNA plasmids can be found at

https://www.addgene.org/browse/article/28189956/ (Čermák et al. 2017).

Plant materials and growth conditions

The *Arabidopsis thaliana* Columbia ecotype (Col-0) was used in these

experiments. The *cmt3-11t* (stock CS16392) genotype was acquired from the

Arabidopsis Biological Resource Center (ABRC). Floral dip transformation was

performed according to the protocol as previously outlined (Zhang et al. 2006).

Transgenic T1 seeds were sown on soil and exposed to BASTA selection to

recover transgenic plants. Plants were grown in a growth chamber with the

following conditions: 16/8 hours light/dark cycle, 22 °C, and 55% humidity.

Characterization of chromatin features

Previously analyzed datasets (BigWig files) for the 23 chromatin features were

downloaded from Plant Chromatin State Database (PCSD) (Y. Liu et al. 2018).

The datasets for wild type Arabidopsis with and without 100 µM 5-azacytidine

treatment was downloaded from Griffin et al. (Griffin, Niederhuth, and Schmitz

2016). For each dataset, the values for the 1kb window (500 bp upstream and

500 bp downstream from the center of gRNA) were calculated using the

Deeptools2 computeMatrix in the reference-point mode. Similarly, the values for

each nucleotide of target sites were calculated using the scale-region mode at

single base resolution. Each dataset was then normalized to allow for

comparison on a scale of 0-1 with 1 indicating the highest level of that feature.

5-azacytidine treatment and luciferase screening

T2 seedlings from self-pollinated T1 Arabidopsis plants were grown on 1% agar

plates containing 0.5 Murashige and Skoog (PhytoTech Labs) and 100 µM 5-

azacytidine (Griffin, Niederhuth, and Schmitz 2016). After two weeks, seedlings

were screened for presence of the transgene using a luciferase reporter. The

luciferase assay procedure was performed using the Bio-Glo™ Luciferase Assay

System (Promega Corp., Madison, WI, USA) in accordance with the

manufacturer's instructions.

Mutation genotyping and characterization of mutation profiles

Genotyping was performed using two methods: genomic PCR followed by

restriction enzyme digestion (CAPS) and the NGS assay using Illumina paired-

end read amplicon sequencing. All tissues for genotyping were collected at two

weeks post germination for the CTAB-base genomic DNA extraction. PCR was

performed using GoTaq Green Mastermix (Promega Corp., Madison, WI, USA)

according to the manufacturer's instructions, with an annealing temperature of 55

°C (CHIL2 and MCsite4) or 60 °C (MCsite5) with an extension time of 1 minute.

Primers to amplify CHLl2, MCsite4, and MCsite5 target sites can be found in

Table S1. Amplicons were then subjected to restriction enzyme digestion using

*BsmA*I (CHIL2), *Alu*I (MCsite4), or *Drd*I (MCsite5) according to the

manufacturer's instructions. PCR amplicons generated with the corresponding

primers were subjected to Illumina paired-end read sequencing (Genewiz Inc.,

South Plainfield, NJ, USA). The raw NGS reads were analyzed using

CRISPResso2 to estimate indel mutation rates (Clement et al. 2019). To analyze

the mutation. To analyze the mutation profiles for each sample, the NGS reads

with indel mutations were extracted from the CRISPResso2 output files with a

2% threshold. The resulting output files were then loaded into R studio (version

4.1.0) for data visualization using ggplot (41). The total read counts for CRISPR-

Cas9 editing frequency and repair profiles can be found in Table S2. Normalized

indel frequencies were calculated by dividing the indel frequency of each MCsite

by the CHlL2 positive control indel frequency within each replicate.

Data availability

All sequencing data analyzed in this manuscript will be available at the National

Center for Biotechnology Information (NCBI) under BioProject Accession

PRJNA795172 (Supplemental Dataset 4).

References

Allen, Felicity, Luca Crepaldi, Clara Alsinet, Alexander J. Strong, Vitalii Kleshchevnikov, Pietro De Angeli, Petra Páleníková, et al. 2018. "Predicting the Mutations Generated by Repair of Cas9-Induced Double-Strand Breaks." *Nature Biotechnology* 37 (1): 64–72.

Čermák, Tomáš, Shaun J. Curtin, Javier Gil-Humanes, Radim Čegan, Thomas J. Y. Kono, Eva Konečná, Joseph J. Belanto, et al. 2017. "A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants." *The Plant Cell.* https://doi.org/10.1105/tpc.16.00922.

Chen, Kunling, Yanpeng Wang, Rui Zhang, Huawei Zhang, and Caixia Gao. 2019. "CRISPR/Cas Genome Editing and Precision Plant Breeding in Agriculture." *Annual Review of Plant Biology* 70 (April): 667–97.

Clement, Kendell, Holly Rees, Matthew C. Canver, Jason M. Gehrke, Rick Farouni, Jonathan Y. Hsu, Mitchel A. Cole, et al. 2019. "CRISPResso2 Provides Accurate and Rapid Genome Editing Sequence Analysis." *Nature Biotechnology* 37 (3): 224–26.

Concordet, Jean-Paul, and Maximilian Haeussler. 2018. "CRISPOR: Intuitive Guide Selection for CRISPR/Cas9 Genome Editing Experiments and Screens." *Nucleic Acids Research* 46 (W1): W242–45.

Crisp, Peter A., Diep R. Ganguly, Aaron B. Smith, Kevin D. Murray, Gonzalo M. Estavillo, Iain Searle, Ethan Ford, et al. 2017. "Rapid Recovery Gene Downregulation during Excess-Light Stress and Recovery in Arabidopsis." *The Plant Cell* 29 (8): 1836–63.

Crisp, Peter A., Alexandre P. Marand, Jaclyn M. Noshay, Peng Zhou, Zefu Lu, Robert J. Schmitz, and Nathan M. Springer. 2020. "Stable Unmethylated DNA Demarcates Expressed Genes and Their Cis-Regulatory Space in Plant Genomes." *Proceedings of the National Academy of Sciences of the United States of America* 117 (38): 23991–0.

Daer, René M., Josh P. Cutts, David A. Brafman, and Karmella A. Haynes. 2017. "The Impact of Chromatin Dynamics on Cas9-Mediated Genome Editing in Human Cells." *ACS Synthetic Biology* 6 (3): 428–38.

Fnu, Sheema, Elizabeth A. Williamson, Leyma P. De Haro, Mark Brenneman, Justin Wray, Montaser Shaheen, Krishnan Radhakrishnan, Suk-Hee Lee, Jac A. Nickoloff, and Robert Hromas. 2011. "Methylation of Histone H3 Lysine 36 Enhances DNA Repair by Nonhomologous End-Joining." *Proceedings of the National Academy of Sciences of the United States of America* 108 (2): 540–45.

Gisler, Santiago, Joana P. Gonçalves, Waseem Akhtar, Johann de Jong, Alexey V. Pindyurin, Lodewyk F. A. Wessels, and Maarten van Lohuizen. 2019. "Multiplexed Cas9 Targeting Reveals Genomic Location Effects and gRNA-Based Staggered Breaks Influencing Mutation Efficiency." *Nature Communications* 10 (1): 1598.

Griffin, Patrick T., Chad E. Niederhuth, and Robert J. Schmitz. 2016. "A Comparative Analysis of 5-Azacytidine- and Zebularine-Induced DNA Demethylation." *G3* 6 (9): 2773–80.

Jacquet, Karine, Amélie Fradet-Turcotte, Nikita Avvakumov, Jean-Philippe Lambert, Céline Roques, Raj K. Pandita, Eric Paquet, et al. 2016. "The TIP60 Complex

Regulates Bivalent Chromatin Recognition by 53BP1 through Direct H4K20me
Binding and H2AK15 Acetylation." *Molecular Cell* 62 (3): 409–21.
Jinek, Martin, Krzysztof Chylinski, Ines Fonfara, Michael Hauer, Jennifer A. Doudna,
and Emmanuelle Charpentier. 2012. "A Programmable Dual-RNA-Guided DNA
Endonuclease in Adaptive Bacterial Immunity." *Science* 337 (6096): 816–21.
Johnson, Lianna M., Magnolia Bostick, Xiaoyu Zhang, Edward Kraft, Ian Henderson,
Judy Callis, and Steven E. Jacobsen. 2007. "The SRA Methyl-Cytosine-Binding
Domain Links DNA and Histone Methylation." *Current Biology: CB* 17 (4): 379–
84.
Kallimasioti-Pazi, Eirini M., Keerthi Thelakkad Chathoth, Gillian C. Taylor, Alison
Meynert, Tracy Ballinger, Martijn J. E. Kelder, Sébastien Lalevée, Ildem Sanli,
Robert Feil, and Andrew J. Wood. 2018. "Heterochromatin Delays CRISPR-Cas9
Mutagenesis but Does Not Influence the Outcome of Mutagenic DNA Repair."
PLoS Biology 16 (12): e2005595.
Lazzarotto, Cicera R., Nikolay L. Malinin, Yichao Li, Ruochi Zhang, Yang Yang,
Gahyun Lee, Eleanor Cowley, et al. 2020. "CHANGE-Seq Reveals Genetic and
Epigenetic Effects on CRISPR–Cas9 Genome-Wide Activity." *Nature
Biotechnology* 38 (11): 1317–27.
Lemos, Brenda R., Adam C. Kaplan, Ji Eun Bae, Alexander E. Ferrazzoli, James
Kuo, Ranjith P. Anand, David P. Waterman, and James E. Haber. 2018.
"CRISPR/Cas9 Cleavages in Budding Yeast Reveal Templated Insertions and
Strand-Specific Insertion/deletion Profiles." *Proceedings of the National Academy
of Sciences of the United States of America* 115 (9): E2040–47.
Liu, Guanwen, Kangquan Yin, Qianwei Zhang, Caixia Gao, and Jin-Long Qiu. 2019.
"Modulating Chromatin Accessibility by Transactivation and Targeting Proximal
dsgRNAs Enhances Cas9 Editing Efficiency in Vivo." *Genome Biology* 20 (1):
145.
Liu, Yue, Tian Tian, Kang Zhang, Qi You, Hengyu Yan, Nannan Zhao, Xin Yi,
Wenying Xu, and Zhen Su. 2018. "PCSD: A Plant Chromatin State Database."
Nucleic Acids Research 46 (D1): D1157–67.
Luijsterburg, Martijn S., Inge de Krijger, Wouter W. Wiegant, Rashmi G. Shah,
Godelieve Smeenk, Anton J. L. de Groot, Alex Pines, et al. 2016. "PARP1 Links
CHD2-Mediated Chromatin Expansion and H3.3 Deposition to DNA Repair by
Non-Homologous End-Joining." *Molecular Cell* 61 (4): 547–62.
Lu, Zefu, Brigitte T. Hofmeister, Christopher Vollmers, Rebecca M. DuBois, and
Robert J. Schmitz. 2016. "Combining ATAC-Seq with Nuclei Sorting for
Discovery of Cis-Regulatory Regions in Plant Genomes." *Nucleic Acids
Research* 45 (6): e41–e41.
Mao, Yanfei, Hui Zhang, Nanfei Xu, Botao Zhang, Feng Gou, and Jian-Kang Zhu.
2013. "Application of the CRISPR-Cas System for Efficient Genome Engineering
in Plants." *Molecular Plant* 6 (6): 2008–11.
Naim, Fatima, Kylie Shand, Satomi Hayashi, Martin O'Brien, James McGree,
Alexander A. T. Johnson, Benjamin Dugdale, and Peter M. Waterhouse. 2020.
"Are the Current gRNA Ranking Prediction Algorithms Useful for Genome Editing
in Plants?" *PloS One* 15 (1): e0227994.

600 Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for
Comparing Genomic Features." *Bioinformatics* 26 (6): 841–42.
Roudier, François, Ikhlak Ahmed, Caroline Bérard, Alexis Sarazin, Tristan Mary-
Huard, Sandra Cortijo, Daniel Bouyer, et al. 2011. "Integrative Epigenomic
Mapping Defines Four Main Chromatin States in Arabidopsis." *The EMBO*
Journal 30 (10): 1928–38.
Schep, Ruben, Eva K. Brinkman, Christ Leemans, Xabier Vergara, Robin H. van der
Weide, Ben Morris, Tom van Schaik, et al. 2021. "Impact of Chromatin Context
on Cas9-Induced DNA Double-Strand Break Repair Pathway Balance."
Molecular Cell 81 (10): 2216–30.e10.
Schmid-Burgk, Jonathan L., Linyi Gao, David Li, Zachary Gardner, Jonathan
Strecker, Blake Lash, and Feng Zhang. 2020. "Highly Parallel Profiling of Cas9
Variant Specificity." *Molecular Cell* 78 (4): 794–800.e8.
Sequeira-Mendes, Joana, Irene Aragüez, Ramón Peiró, Raul Mendez-Giraldez,
Xiaoyu Zhang, Steven E. Jacobsen, Ugo Bastolla, and Crisanto Gutierrez. 2014.
"The Functional Topography of the Arabidopsis Genome Is Organized in a
Reduced Number of Linear Motifs of Chromatin States." *The Plant Cell* 26 (6):
2351–66.
Shen, Wei, Shuai Le, Yan Li, and Fuquan Hu. 2016. "SeqKit: A Cross-Platform and
Ultrafast Toolkit for FASTA/Q File Manipulation." *PloS One* 11 (10): e0163962.
Springer, Nathan M., and Robert J. Schmitz. 2017. "Exploiting Induced and Natural
Epigenetic Variation for Crop Improvement." *Nature Reviews. Genetics* 18 (9):
563–75.
Stroud, Hume, Maxim V. C. Greenberg, Suhua Feng, Yana V. Bernatavichute, and
Steven E. Jacobsen. 2013. "Comprehensive Analysis of Silencing Mutants
Reveals Complex Regulation of the Arabidopsis Methylome." *Cell* 152 (1): 352–
64.
Uusi-Mäkelä, Meri I. E., Harlan R. Barker, Carina A. Bäuerlein, Tomi Häkkinen, Matti
Nykter, and Mika Rämet. 2018. "Chromatin Accessibility Is Associated with
CRISPR-Cas9 Efficiency in the Zebrafish (Danio Rerio)." *PloS One* 13 (4):
e0196238.
Wang, Zhi-Ping, Hui-Li Xing, Li Dong, Hai-Yan Zhang, Chun-Yan Han, Xue-Chen
Wang, and Qi-Jun Chen. 2015. "Egg Cell-Specific Promoter-Controlled
CRISPR/Cas9 Efficiently Generates Homozygous Mutants for Multiple Target
Genes in Arabidopsis in a Single Generation." *Genome Biology* 16 (July): 144.
Wu, Xuebing, David A. Scott, Andrea J. Kriz, Anthony C. Chiu, Patrick D. Hsu, Daniel
B. Dadon, Albert W. Cheng, et al. 2014. "Genome-Wide Binding of the CRISPR
Endonuclease Cas9 in Mammalian Cells." *Nature Biotechnology* 32 (7): 670–76.
Xiang, Xi, Giulia I. Corsi, Christian Anthon, Kunli Qu, Xiaoguang Pan, Xue Liang,
Peng Han, et al. 2021. "Enhancing CRISPR-Cas9 gRNA Efficiency Prediction by
Data Integration and Deep Learning." *Nature Communications* 12 (1): 3238.
Yan, Liuhua, Shaowei Wei, Yaorong Wu, Ruolan Hu, Hongju Li, Weicai Yang, and Qi
Xie. 2015. "High-Efficiency Genome Editing in Arabidopsis Using YAO Promoter-
Driven CRISPR/Cas9 System." *Molecular Plant* 8 (12): 1820–23.
Yarrington, Robert M., Surbhi Verma, Shaina Schwartz, Jonathan K. Trautman, and
Dana Carroll. 2018. "Nucleosomes Inhibit Target Cleavage by CRISPR-Cas9 in

Vivo." *Proceedings of the National Academy of Sciences of the United States of*
America 115 (38): 9351–58.
Zemach, Assaf, and Gideon Grafi. 2003. "Characterization of Arabidopsis Thaliana
Methyl-CpG-Binding Domain (MBD) Proteins." *The Plant Journal: For Cell and*
Molecular Biology 34 (5): 565–72.
Zhang, Xiuren, Rossana Henriques, Shih-Shun Lin, Qi-Wen Niu, and Nam-Hai Chua.
2006. "Agrobacterium-Mediated Transformation of Arabidopsis Thaliana Using
the Floral Dip Method." *Nature Protocols* 1 (2): 641–46.
Zhong, Zhenhui, Suhua Feng, Sascha H. Duttke, Magdalena E. Potok, Yiwei Zhang,
Javier Gallego-Bartolomé, Wanlu Liu, and Steven E. Jacobsen. 2021. "DNA
Methylation-Linked Chromatin Accessibility Affects Genomic Architecture in
Arabidopsis." *Proceedings of the National Academy of Sciences of the United*
States of America 118 (5). https://doi.org/10.1073/pnas.2023347118.

Acknowledgements and Funding

We thank all members of the Zhang Lab for helpful discussion. M.S. was

supported by the Undergraduate Research Opportunities Program (UROP) from

the University of Minnesota. Funding from NSF IOS-1934384 to N.M.S. partially

supported this work. P.A.C. was supported by an ARC Discovery Early Career

Award (DE200101748). F.Z. was supported by the startup fund from Department

of Plant and Microbial Biology, University of Minnesota.

Author Contributions

T.W., P.A.C., N.M.S. and F.Z. designed research; T.W., P.A.C., K.M.R., and M.S.

conducted experiments; T.W., K.M.R., and F.Z. analyzed data; T.W., N.M.S. and

F.Z. wrote the paper.

Conflict of Interest

The authors declare no conflict of interest.

686 **Figures and Tables**
687

688
689
Figure 1. Identification of multicopy CRISPR sites (MCsites) in various chromatin
contexts. *(A)* Bioinformatic pipeline used to identify MCsites. *(B)* Characterization
of the chromatin accessibility as Closed (red) or Open (blue) at each MCsite
using ATAC-Seq data. *(C)* Characterization of DNA methylation domains at each
MCsite. CG-only (red) sites contain greater than 40% mCG; Heterochromatin
(tan) sites contain greater than 40% mCG and mCHG methylation; RdDM (light
blue) sites contain mCG, mCHG, and mCHH, with at least 15% mCHH;
Unmethylated (blue) sites contain less than 10% mCG, mCHG, and mCHH;
Intermediate (black) sites are everything else with data that didn't meet any of the
above criteria.
700
701

702
703
Figure 2. Characterization of CRISPR-Cas9 mutation frequencies and chromatin
features at MCsites. *(A)* The NGS-based workflow to analyze CRISPR-Cas9
mutagenesis efficiency and repair profile at MCsites. *(B)* and *(C)* Normalized
mutation frequency for individual MCsite4 and MCsite5 CRISPR targets. The
standard error (SEM) was calculated for each target site with three replicates. A
one-way analysis of variance and Tukey's multiple comparison test were
performed to group the target sites into H (high), M (moderate), and L (low)
mutagenesis levels. The DNA methylation domain and accessibility features
were listed below each target site. Het stands for Heterochromatin DNA
methylation domain. *(D)* and *(E)* Hierarchical clustering heatmap of distinct
chromatin features (x-axis) for individual MCsite4 or MCsite5 sites (y-axis). H
(high), M (moderate), and L (low) indicated the mutagenesis group for each site.
Each feature was normalized to a scale of 0-1, with 1 indicating the highest level
(the darkest color) and 0 indicating the lowest level (white). The dendrogram
indicated similarity clustering of chromatin features.

719
720

721
722
Figure 3. Comparisons of CRISPR-Cas9 mutagenesis frequencies using genetic
and chemical approaches. *(A)* and *(B)* Normalized mutation frequency were
compared for MCsite4 and MCsite5 sites in wild type and *cmt3* mutant
background. *(C)* Normalized mutation frequencies for MCsite4 sites in wild type
plants were compared with or without 5-azacytidine treatment. *(D)* Normalized
mutagenesis frequencies were compared for the MCsite4 sites in the *cmt3*
mutant plants with or without 5-azacytidine treatment. The MCsite4 sites were
categorized in High (H; site 8), Moderate (M; site 7 and 4) and Low (L; sites 9, 3,
1, 5 and 2) mutagenesis groups. The standard error (SEM) was calculated for
each target site with 3 replicates. The fold changes in mutagenesis were
calculated for the sites in group L. Mann-Whitney U test was also performed with
asterisks indicating significant *p*-values ($p < 0.05$).
735

736
737
Figure 4. Analyses of CRISPR-Cas9 mutation profiles at MCsites. *(A)* Distribution of mutation
outcomes at MCsite4 and MCsite5. The mutations were categorized as 1 bp insertions (+1 bp,
red), 1-5 bp deletions (Δ 1-5 bp, dark blue), 6-10 bp deletions (Δ 6-10 bp, blue), and deletions of
more than 10 bp (Δ >10 bp, light blue). *(B)* and *(C)* The 1 bp insertion frequencies at individual
MCsite4 and MCsite5 targets. The standard error (SEM) was calculated for each target site with
replicates (n=6 for MCsite4; n=3 for MCsite5). Kruskal-Wallis analysis indicated the p-value for
MCsite4 and MCsite5 are 0.17 and 0.016, respectively. *(D)* Correlation plots between the
insertion rate and 3 chromatin markers, H3K4me1, H3.3 and H3.1, with the blue dots for MCsite4
and the red dots for MCsite5. The trendline was plotted with gray indicating the standard error.
The R value and *p*-values were calculated using Spearman's rank correlation coefficient.
748
749
750

751
752
Figure 5. Model depicting the factors that influence CRISPR-Cas9 mutagenesis
efficiency and mutation outcomes. The four key check points were indicated in
the numerical order with red circles. At Point 1, both sequence and chromatin
features are the key determinants for CRISPR-Cas9 recognition and binding
efficiency. At Point 2, sequence and chromatin features are crucial to determine
staggered or blunt cleavage by CRISPR-Cas9. The cleaved products can be
ligated back perfectly or go to Point 3 or 4, where chromatin features likely
influence occurrence of end resection. With end resection, it generates small
deletions. When the end resection is limited, the broken ends will be directly filled
by DNA polymerase (DNA Pol) to generate 1 bp insertions. 1 bp insertions can
be the products from template-dependent repair using the 1 bp 5' overhang by
staggered cleavage, or from template-independent repair by filling in blunt ends.
The staggered cleavage primarily leads to 1 bp insertions, while the blunt cut
mainly results in small deletions as indicated by line thickness.

767
768

Figure 1. Identification of multicopy CRISPR sites (MCsites) in various chromatin contexts. *(A)* Bioinformatic pipeline used to identify MCsites. *(B)* Characterization of the chromatin accessibility as Closed (red) or Open (blue) at each MCsite using ATAC-Seq data. *(C)* Characterization of DNA methylation domains at each MCsite. CG-only (red) sites contain greater than 40% mCG; Heterochromatin (tan) sites contain greater than 40% mCG and mCHG methylation; RdDM (light blue) sites contain mCG, mCHG, and mCHH, with at least 15% mCHH; Unmethylated (blue) sites contain less than 10% mCG, mCHG, and mCHH; Intermediate (black) sites are everything else with data that didn't meet any of the above criteria.

Figure 2. Characterization of CRISPR-Cas9 mutation frequencies and chromatin features at MCsites. *(A)* The NGS-based workflow to analyze CRISPR-Cas9 mutagenesis efficiency and repair profile at MCsites. *(B)* and *(C)* Normalized mutation frequency for individual MCsite4 and MCsite5 CRISPR targets. The standard error (SEM) was calculated for each target site with three replicates. A one-way analysis of variance and Tukey's multiple comparison test were performed to group the target sites into H (high), M (moderate), and L (low) mutagenesis levels. The DNA methylation domain and accessibility features were listed below each target site. Het stands for Heterochromatin DNA methylation domain. *(D)* and *(E)* Hierarchical clustering heatmap of distinct chromatin features (x-axis) for individual MCsite4 or MCsite5 sites (y-axis). H (high), M (moderate), and L (low) indicated the mutagenesis group for each site. Each feature was normalized to a scale of 0-1, with 1 indicating the highest level (the darkest color) and 0 indicating the lowest level (white). The dendrogram indicated similarity clustering of chromatin features.

Figure 3. Comparisons of CRISPR-Cas9 mutagenesis frequencies using genetic and chemical approaches. *(A)* and *(B)* Normalized mutation frequency were compared for MCsite4 and MCsite5 sites in wild type and *cmt3* mutant background. *(C)* Normalized mutation frequencies for MCsite4 sites in wild type plants were compared with or without 5-azacytidine treatment. *(D)* Normalized mutagenesis frequencies were compared for the MCsite4 sites in the *cmt3* mutant plants with or without 5-azacytidine treatment. The MCsite4 sites were categorized in High (H; site 8), Moderate (M; site 7 and 4) and Low (L; sites 9, 3, 1, 5 and 2) mutagenesis groups. The standard error (SEM) was calculated for each target site with 3 replicates. The fold changes in mutagenesis were calculated for the sites in group L. Mann-Whitney U test was also performed with asterisks indicating significant *p*-values ($p < 0.05$).

Figure 4. Analyses of CRISPR-Cas9 mutation profiles at MCsites. *(A)* Distribution of mutation outcomes at MCsite4 and MCsite5. The mutations were categorized as 1 bp insertions (+1 bp, red), 1-5 bp deletions (Δ 1-5 bp, dark blue), 6-10 bp deletions (Δ 6-10 bp, blue), and deletions of more than 10 bp (Δ >10 bp, light blue). *(B)* and *(C)* The 1 bp insertion frequencies at individual MCsite4 and MCsite5 targets. The standard error (SEM) was calculated for each target site with replicates (n=6 for MCsite4; n=3 for MCsite5). Kruskal-Wallis analysis indicated the p-value for MCsite4 and MCsite5 are 0.17 and 0.016, respectively. *(D)* Correlation plots between the insertion rate and 3 chromatin markers, H3K4me1, H3.3 and H3.1, with the blue dots for MCsite4 and the red dots for MCsite5. The trendline was plotted with gray indicating the standard error. The R value and *p*-values were calculated using Spearman's rank correlation coefficient.

Figure 5. Model depicting the factors that influence CRISPR-Cas9 mutagenesis efficiency and mutation outcomes. The four key check points were indicated in the numerical order with red circles. In Point 1, both sequence and chromatin features are the key determinants for CRISPR-Cas9 recognition and binding efficiency. In Point 2, sequence and chromatin features are crucial to determine staggered or blunt cleavage by CRISPR-Cas9. The cleaved products can be ligated back perfectly or go to Point 3 or 4, where chromatin features likely influence occurrence of end resection. With end resection, it generates small deletions. When end resection is limited, the broken ends will be directly filled by DNA polymerase (DNA Pol) to generate 1 bp insertions. 1 bp insertions can be the products from template-dependent repair using the 1 bp 5' overhang by staggered cleavage, or from template-independent repair by filling in blunt ends. The staggered cleavage primarily leads to 1 bp insertions, while the blunt cut mainly results in small deletions as indicated by line thickness.

Parsed Citations

Allen, Felicity, Luca Crepaldi, Clara Alsinet, Alexander J. Strong, Vitalii Kleshchevnikov, Pietro De Angeli, Petra Páleníková, et al. 2018. "Predicting the Mutations Generated by Repair of Cas9-Induced Double-Strand Breaks." Nature Biotechnology 37 (1): 64–72.
  Google Scholar: Author Only Title Only Author and Title

Čermák, Tomáš, Shaun J. Curtin, Javier Gil-Humanes, Radim Čegan, Thomas J. Y. Kono, Eva Konečná, Joseph J. Belanto, et al. 2017. "A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants." The Plant Cell. https://doi.org/10.1105/tpc.16.00922.
  Google Scholar: Author Only Title Only Author and Title

Chen, Kunling, Yanpeng Wang, Rui Zhang, Huawei Zhang, and Caixia Gao. 2019. "CRISPR/Cas Genome Editing and Precision Plant Breeding in Agriculture." Annual Review of Plant Biology 70 (April): 667–97.
  Google Scholar: Author Only Title Only Author and Title

Clement, Kendell, Holly Rees, Matthew C. Canver, Jason M. Gehrke, Rick Farouni, Jonathan Y. Hsu, Mitchel A. Cole, et al. 2019. "CRISPResso2 Provides Accurate and Rapid Genome Editing Sequence Analysis." Nature Biotechnology 37 (3): 224–26.
  Google Scholar: Author Only Title Only Author and Title

Concordet, Jean-Paul, and Maximilian Haeussler. 2018. "CRISPOR: Intuitive Guide Selection for CRISPR/Cas9 Genome Editing Experiments and Screens." Nucleic Acids Research 46 (W1): W242–45.
  Google Scholar: Author Only Title Only Author and Title

Crisp, Peter A., Diep R. Ganguly, Aaron B. Smith, Kevin D. Murray, Gonzalo M. Estavillo, Iain Searle, Ethan Ford, et al. 2017. "Rapid Recovery Gene Downregulation during Excess-Light Stress and Recovery in Arabidopsis." The Plant Cell 29 (8): 1836–63.
  Google Scholar: Author Only Title Only Author and Title

Crisp, Peter A., Alexandre P. Marand, Jaclyn M. Noshay, Peng Zhou, Zefu Lu, Robert J. Schmitz, and Nathan M. Springer. 2020. "Stable Unmethylated DNA Demarcates Expressed Genes and Their Cis-Regulatory Space in Plant Genomes." Proceedings of the National Academy of Sciences of the United States of America 117 (38): 23991–0.
  Google Scholar: Author Only Title Only Author and Title

Daer, René M., Josh P. Cutts, David A. Brafman, and Karmella A. Haynes. 2017. "The Impact of Chromatin Dynamics on Cas9-Mediated Genome Editing in Human Cells." ACS Synthetic Biology 6 (3): 428–38.
  Google Scholar: Author Only Title Only Author and Title

Fnu, Sheema, Elizabeth A. Williamson, Leyma P. De Haro, Mark Brenneman, Justin Wray, Montaser Shaheen, Krishnan Radhakrishnan, Suk-Hee Lee, Jac A. Nickoloff, and Robert Hromas. 2011. "Methylation of Histone H3 Lysine 36 Enhances DNA Repair by Nonhomologous End-Joining." Proceedings of the National Academy of Sciences of the United States of America 108 (2): 540–45.
  Google Scholar: Author Only Title Only Author and Title

Gisler, Santiago, Joana P. Gonçalves, Waseem Akhtar, Johann de Jong, Alexey V. Pindyurin, Lodewyk F. A. Wessels, and Maarten van Lohuizen. 2019. "Multiplexed Cas9 Targeting Reveals Genomic Location Effects and gRNA-Based Staggered Breaks Influencing Mutation Efficiency." Nature Communications 10 (1): 1598.
  Google Scholar: Author Only Title Only Author and Title

Griffin, Patrick T., Chad E. Niederhuth, and Robert J. Schmitz. 2016. "A Comparative Analysis of 5-Azacytidine- and Zebularine-Induced DNA Demethylation." G3 6 (9): 2773–80.
  Google Scholar: Author Only Title Only Author and Title

Jacquet, Karine, Amélie Fradet-Turcotte, Nikita Avvakumov, Jean-Philippe Lambert, Céline Roques, Raj K. Pandita, Eric Paquet, et al. 2016. "The TIP60 Complex Regulates Bivalent Chromatin Recognition by 53BP1 through Direct H4K20me Binding and H2AK15 Acetylation." Molecular Cell 62 (3): 409–21.
  Google Scholar: Author Only Title Only Author and Title

Jinek, Martin, Krzysztof Chylinski, Ines Fonfara, Michael Hauer, Jennifer A. Doudna, and Emmanuelle Charpentier. 2012. "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity." Science 337 (6096): 816–21.
  Google Scholar: Author Only Title Only Author and Title

Johnson, Lianna M., Magnolia Bostick, Xiaoyu Zhang, Edward Kraft, Ian Henderson, Judy Callis, and Steven E. Jacobsen. 2007. "The SRA Methyl-Cytosine-Binding Domain Links DNA and Histone Methylation." Current Biology: CB 17 (4): 379–84.
  Google Scholar: Author Only Title Only Author and Title

Kallimasioti-Pazi, Eirini M., Keerthi Thelakkad Chathoth, Gillian C. Taylor, Alison Meynert, Tracy Ballinger, Martijn J. E. Kelder, Sébastien Lalevée, Ildem Sanli, Robert Feil, and Andrew J. Wood. 2018. "Heterochromatin Delays CRISPR-Cas9 Mutagenesis but Does Not Influence the Outcome of Mutagenic DNA Repair." PLoS Biology 16 (12): e2005595.
  Google Scholar: Author Only Title Only Author and Title

Lazzarotto, Cicera R., Nikolay L. Malinin, Yichao Li, Ruochi Zhang, Yang Yang, Gahyun Lee, Eleanor Cowley, et al. 2020. "CHANGE-Seq Reveals Genetic and Epigenetic Effects on CRISPR–Cas9 Genome-Wide Activity." Nature Biotechnology 38 (11): 1317–27.
  Google Scholar: Author Only Title Only Author and Title

Lemos, Brenda R., Adam C. Kaplan, Ji Eun Bae, Alexander E. Ferrazzoli, James Kuo, Ranjith P. Anand, David P. Waterman, and James E. Haber. 2018. "CRISPR/Cas9 Cleavages in Budding Yeast Reveal Templated Insertions and Strand-Specific Insertion/deletion Profiles." Proceedings of the National Academy of Sciences of the United States of America 115 (9): E2040–47.
  Google Scholar: Author Only Title Only Author and Title

Liu, Guanwen, Kangquan Yin, Qianwei Zhang, Caixia Gao, and Jin-Long Qiu. 2019. "Modulating Chromatin Accessibility by Transactivation and Targeting Proximal dsgRNAs Enhances Cas9 Editing Efficiency in Vivo." Genome Biology 20 (1): 145.

Liu, Yue, Tian Tian, Kang Zhang, Qi You, Hengyu Yan, Nannan Zhao, Xin Yi, Wenying Xu, and Zhen Su. 2018. "PCSD: A Plant Chromatin State Database." Nucleic Acids Research 46 (D1): D1157–67.
  Google Scholar: Author Only Title Only Author and Title

Luijsterburg, Martijn S., Inge de Krijger, Wouter W. Wiegant, Rashmi G. Shah, Godelieve Smeenk, Anton J. L. de Groot, Alex Pines, et al. 2016. "PARP1 Links CHD2-Mediated Chromatin Expansion and H3.3 Deposition to DNA Repair by Non-Homologous End-Joining." Molecular Cell 61 (4): 547–62.
  Google Scholar: Author Only Title Only Author and Title

Lu, Zefu, Brigitte T. Hofmeister, Christopher Vollmers, Rebecca M. DuBois, and Robert J. Schmitz. 2016. "Combining ATAC-Seq with Nuclei Sorting for Discovery of Cis-Regulatory Regions in Plant Genomes." Nucleic Acids Research 45 (6): e41–e41.
  Google Scholar: Author Only Title Only Author and Title

Mao, Yanfei, Hui Zhang, Nanfei Xu, Botao Zhang, Feng Gou, and Jian-Kang Zhu. 2013. "Application of the CRISPR-Cas System for Efficient Genome Engineering in Plants." Molecular Plant 6 (6): 2008–11.
  Google Scholar: Author Only Title Only Author and Title

Naim, Fatima, Kylie Shand, Satomi Hayashi, Martin O'Brien, James McGree, Alexander A. T. Johnson, Benjamin Dugdale, and Peter M. Waterhouse. 2020. "Are the Current gRNA Ranking Prediction Algorithms Useful for Genome Editing in Plants?" PloS One 15 (1): e0227994.
  Google Scholar: Author Only Title Only Author and Title

Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing Genomic Features." Bioinformatics 26 (6): 841–42.
  Google Scholar: Author Only Title Only Author and Title

Roudier, François, Ikhlak Ahmed, Caroline Bérard, Alexis Sarazin, Tristan Mary-Huard, Sandra Cortijo, Daniel Bouyer, et al. 2011. "Integrative Epigenomic Mapping Defines Four Main Chromatin States in Arabidopsis." The EMBO Journal 30 (10): 1928–38.
  Google Scholar: Author Only Title Only Author and Title

Schep, Ruben, Eva K. Brinkman, Christ Leemans, Xabier Vergara, Robin H. van der Weide, Ben Morris, Tom van Schaik, et al. 2021. "Impact of Chromatin Context on Cas9-Induced DNA Double-Strand Break Repair Pathway Balance." Molecular Cell 81 (10): 2216–30.e10.
  Google Scholar: Author Only Title Only Author and Title

Schmid-Burgk, Jonathan L., Linyi Gao, David Li, Zachary Gardner, Jonathan Strecker, Blake Lash, and Feng Zhang. 2020. "Highly Parallel Profiling of Cas9 Variant Specificity." Molecular Cell 78 (4): 794–800.e8.
  Google Scholar: Author Only Title Only Author and Title

Sequeira-Mendes, Joana, Irene Aragüez, Ramón Peiró, Raul Mendez-Giraldez, Xiaoyu Zhang, Steven E. Jacobsen, Ugo Bastolla, and Crisanto Gutierrez. 2014. "The Functional Topography of the Arabidopsis Genome Is Organized in a Reduced Number of Linear Motifs of Chromatin States." The Plant Cell 26 (6): 2351–66.
  Google Scholar: Author Only Title Only Author and Title

Shen, Wei, Shuai Le, Yan Li, and Fuquan Hu. 2016. "SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation." PloS One 11 (10): e0163962.
  Google Scholar: Author Only Title Only Author and Title

Springer, Nathan M., and Robert J. Schmitz. 2017. "Exploiting Induced and Natural Epigenetic Variation for Crop Improvement." Nature Reviews. Genetics 18 (9): 563–75.
  Google Scholar: Author Only Title Only Author and Title

Stroud, Hume, Maxim V. C. Greenberg, Suhua Feng, Yana V. Bernatavichute, and Steven E. Jacobsen. 2013. "Comprehensive Analysis of Silencing Mutants Reveals Complex Regulation of the Arabidopsis Methylome." Cell 152 (1): 352–64.
  Google Scholar: Author Only Title Only Author and Title

Uusi-Mäkelä, Meri I. E., Harlan R. Barker, Carina A. Bäuerlein, Tomi Häkkinen, Matti Nykter, and Mika Rämet. 2018. "Chromatin

Accessibility Is Associated with CRISPR-Cas9 Efficiency in the Zebrafish (Danio Rerio)." PloS One 13 (4): e0196238.
 Google Scholar: Author Only Title Only Author and Title

Wang, Zhi-Ping, Hui-Li Xing, Li Dong, Hai-Yan Zhang, Chun-Yan Han, Xue-Chen Wang, and Qi-Jun Chen. 2015. "Egg Cell-Specific Promoter-Controlled CRISPR/Cas9 Efficiently Generates Homozygous Mutants for Multiple Target Genes in Arabidopsis in a Single Generation." Genome Biology 16 (July): 144.

Wu, Xuebing, David A. Scott, Andrea J. Kriz, Anthony C. Chiu, Patrick D. Hsu, Daniel B. Dadon, Albert W. Cheng, et al. 2014. "Genome-Wide Binding of the CRISPR Endonuclease Cas9 in Mammalian Cells." Nature Biotechnology 32 (7): 670–76.
 Google Scholar: Author Only Title Only Author and Title

Xiang, Xi, Giulia I. Corsi, Christian Anthon, Kunli Qu, Xiaoguang Pan, Xue Liang, Peng Han, et al. 2021. "Enhancing CRISPR-Cas9 gRNA Efficiency Prediction by Data Integration and Deep Learning." Nature Communications 12 (1): 3238.
 Google Scholar: Author Only Title Only Author and Title

Yan, Liuhua, Shaowei Wei, Yaorong Wu, Ruolan Hu, Hongju Li, Weicai Yang, and Qi Xie. 2015. "High-Efficiency Genome Editing in Arabidopsis Using YAO Promoter-Driven CRISPR/Cas9 System." Molecular Plant 8 (12): 1820–23.
 Google Scholar: Author Only Title Only Author and Title

Yarrington, Robert M., Surbhi Verma, Shaina Schwartz, Jonathan K. Trautman, and Dana Carroll. 2018. "Nucleosomes Inhibit Target Cleavage by CRISPR-Cas9 in Vivo." Proceedings of the National Academy of Sciences of the United States of America 115 (38): 9351–58.
 Google Scholar: Author Only Title Only Author and Title

Zemach, Assaf, and Gideon Grafi. 2003. "Characterization of Arabidopsis Thaliana Methyl-CpG-Binding Domain (MBD) Proteins." The Plant Journal: For Cell and Molecular Biology 34 (5): 565–72.
 Google Scholar: Author Only Title Only Author and Title

Zhang, Xiuren, Rossana Henriques, Shih-Shun Lin, Qi-Wen Niu, and Nam-Hai Chua. 2006. "Agrobacterium-Mediated Transformation of Arabidopsis Thaliana Using the Floral Dip Method." Nature Protocols 1 (2): 641–46.
 Google Scholar: Author Only Title Only Author and Title

Zhong, Zhenhui, Suhua Feng, Sascha H. Duttke, Magdalena E. Potok, Yiwei Zhang, Javier Gallego-Bartolomé, Wanlu Liu, and Steven E. Jacobsen. 2021. "DNA Methylation-Linked Chromatin Accessibility Affects Genomic Architecture in Arabidopsis." Proceedings of the National Academy of Sciences of the United States of America 118 (5). https://doi.org/10.1073/pnas.2023347118.
 Google Scholar: Author Only Title Only Author and Title

Acknowledgements and Funding

We thank all members of the Zhang Lab for helpful discussion. M.S. was supported by the Undergraduate Research Opportunities Program (UROP) from the University of Minnesota. Funding from NSF IOS-1934384 to N.M.S. partially supported this work. P.A.C. was supported by an ARC Discovery Early Career Award (DE200101748). F.Z. was supported by the startup fund from Department of Plant and Microbial Biology, University of Minnesota.

Author Contributions

T.W., P.A.C., N.M.S. and F.Z. designed research; T.W., P.A.C., K.M.R., and M.S. conducted experiments; T.W., K.M.R., and F.Z. analyzed data; T.W., N.M.S. and F.Z. wrote the paper.

Conflict of Interest

The authors declare no conflict of interest.