IGF2BP1/IMP1 contributes to autophagy modulation directly via MAP1LC3B

Patrick A. Williams[a], Louis R. Parham[a], Kofi K. Acheampong[b], Charles H. Danan[a], Xianghui Ma[a], Lauren A. Simon[a], Kaitlyn E. Naughton[a], Tatiana Karakasheva[a], Kelly A. Whelan[c,d], Donita C. Brady[e,f], Sydney M. Shaffer[b,g], Kathryn E. Hamilton[a]*

[a]Division of Gastroenterology, Hepatology, and Nutrition; Department of Pediatrics; Children's Hospital of Philadelphia; University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA; [b]Department of Pathology and Laboratory Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. [c]Department of Pathology & Laboratory Medicine, Lewis Katz School of Medicine at Temple University, Philadelphia, PA, USA; [d]Fels Institute for Cancer Research & Molecular Biology, Lewis Katz School of Medicine at Temple University, Philadelphia, PA, USA. [e]Department of Cancer Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA. [f]Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA. [g]Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, USA.

*Corresponding author
Kathryn E. Hamilton, PhD
903 Abramson Research Building
Children's Hospital of Philadelphia
3615 Civic Center Blvd.
Philadelphia, PA 19104, USA
267-426-5266
hamiltonk1@chop.edu
@KHamiltonPhD
ORCID: 0000-0001-9575-2499

Abstract

Homeostatic tissue maintenance requires coordinated regulation of metabolic processes including macroautophagy/autophagy. Autophagy dysregulation underlies numerous human diseases. Our prior work revealed that the RNA binding protein IGF2BP1/IMP1 binds transcripts encoding autophagy-related proteins. Furthermore, *Imp1* deletion in gastrointestinal epithelial cells in mice was associated with enhanced autophagy flux and improved recovery from tissue injury. In the current study, we evaluated molecular mechanisms underlying IMP1 modulation of autophagy. We report increased expression of autophagy protein microtubule associated protein 1 light-chain 3B (*MAP1LC3B*) in cells with *IMP1* deletion compared to control cells. *MAP1LC3B* expression was also increased in cells with mutated IMP1 phosphorylation sites. Luciferase reporter assays demonstrate increased translation of *MAP1LC3B*-3'UTR in cells with *IMP1* deletion. Furthermore, we find that IMP1 co-localizes with *MAP1LC3B* transcripts at homeostasis, and co-localization is decreased in cells where IMP1 phosphorylation sites are mutated. IMP1 localization with *MAP1LC3B* transcripts is also decreased following cell stress. Taken together, our data support a model whereby IMP1 regulates MAP1LC3B translation which can serve as a mechanism to calibrate autophagy levels in the cell. This new mechanism may be particularly important in gastrointestinal epithelial cells, where autophagy is essential for tissue recovery following injury, or in diseases such as inflammatory bowel disease where defective autophagy is implicated.

Abbreviations *IGF2* mRNA binding protein 1 (IMP1); Microtubule Associated Protein 1 Light Chain 3 Beta (MAP1LC3B)

Introduction

Studies of the mammalian gastrointestinal tract suggest diverse roles for the autophagy pathway in both homeostasis and disease. During homeostasis, epithelial cells that line the intestine and colon utilize autophagy to maintain microbial symbiosis, secrete antimicrobial lysozyme via secretory autophagy, maintain tight junctions, and protect from DNA damage via reduction of excessive reactive oxygen species (Asano et al., 2017; Bel et al., 2017; Conway et al., 2013; Gorbunov and Kiang, 2009; Nighot et al., 2015; Sadaghian Sadabad et al., 2015). Furthermore, variants in human autophagy genes were identified as susceptibility genes in inflammatory bowel disease (Rioux et al., 2007). Mice with intestinal epithelial-specific autophagy gene deletion exhibit enhanced susceptibility to inflammatory tissue damage (Adolph et al., 2013; Asano et al., 2017; Cadwell et al., 2008; Conway et al., 2013; Lassen et al., 2014; Trentesaux et al., 2020). Conversely, factors that increase autophagy, such as nutrient deprivation, enhance intestinal/colonic tissue regeneration (Rangan et al., 2019; Yousefi et al., 2018). These data together underscore diverse and important roles for autophagy in gastrointestinal epithelial cells, however, regulation of autophagy in the gut is incompletely understood.

Individual components of the autophagy pathway are regulated at multiple levels (Copetti et al., 2009; Feng et al., 2015; Lipinski et al., 2010; Yee et al., 2009). Emerging studies demonstrate post-transcriptional regulation of autophagy via long non-coding RNAs, microRNAs, and RNA binding proteins (RBPs) (Jing et al., 2015). Cytoplasmic RBPs regulate mRNA stability, localization, and translation by governing interactions with the translation machinery, cytoskeletal network, or other trans-acting factors that ultimately direct temporal and spatial protein translation. As such, RBPs contribute to an essential mechanism for cells to preferentially translate mRNAs in response to external stimuli or stress. Defining how RBPs contribute to autophagy regulation in specific cell types, such as gastrointestinal epithelial cells, could provide a basis for new therapies for gastrointestinal diseases.

Insulin-like growth factor 2 mRNA-binding protein 1 (IGF2BP1/IMP1) is an RBP expressed broadly during development and in human tumors, but its expression in adult tissues is low or absent(Bell et al., 2013). Our recent studies demonstrate upregulation of IMP1 in adult tissue including the intestinal/colonic stem cell niche during tissue damage in mice and in patients with inflammatory bowel disease (Chatterji et al., 2018; Chatterji et al., 2019). Studies in cell lines suggest a primary role for IMP1 to regulate cell proliferation, however, *in vivo* analyses revealed little effect on cell proliferation in mice with intestinal epithelial cell-specific deletion of Imp1 (*Imp1$^{\Delta IEC}$*) at homeostasis. Instead, the intestinal epithelium of *Imp1$^{\Delta IEC}$* mice exhibits increased autophagy flux associated with a decrease in intestinal tissue damage compared to control mice (Chatterji et al., 2018; Chatterji et al., 2019). IMP1 modulation of autophagy is consistent with *in vitro* studies implicating a role for IMP1 during cell stress and studies showing that cytoplasmic IMP1-granules contain transcripts encoding secretory pathway genes that may be related to the autophagy pathway(Jonson et al., 2007; Stohr et al., 2006; Zeng et al., 2020). While our prior data support the notion that IMP1 can act as a modulator of autophagy in gastrointestinal epithelial cells, a direct mechanism remains unclear.

Our goal in the present study was to evaluate direct regulation of autophagy transcripts by IMP1. We evaluated IMP1 regulation of autophagy at the RNA and protein levels and via direct translational regulation using dual-luciferase reporter assays. We examined the relative contribution of phosphorylated IMP1 (p-IMP1), known to regulate transcript binding and translation in other contexts (Dai et al., 2013; Huttelmaier et al., 2005; Lambrianidou et al., 2021), to autophagy protein expression and IMP1 cytoplasmic localization. Finally, we evaluated IMP1 co-localization with autophagy transcripts at homeostasis and in response to arsenite-induced cell stress. Herein, we report new data to support a previously unknown mechanism of autophagy pathway modulation via post-transcriptional regulation of MAP1LC3B by IMP1.

Results

IMP1 knockout cells exhibit increased MAP1LC3B protein expression

Prior work from our group demonstrated that *IMP1* deletion *in vivo* results in increased autophagy flux. In addition, we found via ribonucleoprotein immunoprecipitation (RIP)-qPCR that IMP1 can bind directly to autophagy transcripts *ATG5*, *ATG3*, and *MAP1LC3B* (Chatterji et al., 2019). The downstream effects of IMP1 binding to specific autophagy transcripts in not known. We evaluated the consequence of *IMP1* deletion on MAP1LC3B protein expression by examining *IMP1* wild type (*IMP1-WT*) and *IMP1* knockout (*IMP1-KO*) SW480 colon cancer cells. We used two different protein concentrations for immunoblotting to allow for quantification, with the lower concentration giving for better band resolution. Using an antibody specific to MAP1LC3B, we observed a significant increase the phosphatidylethanolamine(PE)-conjugated form of LC3 (LC3-II) in *IMP1-KO* compared to *IMP1-WT* cells (3.2 ± 0.32 vs 1.0 ± 0.12, $p<0.0001$), an effect that was increased with torin (3 hours), an mTOR inhibitor that induces autophagy (Thoreen et al., 2009). We also observed an increase in LC3-II in cells treated with V-ATPase inhibitor bafilomycin A_1, which inhibits autophagosome-lysosome fusion (Klionsky et al., 2008; Yamamoto et al., 1998), in *IMP1-KO* compared to *IMP1-WT* cells (4.9 ± 0.24 vs 2.8 ± 0.23, $p<0.0001$) (**Figure 1A and 1B**).

An increase in LC3-II in untreated cells can be interpreted as either increased autophagosomes or blocked autophagy, however, the further increase in LC3-II in IMP1-KO cells with bafilomycin A_1 treatment suggests increased autophagosomes/autophagy flux (Klionsky et al., 2021; Mauvezin and Neufeld, 2015; Mizushima and Yoshimori, 2007; Yoshii and Mizushima, 2017). We used a second antibody with reactivity to both MAP1LC3A and MAP1LC3B to evaluate LC3-II with saturating doses of bafilomycin A_1 treatment for 24 hours. In this case, there was little difference between cell lines (**Figure 1C & 1D**) suggesting that the MAP1LC3A isoform is not affected by IMP1 deletion and use of an antibody targeting both MAP1LC3A and MAP1LC3B instead of MAP1LC3B-only leads to a diminution in phenotype with IMP1-KO. Despite the muted phenotype by immunoblot with MAP1LC3A/B antibody detection, immunostaining using the MAP1LC3A/B antibody revealed increased puncta in IMP1-KO versus IMP1-WT cells (**Figure 1F**), supporting an overall effect of IMP1-KO to increase MAP1LC3B. Immunoblot analysis of other IMP1 targets ATG5 and ATG3 revealed no change between *IMP1-WT* and *IMP1-KO* cells (**Supplemental Figure 1A**). Taken together, these data suggest that IMP1-KO has a specific effect on MAP1LC3B compared to MAP1LCA or other its other putative RNA-binding targets, and that this effect may contribute to MAP1LC3 protein localization into puncta.

We next evaluated cells treated with exposure to torin (24 hours). We observed an increase in LC3-II between IMP1-WT and IMP1-KO cells at baseline and with torin treatment (**Supplemental Figure 1A & B**), however, cargo-associated protein p62 appeared higher in untreated IMP1-KO cells compared to IMP1-WT cells, indicated that at this time point, IMP1-KO cells may exhibit stalled autophagy flux. Taken together, results in Figure 1 indicate that at longer timepoints post-plating, autophagy flux could ultimately be saturated, rather than increased, in IMP1-KO compared to IMP1-WT cells. This is intriguing, since our prior studies in mouse organoids from Imp1$^{\Delta IEC}$ mice demonstrate enhanced autophagy flux (Chatterji et al., 2019). As such, it is possible that in longer cell culture intervals, a compensatory mechanism may be activated in IMP1-KO cells. Finally, evaluation of *MAP1LC3B* mRNA levels with real-time RT-PCR revealed no difference in *MAP1LC3B* expression in *IMP1-KO* compared to *IMP1-WT* cells (**Supplemental Figure 1C**), indicating that IMP1-KO is associated with increased MAP1LC3B primarily at the protein level. Taken together, we observe increased LC3-II levels with IMP1-KO via specific effects on MAP1LC3B, and not MAP1LC3A, ATG3, or ATG5, protein levels. Analysis of shorter timepoints post-plating and post-treatment reveal that upregulation may result in enhanced autophagy flux initially, whereas longer timepoints post-plating or post-treatment is associated with stalled autophagy flux based on accumulation of cargo-associated protein p62 (Klionsky et al., 2021).

IMP1 phosphorylation contributes to its modulation of MAP1LC3B protein expression

IMP1 contains a mTORC2-dependent phosphorylation site at serine 181 (S181) and a Src kinase-dependent site at tyrosine 398 (Y396) (**Figure 2A**), which can affect translation and/or localization of IMP1 targets (Dai et al., 2013; Huttelmaier et al., 2005; Lambrianidou et al., 2021). We therefore evaluated the relative contribution of IMP1 phosphorylation (p-IMP1) to its modulation of MAP1LC3B. We generated *IMP1-KO* HEK293 cell lines and introduced the following constructs: *control-GFP* (*IMP1-KO*), *IMP1-WT-GFP*, and phosphorylation mutants: *IMP1-S181A-GFP*, *IMP1-Y396F-GFP*, or double-mutant *IMP1-S181A/Y396F-GFP*. We observed increased MAP1LC3B protein expression in *control-GFP* (*IMP1-KO,*

3.282 ± 0.91), *IMP1-Y396F-GFP* (1.45 ± 0.31) and *IMP1-S181A/Y396F-GFP* (3.95 ± 1.38) mutant cells compared to *IMP1-WT-GFP* (1.0 ± 0.31), or *IMP1-S181A-GFP* (0.83 ±0.45), with statistical significance in *IMP1-Y396F-GFP* versus *IMP1-WT-GFP* cells (**Figure 2B&C**). Taken together, these data suggest that *IMP1-Y396* contributes to IMP1 modulation of MAP1LC3B protein expression.

We next used "Autophagy Assay, Red", a fluorescent probe that is activated when inserted into lipid membranes of autophagosomes and autolysosomes (**Supplemental Figure 2A**) (Czarnomysy et al., 2020), to determine the functional relevance of p-IMP1 for autophagy flux. We treated control-GFP (*IMP1-KO*), *IMP1-WT-GFP*, *IMP1-S181A-GFP*, *IMP1-Y396F-GFP*, or *IMP1-S181A/Y396F-GFP* cells with 100μM chloroquine (CQ) for 3 hours to inhibit lysosomal degradation, where accumulation of autophagic vesicles serves as an indicator for levels of autophagy flux (Mauthe et al., 2018). We observed a greater increase in autophagic vesicle accumulation in control-GFP (*IMP1-KO*), *IMP1-Y396F-GFP*, and *IMP1-S181A/Y396F-GFP* cells compared to *IMP1-WT-GFP* or *IMP1-S181A-GFP* cells, with statistically significant increases in GFP (*IMP1-KO*), *IMP1-Y396F-GFP* cells. These data together support the premise that IMP1-Y396, but not -S181 contributes to IMP1's regulation of MAP1LC3B, which can ultimately contribute to modulation of autophagy flux.

We next evaluated how p-IMP1 mutation affects its cellular localization, since prior studies suggest that IMP1 phosphorylation modulates IMP1 regulation of MYC (Lambrianidou et al., 2021). Imaging of GFP expression revealed primarily cytoplasmic staining in *IMP1-WT-GFP*, *IMP1-Y396F-GFP*, and *IMP1-S181A/Y396F-GFP* cells. *IMP1-S181A-GFP* cells exhibited a decrease in cytoplasmic staining with concomitant GFP expression in the nucleus (**Figure 3A**). Using cell fractionation followed by western blotting, we confirmed reduced cytoplasmic expression in *IMP1-S181A-GFP* cells compared to *IMP1-WT-GFP* cells (**Supplemental Figure 3A-B**). While IMP1 localization differences in *IMP1-S181A-GFP* cells is intriguing, we ultimately utilized *IMP1-WT-GFP* and *IMP1-S181A/Y396F-GFP* cells in subsequent mechanistic studies since *IMP1-S181A* did not affect MAP1LC3B expression.

IMP1 regulates MAP1LC3B in part via direct modulation of MAP1LC3B-3'UTR translation

Our previous ribosome-profiling data suggested that autophagy pathway translation is enhanced in *IMP1-KO* cells (Chatterji et al., 2019). Furthermore, analysis of published enhanced UV crosslinking and immunoprecipitation (eCLIP) studies showed enrichment of IMP1 binding sites in the *MAP1LC3B*-3'UTR (Chatterji et al., 2019; Conway et al., 2016). We therefore evaluated whether *IMP1* deletion was associated with increased translation of *MAP1LC3B* through the 3'UTR using a dual-luciferase reporter system (**Figure 3B**). We found that *IMP1-KO* (1.37 ± 0.1) and *IMP1-S181A/Y396F-GFP* (1.39 ± 0.24) cells transfected with *MAP1LC3B*-3'UTR exhibited increased luciferase reporter activity compared to *IMP1-WT* cells (1.09 ± 0.16; **Figure 3C**). Taken together, these data suggest that IMP1 may modulate MAP1LC3B in part via direct modulation of *MAP1LC3B*-3'UTR.

IMP1 phosphorylation contributes to dynamic localization of IMP1 in response to cell stress

Prior studies indicate that IMP1 can localize to mRNA-containing granules (mRNP granules) including p-bodies (PBs) and stress granules (SGs), which can be sites of translational quiescence and/or mRNA degradation (Bley et al., 2015; Hubstenberger et al., 2017). Under homeostatic conditions, IMP1-WT-GFP signal is mostly diffuse, however, we observed a small number of granules positively stained with the PB marker EDC4 that co-localized with *IMP1-WT-GFP* puncta (**Figure 4A**, vehicle, white arrows). Treatment with sodium arsenite (which elicits oxidative stress and can induce autophagy) (Bolt et al., 2010), results in increased GFP abundance in larger, more punctate granules that exhibit reduced overlap with EDC4+ granules denoted by non-overlapping red and green puncta (Figure 4A, arsenite-treated). GFP expression and localization appeared similar in *IMP1-S181A/Y396F-GFP* cells in both vehicle and sodium arsenite conditions and overlap with EDC4+ puncta were similarly decreased with sodium arsenite treatment (**Figure 4B**). There were fewer apparent EDC4 puncta in *IMP1-S181A/Y396F-GFP* cells in both vehicle and sodium arsenite conditions compared to *IMP1-WT-GFP* cells, indicating that while localization of IMP1 in PBs does not appear to be altered by mutant p-IMP1, there may be a general effect of mutant p-IMP1 on PB dynamics.

IMP1 has been described previously to localize in SGs and to be increased in SGs following cell stress (Bley et al., 2015). We therefore evaluated co-staining of IMP1 with SG marker PABP. As anticipated, a majority of IMP1 appeared to co-localize with PABP-marked granules following sodium arsenite treatment in *IMP1-WT-GFP* cells (**Figure 4C**, white arrows in insets in arsenite-treated cells). We observed reduced overall granule induction in *IMP1-S181A/Y396F-GFP* cells (**Figure 4D**), though the

granules that did form exhibited overlap between GFP and PABP (white arrows in insets with sodium arsenite treatment). Finally, since we observed an overall reduction in granule formation in response to sodium arsenite in mutant p-IMP1 cells, we performed time-lapse imaging of GFP in sodium arsenite-treated cells over 1.5 hours in *IMP1-WT-GFP*, *IMP1-S181A-GFP,* and *IMP1-Y396F-GFP* expressing cells. IMP1-WT-GFP cells exhibited robust cytoplasmic granule formation with sodium arsenite treatment (Movie S1), whereas granule formation appeared modestly reduced, but still intact, in *IMP1-S181A-GFP,* and *IMP1-Y396F-GFP* expressing cells (Movie S2 and S3). Collectively, these data support a role for p-IMP1 in contributing to cytoplasmic granule formation in response to stress, but its localization in specific granules warrants further investigation.

Cytoplasmic IMP1 co-localizes with MAP1LC3B transcript

Due to the complexity of p-IMP1's contribution to its localization in specific granules, we instead decided to evaluate whether IMP1 co-localizes with MAP1LC3B transcripts using single molecule mRNA FISH (smFISH) combined with immunofluorescent imaging of GFP. We observed co-localization of IMP1-GFP with *MAP1LC3B* transcripts (**Figure 5A**, white arrows), but not *ATG5* transcripts (**Figure 5B**). IMP1-S181A/Y396F-GFP cells exhibited reduced GFP co-localization with MAP1LC3B transcripts (**Figure 5C**). Arsenite treatment was associated with re-distribution of *MAP1LC3B* mRNA away from IMP1 granules in IMP1-WT-GFP cells (**Figure 5A**), while IMP1-S181A/Y396F-GFP cells exhibited less overall cytoplasmic IMP1 (Figure 5C). IMP1 did not co-localize with ATG5 mRNA in arsenite-treated cells (**Figure 5D**). We quantified co-localization between IMP1-GFP and *MAP1LC3B* fluorescence using rank weight coefficient and found significantly more overlap between *MAP1LC3B* and IMP1-WT-GFP compared to IMP1-S181A/Y396F-GFP (**Figure 5E**). Finally, we found that *MAP1LC3B* mRNA was distributed closer to the cell nucleus in IMP1-WT-GFP cells compared to IMP1-S181A/Y396F-GFP cells or IMP1-WT-GFP cells treated with arsenite (**Figure 5F**), supporting the notion that mutation of p-IMP1 mimics arsenite-induced cell stress. Taken together, these data suggest that at homeostasis, IMP1 contributes to *MAP1LC3B* transcript localization closer to the nucleus, whereas arsenite-induced stress appears to redistribute *MAP1LC3B* transcripts away from the nucleus, presumably when it is no longer sequestered by IMP1.

Discussion

Studies presented herein define new molecular mechanisms underlying our prior observation that autophagy flux is upregulated in intestinal epithelial cells in mice with *IMP1* deletion. Prior results from our lab suggested direct binding of IMP1 to multiple autophagy transcripts (Chatterji et al., 2019). In the present manuscript, we evaluated autophagy genes at the protein and RNA level in cells with *IMP1* deletion and found a robust increase in microtubule associated protein 1 light-chain 3B (MAP1LC3B) protein compared to other autophagy targets. Real-time RT-PCR results, where *MAP1LC3B* expression was not changed with *IMP1* deletion, suggest that IMP1 modulates MAP1LC3B almost exclusively at the protein level. Furthermore, mutation of the IMP1 Y-396 phosphorylation site, either alone or together with mutation in the IMP1 S-181 phosphorylation site, recapitulated the increase in MAP1LC3B we observed with *IMP1* deletion. We next evaluated the effect of *IMP1* deletion on *MAP1LC3B* translation using a dual luciferase reporter assay. We focused on the 3'UTR of *MAP1LC3B* based on our analysis of eCLIP data published by others (Conway et al., 2016), which showed a propensity of IMP1 binding at *MAP1LC3B*-3'UTR versus the coding sequence or 5'UTR (Chatterji et al., 2019). Our data demonstrate that cells with *IMP1* deletion exhibit higher translational activity for *MAP1LC3B*-3'UTR than in cells with wildtype IMP1 levels. This could be due to, at least in part, to IMP1 localizing out of translationally quiescent processing bodies and away from *MAP1LC3B* transcripts in response to stress, as demonstrated by co-staining with IMP1 with processing body marker EDC4. We used single molecule mRNA FISH to evaluate IMP1 protein localization with *MAP1LCB3* transcripts. We report that IMP1 co-localizes with *MAP1LCB3B* transcripts at homeostasis, and that both stress induction via sodium arsenite or mutation of p-IMP1 sites is associated with decreased co-localization. Furthermore, MAP1LC3B transcripts exhibit different localization in IMP1-WT versus p-IMP1 mutant or arsenite-treated cells, localizing closer to the nucleus in IMP1-WT cells. Taken together, our results support the notion that IMP1 modulates autophagy at homeostasis via direct regulation of *MAP1LC3B* translation and localization in a manner dependent, at least in part, upon p-IMP1 (**Figure 6**).

MAP1LC3 (also known as Atg8 in yeast) is a component of the autophagy cascade that, upon conjugation with phosphatidylethanolamine (PE), is essential for autophagosome formation and expansion (Ichimura et al., 2000; Nakatogawa et al., 2007). There are three human isoforms of MAP1LC3 (LC3A,

LC3B, LC3C) with differing expression patterns in tissue, all of which are associated with autophagic membranes in the cell (He et al., 2003). Our prior published study showed specific binding of IMP1 to the *MAP1LC3B* isoform (Chatterji et al., 2019). While mechanisms regulating expression and localization of MAP1LC3B protein are reported (Drake et al., 2010; Huang et al., 2015), post-transcriptional regulation of MAP1LC3B is only beginning to be elucidated. For example, knockdown of *BCL2-associated athanogene 3* (BAG3), a co-chaperone of the HSP70 system, was associated with decreased MAP1LC3B protein, but not mRNA, although the direct mechanism is not yet known (Rodriguez et al., 2016). Others reported regulation of *MAP1LC3B* transcripts by the microRNA mir204 (Mikhaylova et al., 2012). We find that *IMP1* deletion is associated with increased MAP1LC3B expression primarily at the protein level. Importantly, the increase in LC3-II observed with *IMP1* deletion does not appear to be attributed to impaired autophagy, as treatment with autophagy inhibitor bafilomycin A was associated with accumulation of LC3 and p62 in both control and *IMP1-KO* cells. As such, IMP1 appears to regulate the total amount of MAP1LC3B protein that is expressed, which could allow for more autophagy flux at homeostasis.

IMP1 is regulated in part via phosphorylation by either mTORC2 at S-181 or Src at Y-396 (Dai et al., 2013; Huttelmaier et al., 2005). Others reported recently diffuse cytoplasmic IMP1 in IMP1-S181A cells, and increased granule formation in IMP1-Y396F cells (Lambrianidou et al., 2021). In our hands, we confirm that IMP1-S181 contributes to diffuse cytoplasmic IMP1, however, we found additionally that cytoplasmic IMP1 decreased, and nuclear IMP1 expression increased, in IMP1-S181 mutant cells. More importantly, we did not observe a difference in MAP1LC3B protein expression or autophagy levels in IMP1-S181 mutant cells. Rather, mutation of IMP1-Y396 alone, or together with mutant IMP1-S181, was associated with increased MAP1LC3B expression and increased autophagy, similar to what is observed in *IMP1-KO* cells. Differences in IMP1 localization patterns between the present study and Lambrianidou *et* al. could be due to expression of phospho-mutants on an *IMP1-KO* versus *IMP1-WT* background, respectively. Taken together, we demonstrate a mechanistic link between IMP1 phosphorylation by Src and its regulation of MAP1LC3B protein expression.

Previous reports demonstrate IMP1 expression in processing bodies (PBs), which are typically translationally inactive (Hubstenberger et al., 2017), or in stress granules (SGs), which are dense RNA-protein aggregates that form during cell stress (Guzikowski et al., 2019; Niewidok et al., 2018; Stohr et al., 2006). IMP1-containing granules may also represent an entity distinct from PBs, SGs, or other granules (Jonson et al., 2007). In addition, several studies suggest IMP1 contributes to "post-transcriptional regulons" via its presence in granules containing multiple different transcripts, whereas another recent study suggested IMP1 localization in granules with mRNA singletons (Mateu-Regue et al., 2019). While its localization remains complex and requires further investigation, increased IMP1 expression during cell stress has been shown in multiple studies, including our own, (Bley et al., 2015; Wachter et al., 2013; Zeng et al., 2020). As such, IMP1 may contribute to "triaging" transcripts for translation as a mechanism to respond to cell stress. We find that IMP1 partially overlaps with PB marker EDC4 at homeostasis, but IMP1-expressing puncta are redistributed away from EDC4 granules following stress induction with sodium arsenite. We find fewer EDC4+ puncta in p-IMP1 mutant cells, however mutant p-IMP1 does still co-localizes with EDC4. We observed the same effect for IMP1 with SG marker PABP in arsenite-treated cells-an overall decrease in granules with mutant p-IMP1, but intact co-localization. Time-lapse imaging of IMP1-WT-GFP or p-IMP1 mutant cells confirm only partial impairment of IMP1 accumulation into puncta following sodium arsenite treatment. Prior studies suggest that IMP1 is dispensable for SG assembly (Bley et al., 2015). Since it is known that IMP1 localizes to multiple cytoplasmic granule subtypes, mutation of p-IMP1 could affect dynamic movement of IMP1 to multiple different granules in response to stress, resulting in a net decrease of these granules rather than a specific decrease in SGs (Bley et al., 2015). Taken together, we observed dynamic movement of cytoplasmic IMP1 in response to stress that appears to depend, at least in part, upon p-IMP1.

Finally, our finding that IMP1-WT-GFP colocalizes with *MAP1LC3B* mRNA and not *ATG5* mRNA supports the notion that IMP1 specifically associates with *MAP1LC3B* transcripts. Sodium arsenite-induced stress decreased IMP1 localization with *MAP1LC3B* mRNA, suggesting that during stress, IMP1 no longer complexes with *MAP1LC3B* mRNA. We find reduced IMP1-*MAP1LC3B* co-localization following stress or in p-IMP1 mutant cells. This is intriguing, as recent studies suggest a functional role for localized translation of specific transcripts in polarized epithelium, such as intestinal epithelial cells (Moor et al., 2017). We find that there is more *MAP1LC3B* transcript near the nucleus in IMP1-WT cells compared to p-IMP1 mutant cells or wildtype cells treated with sodium arsenite. Based on these findings in adherent cells, follow-up studies will

evaluate whether IMP1 contributes to specific localization of *MAP1LC3B* mRNA at apical versus basolateral compartments at homeostasis and in response to cell stress.

In summary, we report a new mechanism for post-transcriptional MAP1LC3B regulation. IMP1 sequestration of *MAP1LC3B* in cytoplasmic granules may poise homeostatic cells to respond to stress by protecting *MAP1LC3B* mRNA, and upon stress, releasing *MAP1LC3B* mRNA for translation. Our prior work demonstrated that IMP1 is expressed in a small, yet important, subset of cells in the intestine- namely stem cells and other cells in the intestinal crypt base. Both intestinal stem cells and Paneth cells may utilize the autophagy pathway for different functions (Asano et al., 2017; Bel et al., 2017; Trentesaux et al., 2020). In addition, recent studies demonstrate that autophagy is upregulated with IMP1 deletion in macrophages (Liu et al., 2022). As such, IMP1 modulation of MAP1LC3B could exhibit variable functional consequences depending on cell type expression, cell function, and upstream signaling. Studies evaluating the interplay between autophagy and post-transcriptional mechanisms continue to emerge (reviewed in (Frankel et al., 2017) and (Abildgaard et al., 2020)). Several pathologies, including neurodegeneration and inflammatory bowel disease (IBD) are associated with abnormal or defective autophagy. Future studies in human tissues and/or organoids are required to place the IMP1-*MAP1LC3B* regulatory axis into the context of human diseases. Understanding post-transcriptional mechanisms of autophagy regulation in tissues such as the gastrointestinal tract will provide a basis for understanding human disease pathogenesis, and ultimately, new therapeutic strategies for these diseases.

Acknowledgments
The authors thank Dr. Joseph Baur for helpful discussions and feedback. The authors thank Drs. Sarah Andres and Todd Strochlic for providing plasmid backbones used in the present studies.

Funding
NIH R01DK124369 (KEH); NIH R21ES031533 (KEH); Institutional Development Funds from Children's Hospital of Philadelphia Research Institute (KEH); Penn Center for Molecular Studies in Digestive and Liver Diseases NIH P30DK050306 (KEH); NIH R01DK12115 (KAW); NIH DP5OD028144 (SMS).

Author Contributions
Study concept and design: Williams, Hamilton. Acquisition of data: Williams, Parham, Acheampong, Ma, Simon, Naughton, Shaffer. Analysis and interpretation of data: Williams, Parham, Acheampong, Ma, Simon, Danan, Karakasheva, Whelan, Shaffer, Hamilton. Drafting of the manuscript: Williams, Hamilton. Analysis of the manuscript for intellectual and conceptual content: Williams, Whelan, Brady, Hamilton. Study Supervision: Hamilton.

Declaration of Interests
The authors declare no conflicts.

Methods
Generation of HEK293 and SW480 IMP1-KO cell lines
Indicated cells were seeded into a 6-well plate and grown to 60% confluency under standard conditions: growth medium- DMEM with 4.5 g/L glucose and L-glutamine with 10 mM sodium pyruvate (Corning #10-017-CV), 10% FBS (Peak Serum PS-FB1), and Penn/Strep (Gibco #15140122), and in 5% CO_2. IMP1 deletion was performed using the IGF2BP1 CRISPR/Cas9 knockout plasmid (Santa Cruz Cat# sc-401703) and IGF2BP1 HDR Plasmid (Santa Cruz Cat# sc-401703-HDR) protocol as described by the manufacturer. Cells were selected using 2 ug/mL of puromycin for 2 passages and knockout was confirmed using western blotting.

Generation of cell lysates and western blot
Cells were lysed with lysis buffer (Cell Lysis reagent (Cell Signaling #9803S), Halt Protease Inhibitor Cocktail (Life technologies #78430), 1 mM Sodium Orthovanadate, 10 uM Sodium Fluoride) on ice and spun at 9391 rcf at 4°C for 10 minutes. Running buffer (final concentration 62.5 mM Tris-HCl pH 6.8, 2.5% SDS, 0.002% Bromophenol Blue, 5% B-mercaptoethanol, and 10% glycerol) was added to supernatant and boiled for 6 minutes and loaded onto an SDS-PAGE gel with a percentage of 10%, 12.5%, or 15% acrylamide. Samples were run and transferred onto PVDF membrane using the Bio-rad Trans-blot Turbo system and then blocked in either 5% Milk or BSA in TBS-T (20.7 mM Tris Base, 150.7 NaCl, 0.1% Tween-

20, pH 7.6) for 1 hour at room temperature. Primary antibodies were diluted in 5% Milk or BSA in TBS-T and incubated with gentle rocking at 4°C for 24 hours. Blots were washed with TBS-T and then incubated in appropriate secondary antibody for 1 hour at room temperature with rocking. Following washing, blots were incubated with luminol reagent (Santa Cruz Biotechnology sc-2048) per manufacturer protocol and exposed to autoradiography film. All antibodies are listed in Table 1.

Real- time RT-PCR
Total RNA was isolated from cells using the Quick-RNA Mini-Prep Kit (Zymo Research R1054) and cDNA was made using the High-Capacity cDNA reverse Transcription Kit (Applied Biosystems #4368813). Real-time PCR assays were performed on cDNA using the TAQMAN Fast Gene Expression Universal PCR Master Mix 2X (Applied Biosystems #4352042) per the manufacturer protocol in conjunction with the Applied Biosystems Step-One Plus. Gene expression was calculated using $R=2^{(-DDCt)}$ method. All qPCR assays are listed in Table 2.

Plasmid construct generation
IMP1-GFP was cloned by amplifying PB-FLAG-IMP1 - short 3'UTR (obtained from Dr. Sarah Andres, OHSU) using primers (F'-CCTGGTACCATGAACAAGCTTTACATC and R'ACGCATGAATTCCTTCCTCCGTGCCTGGGCCTG) and was purified from 1% agarose gel. Fragment was digested using EcoRI and KpnI and ligated into pcDNA3-eGFP (Addgene) and sequenced. IMP1-S181A-GFP and IMP1-Y396-GFP were generated by site directed mutagenesis of IMP1-GFP by Genewiz LLC, and IMP1-S181A/Y396F-GFP was generated by site directed mutagenesis of IMP1-S181A-GFP by Genewiz LLC. 3'UTR luciferase constructs were generated by insertion through restriction digestion of a synthetic peptide corresponding to appropriate gene sequence into psiCHECK-2 (Promega C8021) at the XhoI and NotI sites by Genewiz LLC.

Generation of HEK293- IMP1-GFP mutant cells
HEK293-*IMP1-KO* cells described above were used to seed 6-well plates to 60% confluency. Cells were then transfected with pcDNA-GFP, pcDNA-IMP1-GFP, pcDNA-IMP1-S181A-GFP, pcDNA-IMP1-Y396F-GFP, or pcDNA-IMP1-S181A/Y396F-GFP plasmids using lipofectamine 3000 and 2.5 ug of DNA/well for 24 hours in DMEM+10% FBS. Following 24 hours transfection, media was replaced with standard media (DMEM, 10% FBS, Penn/Strep) for 24 hours. Selection with 500 ug/mL G418 occurred for 4-5 days to enrich for a stably expressing population. After selection, cells were sorted for highest 5% GFP expression using Zombie Aqua (BioLegend 423101) and the MoFlo Astrios (Beckman Coulter).

Immunofluorescence imaging
Cells were grown on sterilized glass coverslips in 6-well plates under standard conditions until 70% confluent, washed with 1X PBS and fixed with 4% paraformaldehyde (thermoscientific J19943-K2) for 30 minutes at room temperature. Cells were washed again with PBS and then permeabilized in 0.1% triton X-100 for 5 minutes and then 0.5% triton X-100 for 20 minutes before washing with PBS. Cells blocked in 3% BSA for 1 hour, then incubated in primary at 4°C for 24 hours followed by washing and incubation with secondary antibody for 30 minutes. Cells stained with DAPI and mounted to coverslips using VECTASHIELD Vibrance Antifade Mounting Medium (Vector Laboratories H-1700-2) and sealed with nail polish (Wet'n'Wild) and imaged using a Keyence BZ-X700 microscope. Images were processed using BZ-X Analyzer (Keyence).

Cellular fractionation assay
Cells were grown in 6-well plates until reaching 80% confluency. Cells were trypsinized, pelleted and washed with PBS. Cell pellets were used for Subcellular Protein Fractionation Kit for Cultured Cells (Thermo Scientific Cat# 78840) per the manufacturer instructions and lysates were run on a denaturing gel.

Autophagy Assay Kit (Red) for measuring autophagic flux
To evaluate autophagic flux, HEK293T cells were treated either with DMSO (control) or chloroquine (100uM) for 3 hours before being collected and stained with the Autophagy Probe, Red for subsequent flow cytometric analysis. Autophagy Probe, Red kit (Abcam, ab270790) was performed according to manufacturer's instructions. One vial of the provided Autophagy Assay, Red powder was reconstituted in 100ul of DMSO to form a 250X stock solution. Immediately before staining, a 50X solution of the probe was

prepared in PBS and 10ul of the solution was added to 490uL of each sample cell suspension. The samples were shielded from light and incubated for 30 minutes at 37°C. Following staining, cells were washed and resuspended in 500ul of 1X PBS with 4% FBS and were analysed via flow cytometry. Dead cells were excluded from the analysis with the use of DAPI staining. Samples with less than 50% viability were excluded from analyses. Flow cytometry was performed on the LSRFortessa (BD Bioscience) and analysed with FlowJo software. Data were analyzed by comparing the median fluorescence intensity (MFI) of untreated (control) and chloroquine (CQ) treated samples. The fold change in MFI between each treatment group was calculated and was compared using an ordinary one-way ANOVA with Dunnett's multiple comparison test.

Luciferase translation assay
HEK293 *IMP1-WT* or *KO* were grown in 12-well plates to 60% confluency followed by transfection with previously described siCHECK2 constructs using lipofectamine 3000 reagent (Thermofisher Cat#L300008) according to manufacturer protocol at 0.5 ug of DNA per well for 24 hours in DMEM+10% FBS. Transfected cells were then trypsinized and used to seed 96-well plates at a density of 8.5e4 cells/well and allowed to grow overnight. Dual luciferase assays were then performed using Promega Dual-Luciferase Reporter Assay System (Promega Cat#E1910) following manufacturer protocol. Briefly, cells were lysed in 80 uL 1x Passive Lysis buffer at room temperature for 15 minutes with vigorous agitation. After agitation, 50 uL of lysate were transferred to a 96-well black bottom plate and read using dual injector luminometer set to dispense 50 uL of reconstituted Luciferase assay substrate followed by Stop & Glo buffer. For values, renilla luciferase (has the 3'UTR insert) is normalized to firefly luciferase, which is a built-in control. Expression was than calculated by normalizing those plasmids with insert to empty control plasmid.

Live cell imaging
6-well glass bottom tissue culture plates (MatTek Corporation P06G-1.5-10-F.S.) were coated with 34 ug/mL collagen (Sigma-Aldrich, C5533-5MG) for 1 hour at 37 °C then seeded with cells grown in standard media until 70% confluence. Cells were treated with 100 uM sodium arsenite for 90 minutes in an onstage incubator (37°C, 5% CO_2) in conjunction with the Keyence BZ-X700 Microscope.

Single molecule fluorescent in situ hybridization (smFISH)
Single molecule FISH studies were done as described previously (Shaffer et al., 2013). Briefly, cells were grown on coverslips in 6-well plates until confluent. Cells were then fixed with 4% paraformaldehyde for 10 minutes followed by washing and permeabilization in 70% ethanol overnight at 4C. Cells washed in FISH wash buffer (2x SSC, 10% Formamide) then incubated overnight with probes in hybridization buffer (10% dextran sulfate, 10% formamide, 2x SSC) at 37C at dilution of 1:100. After washing, cells stained with DAPI followed by wash in 2x SSC buffer before being mounted using VECTASHIELD Vibrance Antifade Mounting Medium (Vector Laboratories H-1700-2). Cells imaged using inverted Nikon Ti2-E microscope equipped with a SOLA SE U-nIR light engine (Lumencor), ORCA-Flash 4.0 V3 sCMOS camera (Hamamatsu), with ×60 Plan-Apo λ (MRD01605) or ×20 Plan-Apo λ (Nikon MRD00205). Specific probes are listed in Table 3. Images analyses were performed using ImageJ and CellProfiler (Dunn et al., 2011; McQuin et al., 2018).

Statistical methods
Data were analysed using unpaired, two-tailed student's t-tests, one-way ANOVA, or two-way ANOVA with post-hoc tests (Dunnett's or Tukey's) and p-values indicated in individual figures. Analyses were performed on data from a minimum of three experiments unless otherwise noted, in which case multiple fields are analysed. All values are presented as mean ± standard error of mean (SEM) unless noted otherwise. Specific experimental replicates are described in each figure legend.

References
Abildgaard, M.H., S.H. Brynjolfsdottir, and L.B. Frankel. 2020. The Autophagy-RNA Interplay: Degradation and Beyond. *Trends Biochem Sci*. 45:845-857.
Adolph, T.E., M.F. Tomczak, L. Niederreiter, H.J. Ko, J. Bock, E. Martinez-Naves, J.N. Glickman, M. Tschurtschenthaler, J. Hartwig, S. Hosomi, M.B. Flak, J.L. Cusick, K. Kohno, T. Iwawaki, S. Billmann-Born, T. Raine, R. Bharti, R. Lucius, M.N. Kweon, S.J. Marciniak, A. Choi, S.J. Hagen, S.

Schreiber, P. Rosenstiel, A. Kaser, and R.S. Blumberg. 2013. Paneth cells as a site of origin for intestinal inflammation. *Nature*. 503:272-276.

Asano, J., T. Sato, S. Ichinose, M. Kajita, N. Onai, S. Shimizu, and T. Ohteki. 2017. Intrinsic Autophagy Is Required for the Maintenance of Intestinal Stem Cells and for Irradiation-Induced Intestinal Regeneration. *Cell Rep*. 20:1050-1060.

Bel, S., M. Pendse, Y. Wang, Y. Li, K.A. Ruhn, B. Hassell, T. Leal, S.E. Winter, R.J. Xavier, and L.V. Hooper. 2017. Paneth cells secrete lysozyme via secretory autophagy during bacterial infection of the intestine. *Science*. 357:1047-1052.

Bell, J.L., K. Wachter, B. Muhleck, N. Pazaitis, M. Kohn, M. Lederer, and S. Huttelmaier. 2013. Insulin-like growth factor 2 mRNA-binding proteins (IGF2BPs): post-transcriptional drivers of cancer progression? *Cell Mol Life Sci*. 70:2657-2675.

Bley, N., M. Lederer, B. Pfalz, C. Reinke, T. Fuchs, M. Glass, B. Moller, and S. Huttelmaier. 2015. Stress granules are dispensable for mRNA stabilization during cellular stress. *Nucleic Acids Res*. 43:e26.

Bolt, A.M., R.M. Byrd, and W.T. Klimecki. 2010. Autophagy is the predominant process induced by arsenite in human lymphoblastoid cell lines. *Toxicol Appl Pharmacol*. 244:366-373.

Cadwell, K., J.Y. Liu, S.L. Brown, H. Miyoshi, J. Loh, J.K. Lennerz, C. Kishi, W. Kc, J.A. Carrero, S. Hunt, C.D. Stone, E.M. Brunt, R.J. Xavier, B.P. Sleckman, E. Li, N. Mizushima, T.S. Stappenbeck, and H.W.t. Virgin. 2008. A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature*. 456:259-263.

Chatterji, P., K.E. Hamilton, S. Liang, S.F. Andres, H.R.S. Wijeratne, R. Mizuno, L.A. Simon, P.D. Hicks, S.W. Foley, J.R. Pitarresi, A.J. Klein-Szanto, A.T. Mah, L. Van Landeghem, B.D. Gregory, C.J. Lengner, B.B. Madison, P. Shah, and A.K. Rustgi. 2018. The LIN28B-IMP1 post-transcriptional regulon has opposing effects on oncogenic signaling in the intestine. *Genes Dev*. 32:1020-1034.

Chatterji, P., P.A. Williams, K.A. Whelan, F.C. Samper, S.F. Andres, L.A. Simon, L.R. Parham, R. Mizuno, E.T. Lundsmith, D.S. Lee, S. Liang, H.S. Wijeratne, S. Marti, L. Chau, V. Giroux, B.J. Wilkins, G.D. Wu, P. Shah, G.G. Tartaglia, and K.E. Hamilton. 2019. Posttranscriptional regulation of colonic epithelial repair by RNA binding protein IMP1/IGF2BP1. *EMBO Rep*. 20:e47074.

Conway, A.E., E.L. Van Nostrand, G.A. Pratt, S. Aigner, M.L. Wilbert, B. Sundararaman, P. Freese, N.J. Lambert, S. Sathe, T.Y. Liang, A. Essex, S. Landais, C.B. Burge, D.L. Jones, and G.W. Yeo. 2016. Enhanced CLIP Uncovers IMP Protein-RNA Targets in Human Pluripotent Stem Cells Important for Cell Adhesion and Survival. *Cell Rep*. 15:666-679.

Conway, K.L., P. Kuballa, J.H. Song, K.K. Patel, A.B. Castoreno, O.H. Yilmaz, H.B. Jijon, M. Zhang, L.N. Aldrich, E.J. Villablanca, J.M. Peloquin, G. Goel, I.A. Lee, E. Mizoguchi, H.N. Shi, A.K. Bhan, S.Y. Shaw, S.L. Schreiber, H.W. Virgin, A.F. Shamji, T.S. Stappenbeck, H.C. Reinecker, and R.J. Xavier. 2013. Atg16l1 is required for autophagy in intestinal epithelial cells and protection of mice from Salmonella infection. *Gastroenterology*. 145:1347-1357.

Copetti, T., C. Bertoli, E. Dalla, F. Demarchi, and C. Schneider. 2009. p65/RelA modulates BECN1 transcription and autophagy. *Mol Cell Biol*. 29:2594-2608.

Czarnomysy, R., D. Radomska, A. Muszynska, J.M. Hermanowicz, I. Prokop, A. Bielawska, and K. Bielawski. 2020. Evaluation of the Anticancer Activities of Novel Transition Metal Complexes with Berenil and Nitroimidazole. *Molecules*. 25:2860.

Dai, N., J. Christiansen, F.C. Nielsen, and J. Avruch. 2013. mTOR complex 2 phosphorylates IMP1 cotranslationally to promote IGF2 production and the proliferation of mouse embryonic fibroblasts. *Genes Dev*. 27:301-312.

Drake, K.R., M. Kang, and A.K. Kenworthy. 2010. Nucleocytoplasmic distribution and dynamics of the autophagosome marker EGFP-LC3. *PLoS One*. 5:e9806.

Dunn, K.W., M.M. Kamocka, and J.H. McDonald. 2011. A practical guide to evaluating colocalization in biological microscopy. *Am J Physiol Cell Physiol*. 300:C723-742.

Feng, Y., Z. Yao, and D.J. Klionsky. 2015. How to control self-digestion: transcriptional, post-transcriptional, and post-translational regulation of autophagy. *Trends in cell biology*. 25:354-363.

Frankel, L.B., M. Lubas, and A.H. Lund. 2017. Emerging connections between RNA and autophagy. *Autophagy*. 13:3-23.

Gorbunov, N.V., and J.G. Kiang. 2009. Up-regulation of autophagy in small intestine Paneth cells in response to total-body gamma-irradiation. *The Journal of pathology*. 219:242-252.

Guzikowski, A.R., Y.S. Chen, and B.M. Zid. 2019. Stress-induced mRNP granules: Form and function of processing bodies and stress granules. *Wiley Interdiscip Rev RNA*. 10:e1524.

He, H., Y. Dang, F. Dai, Z. Guo, J. Wu, X. She, Y. Pei, Y. Chen, W. Ling, C. Wu, S. Zhao, J.O. Liu, and L. Yu. 2003. Post-translational modifications of three members of the human MAP1LC3 family and detection of a novel type of modification for MAP1LC3B. *J Biol Chem*. 278:29278-29287.

Huang, R., Y. Xu, W. Wan, X. Shou, J. Qian, Z. You, B. Liu, C. Chang, T. Zhou, J. Lippincott-Schwartz, and W. Liu. 2015. Deacetylation of nuclear LC3 drives autophagy initiation under starvation. *Mol Cell*. 57:456-466.

Hubstenberger, A., M. Courel, M. Benard, S. Souquere, M. Ernoult-Lange, R. Chouaib, Z. Yi, J.B. Morlot, A. Munier, M. Fradet, M. Daunesse, E. Bertrand, G. Pierron, J. Mozziconacci, M. Kress, and D. Weil. 2017. P-Body Purification Reveals the Condensation of Repressed mRNA Regulons. *Mol Cell*. 68:144-157 e145.

Huttelmaier, S., D. Zenklusen, M. Lederer, J. Dictenberg, M. Lorenz, X. Meng, G.J. Bassell, J. Condeelis, and R.H. Singer. 2005. Spatial regulation of beta-actin translation by Src-dependent phosphorylation of ZBP1. *Nature*. 438:512-515.

Ichimura, Y., T. Kirisako, T. Takao, Y. Satomi, Y. Shimonishi, N. Ishihara, N. Mizushima, I. Tanida, E. Kominami, M. Ohsumi, T. Noda, and Y. Ohsumi. 2000. A ubiquitin-like system mediates protein lipidation. *Nature*. 408:488-492.

Jing, Z., W. Han, X. Sui, J. Xie, and H. Pan. 2015. Interaction of autophagy with microRNAs and their potential therapeutic implications in human cancers. *Cancer letters*. 356:332-338.

Jonson, L., J. Vikesaa, A. Krogh, L.K. Nielsen, T. Hansen, R. Borup, A.H. Johnsen, J. Christiansen, and F.C. Nielsen. 2007. Molecular composition of IMP1 ribonucleoprotein granules. *Mol Cell Proteomics*. 6:798-811.

Klionsky, D.J., A.K. Abdel-Aziz, S. Abdelfatah, M. Abdellatif, A. Abdoli, S. Abel, H. Abeliovich, M.H. Abildgaard, Y.P. Abudu, A. Acevedo-Arozena, I.E. Adamopoulos, K. Adeli, T.E. Adolph, A. Adornetto, E. Aflaki, G. Agam, A. Agarwal, B.B. Aggarwal, M. Agnello, P. Agostinis, J.N. Agrewala, A. Agrotis, P.V. Aguilar, S.T. Ahmad, Z.M. Ahmed, U. Ahumada-Castro, S. Aits, S. Aizawa, Y. Akkoc, T. Akoumianaki, H.A. Akpinar, A.M. Al-Abd, L. Al-Akra, A. Al-Gharaibeh, M.A. Alaoui-Jamali, S. Alberti, E. Alcocer-Gomez, C. Alessandri, M. Ali, M.A. Alim Al-Bari, S. Aliwaini, J. Alizadeh, E. Almacellas, A. Almasan, A. Alonso, G.D. Alonso, N. Altan-Bonnet, D.C. Altieri, E.M.C. Alvarez, S. Alves, C. Alves da Costa, M.M. Alzaharna, M. Amadio, C. Amantini, C. Amaral, S. Ambrosio, A.O. Amer, V. Ammanathan, Z. An, S.U. Andersen, S.A. Andrabi, M. Andrade-Silva, A.M. Andres, S. Angelini, D. Ann, U.C. Anozie, M.Y. Ansari, P. Antas, A. Antebi, Z. Anton, T. Anwar, L. Apetoh, N. Apostolova, T. Araki, Y. Araki, K. Arasaki, W.L. Araujo, J. Araya, C. Arden, M.A. Arevalo, S. Arguelles, E. Arias, J. Arikkath, H. Arimoto, A.R. Ariosa, D. Armstrong-James, L. Arnaune-Pelloquin, A. Aroca, D.S. Arroyo, I. Arsov, R. Artero, D.M.L. Asaro, M. Aschner, M. Ashrafizadeh, O. Ashur-Fabian, A.G. Atanasov, A.K. Au, P. Auberger, H.W. Auner, L. Aurelian, et al. 2021. Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition)(1). *Autophagy*. 17:1-382.

Klionsky, D.J., Z. Elazar, P.O. Seglen, and D.C. Rubinsztein. 2008. Does bafilomycin A1 block the fusion of autophagosomes with lysosomes? *Autophagy*. 4:849-850.

Lambrianidou, A., E. Sereti, K. Soupsana, C. Komini, K. Dimas, and T. Trangas. 2021. mTORC2 deploys the mRNA binding protein IGF2BP1 to regulate c-MYC expression and promote cell survival. *Cell Signal*. 80:109912.

Lassen, K.G., P. Kuballa, K.L. Conway, K.K. Patel, C.E. Becker, J.M. Peloquin, E.J. Villablanca, J.M. Norman, T.C. Liu, R.J. Heath, M.L. Becker, L. Fagbami, H. Horn, J. Mercer, O.H. Yilmaz, J.D. Jaffe, A.F. Shamji, A.K. Bhan, S.A. Carr, M.J. Daly, H.W. Virgin, S.L. Schreiber, T.S. Stappenbeck, and R.J. Xavier. 2014. Atg16L1 T300A variant decreases selective autophagy resulting in altered cytokine signaling and decreased antibacterial defense. *Proc Natl Acad Sci U S A*. 111:7741-7746.

Lipinski, M.M., G. Hoffman, A. Ng, W. Zhou, B.F. Py, E. Hsu, X. Liu, J. Eisenberg, J. Liu, J. Blenis, R.J. Xavier, and J. Yuan. 2010. A genome-wide siRNA screen reveals multiple mTORC1 independent signaling pathways regulating autophagy under normal nutritional conditions. *Developmental cell*. 18:1041-1052.

Liu, M., G. Tao, Y. Cao, Y. Hu, and Z. Zhang. 2022. Silencing of IGF2BP1 restrains ox-LDL-induced lipid accumulation and inflammation by reducing RUNX1 expression and promoting autophagy in macrophages. *J Biochem Mol Toxicol*:e22994.

Mateu-Regue, A., J. Christiansen, F.O. Bagger, O. Winther, C. Hellriegel, and F.C. Nielsen. 2019. Single mRNP Analysis Reveals that Small Cytoplasmic mRNP Granules Represent mRNA Singletons. *Cell Rep*. 29:736-748 e734.

Mauthe, M., I. Orhon, C. Rocchi, X. Zhou, M. Luhr, K.J. Hijlkema, R.P. Coppes, N. Engedal, M. Mari, and F. Reggiori. 2018. Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy*. 14:1435-1455.

Mauvezin, C., and T.P. Neufeld. 2015. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-lysosome fusion. *Autophagy*. 11:1437-1438.

McQuin, C., A. Goodman, V. Chernyshev, L. Kamentsky, B.A. Cimini, K.W. Karhohs, M. Doan, L. Ding, S.M. Rafelski, D. Thirstrup, W. Wiegraebe, S. Singh, T. Becker, J.C. Caicedo, and A.E. Carpenter. 2018. CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol*. 16:e2005970.

Mikhaylova, O., Y. Stratton, D. Hall, E. Kellner, B. Ehmer, A.F. Drew, C.A. Gallo, D.R. Plas, J. Biesiada, J. Meller, and M.F. Czyzyk-Krzeska. 2012. VHL-regulated MiR-204 suppresses tumor growth through inhibition of LC3B-mediated autophagy in renal clear cell carcinoma. *Cancer Cell*. 21:532-546.

Mizushima, N., and T. Yoshimori. 2007. How to interpret LC3 immunoblotting. *Autophagy*. 3:542-545.

Moor, A.E., M. Golan, E.E. Massasa, D. Lemze, T. Weizman, R. Shenhav, S. Baydatch, O. Mizrahi, R. Winkler, O. Golani, N. Stern-Ginossar, and S. Itzkovitz. 2017. Global mRNA polarization regulates translation efficiency in the intestinal epithelium. *Science*. 357:1299-1303.

Nakatogawa, H., Y. Ichimura, and Y. Ohsumi. 2007. Atg8, a ubiquitin-like protein required for autophagosome formation, mediates membrane tethering and hemifusion. *Cell*. 130:165-178.

Niewidok, B., M. Igaev, A. Pereira da Graca, A. Strassner, C. Lenzen, C.P. Richter, J. Piehler, R. Kurre, and R. Brandt. 2018. Single-molecule imaging reveals dynamic biphasic partition of RNA-binding proteins in stress granules. *J Cell Biol*. 217:1303-1318.

Nighot, P.K., C.A. Hu, and T.Y. Ma. 2015. Autophagy enhances intestinal epithelial tight junction barrier function by targeting claudin-2 protein degradation. *J Biol Chem*. 290:7234-7246.

Rangan, P., I. Choi, M. Wei, G. Navarrete, E. Guen, S. Brandhorst, N. Enyati, G. Pasia, D. Maesincee, V. Ocon, M. Abdulridha, and V.D. Longo. 2019. Fasting-Mimicking Diet Modulates Microbiota and Promotes Intestinal Regeneration to Reduce Inflammatory Bowel Disease Pathology. *Cell Rep*. 26:2704-2719 e2706.

Rioux, J.D., R.J. Xavier, K.D. Taylor, M.S. Silverberg, P. Goyette, A. Huett, T. Green, P. Kuballa, M.M. Barmada, L.W. Datta, Y.Y. Shugart, A.M. Griffiths, S.R. Targan, A.F. Ippoliti, E.J. Bernard, L. Mei, D.L. Nicolae, M. Regueiro, L.P. Schumm, A.H. Steinhart, J.I. Rotter, R.H. Duerr, J.H. Cho, M.J. Daly, and S.R. Brant. 2007. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nature genetics*. 39:596-604.

Rodriguez, A.E., C. Lopez-Crisosto, D. Pena-Oyarzun, D. Salas, V. Parra, C. Quiroga, T. Morawe, M. Chiong, C. Behl, and S. Lavandero. 2016. BAG3 regulates total MAP1LC3B protein levels through a translational but not transcriptional mechanism. *Autophagy*. 12:287-296.

Sadaghian Sadabad, M., A. Regeling, M.C. de Goffau, T. Blokzijl, R.K. Weersma, J. Penders, K.N. Faber, H.J. Harmsen, and G. Dijkstra. 2015. The ATG16L1-T300A allele impairs clearance of pathosymbionts in the inflamed ileal mucosa of Crohn's disease patients. *Gut*. 64:1546-1552.

Shaffer, S.M., M.T. Wu, M.J. Levesque, and A. Raj. 2013. Turbo FISH: a method for rapid single molecule RNA FISH. *PLoS One*. 8:e75120.

Stohr, N., M. Lederer, C. Reinke, S. Meyer, M. Hatzfeld, R.H. Singer, and S. Huttelmaier. 2006. ZBP1 regulates mRNA stability during cellular stress. *J Cell Biol*. 175:527-534.

Thoreen, C.C., S.A. Kang, J.W. Chang, Q. Liu, J. Zhang, Y. Gao, L.J. Reichling, T. Sim, D.M. Sabatini, and N.S. Gray. 2009. An ATP-competitive mammalian target of rapamycin inhibitor reveals rapamycin-resistant functions of mTORC1. *J Biol Chem*. 284:8023-8032.

Trentesaux, C., M. Fraudeau, C.L. Pitasi, J. Lemarchand, S. Jacques, A. Duche, F. Letourneur, E. Naser, K. Bailly, A. Schmitt, C. Perret, and B. Romagnolo. 2020. Essential role for autophagy protein ATG7 in the maintenance of intestinal stem cell integrity. *Proc Natl Acad Sci U S A*. 117:11136-11146.

Wachter, K., M. Kohn, N. Stohr, and S. Huttelmaier. 2013. Subcellular localization and RNP formation of IGF2BPs (IGF2 mRNA-binding proteins) is modulated by distinct RNA-binding domains. *Biol Chem*. 394:1077-1090.

Yamamoto, A., Y. Tagawa, T. Yoshimori, Y. Moriyama, R. Masaki, and Y. Tashiro. 1998. Bafilomycin A1 prevents maturation of autophagic vacuoles by inhibiting fusion between autophagosomes and lysosomes in rat hepatoma cell line, H-4-II-E cells. *Cell Struct Funct*. 23:33-42.

Yee, K.S., S. Wilkinson, J. James, K.M. Ryan, and K.H. Vousden. 2009. PUMA- and Bax-induced autophagy contributes to apoptosis. *Cell death and differentiation*. 16:1135-1145.

Yoshii, S.R., and N. Mizushima. 2017. Monitoring and Measuring Autophagy. *Int J Mol Sci*. 18:1865.

Yousefi, M., A. Nakauka-Ddamba, C.T. Berry, N. Li, J. Schoenberger, K.P. Simeonov, R.J. Cedeno, Z. Yu, and C.J. Lengner. 2018. Calorie Restriction Governs Intestinal Epithelial Regeneration through Cell-Autonomous Regulation of mTORC1 in Reserve Stem Cells. *Stem Cell Reports*. 10:703-711.

Zeng, W.J., C. Lu, Y. Shi, C. Wu, X. Chen, C. Li, and J. Yao. 2020. Initiation of stress granule assembly by rapid clustering of IGF2BP proteins upon osmotic shock. *Biochim Biophys Acta Mol Cell Res*. 1867:118795.

Figure Legends

Figure 1: IMP1 knockout is associated with increased expression of MAP1LC3 protein. A. Western blot of whole cell lysates from *IMP1-WT* or *IMP1-KO* SW480 cells treated with 20nM torin or 50 nm bafilomycin A1 (Baf) for 3 hours and probed with antibody specific to MAP1LC3B. For MAP1LC3B blots, two concentrations of lysates are presented to facilitate accurate quantification of LC3-II (10ug) and to visualize both LC3-I and LC3-II (20ug), respectively. B. Densitometry analysis of western blots represented in Figure 1A. LC3-II quantification normalized to GAPDH. Data are expressed as mean $\pm$ SEM, N = 4. ***p= 0.0002; ****p<0.0001. C. Western blot of lysates from *IMP1-WT* or *IMP1-KO* SW480 cells treated with bafilomycin A1 at indicated concentrations for 24 hours and probed with antibody that recognizes both MAP1LC3A and MAP1LC3B. D. Densitometry analysis of western blot in Figure 1C. LC3-II quantification normalized to GAPD. Data are presented as mean ± SEM. F. Representative immunofluorescent images of *IMP1-WT* and *IMP1-KO* SW480 cells stained with antibody for MAP1LC3A/B. Note increased MAP1LC3A/B in *IMP1-KO* compared to *IMP1-WT* cells. Images representative of n=3 fields of view for n = 2 independent experiments.

Figure 2: IMP1 phosphorylation (p-IMP1) contributes to regulation of MAP1LC3B expression. A. Schematic of IMP1 phosphorylation sites at serine-181 (S-181) and tyrosine-396 (Y-396) B. Western blot of whole cell lysates from HEK-293 *IMP1-KO* cells with stable expression of indicated GFP constructs: control-GFP (IMP1-KO), IMP1-WT-GFP, IMP1-S181A-GFP, IMP1-Y369F-GFP, IMP1-S181A/Y396F-GFP. Blots probed with IMP1, MAP1LC3B, and GAPDH antibodies. C. Densitometry analysis of western blots represented in Figure 2B. MAP1LC3B normalized to GAPDH. Data are expressed as mean $\pm$ SEM. N=3. **p=0.006. E. Representative flow cytometry plots of Autophagy Assay, Red fluorescence intensity in control (red) and chloroquine treated (3 hours, 100μm; blue) control-GFP (IMP1-KO), IMP1-WT-GFP, IMP1-S181A-GFP, IMP1-Y369F-GFP, and IMP1-S181A/Y396F-GFP cells normalized to mode. E. Quantification of fold change in median fluorescence intensity (MFI) with chloroquine treatment between all five cell lines listed in D. Data are expressed as mean ± SEM. N ≥ 3. *p=0.03.

Figure 3: IMP1 knockout is associated with increased MAP1LC3B-3'UTR translation A. Immunofluorescence of HEK-293 *IMP1-KO* cells with stable expression of indicated GFP constructs shown with IMP1-GFP (Green) and DAPI (blue) shown. Sample cell nuclei are indicated with white dotted lines. B. Schematic of dual-luciferase reporter assay with *MAP1LC3B*-3'UTR cloned into siCHECK2 plasmid. Plasmids were transiently transfected at equal concentrations into control-GFP (*IMP1-KO*), *IMP1-WT-GFP*, and *IMP1-S181A/Y396F-GFP* cells. B. Dual luciferase reporter assays were performed to compare relative luciferase activity of *MAP1LC3B*-3'UTR compared to empty control in each of the cell lines described in B. Renilla luciferase was normalized to built-in control firefly luciferase. Data are presented as mean ± SEM. N ≥ 3. *p=0.03.

Figure 4: IMP1 partially co-localizes with stress granule marker PABP in response to cell stress. Immunofluorescence of HEK-293 *IMP1-KO* cells with stable expression of *IMP1-WT-GFP* or *IMP1-S181A/Y396F-GFP* and treated with 100 uM of sodium arsenite or vehicle control (water) for 45 minutes A. Staining for GFP (green), EDC4 (red), and DAPI (blue) in *IMP1-WT-GFP*-expressing cells. White arrows in insets of vehicle control denote examples of GFP-EDC4 co-localization that is lost with arsenite treatment. B. GFP (green), EDC4 (red), and DAPI (blue) in *S181A/Y396F-GFP*-expressing cells. White arrows in vehicle control denote GFP-EDC4 co-localization that is decreased with arsenite treatment. Note EDC4 granules appear reduced in arsenite treated *IMP1- S181A/Y396F-GFP* compared to *IMP1-WT-GFP* cells. C. Staining for GFP (green), PABP (red), and DAPI (blue). White arrows in arsenite-treated insets denote overlap of GFP and PABP in *IMP1-WT-GFP*-expressing cells. D. GFP (green), PABP (red), and

DAPI (blue) staining in *S181A/Y396F-GFP*-expressing cells. Note an overall decrease in granule formation, but that GFP and PABP+ are still apparent (white arrows in insets). N ≥ 2 independent experiments and images are representative of 3 fields of view per experiment.

Figure 5: IMP1-GFP colocalization with MAP1LC3B transcript is reduced following acute cell stress.
A. Immunofluorescence of HEK-293 *IMP1-KO* cells transfected with IMP1-WT-GFP (green) and *MAP1LC3B* mRNA (red) using probes listed in Table 3. Cells were treated with 100 uM sodium arsenite or water control for 45 minutes. Nuclei are stained with DAPI (blue). White arrows in vehicle-treated cells indicate IMP1-*MAP1LC3B* mRNA co-localization. B. Immunofluorescence in IMP1-WT-GFP cells with *IMP1-WT-GFP* (green), *ATG5* mRNA (red). Note lack of GFP- ATG5 mRNA co-localization. C. Immunofluorescence of HEK-293 *IMP1-KO* cells transfected with IMP1-*S181A/Y396F-GFP* (green) and *MAP1LC3B* mRNA (red). Note a reduction in IMP1-*MAP1LC3B* mRNA co-localization compared to A (vehicle-treated). D. Immunofluorescence of cells described in C with IMP1-*S181A/Y396F-GFP* (green), *ATG5* mRNA (red). E. Correlation analysis of images in 5A & C (vehicle) using CellProfiler to quantify overlap of *MAP1LC3* mRNA and IMP1 fluorescence. Rank weight coefficient analysis was performed on n ≥ 5 high-powered fields across two independent experiments. Data expressed as SEM. **p=0.01. F. Distribution of *MAP1LC3B* within indicated cell lines from perinuclear (1) to cell periphery (4). N ≥ 4 images analysed across two independent experiments using CellProfiler with mean fraction distribution.

Figure 6: Model of IMP1 regulation of MAP1LC3B
Our data support a new mechanism for post-transcriptional regulation of *MAP1LC3B* by IMP1. IMP1 sequestration of *MAP1LC3B* transcripts in cytoplasmic granules may poise homeostatic cells to respond to stress by protecting *MAP1LC3B* mRNA in potentially diverse ribonucleoprotein (RNP) granules, including processing bodies. Upon stress, or with mutation of IMP1 phosphorylation sites (p-IMP1), IMP1-*MAP1LC3B* mRNA co-localization is reduced, where IMP1 is localized to stress granules and MAP1LC3B transcripts are observed further from the nucleus. Our observation that MAP1LC3B protein is increased with IMP1 deletion supports a model in which reduction of IMP1-*MAP1LC3B* mRNA co-localization permits increased translation of MAP1LC3B. Schematic created with BioRender (Biorender.com).

Supplemental Figure 1: A. Western blot of SW480 whole cell lysates from *IMP1-WT* or *IMP1-KO* cells treated with either 20 nM Torin1 or DMSO (control) for 24 hours and probed with antibody that recognizes both MAP1LC3A and MAP1LC3B, as well as other antibodies listed. B. Densitometry analysis of western blots represented in Supplemental Figure 1A. LC3-II was normalized to GAPDH. Data are expressed as mean ± SEM, N = 3. *p=.04. C. Real-time RT-PCR on total RNA from SW480 *IMP1-WT* or *IMP1-KO* cells treated with either 20 nM torin1 or vehicle (DMSO) for 24 hours. Gene expression for *MAP1LC3B* was normalized to GAPDH expression. Data are expressed as mean ± SEM versus *IMP1-WT*. N = 3.

Supplemental Figure 2: Schematic of Autophagy Probe, Red assay in Figure 2D, E.

Supplemental Figure 3: A. Western blot of lysates from fractionated HEK-293 *IMP1-KO* cells with stable expression of *IMP1-WT-GFP* or *IMP1-S181A-GFP* using the Subcellular Protein Fractionation Kit for Cultured Cells. Blots are probed with indicated antibodies to indicate nuclear and cytosolic fractions. Cytosolic Extraction (CEB), Membrane Extraction (MEB), Nuclear Extraction (NEB), Chromatin Bound Extraction (NEB-C), Cytoskeletal Extraction (PEB) lysates are represented. B. Densitometry of western blots indicate significantly more cytosolic than nuclear IMP1 in *IMP1-WT-GFP*, but not *IMP1-S181A-GFP* cells. Cytosolic included CEB and MEB fractions, nuclear included NEB and NEB-C fractions. Data are presented as mean ± SEM with p-values indicated. N = 3.

Movie S1. Time-lapse imaging of arsenite-treated IMP1-WT-GFP cells.
IMP1-WT-GFP cells exhibited robust cytoplasmic granule formation with arsenite treatment. Cells were treated with 100 uM sodium arsenite for 90 minutes. https://doi.org/10.6084/m9.figshare.16304799.

Movie S2. Time-lapse imaging of arsenite-treated IMP1-S181A-GFP cells.
IMP1-S181A-GFP cells exhibited reduced overall granule formation after arsenite treatment compared to *IMP1-WT-GFP* cells. Cells were treated with 100 uM sodium arsenite for 90 minutes. https://doi.org/10.6084/m9.figshare.16304802.

Movie S3. Time-lapse imaging of arsenite-treated IMP1-Y-396F-GFP cells.
IMP1-Y-396F-GFP cells exhibited reduced overall granule formation after arsenite treatment compared to *IMP1-WT-GFP* cells. Cells were treated with 100 uM sodium arsenite for 90 minutes.
https://doi.org/10.6084/m9.figshare.17162717.v1.

Table 1: Antibodies

Antibody	Manufacturer	Catalogue Number	Dilution (application, block)
IMP1	Cell Signaling	D33A2	1:1000 (western blotting, milk)
MAP1LC3A/B	Cell Signaling	12741	1:1000 (western blotting, milk), 1:100 (immunofluorescence)
MAP1LC3B (D11)	Cell Signaling	3868	1:1000 (western blotting, milk)
SQSTM1/p62	Abcam	Ab56416	1:1000 (western blotting, milk)
ATG5	Cell Signaling	12994S	1:1000 (western blotting, milk)
ATG3	Cell Signaling	3415T	1:1000 (western blotting, milk)
pS6RP (Ser235/236)	Cell Signaling	4858S	1:1000 (western blotting, BSA)
S6RP	Cell Signaling	2217S	1:1000 (western blotting, milk)
GAPDH	Santa Cruz	sc-32233	1:2000 (western blotting, milk)
HDAC2	Cell Signaling	2540S	1:1000 (western blotting, milk)
EDC4	Santa Cruz	Sc-374211	1:100 (immunofluorescence)
GFP	Novus Bio	NB100-1614	1:200 (immunofluorescence)
PABP	Santa Cruz	sc-32318	1:100 (immunofluorescence)

Table 2: qPCR assays

Gene target	Assay number
GAPDH	Hs02758991_g1
MAP1LC3B	Hs01076567_g1

Table 3 smFISH Probes

agctgcatttttctggtttt	MAP1LC3B_1
actgatcgatctcagttggt	MAP1LC3B_2
actactgtttcatgatctgt	MAP1LC3B_3
gctcagtttttctgcttgaa	MAP1LC3B_4
ccaaagctgaatgtgctcac	MAP1LC3B_5
gtctctctgaaagcctttaa	MAP1LC3B_6
acttctacccaaaaaggcgg	MAP1LC3B_7
gatctctctagctgagtaga	MAP1LC3B_8
cttcacaactcaggcctaaa	MAP1LC3B_9

aaatgcgtttttgcggggggtt	MAP1LC3B_10
cgtttgtgccaactgtgatg	MAP1LC3B_11
tttacagtcagggccgtttt	MAP1LC3B_12
taggagtcagggaccttcag	MAP1LC3B_13
tgaggactttgggtgtggtt	MAP1LC3B_14
gaaatgtctacccctgcaag	MAP1LC3B_15
tttctccagggtcttttaag	MAP1LC3B_16
ggcactagtttgaactcaga	MAP1LC3B_17
tgctgctttccgtaacaaca	MAP1LC3B_18
tgtactccagagtgacactg	MAP1LC3B_19
tactattttgtgtttctccc	MAP1LC3B_20
agggagtgtgtctgaatgtt	MAP1LC3B_21
tggatcatccacagctttaa	MAP1LC3B_22
gtgggtaccttttcagtaat	MAP1LC3B_23
ctgtctgacctaagagtgat	MAP1LC3B_24
tggtgtgggagaattctgat	MAP1LC3B_25
ttctccacaactgacatgca	MAP1LC3B_26
gtacggctcttttgctatg	MAP1LC3B_27
ctcttgacattagtatctgt	MAP1LC3B_28
atcacaggttgaacctgagg	MAP1LC3B_29

tactgggaagcactagtctt	MAP1LC3B_30
taaatagtgaaccccatgca	MAP1LC3B_31
cacatagggtctaaagtggt	MAP1LC3B_32
acaagggtaagttgtgacct	MAP1LC3B_33
caggtatttcatacacatct	MAP1LC3B_34
ttccccgttacagtaaacag	MAP1LC3B_35
cccagatgcaaagaatctgg	MAP1LC3B_36
tgacaggaactcctttaaca	MAP1LC3B_37
tctaaacgagacagtccagt	MAP1LC3B_38
ctgatcacacggtagttaca	MAP1LC3B_39
gaccactcacatgggatata	MAP1LC3B_40
gacagtttttaattgtgcct	MAP1LC3B_41
cagcgaatacaaccggcaac	ATG5_1
caaagccaaacctactccaa	ATG5_2
gtcattcttccaggagttaa	ATG5_3
tctcgaagcacatctttgtc	ATG5_4
tggaattcgtccaaaccaca	ATG5_5
ctttcagttatctcatcctg	ATG5_6
actttgtcagttaccaacgt	ATG5_7
tgtcttcttgtctcataacc	ATG5_8
ggataatgccatttcagtgg	ATG5_9
gtgatgttccaaggaagagc	ATG5_10
ttgcatccttagatggacag	ATG5_11
cagagttgcttgtgatcttt	ATG5_12
gcccaaaactggtcaaatct	ATG5_13
attccatgagtttccgattg	ATG5_14
aatccatttcttctgcagg	ATG5_15
ctttcagtcgttgtctgata	ATG5_16
ttctttgaggagatctccta	ATG5_17

agaggtgtttccaacattgg	ATG5_18
ggatgattctgttcaggcaa	ATG5_19
acatgctctgataaatccca	ATG5_20
aagtgtttctggtcaggttg	ATG5_21
gccttatctgacatggaatc	ATG5_22
gatgcaaagtaagaccagcc	ATG5_23
ggtttaatgatggcagtgga	ATG5_24
aagtctttcatgtcacagct	ATG5_25
ataccctgtttatttgtta	ATG5_26
aaagttcaaccatggtcacc	ATG5_27
ccaatgtttccactttcatg	ATG5_28
agtaatcctatgccctatat	ATG5_29
gcttggaaggaatgggtttt	ATG5_30
ttagacctggagccaatgaa	ATG5_31
ttcaaaagactggctgccag	ATG5_32
ctgcagtcaagagacaggta	ATG5_33
cccagaggcttggtaataaa	ATG5_34
gcctttttagcgtactcaaat	ATG5_35
accatgatcgtgtctgatac	ATG5_36
tatctggtctttgaccaact	ATG5_37
agattgtatgctgtactggc	ATG5_38
tggcactttcgtttacatca	ATG5_39
gtcagtggattttttatcct	ATG5_40
caatcatattcagaccagcc	ATG5_41
ccccaaaatgaaccgacgaa	ATG5_42
ctgatccttttgctcactaa	ATG5_43
cttcatcacagcttagtgtt	ATG5_44
tgacactggttttgtctgta	ATG5_45
cactcaagcctactgcaaag	ATG5_46
gacttacctgggattagagt	ATG5_47
atttctggttttactcttcc	ATG5_48

Figure 1

Figure 2

Figure 3

A

| IMP1-WT-GFP | IMP1-S181A-GFP | IMP1-Y396F-GFP | IMP1-S181A/Y396F-GFP |

B

atcgctcgagcccgggaattcgtttaaacctagagcggccgctggcc

C

- ● IMP1-KO
- ■ IMP1-WT
- ▲ IMP1-S181A/Y396F

Figure 4

Figure 5

A IMP1-WT-GFP
Vehicle — *MAP1LC3B* mRNA | Arsenite — *MAP1LC3B* mRNA
GFP | GFP
MERGE | MERGE
100 um

B IMP1-WT-GFP
Vehicle — *ATG5* mRNA | Arsenite — *ATG5* mRNA
GFP | GFP
MERGE | MERGE
100 um

C IMP1-S181A/Y396F-GFP
Vehicle — *MAP1LC3B* mRNA | Arsenite — *MAP1LC3B* mRNA
GFP | GFP
MERGE | MERGE
100 um

D IMP1-S181A/Y396F-GFP
Vehicle — *ATG5* mRNA | Arsenite — *ATG5* mRNA
GFP | GFP
MERGE | MERGE
100 um

E Co-localization between IMP1-GFP and *MAP1LC3B* mRNA

Rank weight coefficient

**

IMP1-WT-GFP | IMP1-S181A/Y396F-GFP

F *MAP1LC3B* mRNA distribution

Mean fraction distribution

Distance from the nucleus (perinuclear to periphery)

- IMP1-WT
- IMP1-S181/Y396F
- IMP1-WT + arsenite

Figure 6

Supplemental Figure 1

A

| IMP1 | + | + | – | – |
| Torin | – | + | – | + |

MAP1LC3A/B

IMP1

LC3-I
LC3-II

p62

ATG5

ATG3

pS6RP

S6RP

GAPDH

B   LC3-II protein

C   *MAP1LC3B* mRNA

IMP1-WT
IMP1-KO

Supplemental Figure 2

A

Supplemental Figure 3

A

B