A parametric logistic equation
with light flux and media concentration
for cultivation planning of microalgae

Kazuki Kambe[1†], Yasutaka Hirokawa[2†],
Asuka Koshi[1] and Yutaka Hori[1]

[1]Department of Applied Physics and Physico-Informatics,
Keio University, 3-14-1, Yokohama, Kanagawa, 223-8522 Japan.

[2]CMD Inc., 6-11-40, Soka, Saitama, 340-0002 Japan.

Abstract

Microalgae are considered to be promising producers of bioactive chemicals, feeds, and fuels from carbon dioxide by photosynthesis. Thus, the prediction of microalgal growth profiles is important for the planning of cost-effective and sustainable cultivation-harvest cycles. This paper proposes a mathematical model capable of predicting the effect of light flux into culture and media concentration on the growth profiles of microalgae by incorporating these growth-limiting factors into the logistic equation. The specific form of the equation is derived based on the experimentally measured growth profiles of *Monoraphidium* sp., a microalgal strain isolated by the authors. Using a cross-validation method, it is shown that the proposed model has the ability to predict necessary incident light flux into culture and initial media concentration for harvesting target biomass at target time. Finally, the model-guided cultivation planning is performed and is evaluated by comparing the result with the experimental data.

Keywords: logistic equation, growth profile, microalgae, cultivation planning, light flux, media concentration

---

† These authors contributed equally to this work.

1 Introduction

Photosynthesis is a beneficial reaction that reconverts atmospheric carbon dioxide produced by the consumption of fossil resources into organic carbon. It is one of the desirable solutions for a sustainable future to substitute photosynthetic products for fossil resources. Microalgae, which grow faster and show higher carbon fixation rates than higher plants, are excellent candidates for carbon neutral producer. Some species of microalgae have been used as live feed for fish larvae and are expected as alternative feedstocks for livestock [1] and aquaculture [2] because of their high nutritional quality. Other species of microalgae were reported as superior producers of bioactive chemicals [3] and biofuels [4, 5], which would potentially revolutionize the production of cosmetics, health foods, and fossil-based energy. Among them, *Monoraphidium* genera classified in Selenastraceae are oleaginous microalgae showing high lipid contents and were reported as promising hosts for lipid biofuel production [5–9]. Moreover, recent studies showed that *Monoraphidium* could grow robustly even in wastewater, suggesting that *Monoraphidium* cultivation would become a simultaneous solution for bioremediation and biorefinery [10, 11]. However, a major challenge in practical algal cultivation often lies in the discovery of cultivation conditions that enable continuously stable and cost-effective production over multiple cultivation-harvest cycles though the titer in a single harvest is often optimized. This motivates to develop a model-guided approach to the design of a cultivation plan for long-term cultivation. Specifically, development of a mathematical model capable of predicting the number of cells and harvest time in response to various cultivation conditions will be a key to finding the conditions for sustainable cultivation.

Many existing mathematical models capture the effect of various environmental factors on the dynamic change of the growth rate during the cultivation [12,13]. Examples include the Monod's growth model describing the effect of specific nutrients [14] and the models of temperature-dependent cell growth [15–17]. The effect of light intensity on the growth profile was also modeled in various ways depending on the situations of the cell culture [13]. Although the growth rate is affected by numerous factors including media concentration, light intensity, temperature, carbon dioxide concentration, and toxic byproducts in the media [18], these models consider the effect of only a few factors based on the assumption that cell growth is ultimately limited by those factors. In contrast to these models, the logistic equation incorporates the bulk effect of multiple growth limiting factors such as media concentration and toxic byproducts into a single parameter called the carrying

capacity of the environment [19]. Thus, the equation can be fitted to the growth profiles of a wide range of microorganisms [20–24]. However, since it does not explicitly contain the physical variables, the logistic equation cannot directly be used for the exploration of the growth conditions in the cultivation planning. To overcome the limitations of these models, a recent study proposed a hybrid Logistic-Monod model [25]. This model explicitly takes the media concentration into the variable while the other self-limiting factors are considered in the logistic-type equation, enabling the exploration of cultivation condition while maintaining the flexibility of the model. This idea can be used to build a more advanced hybrid model that includes other important factors for algal cultivation such as light intensity.

This paper proposes an ordinary differential equation (ODE) model of the growth profile of microalgae to enable cultivation planning by model-based exploration of the cultivation parameter space, as shown in Fig. 1 (A). The proposed model extends the logistic equation in a way that explicitly incorporates the dependence of the growth profile on light flux into culture and media concentration, enabling to find a cultivation condition that satisfies various constraints of practical cultivation process such as target biomass and target time. The specific form of the extended logistic equation and its parameters were determined from experimentally measured growth profiles of an originally isolated *Monoraphidium* sp. under different light fluxes into culture and media concentrations. These experimental data were further used for the cross-validation of the proposed model. The cross-validation result showed that the model could replicate various features of the growth profiles, including the peak biomass and its timing, for different cultivation conditions. Finally, we showcase a procedure of cultivation planning, where the initial media concentration and the incident light flux into culture are explored to achieve predefined target biomass and target time based on the simulations and some analytic relation of the proposed model.

2 Experimental conditions and overview of the proposed model

The goal of cultivation planning is to find parameters of cultivation such as the initial media concentration C_0 and the incident light flux into culture E_{in} that achieve a target biomass $N_\star$ at target time $T_\star$. To this end, we build a mathematical model that can

Figure 1: (A) Overview of cultivation planning. The symbols (C_0, E_{in}, N_{max}, $N_\star$ and $T_\star$) are the initial media concentration, the incident light flux into culture, the maximum biomass, the target biomass, and the target time, respectively. (B) Enlarged view of experimental setup. The symbol E_{out} is the transmitted light flux thorough culture.

predict the dynamic biomass N in response to various cultivation parameters as shown in Fig. 1 (A). In this section, we first introduce experimental conditions for building the proposed model, and then outline the overview for the proposed model.

2.1 Experimental conditions

Microalgae isolated from a freshwater pond in Yoshikawa, Saitama, Japan was named ACCB1808. The 18S ribosomal DNA sequence of ACCB1808 was 99.3 %, 99.1 %, and 99.1 % identical to that of *Monoraphidium* sp. LB59, *M. subclavatum* FBCC-A409, and *Monoraphidium* sp. NDMA-11, respectively. This result indicated that ACCB1808 belonged to the genus of *Monoraphidium*.

To measure the growth profile for modeling, ACCB1808 cultures were incubated in 300-mL flasks containing 200 mL of modified BG11 media. The details of experimental conditions including the media composition were summarized in Appendix A. Flasks containing culture were directly placed onto LED light. To strictly regulate the incident light flux into culture, black drawing paper with a 3-cm radius hole (area of 2.826×10^{-3} m^2) was inserted between the flasks and the LED light. The flasks were placed in an enclosure made with the same drawing paper as shown in Fig. 1 (B). The incident

light intensities to flasks measured as photosynthetic photon flux density (PPFD) were adjusted to 1034, 386.7, 184.8, and 96.8 $\mu E \cdot m^{-2} \cdot s^{-1}$. Owing to strict regulation of the illuminated area $(2.826 \times 10^{-3} \ m^2)$, the incident light fluxes into culture E_{in} were calculated as 2.92, 1.09, 0.521, and 0.274 $\mu E \cdot s^{-1}$. The concentration of modified BG11 media C was defined as 1, and the initial media concentrations C_0 were adjusted to 0.5, 0.25, and 0.125 by dilution. ACCB1808 cultures were incubated under 16 conditions consisting of the combinations of four patterns of initial media concentration C_0 and four patterns of incident light fluxes into culture E_{in} as shown in Fig. 2 (A). These experimental data were used for building and evaluating the proposed model in the following sections.

2.2　Overview of the proposed model

The growth profiles of experimental data shown in Fig. 2 (B) indicated that the growth rate and the biomass accumulation were dependent on the incident light flux into culture E_{in} and the initial media concentration C_0, respectively. In initial growth phase, the biomass N increased exponentially. After following linear increase, the growth rate gradually decreased and leveled off. The state of culture used for subculture often affected the initial growth rate in the next cultivation. Specifically, the culture state before stationary phase was preferable for use in subculture. To harvest the culture in the preferable state, it is important to predict not only a target biomass $N_\star$ but also the target time $T_\star$ at which the biomass reaches $N_\star$. Moreover, for practical operation, the target values ($N_\star$ and $T_\star$) have to be decided from various viewpoints such as cost effectiveness and the schedule of operators. A mathematical model would be a desirable tool for exploring cultivation conditions satisfying these target values.

In what follows, we build a mathematical model capable of predicting the growth kinetics of ACCB1808 culture for various cultivation conditions. In particular, we are interested in the growth kinetics in response to light flux into culture E_{in} and media concentration C since these parameters largely affect the growth profile as shown in Fig. 2 (B). Since the increase of biomass decreases available light flux for a single cell even if light flux into culture E_{in} is constant during cultivation, available light flux for a single cell is one of the key variables for the prediction of the growth kinetics. In this paper, light flux being available for a single cell is called light flux per cell L. The effect of light flux per cell on the growth kinetics will be discussed in Section 3.

We develop an ordinary differential equation (ODE) model of the growth kinetics based

on the logistic equation [19]:

$$\frac{\mathrm{d}N(t)}{\mathrm{d}t} = r\left(1 - \frac{N(t)}{\Gamma}\right)N(t), \tag{1}$$

where $N(t)$ is the biomass at time t, r is the maximum specific growth rate, and Γ is the carrying capacity of the environment, or the maximum achievable population size. The first term of the logistic equation represents the growth due to proliferation, and the second term collectively accounts for growth-limiting factors such as toxic byproducts and reactive oxygen.

Unlike the original logistic equation [19], we assume that the maximum specific growth rate r is dependent on the light flux per cell L, and the media concentration C, that is, $r := r(L, C)$. As discussed later in detail, L and C are subject to dynamic change since these two variables are affected by biomass $N(t)$. The combined effect of these factors is incorporated into the maximum specific growth rate $r(L, C)$ by

$$r(L, C) := \mu r_{\text{light}}(L) r_{\text{media}}(C), \tag{2}$$

where μ is a constant, and $r_{\text{light}}(L)$ and $r_{\text{media}}(C)$ are functions of light flux per cell and media concentration. These functions take values between 0 and 1. The specific forms of these functions are defined in Section 3 and 4. The functions $r_{\text{light}}(L)$ and $r_{\text{media}}(C)$ represent the growth-limiting effect due to insufficient absorbance of light flux per cell and insufficient nutrients, respectively. In other words, $r(L, C) \simeq \mu$ when the light flux per cell L and the media concentration C are sufficiently high such that $r_{\text{light}}(L) \simeq 1$, and $r_{\text{media}}(C) \simeq 1$, while $r(L, C) \simeq 0$ when L or C is close to 0.

3   Logistic equation with light flux per cell

In the previous section, the maximum specific growth rate r is defined by the functions of the media concentration C and of the light flux per cell L. When the media concentration C is sufficiently high, the maximum specific growth rate r is only dependent on the light flux per cell L. In this section, we consider this special case and formulate a logistic equation that incorporates the effect of the light flux per cell L, which we call the media-rich model or the MR model for shorthand.

3.1 Modeling of the MR model

We incorporate the effect of the light flux per cell L on the growth kinetics by defining

$$r_{\text{light}}(L) := \frac{L(N(t), E_{\text{in}})}{\lambda_{\text{L}} + L(N(t), E_{\text{in}})},\tag{3}$$

where λ_{L} is a half-velocity constant satisfying $r_{\text{light}}(\lambda_{\text{L}}) = 1/2$. The light flux per cell $L(N(t), E_{\text{in}})$ is dependent on the incident light flux into culture E_{in} and the biomass $N(t)$. The light flux absorbed by the culture is expressed by $E_{\text{in}} - E_{\text{out}}$ where E_{out} is the transmitted light flux through culture as illustrated in Fig. 1 (B) (see details about E_{out} in Appendix B). Thus, the absorbed light flux per cell $L(N(t), E_{\text{in}})$ is written as $(E_{\text{in}} - E_{\text{out}})/N(t)$ by assuming that light flux is uniformly absorbed by cells in culture. Moreover, the attenuation of transmitted light in the culture obeys Lambert-Beer's law [26], which states the exponential decrease of light flux with the path length of light flux and the concentration of the solution (biomass concentration) (see details in Appendix C). Thus, the absorbed light flux per cell $L(N(t), E_{\text{in}})$ is

$$L(N(t), E_{\text{in}}) = \frac{E_{\text{in}} - E_{\text{out}}}{N(t)} = \frac{1 - 10^{-K \cdot N(t)/V \cdot D}}{N(t)} E_{\text{in}},\tag{4}$$

where K is the cell-specific extinction coefficient, D is the culture depth, and V is the culture volume as illustrated in Fig. 1 (B). When the nutrients in the media are sufficient and are not rate-limiting factors, $r_{\text{media}}(C) = 1$ holds. Thus, it follows that

$$\frac{\text{d}N(t)}{\text{d}t} = \mu \frac{L(N(t), E_{\text{in}})}{\lambda_{\text{L}} + L(N(t), E_{\text{in}})} \left(1 - \frac{N(t)}{\Gamma}\right) N(t)\tag{5}$$

by substituting Eq. (3) into Eq. (1). In what follows, Eq. (5) is called the MR model. The MR model allows us to identify some of the parameters of Eqs. (1) and (2) by using experimental data taken under the conditions with sufficient media concentration as shown in the next subsection. The other parameters that appear in $r_{\text{media}}(C)$ can then be identified by subsequent experiments, which will be discussed in Section 4. The two-step parameter identification helps avoid overfitting to a single experimental data.

3.2 Parameter identification of the MR model

Experiments were conducted to identify the parameters K, μ, λ_{L}, and Γ of the MR model (Eq. (5)). Firstly, the extinction coefficient K being specific to ACCB1808 was identified

Table 1: Initial values and identified extinction coefficient K.

Parameter	Unit	Value
V	mL	200
D	cm	3.7
OD_0	-	0.025
K	$\mathrm{mL \cdot cm^{-1} \cdot cell^{-1}}$	-5.1×10^{-9}

by the method of Masuda *et al.* [26]. PPFD was measured in ACCB1808 culture with various biomass concentrations and culture depths (see details in Appendix C). The relative logarithmic PPFD was negatively correlated with biomass concentrations and culture depths, indicating that Lambert-Beer's law was obeyed in ACCB1808 culture. When the units of culture depth and biomass concentration were defined as "cm" and "cell $\cdot$ mL^{-1}", the extinction coefficient K was identified as $K = -5.1 \times 10^{-9}$ mL $\cdot$ cm^{-1} $\cdot$ cell^{-1}.

Using this extinction coefficient, we further performed identification of μ, λ_{L} and Γ in the MR model (Eq. (5)) based on the growth profile of ACCB1808 culture (see details about cultivation conditions in Appendix A). Experiments were conducted under a total of 16 different conditions as shown in Fig. 2 (A), where the growth curves were obtained for the combinations of four different light fluxes into culture, 0.274, 0.521, 1.09 and 2.92 μE $\cdot$ s^{-1}, and four initial media concentrations, 0.125, 0.25, 0.5 and 1. Optical density at 730 nm (OD) was measured as a proxy of biomass $N(t)$ (see details in Appendix C). The parameters were then fitted to the four growth kinetics with the sufficient media concentration, *i.e.,* $C_0 = 1$ in Fig. 2 (C). Specifically, the culture volume V, the culture depth D, the initial optical density OD_0, and the identified extinction coefficient K were set in Eq. (5) as shown in Table 1. When 200 mL culture was put into 300-mL flask, culture depth corresponded 3.7 cm. The culture depth D and the culture volume V were assumed to remain constant during cultivation. For $E_{\mathrm{in}} = 1.09$ and 2.92 μE $\cdot$ s^{-1}, time series data were fitted only for the first 690 and 306 hour, respectively, since the media concentration could become a rate-limiting factor, violating the assumption of the MR model, when the culture reached stationary phase.

The identified parameters are shown in Table 2. This result implied that the maximum specific growth rate $r(L, C) \simeq \mu r_{\mathrm{light}}(L)$ was between 0 and 0.194 hour^{-1} depending on the light flux per cell L when the media concentration was high, *i.e.,* $r_{\mathrm{media}}(C) \simeq 1$. The carrying capacity Γ was identified as 99.9×10^9 cell. The biomass concentration $99.9 \times 10^9 / 200$ cell $\cdot$ mL^{-1} can be converted into OD of 16.6. This indicated that the value

Table 2: List of identified parameters of the proposed model.

Parameter	Unit	Identified value
μ	hour^{-1}	0.194
λ_{L}	$\mu\mathrm{E} \cdot \mathrm{s}^{-1} \cdot \mathrm{cell}^{-1}$	1.90×10^{-6}
Γ	cell	99.9×10^{9}
ξ_{C}	-	0.012
α	cell^{-1}	8.7×10^{-12}

of OD would never exceed 16.6 regardless of the media concentration.

To evaluate the predictive ability of the MR model, the predicted results of the MR model were further compared with the experimental data using leave-one-out cross-validation (LOOCV), where three of the four experimental conditions were grouped together for parameter identification and the other was used for prediction. Specifically, the parameters (μ, λ_{L} and Γ) were fitted to the three growth curves with $C_0 = 1$. Then, the identified model was used to predict the growth kinetics as shown in Fig. 2 (C). Figure 2 (C) shows that the MR model was capable of predicting the growth kinetics before reaching stationary phase, where the decrease in media concentration had little effect on cell growth.

In the next section, the MR model is used to build a full model that captures the effect of both light flux per cell L and media concentration C on cell growth.

4 Logistic equation with light flux per cell and media concentration

A standing assumption of the MR model is that the media concentration $C(t)$ is sufficiently high so that the maximum specific growth rate r is independent of $C(t)$. In this section, we will extend the MR model (Eq. (5)) to remove this assumption and explicitly incorporate the effect of the media concentration on the maximum specific growth rate.

4.1 Modeling of logistic equation with light flux and media concentration

The experimental data in Fig. 2 (B) suggest that the media concentration $C(t)$ does not affect the growth profile just before the stationary phase is reached. Based on this observation, we incorporate the rate-limiting effect of the media concentration using the

Figure 2: (A) Experimental conditions combining initial media concentrations C_0 and light fluxes into culture E_{in}. (B) Time series data for different initial media concentrations C_0 and incident light flux into culture E_{in}. Circles and solid lines show experimental data and predicted results of the full model described in Section 4, respectively. (C) Time series data for the four incident light fluxes into culture E_{in} with $C_0 = 1$. Circles and solid lines show experimental data (repeat of the data in panel (B)) and predicted results of the MR model described in Section 3, respectively.

Monod-type model [14]:

$$r_{\text{media}}(C(t)) := \frac{C(t)}{\xi_{\text{C}} + C(t)},$$

where ξ_{C} is a half-velocity constant. We assume that nutrients in the media are mainly used for cell growth, and the consumption rate is proportional to the growth rate $\mathrm{d}N(t)/\mathrm{d}t$. Consequently, the mathematical model that incorporates the effect of both the light flux per cell $L(N(t), E_{\text{in}})$ and the media concentration $C(t)$ is obtained as

$$\begin{cases} \dfrac{\mathrm{d}N(t)}{\mathrm{d}t} = \mu \dfrac{C(t)}{\xi_{\text{C}} + C(t)} \dfrac{L(N(t), E_{\text{in}})}{\lambda_{\text{L}} + L(N(t), E_{\text{in}})} \left(1 - \dfrac{N(t)}{\Gamma}\right) N(t) & \text{(6a)} \\ \dfrac{\mathrm{d}C(t)}{\mathrm{d}t} = -\alpha \dfrac{\mathrm{d}N(t)}{\mathrm{d}t}, & \text{(6b)} \end{cases}$$

where α is a parameter representing consumption of the media concentration per a unit increase of the biomass, and $L(N(t), E_{\text{in}})$ is defined by Eq. (4). Eq. (6) is called the full model in the following sections. Defining the initial biomass $N(0)$ by N_0, Eq. (6) can equivalently be expressed as

$$\frac{\mathrm{d}N(t)}{\mathrm{d}t} = \mu \frac{C_0 - \alpha(N(t) - N_0)}{\xi_{\text{C}} + C_0 - \alpha(N(t) - N_0)} \cdot$$
$$\frac{1 - 10^{-K \cdot N(t)/V \cdot D}}{\lambda_{\text{L}} \cdot E_{\text{in}}^{-1} \cdot N(t) + 1 - 10^{-K \cdot N(t)/V \cdot D}} \left(1 - \frac{N(t)}{\Gamma}\right) N(t)$$

since Eq. (6b) implies $C(t) = C_0 - \alpha(N(t) - N_0)$.

It should be noted that the parameters K, μ, λ_{L}, and Γ are identified with the MR model as described in Section 3.2. The other parameters ξ_{C} and α can be identified using the 16 experimental data for the different incident light fluxes into culture E_{in} and the initial media concentrations C_0 in Fig. 2 (B).

4.2   Parameter identification of the full model

The parameters ξ_{C} and α in the full model (Eq. (6)) were identified using the 16 experimental data in Fig. 2 (B). Specifically, the initial parameters, V, D, and OD_0, and the identified extinction coefficient K in Table 1 were used. The parameters μ, λ_{L}, and Γ obtained using the MR model in Section 3.2 were set (see Table 2). Then, the least square solution was searched for ξ_{C} and α. The identified parameters are shown in Table 2.

The generalizability of the full model was also evaluated by LOOCV using the experimental conditions matrix in Fig. 2 (A). Specifically, for each combination of the light flux into culture E_{in} and the initial media concentration C_0, the parameters ξ_{C} and α were identified with the other 15 experimental data. Then, the identified model was used to predict the growth kinetics as shown in Fig. 2 (B). The simulated growth kinetics showed agreement with the experimental data in that the dynamics of the growth rate was dependent on the light flux into culture E_{in} before reaching stationary phase, while the maximum biomass was dependent on the initial media concentration C_0. The result of LOOCV in Fig. 2 (B) also suggests that the model can predict the maximum biomass N_{max} or its corresponding OD OD_{max}. These results will be more quantitatively evaluated in the next section along with the demonstration of cultivation planning.

5 Demonstration of cultivation planning

The goal of cultivation planning is to find the initial media concentration $C_0^{\star}$ and the incident light flux into culture $E_{\text{in}}^{\star}$ to achieve a predefined target biomass $N_{\star}$ at target time $T_{\star}$. In a typical cultivation process, cells are harvested before reaching the stationary phase to avoid the carry-over of potentially toxic byproducts in subculture. Thus, $C_0^{\star}$ and $E_{\text{in}}^{\star}$ should be planned so that biomass reaches $N_{\star}$ at $T_{\star}$.

5.1 Prediction of initial media concentration $C_0^{\star}$ for target biomass $N_{\star}$

In a typical cultivation cycle, culture is harvested before reaching stationary phase to maintain a preferable culture state in subculture. In other words, target biomass $N_{\star}$ should be set less than the maximum biomass at stationary phase N_{max}, $e.g.$, $N_{\star} = 0.8 N_{\text{max}}$. Thus, prediction of the maximum biomass N_{max} or its corresponding OD OD_{max} in response to cultivation conditions such as incident light flux into culture E_{in} and initial media concentration C_0 is important in cultivation planning. In theory, N_{max}, or its corresponding OD OD_{max}, can be analytically calculated from Eq. (6) as

$$\text{OD}_{\text{max}} \propto N_{\text{max}} = \min \left(\frac{C_0}{\alpha} + N_0, \ \Gamma \right). \tag{7}$$

Eq. (7) implies that the maximum biomass N_{max} is obtained from an initial media concentration C_0 when $C_0 < \alpha(\Gamma - N_0)$. This analytic solution provides a crude estimation of biomass at stationary phase for a given initial media concentration.

Figure 3: OD_{max} calculated from the analytic solution (Eq. (7)) and measured by experiments in Fig. 2 (B).

Figure 3 shows the maximum OD OD_{max} predicted from Eq. (7) and measured by the experiments in Fig. 2 (B), where the parameters α and Γ in Table 2 and the initial OD_0 in Table 1 corresponding the initial biomass N_0 were used for calculation. The results show that OD_{max} is determined from the initial media concentrations C_0 and is independent of the incident light fluxes into culture E_{in}. Thus, a predefined target biomass $N_\star$ or its corresponding target OD for cultivation planning is obtained by simply selecting the initial media concentration $C_0^\star$, which can be calculated from Eq. (7). Once the initial media concentration $C_0^\star$ is fixed, the time when biomass reaches $N_\star$ can be adjusted by incident light flux into culture E_{in}. In the next subsection, we will give a demonstration to select the incident light flux into culture $E_{in}^\star$ for target time $T_\star$.

5.2 Prediction of incident light flux into culture $E_{in}^\star$ for target time $T_\star$

Once the initial media concentration $C_0^\star$ is selected based on Eq. (7), the next goal is to seek the incident light flux into culture $E_{in}^\star$ that achieves target biomass $N_\star$ at target time $T_\star$. The incident light flux into culture $E_{in}^\star$ is explored by running simulations of the full model (Eq. (6)). Since the target biomass $N_\star$ is often set less than the maximum biomass at stationary phase in practical cultivation, let us suppose, for example, that the target biomass $N_\star$ is around 70–90% of the maximum biomass. Then, the expected harvest time, at which the biomass reaches the target biomass can be computed from the growth kinetics

Figure 4: Span of target time obtained by simulations of the full model (Eq. (6)) and measured by experiments in Fig. 2 (B).

simulated by Eq. (6) for each E_{in}.

Figure 4 shows the time spans in which the biomass reaches 70–90% of its maximum value for different incident light fluxes into culture E_{in}. In Fig. 4, T_1 and T_2 represent the simulated lower and upper bound of harvest time, at which biomass reaches $0.7N_{max}$ and $0.9N_{max}$, respectively. The parameters in Tables 1 and 2 were used for these simulations of the full model (Eq. (6)). The experimental data in Fig. 4 is obtained from the time-series data in Fig. 2 (B).

Figure 4 shows agreement of the computationally predicted time spans with the experimental data. Thus, the incident light flux into culture $E_{in}^{\star}$ that enables to harvest target biomass at target time can be computationally determined based on the full model (Eq.

(6)). Thus, combining with the prediction of the initial media concentration $C_0^\star$ in Section 5.1, one can find cultivation conditions ($C_0^\star$ and $E_{\mathrm{in}}^\star$) satisfying the predefined constraints ($N_\star$ and $T_\star$), fulfilling the goal of cultivation planning.

6   Conclusion

Microalgae are photosynthetic organisms that have high potential as carbon neutral producer and alternative feedstocks for livestock [1] and aquaculture [2]. Prediction of biomass in cultivation of microalgae is a difficult task due to the complex interplay of growth conditions such as light flux, media concentration, and temperature [18]. As a result, experimental conditions for harvesting target biomass at target time, *i.e.,* cultivation plans, are often sought empirically by operators.

This paper has proposed an ODE model for predicting the growth profile of microalgae in response to media concentration and light flux into culture to help the operators with cultivation planning. The proposed model has been built by extending the logistic equation in two steps. We have firstly constructed the MR model (Eq. (5)) that considers only the effect of light flux into culture, assuming that the media concentration is high. Then, we have extended the MR model to incorporate the effect of the media concentration (Eq. (6)). The predictive ability of the proposed model has been evaluated by comparing the predicted growth kinetics with the experimentally obtained growth profile of ACCB1808 (*Monoraphidium* sp.). Finally, model-guided cultivation planning has been shown as a demonstration example, where the initial media concentration and the incident light flux into culture have been planned for harvesting predefined target biomass at target time. The proposed model streamlines the planning process of cultivation cycles that satisfy various practical demands such as cost effectiveness and the schedule of operators.

Data, codes and materials

The program codes used in this study are available at GitHub (`https://github.com/hori-group/logistic_eq_for_cultivation_planning`).

Competing interests

Keio University received a research fund from CMD Inc. for conducting this work. Y. Hirokawa is an employee of CMD Inc..

Authors' contributions

Y. Hirokawa and Y. Hori conceived the study. K.K, A.K, and Y. Hori developed the mathematical model, computational methods, and program codes. Y. Hirokawa performed cultivation experiments and data analysis. K.K., Y. Hirokawa, and Y. Hori drafted the manuscript. All authors gave feedback to the manuscript and gave final approval for publication.

References

[1] Saadaoui I, Rasheed R, Aguilar A, Cherif M, Al Jabri H, Sayadi S, et al. Microalgal-based feed: promising alternative feedstocks for livestock and poultry production. Journal of Animal Science and Biotechnology. 2021;12(1):1-15. doi: 10.1186/s40104-021-00593-z.

[2] Nagappan S, Das P, AbdulQuadir M, Thaher M, Khan S, Mahata C, et al. Potential of microalgae as a sustainable feed ingredient for aquaculture. Journal of Biotechnology. 2021;341:1-20. doi: 10.1016/j.jbiotec.2021.09.003.

[3] Ren Y, Sun H, Deng J, Huang J, Chen F. Carotenoid production from microalgae: biosynthesis, salinity responses and novel biotechnologies. Marine Drugs. 2021;19(12):713. doi: 10.3390/md19120713.

[4] Patel A, Karageorgou D, Rova E, Katapodis P, Rova U, Christakopoulos P, et al. An overview of potential oleaginous microorganisms and their role in biodiesel and omega-3 fatty acid-based industries. Microorganisms. 2020;8(3):434. doi: 10.3390/microorganisms8030434.

[5] Yee W. Microalgae from the Selenastraceae as emerging candidates for biodiesel production: a mini review. World Journal of Microbiology and Biotechnology. 2016;32(4):1-11. doi: 10.1007/s11274-016-2023-6.

[6] Yu X, Zhao P, He C, Li J, Tang X, Zhou J, et al. Isolation of a novel strain of Monoraphidium sp. and characterization of its potential application as biodiesel feedstock. Bioresource Technology. 2012;121:256-62. doi: 10.1016/j.biortech.2012.07.002.

[7] Zhao Y, Li D, Xu JW, Zhao P, Li T, Ma H, et al. Melatonin enhances lipid production in Monoraphidium sp. QLY-1 under nitrogen deficiency conditions via a multi-level mechanism. Bioresource Technology. 2018;259:46-53. doi: 10.1016/j.biortech.2018.03.014.

[8] Song X, Zhao Y, Han B, Li T, Zhao P, Xu JW, et al. Strigolactone mediates jasmonic acid-induced lipid production in microalga Monoraphidium sp. QLY-1 under nitrogen deficiency conditions. Bioresource Technology. 2020;306:123107. doi: 10.1016/j.biortech.2020.123107.

[9] Li X, Zhang X, Zhao Y, Yu X. Cross-talk between gama-aminobutyric acid and calcium ion regulates lipid biosynthesis in Monoraphidium sp. QLY-1 in response to combined treatment of fulvic acid and salinity stress. Bioresource Technology. 2020;315:123833. doi: 10.1016/j.biortech.2020.123833.

[10] Tale M, Ghosh S, Kapadnis B, Kale S. Isolation and characterization of microalgae for biodiesel production from Nisargruna biogas plant effluent. Bioresource technology. 2014;169:328-35. doi: 10.1016/j.biortech.2014.06.017.

[11] Mishra S, Mohanty K. Comprehensive characterization of microalgal isolates and lipid-extracted biomass as zero-waste bioenergy feedstock: an integrated bioremediation and biorefinery approach. Bioresource technology. 2019;273:177-84. doi: 10.1016/j.biortech.2018.11.012.

[12] Bekirogullari M, Figueroa-Torres GM, Pittman JK, Theodoropoulos C. Models of microalgal cultivation for added-value products – a review. Biotechnology Advances. 2020;44:107609. doi: 10.1016/j.biotechadv.2020.107609.

[13] Béchet Q, Shilton A, Guieysse B. Modeling the effects of light and temperature on algae growth: state of the art and critical assessment for productivity prediction during outdoor cultivation. Biotechnology Advances. 2013;31(8):1648-63. doi: 10.1016/j.biotechadv.2013.08.014.

[14] Monod J. The growth of bacterial cultures. Annual review of microbiology. 1949;3(1):371-94. doi: 10.1146/annurev.mi.03.100149.002103.

[15] Ratkowsky DA, Olley J, McMeekin T, Ball A. Relationship between temperature and growth rate of bacterial cultures. Journal of bacteriology. 1982;149(1):1-5. doi: 10.1128/jb.149.1.1-5.1982.

[16] Ratkowsky D, Lowry R, McMeekin T, Stokes A, Chandler R. Model for bacterial culture growth rate throughout the entire biokinetic temperature range. Journal of bacteriology. 1983;154(3):1222-6. doi: 10.1128/jb.154.3.1222-1226.1983.

[17] Bernard O, Rémond B. Validation of a simple model accounting for light and temperature effect on microalgal growth. Bioresource technology. 2012;123:520-7. doi: 10.1016/j.biortech.2012.07.022.

[18] Morales M, Sánchez L, Revah S. The impact of environmental factors on carbon dioxide fixation by microalgae. FEMS microbiology letters. 2018;365(3):fnx262. doi: 10.1093/femsle/fnx262.

[19] Krebs CJ. Ecology: The experimental analysis of distribution and abundance. Pearson Education UK; 2013.

[20] Fujikawa H, Kai A, Morozumi S. A new logistic model for bacterial growth. Food Hygiene and Safety Science. 2003;44(3):155-60. doi: 10.3358/shokueishi.44.155.

[21] Fujikawa H, Kai A, Morozumi S. Improvement of new logistic model for bacterial growth. Food Hygiene and Safety Science. 2004;45(5):250-4. doi: 10.3358/shokueishi.45.250.

[22] Yang Z, Zhao Y, Liu Z, Liu C, Hu Z, Hou Y. A mathematical model of neutral lipid content in terms of initial nitrogen concentration and validation in Coelastrum sp. ha-1 and application in Chlorella sorokiniana. BioMed Research International. 2017;2017. doi: 10.1155/2017/9253020.

[23] Ram Y, Dellus-Gur E, Bibi M, Karkare K, Obolski U, Feldman MW, et al. Predicting microbial growth in a mixed culture from growth curve data. Proceedings of the National Academy of Sciences of the United States of America. 2019;116(29):14698-707. doi: 10.1073/pnas.1902217116.

[24] Peleg M, Corradini MG, Normand MD. The logistic (Verhulst) model for sigmoid microbial growth curves revisited. Food research international. 2007;40(7):808-18. doi: 10.1016/j.foodres.2007.01.012.

[25] Xu P. Analytical solution for a hybrid Logistic-Monod cell growth model in batch and continuous stirred tank reactor culture. Biotechnology and bioengineering. 2020;117(3):873-8. doi: 10.1002/bit.27230.

[26] Masuda A, Horaguchi K, Kosaka S, Ozawa T, Kato M, Murakami K. The reduction of luminous transmittance in a microalgal culture tank with the increase of microalgal density and culture liquid thickness. Eco-engineering. 2006;18(1):3-8. doi: 10.11450/seitaikogaku.18.3 (in Japanese).

A Experimental conditions for cultivation

All ACCB1808 cultures were incubated in a space of which room temperature was controlled at 25°C with continuous aeration including 1.5 % carbon dioxide. LED light was used for growth, and light intensities were altered for incubation conditions. Photosynthetic photon flux density (PPFD) in each condition was measured by using Spectromaster C-7000 (SEKONIC; Tokyo, Japan). Optical density at 730 nm (OD), which is proportional to biomass concentration N/V where N is the biomass and V is the culture volume, was measured by using Spectrophotometer V-700 (JASCO; Tokyo, Japan). Biomass concentration N/V was counted by using microscope (Olympus; Tokyo, Japan) and a Thoma chamber.

For isolation and preculture, liquid and solid BG11 media containing following components (per liter); 1.5 g of $NaNO_3$, 31.4 mg of K_2HPO_4, 73.9 mg of $MgSO_4 \cdot 7H_2O$, 36.8 mg of $CaCl_2 \cdot 2H_2O$, 20.1 mg of Na_2CO_3, 1.12 mg of EDTA 2Na, 6.09 mg of citrate, 10.15 mg of ferric ammonium citrate, 1 mL of A6 solution, was used. A6 solution included following components (per liter); 2.86 g of H_3BO_4, 1.81 g of $MnCl_2 \cdot 4H_2O$, 0.22 g of $ZnSO_4 \cdot 7H_2O$, 0.39 g of $Na_2MoO_4 \cdot 2H_2O$, 0.079 g of $CuSO_4 \cdot 5H_2O$, 0.049 g of $Co(NO_3)_2 \cdot 6H_2O$. To prepare solid media, 15 $g \cdot L^{-1}$ agar was supplemented.

The growth profiles of ACCB1808 in different concentrations (from 73.9 to 739 $mg \cdot L^{-1}$) of $MgSO_4 \cdot 7H_2O$ in BG11 media indicated that 6-fold concentration (443 $mg \cdot L^{-1}$) was appropriate (data not shown). Therefore, $MgSO_4$ fortified media was called modified BG11 media and used for main culture cultivation.

In preculture cultivation, cells were inoculated into 60 mL of BG11 media in a 100-mL test tube and incubated under continuous illumination (40 $\mu E \cdot m^{-2} \cdot s^{-1}$) and aeration including 1.5 % carbon dioxide. Cells of preculture in late logarithmic growth phase were

inoculated into 200 mL modified BG11 media in 300-mL flask to an initial optical density at 730 nm (OD_0) of 0.025.

B   Relation between incident light flux and its density

The incident light flux into culture E_{in} can be calculated by multiplication of incident light flux density e_{in} and illuminated area (2.826×10^{-3} m^2):

$$E_{in} = e_{in} \times 2.826 \times 10^{-3}.$$

Assuming that the incident light linearly passes culture, the transmitted light flux through culture E_{out} is also calculated by

$$E_{out} = e_{out} \times 2.826 \times 10^{-3}.$$

Based on Lambert-Beer's law as mentioned in Appendix C, the light flux per cell $L(N(t), E_{in})$ is obtained as Eq. (4) in Section 3.1.

C   Identification of extinction coefficient K based on Lambert-Beer's law

The measured photosynthetic photon flux density (PPFD) under conditions with different culture depths D and optical densities at 730 nm (ODs) are summarized in Table 3. It should be noted that the unit of culture depth D is centimeter (cm) and that OD is proportional to the biomass concentration N/V, where N is the biomass and V is the culture volume. In Table 3, e_{in} and e_{out} correspond to PPFD in culture solution without cells and in culture with various ODs (OD=0.149–4.78), respectively.

Optical density (OD) can be converted to biomass concentration N/V by

$$\frac{N}{V} = OD \times 30.1 \times 10^6. \tag{8}$$

Thus, OD of 0.149, 0.310, 0.600, 1.17, 2.47 and 4.78 in Table 3 correspond to 4.55×10^6, 10.1×10^6, 17.6×10^6, 34.9×10^6, 69.8×10^6 and 144×10^6 cell$\cdot$mL^{-1}, respectively.

Figures 5 (A) and 5 (B) show relative logarithmic PPFD ($= \log(e_{out}/e_{in})$) for various culture depths D at particular OD (OD=0.149–4.78) and for various ODs at particu-

Table 3: PPFD in different depth D and OD.

| | PPFD ($\mu E \cdot s^{-1} \cdot m^{-2}$) | | | | | | |
| | e_{in} | e_{out} | | | | | |
OD $\diagdown$ D (cm)	0	0.149	0.310	0.600	1.17	2.47	4.78
0	834.3	829.0	834.0	824.5	818.0	842.0	839.5
1	419.3	386.5	365.0	300.5	226.5	155.5	69.0
2	203.3	175.0	157.0	120.0	77.0	40.0	9.5
3	114.3	95.0	82.0	58.0	33.5	11.5	1.0
4	75.0	58.5	48.0	31.0	15.0	3.0	N/A
5	52.0	38.0	29.5	17.5	6.5	0.5	N/A
6	39.0	26.0	20.0	10.0	3.0	N/A	N/A

lar culture depth (1–6 cm), respectively. The relative logarithmic PPFD was negatively proportional to culture depth D and OD. This suggested that Lambert-Beer's law was applicable to the culture solution. In other words,

$$\log_{10}(e_{\text{out}}/e_{\text{in}}) = K \cdot \frac{N}{V} \cdot D = J \cdot \text{OD} \cdot D, \tag{9}$$

where K and J are the extinction coefficients defined for biomass concentration N/V and OD, respectively. The relationship between K and J can be written by $J = K \times 30.1 \times 10^6$, according to Eq. (8).

To obtain the extinction coefficients, least square fittings were performed. Specifically, Eq. (9) is rewritten as

$$\log_{10}(e_{\text{out}}/e_{\text{in}}) = J_{\text{OD}} \cdot D = J_{\text{D}} \cdot \text{OD}$$

where J_{OD} is defined by $J_{\text{OD}} := J \times \text{OD}$ and J_{D} is defined by $J_{\text{D}} := J \times D$. Then, J_{OD} and J_{D} were calculated from the slopes of the fitted lines in Figs. 5 (A) and 5 (B), respectively. The extinct coefficient J was then obtained as $J = -0.153$ from the slope in Fig. 5 (C), which plotted J_{OD} and J_{D}. The value of $J = -0.153$ corresponds to

$$K = \frac{J}{30.1 \times 10^6} = -5.1 \times 10^{-9},$$

which is the extinction coefficient of ACCB1808 in the unit of $mL \cdot cm^{-1} \cdot cell^{-1}$. The identified extinction coefficient $K = -5.1 \times 10^{-9}$ $mL \cdot cm^{-1} \cdot cell^{-1}$ was used for the prediction of the MR model in Section 3.2 and the full model in Section 4.2, and for the

Figure 5: (A) Relationships between relative logarithmic PPFD ($= \log(e_{\mathrm{out}}/e_{\mathrm{in}})$) and culture depths D at particular OD (OD=0.149–4.78). Circles and solid lines show experimental data and slope of the fitted line for J_{OD}, respectively. (B) Relationships between relative logarithmic PPFD and ODs at particular culture depth (1–6 cm). Circles and solid lines show experimental data and slope of the fitted line for J_{D}, respectively. (C) Relation of J_{OD} and J_{D} with depth D and OD. Diamonds and squares show slopes J_{OD} and J_{D} obtained in panels (A) and (B), respectively. Solid line show slope of the fitted line for J.

demonstration of cultivation planning in Section 5.