Mycobacterial Serine/Threonine phosphatase PstP is phospho-regulated and localized to mediate control of cell wall metabolism

Farah Shamma[1], E. Hesper Rego[2] and Cara Boutte [1*]

[1]Department of Biology, University of Texas at Arlington, Arlington, Texas

[2] Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, Connecticut

*Corresponding author, cara.boutte@uta.edu

Key Words: PstP, CwlM, FhaA, Wag31, phospho-regulation, peptidoglycan metabolism, Serine/Threonine Phosphatase

Abstract

The mycobacterial cell wall is profoundly regulated in response to environmental stresses, and this regulation contributes to antibiotic tolerance. The reversible phosphorylation of different cell wall regulatory proteins is a major mechanism of cell wall regulation. Eleven Serine/Threonine protein kinases (STPKs) phosphorylate many critical cell wall-related proteins in mycobacteria. PstP is the sole serine/ threonine phosphatase, but few proteins have been verified as PstP substrates. PstP is itself phosphorylated but the role of its phosphorylation in regulating its activity has been unclear. In this study we aim to discover novel substrates of PstP in *Mycobacterium tuberculosis* (*Mtb*). We show *in vitro* that PstP dephosphorylates two peptidoglycan regulators of *Mtb*, FhaA and Wag31. We also show that a phospho-mimetic mutation of T137 on PstP negatively regulates its catalytic activity against the cell wall regulatory substrates FhaA, Wag31, CwlM, PknB and PknA, and that the corresponding mutation in *Mycobacterium smegmatis (Msmeg)* causes mis-regulation of peptidoglycan *in vivo*. We show that PstP is localized to the septum, which likely restricts its access to certain substrates. These findings on the regulation of PstP provide insight into the control of cell wall metabolism in mycobacteria.

Introduction

Mycobacterium tuberculosis (*Mtb*), which causes Tuberculosis (TB), remains the age-old menace, infecting about 10 million and killing at least a million people worldwide each year (World Health Organization, 2021). TB infections require extended multi-drug antibiotic treatment regimens (Nguyen, 2016). The mycobacterial cell wall contributes to *Mtb*'s inherent antibiotic tolerance (Jarlier and Nikaido, 1994; Sarathy *et al.*, 2013; Batt *et al.*, 2020). The cell wall consists of a covalently linked peptidoglycan-arabinogalactan-mycolic acid layer core surrounding the plasma membrane (Marrakchi *et al.*, 2014). Peptidoglycan is a network of glycan chains cross-linked by small peptides, and is the innermost layer of the wall (Abrahams and Besra, 2021). The wall serves as permeability

barrier (Jarlier and Nikaido, 1990; Hett and Rubin, 2008; Hoffmann *et al.*, 2008;

Sarathy *et al.*, 2013), and is strongly regulated in response to environmental

stresses (Cunningham and Spreadbury, 1998; Betts *et al.*, 2002; Bhamidi *et al.*,

2012; Sarathy *et al.*, 2013), including infection (Sharma *et al.*, 2006; Doerks *et*

al., 2012). The cell wall changes in response to environmental stresses, and

these changes correlate with increased antibiotic tolerance (Xie *et al.*, 2005; Liu

et al., 2016; Sarathy *et al.*, 2017). The expression and activity of many cell wall

enzymes and regulators are also tightly coordinated over the course of the cell

cycle in order to maintain cell wall integrity during elongation and division (Kieser

and Rubin, 2014; Dulberger *et al.*, 2020; Bandekar *et al.*, 2020).

A crucial regulatory mechanism for transmission of environmental signals to

control cell growth is reversible Serine/Threonine (S/T) phosphorylation of a

myriad of proteins (Wang *et al.*, 1998; Juris *et al.*, 2000; Echenique *et al.*, 2004).

Mtb regulates cell physiology and growth (Kang *et al.*, 2005; Gee *et al.*, 2012;

Baer *et al.*, 2014; Boutte *et al.*, 2016) through 11 Serine/Threonine kinases

(STPKs) (PknA-B and PknD-L) and one essential S/T protein phosphatase PstP

(Cole *et al.*, 1998; Bach *et al.*, 2009). PstP is also essential in *Msmeg* (Sharma *et*

al., 2016; DeJesus *et al.*, 2017).

PstP belongs to the protein phosphatase 2C (PP2C) subfamily of the strictly

divalent metal-ion dependent protein serine/threonine phosphatases (Cohen,

1989; Barford, 1996; Chopra *et al.*, 2003), known to regulate prokaryotic and

eukaryotic cell growth and division, sporulation, stress response and metabolic

processes by responding to various environmental signals (Vijay *et al.*, 2000:2;

Irmler and Forchhammer, 2001; Mougous *et al.*, 2007; Lu and Wang, 2008;

Bradshaw and Losick, 2015; Bradshaw *et al.*, 2017). PstP$_{Mtb}$ shares structural

architecture and conserved catalytic residues with the human PP2Cα (Pullen *et*

al., 2004) (Figure 1), which is considered the prototypical member of the PP2C

family. The N-terminal cytoplasmic enzymatic domain of PstP$_{Mtb}$ (Figure 1)

connects to its C-terminal extracellular domain via a transmembrane helix (Pullen

et al., 2004). PstP$_{Mtb}$ is itself phosphorylated on T134, T141, T174 and T290

(Figure 1) and its phosphorylation affects its activity against small molecule

substrates *in vitro* (Sajid *et al.*, 2011) and cell wall metabolism *in vivo* (Shamma

et al., 2021)*.*

Two essential STPKs, PknA and PknB, play important roles in cell wall regulation

by phosphorylating enzymes or regulatory proteins involved in cell division and

growth (Sassetti and Rubin, 2003; Kang *et al.*, 2005; Fernandez *et al.*, 2006;

Kusebauch *et al.*, 2014; Boutte *et al.*, 2016). Phosphorylation of different cell wall

substrates has different effects on their activities. For example, phosphorylation

activates synthesis of peptidoglycan precursors through regulatory protein CwlM

(Boutte *et al.*, 2016), and inhibits synthesis of mycolic acids by inactivating

several mycolic acid biosynthesis enzymes in *Mtb* (Molle *et al.*, 2006; Molle and

Kremer, 2010; Veyron-Churlet *et al.*, 2010; Khan *et al.*, 2010; Slama *et al.*, 2011;

Vilchèze *et al.*, 2014). PstP is suggested to be a global negative regulator of

STPK-mediated phosphorylation (Iswahyudi *et al.*, 2019). Being the only S/T

protein phosphatase (Cole *et al.*, 1998), it seems likely to dephosphorylate many

of the substrates phosphorylated by the 11 STPKs (Prisic and Husson, 2014).

However, only a few proteins have been identified to be substrates of PstP. The

biochemically-verified cell wall substrates of PstP include the STPKs PknA and

PknB (Boitel *et al.*, 2003; Chopra *et al.*, 2003; Durán *et al.*, 2005; Sajid *et al.*,

2011), KasA and KasB, enzymes of the fatty acid synthase –II (FAS-II) system

of mycolic acid biosynthesis (Molle *et al.*, 2006), the arabinogalactan layer

regulator EmbR (Sharma *et al.*, 2006), and CwlM, which regulates the

peptidoglycan layer (Shamma *et al.*, 2021).

CwlM is crucial for peptidoglycan synthesis; it interacts with MurA (Boutte *et al.*,

2016), the first enzyme of peptidoglycan precursor synthesis (Marquardt *et al.*,

1992), and CwlM's phosphorylation is required for MurA's activity (Boutte *et al.*,

2016). So dephosphorylation of CwlM~P by PstP (Shamma *et al.*, 2021) should

halt peptidoglycan synthesis. Phosphorylated CwlM~P also interacts with FhaA

(Turapov *et al.*, 2018), which is involved in modulating peptidoglycan

biosynthesis (Gee *et al.*, 2012; Viswanathan *et al.*, 2015; Viswanathan *et al.*,

2017) and unphosphorylated CwlM interacts with the peptidoglycan precursor

lipid-II flippase MurJ (Turapov *et al.*, 2018), suggesting that CwlM's activity and

interactions with other peptidoglycan regulators and enzymes are phospho-

regulated and that both CwlM and FhaA are part of a phosphorylation-controlled

peptidoglycan regulatory hub (Turapov *et al.*, 2018).

FhaA and Wag31 are both regulators of mycobacterial cell wall metabolism.

FhaA helps maintain cell envelope integrity, cell length and antibiotic tolerance

(Viswanathan *et al.*, 2015; Viswanathan *et al.*, 2017). Available data suggest that

FhaA binds and inhibits the MurJ flippase (Gee *et al.*, 2012) and stabilizes PbpA

(Viswanathan *et al.*, 2017). FhaA is phosphorylated by PknB (Grundner *et al.*,

2005; Prisic *et al.*, 2010; Roumestand *et al.*, 2011), though the role of its

phosphorylation in regulation is not yet clear. Wag31 localizes at the cell poles

(Kang *et al.*, 2005), is essential for polar peptidoglycan synthesis (Kang *et al.*,

2008; Jani *et al.*, 2010) and is phosphorylated by the STPK PknA (Kang *et al.*,

2005). Experiments with phospho-mimetic and phospho-ablative mutations of

*wag31*T73 suggest that its phosphorylation has a role in controlling

peptidoglycan metabolism (Jani *et al.*, 2010). It was not known whether FhaA

and Wag31 were actively dephosphorylated, or if their phosphorylated forms

were removed through protein degradation.

In this study we show that FhaA$_{Mtb}$ and Wag31$_{Mtb}$ are substrates of PstP$_{Mtb}$ *in*

vitro, and that PstP dephosphorylates Wag31 much faster than it does FhaA or

CwlM. PstP's activity against different substrates could be controlled by its

localization, by its own phosphorylation (Sajid *et al.*, 2011) or through interactions

with other regulatory proteins. Here, we show that the T137E phospho-mimetic

mutation of PstP$_{Mtb}$ inhibits PstP$_{Mtb}$'s catalytic activity against the substrates

CwlM, FhaA, Wag31, PknB and PknA *in vitro*. Furthermore, the corresponding

phospho-mimetic mutation in *Msmeg* leads to dis-regulation of PG metabolism.

We also find that PstP is enriched at the cell division site, suggesting that its

access to substrates at the pole might also control its activity. We suggest a

model where PstP's activity against assorted substrates is regulated by substrate

affinity and protein localization, and further toggled by the phosphorylations on

PstP.

Methods and Materials

Bacterial strains and culture conditions

All *Mycobacterium smegmatis* mc^{2}155 ATCC 700084 cultures were grown in 7H9

(Becton, Dickinson, Franklin Lakes, NJ) medium containing 0.003 g/liter catalase,

5 g/liter bovine serum albumin (BSA), 2 g/liter dextrose, 0.2% glycerol, 0.85

g/liter NaCl, and 0.05% Tween 80 and incubated at 37°C until log. phase. 1xPBS

supplemented with 0.05% Tween 80 was used for starvation assays. Cultures

were plated on LB Lennox agar (Fisher BP1427-2).

E. coli Top10, XL1-Blue, and Dh5α strains were used for cloning and *E. coli* BL21

Codon Plus and BL21(DE3) strains were used for expressing proteins. Antibiotic

concentrations for *M. smegmatis* were 25μg/ml kanamycin and 20 μg/ml zeocin.

Antibiotic concentrations for *E. coli* were 50 μg/ml kanamycin, 25 μg/ml zeocin,

20 μg/ml chloramphenicol, and 140 μg/ml ampicillin.

Cloning and Strain construction

N-terminally Histidine (His) tagged cytosolic domains (1-300 amino acids (Gupta

et al., 2009)) of *pstP*$_c$T137E$_{Mtb}$ and *pstP*$_c$T141E$_{Mtb}$, and *fhaA*WT$_{Mtb}$ codon-

optimized for expression in *E. coli* were ordered from IDT, Inc. and cloned in to

pET28a.

The *Msmeg* mc^{2}155 strains Δ*pstP*::lox L5::pCT94-p766tetON6-*pstP*$_{Msmeg}$T134E

and Δ*pstP*::lox L5::pCT94-p766tetON6-*pstP*$_{Msmeg}$WT were generated by L5 allele

swapping as described previously (Pashley and Parish, 2003; Shamma *et al.*,

2021). All vectors and strains are described in supplementary tables 1-3.

A vector that expressed mCherry-2B-PstP$_{Msmeg}$ was constructed using isothermal

assembly and integrated at the tweety phage integration site in *Msmeg*. In brief,

pstP was amplified from *Msmeg* genomic DNA, and subcloned into a plasmid

along with *mCherry2B* and the uv15 ribosomal promoter. The resulting vector

was transformed into *Msmeg*.

Growth Curve Assay

At least three biological replicates of $pstP_{Msmeg}$T134E and WT allelic variants

were grown in 7H9 medium up to the mid-log. phase. The growth curve assays

were performed in non-treated non-tissue culture plates using a plate reader

(BioTek Synergy neo2 multi mode reader) in 200μl of 7H9 medium at 37°C

starting at an Optical Density at 600nm (OD_{600}) of 0.1.

The doubling times of each strain were calculated in the non-linear (curve fit)

method using an exponential growth equation with the least squares (ordinary) fit

in GraphPad Prism (version 7.0d). *P* values were calculated using two-tailed

unpaired t-tests.

Protein purification

His-MBP-PknA$_{Mtb}$ was expressed in *E. coli* BL21(DE3) and all other proteins were

expressed using *E. coli* BL21 Codon Plus cells. The *Mtb* genes of all studied

factors were cloned into expression vectors. His-MBP-PknB$_{Mtb}$, His-SUMO-

CwlM$_{Mtb}$, His-PstP$_c$WT$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$ were expressed and purified as

previously described (Kieser *et al.*, 2015; Shamma *et al.*, 2021). His-

PstP$_c$T137E$_{Mtb}$ and His-PstP$_c$T141E$_{Mtb}$ were expressed and purified using the

same conditions and buffers used to purify His-PstP$_c$WT$_{Mtb}$ and His-

PstP$_c$T174E$_{Mtb}$ (Shamma *et al.*, 2021).

His-Wag31$_{Mtb}$ was induced with 0.5mM IPTG (isopropyl-β-D-

thiogalactopyranoside) for 6h at 18°C, and purified on Ni-NTA resin (G-

Biosciences 786-940) then dialyzed and concentrated. The buffer for His-

Wag31$_{Mtb}$ was 50mM Tris pH 8, 350mM NaCl, 1 mM dithiothreitol (DTT), and

10% glycerol. Imidazole (10mM and 20mM) was added to the buffers for lysis of

the cell pellets and washing the Ni-NTA resin respectively, and 250mM imidazole

was added for elution. The lysis buffer was supplemented with 1mM PMSF and

0.2% Triton X100.

His-MBP-PknA$_{Mtb}$ (1-279 amino acids containing the kinase domain (Baer et $al.$,

2014)) was induced with 0.5mM IPTG at 18°C overnight, purified with 5ml Ni-

NTA resins (BIO-RAD EconoFit Nuvia IMAC, 12009287), then run over the size

exclusion resin (Enrich SEC650, BIO-RAD 780-1650) after dialysis and

concentration to get soluble protein. The buffer for purifying His-MBP-PknA$_{Mtb}$

was 50mM Tris pH 7.5, 150mM NaCl, 20% glycerol. PMSF (1mM) was added to

the buffer while resuspending the cell pellets. Imidazole was used in buffer for

application to (20mM) and for elution from Ni-NTA resin (200mM).

His-FhaA$_{Mtb}$ was induced with 1mM IPTG at 18°C overnight and purified on Ni-

NTA resin (G-Biosciences, 786-940 in 5 ml Bio-Scale Mini Cartridges, BIO-RAD

7324661), then dialyzed, concentrated, and run over size exclusion resin (GE

Healthcare Sephacryl S-200 in HiPrep 26/70 column). His-FhaA$_{Mtb}$ was purified

using the buffers described in (Roumestand et $al.$, 2011).

***In vitro* dephosphorylation assays**

His-SUMO-CwlM$_{Mtb}$ at 0.15 µg/µl was phosphorylated in $vitro$ with His-MBP-

PknB$_{Mtb}$ at 0.015 µg/µl for 30 minutes at room temperature with 0.15µCi/µl ATP

[γ-^{32}P] (3000Ci/mmol, 10mCi/ml, PerkinElmer BLU002A250UC) and 2mM MnCl$_2$ in

buffer (50mM Tris, pH 7.5, 250mM NaCl, 10% glycerol and 1mM DTT). Right

after collecting kinase reaction samples, the leftover mixture was split into equal

parts in different microcentrifuge tubes to carry on phosphatase assays by

adding replicates of His-PstP$_c$WT$_{Mtb}$, His-PstP$_c$T137E$_{Mtb}$, His-PstPcT141E$_{Mtb}$ or

His-PstPcT174E$_{Mtb}$ at 0.012 µg/µl to the kinase reaction mixtures containing 0.12

µg/µl of His-SUMO-CwlM$_{Mtb}$ and incubated in phosphatase buffer (50mM Tris pH

7.5, 10mM MnCl$_2$, and 1mM DTT) at room temperature up to 40 minutes. His-

FhaA*Mtb* at 0.15 µg/µl was phosphorylated *in vitro* with His-MBP-PknB*Mtb* at 0.015

µg/µl for 45 minutes at 37°C with 0.15µCi/µl ATP [γ-^{32}P] in presence of 5mM

MnCl$_2$ in 25mM Tris pH 7.5 and 1mM DTT. Right after collecting 0 minute kinase

reaction sample, the reaction mixture was then split into equal parts and

incubated with either PstP$_c$WT*Mtb* or different phospho-mimetic variants of it at

0.012 µg/µl up to 80 minutes at room temperature in presence of the

phosphatase buffer where the concentration of the substrate was 0.12 µg/µl For

CwlM and FhaA, the amount of kinase and phosphatases was 1/10th the amount

of substrate, by weight.

His-Wag31*Mtb* at 0.15 µg/µl was phosphorylated with His-MBP-PknA*Mtb* at 0.03

µg/µl for 2 hours at room temperature in presence of 0.2µCi/µl ATP [γ-^{32}P], 10mM

MnCl$_2$ and buffer (50mM Tris pH 7.5, 20% glycerol and 150mM NaCl). The amount of

PknA was 1/5th the amount of Wag31 substrate, by weight.

The kinase reaction mixture was equally split in to separate tubes after the 0

second kinase reaction sample was taken. His-PstP$_c$WT*Mtb* or its different

phospho-mimetic variants at 0.0013 µg/µl were added to the mixture containing

0.13 µg/µl substrate and incubated at room temperature for up to 120 seconds in

the phosphatase buffer. The amount of phosphatase was 1/100th the amount of

Wag31 substrate in the phosphatase reactions, by weight.

Control samples with no kinase and with no phosphatase were performed in all

the above-mentioned substrate dephosphorylation assays. All *in vitro*

dephosphorylation assays were carried out separately with two individually

purified batches of WT and phospho-mimetic mutant of PstP as biological

replicates.

Quantification of γ-^{32}P signals

Samples from each *in vitro* biochemical reaction were collected before (0 minute

or second) and after addition of each of the phosphatases at several different

time points. The samples were run on 12% (Mini-Protean TGX, BIO-RAD,

4561046) gels and stained with Coomassie blue solution. After destaining, gels

were dried using the GD200 Vacuum Gel Drying system (Hoefer Inc.) and

imaged with phosphor screens (Molecular Dynamics) and a scanner (Storm 860,

Amersham Biosciences). ImageQuant software (Molecular Dynamics) was used

to process the images.

The protein bands on the gel and the phospho-signals on the autoradiogram

were then quantified with FIJI. The intensities of the substrate protein bands on

the Coomassie-stained gel and their corresponding phospho-signals on the

autoradiograms at each time point were normalized against the respective

protein band and the phospho-signal at 0 minute (His-SUMO-CwlM$_{Mtb}$ and His-

FhaA$_{Mtb}$) or 0 second (His-Wag31$_{Mtb}$) before addition of the phosphatase. These

relative intensities were used to calculate the intensity of phospho-signal over the

amount of substrate for each time point. These values were used to plot the

graph. The P32 signal intensities of the kinases on autoradiograms at each time

point were normalized against the respective phospho-signal on the kinase at 0

minute (His-MBP-PknB$_{Mtb}$) or 0 second (His-MBP-PknA$_{Mtb}$) before addition of the

phosphatase. These relative P32 signal intensity values were then used to plot

the graph. All values were plotted and p-values were calculated using two-tailed,

unpaired Student's t-test in GraphPad Prism (version 7.0d).

Cell staining

At least three biological replicates of *pstP$_{Msmeg}$* T134E and WT allelic variants

were grown in 7H9 medium up to mid log. phase. Cultures were washed and

then starved in 1xPBS with 0.05% Tween 80 (PBST) at OD$_{600=}$ 0.3 for 15

minutes. 100µl of starved culture was then treated with 3µl of 10mM fluorescent

D-amino acid HADA for 10 minutes. Cells were then pelleted and resuspended in

PBST and fixed with 10µl of 16% paraformaldehyde for 10 minutes at room

temperature. Cells were then pelleted and resuspended in PBST before imaging

on agarose pads.

For staining cells in the log. phase, 1µl of 1mM HADA was added to 100µl

culture grown up to the mid-log. phase and incubated at 37°C for 15 minutes.

Stained cells were then pelleted, washed and fixed as described above.

Static microscopy and image analysis

A Nikon Ti-2 widefield epifluorescence microscope with a Photometrics Prime

95B camera and a Plan Apo 100x, 1.45 NA objective lens was used to image

cells. A 350/50nm excitation filter and a 460/50nm emission filter were used to

take the blue fluorescence images for HADA staining. All images were captured

using the NIS Elements software and analyzed using FIJI and MicrobeJ (Ducret

et al., 2016). Appropriate parameters for length, width and area were set in

MicrobeJ to detect cells. All detected dividing (V-snapping) cells in the images

were split at the septum so that each daughter cell could be considered as a

single cell. No overlapping cells in the images were included in the analysis.

Length and mean-intensities of HADA signals of at least 330 cells from each of

*pstP*T134E$_{Msmeg}$ and *pstP*WT$_{Msmeg}$ (at least 110 cells from each of three biological

replicate strains of each genotype) were quantified using MicrobeJ. The values of

the mean intensities of 330 cells of each *pstP* allelic mutant and WT are

represented in the graphs generated using GraphPad Prism (version 7.0d).

The medial intensity profiles of HADA signals in cells from different *pstP* allele

strains in log. phase and starvation analyzed with MicrobeJ were plotted on the

Y-axis over relative positions of cells using the "XStatProfile" plotting feature in

MicrobeJ to show the subcellular localization of fluorescent intensities in cells.

Demographs of HADA signal intensity across cell lengths in log. phase and

starvation were built using the "Demograph" feature of MicrobeJ by plotting the

medial intensity profiles of HADA signals.

Time-lapse microscopy

An inverted Nikon Ti-E microscope was used for the time-lapse imaging. An
environmental chamber was maintained at 37°C. Cells were cultivated in an B04
microfluidics plate from CellAsic; continuously supplied with fresh medium; and
imaged every 15 minutes using a 60x 1.4 N.A. phase contrast objective, with an
intermediate magnification of 1.5x. mCherry2B was excited with a SpectraX
Illuminator at 561-nm. Fluorescence was detected with a bandpass filter of
595/40 and detected with an ORCA Flash Digital sCMOS.

Results

PstP$_{Mtb}$ dephosphorylates FhaA$_{Mtb}$ and Wag31$_{Mtb}$ in vitro

FhaA$_{Mtb}$ is phosphorylated by PknB$_{Mtb}$ in vivo (Prisic et al., 2010) and in vitro
(Grundner et al., 2005; Roumestand et al., 2011) on T116 (Roumestand et al.,
2011). Wag31$_{Mtb}$ is phosphorylated at T73 by PknA$_{Mtb}$ (Kang et al., 2005). To
determine if FhaA$_{Mtb}$ and Wag31$_{Mtb}$ are substrates of PstP$_{Mtb}$, we performed in
vitro dephosphorylation assays with purified full-length His-FhaA$_{Mtb}$, His-
Wag31$_{Mtb}$, N-terminally His-MBP tagged cytosolic kinase domains of PknA$_{Mtb}$ and
PknB$_{Mtb}$, and the cytosolic region of PstP$_{Mtb}$ containing the catalytic domain (His-
PstP$_c$WT$_{Mtb}$).

We phosphorylated His-FhaA$_{Mtb}$ with His-MBP-PknB$_{Mtb}$ and ATP [γ-^{32}P]. After
addition of 1/10th the amount of His-PstP$_c$WT$_{Mtb}$, the phosphorylation on His-
FhaA$_{Mtb}$~P decreases by ~80% in 80 minutes (Figure 2a,b). Our control assay
with no phosphatase shows that the phosphorylation on His-FhaA$_{Mtb}$~P is stable
over time in the phosphatase buffer (Figure 2a, b). These data show that the cell
wall regulator FhaA$_{Mtb}$ is a substrate of PstPWT$_{Mtb}$ in vitro.

We performed similar phosphatase assays to determine if PstP$_c$WT$_{Mtb}$ can
dephosphorylate Wag31$_{Mtb}$. We first phosphorylated His-Wag31$_{Mtb}$ with His-MBP-

PknA$_{Mtb}$ and ATP [γ-^{32}P], then added His-PstP$_c$WT$_{Mtb}$. With 1/100th the amount of

PstP compared to Wag31, Wag31$_{Mtb}$ phosphorylation decreased by about ~90%

in 2 minutes (Figure 3a,b), only in the presence of His-PstP$_c$WT$_{Mtb}$. These data

show that Wag31$_{Mtb}$ is a substrate of PstPWT$_{Mtb}$ *in vitro*.

Our previous biochemical data that demonstrated PstP$_{Mtb}$ dephosphorylates

CwlM$_{Mtb}$ (Shamma *et al.*, 2021) is reproducible in our *in vitro* assays with ATP [γ-

^{32}P] where we observe a 97% decrease in phosphorylation on His-SUMO-CwlM$_{Mtb}$~P

over a time period of 40 minutes after addition of His-PstP$_c$WT$_{Mtb}$ (Figure 4a, b). In all

the assays, we see no phosphorylation signals on the WT and phospho-mimetic

PstPs on the autoradiograms (data not shown) since PstP can quickly

dephosphorylate itself (Sajid *et al.*, 2011) and the purified forms are thus

unphosphorylated.

Because measuring the rates of dephosphorylation by PstP against different

substrates required different PstP concentrations and timing, we calculated the

initial linear rate of dephosphorylation per unit concentration of PstP, in order to

compare PstP's activity against different substrates. This comparison indicates

that His-PstP$_c$WT$_{Mtb}$ dephosphorylates His-Wag31$_{Mtb}$ more than 500 times faster

than His-SUMO-CwlM$_{Mtb}$ and about 2500 times faster than His-FhaA$_{Mtb}$ (Figure

4c).

Phospho-mimetic mutation at PstPT137$_{Mtb}$ inhibits dephosphatase activity

PstP$_{Mtb}$ is phosphorylated on the conserved threonine (T) residues 137, 141 and

174 (Figure 1), and when phosphorylated, shows increased activity against small

molecule substrates *in vitro* (Sajid *et al.*, 2011). We hypothesized that

phosphorylation of the threonine residues of PstP individually might help regulate

PstP's activity through switching its catalytic activity against different substrates

or by affecting substrate affinity. To test this, we made phospho-mimetic (T to E)

(Cottin *et al.*, 1999) mutants of PstP$_{Mtb}$ at each of the three conserved

phosphorylation sites of PstP$_{Mtb}$ in the catalytic domain and purified them (His-

PstP$_c$T137E$_{Mtb}$, His-PstP$_c$T141E$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$). We previously

showed that *in vitro* His-PstP$_c$WT$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$ dephosphorylate

His-SUMO-CwlM$_{Mtb}$ at the same rate (Shamma *et al.*, 2021). We performed

separate *in vitro* phosphatase assays with the substrates His-FhaA $_{Mtb}$, His-

Wag31$_{Mtb}$ and His-SUMO-CwlM$_{Mtb}$ with the phospho-mimetic versions of PstP.

His-PstP$_c$T137E$_{Mtb}$ is completely inactive *in vitro* against all three substrates

(Figures 2,3,4). This suggests that phosphorylation on T137 of PstP may inhibit

its activity against multiple substrates.

His-MBP-PknB$_{Mtb}$ and His-MBP-PknA$_{Mtb}$, the STPKs used in these assays, are

also confirmed substrates of PstP (Sajid *et al.*, 2011). PknA and PknB are

activated to phosphorylate their own substrates when they themselves are

phosphorylated at their activation loops (Baer *et al.*, 2014). Because we already

had these kinases in our phosphatase assays, we were able to measure

dephosphorylation of His-MBP-PknB$_{Mtb}$ and His-MBP-PknA$_{Mtb}$ by PstP in the

context of a 10-fold and 5-fold excess of His-SUMO-CwlM$_{Mtb}$ or His-Wag31$_{Mtb}$,

respectively. We found that His-PstP$_c$T137E$_{Mtb}$ dephosphorylates both His-

MBP-PknB$_{Mtb}$ (Figure 5a,b) and His-MBP-PknA$_{Mtb}$ (Figure 5c,d) more slowly than

His-PstP$_c$WT$_{Mtb}$ and the other PstP mutants. This shows that His-PstP$_c$T137E$_{Mtb}$

is inhibited against at least five different substrates.

375　　Phospho-mimetic mutations at PstPT141$_{Mtb}$ and PstPT174$_{Mtb}$ moderately

376　　inhibit Wag31's dephosphorylation

His-PstP$_c$T141E$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$ dephosphorylate His-FhaA$_{Mtb}$ (Figure

2,4c) and His-SUMO-CwlM$_{Mtb}$ (Figure 4) at about the same rate as His-

PstP$_c$WT$_{Mtb}$, but dephosphorylate His-Wag31$_{Mtb}$ about half as fast as His-

PstP$_c$WT$_{Mtb}$ (Figure 3, 4c). These biochemical data support our hypothesis that

phosphorylation on PstP may modulate its activity by affecting its substrate

specificity. Phosphorylation at T141 and T174 appear to inhibit binding or

catalysis of Wag31 but not FhaA or CwlM.

Phospho-site T134 of PstP*Msmeg* regulates peptidoglycan metabolism in

starvation

Phosphorylation of CwlM allows it to activate the peptidoglycan precursor

enzyme MurA, thereby licensing peptidoglycan synthesis (Boutte *et al.*, 2016).

Phosphorylation of CwlM also affects its binding to FhaA and the lipid II flippase

MurJ (Turapov *et al.*, 2018). CwlM is quickly dephosphorylated in the transition to

stasis (Boutte *et al.*, 2016) likely by PstP (Shamma *et al.*, 2021). The function of

phosphorylation on FhaA and Wag31 is not yet clear, but available data (Jani *et*

al., 2010; Gee *et al.*, 2012; Viswanathan *et al.*, 2016) suggest it could also affect

peptidoglycan metabolism. Dephosphorylation of these regulators could be

important for downregulation of peptidoglycan metabolism in stasis.

As the phospho-mimetic mutation at PstPT137*Mtb* inhibits PstP's

dephosphorylation of three peptidoglycan regulatory proteins *in vitro* (Figure

2,3,4), we sought to determine if the corresponding phosphosite on PstP (T134)

might affect peptidoglycan metabolism *in vivo* in *Msmeg*. We made *Msmeg*

strains with a phospho-mimetic *pstP*T134E*Msmeg* allele replacing the wild-type

pstP. We reported previously that the biological replicates of *pstP*T134E*Msmeg*

mutant strains had inconsistent doubling times (Shamma *et al.*, 2021) probably

due to potential suppressor mutations. We confirmed that the slow growing

clones could, after passaging, accumulate suppressor mutations that allowed

them to grow as fast as the *pstP*WT strain. To characterize the original

*pstP*T134E*Msmeg* strain, we isolated numerous clones from fresh transformations,

and measured the doubling times of each before continuing with experiments.

We show data only from clones that grew at the slower rate.

We used three biological replicate *pstP*T134E*Msmeg* strains that had consistently

slower growth than the *pstP*WT*Msmeg* strains (Figure 6a, red bar) in our next

experiments. We stained both log.phase and starved *pstP* T134E*Msmeg* and

WT*Msmeg* strains using a fluorescent D-amino acid HADA (Kuru *et al.*, 2012) as a

reporter of active peptidoglycan metabolism. We saw a slight increase in length

(Figure 6b,d and Supplementary Figure S2a,b) and HADA intensity (Figure 6c,d

and Supplementary Figure S2a,b) in the *pstP* T134E*Msmeg* strains in the log.

phase compared to the WT*Msmeg* strains. After starvation, these differences

increased (Figure 6b,c,e and Supplementary Figure S2c,d). WT*Msmeg* strains

quickly downregulate peptidoglycan metabolism upon starvation (Boutte *et al.*,

2016), as the decreased HADA staining (Figure 6c,d,e,f,g) shows. The starved

*pstP*T134E*Msmeg* strains show significantly higher HADA signal than the WT*Msmeg*

(Figure 6c, top right panels of d,e and Supplementary Figure S2b,d) indicating a

defect in downregulating peptidoglycan metabolism. This suggests that the

presence of a constant negative charge on phosphosite T134 of PstP*Msmeg*

impairs downregulation of peptidoglycan synthesis, likely by inhibiting PstP's

dephosphorylation of peptidoglycan regulators (Figure 2,3,4).

Our data showing that the *pstP*T134E*Msmeg* strains have increased length (Figure

6b) and higher HADA intensity (Figure 6c,d,f and Supplementary Figure S2b) in

the log. phase support the notion (Jani *et al.*, 2010; Turapov *et al.*, 2018) that

peptidoglycan regulation involves a balance in phosphorylated and

dephosphorylated forms of its regulatory substrates and PstP's phospho-

regulation helps toggle its activity against substrates in log. phase. The

misregulation of peptidoglycan metabolism (Figure 6c,d,e,f,g and Supplementary

Figure S2b,d) may account for the slower growth of the *pstP*T134E*Msmeg* strains

that we see in our data (Figure 6a), though it is likely that other processes

beyond peptidoglycan metabolism are also misregulated, as PstP likely has

many substrates involved in diverse processes (Molle *et al.*, 2006; Iswahyudi *et*

al., 2019).

PstP localizes at the septum in exponentially growing *Msmeg* cells

We sought to determine if PstP's activity against its diverse substrates could be regulated through protein localization. We made an *Msmeg* strain carrying an additional copy of *pstP$_{Msmeg}$* fused to *mCherry2B*. By time-lapse fluorescence microscopy, we observed that mCherry2B-PstP$_{Msmeg}$ appears to be uniformly distributed along the membrane throughout most of the cell cycle, but becomes localized at the septum before cell separation (Figure 7). These data suggest that PstP is also regulated by its access to substrates at different sites in the cell.

Discussion

Previous works have suggested that PstP plays a crucial role as a negative regulator of protein Serine/ Threonine phosphorylation (Boitel *et al.*, 2003; Molle *et al.*, 2006; Sajid *et al.*, 2011; Boutte *et al.*, 2016; Iswahyudi *et al.*, 2019; Dulberger *et al.*, 2020; Shamma *et al.*, 2021). Our findings indicate that PstP is regulated by its own phosphorylation (Figure 2,3,4,5) to regulate cell wall metabolism (Figure 6), and that it may also be regulated by its localization (Figure 7).

We see that PstPWT$_{Mtb}$ dephosphorylates CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$ at different rates *in vitro* (Figure 4c), with Wag31$_{Mtb}$ being the fastest to be dephosphorylated and FhaA the slowest (Figure 4c). We showed that PstP localizes at the septum in the exponentially growing *Msmeg* cells (Figure 7). Both FhaA and Wag31 localize at the poles and septum (Gee *et al.*, 2012; Hannebelle *et al.*, 2020). Our localization data (Figure 7) suggest that both Wag31 and FhaA (Figure 2, 3, 4c) may be more highly phosphorylated at the poles than at the septum. PstP's rapid dephosphorylation of Wag31 (Figure 3, 4c) suggests that it is likely to be nearly completely unphosphorylated at the septum.

Our data in Figure 2, 3 and 4 show that negative charges at PstPT141$_{Mtb}$ and T174$_{Mtb}$ slow dephosphorylation of Wag31$_{Mtb}$, but not FhaA$_{Mtb}$ and CwlM$_{Mtb}$. Phosphorylation site T141 is on the flap subdomain and T174 is on the β-sheet

core (β8) on PstP$_{Mtb}$ (Pullen *et al.*, 2004), but both are facing into the same

groove between these domains (Figure 1). T137 is the first residue of the flap

subdomain (Pullen *et al.*, 2004), and it faces away from the groove with the other

phospho-sites (Figure 1). We propose that the T141/ T174 groove may contribute

to toggling specificity of PstP against certain substrates, depending on the

phosphorylation status.

Our *in vitro* data in Figure 2,3, 4 and 5 show that PstPT137E$_{Mtb}$ inhibits its activity

against CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$, and also against the STPKs PknA and

PknB that phosphorylate them, all of which are involved in regulation of

peptidoglycan metabolism (Thakur and Chakraborti, 2006; Kang *et al.*, 2008; Jani

et al., 2010; Gee *et al.*, 2012; Prisic and Husson, 2014; Boutte *et al.*, 2016;

Viswanathan *et al.*, 2016; Viswanathan *et al.*, 2017; Carette *et al.*, 2018; Turapov

et al., 2018). We expected the corresponding phospho-mimetic mutant

*pstP*T134E$_{Msmeg}$ allele to cause mis-regulation of peptidoglycan synthesis *in vivo*.

Our *in vivo* data show that the *pstP*T134E$_{Msmeg}$ strains grow more slowly, and

have an increase in cell length and HADA staining in log. phase and starvation

(Figure 6). We see relatively mild defects in peptidoglycan metabolism in these

strains in log. phase (Figure 6b,c), which suggests that phospho-site T134 on

PstP may not be important for controlling peptidoglycan metabolism in active

growth. Alternatively, the function of the negative charge at T134 may be

different in the cell: phosphorylation here could help recruit an interacting protein

that could alter PstP's activity in a different way. As expected, however, we do

see greater defects in peptidoglycan metabolism during starvation (Figure

6c,d,e,f,g), suggesting that dephosphorylation of CwlM, FhaA, Wag31, PknB or

PknA, and/or other substrates, is more important in the transition to stasis than in

log. phase. The cells still largely downregulate peptidoglycan metabolism,

indicating that other regulatory systems contribute to this response, or that

*pstP*T134E has different effects *in vivo* than *in vitro*.

Although our findings are limited to PstP's activity on certain substrates only,

altogether they prove that the individual phosphosites are important in regulating

PstP. *In vitro*, we see that a negative charge on T137 of PstP*Mtb* inhibits its
activity against three PG regulators (Figure 2,3,4), as well as the essential
kinases PknB and PknA when they are mixed together (Figure 5). In the cell, the
substrates and kinases also co-exist, and PstP may exert its effect directly by
dephosphorylating the substrates or indirectly by acting on the kinase (Bhaskara
et al., 2019), or both simultaneously. There are likely other mechanisms of cell
wall regulation involved in the complex *in vivo* scenario like protein localization or
PstP's interaction with other regulatory proteins that may contribute to PstP's
regulation as well. We do not know the conditions under which PstP is
phosphorylated, or the combinations of PstP's phosphorylations that exist in the
cell. So the *in vitro* data may not mirror the deep complexity of the STPK-
substrate-PstP network *in vivo*.

In this work we show that PstP is regulated by both phosphorylation and
localization. This work demonstrates the importance of protein
dephosphorylation, not just phosphorylation, in regulating the activity of proteins
involved in cell wall metabolism in mycobacteria. Future work that addresses
when all of these important phospho-sites are phosphorylated and
dephosphorylated will help us to understand how all these protein regulators
contribute to coordinating the construction and integrity of the cell wall in
changing conditions.

Acknowledgement:

This work was supported by NIH grant R01AI148917 to CCB and R01AI148255
to EHR.

521
522
523
524

References:

Abrahams, K.A., and Besra, G.S. (2021) Synthesis and recycling of the mycobacterial cell envelope. *Current Opinion in Microbiology* **60**: 58–65 https://linkinghub.elsevier.com/retrieve/pii/S1369527421000187. Accessed February 17, 2022.

Bach, H., Wong, D., and Av-Gay, Y. (2009) Mycobacterium tuberculosis PtkA is a novel protein tyrosine kinase whose substrate is PtpA. *Biochemical Journal* **420**: 155–162 https://portlandpress.com/biochemj/article/420/2/155/44900/Mycobacterium-tuberculosis-PtkA-is-a-novel-protein. Accessed November 15, 2021.

Baer, C.E., Iavarone, A.T., Alber, T., and Sassetti, C.M. (2014) Biochemical and Spatial Coincidence in the Provisional Ser/Thr Protein Kinase Interaction Network of *Mycobacterium tuberculosis*. *J Biol Chem* **289**: 20422–20433 http://www.jbc.org/lookup/doi/10.1074/jbc.M114.559054. Accessed October 23, 2019.

Bandekar, A.C., Subedi, S., Ioerger, T.R., and Sassetti, C.M. (2020) Cell-Cycle-Associated Expression Patterns Predict Gene Function in Mycobacteria. *Current Biology* **30**: 3961-3971.e6 https://linkinghub.elsevier.com/retrieve/pii/S0960982220310952. Accessed September 27, 2021.

Barford, D. (1996) Molecular mechanisms of the protein serine/threonine phosphatases. *Trends Biochem Sci* **21**: 407–412.

Batt, S.M., Burke, C.E., Moorey, A.R., and Besra, G.S. (2020) Antibiotics and resistance: the two-sided coin of the mycobacterial cell wall. *The Cell Surface* **6**: 100044 https://linkinghub.elsevier.com/retrieve/pii/S2468233020300116. Accessed October 1, 2020.

Betts, J.C., Lukey, P.T., Robb, L.C., McAdam, R.A., and Duncan, K. (2002) Evaluation of a nutrient starvation model of Mycobacterium tuberculosis persistence by gene and protein expression profiling. *Molecular Microbiology* **43**: 717–731 https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2958.2002.02779.x. Accessed February 8, 2019.

Bhamidi, S., Shi, L., Chatterjee, D., Belisle, J.T., Crick, D.C., and McNeil, M.R. (2012) A bioanalytical method to determine the cell wall composition of Mycobacterium tuberculosis grown in vivo. *Analytical Biochemistry* **421**: 240–249 https://linkinghub.elsevier.com/retrieve/pii/S0003269711007123. Accessed February 5, 2020.

Bhaskara, G.B., Wong, M.M., and Verslues, P.E. (2019) The flip side of phospho-signalling: Regulation of protein dephosphorylation and the protein phosphatase 2Cs. *Plant Cell Environ* **42**: 2913–2930 https://onlinelibrary.wiley.com/doi/abs/10.1111/pce.13616. Accessed April 27, 2020.

Boitel, B., Ortiz-Lombardía, M., Durán, R., Pompeo, F., Cole, S.T., Cerveñansky, C., and
Alzari, P.M. (2003) PknB kinase activity is regulated by phosphorylation in two Thr
residues and dephosphorylation by PstP, the cognate phospho-Ser/Thr phosphatase, in
Mycobacterium tuberculosis: Mycobacterial serine/threonine kinase and phosphatase.
Molecular Microbiology **49**: 1493–1508 http://doi.wiley.com/10.1046/j.1365-
2958.2003.03657.x. Accessed February 6, 2019.

Boutte, C.C., Baer, C.E., Papavinasasundaram, K., Liu, W., Chase, M.R., Meniche, X., *et*
al. (2016) A cytoplasmic peptidoglycan amidase homologue controls mycobacterial cell
wall synthesis. *eLife* **5**: e14590 https://elifesciences.org/articles/14590. Accessed
September 19, 2019.

Bradshaw, N., Levdikov, V., Zimanyi, C., Gaudet, R., Wilkinson, A., and Losick, R.
(2017) A widespread family of serine/threonine protein phosphatases shares a common
regulatory switch with proteasomal proteases. *eLife* **6**.

Bradshaw, N., and Losick, R. (2015) Asymmetric division triggers cell-specific gene
expression through coupled capture and stabilization of a phosphatase. *eLife* **4**: e08145
https://elifesciences.org/articles/08145. Accessed September 19, 2019.

Carette, X., Platig, J., Young, D.C., Helmel, M., Young, A.T., Wang, Z., *et al.* (2018)
Multisystem Analysis of *Mycobacterium tuberculosis* Reveals Kinase-Dependent
Remodeling of the Pathogen-Environment Interface. *mBio* **9**
http://mbio.asm.org/lookup/doi/10.1128/mBio.02333-17. Accessed February 15, 2019.

Chopra, P., Singh, B., Singh, R., Vohra, R., Koul, A., Meena, L.S., *et al.* (2003)
Phosphoprotein phosphatase of Mycobacterium tuberculosis dephosphorylates serine-
threonine kinases PknA and PknB. *Biochem Biophys Res Commun* **311**: 112–120.

Cohen, P. (1989) The structure and regulation of protein phosphatases. *Annu Rev*
Biochem **58**: 453–508.

Cole, S.T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., *et al.* (1998)
Deciphering the biology of Mycobacterium tuberculosis from the complete genome
sequence. **396**: 27.

Cottin, V., Linden, A.V., and Riches, D.W.H. (1999) Phosphorylation of Tumor Necrosis
Factor Receptor CD120a (p55) by p42mapk/erk2 Induces Changes in Its Subcellular
Localization. 14.

Cunningham, A.F., and Spreadbury, C.L. (1998) Mycobacterial Stationary Phase Induced
by Low Oxygen Tension: Cell Wall Thickening and Localization of the 16-Kilodalton ꞷ-
Crystallin Homolog. *J BACTERIOL* **180**: 8.

DeJesus, M.A., Gerrick, E.R., Xu, W., Park, S.W., Long, J.E., Boutte, C.C., *et al.* (2017)
Comprehensive Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via

Saturating Transposon Mutagenesis. *mBio* **8**: e02133-16, /mbio/8/1/e02133-16.atom
https://mbio.asm.org/content/8/1/e02133-16. Accessed August 1, 2020.

Doerks, T., Noort, V. van, Minguez, P., and Bork, P. (2012) Annotation of the M.
tuberculosis Hypothetical Orfeome: Adding Functional Information to More than Half of
the Uncharacterized Proteins. *PLoS ONE* **7**: e34302
https://dx.plos.org/10.1371/journal.pone.0034302. Accessed November 14, 2021.

Ducret, A., Quardokus, E.M., and Brun, Y.V. (2016) MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. *Nature Microbiology* **1**: 16077
https://www.nature.com/articles/nmicrobiol201677. Accessed May 3, 2019.

Dulberger, C.L., Rubin, E.J., and Boutte, C.C. (2020) The mycobacterial cell envelope —
a moving target. *Nat Rev Microbiol* **18**: 47–59 http://www.nature.com/articles/s41579-
019-0273-7. Accessed May 20, 2021.

Durán, R., Villarino, A., Bellinzoni, M., Wehenkel, A., Fernandez, P., Boitel, B., *et al.*
(2005) Conserved autophosphorylation pattern in activation loops and juxtamembrane
regions of Mycobacterium tuberculosis Ser/Thr protein kinases. *Biochemical and*
Biophysical Research Communications **333**: 858–867
https://linkinghub.elsevier.com/retrieve/pii/S0006291X05011848. Accessed September
19, 2019.

Echenique, J., Kadioglu, A., Romao, S., Andrew, P.W., and Trombe, M.-C. (2004)
Protein Serine/Threonine Kinase StkP Positively Controls Virulence and Competence in
Streptococcus pneumoniae. Infect Immun **72**: 2434–2437
https://journals.asm.org/doi/10.1128/IAI.72.4.2434-2437.2004. Accessed November 15,
2021.

Fernandez, P., Saint-Joanis, B., Barilone, N., Jackson, M., Gicquel, B., Cole, S.T., and
Alzari, P.M. (2006) The Ser/Thr Protein Kinase PknB Is Essential for Sustaining
Mycobacterial Growth. *Journal of Bacteriology* **188**: 7778–7784
http://jb.asm.org/cgi/doi/10.1128/JB.00963-06. Accessed February 6, 2019.

Gee, C.L., Papavinasasundaram, K.G., Blair, S.R., Baer, C.E., Falick, A.M., King, D.S.,
et al. (2012) A Phosphorylated Pseudokinase Complex Controls Cell Wall Synthesis in
Mycobacteria. *Sci Signal* **5**: ra7–ra7 https://stke.sciencemag.org/content/5/208/ra7.
Accessed June 25, 2019.

Grundner, C., Gay, L.M., and Alber, T. (2005) *Mycobacterium tuberculosis*
serine/threonine kinases PknB, PknD, PknE, and PknF phosphorylate multiple FHA
domains. *Protein Sci* **14**: 1918–1921 http://doi.wiley.com/10.1110/ps.051413405.
Accessed September 19, 2019.

Gupta, M., Sajid, A., Arora, G., Tandon, V., and Singh, Y. (2009) Forkhead-associated
Domain-containing Protein Rv0019c and Polyketide-associated Protein PapA5, from
Substrates of Serine/Threonine Protein Kinase PknB to Interacting Proteins of
Mycobacterium tuberculosis. *Journal of Biological Chemistry* **284**: 34723–34734

637 https://linkinghub.elsevier.com/retrieve/pii/S0021925820375967. Accessed May 26,
638 2021.

639 Hannebelle, M.T.M., Ven, J.X.Y., Toniolo, C., Eskandarian, H.A., Vuaridel-Thurre, G.,
640 McKinney, J.D., and Fantner, G.E. (2020) A biphasic growth model for cell pole
641 elongation in mycobacteria. *Nat Commun* **11**: 452
642 http://www.nature.com/articles/s41467-019-14088-z. Accessed January 27, 2020.

643 Hett, E.C., and Rubin, E.J. (2008) Bacterial Growth and Cell Division: a Mycobacterial
644 Perspective. *Microbiology and Molecular Biology Reviews* **72**: 126–156
645 http://mmbr.asm.org/cgi/doi/10.1128/MMBR.00028-07. Accessed September 19, 2019.

646 Hoffmann, C., Leis, A., Niederweis, M., Plitzko, J.M., and Engelhardt, H. (2008)
647 Disclosure of the mycobacterial outer membrane: Cryo-electron tomography and vitreous
648 sections reveal the lipid bilayer structure. *Proceedings of the National Academy of*
649 *Sciences* **105**: 3963–3967 http://www.pnas.org/cgi/doi/10.1073/pnas.0709530105.
650 Accessed November 14, 2021.

651 Irmler, A., and Forchhammer, K. (2001) A PP2C-type phosphatase dephosphorylates the
652 PII signaling protein in the cyanobacterium Synechocystis PCC 6803. *Proceedings of the*
653 *National Academy of Sciences* **98**: 12978–12983
654 http://www.pnas.org/cgi/doi/10.1073/pnas.231254998. Accessed September 19, 2019.

655 Iswahyudi, Mukamolova, G.V., Straatman-Iwanowska, A.A., Allcock, N., Ajuh, P.,
656 Turapov, O., and O'Hare, H.M. (2019) Mycobacterial phosphatase PstP regulates global
657 serine threonine phosphorylation and cell division. *Sci Rep* **9**: 8337
658 http://www.nature.com/articles/s41598-019-44841-9. Accessed June 14, 2019.

659 Jani, C., Eoh, H., Lee, J., Hamasha, K., Sahana, M., Han, J.-S., *et al.* (2010) Regulation
660 of Polar Peptidoglycan Biosynthesis by Wag31 Phosphorylation in Mycobacteria. *BMC*
661 *Microbiol* **10**: 327 http://bmcmicrobiol.biomedcentral.com/articles/10.1186/1471-2180-
662 10-327. Accessed May 10, 2019.

663 Jarlier, V., and Nikaido, H. (1990) Permeability barrier to hydrophilic solutes in
664 Mycobacterium chelonei. *J Bacteriol* **172**: 1418–1423
665 https://journals.asm.org/doi/10.1128/jb.172.3.1418-1423.1990. Accessed November 14,
666 2021.

667 Jarlier, V., and Nikaido, H. (1994) Mycobacterial cell wall: Structure and role in natural
668 resistance to antibiotics. *FEMS Microbiology Letters* 11–18.

669 Juris, S.J., Rudolph, A.E., Huddler, D., Orth, K., and Dixon, J.E. (2000) A distinctive role
670 for the Yersinia protein kinase: Actin binding, kinase activation, and cytoskeleton
671 disruption. *Proceedings of the National Academy of Sciences* **97**: 9431–9436
672 http://www.pnas.org/cgi/doi/10.1073/pnas.170281997. Accessed November 15, 2021.

673 Kang, C.-M., Abbott, D.W., Park, S.T., Dascher, C.C., Cantley, L.C., and Husson, R.N.
674 (2005) The Mycobacterium tuberculosis serine/threonine kinases PknA and PknB:

substrate identification and regulation of cell shape. *Genes Dev* **19**: 1692–1704
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1176007/. Accessed July 15, 2019.

Kang, C.-M., Nyayapathy, S., Lee, J.-Y., Suh, J.-W., and Husson, R.N. (2008) Wag31, a
homologue of the cell division protein DivIVA, regulates growth, morphology and polar
cell wall synthesis in mycobacteria. *Microbiology* **154**: 725–735
http://mic.microbiologyresearch.org/content/journal/micro/10.1099/mic.0.2007/014076-
0. Accessed February 6, 2019.

Khan, S., Nagarajan, S.N., Parikh, A., Samantaray, S., Singh, A., Kumar, D., *et al.* (2010)
Phosphorylation of Enoyl-Acyl Carrier Protein Reductase InhA Impacts Mycobacterial
Growth and Survival. *Journal of Biological Chemistry* **285**: 37860–37871
http://www.jbc.org/lookup/doi/10.1074/jbc.M110.143131. Accessed February 6, 2019.

Kieser, K.J., Baranowski, C., Chao, M.C., Long, J.E., Sassetti, C.M., Waldor, M.K., *et al.*
(2015) Peptidoglycan synthesis in Mycobacterium tuberculosis is organized into
networks with varying drug susceptibility. *PNAS* **112**: 13087–13092
https://www.pnas.org/content/112/42/13087. Accessed March 12, 2019.

Kieser, K.J., and Rubin, E.J. (2014) How sisters grow apart: mycobacterial growth and
division. *Nature Reviews Microbiology* **12**: 550–562
http://www.nature.com/articles/nrmicro3299. Accessed February 6, 2019.

Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., *et al.* (2012) In Situ
Probing of Newly Synthesized Peptidoglycan in Live Bacteria with Fluorescent D -
Amino Acids. *Angewandte Chemie International Edition* **51**: 12519–12523
http://doi.wiley.com/10.1002/anie.201206749. Accessed February 6, 2019.

Kusebauch, U., Ortega, C., Ollodart, A., Rogers, R.S., Sherman, D.R., Moritz, R.L., and
Grundner, C. (2014) Mycobacterium tuberculosis supports protein tyrosine
phosphorylation. *Proceedings of the National Academy of Sciences* **111**: 9265–9270
http://www.pnas.org/cgi/doi/10.1073/pnas.1323894111. Accessed November 15, 2021.

Liu, Y., Tan, S., Huang, L., Abramovitch, R.B., Rohde, K.H., Zimmerman, M.D., *et al.*
(2016) Immune activation of the host cell induces drug tolerance in *Mycobacterium*
tuberculosis both in vitro and in vivo. *J Exp Med* **213**: 809–825
http://www.jem.org/lookup/doi/10.1084/jem.20151248. Accessed May 9, 2019.

Lu, G., and Wang, Y. (2008) FUNCTIONAL DIVERSITY OF MAMMALIAN TYPE
2C PROTEIN PHOSPHATASE ISOFORMS: NEW TALES FROM AN OLD FAMILY:
Function of PP2C isoforms. *Clinical and Experimental Pharmacology and Physiology*
35: 107–112 https://onlinelibrary.wiley.com/doi/10.1111/j.1440-1681.2007.04843.x.
Accessed November 19, 2021.

Marquardt, J.L., Siegele, D.A., Kolter, R., and Walsh, C.T. (1992) Cloning and
sequencing of Escherichia coli murZ and purification of its product, a UDP-N-
acetylglucosamine enolpyruvyl transferase. *J Bacteriol* **174**: 5748–5752

https://journals.asm.org/doi/10.1128/jb.174.17.5748-5752.1992. Accessed November 15,
2021.

Marrakchi, H., Lanéelle, M.-A., and Daffé, M. (2014) Mycolic Acids: Structures,
Biosynthesis, and Beyond. *Chemistry & Biology* **21**: 67–85
http://www.sciencedirect.com/science/article/pii/S1074552113004201. Accessed
February 6, 2019.

Molle, V., Brown, A.K., Besra, G.S., Cozzone, A.J., and Kremer, L. (2006) The
Condensing Activities of the Mycobacterium tuberculosis Type II Fatty Acid Synthase
Are Differentially Regulated by Phosphorylation. *Journal of Biological Chemistry* **281**:
30094–30103 https://linkinghub.elsevier.com/retrieve/pii/S0021925819339158. Accessed
May 26, 2021.

Molle, V., and Kremer, L. (2010) Division and cell envelope regulation by Ser/Thr
phosphorylation: *Mycobacterium* shows the way. *Molecular Microbiology* **75**: 1064–
1077 http://doi.wiley.com/10.1111/j.1365-2958.2009.07041.x. Accessed February 6,
2019.

Mougous, J.D., Gifford, C.A., Ramsdell, T.L., and Mekalanos, J.J. (2007) Threonine
phosphorylation post-translationally regulates protein secretion in Pseudomonas
aeruginosa. *Nat Cell Biol* **9**: 797–803 http://www.nature.com/articles/ncb1605. Accessed
November 19, 2021.

Nguyen, L. (2016) Antibiotic resistance mechanisms in M. tuberculosis: an update. *Arch*
Toxicol **90**: 1585–1604 http://link.springer.com/10.1007/s00204-016-1727-6. Accessed
September 19, 2019.

Pashley, C.A., and Parish, T. (2003) Efficient switching of mycobacteriophage L5-based
integrating plasmids in *Mycobacterium tuberculosis*. *FEMS Microbiology Letters* **229**:
211–215 https://academic.oup.com/femsle/article-lookup/doi/10.1016/S0378-
1097(03)00823-1. Accessed February 6, 2019.

Prisic, S., Dankwa, S., Schwartz, D., Chou, M.F., Locasale, J.W., Kang, C.-M., *et al.*
(2010) Extensive phosphorylation with overlapping specificity by Mycobacterium
tuberculosis serine/ threonine protein kinases. *Proceedings of the National Academy of*
Sciences **107**: 7521–7526.

Prisic, S., and Husson, R.N. (2014) Mycobacterium tuberculosis Serine/Threonine
Protein Kinases. *Microbiology Spectrum* **2**
http://www.asmscience.org/content/journal/microbiolspec/10.1128/microbiolspec.MGM2
-0006-2013. Accessed February 6, 2019.

Pullen, K.E., Ng, H.-L., Sung, P.-Y., Good, M.C., Smith, S.M., and Alber, T. (2004) An
Alternate Conformation and a Third Metal in PstP/Ppp, the M. tuberculosis PP2C-Family
Ser/Thr Protein Phosphatase. *Structure* **12**: 1947–1954
https://linkinghub.elsevier.com/retrieve/pii/S096921260400334X. Accessed May 26,
2021.

Roumestand, C., Leiba, J., Galophe, N., Margeat, E., Padilla, A., Bessin, Y., *et al.* (2011)
Structural Insight into the Mycobacterium tuberculosis Rv0020c Protein and Its
Interaction with the PknB Kinase. *Structure* **19**: 1525–1534
http://www.sciencedirect.com/science/article/pii/S096921261100253X. Accessed March
27, 2019.

Sajid, A., Arora, G., Gupta, M., Upadhyay, S., Nandicoori, V.K., and Singh, Y. (2011)
Phosphorylation of Mycobacterium tuberculosis Ser/Thr Phosphatase by PknA and PknB.
PLoS ONE **6**: e17871 https://dx.plos.org/10.1371/journal.pone.0017871. Accessed
December 9, 2021.

Sarathy, J., Dartois, V., Dick, T., and Gengenbacher, M. (2013) Reduced Drug Uptake in
Phenotypically Resistant Nutrient-Starved Nonreplicating Mycobacterium tuberculosis.
Antimicrobial Agents and Chemotherapy **57**: 1648–1653
http://aac.asm.org/lookup/doi/10.1128/AAC.02202-12. Accessed February 6, 2019.

Sarathy, J.P., Via, L.E., Weiner, D., Blanc, L., Boshoff, H., Eugenin, E.A., *et al.* (2017)
Extreme Drug Tolerance of *Mycobacterium tuberculosis* in Caseum. *Antimicrobial*
Agents and Chemotherapy **62** http://aac.asm.org/lookup/doi/10.1128/AAC.02266-17.
Accessed February 6, 2019.

Sassetti, C.M., and Rubin, E.J. (2003) Genetic requirements for mycobacterial survival
during infection. *Proceedings of the National Academy of Sciences* **100**: 12989–12994
http://www.pnas.org/cgi/doi/10.1073/pnas.2134250100. Accessed August 1, 2020.

Shamma, F., Quintanilla, S., Bandekar, A., Sassetti, C., and Boutte, C. (2021)
Phosphorylation on PstP Regulates Cell Wall Metabolism and Antibiotic Tolerance in
Mycobacterium smegmatis. *Journal of Bacteriology* **203**: e00563-20.

Sharma, A.K., Arora, D., Singh, L.K., Gangwal, A., Sajid, A., Molle, V., *et al.* (2016)
Serine/Threonine Protein Phosphatase PstP of *Mycobacterium tuberculosis* Is Necessary
for Accurate Cell Division and Survival of Pathogen. *Journal of Biological Chemistry*
291: 24215–24230 http://www.jbc.org/lookup/doi/10.1074/jbc.M116.754531. Accessed
February 6, 2019.

Sharma, K., Gupta, M., Krupa, A., Srinivasan, N., and Singh, Y. (2006) EmbR, a
regulatory protein with ATPase activity, is a substrate of multiple serine/threonine
kinases and phosphatase in *Mycobacterium tuberculosis*. *FEBS Journal* **273**: 2711–2721
http://doi.wiley.com/10.1111/j.1742-4658.2006.05289.x. Accessed May 26, 2021.

Slama, N., Leiba, J., Eynard, N., Daffé, M., Kremer, L., Quémard, A., and Molle, V.
(2011) Negative regulation by Ser/Thr phosphorylation of HadAB and HadBC
dehydratases from Mycobacterium tuberculosis type II fatty acid synthase system.
Biochemical and Biophysical Research Communications **412**: 401–406
https://linkinghub.elsevier.com/retrieve/pii/S0006291X11012770. Accessed March 31,
2019.

Thakur, M., and Chakraborti, P.K. (2006) GTPase Activity of Mycobacterial FtsZ Is
Impaired Due to Its Transphosphorylation by the Eukaryotic-type Ser/Thr Kinase, PknA.
J Biol Chem **281**: 40107–40113
http://www.jbc.org/lookup/doi/10.1074/jbc.M607216200. Accessed May 8, 2019.

Turapov, O., Forti, F., Kadhim, B., Ghisotti, D., Sassine, J., Straatman-Iwanowska, A., *et*
al. (2018) Two Faces of CwlM, an Essential PknB Substrate, in Mycobacterium
tuberculosis. *Cell Rep* **25**: 57-67.e5
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6180346/. Accessed July 22, 2020.

Veyron-Churlet, R., Zanella-Cléon, I., Cohen-Gonsaud, M., Molle, V., and Kremer, L.
(2010) Phosphorylation of the *Mycobacterium tuberculosis* β-Ketoacyl-Acyl Carrier
Protein Reductase MabA Regulates Mycolic Acid Biosynthesis. *Journal of Biological*
Chemistry **285**: 12714–12725 http://www.jbc.org/lookup/doi/10.1074/jbc.M110.105189.
Accessed February 6, 2019.

Vijay, K., Brody, M.S., Fredlund, E., and Price, C.W. (2000) A PP2C phosphatase
containing a PAS domain is required to convey signals of energy stress to the sB
transcription factor of Bacillus subtilis. *Molecular Microbiology* 9.

Vilchèze, C., Molle, V., Carrère-Kremer, S., Leiba, J., Mourey, L., Shenai, S., *et al.*
(2014) Phosphorylation of KasB Regulates Virulence and Acid-Fastness in
Mycobacterium tuberculosis. *PLoS Pathogens* **10**: e1004115
http://dx.plos.org/10.1371/journal.ppat.1004115. Accessed February 6, 2019.

Viswanathan, G., Joshi, S.V., Sridhar, A., Dutta, S., and Raghunand, T.R. (2015)
Identifying novel mycobacterial stress associated genes using a random mutagenesis
screen in Mycobacterium smegmatis. *Gene* **574**: 20–27
https://linkinghub.elsevier.com/retrieve/pii/S037811191500904X. Accessed November
11, 2021.

Viswanathan, G., Yadav, S., Joshi, S.V., and Raghunand, T.R. (2016) Insights into the
function of FhaA, a cell-division associated protein in mycobacteria. *FEMS Microbiology*
Letters fnw294 https://academic.oup.com/femsle/article-
lookup/doi/10.1093/femsle/fnw294. Accessed February 6, 2019.

Viswanathan, G., Yadav, S., and Raghunand, T.R. (2017) Identification of Mycobacterial
Genes Involved in Antibiotic Sensitivity: Implications for the Treatment of Tuberculosis
with β-Lactam-Containing Regimens. *Antimicrobial Agents and Chemotherapy* **61**
http://aac.asm.org/lookup/doi/10.1128/AAC.00425-17. Accessed February 6, 2019.

Wang, J., Li, C., Yang, H., Mushegian, A., and Jin, S. (1998) A Novel Serine/Threonine
Protein Kinase Homologue of *Pseudomonas aeruginosa* Is Specifically Inducible within
the Host Infection Site and Is Required for Full Virulence in Neutropenic Mice. *J*
Bacteriol **180**: 6764–6768 https://journals.asm.org/doi/10.1128/JB.180.24.6764-
6768.1998. Accessed November 15, 2021.

World Health Organization (2021) *Global tuberculosis report 2021*. World Health
Organization, Geneva. https://apps.who.int/iris/handle/10665/346387. Accessed
November 14, 2021.

Xie, Z., Siddiqi, N., and Rubin, E.J. (2005) Differential Antibiotic Susceptibilities of
Starved Mycobacterium tuberculosis Isolates. *Antimicrobial Agents and Chemotherapy*
49: 4778–4780 http://aac.asm.org/cgi/doi/10.1128/AAC.49.11.4778-4780.2005.
Accessed February 6, 2019.

PstP T137 paper 2022 Figures

838
839
840
Figure 1 Phosphorylation sites are not near the active site in PstP

Schematic of the crystal structure of PstP from *M. tuberculosis* (PstP$_{Mtb}$WT)
(Pullen *et al.*, 2004) (PDB code 1TXO) showing the conserved threonines (T)
which are phosphorylated by the kinases PknA and PknB (Sajid *et al.*, 2011).
Highlighted on the structure are: red - PstP$_{Mtb}$T137 (PstPT134 in PstP$_{Msmeg}$); blue
- PstP$_{Mtb}$T141 (PstPT138 in PstP$_{Msmeg}$); green - PstP$_{Mtb}$T174 (PstPT171 in
PstP$_{Msmeg}$); yellow - Mn^{2+} present at the conserved two-metal center typically
seen in the PP2C subfamily of protein Serine/Threonine phosphatases, which
are part of the active site core; pink - the unique third Mn^{2+} ion bound in the
catalytic core of PstP$_{Mtb}$ but not present in the human PP2Cα which sits in a
groove created by the flap subdomain next to the active site; orange - Ser60 and
Asp118 that act as ligands for the third Mn ion; lime green - residues of the
catalytic sites surrounding the two-metal center. Arrows point to position of the
OH-groups on T137, T141 and T174 that are phosphorylated.

FIGURE 2 Dephosphorylation of FhaA by PstP and phospho-mimetic isoforms

(a) Autoradiograms and corresponding Coomassie stained SDS-gels of *in vitro* phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-FhaA$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$137E$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no phosphatase control reaction (bottom panel). One set of representative images for each assay is shown here from two individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-FhaA$_{Mtb}$ on autoradiogram over amount of His-FhaA$_{Mtb}$ on the SDS-gel. P values were calculated using two-tailed unpaired t-test.

P-values of T137E at 0 min *vs.* T137E 80 min =0.7598, WT *vs.* T137E at 80 min= 0.0487, WT *vs.* T141E at 80 min =0.3712, WT *vs.* T174E at 80 min=0.2000. The error bars represent standard error of mean.

874
875
876 **FIGURE 3 Dephosphorylation of Wag31$_{Mtb}$ by PstP and phospho-mimetic**
877 **isoforms**
878

(a) Autoradiograms and their respective Coomassie stained SDS-gels of *in vitro*
phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-Wag31$_{Mtb}$
and His-PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$WT$_{Mtb}$ (second panel), His-
PstP$_c$T141E$_{Mtb}$ (third panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no
phosphatase control reaction (bottom panel). One set of representative images
for each reaction is shown here from two individual assays performed with
individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-Wag31$_{Mtb}$ on
autoradiogram over amount of His-Wag31$_{Mtb}$ on the SDS-gel. P values were
calculated using two-tailed unpaired t-test.

P-values of T137E at 0 sec *vs.* 15 sec = 0. 0.1947 and 0 sec *vs.* 120sec= 0.7493,
WT *vs.* T137E at 120 sec= 0.0130, WT *vs.* T141E at 120 sec =0.0106 and WT
vs. T174E at 120 sec = 0.170. The error bars represent standard error of mean.

FIGURE 4 Dephosphorylation of CwlM$_{Mtb}$ by PstP$_{Mtb}$ its phospho-mimetic isoforms and linear rate of dephosphorylation of CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$

(a) Autoradiograms and their respective Coomassie stained SDS-gels of *in vitro* phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-SUMO-CwlM$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no phosphatase control reaction (bottom panel). One set of representative images for each reaction is shown here from two individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-SUMO-CwlM$_{Mtb}$ on autoradiogram over amount of His-SUMO-CwlM$_{Mtb}$ on the SDS-gel. P values were calculated using two-tailed unpaired t-test. P-values of T137E at 0 min *vs.* 40 min= 0.1107, WT *vs.* T137E at 40 min= 0.0127, WT *vs.* T141E at 40 min=0.3683 and WT *vs.* T174E at 40 min= 0.5918. The error bars represent standard error of mean.

(c) Quantification of linear rate of change in [γ-^{32}P] signal on 10μg of His-SUMO-CwlM$_{Mtb}$, His-FhaA$_{Mtb}$ and His-Wag31$_{Mtb}$ in presence of His-PstP$_c$WT$_{Mtb}$ and its phospho-mimetic mutants per minute. P values were calculated using two-tailed unpaired t-test. **, P value = 0.0026 and *, P value 0306. The error bars represent standard error of mean.

FIGURE 5 Dephosphorylation of PknB_{Mtb} and PknA by PstPWT_{Mtb} and its phospho-mimetic variants

(a) Autoradiograms showing $[\gamma\text{-}^{32}P]$-phospho-signals on His-MBP-PknB_{Mtb} in the *in vitro* phosphatase assays with His-SUMO-CwlM_{Mtb} and His-PstP$_c$WT_{Mtb} (first panel), His-PstP$_c$WT_{Mtb} (second panel), His-PstP$_c$T141E_{Mtb} (third panel) and His-PstP$_c$T174E_{Mtb} (fourth panel), where His-MBP-PknB_{Mtb} was used to phosphorylate His-SUMO-CwlM_{Mtb}. One set of representative images for each reaction is shown here from two individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of $[\gamma\text{-}^{32}P]$ on His-MBP-PknB_{Mtb} on autoradiogram. P values were calculated using two-tailed unpaired t-test. P value of T137E at 0 min *vs.* 40 min= 0.0523. *, P value= 0.0375 (WT *vs.*T137E at 40 min); *, P value=0.0314 (T141E *vs.*T137E at 40 min); *, P value= 0.0217 (T174E *vs.*T137E at 40 min). The error bars represent standard error of mean.

(c) Autoradiograms showing [γ-^{32}P]-phospho-signals on His-MBP-PknA$_{Mtb}$ in the *in*
vitro phosphatase assays with His-Wag31$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel),
His-PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel) and His-
PstP$_c$T174E$_{Mtb}$ (fourth panel), where His-Wag31$_{Mtb}$ was first phosphorylated with
His-MBP-PknA$_{Mtb}$. One set of representative images for each reaction is shown
here from two individual assays performed with individually purified batches of
phosphatases.

(d) Quantification of relative intensities of [γ-^{32}P] on His-MBP-PknA$_{Mtb}$ on
autoradiogram. P values were calculated using two-tailed unpaired t-test. P value
of T137E at 0 sec *vs.* 120 sec= 0.0643. *, P value= 0.0163 (WT *vs.* T137E at 120
sec). The error bars represent standard error of mean.

943
944

945
Figure 6 The phosphosite on T134 on PstP$_{Msmeg}$ affects PG metabolism

(a) Doubling times of *Msmeg* strains carrying the *pstP_Msmeg* WT and the phospho-
mimetic T134E alleles. Each bar represents the mean of individual average
doubling times of three biological replicates of each genotype from two different
experiments on different dates. The error bars represent mean with standard
deviation. **, P value= 0.0026.

(b) Quantification of cell lengths of isogenic *pstP_Msmeg* allele strains (WT and
T134E) grown in 7H9 in log. phase and starved in PBS with Tween80. At least
100 cells from each of three biological replicates were quantified. P values were
calculated by unpaired t-test. *, P value = 0.0122, ****, P value= 0.000000031.

(c) Quantification of mean intensities of HADA of *pstP_Msmeg* allele strains (WT and
T134E) grown in 7H9 in log. phase and starved in PBS with Tween80. Cells were
stained after 15 minutes of starvation and washed and fixed after another 15
minutes (starved for a total of 30 minutes). At least 100 cells from each of three
biological replicates were quantified. P values were calculated by unpaired t-test.
*, P value = 0.0109, ****, P value= <0.000000000000001.

(d) and (e) Representative HADA and phase images of cells from *pstP_Msmeg* allele
strains (WT and T134E) grown in 7H9 in log. phase (d) and after 30 minutes of
starvation in PBS with Tween80 (e).

(f) and (g) Intensity profiles of HADA signals in cells of *pstP_Msmeg* allele strains
(WT and T134E) in log. phase (f) and after starving for 30 minutes in PBS
supplemented with Tween80 (g). The shaded regions denote standard deviations
and the solid lines represent the mean intensity values. Signal intensities from at
least 110 cells from each of three biological replicates of every *pstP_Msmeg* allelic
variant genotype were quantified and used to analyze the signal profiles in
MicrobeJ. 0 to 1 on the X-axes represent cells from the old pole to the new pole,
respectively.

Figure 7 Localization of PstP WT _Msmeg_

Msmeg cells expressing mCherry2B-PstP were imaged over time in a CellAsic
Microfluidic device at 15-minute intervals. Intensity is seen at the midcell (white

arrows) for several frames before cell separation can be visualized by phase
contrast microscopy (yellow arrows).

984 **Supplemental Figures:**

985
986

987 **Supplementary Figure S1: Dephosphorylation of His-Wag31$_{Mtb}$ by His-**
988 **PstP$_c$WT$_{Mtb}$.** Autoradiogram and the respective Coomassie stained SDS-gel of *in*
989 *vitro* phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-
990 Wag31$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$. Ten times more substrate than the phosphatase
991 was used in the assay.

993
994

Supplementary Figure S2: Demographic profiles of HADA signal intensity in *pstP_Msmeg* strains in growth and starvation. (a) and (b) Demographs showing intensities of the fluorescent dye HADA signal in individual cells from *pstP_Msmeg*WT allele strains (a) and *pstP_Msmeg*T134E allele strains (b) in log. phase. (c) and (d) Demographs showing intensities of the fluorescent dye HADA signal in individual cells from *pstP_Msmeg*WT allele strains (c) and *pstP_Msmeg*T134E allele strains (d) starved in PBS with Tween80 for a total time period of 30 minutes.

Biological triplicates of each *pstP_Msmeg* allelic variant were analyzed. Signal intensities from at least 340 cells from each *pstP* allelic variant (at least 100 cells from each biological triplicate of each genotype) in log. phase and starvation were plotted in the demographs.