Mycobacterial Serine/Threonine phosphatase PstP is phospho-regulated and localized to mediate control of cell wall metabolism

Farah Shamma[1], E. Hesper Rego[2] and Cara C. Boutte [1*]

[1]Department of Biology, University of Texas at Arlington, Arlington, Texas

[2] Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, Connecticut

*Corresponding author, cara.boutte@uta.edu

ABSTRACT

The mycobacterial cell wall is profoundly regulated in response to environmental stresses, and this regulation contributes to antibiotic tolerance. The reversible phosphorylation of different cell wall regulatory proteins is a major mechanism of cell wall regulation. Eleven Serine/Threonine protein kinases (STPKs) phosphorylate many critical cell wall-related proteins in mycobacteria. PstP is the sole serine/ threonine phosphatase, but few proteins have been verified as PstP substrates. PstP is itself phosphorylated but the role of its phosphorylation in regulating its activity has been unclear. In this study we aim to discover novel substrates of PstP in *Mycobacterium tuberculosis* (*Mtb*). We show *in vitro* that PstP dephosphorylates two regulators of peptidoglycan in *Mtb*, FhaA and Wag31. We also show that a phospho-mimetic mutation of T137 on PstP negatively regulates its catalytic activity against the cell wall regulators FhaA, Wag31, CwlM, PknB and PknA, and that the corresponding mutation in *Mycobacterium smegmatis (Msmeg)* causes mis-regulation of peptidoglycan *in vivo*. We show that PstP is localized to the septum, which likely restricts its access to certain substrates. These findings on the regulation of PstP provide insight into the control of cell wall metabolism in mycobacteria.

INTRODUCTION

Mycobacterium tuberculosis (*Mtb*), which causes Tuberculosis (TB), remains a public health menace, infecting about 10 million and killing at least a million people worldwide each year (World Health Organization, 2021). TB infections require extended multi-drug antibiotic treatment regimens (Nguyen, 2016). The mycobacterial cell wall contributes to *Mtb*'s inherent antibiotic tolerance (Jarlier and Nikaido, 1994; Sarathy *et al.*, 2013; Batt *et al.*, 2020). The wall serves as permeability barrier (Jarlier and Nikaido, 1990; Hett and Rubin, 2008; Hoffmann *et al.*, 2008; Sarathy *et al.*, 2013), and is strongly regulated in response to stresses (Cunningham and Spreadbury, 1998; Betts *et al.*, 2002; Bhamidi *et al.*, 2012; Sarathy *et al.*, 2013), including infection (Sharma *et al.*, 2006). The cell

wall changes in stress are associated with increased antibiotic tolerance (Xie *et*
al., 2005; Liu *et al.*, 2016; Sarathy *et al.*, 2017). The expression and activity of
many cell wall enzymes and regulators are also tightly coordinated over the
course of the cell cycle in order to maintain cell wall integrity during elongation
and division (Kieser and Rubin, 2014; Dulberger *et al.*, 2020; Bandekar *et al.*,
2020).

The cell wall consists of three covalently linked layers: a peptidoglycan layer
surrounding the plasma membrane, an arabinogalactan layer and a mycolic acid
layer (Marrakchi *et al.*, 2014). Peptidoglycan is a network of glycan chains cross-
linked by small peptides, and is the innermost layer of the wall (Abrahams and
Besra, 2021). Reversible Serine/Threonine (S/T) phosphorylation of cell wall
enzymes and regulators is critical for regulation of cell wall metabolism in
response to environmental signals (Wang *et al.*, 1998; Juris *et al.*, 2000;
Echenique *et al.*, 2004). There are 11 Serine/Threonine kinases (STPKs) (PknA-
B and PknD-L) and one essential S/T protein phosphatase PstP in *Mtb* (Kang *et*
al., 2005; Gee *et al.*, 2012; Baer *et al.*, 2014; Boutte *et al.*, 2016)(Cole *et al.*,
1998; Bach *et al.*, 2009). PstP is also essential in *Msmeg* (Sharma *et al.*, 2016;
DeJesus *et al.*, 2017).

PstP belongs to the protein phosphatase 2C (PP2C) subfamily of divalent metal-
ion dependent protein serine/threonine phosphatases (Cohen, 1989; Barford,
1996; Chopra *et al.*, 2003), which regulate prokaryotic and eukaryotic cell growth
and division, sporulation, stress response and metabolic processes in response
to various environmental signals (Vijay *et al.*, 2000:2; Irmler and Forchhammer,
2001; Mougous *et al.*, 2007; Lu and Wang, 2008; Bradshaw and Losick, 2015;
Bradshaw *et al.*, 2017). PstP$_{Mtb}$ shares structural architecture and conserved
catalytic residues with the human PP2Cα (Pullen *et al.*, 2004) (Figure 1), the
PP2C family prototype. The N-terminal cytoplasmic enzymatic domain of PstP$_{Mtb}$
(Figure 1) connects to its C-terminal extracellular domain via a transmembrane
helix (Pullen *et al.*, 2004). PstP$_{Mtb}$ is itself phosphorylated on T137, T141, T174
and T290 (Figure 1 (Sajid *et al.*, 2011). Phosphorylation on all these sites

together affects PstP activity against small molecule substrates *in vitro* (Sajid *et*
al., 2011). And phosphorylation on T174 affects cell wall metabolism *in vivo* in
Msmeg (Shamma *et al.*, 2021)*.*

As PstP is the only mycobacterial S/T protein phosphatase (Cole *et al.*, 1998), it
likely dephosphorylates many of the substrates phosphorylated by the 11 STPKs
(Prisic and Husson, 2014). However, only a few proteins are known PstP
substrates: the STPKs PknA and PknB (Boitel *et al.*, 2003; Chopra *et al.*, 2003;
Durán *et al.*, 2005; Sajid *et al.*, 2011), mycolic acid biosynthesis enzymes KasA
and KasB (Molle *et al.*, 2006), the arabinogalactan regulator EmbR (Sharma *et*
al., 2006), and CwlM (Shamma *et al.*, 2021), which in its phosphorylated form is
an essential activator of MurA (Boutte *et al.*, 2016), the first enzyme of
peptidoglycan precursor synthesis (Marquardt *et al.*, 1992). So
dephosphorylation of CwlM~P by PstP (Shamma *et al.*, 2021) should halt
peptidoglycan synthesis. CwlM~P also interacts with FhaA (Turapov *et al.*, 2018),
which is involved in modulating peptidoglycan biosynthesis (Gee *et al.*, 2012;
Viswanathan *et al.*, 2015; Viswanathan *et al.*, 2017) and unphosphorylated CwlM
interacts with the peptidoglycan precursor lipid-II flippase MurJ (Turapov *et al.*,
2018), suggesting that CwlM may regulate multiple steps in peptidoglycan
synthesis (Turapov *et al.*, 2018).

FhaA and Wag31 are both regulators of mycobacterial cell wall metabolism.
FhaA helps maintain cell envelope integrity, cell length and antibiotic tolerance
(Viswanathan *et al.*, 2015; Viswanathan *et al.*, 2017). FhaA stabilizes PbpA
(Viswanathan *et al.*, 2017) and binds and may inhibit the MurJ flippase (Gee *et*
al., 2012). FhaA is phosphorylated by PknB (Grundner *et al.*, 2005; Prisic *et al.*,
2010; Roumestand *et al.*, 2011), though the role of its phosphorylation is not yet
clear. Wag31 localizes at the cell poles (Kang *et al.*, 2005), is essential for polar
peptidoglycan synthesis (Kang *et al.*, 2008; Jani *et al.*, 2010) and is
phosphorylated by the STPK PknA (Kang *et al.*, 2005). Wag31's phosphorylation
may also have a role in controlling peptidoglycan metabolism (Jani *et al.*, 2010).

It was not known whether FhaA and Wag31 were actively dephosphorylated, or if

their phosphorylated forms were removed through protein degradation.

In this study we show that FhaA$_{Mtb}$ and Wag31$_{Mtb}$ are substrates of PstP$_{Mtb}$ *in*

vitro, and that PstP dephosphorylates Wag31 much faster than it does FhaA or

CwlM. We show that the T137E phospho-mimetic mutation of PstP$_{Mtb}$ inhibits

PstP$_{Mtb}$'s catalytic activity against the substrates CwlM, FhaA, Wag31, PknB and

PknA *in vitro*. Furthermore, the corresponding phospho-mimetic mutation in

Msmeg leads to misregulation of peptidoglycan metabolism. We also find that

PstP is enriched at the cell division site, suggesting that its access to substrates

might also control its activity. We suggest a model where PstP's activity against

assorted substrates is regulated by substrate affinity and protein localization, and

further toggled by the phosphorylations on PstP.

RESULTS

PstP$_{Mtb}$ dephosphorylates FhaA$_{Mtb}$ and Wag31$_{Mtb}$ *in vitro*

FhaA$_{Mtb}$ is phosphorylated by PknB$_{Mtb}$ *in vivo* (Prisic *et al.*, 2010) and *in vitro*

(Grundner *et al.*, 2005; Roumestand *et al.*, 2011) on T116 (Roumestand *et al.*,

2011). Wag31$_{Mtb}$ is phosphorylated at T73 by PknA$_{Mtb}$ (Kang *et al.*, 2005). To

determine if FhaA$_{Mtb}$ and Wag31$_{Mtb}$ are substrates of PstP$_{Mtb}$, we purified full-

length His-FhaA$_{Mtb}$, His-Wag31$_{Mtb}$ and phosphorylated them by by His-MBP

fusions to the kinase domains of PknA$_{Mtb}$ and PknB$_{Mtb}$. We then added the

cytosolic domain of PstP$_{Mtb}$ containing the catalytic domain (His-PstP$_c$WT$_{Mtb}$) and

measured changes in phosphorylation. All the proteins in this study were purified

first by nickel column and then by size exclusion chromatography, to ensure

proper folding. In all the kinase reaction mixtures in this study, the molar

concentrations of the substrates were in excess over the ATP [γ-^{32}P], so that

there is no leftover ATP [γ-^{32}P] to allow re-phosphorylation once His-PstP$_c$ is

added to the reaction mixture for the phosphatase assay. This experimental

setup allows us to measure dephosphorylation in the presence of the kinases

without re-phosphorylation happening simultaneously.

We phosphorylated His-FhaA$_{Mtb}$ with His-MBP-PknB$_{Mtb}$ and ATP [γ-^{32}P]. After

addition of His-PstP$_c$WT$_{Mtb}$, at 1/10th the concentration of the His-FhaA$_{Mtb,}$ the

phosphorylation on His-FhaA$_{Mtb}$~P decreased by ~80% in 80 minutes (Figure

2a,b). Our control assay with no phosphatase shows that the phosphorylation on

His-FhaA$_{Mtb}$~P is stable over time in the phosphatase buffer (Figure 2a, b). We

then phosphorylated His-Wag31$_{Mtb}$ with His-MBP-PknA$_{Mtb}$ and ATP [γ-^{32}P], and

added His-PstP$_c$WT$_{Mtb}$. With 1/100th the amount of PstP compared to Wag31,

Wag31$_{Mtb}$ phosphorylation decreased by about ~90% in 2 minutes (Figure 3a,b),

only in the presence of His-PstP$_c$WT$_{Mtb}$. These data show that FhaA$_{Mtb}$ and

Wag31$_{Mtb}$ are substrates of PstPWT$_{Mtb}$ *in vitro*.

Our previous data that demonstrated PstP$_{Mtb}$ dephosphorylates CwlM$_{Mtb}$

(Shamma *et al.*, 2021) is reproducible in our *in vitro* assays with ATP [γ-^{32}P] where

we observe a 97% decrease in phosphorylation on His-SUMO-CwlM$_{Mtb}$~P 40 minutes

after addition of 1/10th the amount of His-PstP$_c$WT$_{Mtb}$ (Figure 4a, b). In all the assays,

we see no phosphorylation signals on the WT and phospho-mimetic PstPs on the

autoradiograms (Supplementary figure S3). This is likely because PstP can

quickly dephosphorylate itself (Sajid *et al.*, 2011) and the purified forms are thus

unphosphorylated.

Because measuring the rates of dephosphorylation by PstP against different

substrates required different PstP concentrations and timing, we calculated the

initial linear rate of dephosphorylation per unit concentration of PstP, in order to

compare PstP's activity against different substrates. This comparison indicates

that His-PstP$_c$WT$_{Mtb}$ dephosphorylates His-Wag31$_{Mtb}$ more than 500 times faster

than His-SUMO-CwlM$_{Mtb}$ and about 2500 times faster than His-FhaA$_{Mtb}$ (Figure

4c).

Phospho-mimetic mutation at PstPT137$_{Mtb}$ inhibits phosphatase activity

PstP$_{Mtb}$ is phosphorylated on the conserved threonine (T) residues 137, 141 and

174 (Figure 1), and when phosphorylated, shows increased activity against small

molecule substrates *in vitro* (Sajid *et al.*, 2011). We hypothesized that

phosphorylation of the threonine residues of PstP individually might help regulate

PstP's activity by either affecting its catalytic activity broadly, or by specifically

affecting substrate affinity or activity against specific substrates. To test this, we

made phospho-mimetic (T to E) (Cottin *et al.*, 1999) mutants of $PstP_{Mtb}$ at each of

the three conserved phosphorylation sites of $PstP_{Mtb}$ in the catalytic domain and

purified them (His-$PstP_c$T137E$_{Mtb}$, His-$PstP_c$T141E$_{Mtb}$ and His-$PstP_c$T174E$_{Mtb}$).

We previously showed that *in vitro* His-$PstP_c$WT$_{Mtb}$ and His-$PstP_c$T174E$_{Mtb}$

dephosphorylate His-SUMO-CwlM$_{Mtb}$ at the same rate (Shamma *et al.*, 2021).

We performed separate *in vitro* phosphatase assays with the substrates His-

FhaA$_{Mtb}$, His-Wag31$_{Mtb}$ and His-SUMO-CwlM$_{Mtb}$ with the phospho-mimetic

versions of PstP. All three substrates exhibit stable phosphorylation with His-

$PstP_c$T137E$_{Mtb}$, indicating that it is inactive *in vitro* against all three substrates

(Figures 2,3,4). This suggests that phosphorylation on T137 of PstP may inhibit

its activity against multiple substrates.

His-MBP-PknB$_{Mtb}$ and His-MBP-PknA$_{Mtb}$, the STPKs used in these assays, are

also confirmed substrates of PstP (Sajid *et al.*, 2011). PknA and PknB are

activated to phosphorylate their own substrates when they themselves are

phosphorylated at their activation loops (Baer *et al.*, 2014). Because we already

had these kinases in our phosphatase assays, we were able to measure

dephosphorylation of His-MBP-PknB$_{Mtb}$ and His-MBP-PknA$_{Mtb}$ by PstP in the

context of a 10-fold and 5-fold excess of His-SUMO-CwlM$_{Mtb}$ or His-Wag31$_{Mtb}$,

respectively. We found that His-$PstP_c$T137E$_{Mtb}$ dephosphorylates both His-MBP-

PknB$_{Mtb}$ (Figure 5a,b) and His-MBP-PknA$_{Mtb}$ (Figure 5c,d) more slowly than His-

$PstP_c$WT$_{Mtb}$ and the other PstP mutants. This shows that His-$PstP_c$T137E$_{Mtb}$ is

inhibited against at least five different substrates.

Phospho-mimetic mutations at PstPT141$_{Mtb}$ and PstPT174$_{Mtb}$ moderately

inhibit Wag31's dephosphorylation

His-PstP$_c$T141E$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$ dephosphorylate His-FhaA$_{Mtb}$ (Figure

2,4c) and His-SUMO-CwlM$_{Mtb}$ (Figure 4) at about the same rate as His-

PstP$_c$WT$_{Mtb}$, but dephosphorylate His-Wag31$_{Mtb}$ about half as fast as His-

PstP$_c$WT$_{Mtb}$ (Figure 3,4c). These biochemical data support our hypothesis that

phosphorylation on PstP may modulate its substrate specificity. Phosphorylation

at T141 and T174 appear to inhibit binding or catalysis of Wag31 but not FhaA or

CwlM.

A negative charge on T134 of PstP$_{Msmeg}$ regulates peptidoglycan

metabolism in starvation

Phosphorylation of CwlM allows it to activate the peptidoglycan precursor

enzyme MurA, thereby licensing peptidoglycan synthesis (Boutte *et al.*, 2016).

Phosphorylation of CwlM also affects its binding to FhaA and the lipid II flippase

MurJ (Turapov *et al.*, 2018). CwlM is quickly dephosphorylated in the transition to

stasis (Boutte *et al.*, 2016) likely by PstP (Shamma *et al.*, 2021). The function of

phosphorylation on FhaA and Wag31 is not yet clear, but available data (Jani *et*

al., 2010; Gee *et al.*, 2012; Viswanathan *et al.*, 2016) suggest it could also affect

peptidoglycan metabolism. Dephosphorylation of these regulators could be

important for downregulation of peptidoglycan metabolism in stasis.

As the phospho-mimetic mutation at PstPT137$_{Mtb}$ inhibits PstP's

dephosphorylation of three peptidoglycan regulatory proteins *in vitro* (Figure

2,3,4), we sought to determine if the corresponding phospho-site on PstP (T134)

might affect peptidoglycan metabolism *in vivo* in *Msmeg*. Although this site is not

known to be phosphorylated in PstP$_{Msmeg}$ (Iswahyudi *et al.*, 2019; Le *et al.*, 2020),

the PstP proteins are 78% identical between *Msmeg* and *Mtb* and most of PstP's

substrates are also conserved, so we reasoned that performing experiments with

a negative charge at residue T134 on PstP in *Msmeg* will provide a preliminary

approximation of what the effect of phosphorylation on this residue might be in

Mtb. We made *Msmeg* strains with a phospho-mimetic *pstP*T134E*Msmeg* allele and

a phospho-ablative *pstP*T134A*Msmeg* allele replacing the wild-type *pstP*. We

verified the expression of PstPT134E*Msmeg* and T134A in these strains under

normal growth condition (Figure 6h). We found that PstPT134E*Msmeg* was stable

but PstPT134A*Msmeg* was not, so we did not use the *pstP*T134A*Msmeg* allele strains

for further experiments. We reported previously that the biological replicates of

*pstP*T134E*Msmeg* mutant strains had inconsistent doubling times (Shamma *et al.*,

2021) probably due to potential suppressor mutations. We confirmed that the

slow growing clones could, after passaging, accumulate suppressor mutations

that allowed them to grow as fast as the *pstP*WT*Msmeg* strain. To characterize the

original *pstP*T134E*Msmeg* strain, we isolated numerous clones from fresh

transformations, and measured the doubling times of each before continuing with

experiments. We show data only from clones that grew at the slower rate.

We stained both log. phase and starved *pstP*T134E*Msmeg* and WT*Msmeg* strains

using a fluorescent D-amino acid HADA (Kuru *et al.*, 2012) as a reporter of active

peptidoglycan metabolism. We saw a slight increase in length (Figure 6b,d and

Supplementary Figure S2a,b) and HADA intensity (Figure 6c,d and

Supplementary Figure S2a,b) in the *pstP*T134E*Msmeg* strains in the log. phase

compared to the WT*Msmeg* strains. After starvation, these differences increased

(Figure 6b,c,e and Supplementary Figure S2c,d). WT*Msmeg* strains quickly

downregulate peptidoglycan metabolism upon starvation (Boutte *et al.*, 2016), as

the decreased HADA staining (Figure 6c,d,e,f,g) shows. The starved

*pstP*T134E*Msmeg* strains show significantly higher HADA signal than the WT*Msmeg*

(Figure 6c, top right panels of d,e and Supplementary Figure S2b,d) indicating a

defect in downregulating peptidoglycan metabolism. This suggests that the

presence of a constant negative charge on phosphosite T134 of PstP*Msmeg*

impairs downregulation of peptidoglycan synthesis, likely by inhibiting PstP's

dephosphorylation of peptidoglycan regulators (Figure 2,3,4).

The mis-regulation of peptidoglycan metabolism (Figure 6c,d,e,f,g and

Supplementary Figure S2b,d) may account for the slower growth of the

*pstP*T134E*Msmeg* strains that we see in our data (Figure 6a), though it is likely that

other processes beyond peptidoglycan metabolism are also misregulated, as

PstP likely has many substrates involved in diverse processes (Molle *et al.*, 2006;

Iswahyudi *et al.*, 2019).

PstP localizes at the septum in exponentially growing *Msmeg* cells

We sought to determine if PstP's activity against its diverse substrates could be

regulated through protein localization. We made an *Msmeg* strain carrying an

additional copy of *pstPWT*$_{Msmeg}$ fused to *mCherry2B*. By time-lapse fluorescence

microscopy, we observed that mCherry2B-PstPWT$_{Msmeg}$ appears to be uniformly

distributed along the membrane throughout most of the cell cycle, but becomes

localized at the septum before cell separation (Figure 7). These data suggest that

PstP is also regulated by its access to substrates at different sites in the cell.

Discussion

Our findings indicate that PstP has different basal activity rates against different

substrates (Figure 4c), is regulated by its own phosphorylation (Figure 2,3,4,5) to

regulate cell wall metabolism (Figure 6), and that it may also be regulated by its

localization (Figure 7).

We see that PstPWT$_{Mtb}$ dephosphorylates CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$ at

different rates *in vitro* (Figure 4c), with Wag31$_{Mtb}$ being the fastest to be

dephosphorylated and FhaA the slowest (Figure 4c). We showed that PstP

localizes at the septum, or cell division site, in the exponentially growing *Msmeg*

cells (Figure 7). Both FhaA and Wag31 localize at the poles and septum (Gee *et*

al., 2012; Hannebelle *et al.*, 2020). Our localization data (Figure 7) suggest that

co-localization of PstP with these substrates at the septum may cause both

Wag31 and FhaA (Figure 2, 3, 4c) to be more highly phosphorylated at the cell

poles than at the septum. As PstP dephosphorylates Wag31 so rapidly (Figure 3,

4c), we hypothesize that Wag31 is likely to be largely unphosphorylated at the

septum, and may only be phosphorylated at the poles.

Our data in Figure 2, 3 and 4 show that negative charges at PstPT141$_{Mtb}$ and

T174$_{Mtb}$ slow dephosphorylation of Wag31$_{Mtb}$, but not FhaA$_{Mtb}$ and CwlM$_{Mtb}$.

Phosphorylation site T141 is on the flap subdomain and T174 is on the β-sheet

core (β8) on PstP$_{Mtb}$ (Pullen *et al.*, 2004), but both are facing into the same

groove between these domains (Figure 1). T137 is the first residue of the flap

subdomain (Pullen *et al.*, 2004), and it faces away from the groove with the other

phospho-sites (Figure 1). We propose that the T141/ T174 groove may contribute

to toggling specificity of PstP against certain substrates, depending on the

phosphorylation status.

A previous study showed that PstP$_{cMtb}$, phosphorylated by co-expression with

PknA$_{Mtb}$ or PknB$_{Mtb}$, showed higher activity than unphosphorylated PstP against

a non-biological small molecule substrate *in vitro*, while PstP$_c$T141E$_{Mtb}$ showed

the same activity as the wild-type against the small molecule (Sajid *et al.*, 2011).

However, the status of phosphorylation of the other individual phospho-sites on

PstP$_{Mtb}$ was not examined in that study. Our study, in contrast, sheds light on the

effect of individual phospho-sites against biological protein substrates. Our

findings (Figure 2,3, and 4) show that the phospho-mimetic PstP$_c$T137E$_{Mtb}$ has

an inhibitory effect whereas the phospho-mimetic variants PstP$_c$T141E$_{Mtb}$ and

PstP$_c$T174E$_{Mtb}$ play a role in substrate specificity.

Our *in vitro* data in Figure 2, 3, 4 and 5 show that PstPT137E$_{Mtb}$ inhibits its

activity against CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$, and also against the STPKs

PknA and PknB that phosphorylate them, all of which are involved in regulation

of peptidoglycan metabolism (Thakur and Chakraborti, 2006; Kang *et al.*, 2008;

Jani *et al.*, 2010; Gee *et al.*, 2012; Prisic and Husson, 2014; Boutte *et al.*, 2016;

Viswanathan *et al.*, 2016; Viswanathan *et al.*, 2017; Carette *et al.*, 2018; Turapov

et al., 2018). The corresponding residue in PstP$_{Msmeg}$ is T134. Although this

residue may not be phosphorylated in PstP$_{Msmeg}$ (Iswahyudi *et al.*, 2019; Le *et al.*,

2020), we sought to determine if a negative charge at this site would still affect

peptidoglycan metabolism in *Msmeg*, as we would expect from the changes in

dephosphorylation of the peptidoglycan regulators from *Mtb.* Our *in vivo* data

show that the *pstP*T134E$_{Msmeg}$ strains grow more slowly, and have an increase in

cell length and HADA staining, especially in starvation (Figure 6). This suggests

that dephosphorylation of CwlM, FhaA, Wag31, PknB or PknA, is important in the

transition to stasis and that a negative charge at the T134$_{Msmeg}$/T137$_{Mtb}$ site would

inhibit the ability of PstP to properly regulate the cell wall through these and other

substrates. In this study, we focus on the residues that are phosphorylated in the

PstP protein from *Mtb* (Sajid *et al.*, 2011); while the experiments in *Msmeg* do not

predict what effects phosphorylation on PstPT137 would have in *Mtb*, they at

least indicate that some effect on peptidoglycan metabolism is probable.

Other studies have shown that PstP is an important regulator of cell division and

cell length (Sharma *et al.*, 2016). Our work shows that PstP localizes to the cell

division site, which corroborates its importance in septation (Figure 7). We also

show that the PstPT134E$_{Msmeg}$ mutation affects cell length (Figure 6b), which

could be mediated by changes in division or elongation.

Although our findings are limited to PstP's activity on certain substrates only, they

show that the individual phosphosites are important in regulating PstP. *In vitro*,

we see that a negative charge on T137 of PstP$_{Mtb}$ inhibits its activity against three

PG regulators (Figure 2,3,4), as well as the essential kinases PknB and PknA

when they are mixed together (Figure 5). In the cell, the substrates and kinases

also co-exist, and PstP may exert its effect directly by dephosphorylating the

substrates or indirectly by acting on the kinase (Bhaskara *et al.*, 2019), or both

simultaneously. There are likely other mechanisms of cell wall regulation involved

in the complex *in vivo* scenario like protein localization or PstP's interaction with

other regulatory proteins that may contribute to PstP's regulation as well. We do

not know the conditions under which PstP is phosphorylated, or the combinations

of PstP's phosphorylations that exist in the cell. So the *in vitro* data only gives us

a hint of the deep complexity of the STPK-substrate-PstP network *in vivo*.

In this work we show that PstP is regulated by both phosphorylation and

localization. This work demonstrates the importance of protein

dephosphorylation, not just phosphorylation, in regulating the activity of proteins

involved in cell wall metabolism in mycobacteria. Future work that addresses

when all of these important phospho-sites are phosphorylated and

dephosphorylated will help us to understand how all these protein regulators

contribute to coordinating the construction and integrity of the cell wall in

changing conditions.

MATERIALS AND METHODS

Bacterial strains and culture conditions

All *Mycobacterium smegmatis* mc^{2}155 ATCC 700084 cultures were grown in 7H9

(Becton, Dickinson, Franklin Lakes, NJ) medium containing 0.003 g/liter catalase,

5 g/liter bovine serum albumin (BSA), 2 g/liter dextrose, 0.2% glycerol, 0.85

g/liter NaCl, and 0.05% Tween 80 and incubated at 37°C until log. phase. 1xPBS

supplemented with 0.05% Tween 80 was used for starvation assays. Cultures

were plated on LB Lennox agar (Fisher BP1427-2).

E. coli Top10, XL1-Blue, and Dh5α strains were used for cloning and *E. coli* BL21

Codon Plus and BL21(DE3) strains were used for expressing proteins. Antibiotic

concentrations for *M. smegmatis* were 25 µg/ml kanamycin and 20 µg/ml zeocin.

Antibiotic concentrations for *E. coli* were 50 µg/ml kanamycin, 25 µg/ml zeocin,

20 µg/ml chloramphenicol, and 140 µg/ml ampicillin.

Cloning and Strain construction

N-terminally Histidine (His) tagged cytosolic domains (1-300 amino acids (Gupta

et al., 2009)) of *pstP*$_c$T137E$_{Mtb}$ and *pstP*$_c$T141E$_{Mtb}$, and *fhaA*WT$_{Mtb}$ codon-

optimized for expression in *E. coli* were ordered from IDT, Inc. and cloned in to

pET28a.

The *Msmeg* mc^{2}155 strains Δ*pstP*::lox L5::pCT94-p766tetON6-*pstP*T134E$_{Msmeg}$

and Δ*pstP*::lox L5::pCT94-p766tetON6-*pstP*WT$_{Msmeg}$ were generated by L5 allele

swapping as described previously (Pashley and Parish, 2003; Shamma *et al.*,

2021). All vectors and strains are described in supplementary tables 1-3.

A vector that expressed mCherry-2B-PstPWT$_{Msmeg}$ was constructed using

isothermal assembly and integrated at the tweety phage integration site in

Msmeg. In brief, *pstP*WT was amplified from *Msmeg* genomic DNA, and

subcloned into a plasmid along with *mCherry2B* and the uv15 ribosomal

promoter. The resulting vector was transformed into *Msmeg*.

Growth Curve Assay

At least three biological replicates of *pstP*T134E$_{Msmeg}$ and WT allelic variants

were grown in 7H9 medium up to the mid-log. phase. The growth curve assays

were performed in non-treated non-tissue culture plates using a plate reader

(BioTek Synergy neo2 multi mode reader) in 200µl of 7H9 medium at 37°C

starting at an Optical Density at 600nm (OD$_{600}$) of 0.1.

The doubling times of each strain were calculated in the non-linear (curve fit)

method using an exponential growth equation with the least squares (ordinary) fit

in GraphPad Prism (version 7.0d). *P* values were calculated using two-tailed

unpaired t-tests.

Protein purification

His-MBP-PknA$_{Mtb}$ was expressed in *E. coli* BL21(DE3) and all other proteins were

expressed using *E. coli* BL21 Codon Plus cells. The *Mtb* genes of all studied

factors were cloned into expression vectors. His-MBP-PknB$_{Mtb}$, His-SUMO-

CwlM$_{Mtb}$, His-PstP$_c$WT$_{Mtb}$ and His-PstP$_c$T174E$_{Mtb}$ were expressed and purified as

previously described (Kieser *et al.*, 2015; Shamma *et al.*, 2021). His-

PstP$_c$T137E$_{Mtb}$ and His-PstP$_c$T141E$_{Mtb}$ were expressed and purified using the

same conditions and buffers used to purify His-PstP$_c$WT$_{Mtb}$ and His-

PstP$_c$T174E$_{Mtb}$ (Shamma *et al.*, 2021). We purified His-PstP$_c$WT$_{Mtb}$ and its

phospho-mimetic variants using nickel affinity chromatography followed by Size

Exclusion chromatography to ensure elution of the correctly folded proteins in the

eluate. His-PstP$_c$WT$_{Mtb}$ and all its phospho-mimetic variants are 34.1KDa and all

of them eluted at the same column volume when run on the size exclusion resin

(GE Healthcare Sephacryl S-200 in HiPrep 26/70 column). The purification

chromatograms were analysed with a standard curve made from running protein

standards (BIORAD) on the same size exclusion resin. All the PstP$_c$ variants

came out at a column volume where proteins weighing ~35KDa elute according

to our standard curve. This indicates that all the PstP variants fold like the wild-

type.

His-Wag31$_{Mtb}$ was induced with 0.5mM IPTG (isopropyl-β-D-

thiogalactopyranoside) for 6h at 18°C, and purified on Ni-NTA resin (G-

Biosciences 786-940) then dialyzed and concentrated. The buffer for His-

Wag31$_{Mtb}$ was 50mM Tris pH 8, 350mM NaCl, 1 mM dithiothreitol (DTT), and

10% glycerol. Imidazole (10mM and 20mM) was added to the buffers for lysis of

the cell pellets and washing the Ni-NTA resin respectively, and 250mM imidazole

was added for elution. The lysis buffer was supplemented with 1mM PMSF and

0.2% Triton X100.

His-MBP-PknA$_{Mtb}$ (1-279 amino acids containing the kinase domain (Baer *et al.*,

2014)) was induced with 0.5mM IPTG at 18°C overnight, purified with 5ml Ni-

NTA resins (BIO-RAD EconoFit Nuvia IMAC, 12009287), then run over the size

exclusion resin (Enrich SEC650, BIO-RAD 780-1650) after dialysis and

concentration to get soluble protein. The buffer for purifying His-MBP-PknA$_{Mtb}$

was 50mM Tris pH 7.5, 150mM NaCl, 20% glycerol. PMSF (1mM) was added to

the buffer while resuspending the cell pellets. Imidazole was used in buffer for

application to (20mM) and for elution from Ni-NTA resin (200mM).

His-FhaA$_{Mtb}$ was induced with 1mM IPTG at 18°C overnight and purified on Ni-

NTA resin (G-Biosciences, 786-940 in 5 ml Bio-Scale Mini Cartridges, BIO-RAD

7324661), then dialyzed, concentrated, and run over size exclusion resin (GE

Healthcare Sephacryl S-200 in HiPrep 26/70 column). His-FhaA$_{Mtb}$ was purified

using the buffers described in (Roumestand *et al.*, 2011).

***In vitro* dephosphorylation assays**

His-SUMO-CwlM$_{Mtb}$ at 0.15 µg/µl (2.64 µM) was phosphorylated *in vitro* with His-

MBP-PknB$_{Mtb}$ at 0.015 µg/µl for 30 minutes at room temperature with 0.15µCi/µl

ATP [γ-^{32}P] (5x10^{-8} µmol) (3000Ci/mmol, 10mCi/ml, PerkinElmer BLU002A250UC)

and 2mM MnCl$_2$ in buffer (50mM Tris, pH 7.5, 250mM NaCl, 10% glycerol and

1mM DTT). Right after collecting kinase reaction samples, the leftover mixture

was split into equal parts in different microcentrifuge tubes to carry on

phosphatase assays by adding replicates of His-PstP$_c$WT$_{Mtb}$, His-PstP$_c$T137E$_{Mtb}$,

His-PstPcT141E$_{Mtb}$ or His-PstPcT174E$_{Mtb}$ at 0.012 µg/µl to the kinase reaction

mixtures containing 0.12 µg/µl of His-SUMO-CwlM$_{Mtb}$ and incubated in

phosphatase buffer (50mM Tris pH 7.5, 10mM MnCl$_2$, and 1mM DTT) at room

temperature up to 40 minutes. His-FhaA$_{Mtb}$ at 0.15 µg/µl (2.54 µM) was

phosphorylated *in vitro* with His-MBP-PknB$_{Mtb}$ at 0.015 µg/µl for 45 minutes at

37°C with 0.15µCi/µl ATP [γ-^{32}P] in presence of 5mM MnCl$_2$ in 25mM Tris pH 7.5

and 1mM DTT. Right after collecting 0 minute kinase reaction sample, the

reaction mixture was then split into equal parts and incubated with either

PstP$_c$WT$_{Mtb}$ or different phospho-mimetic variants of it at 0.012 µg/µl up to 80

minutes at room temperature in presence of the phosphatase buffer where the

concentration of the substrate was 0.12 µg/µl For CwlM and FhaA, the amount of

kinase and phosphatases was 1/10th the amount of substrate, by weight.

His-Wag31$_{Mtb}$ at 0.15 µg/µl (4.90 µM) was phosphorylated with His-MBP-PknA$_{Mtb}$

at 0.03 µg/µl for 2 hours at room temperature in presence of 0.2µCi/µl ATP [γ-^{32}P]

(6.67x10^{-8} µM), 10mM MnCl$_2$ and buffer (50mM Tris pH 7.5, 20% glycerol and 150mM

NaCl). The amount of PknA was 1/5th the amount of Wag31 substrate, by weight.

The kinase reaction mixture was equally split into separate tubes after the 0

second kinase reaction sample was taken. His-PstP$_c$WT$_{Mtb}$ or its different

phospho-mimetic variants at 0.0013 µg/µl were added to the mixture containing

0.13 µg/µl substrate and incubated at room temperature for up to 120 seconds in

the phosphatase buffer. The amount of phosphatase was 1/100th the amount of

Wag31 substrate in the phosphatase reactions, by weight.

Control samples with no kinase and with no phosphatase were performed in all

the above-mentioned substrate dephosphorylation assays. All *in vitro*

dephosphorylation assays were carried out separately with two individually

purified batches of WT and phospho-mimetic mutant of PstP as biological

replicates. In all assays, the substrates were in molar excess over the ATP, so

that all the ATP will have been consumed before the phosphatase is added, and

re-phosphorylation by the kinases should not occur.

Quantification of γ-^{32}P signals

Samples from each *in vitro* biochemical reaction were collected before (0 minute

or second) and after addition of each of the phosphatases at several different

time points. The samples were run on 12% (Mini-Protean TGX, BIO-RAD,

4561046) gels and stained with Coomassie blue solution. After destaining, gels

were dried using the GD200 Vacuum Gel Drying system (Hoefer Inc.) and

imaged with phosphor screens (Molecular Dynamics) and a scanner (Storm 860,

Amersham Biosciences). ImageQuant software (Molecular Dynamics) was used

to process the images.

The protein bands on the gel and the phospho-signals on the autoradiogram

were then quantified with FIJI. The intensities of the substrate protein bands on

the Coomassie-stained gel and their corresponding phospho-signals on the

autoradiograms at each time point were normalized against the respective

protein band and the phospho-signal at 0 minute (His-SUMO-CwlM$_{Mtb}$ and His-

FhaA$_{Mtb}$) or 0 second (His-Wag31$_{Mtb}$) before addition of the phosphatase. These

relative intensities were used to calculate the intensity of phospho-signal over the

amount of substrate for each time point. These values were used to plot the

graph. The γ-^{32}P signal intensities of the kinases on autoradiograms at each time

point were normalized against the respective phospho-signal on the kinase at 0

minute (His-MBP-PknB$_{Mtb}$) or 0 second (His-MBP-PknA$_{Mtb}$) before addition of the

phosphatase. These relative γ-^{32}P signal intensity values were then used to plot

the graph in GraphPad Prism (version 7.0d). All values were plotted and p-values

were calculated using two-tailed, unpaired Student's t-test in GraphPad Prism

(version 7.0d).

Cell staining

At least three biological replicates of $pstP$T134E$_{Msmeg}$ and WT$_{Msmeg}$ allelic variants

were grown in 7H9 medium up to mid log. phase. Cultures were washed and

then starved in 1xPBS with 0.05% Tween 80 (PBST) at OD$_{600=}$ 0.3 for 15

minutes. 100µl of starved culture was then treated with 3µl of 10mM fluorescent

D-amino acid HADA for 10 minutes. Cells were then pelleted and resuspended in

PBST and fixed with 10µl of 16% paraformaldehyde for 10 minutes at room

temperature. Cells were then pelleted and resuspended in PBST before imaging

on agarose pads.

For staining cells in the log. phase, 1µl of 1mM HADA was added to 100µl

culture grown up to the mid-log. phase and incubated at 37°C for 15 minutes.

Stained cells were then pelleted, washed and fixed as described above.

Static microscopy and image analysis

A Nikon Ti-2 widefield epifluorescence microscope with a Photometrics Prime

95B camera and a Plan Apo 100x, 1.45 NA objective lens was used to image

cells. A 350/50nm excitation filter and a 460/50nm emission filter were used to

take the blue fluorescence images for HADA staining. All images were captured

using the NIS Elements software and analyzed using FIJI and MicrobeJ (Ducret

et al., 2016). Appropriate parameters for length, width and area were set in

MicrobeJ to detect cells. All detected dividing (V-snapping) cells in the images

were split at the septum so that each daughter cell could be considered as a

single cell. No overlapping cells in the images were included in the analysis.

Length and mean-intensities of HADA signals of at least 330 cells from each of

*pstP*T134E*Msmeg* and *pstP*WT*Msmeg* (at least 110 cells from each of three biological

replicate strains of each genotype) were quantified using MicrobeJ. The values of

the mean intensities of 330 cells of each *pstP* allelic mutant and WT are

represented in the graphs generated using GraphPad Prism (version 7.0d).

The medial intensity profiles of HADA signals in cells from different *pstP* allele

strains in log. phase and starvation analyzed with MicrobeJ were plotted on the

Y-axis over relative positions of cells using the "XStatProfile" plotting feature in

MicrobeJ to show the subcellular localization of fluorescent intensities in cells.

Demographs of HADA signal intensity across cell lengths in log. phase and

starvation were built using the "Demograph" feature of MicrobeJ by plotting the

medial intensity profiles of HADA signals.

Time-lapse microscopy

An inverted Nikon Ti-E microscope was used for the time-lapse imaging. An

environmental chamber was maintained at 37°C. Cells were cultivated in an B04

microfluidics plate from CellAsic; continuously supplied with fresh medium; and

imaged every 15 minutes using a 60x 1.4 N.A. phase contrast objective, with an

intermediate magnification of 1.5x. mCherry2B was excited with a SpectraX

Illuminator at 561-nm. Fluorescence was detected with a bandpass filter of 595/40

and detected with an ORCA Flash Digital sCMOS.

Western Blotting

To verify expression of PstPWT*Msmeg*, T134A*Msmeg* and T134E*Msmeg*, cultures were

grown in 10ml of 7H9 medium to OD_{600} = 0.8, pelleted and resuspended in

500μl PBS with 1mM phenylmethylsulfonyl fluoride (PMSF), and lysed

(MiniBeadBeater-16, model 607, Biospec). The supernatants of the cell lysates

were run on 12% resolving Tris-glycine gels and then transferred onto

polyvinylidene difluoride (PVDF) membranes (GE Healthcare). To detect PstP-

strep-tag II, rabbit anti-Strep-tag II antibody (1:1,000, Abcam, ab76949) in Tris-

buffered saline supplemented with Tween 20 (TBST) and 0.5% milk and goat

anti-rabbit IgG (H1L) horseradish peroxidase (HRP)-conjugated secondary

antibody (1:10,000, Thermo Fisher Scientific 31460) in TBST were used.

Acknowledgement:

This work was supported by NIH grant R01AI148917 to CCB and R01AI148255

to EHR.

References:

Abrahams, K.A., and Besra, G.S. (2021) Synthesis and recycling of the mycobacterial
cell envelope. *Current Opinion in Microbiology* **60**: 58–65.

Bach, H., Wong, D., and Av-Gay, Y. (2009) Mycobacterium tuberculosis PtkA is a novel
protein tyrosine kinase whose substrate is PtpA. *Biochemical Journal* **420**: 155–162.

Baer, C.E., Iavarone, A.T., Alber, T., and Sassetti, C.M. (2014) Biochemical and Spatial
Coincidence in the Provisional Ser/Thr Protein Kinase Interaction Network of
Mycobacterium tuberculosis. *J Biol Chem* **289**: 20422–20433.

Bandekar, A.C., Subedi, S., Ioerger, T.R., and Sassetti, C.M. (2020) Cell-Cycle-
Associated Expression Patterns Predict Gene Function in Mycobacteria. *Current Biology*
30: 3961-3971.e6.

Barford, D. (1996) Molecular mechanisms of the protein serine/threonine phosphatases.
Trends Biochem Sci **21**: 407–412.

Batt, S.M., Burke, C.E., Moorey, A.R., and Besra, G.S. (2020) Antibiotics and resistance:
the two-sided coin of the mycobacterial cell wall. *The Cell Surface* **6**: 100044.

Betts, J.C., Lukey, P.T., Robb, L.C., McAdam, R.A., and Duncan, K. (2002) Evaluation
of a nutrient starvation model of Mycobacterium tuberculosis persistence by gene and
protein expression profiling. *Molecular Microbiology* **43**: 717–731.

Bhamidi, S., Shi, L., Chatterjee, D., Belisle, J.T., Crick, D.C., and McNeil, M.R. (2012)
A bioanalytical method to determine the cell wall composition of Mycobacterium
tuberculosis grown in vivo. *Analytical Biochemistry* **421**: 240–249.

Bhaskara, G.B., Wong, M.M., and Verslues, P.E. (2019) The flip side of phospho-
signalling: Regulation of protein dephosphorylation and the protein phosphatase 2Cs.
Plant Cell Environ **42**: 2913–2930.

Boitel, B., Ortiz-Lombardía, M., Durán, R., Pompeo, F., Cole, S.T., Cerveñansky, C., and
Alzari, P.M. (2003) PknB kinase activity is regulated by phosphorylation in two Thr
residues and dephosphorylation by PstP, the cognate phospho-Ser/Thr phosphatase, in
Mycobacterium tuberculosis: Mycobacterial serine/threonine kinase and phosphatase.
Molecular Microbiology **49**: 1493–1508.

Boutte, C.C., Baer, C.E., Papavinasasundaram, K., Liu, W., Chase, M.R., Meniche, X., *et*
al. (2016) A cytoplasmic peptidoglycan amidase homologue controls mycobacterial cell
wall synthesis. *eLife* **5**: e14590.

Bradshaw, N., Levdikov, V., Zimanyi, C., Gaudet, R., Wilkinson, A., and Losick, R.
(2017) A widespread family of serine/threonine protein phosphatases shares a common
regulatory switch with proteasomal proteases. *eLife* **6**.

Bradshaw, N., and Losick, R. (2015) Asymmetric division triggers cell-specific gene
expression through coupled capture and stabilization of a phosphatase. *eLife* **4**: e08145.

Carette, X., Platig, J., Young, D.C., Helmel, M., Young, A.T., Wang, Z., *et al.* (2018)
Multisystem Analysis of *Mycobacterium tuberculosis* Reveals Kinase-Dependent
Remodeling of the Pathogen-Environment Interface. *mBio* **9**
http://mbio.asm.org/lookup/doi/10.1128/mBio.02333-17. Accessed February 15, 2019.

Chopra, P., Singh, B., Singh, R., Vohra, R., Koul, A., Meena, L.S., *et al.* (2003)
Phosphoprotein phosphatase of Mycobacterium tuberculosis dephosphorylates serine-
threonine kinases PknA and PknB. *Biochem Biophys Res Commun* **311**: 112–120.

Cohen, P. (1989) The structure and regulation of protein phosphatases. *Annu Rev*
Biochem **58**: 453–508.

Cole, S.T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., *et al.* (1998)
Deciphering the biology of Mycobacterium tuberculosis from the complete genome
sequence. **396**: 27.

Cottin, V., Linden, A.V., and Riches, D.W.H. (1999) Phosphorylation of Tumor Necrosis
Factor Receptor CD120a (p55) by p42mapk/erk2 Induces Changes in Its Subcellular
Localization. 14.

Cunningham, A.F., and Spreadbury, C.L. (1998) Mycobacterial Stationary Phase Induced
by Low Oxygen Tension: Cell Wall Thickening and Localization of the 16-Kilodalton ␣-
Crystallin Homolog. *J BACTERIOL* **180**: 8.

DeJesus, M.A., Gerrick, E.R., Xu, W., Park, S.W., Long, J.E., Boutte, C.C., *et al.* (2017)
Comprehensive Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via
Saturating Transposon Mutagenesis. *mBio* **8**: e02133-16, /mbio/8/1/e02133-16.atom.

Ducret, A., Quardokus, E.M., and Brun, Y.V. (2016) MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. *Nature Microbiology* **1**: 16077.

Dulberger, C.L., Rubin, E.J., and Boutte, C.C. (2020) The mycobacterial cell envelope —
a moving target. *Nat Rev Microbiol* **18**: 47–59.

Durán, R., Villarino, A., Bellinzoni, M., Wehenkel, A., Fernandez, P., Boitel, B., *et al.*
(2005) Conserved autophosphorylation pattern in activation loops and juxtamembrane
regions of Mycobacterium tuberculosis Ser/Thr protein kinases. *Biochemical and*
Biophysical Research Communications **333**: 858–867.

Echenique, J., Kadioglu, A., Romao, S., Andrew, P.W., and Trombe, M.-C. (2004)
Protein Serine/Threonine Kinase StkP Positively Controls Virulence and Competence in
Streptococcus pneumoniae. Infect Immun **72**: 2434–2437.

Gee, C.L., Papavinasasundaram, K.G., Blair, S.R., Baer, C.E., Falick, A.M., King, D.S.,
et al. (2012) A Phosphorylated Pseudokinase Complex Controls Cell Wall Synthesis in
Mycobacteria. *Sci Signal* **5**: ra7–ra7.

Grundner, C., Gay, L.M., and Alber, T. (2005) *Mycobacterium tuberculosis*
serine/threonine kinases PknB, PknD, PknE, and PknF phosphorylate multiple FHA
domains. *Protein Sci* **14**: 1918–1921.

Gupta, M., Sajid, A., Arora, G., Tandon, V., and Singh, Y. (2009) Forkhead-associated
Domain-containing Protein Rv0019c and Polyketide-associated Protein PapA5, from
Substrates of Serine/Threonine Protein Kinase PknB to Interacting Proteins of
Mycobacterium tuberculosis. *Journal of Biological Chemistry* **284**: 34723–34734.

Hannebelle, M.T.M., Ven, J.X.Y., Toniolo, C., Eskandarian, H.A., Vuaridel-Thurre, G.,
McKinney, J.D., and Fantner, G.E. (2020) A biphasic growth model for cell pole
elongation in mycobacteria. *Nat Commun* **11**: 452.

Hett, E.C., and Rubin, E.J. (2008) Bacterial Growth and Cell Division: a Mycobacterial
Perspective. *Microbiology and Molecular Biology Reviews* **72**: 126–156.

Hoffmann, C., Leis, A., Niederweis, M., Plitzko, J.M., and Engelhardt, H. (2008)
Disclosure of the mycobacterial outer membrane: Cryo-electron tomography and vitreous
sections reveal the lipid bilayer structure. *Proceedings of the National Academy of*
Sciences **105**: 3963–3967.

Irmler, A., and Forchhammer, K. (2001) A PP2C-type phosphatase dephosphorylates the
PII signaling protein in the cyanobacterium Synechocystis PCC 6803. *Proceedings of the*
National Academy of Sciences **98**: 12978–12983.

Iswahyudi, Mukamolova, G.V., Straatman-Iwanowska, A.A., Allcock, N., Ajuh, P.,
Turapov, O., and O'Hare, H.M. (2019) Mycobacterial phosphatase PstP regulates global
serine threonine phosphorylation and cell division. *Sci Rep* **9**: 8337.

Jani, C., Eoh, H., Lee, J., Hamasha, K., Sahana, M., Han, J.-S., *et al.* (2010) Regulation
of Polar Peptidoglycan Biosynthesis by Wag31 Phosphorylation in Mycobacteria. *BMC*
Microbiol **10**: 327.

Jarlier, V., and Nikaido, H. (1990) Permeability barrier to hydrophilic solutes in
Mycobacterium chelonei. *J Bacteriol* **172**: 1418–1423.

Jarlier, V., and Nikaido, H. (1994) Mycobacterial cell wall: Structure and role in natural
resistance to antibiotics. *FEMS Microbiology Letters* 11–18.

Juris, S.J., Rudolph, A.E., Huddler, D., Orth, K., and Dixon, J.E. (2000) A distinctive role
for the Yersinia protein kinase: Actin binding, kinase activation, and cytoskeleton
disruption. *Proceedings of the National Academy of Sciences* **97**: 9431–9436.

Kang, C.-M., Abbott, D.W., Park, S.T., Dascher, C.C., Cantley, L.C., and Husson, R.N.
(2005) The Mycobacterium tuberculosis serine/threonine kinases PknA and PknB:
substrate identification and regulation of cell shape. *Genes Dev* **19**: 1692–1704.

Kang, C.-M., Nyayapathy, S., Lee, J.-Y., Suh, J.-W., and Husson, R.N. (2008) Wag31, a
homologue of the cell division protein DivIVA, regulates growth, morphology and polar
cell wall synthesis in mycobacteria. *Microbiology* **154**: 725–735.

Kieser, K.J., Baranowski, C., Chao, M.C., Long, J.E., Sassetti, C.M., Waldor, M.K., *et al.*
(2015) Peptidoglycan synthesis in Mycobacterium tuberculosis is organized into
networks with varying drug susceptibility. *PNAS* **112**: 13087–13092.

Kieser, K.J., and Rubin, E.J. (2014) How sisters grow apart: mycobacterial growth and
division. *Nature Reviews Microbiology* **12**: 550–562.

Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., *et al.* (2012) In Situ
Probing of Newly Synthesized Peptidoglycan in Live Bacteria with Fluorescent D -
Amino Acids. *Angewandte Chemie International Edition* **51**: 12519–12523.

Le, N.-H., Locard-Paulet, M., Stella, A., Tomas, N., Molle, V., Burlet-Schiltz, O., *et al.*
(2020) The protein kinase PknB negatively regulates biosynthesis and trafficking of
mycolic acids in mycobacteria. *J Lipid Res* jlr.RA120000747.

Liu, Y., Tan, S., Huang, L., Abramovitch, R.B., Rohde, K.H., Zimmerman, M.D., *et al.*
(2016) Immune activation of the host cell induces drug tolerance in *Mycobacterium*
tuberculosis both in vitro and in vivo. *J Exp Med* **213**: 809–825.

Lu, G., and Wang, Y. (2008) FUNCTIONAL DIVERSITY OF MAMMALIAN TYPE
2C PROTEIN PHOSPHATASE ISOFORMS: NEW TALES FROM AN OLD FAMILY:
Function of PP2C isoforms. *Clinical and Experimental Pharmacology and Physiology*
35: 107–112.

Marquardt, J.L., Siegele, D.A., Kolter, R., and Walsh, C.T. (1992) Cloning and
sequencing of Escherichia coli murZ and purification of its product, a UDP-N-
acetylglucosamine enolpyruvyl transferase. *J Bacteriol* **174**: 5748–5752.

Marrakchi, H., Lanéelle, M.-A., and Daffé, M. (2014) Mycolic Acids: Structures,
Biosynthesis, and Beyond. *Chemistry & Biology* **21**: 67–85.

Molle, V., Brown, A.K., Besra, G.S., Cozzone, A.J., and Kremer, L. (2006) The
Condensing Activities of the Mycobacterium tuberculosis Type II Fatty Acid Synthase
Are Differentially Regulated by Phosphorylation. *Journal of Biological Chemistry* **281**:
30094–30103.

Mougous, J.D., Gifford, C.A., Ramsdell, T.L., and Mekalanos, J.J. (2007) Threonine
phosphorylation post-translationally regulates protein secretion in Pseudomonas
aeruginosa. *Nat Cell Biol* **9**: 797–803.

Nguyen, L. (2016) Antibiotic resistance mechanisms in M. tuberculosis: an update. *Arch*
Toxicol **90**: 1585–1604.

Pashley, C.A., and Parish, T. (2003) Efficient switching of mycobacteriophage L5-based
integrating plasmids in *Mycobacterium tuberculosis*. *FEMS Microbiology Letters* **229**:
211–215.

Prisic, S., Dankwa, S., Schwartz, D., Chou, M.F., Locasale, J.W., Kang, C.-M., *et al.*
(2010) Extensive phosphorylation with overlapping specificity by Mycobacterium
tuberculosis serine/ threonine protein kinases. *Proceedings of the National Academy of*
Sciences **107**: 7521–7526.

Prisic, S., and Husson, R.N. (2014) Mycobacterium tuberculosis Serine/Threonine
Protein Kinases. *Microbiology Spectrum* **2**
http://www.asmscience.org/content/journal/microbiolspec/10.1128/microbiolspec.MGM2
-0006-2013. Accessed February 6, 2019.

Pullen, K.E., Ng, H.-L., Sung, P.-Y., Good, M.C., Smith, S.M., and Alber, T. (2004) An
Alternate Conformation and a Third Metal in PstP/Ppp, the M. tuberculosis PP2C-Family
Ser/Thr Protein Phosphatase. *Structure* **12**: 1947–1954.

Roumestand, C., Leiba, J., Galophe, N., Margeat, E., Padilla, A., Bessin, Y., *et al.* (2011)
Structural Insight into the Mycobacterium tuberculosis Rv0020c Protein and Its
Interaction with the PknB Kinase. *Structure* **19**: 1525–1534.

Sajid, A., Arora, G., Gupta, M., Upadhyay, S., Nandicoori, V.K., and Singh, Y. (2011)
Phosphorylation of Mycobacterium tuberculosis Ser/Thr Phosphatase by PknA and PknB.
PLoS ONE **6**: e17871.

Sarathy, J., Dartois, V., Dick, T., and Gengenbacher, M. (2013) Reduced Drug Uptake in
Phenotypically Resistant Nutrient-Starved Nonreplicating Mycobacterium tuberculosis.
Antimicrobial Agents and Chemotherapy **57**: 1648–1653.

Sarathy, J.P., Via, L.E., Weiner, D., Blanc, L., Boshoff, H., Eugenin, E.A., *et al.* (2017)
Extreme Drug Tolerance of *Mycobacterium tuberculosis* in Caseum. *Antimicrobial*
Agents and Chemotherapy **62** http://aac.asm.org/lookup/doi/10.1128/AAC.02266-17.
Accessed February 6, 2019.

Shamma, F., Quintanilla, S., Bandekar, A., Sassetti, C., and Boutte, C. (2021)
Phosphorylation on PstP Regulates Cell Wall Metabolism and Antibiotic Tolerance in
Mycobacterium smegmatis. *Journal of Bacteriology* **203**: e00563-20.

Sharma, A.K., Arora, D., Singh, L.K., Gangwal, A., Sajid, A., Molle, V., *et al.* (2016)
Serine/Threonine Protein Phosphatase PstP of *Mycobacterium tuberculosis* Is Necessary
for Accurate Cell Division and Survival of Pathogen. *Journal of Biological Chemistry*
291: 24215–24230.

Sharma, K., Gupta, M., Krupa, A., Srinivasan, N., and Singh, Y. (2006) EmbR, a
regulatory protein with ATPase activity, is a substrate of multiple serine/threonine
kinases and phosphatase in *Mycobacterium tuberculosis*. *FEBS Journal* **273**: 2711–2721.

Thakur, M., and Chakraborti, P.K. (2006) GTPase Activity of Mycobacterial FtsZ Is
Impaired Due to Its Transphosphorylation by the Eukaryotic-type Ser/Thr Kinase, PknA.
J Biol Chem **281**: 40107–40113.

Turapov, O., Forti, F., Kadhim, B., Ghisotti, D., Sassine, J., Straatman-Iwanowska, A., *et*
al. (2018) Two Faces of CwlM, an Essential PknB Substrate, in Mycobacterium
tuberculosis. *Cell Rep* **25**: 57-67.e5.

Vijay, K., Brody, M.S., Fredlund, E., and Price, C.W. (2000) A PP2C phosphatase
containing a PAS domain is required to convey signals of energy stress to the sB
transcription factor of Bacillus subtilis. *Molecular Microbiology* 9.

Viswanathan, G., Joshi, S.V., Sridhar, A., Dutta, S., and Raghunand, T.R. (2015)
Identifying novel mycobacterial stress associated genes using a random mutagenesis
screen in Mycobacterium smegmatis. *Gene* **574**: 20–27.

Viswanathan, G., Yadav, S., Joshi, S.V., and Raghunand, T.R. (2016) Insights into the
function of FhaA, a cell-division associated protein in mycobacteria. *FEMS Microbiology*
Letters fnw294.

Viswanathan, G., Yadav, S., and Raghunand, T.R. (2017) Identification of Mycobacterial
Genes Involved in Antibiotic Sensitivity: Implications for the Treatment of Tuberculosis

with β-Lactam-Containing Regimens. *Antimicrobial Agents and Chemotherapy* **61**
http://aac.asm.org/lookup/doi/10.1128/AAC.00425-17. Accessed February 6, 2019.

Wang, J., Li, C., Yang, H., Mushegian, A., and Jin, S. (1998) A Novel Serine/Threonine
Protein Kinase Homologue of *Pseudomonas aeruginosa* Is Specifically Inducible within
the Host Infection Site and Is Required for Full Virulence in Neutropenic Mice. *J*
Bacteriol **180**: 6764–6768.

World Health Organization (2021) *Global tuberculosis report 2021*. World Health
Organization, Geneva. https://apps.who.int/iris/handle/10665/346387. Accessed
November 14, 2021.

Xie, Z., Siddiqi, N., and Rubin, E.J. (2005) Differential Antibiotic Susceptibilities of
Starved Mycobacterium tuberculosis Isolates. *Antimicrobial Agents and Chemotherapy*
49: 4778–4780.

Figure Legends
748

Flap
subdomain

749
750
Figure 1 Phosphorylation sites are not near the active site in PstP

Schematic of the crystal structure of PstP from *M. tuberculosis* (PstPWT$_{Mtb}$) (Pullen
et al., 2004) (PDB code 1TXO) showing the conserved threonines (T) which are
phosphorylated by the kinases PknA and PknB (Sajid *et al.*, 2011). Highlighted on
the structure are: red - T137 (T134 in PstP$_{Msmeg}$); blue - T141 (T138 in PstP$_{Msmeg}$);
green - T174 (T171 in PstP$_{Msmeg}$); yellow - Mn^{2+} present at the conserved two-
metal center typically seen in the PP2C subfamily of protein Serine/Threonine
phosphatases, which are part of the active site core; pink - the unique third Mn^{2+}
ion bound in the catalytic core of PstP$_{Mtb}$ but not present in the human PP2Cα
which sits in a groove created by the flap subdomain next to the active site; orange
- Ser60 and Asp118 that act as ligands for the third Mn ion; lime green - residues
of the catalytic sites surrounding the two-metal center. Arrows point to position of
the OH-groups on T137, T141 and T174 that are phosphorylated.

Time 0 15 30 45 60 90 120 sec

FIGURE 2 Dephosphorylation of FhaA by PstP and phospho-mimetic isoforms.

(a) Autoradiograms and Coomassie-stained SDS gels of *in vitro* phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-FhaA$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$137E$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no phosphatase control reaction (bottom panel). The ratio of each phosphatase to His-FhaA$_{Mtb}$ in each of the assays was 1:10. One set of representative images for each assay is shown here from two individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-FhaA$_{Mtb}$ on autoradiogram over the relative amount of His-FhaA$_{Mtb}$ on the SDS gel, average between two replicate experiments. The relative values refer to the normalized values of intensity of [γ-^{32}P] signal and the protein bands on gel at each time point (0 to 80min) with the 0 time point. P values were calculated using two-tailed unpaired t-test.

P-values of T137E at 0 min *vs.* T137E 80 min =0.7598, WT *vs.* T137E at 80 min= 0.0487, WT *vs.* T141E at 80 min =0.3712, WT *vs.* T174E at 80 min=0.2000. The error bars represent standard error of mean.

783 Time 0   15 30 45 60 90 120 sec

784
785
786 **FIGURE 3 Dephosphorylation of Wag31**$_{Mtb}$ **by PstP and phospho-mimetic**
787 **isoforms.**
788

(a) Autoradiograms and Coomassie-stained SDS gels of *in vitro* phosphatase
assays performed with ATP-[γ-^{32}P]-phosphorylated His-Wag31$_{Mtb}$ and His-
PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third
panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no phosphatase control reaction
(bottom panel). The ratio of each phosphatase to His-Wag31$_{Mtb}$ in the assays was
1:100. One set of representative images for each reaction is shown here from two
individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-Wag31$_{Mtb}$ on
autoradiograms over the relative amount of His-Wag31$_{Mtb}$ on the SDS gel, average
between two replicate experiments. The relative values are the normalized values
of intensity of [γ-^{32}P] signal and the protein bands on gel at each time point (0 to
120 sec) with the 0 time point. P values were calculated using two-tailed unpaired
t-test.

P-values of T137E at 0 sec *vs.* 15 sec = 0. 0.1947 and 0 sec *vs.* 120sec= 0.7493,

WT *vs.* T137E at 120 sec= 0.0130, WT *vs.* T141E at 120 sec =0.0106 and WT *vs.*

T174E at 120 sec = 0.170. The error bars represent standard error of mean.

(a) Dephosphorylation of His-SUMO-CwlM$_{Mtb}$

His-PstP$_c$WT$_{Mtb}$

His-PstP$_c$T137E$_{Mtb}$

His-PstP$_c$T141E$_{Mtb}$

His-PstP$_c$T174E$_{Mtb}$

no phosphatase control

Time 0   1   3   5   10   20   40 min

(b) Quantification of phosphorylation of His-SUMO-CwlM$_{Mtb}$

● +His-PstP$_c$WT$_{Mtb}$
■ +His-PstP$_c$T137E$_{Mtb}$
▲ +His-PstP$_c$T141E$_{Mtb}$
▼ +His-PstP$_c$T174E$_{Mtb}$
◆ no phosphatase control

Relative (γ-^{32}P signal/amount of His-SUMO-CwlM$_{Mtb}$)

Time (min)

(c) Linear rate of dephosphorylation of substrates

His-SUMO-CwlM$_{Mtb}$ His-FhaA$_{Mtb}$ His-Wag31$_{Mtb}$

Δphospho-signal of 10μg substrate/μg PstP/minute

FIGURE 4 Dephosphorylation of CwlM$_{Mtb}$ by PstP$_{Mtb}$ its phospho-mimetic isoforms and linear rate of dephosphorylation of CwlM$_{Mtb}$, FhaA$_{Mtb}$ and Wag31$_{Mtb}$

(a) Autoradiograms and Coomassie-stained SDS gels of *in vitro* phosphatase assays performed with ATP-[γ-^{32}P]-phosphorylated His-SUMO-CwlM$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel), His-PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel), His-PstP$_c$T174E$_{Mtb}$ (fourth panel) and no phosphatase control reaction (bottom panel). The ratio of each phosphatase to His-SUMO-CwlM$_{Mtb}$ was 1:10. One set of representative images for each reaction is shown here from two individual assays performed with individually purified batches of phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-SUMO-CwlM$_{Mtb}$ on autoradiogram over the relative amount of His-SUMO-CwlM$_{Mtb}$ on the SDS gel, average between two replicate experiments. The relative values are the normalized values of intensity of [γ-^{32}P] signal and the protein bands on gel at each time point (0 to 40min) with the 0 time point. P values were calculated using two-tailed unpaired t-test. P-values of T137E at 0 min *vs.* 40 min= 0.1107, WT *vs.* T137E at 40 min= 0.0127, WT *vs.* T141E at 40 min=0.3683 and WT *vs.* T174E at 40 min= 0.5918. The error bars represent standard error of mean.

(c) Quantification of linear rate of change in [γ-^{32}P] signal on 10µg of His-SUMO-CwlM$_{Mtb}$, His-FhaA$_{Mtb}$ and His-Wag31$_{Mtb}$ by 1µg His-PstP$_c$WT$_{Mtb}$ and its phospho-mimetic mutants per minute. P values were calculated using two-tailed unpaired t-test. **, *P* value = 0.0026 and *, *P* value 0.0306. The error bars represent standard error of mean.

FIGURE 5 Dephosphorylation of PknB$_{Mtb}$ and PknA by PstPWT$_{Mtb}$ and its
phospho-mimetic variants

(a) Autoradiograms showing [γ-^{32}P]-phospho-signals on His-MBP-PknB$_{Mtb}$ in the *in*
vitro phosphatase assays with His-SUMO-CwlM$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first
panel), His-PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel) and His-
PstP$_c$T174E$_{Mtb}$ (fourth panel), where His-MBP-PknB$_{Mtb}$ was used to phosphorylate
His-SUMO-CwlM$_{Mtb}$. One set of representative images for each reaction is shown
here from two individual assays performed with individually purified batches of
phosphatases.

(b) Quantification of relative intensities of [γ-^{32}P] on His-MBP-PknB$_{Mtb}$ on
autoradiogram, average between two replicate experiments. The relative values
refer to the normalized values of intensity of [γ-^{32}P] signal at each time point with
the 0 time point. P values were calculated using two-tailed unpaired t-test. P value
of T137E at 0 min *vs.* 40 min= 0.0523. *, *P* value= 0.0375 (WT *vs.* T137E at 40
min); *, *P* value=0.0314 (T141E *vs.*T137E at 40 min); *, *P* value= 0.0217 (T174E
*vs.*T137E at 40 min). The error bars represent standard error of mean.

(c) Autoradiograms showing [γ-^{32}P]-phospho-signals on His-MBP-PknA$_{Mtb}$ in the *in*
vitro phosphatase assays with His-Wag31$_{Mtb}$ and His-PstP$_c$WT$_{Mtb}$ (first panel), His-
PstP$_c$WT$_{Mtb}$ (second panel), His-PstP$_c$T141E$_{Mtb}$ (third panel) and His-
PstP$_c$T174E$_{Mtb}$ (fourth panel), where His-Wag31$_{Mtb}$ was first phosphorylated with
His-MBP-PknA$_{Mtb}$. One set of representative images for each reaction is shown
here from two individual assays performed with individually purified batches of
phosphatases.

(d) Quantification of relative intensities of [γ-^{32}P] on His-MBP-PknA$_{Mtb}$ on
autoradiogram. P values were calculated using two-tailed unpaired t-test. The
relative values refer to the normalized values of intensity of [γ-^{32}P] signal at each
time point with the 0 time point. P value of T137E at 0 sec *vs.* 120 sec= 0.0643. *,
P value= 0.0163 (WT *vs.* T137E at 120 sec). The error bars represent standard
error of mean.

Figure 6 The phosphosite on T134 on PstP_{Msmeg} affects PG metabolism

(a) Doubling times of *Msmeg* strains carrying the *pstP*_{Msmeg} WT and the phospho-mimetic T134E (equivalent to T137 on PstP_{Mtb}) alleles. Each bar represents the mean of individual average doubling times of three biological replicates of each genotype from two different experiments on different dates. The error bars

represent mean with standard deviation. **, *P* value= 0.0026.

(b) Quantification of cell lengths of isogenic *pstP~Msmeg~* allele strains (WT and
T134E) grown in 7H9 in log. phase and starved in PBS with Tween80. At least 100
cells from each of three biological replicates were quantified. P values were
calculated by unpaired t-test. *, *P* value = 0.0122, ****, *P* value= 0.000000031.

(c) Quantification of mean intensities of HADA of *pstP~Msmeg~* allele strains (WT and
T134E) grown in 7H9 in log. phase and starved in PBS with Tween80. Cells were
stained after 15 minutes of starvation and washed and fixed after another 15
minutes (starved for a total of 30 minutes). At least 100 cells from each of three
biological replicates were quantified. P values were calculated by unpaired t-test.
*, *P* value = 0.0109, ****, *P* value= <0.000000000000001.

(d) and (e) Representative HADA and phase images of cells from *pstP~Msmeg~* allele
strains (WT and T134E) grown in 7H9 in log. phase (d) and after 30 minutes of
starvation in PBS with Tween80 (e).

(f) and (g) Intensity profiles of HADA signals in cells of *pstP~Msmeg~* allele strains (WT
and T134E) in log. phase (f) and after starving for 30 minutes in PBS supplemented
with Tween80 (g). The shaded regions denote standard deviations and the solid
lines represent the mean intensity values. Signal intensities from at least 110 cells
from each of three biological replicates of every *pstP~Msmeg~* allelic variant genotype
were quantified and used to analyze the signal profiles in MicrobeJ. 0 to 1 on the
X-axes represent cells from the old pole to the new pole, respectively.

(h) α-strep western blots of strep-tagged PstP*~Msmeg~* in *pstP~Msmeg~*WT,
*pstP~Msmeg~*T134E and *pstP~Msmeg~*T134A allele strains in log. phase. A non-specific
band is shown as a loading control.

Figure 7 PstPWT*~Msmeg~* localizes at mid-cell.

Msmeg cells expressing mCherry2B-PstPWT*~Msmeg~* were imaged over time in a

CellAsic Microfluidic device at 15-minute intervals. Intensity is seen at the mid-
cell (white arrows) for several frames before cell separation can be visualized by
phase contrast microscopy (yellow arrows).