1 **Calcium homeostasis disruption initiates rapid growth after micro-fragmentation in the**
2 **scleractinian coral *Porites lobata***

4   Colin Lock[1]*, Bastian Bentlage[1], Laurie J Raymundo[1]

5   [1]Marine Laboratory, University of Guam, Mangilao, GU 96913, USA

7   *Corresponding author:*
8   PO Box 5401, Marine Laboratory, University of Guam, Mangilao, Guam, USA 96923
9   clockdive@gmail.com

Abstract

Coral reefs are ecosystems under increasing threat from global climate change. Coral restoration

is a tool for preserving biological and ecological function of coral reefs by mitigating coral loss

and maintaining the structural integrity and complexity of reefs. To generate the necessary stock

for coral restoration, larger coral colonies are usually fragmented to generate smaller specimens

for outplanting, taking advantage of the high regenerative ability of corals. In this study, we

utilized RNA-seq technology to understand the physiological responses of *Porites lobata*

colonies to physical fragmentation and outplanting, which have thus far not been characterized.

Our results demonstrate that P. lobata fragments undergoing physical injury recover through two

distinct phases: rapid wound regeneration of the cut margins, followed by a slower growth phase

that cements the colony to the substrate. Our study found *rapid* physiological responses to acute

physical injury and outplanting in the coral host that involved significantly increased energy

production, calcium homeostasis disruption, and ER stress leading to increased antioxidant

expression and rates of protein turnover. Our results suggest that phosphoinositide-mediated

acute calcium homeostasis disruption stimulates wound recovery processes in response to

physical injury. Contrary to other coral transcriptomic experiments, symbiont gene expression

revealed extremely low gene differences in response to fragmentation, growth, and outplanting.

These results provide insight into the physiological mechanisms that allow for rapid wound

healing and stabilization in response to physical injury in corals.

1 | INTRODUCTION

Global coral reef decline has become so pronounced that passive conservation strategies that rely on natural recovery of reefs, such as the designation of marine protected areas, may be inadequate to preserve the biodiversity and ecosystem services tropical coral reefs provide (Rinkevich, 2005; Forsman et al., 2006). To address this concern, active coral restoration is rapidly becoming an integral management tool to mitigate coral loss and maintain the structural integrity and complexity of reefs (Kojis & Quinn, 2001; Baums, 2008; Edwards, 2008; Barton et al., 2017). Procuring fragments from healthy coral colonies and propagating these to establish coral nursery stock for outplanting in restoration efforts is a commonly used approach in coral reef restoration (Epstein et al., 2001; Rinkevich, 2005; Edwards & Gomez, 2011). After nursery stock has reached a stable size and thus an increased chance of long-term survival (Lirman, 2000; Rinkevich, 2005; Edwards & Gomez, 2011), fragments may be cut from the coral stock. These fragments are then firmly affixed to degraded reefs, while sustaining the initial broodstock for future outplanting (Epstein et al., 2001). While fragmentation has been successfully used as a coral reef restoration technique for corals that fragment naturally, such as species of *Acropora*, (Lirman, 2000; Highsmith, 2007; Lirman et al., 2010), micro-fragmentation of corals that do not naturally propagate asexually via fragmentation, such as species of *Porites*, has been a more recent development (Forsman et al., 2006, 2015; Page et al., 2018). Micro-fragmentation involves cutting corals into minute pieces and growing them in a protected nursery, during which corals grow to a size suitable for outplanting (Forsman et al., 2006, 2015; Page et al., 2018). Microfragmentation allows the generation of a large number of fragments for stress experiments or outplanting, while minimizing impacts to source colonies.

Acute physical injury, such as that caused by fragmentation, may increase mortality

(Lirman, 2000; Forsman et al., 2006) and lead to a loss of fecundity (Harrison & Wallace, 1990;

Zakai et al., 2000; Okubo et al., 2007), limiting dispersal and recruitment in reefs restored in this

manner until coral colonies reach mature sizes (Kozłowski & Wiegert, 1986; Ward, 1995; Smith

& Hughes, 1999). Curiously, micro-fragments of corals display short-term accelerated growth

immediately following initial fragmentation and resulting injury to tissues (Forsman et al., 2015;

Page et al., 2018). The mechanisms underlying this phenotypic response remain poorly

understood. Early studies on corals that focused on recovery from physical injury suggested that

energy required to heal lesions was sourced almost exclusively from adjacent cells (Bak &

Yvonne, 1980; Bak, 1983; Meesters et al., 1994), which is consistent with a lack of cellular

specialization and postulated low levels of colony integration in corals. However, Oren et al.

(2012) demonstrated that lesions in the merulinid *Dipsastrea favus* healed more effectively in

large colonies compared to smaller ones. Further, even polyps distant from the lesion showed

reduced fecundity, indicating that resources for regeneration were translocated across the colony

rather than just from cells bordering the lesion. Such findings indicate that coral colonies are

more highly integrated than traditionally thought.

Massive *Porites* spp. have a documented high regeneration capacity, which may be

related to the integration of resources throughout their often large colonies (e.g., Lough &

Barnes, 2000). Indeed, *Porites* spp. have been shown to heal from tissue damage induced by

micro-fragmentation within days, initiating calcification along fragment margins within a week

(Forsman et al., 2006, 2015). However, the physiological mechanisms that underlie this

response, which is tied to the success of restoration efforts, have not been thoroughly explored.

Here, we employed a time-series transcriptomic approach to shed light on the cellular-level

mechanisms that drive lesion healing and accelerated growth in micro-fragments of *Porites*

lobata. Contrary to other stress events that dampen energy production, suppress calcification,

and increase mortality in corals (e.g., Desalvo et al., 2008; Császár et al., 2009; DeSalvo et al.,

2012; Bay et al., 2013; Tarrant et al., 2014; Maor-Landaw & Levy, 2016; Aguilar et al., 2019),

we provide evidence from gene expression profiles that indicate a temporary increase in energy

metabolism and a phosphoinositide-mediated disruption of calcium-homeostasis that likely

impacts cell cycle regulation and growth. The endoplasmic reticulum (ER) plays an important

role in regulating calcium homeostasis and mitigating the impacts that increased mitochondrial

reactive oxygen production has on protein synthesis during physiological stress. We provide a

model of the cellular processes that are likely driving the increase in growth rates of *Porites* spp.

in the days and weeks following micro-fragmentation.

2 | MATERIALS AND METHODS

2.1 | Sampling of source colonies

Six *Porites lobata* colonies, ranging in size from 15-25cm, were sourced from the Luminao reef

flat, located on the western coast of Guam. All colonies were collected from ~2 m depth and

were at least 20 m apart to minimize the possibility that they were of clonal origin. Colonies

were immediately transported in fresh seawater to the Marine Laboratory at the University of

Guam and allowed to acclimate for four weeks in a flow-through seawater tank under 70%

shade. Oscillating water motion was maintained with OW-40 wavemakers (Zhongshan Jebao

Electronic Appliance Co, Beijing, China).

2.2 | DNA barcoding and algal symbiont profiling

Because *in situ* species-level identification is difficult in massive *Porites* spp., colonies with similar gross morphologies were sampled and species identifications were then verified using DNA barcoding. DNA was extracted using the GenCatch genomic DNA extraction kit (Epoch Life Science, Sugar Land, TX) following the manufacturer's protocol for tissue samples. Mitochondrial markers mt-16 (Cox3-Cox2) and mt-20 (ND5-tRNA-Trp-ATP8-Cox1) (Paz-García et al., 2016) were amplified via PCR using 0.3uM primer, 0.3mM dNTP, 14.25 uL Water, 14.25ul HiFi Fidelity Buffer (5x), 2.5 units Taq (HIFI kapa 1U) and denaturation at 94°C for 120 s, followed by 30 cycles of 94°C for 30 s, 54°C for 30 s, 72°C for 60 s, with a further extension step of 72°C for 300 s. PCR products were sequenced in both directions using Sanger sequencing and resulting sequences assembled using the overlap-layout-consensus algorithm implemented in Geneious Prime (Biomatters, Auckland, New Zealand). Following assembly, consensus sequences for each specimen were aligned using MUSCLE (v. 3.8; Edgar, 2004) and sequences compared to NCBI's GenBank nucleotide sequence database.

To determine the dominant clade of Symbiodinaceae associated with each sample, transcriptomic samples from each source colony were mapped against symbiont transcriptomes for the four Symbiodiniaceae genera that associate with scleractinian corals (*Symbiodinium, Breviolum, Cladocopium, Durusdinium*; LaJeunesse et al., 2018). The number of high-quality mapped reads was then used to calculate relative abundances of Symbiodinaceae genera, following the approach of Manzello et al., (2019).

2.3 | Micro-fragmentation, *ex situ* husbandry, and outplanting

After four weeks of acclimation in flow-through seawater tanks, colonies were cut into ~1.5 cm^2

micro-fragments using a seawater-cooled C-40 diamond band saw (Gryphon Corporation,

Sylmar, CA), glued onto individual tiles, and randomly assigned to one of two replicate tanks.

Micro-fragments were created from coral tissue at least 2 cm from the growing edge of colonies

to reduce variability in growth rates and gene expression differences between fragments. Excess

skeleton was removed from the base of each fragment to create micro-fragments of the same

height, which were then attached to ceramic tiles using cyanoacrylate gel and labeled according

to source colonies. In total, 36 fragments were cut (n=6/colony), to be destructively sampled for

transcriptomics at predetermined time points and an additional 42 fragments were prepared (n=7/

colony) to document growth rates. After micro-fragments had grown for two months in the two

flow-through tanks, 42 micro-fragments were affixed to natural reef substrate using Splash Zone

epoxy (Z-spar, Périgny, France), in three plots at Asan Beach National Park, within a Marine

Preserve on the western coast of Guam.

2.4 | Growth rates

Weekly maintenance of the micro-fragments in *ex-situ* culture included removal of algae and

detritus from tiles and tanks, and visual inspection of all tiles to identify and remove potential

nudibranch predators. Water temperature was monitored using HOBO Tidbit temperature

loggers (Onset, Bourne, MA, USA). Weekly photographs of each fragment (top-down and side-

views from all four sides of each tile) were taken using an Olympus TG-5 (Olympus, Shinjuku,

Tokyo, Japan) in an underwater housing mounted to a PVC stand with attached scale bar. Micro-

fragment surface area was estimated from photographs using ImageJ version 1.43u (National

Institutes of Health, Bethesda, MD), combining surface areas measured from top and side views

for each micro-fragment. Growth was determined via the change in surface area from one week

to the next and normality of growth rate distributions was tested using a Shapiro-Wilks test.

Growth was then binned into two phases, Phase A from T1 (week 0) to T3 (week 3) during

which cut margins were overgrown by new tissue; and Phase B from T3 (week 2) to T5 (week 8)

during which coral tissue and skeleton were deposited onto tiles. To account for the non-

independence of time-series, Repeated Measures ANOVA and Student's t tests were

implemented in R studio to test for significance between these two growth phases, as well as

differences between colonies and rearing tanks.

2.4 | Transcriptome sequencing, assembly, and annotation

One micro-fragment per colony was destructively sampled for RNA extraction at five time

points (Figure 1): after colony collection and acclimation to experimental tanks but prior to

micro-fragmentation (baseline; T1); 24 hr after micro-fragmentation (initial stress; T2); at the

first signs of calcification along cut margins (regeneration; T3); at 2 mo of *ex situ* growth

(recovery; T4); and one day after outplanting (transplantation stress; T5). Tissue samples from

the tank experiment (T1-T4) were placed into sterile flat wire sample bags and immediately flash

Figure 1: The timeline of transcriptomic sampling

frozen using liquid nitrogen; tissue samples were stored at -80°C until extraction. Transcriptomic

samples from the outplanted fragments (T5) were field-preserved in RNAlater and subsequently

stored at -80°C until extraction. Total RNA was extracted using an E.Z.N.A. Plant RNA kit

(Omega Bio-Tek, Norcross, GA). DNA was removed from RNA extracts using a DNase 1

digest. After extraction, RNA was quantified fluorometrically using a Qubit RNA HS Assay kit

(Life Technologies, Carlsbad, CA) and RNA integrity verified using a BioAnalyzer (Agilent

Technologies, Santa Clara, CA). RNA sequencing libraries were constructed using the NEBNext

Ultra RNA Library Prep Kit (New England Biolabs, Ipswich, MA) following manufacturer's

protocol. Sequencing libraries were multiplexed and 150bp paired-end reads generated on an

NextSeq 500 sequencer (Illumina, San Diego, CA) using the Illumina NextSeq 500/550 High

Output Kit v2.5 (Illumina, San Diego, CA).

175   Adapter sequences and bases with a Phred-scaled quality score of less than 30 were

trimmed using Trim Galore (Martin, 2011). Prior to *de novo* transcriptome assembly, all reads

were combined and then normalized using in-silico normalization as implemented in Trinity

v2.10.0 (Grabherr et al., 2013) with a max coverage set to 50; this normalized read set was then

used for transcriptome assembly with Trinity. TransDecoder v5.5.0 (Haas et al., 2013) was used

to predict open reading frames (ORFs); transcripts for which no ORF was predicted were

removed from further analysis. Proteomes from each major bacterial clade (cf. Schulz et al.,

2017), fungi, stramenopiles, poriferans, arthropods, molluscs, and annelids were obtained from

NCBI's GenBank and concatenated into a non-target (alien) BLAST database. Target cnidarian

(corals and other cnidarians) and Symbiodinaceae protein sequences were obtained from the

Uniprot database. Alien Index (Ryan, 2014) was used in conjunction with protein BLAST

searches of our predicted ORFs (proteome) against non-target, alien, and target protein databases

to identify and remove likely contaminant sequences from our *de novo* assembled transcriptome.

This strategy yielded a meta-transcriptome consisting of coral and Symbiodinaceae transcripts

and predicted ORFs. This meta-transcriptome was then annotated with gene ontology (GO)

terms (Ashburner et al., 2004) retaining the top hit from BLAST searches (e-value <= 1e-5)

against the combined cnidarian and Symbiodinaceae Uniprot database. The reference meta-

transcriptome was then parsed into *Porites lobata* host and Symbiodniaceae symbiont

transcriptomes for differential gene analysis.

The Benchmarking Universal SingleCopy Orthologs (BUSCO) (Simão et al., 2015)

pipeline was used to determine the completeness of both the *P. lobata* and Symbiodinaceae

transcriptomes. BUSCO runs were performed on the transcript level, comparing transcriptome

assemblies against the metazoan database for *P. lobata* and alveolates for Symbiodinaceae. In *de*

novo transcriptome assemblies, it is common to find several predicted isoforms for the same

gene, which can lead to high duplication rates in BUSCO results. To adjust for such artificially

inflated duplication rates, transcripts originating from the same gene graph in the assembly that

are inferred to originate from the same gene were counted as a single hit in BUSCO.

2.5 | Differential gene expression and gene ontology

Transcript and gene abundances were inferred using the fast k-mer hashing and pseudoalignment

algorithm implemented in Kallisto (v0.46.2; Pimentel et al., 2017). The R package Sleuth (v.

0.30.0; Pimentel et al., 2017) was used to identify differentially expressed transcripts (*q*-value <

0.01). Significance levels of transcripts belonging to the same gene (see BUSCO analysis under

2.4) were aggregated (*p*-value aggregation method; Pimentel et al., 2017) to infer differential

expression at the gene, rather than transcript, level. Time series differential analysis based on

natural spline models (i.e., a likelihood ratio test) was used to discern patterns of expression

across all time points. A natural spline model (df = 4) was used to fit "knots" along the

observations of the time axis (full model) to determine if gene expression followed a non-random

pattern compared to a model of random variation (null model). A likelihood ratio test, as

implemented in Sleuth, was employed to compare full and null models for each gene to identify

differentially expressed genes along the time series.

Genes identified as differentially expressed were clustered hierarchically to identify

likely co-expressed genes. Clusters of genes with similar expression patterns were identified

after visual examination of the clustering dendrogram and resulting heatmap. These clusters

formed the basis for a Gene ontology (GO) enrichment analysis that was performed using

REVIGO version (Supek et al., 2011). In addition, pairwise Wald tests, as implemented in

Sleuth, were used to identify differentially expressed genes between time points.

3 | RESULTS

3.1 | Coral host and algal symbiont identities

Coral mitochondrial sequences (NCBI Accession nos. OM858841-OM858852) were invariant

across the almost 2,250bp sequenced apart from a single nucleotide polymorphism in specimen

five. Sequence similarity searches against NCBI's GenBank identified the mitochondrial genome

of *Porites lobata* (KU572435;Tisthammer et al., 2016) as the most similar publicly available

sequence data. Query coverage was 100% for all specimens and loci, with similarities greater or

equal to 99.9%. Based on transcript mapping to the putative Symbiodiniaceae transcriptomes, the

algal symbionts for all samples were primarily (>93%) from the genus *Cladocopium*

(Supplementary Figure 1).

3.2 | Micro-fragment growth rates

All 42 *P. lobata* micro-fragments survived the *ex-situ* phase of the experiment without signs of disease, predation, or bleaching. On average, micro-fragments increased in surface area by 355.4% over the 8-week duration of the experiment in the flow-through seawater tanks (Figure 2). Growth rates were normally distributed (Shapiro-Wilk's test; p-value = 0.0665) and growth rates between replicate tanks were not significantly different (t=-0.52893; p-value = 0.273). Two distinct growth phases were identified (Repeated Measures ANOVA, df = 92; p-value = 0.000186): Phase A (weeks 0 – 2) characterized with rapid growth of tissue over cut margins of the micro-fragment and Phase B (weeks 3 – 8) characterized by new growth deposited onto the tile. Significant variation in growth rates during phase B was found between colonies (p-value = 0.002141), with colonies 1 through 4 continuing to grow rapidly while colonies 5 and 6 showed reduced growth rates after initial wound healing (Figure 2). Two weeks after outplanting, the newly grown tissue of half of all outplanted micro-fragments (16/32) showed signs of bleaching despite growing in a similar temperature regime as that of the *ex-situ* tanks. No signs of disease or predation were observed during the *ex-situ* culture or after outplanting.

Figure 2: The average (Mean±SE) weekly surface for the 42 *Porites* micro-fragments over the 8-week

experiment. T1-T5 represent the transcriptomic sampling points. Colored lines represent individual colony growth,

and the black line represents mean micro-frag growth.

3.3 | Reference assembly

Illumina sequencing generated between 6.9 and 54 million reads per sample with an average of

35 million reads per sample (Supplemental Table 1). Due to a low number of reads generated at

some timepoints, colony 5 (Supplemental Table 2) was removed from further analysis. Trinity *de*

novo transcriptome assembly generated 1,963,624 transcripts, which were filtered and annotated

to produce a reference transcriptome composed of 60,475 coral and 17,994 symbiont gene

models (Supplementary Table 1). BUSCO analysis indicated that the reference for the coral host

was almost complete (C: 93.7%; S: 13.0%; D: 80.7% [adjusted D: 4.5%]; F: 2.5%; M: 3.8%; n:

954) while the symbiont reference was roughly 70% complete (C: 69.6%; S: 39.2%; D: 30.4%

[adjusted D:23.4%]; F: 4.1%; M: 26.3%; n: 171) (Supplementary Table 3).

3.4 | Time series gene co-expression

265   A total of 2,282 coral genes and 44 symbiont genes were identified as differentially expressed by

266   the spline regression analysis (*p*-value < 0.01; Figure 3). Hierarchical clustering of significantly

267   differentially expressed genes clustered the samples by time point replicate and then by stress

268   event vs. non-stress event (columns in Figure 3).

Figure 3: Heat map of significantly differentially expressed genes identified by spline regression analysis. The dendrogram on the left corresponds to clustering of genes (rows) by expression differences across samples (columns). The first number of the column name corresponds to colony replicate and the second refers to the sampling timepoint. The gene expression patterns cluster the samples into two main clusters: the stress events (T2; fragmentation and T5; outplanting) and the more stable growth points (T1, T3, & T4). The color scale corresponds to the average number of reads mapping to each base across the whole gene in that sample.

The dendrogram of differentially expressed genes (rows in Figure 3) indicated six major patterns

of expression through the time series (Figure 4). Three of these patterns (4-6) consisted of

relatively few genes and produced no significant GO enrichment results and were thus removed

from further analysis. The remaining three clusters consisted of genes that were upregulated at

the stress events (T2 & T5; Figure 4b), genes that were downregulated during the stress events

(Figure 4c), and genes that were downregulated during the rapid initial Growth Phase A (Figure

4d). GO categories that were significantly upregulated at both stress events (Figure 4a, 1340

genes) include formation of cytoplasmic translation initiation complex (GO:0001732), protein

folding (GO:0006457), cell redox homeostasis (GO:0045454), generation of precursor

metabolites and energy (GO:0006091), ribose phosphate metabolism (GO:0019693), and

pigment metabolism (GO:0042440). Relevant GO categories that were downregulated at both

stress events (Figure 4c, 229 genes) included aromatic amino acid family catabolism

(GO:0009074), and protein localization to endoplasmic reticulum (GO:0070972). GO categories

that were significantly downregulated when the micro-frags were recently cut and rapidly

growing (Figure 4d, 506 genes) included cellular response to organic substance (GO:0071310),

nitrogen compound metabolism (GO:0006807), localization (GO:0051179), and organelle

assembly (GO:0070925) (See Supplementary text for a full list of differentially expressed GO

categories).

3.5 | Pairwise gene expression analysis

While analyzing coral host gene expression profiles during the transition between states, the

largest number of differentially expressed genes was identified immediately following

fragmentation (T1 *versus* T2) (Table 1). The transition between T2 and T3 yielded few

differentially expressed genes. Additionally, the comparison between the baseline (T1) and two

weeks of growth (T3) yielded few differentially expressed genes. The response to fragmentation

(T1 *versus* T2) yielded twice as many differentially expressed genes as the response to

outplanting (T4 *versus* T5). The list of significantly differentially expressed genes was manually

curated to identify crucial metabolic genes that represent fairly complete pathways and genes

that have been previously identified as part of the cnidarian stress response (Supplemental Table

4). Several crucial antioxidant enzymes were upregulated in response to stress in this study,

including Thioredoxin, Peroxiredoxin-1, Peptide-methionine (S)-S-oxide reductase (EC

1.8.4.11), and Glutathione transferase (EC 2.5.1.18) (Supplemental Table 4). Heat shock proteins

were upregulated in response to fragmentation and outplanting, such as HSP70-1, HSP90, 97

kDa heat shock protein, HSP40s, and 71kDa heat shock proteins (Supplemental Table 4). A

variety of protein degradation and biosynthesis enzymes were identified as differentially

expressed in response to both stress events (Supplemental Table 4), including Ubiquitin-

Table 1: The number of differentially expressed genes between all pairwise timepoints for coral host and symbiont.

Coral genes	T1(Baseline)	T2(Immediate response)	T3 (2 weeks of growth)	T4 (2 months of growth)
T1(Baseline)				
T2 (Immediate response)	1648			
T3 (2 weeks of growth)	73	64		
T4 (2 months of growth)	116	1794	2	
T5 (Outplanting)	1753	1207	299	820
Symbiodinium genes				
T1 (Baseline)				
T2 (Immediate response)	7			
T3 (2 weeks of growth)	0	7		
T4 (2 months of growth)	2	42	5	
T5 (Outplanting)	11	210	33	2

associated ligases, hydrolases, conjugation factors, and transferases, translation initiation factors, ribosomal biogenesis proteins, tRNA ligases, deacylases, synthetases, and hydrolases. Calcium homeostasis genes were significantly upregulated in response to fragmentation, such as Calreticulin, calumenin, Voltage-dependent L-type calcium channel subunit alpha, Calcium-transporting ATPase, and many genes which cause the release of calcium into the via the phosphatidylinositol signaling pathway. Energy metabolism genes relating to glycolysis, gluconeogenesis, the citric acid cycle, and electron transport chain were upregulated immediately following fragmentation (See supplementary text for detailed list of differentially expressed genes; Supplemental Table 4).

4 | DISCUSSION

Growth and calcification rates are negatively correlated with size in corals (Chadwick-Furman et

al., 2000) and both increase in response to fragmentation (Lirman et al., 2010; Forsman et al.,

2015). Rapid regeneration is crucial to coral colony survival, as it prevents possible infections by

pathogens and damage from competitors, and ensures the physical integrity of the coral colony

by cementing fragments to each other and the substrate after breakage or fracture. Here, we show

that fragmentation triggers a disruption in calcium homeostasis and initiates rapid wound healing

and recovery in *Porites lobata*. The fragmentation response is characterized by two distinct

phases: (1) rapid tissue regeneration at wound margins, followed by (2) a slower growth phase

during which the colony is stabilized by new tissue deposited onto the substrate (Figure 2). The

initial increase in growth rate following fragmentation of coral colonies has been demonstrated

previously (Page et al., 2018), but the underlying cellular-level processes driving the accelerated

growth have remained largely unexplored and are crucial for understanding coral regeneration.

The initial stress response to both outplanting and fragmentation yielded specific genes

(Supplemental Table 4) and GO terms (Figure 4a) associated with ER stress, protein turnover,

and antioxidant production, which is consistent with a generalized stress response seen in other

cnidarians (Maor-Landaw & Levy, 2016). However, fragmentation initiated some unique

processes, such as upregulation of phosphoinositide-mediated acute calcium homeostasis

disruption and a variety of electron transport genes, which we believe stimulate and provide

immediate energy for the rapid growth of the coral colony in a push for wound healing, and

ultimately, survival.

Overall, the immediate fragmentation response was characterized by the largest number

of differentially regulated genes (1648; q-value = 0.01) in our experiment, indicating that

fragmentation elicits a series of immediate and unique physiological responses (Table 1). During

this initial phase, the margins of the fragment were sealed quickly by new tissue with growth

slowing once margins were sealed. Concomitantly with this slowing in growth, gene expression

stabilized with only two genes differentially expressed when comparing the two-week time point

(T3) to the two-month time point (T4), as well as the baseline (T1) to T3 (Table1). We attribute

this finding to the recovery of the coral host from the initial stress response to fragmentation and

an indication of the return to metabolic homeostasis during this period.

Figure 5: Cell diagram of the integration of calcium homeostasis disruption (A), Endoplasmic-Reticulum

(ER) stress (B), protein anabolism (D) increased energy production (C), and protein degradation (D) found

upregulated 24 hours after physical injury. 5A: Phosphoinositide signaling releases the secondary intracellular messenger, inositol (1,4,5) trisphosphate (IP3), which binds to ligand-gated calcium ion channels of the ER causing the release of calcium into the cytosol. Calcium disruption is sensed by a variety of molecules including calmodulin (upregulated in this study), which leads to enzymatic activation and ultimately, cell cycle regulation. Calmodulin also stimulates the reuptake of Ca^{2+} into the ER to prevent prolonged calcium homeostasis disruption and cell death. 5B: Calcium disruption within the ER leads to the Unfolded Protein Response (UPR). Resident proteins upregulated in this study, such as calreticulin, calumenin, and Heat Shock Proteins (HSPs), assist in the folding of proteins and degradation of terminally misfolded proteins. Calreticulin prevents the export of misfolded proteins to the golgi apparatus. 5C: Ribosomal proteins, translation initiation factors, and tRNA enzymes were significantly upregulated and rapidly produced new proteins in response to fragmentation. 5D: To meet energetic demands of the cell under stress, expression of electron transport chain proteins increased. Increased oxidative phosphorylation causes Reactive Oxygen Species (ROS) to leak from the mitochondria, causing cellular damage to lipids, DNA, and proteins. Antioxidant molecules upregulated in this study (thioredoxin, glutathione transferase, peroxiredoxin, ferritin) scavenge ROS and assist in protein refolding. Cytosolic HSPs refold damaged proteins and assist in their degradation if they are terminally misfolded. 5E. Terminally misfolded proteins are chaperoned to the proteasome by HSPs after they are tagged for destruction by Ubiquitin-conjugating enzymes. The Ubiquitin/Proteosome system integrates with other signal transduction molecules to regulate cell cycle. Amino acid transporters shuttle degraded polypeptides to ribosomes for protein anabolism.

4.1 | Stress response and energy metabolism

When organisms are presented with environmental stress, the balance between ROS production and antioxidant defenses is disrupted. This frequently leads to an increase in production of antioxidant compounds to prevent significant cellular damage to lipids, proteins, DNA and, ultimately, cell death via apoptosis (Gorman et al., 1999; Scandalios, 2002). Energy generation is crucial in providing necessary resources to enact a shift in metabolism and survival in stressful conditions. However, many cnidarian transcriptomic studies find decreases in energy generation

during stress (Veal et al., 2002; Császár et al., 2009; Tarrant et al., 2014; Maor-Landaw & Levy,

2016; Aguilar et al., 2019), which may be related to the breakdown of symbiosis. In contrast,

both our spline regression analysis (Figures 3 and 4) and pairwise comparisons (Figure 5 &

Supplemental Table 4) provided significant evidence of increased cellular energy generation

(oxidative phosphorylation, electron transport chain cytochromes, glycolysis, and citric acid

cycle enzymes) and lack a detectable gene expression response in the symbiont following coral

fragmentation. The maintenance of symbiosis and continued energy production likely provide

the resources necessary for antioxidant defenses, cellular protein maintenance, and rapid wound

healing (Figure 5). Many of the metabolic pathways identified as differentially expressed in this

study (misfolded protein response, HSPs, cell-redox homeostasis, cytoskeleton rearrangement,

and calcium homeostasis disruption) coincide with the generalized oxidative stress response

described for cnidarians regardless of the type of stress (Veal et al., 2002; Császár et al., 2009;

Tarrant et al., 2014; Maor-Landaw & Levy, 2016; Aguilar et al., 2019). However, fragmentation

of *P. lobata* appears to elicit several unique responses, such as phosphoinositide-mediated acute

calcium homeostasis disruption, energy production, and low gene expression differences in the

symbiont.

Our study found several antioxidant molecules (heat-shock proteins, thioredoxins,

ferritin) (Supplemental Table 4), as well as GO categories associated with cell-redox

homeostasis (Figure 4b; Pattern 1), were upregulated in response to both fragmentation and

outplanting. The upregulation of genes that participate in the thioredoxin oxidoreductase system

(thioredoxin, peroxiredoxin-1, Peptide-methionine (S)-S-oxide reductase (EC 1.8.4.11), and

glutathione transferase), ferritin, and ion transporters in response to both fragmentation and

outplanting further suggest significant increase in antioxidant and protein repair mechanisms to

combat oxidative damage from increased energy production necessary for wound regeneration

and survival.

Antioxidant molecules that were once thought of as unique biomarkers for heat stress,

such as heat-shock proteins (HSPs), are now understood to be upregulated in response to a

variety of stressors, including physical injury in this study. Several studies have suggested that

the absence of a HSP response is linked with the initiation of apoptosis pathways (Feder &

Hofmann, 1999; Gorman et al., 1999; Samali et al., 1999). Thus, the presence of upregulated

HSPs indicates a push for survival. In response to physical injury, *P. lobata* colonies

demonstrated a diverse upregulation of mitochondrial, cytosol, and ER resident HSPs (HSP70,

HSP90, 95kDA HSP, 10kDA HSP, DnaJ-like HSP40s). Arguably the most studied and

ubiquitous HSP, HSP70 (Samali et al., 1999), was the strongest upregulated oxidative stress-

associated gene in response to fragmentation ($\log_2$FC 2.95; FDR $1.2e^{-9}$). The upregulation of 97

kDA HSP in response to fragmentation and subsequent downregulation after two weeks of

growth may suggest that oxidative damage to proteins begins to subside within two weeks

following physical injury. Additionally, no antioxidant enzymes were upregulated after either

two weeks or two months of growth, suggesting that the increased ROS, and therefore

antioxidant demands, diminished within a few weeks of physical injury. Although the

thioredoxin system and HSPs remain the most studied and indicative biomarkers for cellular

oxidative stress, further study is required to understand their specific roles in protein folding,

assembly, regulation, and degradation in cnidarians (Feder & Hofmann, 1999; Maor-Landaw &

Levy, 2016).

4.2 | Endoplasmic reticulum stress response and protein turnover

The endoplasmic reticulum (ER) plays crucial roles in controlling protein quality, facilitating the

degradation of misfolded proteins and sensing homeostasis changes such as the release of Ca^{2+}

into the cytosol. When significant protein damage and calcium homeostasis disruption occur, ER

stress pathways are stimulated leading to increased protein stabilization, degradation, and

synthesis required to preserve cellular function (Bahar et al., 2016). Significant ER stress during

both fragmentation and outplanting was indicated by the upregulation of GO terms associated

with protein turnover (protein catabolism, anabolism, folding) (Figure 4b), chaperones involved

in protein stabilization and refolding (HSPs, thioredoxin, calreticulin and calumenin), and

ubiquitin-associated degradation enzymes (Supplemental Table 4). We found that GO terms

associated with amino acid catabolism (Figure 4b) were downregulated in response to

fragmentation and outplanting, which suggests that amino acids were preserved to provide the

necessary building blocks for increased protein anabolism as corals respond to stress events

(Klasing, 2009).

Interestingly, we did not find any evidence for the upregulation of specific ER-stress

signal transducers known that have been described from other organisms (i.e., inositol-requiring

protein-1 (IRE1), activating transcription factor-6 (ATF6), or protein kinase RNA (PKR)-like ER

kinase (PERK); Ron & Walter, 2007). Similar to calcium homeostasis disruption (see below), the

duration and severity of ER stress determines if an adaptive survival response persists or

apoptosis is initiated (Szegezdi et al., 2006; Bahar et al., 2016). The fact that ER-stress signal

transducers were not upregulated one day after outplanting suggests that the fragments were

already recovering from initial transplantation shock and stress. Decreased rates of translation

initiation is one of the earliest stages of significant ER stress, yet we found significant

459   upregulation of translation initiation factors (17/17 identified) one day after fragmentation

460   (Supplemental Table 4). This suggests that corals were recovering from initial ER stress as early

461   as one day after physical injury, upregulating antioxidant defenses, protein stabilizing molecules,

462   and, ultimately, rapid wound regeneration in a push for survival.

4.3 | Calcium homeostasis disruption and its role in rapid regeneration

Calcium signaling pathways are ubiquitous signal transduction systems, which regulate a broad

variety of cellular processes, such as metabolism, apoptosis, cell proliferation, cell to cell

communication, gene expression, and secretion (Stefan, 2020). Many cnidarian stress

experiments report calcium homeostasis disruption, but few have elucidated the cell signaling

mechanisms causing this disruption (Pinzón et al., 2015; Oakley et al., 2017; Aguilar et al.,

2019). Among signaling molecules, membrane-bound phospholipids of the phosphoinositide

family play key roles in regulating the activity of proteins within the cell (reviewed in

Falkenburger et al., 2010). Phosphoinositide signaling networks are composed of a series of

transmembrane G-coupled proteins that regulate diverse functions in the cell, including the

release of Ca^{2+} from the ER via the binding of inositol 1,4,5-trisphosphate (IP_3) to ligand-gated

ion channel receptors in the membrane of the ER (Berridge & Irvine, 1989; Stefan, 2020).

Although well characterized in other organisms, the specific roles of phosphoinositide signaling

in cnidarians remain uncharacterized and the differential expression of these complexes have

been reported rarely (Oakley et al., 2017; Aguilar et al., 2019). For example, phosphoinositide

phospholipase C, a crucial enzyme that catalyzes the release of IP_3, was downregulated in

response to thermal stress in the anemone model system *Aiptasia* (Oakley et al., 2017). We found

that fragmentation caused the upregulation of rate-limiting components of the phosphoinositide

signaling pathway and several growth receptors known to stimulate this pathway, including

Phosphoinositide phospholipase C and tyrosine kinase growth factor receptors (Supplemental

Table 4). We hypothesize that this response was related to the onset of rapid growth and

increased calcification rates following fragmentation.

While initial calcium homeostasis disruption can stimulate survival pathways, sustained

levels of calcium in the cytosol lead to the initiation of apoptosis (Orrenius et al., 2003; Bagur &

Hajnóczky, 2017). Voltage-dependent and ATPase calcium pumps maintain cytosolic Ca^{2+} levels

by sequestering calcium in the ER or secreting it extracellularly. We found the genes coding for

calcium ATPase pumps upregulated in response to fragmentation, concomitant with calcium

homeostasis disruption genes belonging to the phosphoinositide signaling pathway

(Supplemental Table 4). The concurrent upregulation of calcium-releasing mechanisms

(phosphoinositide signaling) and calcium-dependent ATPase pumps immediately following

fragmentation during a period of accelerated growth (Figure 5) suggests that calcium

homeostasis is acutely disrupted by physical injury, stimulating cell proliferation and wound

regeneration. To promote cell proliferation rather than trigger apoptosis, calcium is released into

the cytosol, likely via the phosphoinositide signaling pathway, to stimulate calmodulin-mediated

control of the cell cycle. Excess cytosolic Ca^{2+} is then translocated against a gradient using

ATPase pumps to restore calcium levels in the ER lumen (Figure 5).

Ca^{2+} increases in the cytosol are sensed by various Ca^{2+}-binding proteins and we found

that the following genes encoding for Ca^{2+}-binding proteins were differentially expressed:

calmodulin (CaM), calreticulin, and calumenin (Supplemental Table 4). Ca^{2+}-binding proteins

are well-known for their role in initiating signaling cascades that ultimately lead to shifts in

metabolism. Contrary to the findings of other cnidarian stress experiments (e.g., DeSalvo et al.,

2010; Aguilar et al., 2019), CaM was upregulated in response to fragmentation and subsequently

downregulated after two weeks of rapid growth. This corroborates our hypothesis that Ca^{2+}

concentrations were returning to stable levels in a push for cellular survival. We suggest that

CaM was responsible for the initial calcium homeostasis disruption that triggered rapid

regeneration after injury by fragmentation in our experiment. Interestingly, CaM has been

identified as a regulator of coral larval settlement and metamorphosis (Reyes-Bermudez et al.,

2012, 2016), a period in early development characterized by rapid growth.

In addition, we found that the genes coding for the ER-resident proteins calreticulin and

calumenin which, together with HSP40 (DnaJ chaperone), prevent the export of misfolded

proteins, were upregulated in response to fragmentation (Supplemental Table 4). Increased

mitochondrial ROS production (see *stress response and energy metabolism* above) increases

oxidative stress that may disrupt protein folding processes, elevating ER stress. We identified the

mechanisms that mitigate the impacts of elevated ROS production caused by increased metabolic

activity in corals following fragmentation. In particular, calreticulin prevents the export of

misfolded proteins from the ER to the Golgi apparatus. Instead, misfolded or damaged proteins

are exported to proteasomes for degradation and recycling. Taken together, our gene expression

results put the ER at the center of CaM-mediated cell cycle modifications and mitigation of ROS

stress impacts on protein synthesis in the aftermath of fragmentation (Figure 5).

4.4 | Impacts of physical injury on the host-symbiont relationship

Fewer than ten genes were differentially regulated in the zooxanthellate symbiont community

between time points (Table 1). The absence of differentially expressed symbiont antioxidant

genes suggests that ROS are unlikely to affect the endosymbionts during coral regeneration.

Additionally, we found no evidence of nitric oxide (NO) homeostasis breakdown within the

host's response to physical fragmentation, which has been previously correlated with the

breakdown of symbiosis in response to coral bleaching (Perez & Weis, 2006; DeSalvo et al.,

2010; Maor-Landaw & Levy, 2016). Together, these results suggest that the coral-zooxanthellae

symbiosis is far less affected by physical injury than it is by other environmental stressors.

Continued efficacy of the host-symbiotic relationship may help meet the energetic needs of the

coral host for cellular protein maintenance, antioxidant defense mechanisms, and rapid wound

regeneration in response to physical injury.

4.5 | Physiology of fragmentation response

This study provides a model of the physiological mechanisms that enable rapid recovery and

stabilization of corals in response to physical injury (Figure 5). The response of *Porites lobata* to

fragmentation and outplanting displays many similarities with the response of corals to other

stress events, but we documented several important differences. Environmental stress events

usually lead to decreased energy production (Maor-Landaw & Levy, 2016), reduced calcification

rates (Anthony et al., 2008; Crook et al., 2013; D'Olivo & McCulloch, 2017), and increased

mortality (Glynn, 1990) in corals. By contrast, we found that the acute response to physical

injury by fragmentation was characterized by upregulation of genes linked to increased energy

production. Further, phosphoinositide-mediated calcium homeostasis disruption and ER stress

led to the increased expression of antioxidant defense mechanisms and increased rates of protein

turnover. Phosphoinositide-mediated acute calcium homeostasis disruption is a likely key

mechanism stimulating the wound healing and recovery processes that are evident from the rapid

growth of fragmented corals documented here and by others previously (e.g., Page et al., 2018).

The severity and length of ER stress and calcium homeostasis disruption often determines the

fate of cells in response to environmental change: survival or initiation of apoptosis (Orrenius et

al., 2003; Bahar et al., 2016). The concurrent upregulation of calcium disruption and

sequestering signals, translation initiation factors and ER stress-associated molecules (protein

chaperones and ubiquitin-degradation enzymes), as well as heat shock proteins, in response to

physical injury indicates that fragmented corals push for survival and recovery very quickly

following injury. In our experiment, this rapid immediate response ceased after two weeks when

the exposed margins of the coral skeleton were sealed again with new tissue. All of these

responses are consistent with a life history strategy that utilizes fragmentation as a means of

asexual propagation and broaden our understanding of this fundamental cnidarian process.

ACKNOWLEDGEMENTS

This work received support from the University of Guam Sea Grant Master's Thesis Research

Supplemental Grants and the National Science Foundation's Established Program to Stimulate

Competitive Research (OIA-1457769 and OIA-1946352).

AUTHOR CONTRIBUTIONS

CL, BB, & LJR conceived the project. CL performed the experiments and collected the data. CL

and BB analyzed the data. All authors contributed equally to writing the paper.

ORCID

Colin Lock https://orcid.org/0000-0002-8837-5936

*Bastian Bentlag*e https://orcid.org/0000-0002-9554-3125

Laurie Raymundo https://orcid.org/0000-0003-3446-0390

5 | References

Aguilar, C., Raina, J. B., Fôret, S., Hayward, D. C., Lapeyre, B., Bourne, D. G., & Miller, D. J.
576   (2019). Transcriptomic analysis reveals protein homeostasis breakdown in the coral
577   Acropora millepora during hypo-saline stress. *BMC Genomics*, *20*(1), 1–13.
578   https://doi.org/10.1186/s12864-019-5527-2

579   Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S., & Hoegh-Guldberg, O. (2008). Ocean
580   acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of*
581   *the National Academy of Sciences of the United States of America*, *105*(45), 17442–17446.
582   https://doi.org/10.1073/pnas.0804478105

583   Bagur, R., & Hajnóczky, G. (2017). Intracellular Ca2+ Sensing: Its Role in Calcium
584   Homeostasis and Signaling. *Molecular Cell*, *66*(6), 780–788.
585   https://doi.org/10.1016/j.molcel.2017.05.028

586   Bahar, E., Kim, H., & Yoon, H. (2016). ER stress-mediated signaling: Action potential and Ca2+
587   as key players. *International Journal of Molecular Sciences*, *17*(9), 1–22.
588   https://doi.org/10.3390/ijms17091558

589   Bak, R. P. M. (1983). Neoplasia, regeneration and growth in the reef-building coral Acropora
590   palmata. *Marine Biology*, *77*(3), 221–227. https://doi.org/10.1007/BF00395810

591   Bak, R. P. M., & Yvonne, S.-V. E. (1980). Regeneration of Superficial Damage in the
592   Scleractinian Corals Agaricia Agaricites F. Purpurea and Porites Astreoides. *Bulletin of*
593   *Marine Science*, *30*(4), 883–887. Retrieved from
594   http://www.ingentaconnect.com/content/umrsmas/bullmar/1980/00000030/00000004/art000
595   10

596   Barton, J. A., Willis, B. L., & Hutson, K. S. (2017). Coral propagation: a review of techniques
597   for ornamental trade and reef restoration. *Reviews in Aquaculture*, *9*(3), 238–256.
598   https://doi.org/10.1111/raq.12135

599   Baums, I. B. (2008). A restoration genetics guide for coral reef conservation. *Molecular Ecology*,
600   *17*(12), 2796–2811. https://doi.org/10.1111/j.1365-294X.2008.03787.x

601   Bay, L. K., Guérécheau, A., Andreakis, N., Ulstrup, K. E., & Matz, M. V. (2013). Gene
602   Expression Signatures of Energetic Acclimatisation in the Reef Building Coral Acropora
603   millepora. *PLoS ONE*, *8*(5), e61736. https://doi.org/10.1371/journal.pone.0061736

604   Berridge, M. J., & Irvine, R. F. (1989). Inositol phosphates and cell signalling. *Nature*,
605   *341*(6239), 197–205. https://doi.org/10.1038/341197a0

606   Chadwick-Furman, N. E., Goffredo, S., & Loya, Y. (2000). Growth and population dynamic
607   model of the reef coral Fungia granulosa Klunzinger, 1879 at Eilat, northern Red Sea.

Journal of Experimental Marine Biology and Ecology, *249*(2), 199–218.
https://doi.org/10.1016/S0022-0981(00)00204-5

Crook, E. D., Cohen, A. L., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A. (2013). Reduced
calcification and lack of acclimatization by coral colonies growing in areas of persistent
natural acidification. *Proceedings of the National Academy of Sciences*, *110*(27), 11044–
11049. https://doi.org/10.1073/pnas.1301589110

Császár, N. B. M., Seneca, F. O., & Van Oppen, M. J. H. (2009). Variation in antioxidant gene
expression in the scleractinian coral Acropora millepora under laboratory thermal stress.
Marine Ecology Progress Series, *392*(May 2014), 93–102.
https://doi.org/10.3354/meps08194

D'Olivo, J. P., & McCulloch, M. T. (2017). Response of coral calcification and calcifying fluid
composition to thermally induced bleaching stress. *Scientific Reports*, *7*(1), 1–15.
https://doi.org/10.1038/s41598-017-02306-x

DeSalvo, M. K., Estrada, A., Sunagawa, S., & Medina, M. (2012). Transcriptomic responses to
darkness stress point to common coral bleaching mechanisms. *Coral Reefs*, *31*(1), 215–228.
https://doi.org/10.1007/s00338-011-0833-4

DeSalvo, M. K., Sunagawa, S., Voolstra, C., & Medina, M. (2010). Transcriptomic responses to
heat stress and bleaching in the elkhorn coral Acropora palmata. *Marine Ecology Progress*
Series, *402*, 97–113. https://doi.org/10.3354/meps08372

Desalvo, M. K., Voolstra, C. R., Sunagawa, S., Schwarz, J. A., Stillman, J. H., Coffroth, M. A.,
… Medina, M. (2008). Differential gene expression during thermal stress and bleaching in
the Caribbean coral Montastraea faveolata. *Molecular Ecology*, *17*(17), 3952–3971.
https://doi.org/10.1111/j.1365-294X.2008.03879.x

Edwards, A. J. (2008). *Reef Rehabilitation Manual. Coral Reef Targeted Research & Capacity*
Building for Management Program: St Lucia, Australia. Construction (Vol. 501). The Coral
Reef Targeted Research & Capacity Building for Management Program.

Edwards, A. J., & Gomez, E. D. (2011). Reef Restoration Concepts and Guidelines: making
sensible management choices in the face o f uncertainty. *Encyclopedia of Earth Sciences*
Series, *Part 2*, 889–896. https://doi.org/10.1007/978-90-481-2639-2_140

Epstein, N., Bak, R. P. M., & Rinkevich, B. (2001). Strategies for gardening denuded coral reef
areas: The applicability of using different types of coral material for reef restoration.
Restoration Ecology, *9*(4), 432–442. https://doi.org/10.1046/j.1526-100X.2001.94012.x

Falkenburger, B. H., Jensen, J. B., & Hille, B. (2010). Kinetics of M 1 muscarinic receptor and G
protein signaling to phospholipase C in living cells. *Journal of General Physiology*, *135*(2),
81–97. https://doi.org/10.1085/jgp.200910344

Feder, M. E., & Hofmann, G. E. (1999). Heat-shock proteins, molecular chaperones, and the
stress response: Evolutionary and ecological physiology. *Annual Review of Physiology*, *61*,
243–282. https://doi.org/10.1146/annurev.physiol.61.1.243

Forsman, Z. H., Page, C. A., Toonen, R. J., & Vaughan, D. (2015). Growing coral larger and
faster: micro-colony-fusion as a strategy for accelerating coral cover. *PeerJ*, *3*, e1313.

https://doi.org/10.7717/peerj.1313

Forsman, Z. H., Rinkevich, B., & Hunter, C. L. (2006). Investigating fragment size for culturing
reef-building corals (Porites lobata and P. compressa) in ex situ nurseries.
https://doi.org/10.1016/j.aquaculture.2006.06.040

Glynn, P. W. (1990). Coral Mortality and Disturbances to Coral Reefs in the Tropical Eastern
Pacific. *Elsevier Oceanography Series*, *52*(C), 55–126. https://doi.org/10.1016/S0422-
9894(08)70033-3

Gorman, A. M., Heavey, B., Creagh, E., Cotter, T. G., & Samali, A. (1999). Antioxidant-
mediated inhibition of the heat shock response leads to apoptosis. *FEBS Letters*, *445*(1), 98–
102. https://doi.org/10.1016/S0014-5793(99)00094-0

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., … Regev,
A. (2013). De novo transcript sequence reconstruction from RNA-seq using the Trinity
platform for reference generation and analysis. *Nature Protocols*, *8*(8), 1494–1512.
https://doi.org/10.1038/nprot.2013.084

Harrison, P. L., & Wallace, C. C. (1990). A review of reproduction, larval dispersal and
settlement of scleractinian corals. In *Chapter 7 in Ecosystems of the World 25 Coral Reefs
(ed. Z. Dubinsky).* (pp. 133–196). Retrieved from
https://www.researchgate.net/publication/304784746_Harrison_PL_and_Wallace_C_C_199
0_A_review_of_reproduction_larval_dispersal_and_settlement_of_scleractinian_corals_Ch
apter_7_in_Ecosystems_of_the_World_25_Coral_Reefs_ed_Z_Dubinsky_Amsterdam_Else
vier_p

Highsmith, R. (2007). Reproduction by Fragmentation in Corals. *Marine Ecology Progress
Series*, *7*, 207–226. https://doi.org/10.3354/meps007207

Klasing, K. C. (2009). Minimizing amino acid catabolism decreases amino acid requirements.
Journal of Nutrition, *139*(1), 11–12. https://doi.org/10.3945/jn.108.099341

Kojis, B. L., & Quinn, N. J. (2001). The importance of regional differences in hard coral
recruitment rates for determining the need for coral restoration. *Bulletin of Marine Science*,
69(2), 967–974. https://doi.org/10.1098/rsbl.2003.0087

Kozłowski, J., & Wiegert, R. G. (1986). Optimal allocation of energy to growth and
reproduction. *Theoretical Population Biology*, *29*(1), 16–37. https://doi.org/10.1016/0040-
5809(86)90003-1

LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C.
R., & Santos, S. R. (2018). Systematic Revision of Symbiodiniaceae Highlights the
Antiquity and Diversity of Coral Endosymbionts. *Current Biology*, *28*(16), 2570-2580.e6.
https://doi.org/10.1016/j.cub.2018.07.008

Lirman, D. (2000). Fragmentation in the branching coral Acropora palmata (Lamarck): Growth,
survivorship, and reproduction of colonies and fragments. *Journal of Experimental Marine
Biology and Ecology*, *251*(1), 41–57. https://doi.org/10.1016/S0022-0981(00)00205-7

Lirman, D., Thyberg, T., Herlan, J., Hill, C., Young-Lahiff, C., Schopmeyer, S., … Drury, C.
(2010). Propagation of the threatened staghorn coral Acropora cervicornis: Methods to

minimize the impacts of fragment collection and maximize production. *Coral Reefs*, *29*(3),
729–735. https://doi.org/10.1007/s00338-010-0621-6

Lough, J. M., & Barnes, D. J. (2000). Environmental controls on growth of the massive coral
Porites. *Journal of Experimental Marine Biology and Ecology*, *245*(2), 225–243.
https://doi.org/10.1016/S0022-0981(99)00168-9

Manzello, D. P., Matz, M. V., Enochs, I. C., Valentino, L., Carlton, R. D., Kolodziej, G., …
Jankulak, M. (2019). Role of host genetics and heat-tolerant algal symbionts in sustaining
populations of the endangered coral Orbicella faveolata in the Florida Keys with ocean
warming. *Global Change Biology*, *25*(3), 1016–1031. https://doi.org/10.1111/gcb.14545

Maor-Landaw, K., & Levy, O. (2016). Survey of Cnidarian Gene Expression Profiles in
Response to Environmental Stressors: Summarizing 20 Years of Research, What Are We
Heading for? In *The Cnidaria, Past, Present and Future* (pp. 523–543).
https://doi.org/10.1007/978-3-319-31305-4

Meesters, E. H., Noordeloos, M., & Bak, R. P. M. (1994). Damage and regeneration: Links to
growth in the reef-building coral Montastrea annularis. *Marine Ecology Progress Series*,
112(1–2), 119–128. https://doi.org/10.3354/meps112119

Oakley, C. A., Durand, E., Wilkinson, S. P., Peng, L., Weis, V. M., Grossman, A. R., & Davy, S.
K. (2017). Thermal Shock Induces Host Proteostasis Disruption and Endoplasmic
Reticulum Stress in the Model Symbiotic Cnidarian Aiptasia. *Journal of Proteome*
Research, *16*(6), 2121–2134. https://doi.org/10.1021/acs.jproteome.6b00797

Okubo, N., Motokawa, T., & Omori, M. (2007). When fragmented coral spawn? Effect of size
and timing on survivorship and fecundity of fragmentation in Acropora formosa. *Marine*
Biology, *151*(1), 353–363. https://doi.org/10.1007/s00227-006-0490-2

Oren, U., Benayahu, Y., Lubinevsky, H., & Loya, Y. (2012). Colony Integration during
Regeneration in the Stony Coral Favia favus, *82*(3), 802–813.

Orrenius, S., Zhivotovsky, B., & Nicotera, P. (2003). Regulation of cell death: The calcium-
apoptosis link. *Nature Reviews Molecular Cell Biology*, *4*(7), 552–565.
https://doi.org/10.1038/nrm1150

Page, C. A., Muller, E. M., & Vaughan, D. E. (2018). Microfragmenting for the successful
restoration of slow growing massive corals. *Ecological Engineering*, *123*, 86–94.
https://doi.org/10.1016/j.ecoleng.2018.08.017

Perez, S., & Weis, V. (2006). Nitric oxide and cnidarian bleaching: An eviction notice mediates
breakdown of a symbiosis. *Journal of Experimental Biology*, *209*(14), 2804–2810.
https://doi.org/10.1242/jeb.02309

Pimentel, H., Bray, N. L., Puente, S., Melsted, P., & Pachter, L. (2017). Differential analysis of
RNA-seq incorporating quantification uncertainty. *Nature Methods*, *14*(7), 687–690.
https://doi.org/10.1038/nmeth.4324

Pinzón, J. H., Kamel, B., Burge, C. A., Harvell, C. D., Medina, M., Weil, E., & Mydlarz, L. D.
(2015). Whole transcriptome analysis reveals changes in expression of immune-related
genes during and after bleaching in a reef-building coral. *Royal Society Open Science*, *2*(4).

https://doi.org/10.1098/rsos.140214

Reyes-Bermudez, A., Miller, D. J., & Sprungala, S. (2012). The Neuronal Calcium Sensor
Protein Acrocalcin: A Potential Target of Calmodulin Regulation during Development in
the Coral Acropora millepora. *PLoS ONE*, *7*(12).
https://doi.org/10.1371/journal.pone.0051689

Reyes-Bermudez, A., Villar-Briones, A., Ramirez-Portilla, C., Hidaka, M., & Mikheyev, A. S.
(2016). Developmental progression in the coral acropora digitifera is controlled by
differential expression of distinct regulatory gene networks. *Genome Biology and Evolution*,
8(3), 851–870. https://doi.org/10.1093/gbe/evw042

Rinkevich, B. (2005). Conservation of Coral Reefs through Active Restoration Measures: Recent
Approaches and Last Decade Progress. https://doi.org/10.1021/ES0482583

Ron, D., & Walter, P. (2007). Signal integration in the endoplasmic reticulum unfolded protein
response. *Nature Reviews Molecular Cell Biology*, *8*(7), 519–529.
https://doi.org/10.1038/nrm2199

Ryan, J. (2014). Alien Index: identify potential non-animal transcripts or horizontally transferred
genes in animal transcriptomes. https://doi.org/http://dx.doi.org/10.5281/zenodo.21029

Samali, A., Holmberg, C. I., Sistonen, L., & Orrenius, S. (1999). Thermotolerance and cell death
are distinct cellular responses to stress: Dependence on heat shock proteins. *FEBS Letters*,
461(3), 306–310. https://doi.org/10.1016/S0014-5793(99)01486-6

Scandalios, J. G. (2002). Oxidative stress responses - What have genome-scale studies taught us?
Genome Biology, *3*(7), 1–6. https://doi.org/10.1186/gb-2002-3-7-reviews1019

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics*, *31*(19), 3210–3212.
https://doi.org/10.1093/bioinformatics/btv351

Smith, & Hughes, T. P. (1999). Experimental assessment of coral fragments, *235*, 147–164.

Stefan, C. J. (2020). Endoplasmic reticulum–plasma membrane contacts: Principals of
phosphoinositide and calcium signaling. *Current Opinion in Cell Biology*, *63*, 125–134.
https://doi.org/10.1016/j.ceb.2020.01.010

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). Revigo summarizes and visualizes long
lists of gene ontology terms. *PLoS ONE*, *6*(7). https://doi.org/10.1371/journal.pone.0021800

Szegezdi, E., Logue, S. E., Gorman, A. M., & Samali, A. (2006). Mediators of endoplasmic
reticulum stress-induced apoptosis. *EMBO Reports*, *7*(9), 880–885.
https://doi.org/10.1038/sj.embor.7400779

Tarrant, A. M., Reitzel, A. M., Kwok, C. K., & Jenny, M. J. (2014). Activation of the cnidarian
oxidative stress response by ultraviolet radiation, polycyclic aromatic hydrocarbons and
crude oil. *Journal of Experimental Biology*, *217*(9), 1444–1453.
https://doi.org/10.1242/jeb.093690

Tisthammer, K. H., Forsman, Z. H., Sindorf, V. L., Massey, T. L., Bielecki, C. R., & Toonenb,
R. J. (2016). The complete mitochondrial genome of the lobe coral Porites lobata
(Anthozoa: Scleractinia) sequenced using ezRAD. *Mitochondrial DNA Part B: Resources*,
1(1), 247–249. https://doi.org/10.1080/23802359.2016.1157770

Veal, E. A., Mark Toone, W., Jones, N., & Morgan, B. A. (2002). Distinct roles for glutathione
S-transferases in the oxidative stress response in Schizosaccharomyces pombe. *Journal of*
Biological Chemistry, *277*(38), 35523–35531. https://doi.org/10.1074/jbc.M111548200

Ward, S. (1995). The effect of damage on the growth, reproduction and storage of lipids in the
scleractinian coral Pocillopora damicornis (Linnaeus). *Journal of Experimental Marine*
Biology and Ecology, *187*(2), 193–206. https://doi.org/10.1016/0022-0981(94)00180-L

Zakai, D., Levy, O., & Chadwick-Furman, N. E. (2000). Experimental fragmentation reduces
sexual reproductive output by the reef-building coral Pocillopora damicornis. *Coral Reefs*,
19(2), 185–188. https://doi.org/10.1007/s003380000091