1   **Cellular and Circuit Organization of the Locus Coeruleus of Adult Mice**

3   Andrew McKinney[1,2,3], Ming Hu[2,3], Amber Hoskins[4], Arian Mohammadyar[4], Nabeeha Naeem[4], Junzhan

4   Jing[2,3], Saumil S. Patel[2], Bhavin R. Sheth[5,6]*, Xiaolong Jiang[1,2,3,7]*

5
[1]Neuroscience Graduate Program, Baylor College of Medicine, Houston, TX, USA

[2]Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA

[3]Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital, Houston, TX, USA

[4]University of Houston, Houston, TX, USA

[5]Department of Electrical & Computer Engineering, University of Houston, Houston, TX, USA

[6]Center for NeuroEngineering and Cognitive Science, University of Houston, Houston, TX, USA

[7]Department of Ophthalmology, Baylor College of Medicine, Houston, TX, USA

*Corresponding authors

Abstract

The locus coeruleus (LC) houses the vast majority of noradrenergic neurons in the brain and regulates

many fundamental functions including fight and flight response, attention control, and sleep/wake

cycles. While efferent projections of the LC have been extensively investigated, little is known about its

local circuit organization. Here, we performed large-scale multi-patch recordings of noradrenergic

neurons in adult mouse LC to profile their morpho-electric properties while simultaneously examining

their interactions. LC noradrenergic neurons are diverse and could be classified into two major morpho-

electric types. While fast excitatory synaptic transmission among LC noradrenergic neurons was not

observed in our preparation, these mature LC neurons connected via gap junction at a rate similar to

their early developmental stage and comparable to other brain regions. Most electrical connections

form between dendrites and are restricted to narrowly spaced pairs or small clusters of neurons of the

same type. In addition, more than two electrically coupled cell pairs were often identified across a

cohort of neurons from individual multi-cell recording sets that followed a chain-like organizational

pattern. The assembly of LC noradrenergic neurons thus follows a spatial and cell type-specific wiring

principle that may be imposed by a unique chain-like rule.

Introduction

The locus coeruleus (LC) is a small pontine nucleus that houses the vast majority of norepinephrine

(NE)-producing neurons in the brain and sends widespread noradrenergic projections to all major

divisions of the central nervous system (CNS) (Schwarz & Luo, 2015). By releasing NE across the

CNS, the LC participates in many cognitive processes and behaviors crucial for survival, including fight

and flight response, attentional control, and sleep/wake cycles, while its dysregulation is implicated in

various neuropsychiatric conditions including depression, anxiety, autism spectrum disorder, and post-

traumatic stress disorder (Bast, Poustka, & Freitag, 2018; Fortress et al., 2015; Southwick et al., 1999;

Weinshenker, 2018). Given its small size, the LC has been long considered as a homogeneous cluster

of NE-producing neurons (LC/NE neurons) exerting global, uniform influence over all CNS divisions

(Sara & Bouret, 2012; Schwarz & Luo, 2015). However, increasing evidence supports LC/NE neurons

are heterogeneous in terms of the anatomical projections, morpho-electric profiles, and specific

behavioral functions they involve (Bari et al., 2020; Hirschberg, Li, Randall, Kremer, & Pickering, 2017;

Li et al., 2016). Particularly, recent tracing studies coupled with behavioral paradigms suggest the LC is

composed of multiple specialized noradrenergic modules. Each module comprises non-overlapping

LC/NE neurons with specific electrophysiological properties and projection targets, thus participating in

distinct functions of the circuits within its efferent domain (Chandler, Gao, & Waterhouse, 2014; Li et al.,

2016; N. K. B. Totah, Logothetis, & Eschenko, 2019; Uematsu et al., 2017). Interestingly, LC/NE

neurons within each module are also able to switch from a discrete, patterned coding mode to a global

broadcast mode to cope with the more demanding situation (Uematsu et al., 2017). Such functional

modularity in the LC and their context-dependent adaptivity suggest LC neurons are functionally

heterogeneous and may interact with each other in a complex and subtle way within the local circuit to

coordinate the division of labor. To understand how each LC module is functionally specialized and how

distinct LC modules interact to achieve specific functions, it is essential to dissect the local circuit

organization of the LC in terms of its cellular diversity and wiring logic.

LC/NE neurons belong to monoaminergic systems, and their axon terminals are believed to be

"asynaptic" free nerve endings from which NE is released to modulate the activity of nearby fast-acting

synapses releasing glutamate or GABA in the form of "volume transmission". Increasing evidence

suggests monoamine neurons, including LC/NE neurons, could co-release glutamate to engage in fast

neurotransmission as well (S. I. Fung et al., 1994; Liu, Fung, Reddy, & Barnes, 1995; Yang et al.,

2021). It appears that NE can be released locally in the LC to activate α2 adrenoceptor (α2AR) on the

same or adjacent LC/NE neurons to initiate a negative feedback mechanism regulating their excitability

(Aghajanian & VanderMaelen, 1982; Egan, Henderson, North, & Williams, 1983; Huang et al., 2007;

Williams, Henderson, & North, 1985), but it remains to be illustrated if these neurons can interact with

each other via fast neurotransmission. In addition, it is long believed that LC/NE neurons form

extensive gap junctions to synchronize their activity as a global broadcast. However, gap junctions

among LC/NE neurons appear to be prevalent only in neonates and decline significantly with age

(Christie & Jelinek, 1993; Christie, Williams, & North, 1989; Travagli, Dunwiddie, & Williams, 1995). It

remains to be determined with direct demonstration if electrical coupling persists within mature LC/NE

neurons to mediate their synchronization (Ishimatsu & Williams, 1996; Patrone, Bicego, Hartzler,

Putnam, & Gargaglioni, 2014; Rash et al., 2007; Travagli et al., 1995).

To answer all these questions, here we adopted several strategies for studying cortical circuits,

including adult tissue slicing method and simultaneous multi-patch recording (Jiang et al., 2015; Ting,

Daigle, Chen, & Feng, 2014), to the brainstem to interrogate LC local circuit in adult mice. By profiling

the morpho-electric properties of each LC/NE neuron and examining their connectivity at an

unprecedented scale and level of detail (>700 neurons), we have uncovered and defined two major

morpho-electric noradrenergic cell types in the LC and deciphered the major mechanism governing

their interactions at single-cell resolution. Our results indicate that at least in our horizontal slice

preparation, no glutamatergic excitatory transmission was detected among LC neurons. Instead, we

found electrical coupling is the major synaptic mechanism by which mature LC neurons communicate

with each other, and their assembly follows a spatial and cell type-specific wiring principle that may be

imposed by a unique chain-like rule.

Results

Electrophysiological and morphological heterogeneity of LC/NE neurons

Each brainstem slice prepared from adult mice with an NMDG-based slicing protocol contained

numerous healthy LC neurons (Jiang et al., 2015; Kuo et al., 2020; Ting et al., 2014), allowing for

simultaneous patch-

clamp recordings of up

to eight mature LC

neurons (Fig. 1A). To

ensure only

noradrenergic neurons

(henceforth LC/NE

neurons) were studied

(i.e. GABAergic neurons

in LC were

excluded)(Breton-

Provencher & Sur,

2019), only

fluorescence-positive

neurons were targeted

for recordings in each

slice prepared from

DBH-Cre: Ai9 mice

(Fig.1A). To put their

electrophysiology in a

broader context of slice

electrophysiology, we

also recorded pyramidal

Fig. 1. **Electrophysiological properties of LC/NE neurons**. (**A**) Horizontal slices prepared from DBH-cre/ai9 mice for multi-cell patch-clamp recordings of LC/NE neurons. (**B**) Spontaneous firing of LC-NE neurons at rest and their more depolarized resting membrane potential (RMP) compared to hippocampal CA1 pyramidal cells (PC) and parvalbumin-expressing interneurons (PV). The representative trace for each cell type at rest is shown on the left. Right: average RMP (top, n=20 for CA1-PC and CA1-PV, n= 289 for LC/NE, CA1-PC vs LC/NE, $p<0.0001$; CA1-PV vs LC/NE, $p<0.0001$) and average firing frequency (bottom, n=20 for CA1-PC and for CA1-PV; n= 30 for LC/NE; CA1-PC vs LC/NE, $p<0.001$; CA1-PV vs LC/NE, $p<0.001$) of LC/NE neurons compared to hippocampal CA1-PC and CA-PV neurons. (**C**) The membrane potential response to the step stimulation protocol (10pA incremental) of LC/NE neurons compared to CA1-PC and CA1-PV neurons. (**D**) Individual action potential (AP) of three types of neurons (top) and tSNE of their intrinsic electrophysiology properties (bottom).

cells (CA1-PCs) and parvalbumin-expressing interneurons (CA1-PVs) from the hippocampal CA1, two

cell types commonly recorded and studied in brain slices, using the same conditions. LC/NE neurons at

rest had a highly depolarized resting membrane potential (RMP, -43.0 ± 0.1 mV, n = 289), and all

neurons were spontaneously active at rest (1.6 ± 0.4 Hz, n = 30; Fig.1B), similar to previous reports on

LC neurons (Andrzejewski, Muckenhoff, Scheid, & Ballantyne, 2001; Ballantyne, Andrzejewski,

Muckenhoff, & Scheid, 2004; Kuo et al., 2020), but different from hippocampal neurons (Fig. 1B). In

response to negative current injection, LC/NE neurons were charged much more slowly with a higher

input resistance compared to both types of hippocampal neurons (Fig.1C). In response to prolonged

suprathreshold positive current injection, the majority of LC/NE neurons exhibited a characteristic

pause after their first action potentials (APs) followed by a series of APs, which departed significantly

from stereotypical firing patterns of hippocampal neurons (Fig. 1C). LC/NE neurons were also unique in

their single APs, which were much wider with a lower amplitude, followed by a deep and prolonged

afterhyperpolarization compared to hippocampal neurons (Fig. 1C, D). To illustrate strikingly distinct

properties of LC/NE neurons, we extracted 13 electrophysiological features from three groups of

neurons (see Methods) and plotted them in a t-SNE-map. This t-SNE-based clustering approach clearly

separated LC neurons from the two types of hippocampal neurons (Fig.1D), highlighting the unique

electrophysiological properties of LC neurons.

We recovered the morphology of each recorded LC neuron to characterize their morphological

features (Fig. 2A, 2C, Fig.2S1). Most LC neurons were large cells with smooth dendrites (aspiny),

similar to CA1-PVs in the hippocampus, but different from CA1-PCs with spiny dendrites (Fig.2S1). In

addition, their dendritic trees were more tortuous with less branch order than hippocampal neurons

(Fig.2S1). LC/NE neurons had diverse somatodendritic shapes, based on which we could group them

into two major types: fusiform cells (FF) and multipolar cells (MP) following previous nomenclature

(Swanson, 1976). FFs had an elongated, usually spindle-shaped large soma with two, or rarely three,

primary dendrites originating from opposite poles of the soma (Fig. 2C, 2D, 2F). The second type of

neurons, MP, had a triangular or nearly round soma with more than three primary dendrites that could

originate from any side of their soma (Fig. 2A, 2B, 2F). The dendrites of MPs also branched more

profusely around the soma compared to that of FF (Fig.2A, 2B, 2G; Fig.2S2). To support LC neuron

classification, we reconstructed a subset of LC/NE neurons and quantified their soma bi-polarity (i.e.

the ratio of the major to the minor axis of a fitted ellipse of the soma; the larger the ratio, the more bi-

polarity the soma exhibits) and

the bi-polarity index of dendritic

arborization (the proportion of

the dendrites originating around

the major axis of the fitted

ellipse; the larger the proportion,

the more bi-polarity the dendritic

arborizations exhibits) (Fig. 2E),

and then plotted these two

parameters against each other.

Two types of neurons were well

separated in this plot (Fig. 2E).

We also trained a non-linear

neural network-based classifier

to distinguish cell types and the

classifier could separate two

types with a performance

accuracy of 100% (Fig.2S3C). Of

note, the performance using

soma shape alone was almost

as good as the performance

using two parameters (Fig.2S3A-

B), suggesting soma shape is a

reliable criterion to distinguish

two types of neurons. These

Fig 2.Two morphological cell types of LC/NE neurons. (**A**) A micrograph shows seven morphologically-recovered LC/NE neurons, most of which are multipolar cells (MP). (**B**) Morphological reconstruction of six MPs. The dendrite and soma in gray and the axon in red. The axon is shown only partially, and the arrows indicate the trajectory of the major axonal branch. The triangles indicate the origin of the axon. (**C**) A micrograph shows seven morphologically-recovered LC/NE neurons, most of which are fusiform cells (FF). (**D**) Morphological reconstruction of six FFs. The dendrite and soma in dark blue, and the axon in orange. The axon is shown only partially, and the arrows indicate the trajectory of the major axonal branch. The triangles indicate the origin of the axon. (**E**) Plotting the bi-polarity index of the soma (x-axis) against the bi-polarity index of dendritic arborization (y-axis; n = 31 for FF and n=39 for MP). (**F**) The dendritic length as a function of dendritic orientation of FF and MP. Each radius represents the dendritic length from the soma center at a certain degree. Each morphology is rotated to ensure its major soma axis aligns with the x-axis (90⁰-270⁰) in the 2D space. Zero in the plot indicates the center of the soma (n = 31 for FF, n = 39 for MP). (**G**) Left: Sholl analysis of dendritic arbors of MP and FF (counting the number of dendritic intersections that occur at fixed distances from the soma in concentric circles). FF vs. MP: $F(1) = 132.9$; $p < 0.0001$ with two-way ANOVA, n = 32 for FF and n = 36 for MP. Right: Sholl analysis of axonal arbors of MP and FF. FF vs MP: $F(1) = 87.3$; $p < 0.0001$, n = 25 for FF and n = 31 for MP. (**H**) The membrane responses to increasing current step injections in FF and MP (FF: n = 39; MP, n = 37; $F(1) = 29.5$; *$p < 0.001$ with two-way ANOVA). (**I**) Electrophysiological differences between cell types. Top: two representative firing traces in response to suprathreshold depolarizing current injections into MP and FF (left), and individual AP from MP and FF (right). Bottom: Average AP amplitude (left; FF, n = 19; MP, n = 41; **$p < 0.01$), average AP half-width (middle; FF, n = 19; MP, n = 41; **$p < 0.01$), and Upstroke/downstroke ratio of AP (right; FF, n = 19; MP, n = 41; *$p < 0.05$). Also see figure 2 supplement 1-4 (Fig.2S1-3, Fig.2S5-Table).

165 analyses supported our manual classification based on the somatodendritic shape.

In addition to somatodendritic differences, MP and FF differed in their axonal arborization as well. The majority of LC/NE neurons had a visible thick axon that originated from either the soma or dendrite (see below), and then arborized away from the fourth ventricle and were inevitably severed due to the slicing procedure (as evidenced by a characteristic retraction ball) (Blythe, Wokosin, Atherton, & Bevan, 2009). FFs generally had the axon originating from one of the primary dendrites (~95%, 37 out of 39 reconstructed FF neurons, Fig. 2D, Fig. 2S2B), suggesting most FFs have privileged dendrites (i.e. synapses made on these axon-bearing dendrites are in an electrotonically privileged position in evoking action potentials)(Hausser, Stuart, Racca, & Sakmann, 1995), while most MPs had their axon originating from the soma (~80%, 37 out of 46 reconstructed MPs; Fig. 2B, Fig. 2S2A). In addition, MPs had less local axonal arborization as compared to FFs (Fig. 2G, Fig. 2S2). The axon in many MPs had the first branching point located > 400 μm away from their origin, while the axon in FFs branched much earlier and formed denser local axonal arborization than MPs (Fig. 2G, Fig. 2S2).

FFs and MPs had different population sizes (MP: ~62%, 416 out of 666 recovered LC/NE neurons; FF: ~38%, 250 out of 666 LC/NE neurons). Given that FFs constitute ~ 38% of total LC/NE neurons, the proportion of FFs in each multi-cell recording set (the cohort of >3 LCs neurons from each slice) should fall most frequently within the range of 30-40% if FFs are evenly distributed across the entire LC. However, this was not the case in the histogram of FF proportion (Fig. 2S4) across all recording sets. Instead, the FF proportions were heavily biased toward either the range of 0-20% or of 50-100%, suggesting that FFs were not evenly distributed across the entire LC, but preferentially localized in certain anatomical positions. Indeed, we often recorded a cohort of neurons in which most LC/NE neurons were FFs (Fig.2C). The same was true for the proportions of MP (Fig.2A, Fig. 2S4). This sampling bias might reflect the anatomical preference of different types of LC/NE neurons as reported in rat LC (Loughlin, Foote, & Grzanna, 1986; Swanson, 1976).

To examine if MPs and FFs had different intrinsic membrane properties and firing patterns, we extracted and compared eight electrophysiological parameters between the two types (Fig.2S5-Table). FFs had a narrower spike with a larger amplitude than MPs (Fig. 2I, Fig.2S5-Table). The spikes from the two cell types also had different temporal dynamics (measured at the first spike evoked by our step

193   protocol, Fig. 2I, Fig.2S5-Table). In addition, FFs could be evoked more spikes with large

194   suprathreshold depolarizing current injections than MPs (Fig. 2H).

195   **Synaptic transmission between LC/NE neurons**

Fig. 3. **Chemical and electrical transmission between LC/NE neurons**. (**A**) Simultaneous patch-clamp recording of eight LC neurons with their morphology *post hoc* recovered. Cell 7 and Cell 2 are electrically coupled. (**B**) The membrane responses of eight LC neurons to a hyperpolarizing current injection (bottom trace) and suprathreshold depolarizing current injection(upper trace) and pairwise testing of synaptic connection (or any other type of interactions) between eight LC neurons. Red traces indicate action potential (AP) evoked in each presynaptic cell and the average traces of the potential postsynaptic responses are shown in black. No visible membrane hyperpolarization or depolarization following each AP was detected in any cell pair. Vertical scale bar: 2nA for current injection, 10mV for the traces in postsynaptic cells, 50 mV for AP evoked in presynaptic cells. The traces for electrical coupling not shown. (**C**) Inter-soma distance of all tested cell pairs. The majority of cell pairs have an inter-soma distance < 100 µm. (**D**) The proportion of chemical connections (CC), electrical connections (EC), and non-connection (NC) out of all tested connection pairs. No chemical connection was detected with single AP or a train of APs (5 APs at 20 Hz) across all tested pairs, but electrical connections were sparsely identified in a small subset of cell pairs.

196   LC/NE neurons may co-release glutamate to engage in fast excitatory synaptic transmissions (S. I.

197   Fung et al., 1994; Fung, Reddy, Liu, Wang, & Barnes, 1994; Yang et al., 2021). To directly test this

connection scenario, we examined synaptic connections between any LC/NE cell pair once

simultaneous whole-cell recordings of LC neurons were established (Fig. 3A, B). Single action

potentials (APs) or a train of APs (5 APs at 20Hz) were evoked in presynaptic neurons, while

membrane potentials of other simultaneously-recorded LC neurons were monitored. An excitatory

synaptic connection is identified if the excitatory postsynaptic potentials (EPSPs) in potential

postsynaptic neurons are time-locked to presynaptic APs (Jiang et al., 2015; Scala et al., 2019). We

tested more than 1500 LC neuron pairs, most of which have an inter-soma distance < 100 µm (Fig.

3C), but no single excitatory synaptic connection was detected (Fig. 3D) despite prominent

spontaneous EPSPs present in each potential postsynaptic neuron. In addition, NE released from

LC/NE neurons may activate α2AR on adjacent LC/NE neurons or themselves to evoke

hyperpolarization. However, no visible hyperpolarization in potential postsynaptic cells following single

presynaptic APs or a train of APs was detected in these tested pairs as well, suggesting NE released

from single LC neurons was not sufficient enough to activate the α2ARs.

With our multi-patch recordings, we also examined electrical coupling between any cell pair. Out

of 1503 cell pairs tested, we identified sparse electrical connections among LC/NE neurons from adult

animals (Fig. 3D). The presence of electrical coupling was tested by recording membrane responses in

simultaneously recorded LC/NE neurons following the injection of prolonged depolarizing or

hyperpolarizing current pulses (600 ms) in one of the cells. An example of a coupled pair is illustrated

in Figure 4A-C, in which a current pulse in presynaptic Cell 1 evoked a membrane response of the

same sign in postsynaptic Cell 4, albeit of much lower amplitude. The same occurred when a current

pulse was injected into Cell 4. Other tested pairs did not show evidence of electrical coupling in either

direction (Fig. 4C).

The steady-state coupling coefficient, defined as the voltage change at the postsynaptic cell

divided by the voltage change at the presynaptic cell measured at the last 10-20 ms period of the

current step (see Methods), was calculated for each recorded pair. A pair of LC/NE neurons was

considered to be electrically coupled when the steady-state coupling coefficient was ≥ 0.005, a criterion

determined by the noise of the recording conditions (Curti, Hoge, Nagy, & Pereda, 2012). According to

 this criterion, 65 of the 1503 tested pairs across the inter-soma distance range of 16-531 µm were

Fig. 4. Electrical coupling between LC/NE neurons. (**A**) A micrograph shows eight morphologically-recovered LC/NE neurons. (**B**) The relative position of eight neurons (with an assigned number) and their connections. (**C**) Pairwise testing between eight cells as shown in **A** reveals electrical coupling between Cell 1 and Cell 4. No sign of electrical coupling between the rest of cell pairs. (**D**) Estimates of the coupling coefficient in both directions (R1 and R2) are plotted against each other, indicating that most contacts lack significant rectification. The data are fitted with a straight-line function ($R^2 = 0.75$, $p < 0.001$). (**E**) Electrical connectivity rate as a function of the inter-soma distance (ISD). (**F**) Electrical connectivity rate at different ages (PND: postnatal day). N.S.: no significant difference. (**G**) Electrical connectivity rate between neurons of same type or different types across all cell pairs. ***$p < 0.0001$. (**H**) Electrical connectivity rate as a function of inter-soma distances (ISD) imposed by cell types. (**I**) Coupling coefficients in two homotypic types of electrical coupling (within-cell type) and one heterotypic type of electrical coupling (among the different cell types). *$p < 0.05$.

electrically coupled (4.3%; Fig. 3D, Fig. 4F, also see below). For each coupled pair, the coupling

coefficients were estimated and expressed as the average of the values in both directions. The steady-

state coupling coefficient averaged 0.040 ± 0.028 (SD) (range: 0.0051– 0.13; n = 65) (Fig.4D, I). The

strength of electrical transmission in the vast majority of coupled pairs was bidirectional and symmetric

(Fig.4C, D). Estimates of the coupling coefficient in both directions for each pair showed a positive

correlation with a slope of 1.03 (Fig. 4D), not significantly different from 1 (p = 0.8; Fig. 4D), indicating

that the conductance of putative gap junctions between LC/NE neurons was largely nonrectifying.

Electrical coupling generally occurs between narrowly-spaced cell pairs. Given the wide range of

the inter-soma distance of tested pairs, a low coupling rate in our dataset was not unexpected. To

examine how electrical coupling among LC neurons was restrained by their inter-soma distance, we

measured the inter-soma distances of all tested cell pairs to derive their electrical coupling rate as a

function of the inter-soma distance. With an inter-soma distance < 40 μm, the coupling rate between LC

neurons was ~12%, indicating electrical connections were prominent among LC neurons in adult mice.

However, the coupling rate dropped quickly as the inter-soma distance increased from 40 μm (Fig.4E).

Interestingly, while the coupling rate was much lower in cell pairs with the inter-soma distance > 40 μm,

it was much more resistant to the increase of the inter-soma distance than the rate in cell pairs with the

inter-soma distance < 40 μm (Fig. 4E). The coupling rate remained relatively constant when the inter-

soma distance increased from 40 μm to 240 μm before dropping to zero, with a second small peak

around 140 μm (Fig. 4E).

Previous reports suggest electrical connections among LC/NE neurons decline significantly with

age (Christie & Jelinek, 1993; Christie et al., 1989; Travagli et al., 1995). To examine if this is the case,

we performed additional multi-cell recordings of LC neurons from mice at the age of PND 13-21 (Fig.

4F). Surprisingly, the coupling rate at this young age was very similar to the rate found in adults (4.8%;

8 connections out of 165, p > 0.05, Fig. 4F). The coupling rate was still very similar between the two

ages when the inter-soma distance was controlled (Fig. 4F).

Electrical transmission by morphological cell type in LC

Electrical coupling generally occurs between cell pairs of the same cell type in many brain regions

(Bennett & Zukin, 2004; Hidaka, Akahori, & Kurosawa, 2004). To examine if electrical coupling between

LC/NE neurons was also cell-type specific, we examined all tested cell pairs with recovered

morphology and assigned each cell to either FF or MP, and then derived the coupling rates as a

function of cell type (Fig.3-6). Of all FF–FF pairs (i.e. both cells are FF), we found 8.7% were electrically

coupled (Fig. 4G); of all MP–MP pairs (i.e. both cells are MP), 5.3% were electrically coupled (Fig. 4G).

By contrast, only 0.8% of FF–MP pairs were coupled (Fig. 4G; $p < 0.0001$ for both comparisons). While

the coupling rate of FF–FF pairs was slightly higher than that of MP–MP pairs, there was no significant

difference between these two ($p = 0.08$). Given that the coupling rate is sensitive to the inter-soma

distance, we then compared their rates within the same inter-soma distance range to rule out the

possibility that the rate differences we observed were due to the differences in the inter-soma distance.

Interestingly, with the inter-soma distance of up to 40 µm, no connection was identified in FF–MP pairs

(0 out of 55 tested pairs). In contrast, the coupling rates for FF–FF and MP–MP pairs were much higher

(11 out of 35 tested pairs, 31.4% for FF–FF, $p < 0.01$ compared with FF–MP; 7 out of 65, 10.7% for

MP–MP, $p < 0.01$ compared with FF–MP; Fig. 4H). In addition, the coupling rate for FF–FF pairs was

significantly higher than that for MP–MP pairs over this inter-soma distance range ($p = 0.01$, Fig. 4H).

With the inter-soma distance range of 40-80 µm, the coupling rates for both MP–MP and FF–FF pairs

were also significantly higher than that for FF–MP pairs ($p < 0.01$ for both comparisons, Fig. 4H), but

there was no significant difference between the MP–MP coupling rate and FF–FF coupling rate ($p = $

0.67). Electrical coupling in FF–MP pairs most occurred in those cell pairs with the inter-soma distance

> 80 µm, but their rate even in this inter-soma distance range was lower than the rates for MP–MP and

FF–FF pairs (FF–FF vs FF–MP: 6.3% vs 1.4%, $p < 0.01$; MP–MP vs FF–MP: 3.9% vs 1.4%, $p = 0.05$,

Fig. 4H). These results indicate that electrical connections in LC neurons generally form among cell

pairs of the same cell type (Fig. 3-6). Within the same cell type, a FF is more likely to connect with

another FF than a MP with another MP, particularly if the neurons are located nearby. In addition,

electrical coupling in FF–FF pairs had higher connection strength (i.e. higher coupling coefficients) than

electrical coupling in MP–MP pairs (Fig. 4I).

Putative electrical synapses in LC/NE neurons

We reason electrical synapses formed between FFs (FF–FF) and those formed between MPs (MP–

MP) may locate in different somatodendritic domains (soma vs. dendrite), which may contribute to their

282 coupling coefficient difference (Fig. 4I). To test this hypothesis, we took advantage of high-quality

283 morphology recovered from each LC neuron (Figs. 2-6) to examine putative electrical synapses

Fig.5. Putative electrical synapses in LC/NE neurons. (**A**) Left: a microphotograph depicting six morphologically-recovered LC/NE neurons (simultaneously recorded), among which Cell a and Cell b were identified as an electrically-coupled pair. Multiple putative dendrodendritic electrical synapses can be identified after their morphology was reconstructed (as labeled 1 from 8). Right: a high-magnification view of each putative electrical synapse as shown on the left. (**B**) Top: a microphotograph depicting seven morphologically-recovered LC/NE neurons, among which Cell a and Cell b were identified as a electrically-coupled pair. A putative somatodendritic electrical synapse was identified after their morphology was reconstructed (as labeled as 1). Bottom: a high-magnification view of the putative electrical synapse as shown on the top. (**C**) Top: a microphotograph depicting six morphologically-recovered LC/NE neurons, among which Cell a and Cell b were identified as a electrically-coupled pair. A putative dendrodendritic electrical synapse was identified (as labeled as 1). Bottom: a high-magnification view of this putative electrical synapse as shown on the top. (**D**) Dendritic distances from the putative electrical synapse to the soma of Neuron 1 (Soma 1) are plotted against the distance to the soma of Neuron 2 (Soma 2) for each putative electrical synapse formed between two LC neurons. Correlation coefficient r = 0.79, $p < 0.01$. The slope of the straight line is 0.66. (**E**) The putative synapse number was plotted against the coupling coefficient for each pair. r = 0.86, $p < 0.05$. (**F**) The distances from each putative synapse to two somata were averaged and then plotted against the coupling coefficient of each pair. The coupling coefficient was normalized if a electrically-coupled pair had multiple putative synapses. r= -0.19. (**G**) Comparing the dendritic distance of putative synapses to the soma and the putative synapse number between two homotypic electrical coupling types (FF-FF vs. MP-MP*: p =* 0.19 in the dendritic distance; $p = 0.43$ in the putative synapse number).

between any coupled pairs. We traced the entire somatodendritic domain of a coupled pair under a

100X oil immersion objective lens and scrutinized those areas where two neurons have overlapping

somatodendritic structures. In those coupled cell pairs, a dendritic branch from one cell merged with a

dendritic segment from another cell at one or multiple crossing points (Fig. 5A-C), whereas we rarely

found these close contacts in uncoupled cell pairs. We thus believed that these contacts were putative

electrical synapses among coupled pairs.

For many coupled pairs, only one putative synapse could be identified (Fig. 5C). More than two

putative synaptic contacts could be identified in some coupled pairs as well (Fig. 5A) – these were more

frequent in FF–FF pairs (Fig. 5G). This may be one reason for a higher connection strength in FF–FF

coupled pairs as compared to MP–MP coupled pairs (Fig. 4I). Indeed, putative synaptic numbers and

coupling coefficients showed a positive correlation (r = 0.86, Fig. 5E). The vast majority of putative

synapses were found to form between dendrites: this could be a single crossing point between two

dendrites from a coupled pair (Fig. 5C), or multiple merging sites between an entire dendrite from one

cell and the entire dendrite from another cell (in parallel with each other along their course to the

terminals; Fig. 5A). Only two coupled pairs (one FF–FF and one MP–MP) had putative synapses

formed between the dendrite and the soma (Fig. 5B). No putative synapses forming between two

somata were identified in our data. Put together, our data suggest there was no preferential synaptic

location for either FF–FF coupled pairs or MP-MP coupled pairs.

The dendritic branches where the putative synapse was localized were generally first or second-

order branches and the putative synapses were most frequently located in the middle of the entire

dendritic tree (average dendritic distance to soma: 155 µm, Fig. 5G). When one cell pair had multiple

contacts, these contacts were generally confined to the same major dendrite from each cell; some of

them were even confined to the same dendritic branch of the same order of each cell. For each putative

synapse, the dendric distance to one soma was linearly related to the distance to another soma with an

r of 0.86, indicating that the putative synapses sit a similar dendritic distance away from the two somata

of the coupled pair (Fig. 5D). The average dendritic distance of putative synapses from FF–FF pairs

was 168 µm, tending to be slightly greater than the distance for MP–MP pairs (140.0 µm, Fig. 5G). In

addition, the dendritic distance of the putative synapse appeared to negatively correlate with its

coupling coefficient (Fig. 5F).

Connectivity patterns of electrical coupling beyond two LC neurons

Multi-cell recordings allowed for examining electrical coupling beyond two cells. In many individual

recording sets, we could identify multiple electrical connections among a cohort of cells. An example of

such a case is illustrated in Figure 6A-C, in which a current pulse in presynaptic Cell 6 evoked a

membrane response of the same sign in postsynaptic Cell 1 and 8. A current pulse in presynaptic Cell

1 or 8 evoked a membrane response of the same sign in postsynaptic Cell 6, but not 8 or 1

respectively. This indicates Cell 1 and Cell 8 both electrically connect with Cell 6, but Cell 1 and Cell 8

do not directly connect with each other. This result also indicates that electrical signals from one cell

hardly pass to other cells via second-order coupling if they are not electrically coupled in the first order.

Other tested pairs did not show evidence of electrical coupling in either direction (Fig. 6C).

Out of our all multiple-cell recording sets (> 3 cells), 15 individual recording sets have more than

two electrical connections (or electrically-coupled pairs) identified. In these recording sets, one coupled

pair often shared the same coupling partner with another coupled pair as illustrated in Figure 6A-C (i.e.

A↔B↔C; each letter represents a cell. Fig. 6D). This is surprising since two coupled pairs should be

less likely to share the same partner if each cell is coupled with other cells in the network at the same

rate. However, two coupled pairs without a sharing partner (i.e. A↔B, C↔D) were rare in our recording

sets (Fig. 6D). These data suggest each cell is not equally coupled with other cells in the network;

instead, electrically-coupled cells tend to cluster together and form a chain-like pattern (i.e. A↔B↔C).

To test this hypothesis, we started by assessing how well a randomly connected network describes our

dataset (Erdos & Rényi, 1984). In this model, the existence of an electrical connection between any two

neurons was independently chosen with a uniform probability p (i.e. the overall connectivity rate across

all of our recordings: 0.0438; Fig. 3D). We first tested this random connectivity model in predicting the

patterns of two electrical connections in our recording sets (Model 1, see Methods). Specifically, based

on random pairwise connectivity, we predicted the numbers of a chain-like pattern of two connections

(A↔B↔C) versus a non-chain-like pattern of two connections (A↔B, C↔D) using our recording sets of

3–8 neurons. We found that the observed number of the chain-like connection pattern (n = 16, Fig. 6D)

was higher than the numbers predicted by the random connectivity model (n = 12.9, 95% confidence

interval (C.I) = [6, 21]), whereas the observed number of the non-chain-like patterns (n = 5, Fig. 6D)

was significantly less than the numbers predicted by the model (n = 11.6, 95% C.I = [6, 19], *p* = 0.01).

Fig.6. **Multiple electrically-coupled pairs identified in a cohort of LC/NE neurons**. (**A**) A micrograph shows six morphologically-recovered LC/NE neurons. (**B**) The relative position of six neurons (with assigned number) depicted as above and their connections. (**C**) Pairwise testing between six cells as shown in **A** reveals electrical coupling between Cell 1 and Cell 6, and between Cell 6 and Cell 8. No sign of electrical coupling between the rest of cell pairs. (**D**) The counts of multiple electrically-coupled pairs identified from individual multi-cell recording sets with all possible arranging patterns (top). (**E**) The cell-type identity of each cell in those inter-connected coupled pairs presented in D, and the frequency for each possible combination.

We calculated the ratio of the chain-like pattern number to the non-chain-like pattern number to reflect

the clustering degree of two electrical connections, which we henceforth called the cluster index (Fig.

7A). The cluster index in our experimental dataset was 3.2, which was significantly higher than the

number predicted by the model (1.1, *p* = 0.001). We then tested if differences in the organizing pattern

of two connections (observed vs. predicted) can be better explained by the cell type-specific

connectivity rule as we deciphered (see Fig. 4G) by incorporating cell type information into the model.

That is to say, pairwise connection probabilities for FF–FF, FF–MP, and MP–MP pairs (see Fig. 4G)

were used to obtain a new model prediction. With this revised model (Model 2, See Methods), the

observed numbers on average of the chain-like pattern again remained higher than the predicted

numbers (n = 12.6, 95% C.I. = [7, 18]), and the observed number of the non-chain-like pattern was still

significantly lower than the numbers (n = 11.2, 95% C.I. = [6,16], $p < 0.001$) predicted by the model

(Fig. 7B). The cluster index predicted by Model 2 was very similar to that predicted by Model 1, and

significantly lower than the observed index (Fig. 7A), indicating that the cell type-specific rule can not

explain our dataset. Given that neurons of the same type tended to be near each other (Fig. 2S4) and

electrical connections are highly sensitive to the inter-soma distance (see Fig. 4H), we further revised

our model (Model 3, See Methods) by zeroing out connections between cell pairs that were greater

than 80 µm, and by using the FF–FF and MP–MP connection rate for cell pairs within 40–80 µm apart

and FF–MP connection rate for cell pairs within 80 µm apart. The results did not fundamentally change:

the observed number of the chain-like pattern was not significantly different from the predicted value (n

= 19.4), but the observed number of the non-chain-like pattern (n = 4) was still significantly lower than

that (n = 15.9) predicted by the model. The cluster index predicted by Model 3 was still significantly

lower than the observed index (Fig. 7A), indicating additional spatial connectivity rules could not explain

our dataset as well. In summary, the clustering of two connections in our dataset (a chain-like pattern)

can not be explained by the random model and spatial and cell type-specific rules as shown above,

suggesting an additional, possible chain-like rule dictating the organization of electrical connections in

the LC network (note that all three models shared the assumption that the probability for two neurons to

form an electrical connection is independent of all other pairwise connections, an assumption that the

new rule will perhaps have to violate). Indeed, in a proof-of-concept new model (Model 4, appendix 1)

where the said assumption in Model 1-3 was violated (i.e. a cell's connectivity being scaled up or down

contingent on whether or not it has an existing electrical connection), the observed patterns of two

connections can be well predicted by the model (Appendix 1), supporting that electrical connection

among LC neurons are not random, but incline to cluster together in LC (in addition to following cell

type and spatial rules). Such an underlying new rule could impose on both cell types, and thus the most

frequent patterns of two electrical connections in our dataset are FF–FF–FF and MP–MP–MP, while

other combination patterns are either rare or not observed (Fig. 6E).

Frequent observation of the clustered, chain-like organization of two connections suggests that

an electrically-coupled neuronal network may be wired as a chain-like network structure (Fig. 7C).

Fig.7. Electrically-coupled homotypic LC subnetworks. (**A**) In two electrical connections between LC neurons, comparing observed ratio (chain/non-chain pattern) vs the ratios predicted by Model 1 (random connectivity), Model 2 (cell type-specific) and Model 3 (cell type and spatial-specific) suggests a chain-like rule dictating electrical connections. (**B**) In three electrical connections between LC neurons, observed ratio (chain/non-chain pattern) vs the ratios predicted by Model 1(random connectivity), Model 2 (cell type-specific) and model 3 suggests a chain-like rule dictating electrical connections. Error bars are standard deviations estimated by bootstrap method. ***$p \leq 0.001$, *$p < 0.05$. (C) A sagittal view of the LC with predominant cell types for each subdivision (adopted from Laughlin et al. 1986) and the proposed electrically-coupled homotypic subnetworks in the LC. Spindle-shaped: fusiform cell; round-shaped: multipolar cells. Each electrical connection form between cell pairs of same type and each connected pair is connected with each other like a chain within each LC subregion.

Indeed, when we examined the organizing pattern of three electrical connections in those individual

recording sets, the most frequent was still the chain-like pattern (A↔B↔C↔D, Fig. 6D), while other

possible organizing patterns including circular-like and hub cell-like patterns were rarely observed in our

dataset (Fig. 6D). We then extended our models to predict how three electrical connections are

organized. Our random connectivity model (Model 1) predicted 2.0 chain-like patterns of triple

connections on average (95% C.I. = [0,6]) across all our recordings which was the same as the actual

observed number, while it predicted 3.0 non-chain-like triple connections on average (95% C.I. = [0, 7]),

which was greater than the observed number (1). We again used the ratio of chain-like to non-chain-

like patterns to measure the chain-like tendency in our dataset and compare it with model predictions.

The observed ratio in our dataset was 2, significantly higher than the predicted ratio (0.68 ± 0.25, $p =$

0.001, Fig. 7B). Our Model 2 (see above) also predicted 2.2 chain-like pattern of triple connections on

average (95% C.I. = [0,6]); however, it predicted 4.9 non-chain-like triple connections (95% C.I. = [2,8])

which was significantly greater than the observed number (n = 1, $p =$ 0.01). Again, the observed ratio

was significantly higher than the ratio predicted by Model 2 (0.52 ± 0.47, $p <$ 0.05, Fig.7B). Similar

results were obtained using our Model 3 (see above, incorporating inter-neuronal distance), which

predicted 6.8 of the chain-like patterns of triple connections on average and 9.3 non-chain-like triple

connections on average. Once again, the observed ratio was significantly higher than the ratio

predicted by the third and final model (0.76 ± 0.24, $p <$ 0.001, Fig.7B). All simulations indicate that, in

addition to spatial and cell type-specific rule, an additional chain-like rule may apply for the organization

of three electrical connections.

Discussion

In this study, we overcame the technical barriers to record up to 8 noradrenergic neurons

simultaneously from adult mouse locus coeruleus (LC). Simultaneous multi-cell patch-clamp recordings

(i.e. recording more than two cells) have been developed to study cortical circuits for a while, but it

remains a challenge to apply this technique to small nuclei in the brainstem such as the LC given the

space restraint and limited cell number in each brain slice. By optimizing slice quality and adapting our

multi-patching system to small brainstem slices, we could simultaneously record LC/NE neurons in vitro

in the cohort of an average of six cells from adult mice. This technical development allowed for profiling

the cellular diversity of LC/NE neurons while simultaneously mapping their connections at a single-cell

resolution. In fact, this study may represent the first multi-cell patch recordings of a monoamine

modulatory system in the brain, allowing for examining how monoaminergic neurons interact with each

other at the single-cell level for the first time. Our results indicate noradrenergic neurons in the LC could

be classified into at least two major morpho-electric cell types, and neurons within each cell type form a

homotypic network module via gap junctions. Given that each of these cell types may have preferential

anatomical locations in LC and different projection targets, each electrically-coupled homotypic network

may coordinate/synergize their input/output as a whole to engage in distinct functions of the circuits

within its efferent domain. In addition, our large dataset has allowed us to directly test several long-

standing questions regarding local circuit interaction in the LC, including α2AR-mediated actions and

fast glutaminergic synaptic connection at the single-cell level.

Morpho-electric diversity of LC/NE neurons

Morphological heterogeneity of LC neurons has been noticed since the very early LC studies. The

anterior horn of the LC and the ventral part of the LC core appears to be dominated by cells with a

stellate multipolar soma, whereas its posterior horn and the dorsal edge of the LC core are dominated

by cells with a fusiform-shaped soma (Loughlin et al., 1986; Swanson, 1976). In addition, LC neurons

with distinct soma shapes preferentially project to different brain areas (Loughlin et al., 1986;

Waterhouse, Lin, Burne, & Woodward, 1983), warranting further exploration of the morphological

diversity of LC neurons and their functional relevance. To this end, we recovered the morphology of a

large number of LC/NE neurons via biocytin staining to characterize their morphological features,

providing a quantitative, less ambiguous classification of LC/NE neurons. We confirmed that the soma

shape of LC/NC neurons could be either fusiform, round, or pyramid (Loughlin et al., 1986; Swanson,

1976). Cells with fusiform-shaped soma have a polarized dendritic arborization (bitufted or bipolar),

while cells with round or pyramid-like soma have much less polarized dendrite arborization. These

differences allowed for classifying LC neurons into two morphological cell types – multipolar cell (MP)

and fusiform cell (FF), which was substantiated by a machine learning algorithm. In addition to having

distinct somatodendritic shapes, MP and FF cells have different local axonal arborization as well,

further supporting our classification. Finally, MPs and FFs have distinct intrinsic electrophysiological

properties. While mature LC/NE neurons as a whole exhibit the stereotypical physiological properties that clearly differentiate them from hippocampal neurons, there are subtle differences between MP and FF in their individual action potentials (APs): MPs had a longer AP with smaller amplitude than FFs. Furthermore, FFs could sustain higher frequency firing than MPs with large suprathreshold depolarizing current injections. The fact that FFs fire narrower spikes than MPs suggests FF may correspond to "narrow" units identified during extracellular recordings of LC neurons, while MP may correspond to those "wide" units (Su & Cohen, 2022; N. K. Totah, Neves, Panzeri, Logothetis, & Eschenko, 2018). However, "narrow" units are heavily biased toward the ventral part of the LC (Su & Cohen, 2022; N. K. Totah et al., 2018), while FFs as we described here may be dorsally biased (Loughlin et al., 1986). In addition, FFs do not exhibit a higher spontaneous firing rate than MPs (Fig. 2H), inconsistent with a higher firing rate found for extracellularly recorded "narrow" units (N. K. Totah et al., 2018). Furthermore, there is a two-fold difference in spike width between "narrow" and "wide "units (N. K. Totah et al., 2018), but the difference in spike duration (including the rise phase of first derivatives of AP) between FF and MP is subtle despite being significant (Li et al., 2016). Therefore, there is no consistent evidence supporting that FF cells we defined here correspond to those "narrow" units, and MPs correspond to "wild" units. Likely, each single unit type is not confined to a specific morphological type, or both single unit types may be from the same MP type reflecting within-MP functional heterogeneity. The latter may be more likely given that MPs are a predominant LC cell type and they are more heterogeneous than FF in terms of soma size and spatial distribution (Loughlin et al., 1986), as well as electrophysiology properties (including AP half-width; data not shown). Nevertheless, it is unknown how well in vitro intracellular waveforms correlate with in vivo extracellular ones and thus we cannot rule out the possibility that FF cells correspond to those "narrow" units. Establishing a real correspondence between cell types as we discovered here and single unit types is important for the field, warranting a new systematic study (Su & Cohen, 2022; N. K. Totah et al., 2018; N. K. B. Totah et al., 2019).

The two cell types have distinct morphology and electrophysiology, suggesting they may participate in distinct functions of the circuits within their efferent domains. Indeed, previous studies

indicate that fusiform-shaped cells in the posterior horn of the LC and the dorsal edge of the LC core preferentially project to the neocortex and hippocampus, while multipolar neurons in its anterior horn and/or the ventral part of the LC core preferentially project to the spinal cord, cerebellum, and hypothalamus (Loughlin et al., 1986; Waterhouse et al., 1983). While it is difficult to register each cell in our slice recordings to their anatomical location *in vivo*, we reason that FF cells as we defined here correspond to those fusiform-shaped cells in the posterior horn of the LC and/or the dorsal edge of the LC core, while MP cells may correspond to those multipolar cells in the anterior horn and/or the ventral part of the core. If this is true, these two types as we defined here might form separate LC subnetworks with distinct projection domains, each performing a non-overlapping function in the brain. In addition, FF and MP may be further classified into different subtypes based on the locations, each of which has different preferred projection targets as well and thus form distinct MP and FF subnetworks (Loughlin et al., 1986). This may be particularly the case for MP (also see below). To explore the functional relevance of these two major morphological types or the possible subtypes within each morphological type, identifying a specific genetic marker for each cell type or subtype is required. Prior studies indicate some LC neurons could express neuropeptide Y (NPY) and/or galanin. However, it appears neither NPY-expressing nor galanin-expressing LC neurons are confined to a specific morphological type (Holets, Hokfelt, Rokaeus, Terenius, & Goldstein, 1988; Melander et al., 1986), suggesting neither NPY nor galanin is a good candidate to differentiate the two LC morphological types. In this regard, relatively new approaches such as single-cell RNA-sequencing (scRNA-seq) are being applied to the LC, which have the potential to identify the specific marker genes for each LC type or subtype (Luskin et al., 2022; Scala et al., 2020; Tasic et al., 2016; Zeisel et al., 2015). With type-specific marker genes identified with scRNA-seq, one can leverage an intersectional genetic strategy employing a dual recombinase system using both Cre recombinase as well as a second recombinase, flippase (Flp), to express the reporter genes in each specific LC neuronal type (or subtype). For this purpose, triple transgenic mice could be generated by crossing the Cre/Flp or Cre/Dre double-dependent reporters crossed with *Dbh*-cre (labeling all noradrenergic neurons) and an Flp driver line (expressing Flp in the

marker gene locus) to label and manipulate each cell type (or subtype) while animals are trained to

perform behavioral tasks (Lusk et al., 2022; Madisen et al., 2015).

Chemical transmission between LC/NE neurons

The physiologic function of monoamine neurons is generally considered to be modulatory given that

most monoamine axon terminals in the CNS are "asynaptic" free nerve endings thought to mediate

"volume transmission". However, many lines of evidence suggest monoamine axon terminals, including

noradrenergic terminals, co-release glutamate. At least a subset of their axonal terminals makes the

junctional or conventional-looking synaptic contacts onto targeted neurons, possibly mediating rapid

glutamatergic synaptic transmission (S. I. Fung et al., 1994; Liu et al., 1995; Trudeau, 2004; Yang et al.,

2021). While evidence for this emerging concept of monoamine function is pretty strong (Yang et al.,

2021), a direct demonstration of the glutamatergic nature of synapses established by monoamine

neurons is still lacking. We thus took advantage of our unprecedented multi-cell recordings of

monoamine neurons to examine if there is any glutamatergic synaptic transmission between LC/NE

neurons. Given the high-throughput nature of our system, we were able to test more than 1,500 cell

pairs, including those narrowly spaced cell pairs (< 40 μm apart). However, we did not find evidence of

glutamatergic synaptic transmission between LC/NE neurons in our horizontal slice preparation. It will

be interesting to examine whether this type of connection, if it does exist among LC neurons, can be

more readily detected in coronal slices where peri-LC glutamate transmission may be somewhat more

spared than in horizontal slices. Nevertheless, we believe this type of connection, if it does exist, is very

sparse, given that only a small subset of all terminals established by monoamine neurons may actually

be specialized for the synaptic release of glutamate (Trudeau, 2004). In addition, our result does not

rule out the possibility that a subset of noradrenergic axonal terminals does make glutamatergic

synapses onto other target regions, including the spinal cord and the parabrachial nucleus neurons (S.

I. Fung et al., 1994; Liu et al., 1995; Yang et al., 2021).

Monoamine release is controlled by their autoreceptors in the presynaptic terminals. The same

receptors are often expressed in the somatodendritic domains of monoamine neurons, activation of

which increases potassium conductance as seen in their terminals (Courtney & Ford, 2014). Alpha2

adrenoceptors (α2AR) are such autoreceptors for NE and are expressed in the somata of LC/NE

neurons as well. While α2AR-mediated action on LC neurons is well established, it is unclear under

what conditions these receptors are activated and what is their natural agonist. These receptors may be

expressed in the soma to sense NE locally released from other or the same LC/NE neurons to engage

in lateral inhibition or autoinhibition, and/or epinephrine released from the medulla oblongata

(Cedarbaum & Aghajanian, 1976; Ho¨kfelt, Fuxe, Goldstein, & Johansson, 1974). It appears that NE

could be released locally from local axonal terminals or somatodendritic domains of LC neurons

following a single AP, but it is unknown if NE released from individual neurons is sufficient enough to

activate α2AR to evoke lateral inhibition on adjacent neurons. After testing more than 1500 cell pairs,

we did not find any detectable hyperpolarization following the stimulation of a single LC/NE neuron.

While other unknown reasons may contribute to the absence of lateral inhibition, a plausible

explanation is that NE released from individual LC/NE neurons is not sufficient to activate α2AR to

evoke detectable inhibition on LC/NE neurons at least in our in vitro preparation. A recent study

suggests that α2AR-mediated lateral inhibition is a frequency-dependent population-wide phenomenon,

which can be readily detected when many neurons are simultaneously activated via an optogenetic

approach (Davy et al., 2022). In addition, α2AR-mediated lateral inhibition appears to be a specific

network interaction that is prevalent among LC neurons targeting distinct brain regions (module-

specific), rather than a non-specific, spatial "surround" inhibition (Davy et al., 2022). All these could

explain why lateral inhibition could be not readily detected at the single-cell level in our dataset where

each LC neuron was randomly targeted and individually tested for this type of interaction.

Electrical connection between LC/NE neurons

Several previous studies suggest that electrical coupling (EC) in the LC is negligible or absent in adult

animals (Christie & Jelinek, 1993; Christie et al., 1989; Travagli et al., 1995). This is surprising given

that electrical coupling is an appealing mechanism for LC/NE neurons to synchronize to cope with the

global demand for NE signaling. In contrast to these studies, our multi-cell recordings provide a direct

demonstration of electrical coupling established by mature LC/NE neurons with a coupling rate

consistent with a previous report from extracellular recording (N. K. Totah et al., 2018), substantiating

prior evidence that electrical coupling is still an important mechanism for LC/NE neurons to interact with

each other in adult animals (Alvarez, Chow, Van Bockstaele, & Williams, 2002; Rash et al., 2007; Van

Bockstaele, Garcia-Hernandez, Fox, Alvarez, & Williams, 2004). An electrical coupling rate ranging

from 15% to 80% has been reported in many brain regions based on pairwise electrophysiological

testing of narrowly spaced cell pairs of the same cell type from juvenile animals (Curti et al., 2012;

Devor & Yarom, 2002; Long, Landisman, & Connors, 2004; Tamas, Buhl, Lorincz, & Somogyi, 2000).

While these rates were often reported without the inter-soma distance information, several studies

indicate the coupling rate often drops to zero once the inter-soma distance increases to more than 40

µm (Beierlein, Gibson, & Connors, 2000; Curti et al., 2012; Galarreta & Hestrin, 2001, 2002; Gibson,

Beierlein, & Connors, 1999; Tamas et al., 2000). These spatial and cell type-specific rules appear to

apply to electrical coupling among mature LC neurons as well. While the overall coupling rate is low

among all LC cell pairs, it is much higher when cell pairs are narrowly spaced (with inter-soma distance

< 40 µm), particularly for FF-FFs. Interestingly, when a cell pair is more than 40 µm apart, the possibility

of finding an electrical connection, despite being very low, is much less distance-dependent, possibly

due to the dominant dendrodendritic form of electrical synapse formed between these neurons (see

below). In addition, electrical coupling appears to decrease with age in most brain structures (Peinado,

Yuste, & Katz, 1993), including in LC (Christie & Jelinek, 1993; Christie et al., 1989; Travagli et al.,

1995). However, a direct comparison of the coupling rate in adults to juvenile animals did not reveal a

significant difference between the two ages, indicating that electrical coupling in the LC may be less

subject to developmental change than previously thought.

It is well documented that gap junctions generally form between cell pairs of the same type

(homotypic gap junctions), while gap junctions between different cell types (heterotypic gap junctions)

are rare or absent in most brain regions (Peinado et al., 1993). This rule also applies to LC neurons:

most electrical couplings occur between cell pairs of the same noradrenergic type, while electrical

coupling is rare between cell pairs of the different cell types. This is still true when the inter-soma

distance is controlled. Comparing the spatial profiles of homotypic gap junctions versus heterotypic gap

junctions, we found that neurons appear to form gap junctions exclusively with narrowly spaced,

adjacent LC neurons of the same type (homotypic connection), while the heterotypic connections,

despite being very rare, typically form between cell pairs that are more spatially apart. The result

suggests that each LC neuron is intricately connected with their very close neighboring neurons of the

same type to constitute cell type-specific electrical subnetworks (FF subnetwork vs. MP subnetwork),

each of which may synchronize independently via extensive gap junctions. This finding provides a

plausible cellular mechanism for prior observations that LC pairwise correlated firing and LC ensembles

identified in extracellular recordings are cell-type specific (Noei, Zouridis, Logothetis, Panzeri, & Totah,

2022; N. K. Totah et al., 2018). Of note, the coupling rate between FF–FF pairs is more than threefold

higher than that between MP-MP pairs, suggesting that the MP electrical subnetwork may be smaller in

terms of cell population than the FF subnetwork. Alternatively, a lower coupling rate among MPs may

suggest that MPs are more heterogeneous than FFs. Indeed, as discussed above, LC neurons with

multipolar-shaped soma are widely distributed across multiple LC subregions and more heterogeneous

than those fusiform cells. In addition, putative MPs in distinct LC subregions appear to have different

preferred projection targets (Loughlin et al., 1986; Waterhouse et al., 1983). Therefore, MP cells in

each LC subregion may only electrically connect with MP cells in the same LC subregion, forming

subregion-specific MP subnetworks that may perform distinct functions (Fig. 7C). Given that FFs and

MPs (or their subgroups) may have different preferential locations and distinct projection targets, within-

cell type electrical coupling is likely to promote the spread of electrical and chemical signals and spike

synchronization within each subnetwork, exerting their synergetic effect on specific target regions.

While close appositions between two neurons do not necessarily mean electrical synapses, such

contact sites identified under the light microscope could be reliably confirmed as real electrical

synapses by follow-up electron microscope studies (Fukuda & Kosaka, 2000, 2003; Tamas et al., 2000;

Vervaeke, Lorincz, Nusser, & Silver, 2012). Our examination of close appositions between electrically-

coupled pairs indicates more than 50% of electrically-coupled pairs have only one putative synapse,

and multiple contact sites between pairs are less frequent. Almost all putative synapses were found to

occur between dendrites, corroborating a previous LC study in rats (Ishimatsu & Williams, 1996). In

fact, dendro-dendritic synapses appear to be a dominant form of electrical synapses across brain

regions (Fukuda & Kosaka, 2000, 2003; Hidaka et al., 2004; Sotelo, Llinas, & Baker, 1974; Vervaeke et

al., 2010; Waterhouse et al., 1983), while somato-dendritic or somato-somatic ones are much less

common (Curti et al., 2012; Tamas et al., 2000). In addition, electrical synapses in dendrites are

localized further away from the soma as animals get older (Fukuda & Kosaka, 2000, 2003). Consistent

with these previous findings, putative synapses in mature LC were located most frequently in the

middle part of the entire dendritic trees of the electrically-coupled cell pair, with a similar dendritic

distance to each soma. Given electrical synapses being located at such distal sites, fast signals like

action potentials in prejunctional cells could be low-pass filtered and highly attenuated when

propagating to postjunctional cells, giving rise to small-amplitude coupled potentials (like spikelet) at the

somata of post-junctional neurons. How could such a small coupled potential drive synchronized firing

and network oscillations under natural stimulation conditions? Recent work has shown those gap

junctions found at distal dendritic sites can interact locally with chemical synaptic inputs to drive

synchronized spiking. While coupled potentials are small at the soma, these electrical signals are

expected to be larger near gap junctions, where they enable excitatory synaptic charge to spread into

the dendrites of neighboring coupled neurons (Vervaeke et al., 2012), and/or sum with chemical

synaptic inputs to generate dendritic spikes that actively propagate with little delay to the soma and

drive action potentials with high fidelity. The former may promote the synchronized response of an

electrically-coupled neuronal population to certain specific synaptic inputs, while the latter allows weak

distal gap junction inputs to synchronize somatic action potentials, promoting synchronized spiking of

electrically-coupled neurons upon common chemical synaptic input (Migliore, Hines, & Shepherd, 2005;

Schoppa & Westbrook, 2002; Trenholm et al., 2014). It will be interesting to investigate in future studies

the mechanism(s) that LC neurons deploy to leverage their gap junctions to promote inter-neuronal

synchrony.

The network of electrically-coupled LC/NE neurons

Electrical coupling is not just about connections between two neurons, instead it connects a group of

neurons into an electrically-coupled network that coordinates/synergizes their activity as a whole. Thus,

two questions ensue: what is the population size of each network, and how does an individual neuron in

each network connect to other neurons of the same network? Despite extensive studies of gap

junctions across almost all brain regions, these questions remain largely unexplored due to technical

limitations. Gap junction studies with the dye-coupling approach may provide an insight into these

questions (Curti et al., 2012; Long et al., 2004), but the neuron clusters labeled by this approach could

result from a second-order coupling, in addition to first-order coupling – these couplings are not

otherwise discriminable – greatly compounding interpretation. In addition, the relative impermeability of

different dyes to different kinds of electrical synapses results in many inconsistent results. In this

regard, our multi-patch recordings confer some technical advantages over the dye-based approach.

First, given gap junctions being located at the dendrite far away from the soma, somatic membrane

potentials in a pre-junctional cell are likely to be highly attenuated by the time they reach a post-

junctional cell. It is reasonable to conclude that such highly attenuated coupled potential from the first-

order coupling can hardly propagate any further into cells through second-order coupling. Thus, the gap

junctions with a dominant dendrodendritic form identified with pair-wise recordings should be essentially

first-order gap junctions, but this assertion has never been directly confirmed as yet. Our multi-cell

recording sets allowed for directly testing said assertion and ruled out the possibility that gap junctions

identified with pair-wise patch recordings in LC could result from a second-order coupling. However,

caution should be taken in interpreting the results if electrical synapses are located in the soma (Curti et

al., 2012), assuming the signals passing through somatosomatic electrical synapses are less

attenuated. Given that the electrical couplings we identified in LC neurons are confirmed as first-order,

we can then examine the real organization of electrical connections beyond two cells in the context of

the network. Second, our multiple-cell recording allows for identifying multiple electrically-coupled pairs

across a cohort of up to eight cells, providing the first glimpse into how each neuron interacts with the

rest of the neurons in an electrically-coupled neuronal network.

With the support from the simulation with a random connectivity model, a chain-like pattern of two

connections (A↔B↔C) is overrepresented in our dataset. One might argue that said

overrepresentation may be accounted for by spatial and cell-type specific connectivity rules that were

empirically determined by us here. However, revising our model by adding these connectivity rules did

not significantly improve simulation results, and a chain-like pattern of two connections (A↔B↔C)

remains unexplained. All simulations suggest that, in addition to spatial and cell type-specific rules in

governing LC electrical connections, there is an additional chain-like connectivity rule that may

contribute to the overrepresentation of a chain-like pattern of two connections in our dataset.

Furthermore, while the number of three connections is low, a chain-like pattern (A↔B↔C↔D) is still a

dominant pattern of three connections in our recording sets. Our simulation on the pattern of three

connections yielded similar results and supported the idea of overrepresentation of a chain-like

electrical connectivity pattern in LC/NE neurons. In addition, the overrepresentation can not be

explained by the spatial and cell type-specific connectivity rules, further suggesting an additional chain-

like connectivity rule for the organization of three connections. While we could not test if this additional

connectivity rule may apply to four or more connections for now (no four connections were found in our

dataset), the chain-like connectivity rule may be a general wiring principle dictating a whole electrically-

coupled network composed of hundreds of neurons (Fig. 7C). Indeed, previous dye-coupled studies

suggest the same connectivity rule in many brain regions and cell types, including LC (Christie &

Jelinek, 1993; Curti et al., 2012; Lee, Patrick, Richardson, & Connors, 2014; Long et al., 2004). If a

chain-like organizing pattern is true, then the dye injection of a single cell can only label the maximum

of two additional cells if only first-order coupled cells are labeled. Indeed, the cluster of labeled cells in

these studies is often small with a maximum of two additional labeled cells (Christie & Jelinek, 1993;

Curti et al., 2012). Some studies do label more than two additional cells, but additional cells may be

labeled via second-order coupling with a long dye injection time (Devor & Yarom, 2002; Long et al.,

2004). The most convincing evidence for this rule comes from the mesencephalic trigeminal nucleus,

where gap junctions between five neurons have been elegantly shown to arrange like a chain-like

pattern across neurons (Curti et al., 2012). However, the interpretation of these data is restrained by

several issues inherent to this approach (see above), and whether or not this chain-like connection

pattern applies to the whole LC electrically-coupled network needs to be further explored in the LC, as

well as in many other brain regions. In addition, the other two possible EC organization patterns,

circular (Fig. 6D, right panel 4[th] cartoon) or hub-like connection (see Fig. 6D, right panel 3[rd] cartoon)

patterns, were not observed in our dataset, suggesting that electrically-coupled pairs in LC may not be

organized as these two patterns. However, simulated results on current data could not rule out these

two patterns. Model 3 simulations predicted 2.45 hub-like connections on average (95% CI: [0,5]) and

predicted 1.39 circular-like connections on average (95% CI: [0,4]). While both average predicted

numbers are numerically higher than the observed numbers of each (0), the observed numbers are still

within statistical bounds. More advanced recording techniques may be able to answer these questions

in the future.

685
Acknowledgments

We thank AM's thesis committee members for their suggestions and comments, as well as the

comments and support from the members of Jiang Lab. We acknowledge funding from NINDS R01

NS101596 (XJ, AM, MH), NIMH R01 MH109556 (JJ, XJ, MH), as well as Training Grant T32 EY07001

(AM) to support this work. Research reported in this publication was also supported by the Eunice

Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of

Health under Award Number P50HD103555 for use of the Microscopy Core facilities. The content is

solely the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health.

Author contributions

A. McKinney, X. Jiang designed the experiments. A. McKinney, X. Jiang, J. Jing performed the

experiments. X. Jiang, A. McKinney, M. Hu, S. Patel, BR. Sheth, A. Hoskins, A. Mohammadyar, N.

Naeem analyzed the data. X. Jiang wrote the manuscript with input from other authors and supervised

all aspects of the project.

Disclosure of Conflicts of Interest

The authors declare no conflicts of interest.

Data and Code Availability

All data generated or analyzed during this study are included in the manuscript and supporting files. All

data and Custom analysis codes supporting the findings of this study are available at

doi:10.5061/dryad.kh1893283.

Materials and methods

Animals

All experiments were performed according to the guidelines of the Institutional Animal Care and Use

Committee (IACUC) of Baylor College of Medicine. Experiments on adult male and female mice

(median age: 79, full range: 55--175 days) were performed using DBH-Cre/Ai9 (n = 121) or wild-type (n

= 15). Crossing DBH-Cre (JAX stock No: 033951) mice with Ai9 reporter mice globally labels all

noradrenergic neurons including LC neurons (Rukhadze et al., 2017). Additional younger DBH-Cre/Ai9

mice (P13-21, n = 13) were used to study connectivity between LC neurons at the juvenile age for

comparison with the adult age.

Slice preparation

Slice preparation from adult mouse brainstem follows an N-Methyl-D-glucamine (NMDG) slicing

protocol (Jiang et al., 2015; Ting et al., 2014). Briefly, animals were deeply anesthetized using 3%

isoflurane. After decapitation, the brain was removed and placed into cold (0−4 °C) oxygenated NMDG

solution containing 93 mM NMDG, 93 mM HCl, 2.5 mM KCl, 1.2 mM NaH_2PO_4, 30 mM $NaHCO_3$, 20

mM HEPES, 25 mM glucose, 5 mM sodium ascorbate, 2 mM Thiourea, 3 mM sodium pyruvate, 10mM

$MgSO_4$ and 0.5 mM $CaCl_2$, pH 7.35 (all from SIGMA-ALDRICH). Horizontal slices were prepared using

a vibratome (200 μm thick) with zirconia ceramic blades. The brain slices were kept at 37.0 ± 0.5 °C in

oxygenated NMDG solution for 10 minutes, and then were transferred to an artificial cerebrospinal fluid

(ACSF) containing 125 mM NaCl, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 25 mM $NaHCO_3$, 1 mM $MgCl_2$, 25

mM glucose, and 2 mM $CaCl_2$ (pH 7.4) for at least 1 hour prior to the beginning of recordings. During

the recording sessions, the slices were submerged in a custom chamber and were stabilized with a fine

nylon net attached to a custom-designed platinum ring. This recording chamber was continuously

perfused with oxygenated ACSF throughout the recording session. Brainstem slices from juvenile mice

were prepared following a similar protocol except for the cutting solution (with regular ACSF, not

NMDG). For the recording of hippocampal neurons, the parasagittal cortical slices were prepared

following the same protocol for adult animals.

Electrophysiology

Whole-cell recordings were performed as described previously (Jiang et al., 2015; Scala et al., 2020;

Scala et al., 2019). Briefly, patch pipettes (4−7 MΩ) were filled with an internal solution containing 120

mM potassium gluconate, 10 mM HEPES, 4 mM KCl, 4 mM MgATP, 0.3 mM Na_3GTP, 10 mM sodium

phosphocreatine and 0.5% biocytin (pH 7.25). Simultaneous whole-cell recordings from up to 8 neurons

were performed using two Quadro EPC 10 amplifiers (HEKA Electronic, Germany). PatchMaster

(HEKA) and custom-written Matlab-based programs (Mathworks) were used to operate the recording

system and perform online and offline data analysis. In current-clamp recordings, neurons were first

current clamped at ~-40 pA to prevent its spontaneous firings (see results). Action potentials were

evoked by current injection into presynaptic neurons at 2 nA for 2 ms at 0.05 Hz for 20−50 trials, and

the average of the sweeps in postsynaptic cells was used to detect synaptic connections. The spike

threshold and firing pattern in response to sustained depolarizing currents were recorded for each

neuron by injecting increasing current steps (+10 pA).

For coupling coefficient (CC) calculations in gap junctions, hyperpolarizing current pulses of 600

ms duration were injected into one cell and resulting voltage deflections were measured in both cells.

The voltage deflections were measured at the last 10-20 ms of current injections given the large time

constant of the membrane potential responses of two cells (see Results). The coupling coefficient from

cell 1 to cell 2 (CC1) was defined as V2/V1 and the coupling coefficient in the opposite direction from

cell 2 to cell 1 (CC2) was defined as V1/ V2, where V1 is the voltage deflection in one cell and V2 is the

corresponding voltage deflection in the other cell. To improve the signal-to-noise ratio, a total of 4 to 30

single responses were often generated and averaged. For each of the coupled pairs, the mean

coupling coefficient was calculated as the average from the values in both directions.

Electrophysiological features extraction

A custom-made Python software kit named as Patchview was developed to analyze the

electrophysiology data offline (Ming Hu, 2022). We extracted 13 electrophysiological properties from

each neuron, following a similar method as used in previous studies (Gouwens et al., 2019; Scala et al.,

2021). The rheobase refers to the minimal stimulation current to elicit any action potentials. To extract

the properties of action potential (AP), we used the first AP elicited by our step stimulation protocol. To

calculate the threshold of the AP, the trace within 20 ms preceding the AP peak was used to calculate

the third derivative first and the data point with the maximal third derivative value was defined as the

threshold of the AP. The AP amplitude is the difference between the threshold potential and the peak

potential of the AP. The AP width was defined as that at half-height. The time range from the threshold

point to the peak is 'rising phase time' while 'decay phase time' is the time range from the peak to the

potential equal to the threshold. The afterhyperpolarization was defined as the voltage difference

between the threshold and the minimal value of the trough after the AP. The time range from the

stimulation onset to the threshold of the first AP at the rheobase sweep was defined as the AP delay

time.

To calculate the time constant τ, we fitted each hyperpolarizing trace (from the stimulation onset

to the minimal potential within the first-half stimulation period) with an exponential function as follows:

774
$$f(t) = ae^{(-t/\tau)} + C$$

Here, *t* is the time relative to the stimulation onset, *a* and *C* are two constants, *f(t)* is the current value at

time *t*, and τ is the membrane constant. Then the τ was calculated for each trace and averaged across

all traces.

To calculate the maximal firing rate, the depolarizing trace with the maximal number of APs

elicited by the step protocol was used to count the number of APs, which was then divided by the

stimulation duration (600 ms). To calculate the adaptive index, the inter-spike intervals (ISIs) were first

measured, and then the ratio of the last and the first ISIs was calculated. The CV was calculated as the

ratio of the standard deviation of ISIs and the mean of ISIs. To compare the first AP and the

subsequent ones, we calculated the ratio of their AP amplitude and the ratio of their half-widths. The

amplitude and half-width of the first AP were first calculated and then divided by the mean of these

values of the subsequent APs.

To calculate the input resistance, we measured the steady-state membrane potentials in the

traces elicited by the first five negative currents in the step protocol, and then plotted these values (y-

axis) against each of the corresponding currents (x-axis). We fitted the plot with a linear function and

the slope of the fitting line is the input resistance, and the potential at the x-axis intersection is the RMP.

The RMPs were reported without correcting junction potentials.

Morphological reconstructions and analysis

Morphological reconstructions and analysis were performed using a 100X oil immersion objective lens

and camera lucida system using Neurolucida (MicroBrightField), following the same procedure as

previously described (Jiang et al., 2015; Scala et al., 2020; Scala et al., 2019). Briefly, neurons were

filled with biocytin during the recording, and slices were fixed right after recordings by immersion in

freshly prepared 2.5% glutaraldehyde (from Electron Microscopy Science Cat. no. 16220), 4%

paraformaldehyde (from SIGMA-ALDRICH Cat.no. P6148) in 0.1 M phosphate-buffered saline at 4°C

for 7-10 days, and then processed with the avidin-biotin-peroxidase method to reveal cell morphology.

Slices were then mounted in an aqueous Mowiol mounting solution (SIGMA-ALDRICH) and the

morphologically recovered cells were examined, reconstructed, and analyzed using a 100X oil-

immersion objective lens and camera lucida system. Tissue shrinkage due to the fixation procedure

was not compensated. The shrinkage of the tissue surrounding the biocytin-stained cells is about 10-

20%, consistent with previous studies (Markram, Lubke, Frotscher, Roth, & Sakmann, 1997; Scala et

al., 2019). The reconstructions were not corrected for tissue shrinkage.

Dendritic arbor and axonal structure of reconstructed morphologies of n = 70 cells were analyzed

using Neurolucida software. Additional morphological analysis was performed using custom Python

code.

Data Analysis

Data analysis was performed using custom-written MATLAB software and Python routines. For

statistical comparisons, the normality of the data was first tested using the Shapiro–Wilk test. If

compared groups show normal distributions, Student's t-test or the paired t-test was used; otherwise,

the non-parametric paired Wilcoxon-sign rank test and Mann–Whitney test were used. One-way or two-

way ANOVA was also used for comparing morpho-electric features of different types of neurons.

Data Simulation

Model 1: random connectivity model

First, the overall pairwise connection probability (P) was obtained from the recordings and is the ratio of

the observed number of electrical connections to the total number of potential electrical connections

tested across all recordings, regardless of the neuronal type and inter-neuronal distance. Expected

numbers of double or 2-connections were calculated based on pairwise connection probabilities. For

instance, if there were four neurons A, B, C, and D in a particular recording/slice, then the expected

value of a non-chain-like 2-connection AB-CD was calculated to be equal to P(pairwise connection) *

P(pairwise connection). The expected number of chain-like and non-chain-like 2-connections for a

given slice was the sum of the expected values of all 2-connections for the recording. The grand total

expected number of 2-connections of a given type was the sum of the corresponding expected number

values over all our recordings. Similar simulations were conducted for triple or 3-connections, and the

product of three pairwise connections was used for all calculations.

Model 2: Differentiated neuronal types

For each individual recording (slice) for which the slice was fixed and neuronal types (FF – fusiform cell

type; MP – multipolar cell type) were identified, we calculated the expected numbers of chain-like and

non-chain-like 2- and 3-connections in the following way. For a given pair of neurons from a given

recording slice, its corresponding pairwise connection probability (FF-FF, FF-MP , or MP-MP depending

on the morphological cell types of the neurons constituting the pair) across all recordings, which was

computed earlier, was used for all calculations (Fig.4G). Expected numbers of 2-connections were

calculated based on pairwise connection probabilities. For instance, if neurons A, B, C, and D in a

particular recording/slice were FF, FF, MP, and MP neuronal types respectively, then the expected

value of a chain-like 2-connection AB-BC was calculated to be equal to P(FF-FF connection) * P(FF-

MP connection). Similar calculations were performed for all possible 2-connections (e.g. chain-like 2-

connection AB-CD and non-chain-like connections AB-CD in the above example). The expected

number of chain-like and non-chain-like 2-connections for a given slice was then the sum of the

individual probabilities thus calculated. The grand total expected number of 2-connections of a given

type was computed by summing the corresponding expected number values over all our recordings.

Calculations to compute the expected values of triple or 3-connections were similar, with the exception

that the expected value was now a product of three, and not two, pairwise connections; in the above

example, P(AB-BC-CD) = P(FF-FF connection) * P(FF-MP connection) * P (MP-MP connection).

Model 3: Differentiated neuronal types, short-range connectivity

In addition to the constraints specified in Model 2, namely pairwise connection probabilities based on

neuronal identity, the following additional constraints were added to Model 3. First, because

connections between neuronal pairs were largely non-existent for inter-neuronal distances > 80 μm, all

neuronal pairs in the model that were > 80 μm apart were taken to have no connection. Second,

because neurons of the same type clustered together in the recordings, FF-FF and MP-MP connection

probabilities for short distances (16-40 μm) were used, and because neurons of different types were

further apart in the recordings, FF-MP connection probability for 0-80 mm distance was used in the

model. The remaining computations were performed in accord with those of the previous model.

Monte-Carlo simulations

On each of 1000 runs of the simulation, we counted the number of 2- and 3-connections from all

simulated recordings. Each simulated recording matched one-to-one a biological recording in terms of

the number and type of neurons. A random number generated by the random number generator

determined whether a connection existed between each neuron in the corresponding simulated

recording. If the random number was lower than the expected value of the connection (different values

based on whether the corresponding connection in the recording was FF-FF/FF-MP/MP-MP), the

simulated connection was considered to exist. For each run, simulated double and triple connections

were obtained for each recording and the total simulated number was obtained by summing over all the

simulated recordings. Summary statistics were obtained from the simulated data from 1000 such runs:

i) the mean, standard deviation, and 95% confidence intervals for the simulated numbers of chain-like

and non-chain-like 2- and 3-connections, and ii) the probability that the ratio of the simulated chain-like

over non-chain-like 2-connections (3-connections) was greater than the ratio of the observed number of

chain-like over non-chain-like 2-connections (3-connections). All simulations were conducted on

MATLAB (R2021a, The Mathworks, Natick, MA).

References

Aghajanian, G. K., & VanderMaelen, C. P. (1982). alpha 2-adrenoceptor-mediated hyperpolarization of locus coeruleus neurons: intracellular studies in vivo. *Science, 215*(4538), 1394-1396. doi:10.1126/science.6278591

Alvarez, V. A., Chow, C. C., Van Bockstaele, E. J., & Williams, J. T. (2002). Frequency-dependent synchrony in locus ceruleus: role of electrotonic coupling. *Proc Natl Acad Sci U S A, 99*(6), 4032-4036. doi:10.1073/pnas.062716299

Andrzejewski, M., Muckenhoff, K., Scheid, P., & Ballantyne, D. (2001). Synchronized rhythms in chemosensitive neurons of the locus coeruleus in the absence of chemical synaptic transmission. *Respir Physiol, 129*(1-2), 123-140. doi:10.1016/s0034-5687(01)00300-0

Ballantyne, D., Andrzejewski, M., Muckenhoff, K., & Scheid, P. (2004). Rhythms, synchrony and electrical coupling in the Locus coeruleus. *Respir Physiol Neurobiol, 143*(2-3), 199-214. doi:10.1016/j.resp.2004.07.018

Bari, A., Xu, S., Pignatelli, M., Takeuchi, D., Feng, J., Li, Y., & Tonegawa, S. (2020). Differential attentional control mechanisms by two distinct noradrenergic coeruleo-frontal cortical pathways. *Proc Natl Acad Sci U S A, 117*(46), 29080-29089. doi:10.1073/pnas.2015635117

Bast, N., Poustka, L., & Freitag, C. M. (2018). The locus coeruleus-norepinephrine system as pacemaker of attention - a developmental mechanism of derailed attentional function in autism spectrum disorder. *Eur J Neurosci, 47*(2), 115-125. doi:10.1111/ejn.13795

Beierlein, M., Gibson, J. R., & Connors, B. W. (2000). A network of electrically coupled interneurons drives synchronized inhibition in neocortex. *Nat Neurosci, 3*(9), 904-910. doi:10.1038/78809

Bennett, M. V., & Zukin, R. S. (2004). Electrical coupling and neuronal synchronization in the Mammalian brain. *Neuron, 41*(4), 495-511. doi:10.1016/s0896-6273(04)00043-1

Blythe, S. N., Wokosin, D., Atherton, J. F., & Bevan, M. D. (2009). Cellular mechanisms underlying burst firing in substantia nigra dopamine neurons. *J Neurosci, 29*(49), 15531-15541. doi:10.1523/JNEUROSCI.2961-09.2009

Breton-Provencher, V., & Sur, M. (2019). Active control of arousal by a locus coeruleus GABAergic circuit. *Nat Neurosci, 22*(2), 218-228. doi:10.1038/s41593-018-0305-z

Cedarbaum, J. M., & Aghajanian, G. K. (1976). Noradrenergic neurons of the locus coeruleus: inhibition by epinephrine and activation by the alpha-antagonist piperoxane. *Brain Res, 112*(2), 413-419. doi:10.1016/0006-8993(76)90297-3

Chandler, D. J., Gao, W.-J., & Waterhouse, B. D. (2014). Heterogeneous organization of the locus coeruleus projections to prefrontal and motor cortices. *Proceedings of the National Academy of Sciences, 111*(18), 6816-6821. doi:10.1073/pnas.1320827111

Christie, M. J., & Jelinek, H. F. (1993). Dye-coupling among neurons of the rat locus coeruleus during postnatal development. *Neuroscience, 56*(1), 129-137. doi:10.1016/0306-4522(93)90568-z

Christie, M. J., Williams, J. T., & North, R. A. (1989). Electrical coupling synchronizes subthreshold activity in locus coeruleus neurons in vitro from neonatal rats. *J Neurosci, 9*(10), 3584-3589.

Courtney, N. A., & Ford, C. P. (2014). The timing of dopamine- and noradrenaline-mediated transmission reflects underlying differences in the extent of spillover and pooling. *J Neurosci, 34*(22), 7645-7656. doi:10.1523/JNEUROSCI.0166-14.2014

Curti, S., Hoge, G., Nagy, J. I., & Pereda, A. E. (2012). Synergy between electrical coupling and membrane properties promotes strong synchronization of neurons of the mesencephalic trigeminal nucleus. *J Neurosci, 32*(13), 4341-4359. doi:10.1523/JNEUROSCI.6216-11.2012

Davy, O., Perrins, R., Lavigne, M., Kremer, E., Tsaneva-Atanasova, K., Ashby, M., & Pickering, A. E. (2022). Noradrenergic cross-modular reciprocal inhibition within the locus coeruleus. *bioRxiv*, 2022.2009.2007.506929. doi:10.1101/2022.09.07.506929

Devor, A., & Yarom, Y. (2002). Electrotonic coupling in the inferior olivary nucleus revealed by simultaneous double patch recordings. *J Neurophysiol, 87*(6), 3048-3058. doi:10.1152/jn.2002.87.6.3048

Egan, T. M., Henderson, G., North, R. A., & Williams, J. T. (1983). Noradrenaline-mediated synaptic inhibition in rat locus coeruleus neurones. *J Physiol, 345*, 477-488. doi:10.1113/jphysiol.1983.sp014990

Erdos, P., & Rényi, A. (1984). On the evolution of random graphs. *Transactions of the American Mathematical Society, 286*, 257-257.

Fortress, A. M., Hamlett, E. D., Vazey, E. M., Aston-Jones, G., Cass, W. A., Boger, H. A., & Granholm, A. C. (2015). Designer receptors enhance memory in a mouse model of Down syndrome. *J Neurosci, 35*(4), 1343-1353. doi:10.1523/JNEUROSCI.2658-14.2015

Fukuda, T., & Kosaka, T. (2000). Gap junctions linking the dendritic network of GABAergic interneurons in the hippocampus. *J Neurosci, 20*(4), 1519-1528.

Fukuda, T., & Kosaka, T. (2003). Ultrastructural study of gap junctions between dendrites of parvalbumin-containing GABAergic neurons in various neocortical areas of the adult rat. *Neuroscience, 120*(1), 5-20. doi:10.1016/s0306-4522(03)00328-2

Fung, S. I., Chan, J. Y., Manzoni, D., White, S. R., Lai, Y. Y., Strahlendorf, H. K., . . . Barnes, C. D. (1994). Cotransmitter-mediated locus coeruleus action on motoneurons. *Brain Res Bull, 35*(5-6), 423-432. doi:10.1016/0361-9230(94)90155-4

Fung, S. J., Reddy, V. K., Liu, R. H., Wang, Z., & Barnes, C. D. (1994). Existence of glutamate in noradrenergic locus coeruleus neurons of rodents. *Brain Res Bull, 35*(5-6), 505-512. doi:10.1016/0361-9230(94)90164-3

Galarreta, M., & Hestrin, S. (2001). Electrical synapses between GABA-releasing interneurons. *Nat Rev Neurosci, 2*(6), 425-433. doi:10.1038/35077566

Galarreta, M., & Hestrin, S. (2002). Electrical and chemical synapses among parvalbumin fast-spiking GABAergic interneurons in adult mouse neocortex. *Proc Natl Acad Sci U S A, 99*(19), 12438-12443. doi:10.1073/pnas.192159599

Gibson, J. R., Beierlein, M., & Connors, B. W. (1999). Two networks of electrically coupled inhibitory neurons in neocortex. *Nature, 402*(6757), 75-79. doi:10.1038/47035

Gouwens, N. W., Sorensen, S. A., Berg, J., Lee, C., Jarsky, T., Ting, J., . . . Koch, C. (2019). Classification of electrophysiological and morphological neuron types in the mouse visual cortex. *Nat Neurosci, 22*(7), 1182-1195. doi:10.1038/s41593-019-0417-0

Hausser, M., Stuart, G., Racca, C., & Sakmann, B. (1995). Axonal initiation and active dendritic propagation of action potentials in substantia nigra neurons. *Neuron, 15*(3), 637-647. doi:10.1016/0896-6273(95)90152-3

Hidaka, S., Akahori, Y., & Kurosawa, Y. (2004). Dendrodendritic electrical synapses between mammalian retinal ganglion cells. *J Neurosci, 24*(46), 10553-10567. doi:10.1523/JNEUROSCI.3319-04.2004

Hirschberg, S., Li, Y., Randall, A., Kremer, E. J., & Pickering, A. E. (2017). Functional dichotomy in spinal- vs prefrontal-projecting locus coeruleus modules splits descending noradrenergic analgesia from ascending aversion and anxiety in rats. *Elife, 6*. doi:10.7554/eLife.29808

Ho¨kfelt, T., Fuxe, K., Goldstein, M., & Johansson, O. (1974). Immunohistochemical evidence for the existence of adrenaline neurons in the rat brain. *Brain Research, 66*, 235-251.

Holets, V. R., Hokfelt, T., Rokaeus, A., Terenius, L., & Goldstein, M. (1988). Locus coeruleus neurons in the rat containing neuropeptide Y, tyrosine hydroxylase or galanin and their efferent projections to the spinal cord, cerebral cortex and hypothalamus. *Neuroscience, 24*(3), 893-906. doi:10.1016/0306-4522(88)90076-0

Huang, H. P., Wang, S. R., Yao, W., Zhang, C., Zhou, Y., Chen, X. W., . . . Zhou, Z. (2007). Long latency of evoked quantal transmitter release from somata of locus coeruleus neurons in rat pontine slices. *Proc Natl Acad Sci U S A, 104*(4), 1401-1406. doi:10.1073/pnas.0608897104

Ishimatsu, M., & Williams, J. T. (1996). Synchronous activity in locus coeruleus results from dendritic interactions in pericoerulear regions. *J Neurosci, 16*(16), 5196-5204.

Jiang, X., Shen, S., Cadwell, C. R., Berens, P., Sinz, F., Ecker, A. S., . . . Tolias, A. S. (2015). Principles of connectivity among morphologically defined cell types in adult neocortex. *Science, 350*(6264), aac9462. doi:10.1126/science.aac9462

Kuo, C. C., Hsieh, J. C., Tsai, H. C., Kuo, Y. S., Yau, H. J., Chen, C. C., . . . Min, M. Y. (2020). Inhibitory interneurons
regulate phasic activity of noradrenergic neurons in the mouse locus coeruleus and functional
implications. *J Physiol, 598*(18), 4003-4029. doi:10.1113/JP279557
Lee, S. C., Patrick, S. L., Richardson, K. A., & Connors, B. W. (2014). Two functionally distinct networks of gap
junction-coupled inhibitory neurons in the thalamic reticular nucleus. *J Neurosci, 34*(39), 13170-13182.
doi:10.1523/JNEUROSCI.0562-14.2014
Li, Y., Hickey, L., Perrins, R., Werlen, E., Patel, A. A., Hirschberg, S., . . . Pickering, A. E. (2016). Retrograde
optogenetic characterization of the pontospinal module of the locus coeruleus with a canine adenoviral
vector. *Brain Res, 1641*(Pt B), 274-290. doi:10.1016/j.brainres.2016.02.023
Liu, R. H., Fung, S. J., Reddy, V. K., & Barnes, C. D. (1995). Localization of glutamatergic neurons in the
dorsolateral pontine tegmentum projecting to the spinal cord of the cat with a proposed role of
glutamate on lumbar motoneuron activity. *Neuroscience, 64*(1), 193-208.
doi:https://doi.org/10.1016/0306-4522(94)00354-8
Long, M. A., Landisman, C. E., & Connors, B. W. (2004). Small clusters of electrically coupled neurons generate
synchronous rhythms in the thalamic reticular nucleus. *J Neurosci, 24*(2), 341-349.
doi:10.1523/JNEUROSCI.3358-03.2004
Loughlin, S. E., Foote, S. L., & Grzanna, R. (1986). Efferent projections of nucleus locus coeruleus: morphologic
subpopulations have different efferent targets. *Neuroscience, 18*(2), 307-319. doi:10.1016/0306-
4522(86)90156-9
Lusk, S. J., McKinney, A., Hunt, P. J., Fahey, P. G., Patel, J., Chang, A., . . . Ray, R. S. (2022). A CRISPR toolbox for
generating intersectional genetic mouse models for functional, molecular, and anatomical circuit
mapping. *BMC Biol, 20*(1), 28. doi:10.1186/s12915-022-01227-0
Luskin, A. T., Li, L., Fu, X., Barcomb, K., Blackburn, T., Li, E. M., . . . Bruchas, M. R. (2022). A diverse network of
pericoerulear neurons control arousal states. *bioRxiv*, 2022.2006.2030.498327.
doi:10.1101/2022.06.30.498327
Madisen, L., Garner, A. R., Shimaoka, D., Chuong, A. S., Klapoetke, N. C., Li, L., . . . Zeng, H. (2015). Transgenic
mice for intersectional targeting of neural sensors and effectors with high specificity and performance.
Neuron, 85(5), 942-958. doi:10.1016/j.neuron.2015.02.022
Markram, H., Lubke, J., Frotscher, M., Roth, A., & Sakmann, B. (1997). Physiology and anatomy of synaptic
connections between thick tufted pyramidal neurones in the developing rat neocortex. *J Physiol, 500 (Pt*
2), 409-440. doi:10.1113/jphysiol.1997.sp022031
Melander, T., Hokfelt, T., Rokaeus, A., Cuello, A. C., Oertel, W. H., Verhofstad, A., & Goldstein, M. (1986).
Coexistence of galanin-like immunoreactivity with catecholamines, 5-hydroxytryptamine, GABA and
neuropeptides in the rat CNS. *J Neurosci, 6*(12), 3640-3654.
Migliore, M., Hines, M. L., & Shepherd, G. M. (2005). The role of distal dendritic gap junctions in synchronization
of mitral cell axonal output. *J Comput Neurosci, 18*(2), 151-161. doi:10.1007/s10827-005-6556-1
Ming Hu, X. J. (2022). PatchView: A Python Package for Patch-clamp Data Analysis and Visualization. *Journal of*
Open Source Software, 7(78), 4706.
Noei, S., Zouridis, I. S., Logothetis, N. K., Panzeri, S., & Totah, N. K. (2022). Distinct ensembles in the
noradrenergic locus coeruleus are associated with diverse cortical states. *Proceedings of the National*
Academy of Sciences, 119(18), e2116507119. doi:doi:10.1073/pnas.2116507119
Patrone, L. G., Bicego, K. C., Hartzler, L. K., Putnam, R. W., & Gargaglioni, L. H. (2014). Cardiorespiratory effects
of gap junction blockade in the locus coeruleus in unanesthetized adult rats. *Respir Physiol Neurobiol,*
190, 86-95. doi:10.1016/j.resp.2013.09.001
Peinado, A., Yuste, R., & Katz, L. C. (1993). Gap Junctional Communication and the Development of Local Circuits
in Neocortex. *Cerebral Cortex, 3*(5), 488-498. doi:10.1093/cercor/3.5.488
Rash, J. E., Olson, C. O., Davidson, K. G., Yasumura, T., Kamasawa, N., & Nagy, J. I. (2007). Identification of
connexin36 in gap junctions between neurons in rodent locus coeruleus. *Neuroscience, 147*(4), 938-956.
doi:10.1016/j.neuroscience.2007.04.061

Rukhadze, I., Carballo, N. J., Bandaru, S. S., Malhotra, A., Fuller, P. M., & Fenik, V. B. (2017). Catecholaminergic A1/C1 neurons contribute to the maintenance of upper airway muscle tone but may not participate in NREM sleep-related depression of these muscles. *Respir Physiol Neurobiol, 244*, 41-50. doi:10.1016/j.resp.2017.07.001

Sara, S. J., & Bouret, S. (2012). Orienting and reorienting: the locus coeruleus mediates cognition through arousal. *Neuron, 76*(1), 130-141. doi:10.1016/j.neuron.2012.09.011

Scala, F., Kobak, D., Bernabucci, M., Bernaerts, Y., Cadwell, C. R., Castro, J. R., . . . Tolias, A. S. (2020). Phenotypic variation of transcriptomic cell types in mouse motor cortex. *Nature*. doi:10.1038/s41586-020-2907-3

Scala, F., Kobak, D., Bernabucci, M., Bernaerts, Y., Cadwell, C. R., Castro, J. R., . . . Tolias, A. S. (2021). Phenotypic variation of transcriptomic cell types in mouse motor cortex. *Nature, 598*(7879), 144-150. doi:10.1038/s41586-020-2907-3

Scala, F., Kobak, D., Shan, S., Bernaerts, Y., Laturnus, S., Cadwell, C. R., . . . Tolias, A. S. (2019). Layer 4 of mouse neocortex differs in cell types and circuit organization between sensory areas. *Nat Commun, 10*(1), 4174. doi:10.1038/s41467-019-12058-z

Schoppa, N. E., & Westbrook, G. L. (2002). AMPA autoreceptors drive correlated spiking in olfactory bulb glomeruli. *Nat Neurosci, 5*(11), 1194-1202. doi:10.1038/nn953

Schwarz, L. A., & Luo, L. (2015). Organization of the locus coeruleus-norepinephrine system. *Curr Biol, 25*(21), R1051-R1056. doi:10.1016/j.cub.2015.09.039

Sotelo, C., Llinas, R., & Baker, R. (1974). Structural study of inferior olivary nucleus of the cat: morphological correlates of electrotonic coupling. *J Neurophysiol, 37*(3), 541-559. doi:10.1152/jn.1974.37.3.541

Southwick, S. M., Bremner, J. D., Rasmusson, A., Morgan, C. A., 3rd, Arnsten, A., & Charney, D. S. (1999). Role of norepinephrine in the pathophysiology and treatment of posttraumatic stress disorder. *Biol Psychiatry, 46*(9), 1192-1204. doi:10.1016/s0006-3223(99)00219-x

Su, Z., & Cohen, J. Y. (2022). Two types of locus coeruleus norepinephrine neurons drive reinforcement learning. *bioRxiv*, 2022.2012.2008.519670. doi:10.1101/2022.12.08.519670

Swanson, L. W. (1976). The locus coeruleus: a cytoarchitectonic, Golgi and immunohistochemical study in the albino rat. *Brain Res, 110*(1), 39-56. doi:10.1016/0006-8993(76)90207-9

Tamas, G., Buhl, E. H., Lorincz, A., & Somogyi, P. (2000). Proximally targeted GABAergic synapses and gap junctions synchronize cortical interneurons. *Nat Neurosci, 3*(4), 366-371. doi:10.1038/73936

Tasic, B., Menon, V., Nguyen, T. N., Kim, T. K., Jarsky, T., Yao, Z., . . . Zeng, H. (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat Neurosci, 19*(2), 335-346. doi:10.1038/nn.4216

Ting, J. T., Daigle, T. L., Chen, Q., & Feng, G. (2014). Acute brain slice methods for adult and aging animals: application of targeted patch clamp analysis and optogenetics. *Methods Mol Biol, 1183*, 221-242. doi:10.1007/978-1-4939-1096-0_14

Totah, N. K., Neves, R. M., Panzeri, S., Logothetis, N. K., & Eschenko, O. (2018). The Locus Coeruleus Is a Complex and Differentiated Neuromodulatory System. *Neuron, 99*(5), 1055-1068 e1056. doi:10.1016/j.neuron.2018.07.037

Totah, N. K. B., Logothetis, N. K., & Eschenko, O. (2019). Noradrenergic ensemble-based modulation of cognition over multiple timescales. *Brain Res, 1709*, 50-66. doi:10.1016/j.brainres.2018.12.031

Travagli, R. A., Dunwiddie, T. V., & Williams, J. T. (1995). Opioid inhibition in locus coeruleus. *J Neurophysiol, 74*(2), 518-528.

Trenholm, S., McLaughlin, A. J., Schwab, D. J., Turner, M. H., Smith, R. G., Rieke, F., & Awatramani, G. B. (2014). Nonlinear dendritic integration of electrical and chemical synaptic inputs drives fine-scale correlations. *Nat Neurosci, 17*(12), 1759-1766. doi:10.1038/nn.3851

Trudeau, L. E. (2004). Glutamate co-transmission as an emerging concept in monoamine neuron function. *J Psychiatry Neurosci, 29*(4), 296-310.

Uematsu, A., Tan, B. Z., Ycu, E. A., Cuevas, J. S., Koivumaa, J., Junyent, F., . . . Johansen, J. P. (2017). Modular organization of the brainstem noradrenaline system coordinates opposing learning states. *Nat Neurosci, 20*(11), 1602-1611. doi:10.1038/nn.4642

Van Bockstaele, E. J., Garcia-Hernandez, F., Fox, K., Alvarez, V. A., & Williams, J. T. (2004). Expression of
connexins during development and following manipulation of afferent input in the rat locus coeruleus.
Neurochem Int, 45(2-3), 421-428. doi:10.1016/j.neuint.2003.08.019
Vervaeke, K., Lorincz, A., Gleeson, P., Farinella, M., Nusser, Z., & Silver, R. A. (2010). Rapid desynchronization of
an electrically coupled interneuron network with sparse excitatory synaptic input. *Neuron, 67*(3), 435-
451. doi:10.1016/j.neuron.2010.06.028
Vervaeke, K., Lorincz, A., Nusser, Z., & Silver, R. A. (2012). Gap junctions compensate for sublinear dendritic
integration in an inhibitory network. *Science, 335*(6076), 1624-1628. doi:10.1126/science.1215101
Waterhouse, B. D., Lin, C. S., Burne, R. A., & Woodward, D. J. (1983). The distribution of neocortical projection
neurons in the locus coeruleus. *J Comp Neurol, 217*(4), 418-431. doi:10.1002/cne.902170406
Weinshenker, D. (2018). Long Road to Ruin: Noradrenergic Dysfunction in Neurodegenerative Disease. *Trends
Neurosci, 41*(4), 211-223. doi:10.1016/j.tins.2018.01.010
Williams, J. T., Henderson, G., & North, R. A. (1985). Characterization of alpha 2-adrenoceptors which increase
potassium conductance in rat locus coeruleus neurones. *Neuroscience, 14*(1), 95-101. doi:10.1016/0306-
4522(85)90166-6
Yang, B., Sanches-Padilla, J., Kondapalli, J., Morison, S. L., Delpire, E., Awatramani, R., & Surmeier, D. J. (2021).
Locus coeruleus anchors a trisynaptic circuit controlling fear-induced suppression of feeding. *Neuron*.
doi:10.1016/j.neuron.2020.12.023
Zeisel, A., Munoz-Manchado, A. B., Codeluppi, S., Lonnerberg, P., La Manno, G., Jureus, A., . . . Linnarsson, S.
(2015). Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-
seq. *Science, 347*(6226), 1138-1142. doi:10.1126/science.aaa1934

Figure Supplements
Figure 2-figure supplement 1   (Fig.2S1)

Fig.2S1. Dendrites of LC/NE neurons compared to CA1 hippocampal pyramidal neurons (CA1-PCs) and PV-expressing interneuron (CA1-PV). (**A**) Left: A photomicrograph depicting the morphological recovery of three LC/NE neurons; Right: high-magnification view of a dendritic section (squared on the left with red dash line) of one of the same LC/NE neurons. (**B**) Left: A photomicrograph depicting the morphological recovery of a CA1-PC; Right: high-magnification view of dendritic sections (squared on the left with red dash line) of the same CA1-PC. (C) Left: A photomicrograph depicting the morphological recovery of a CA1-PV; Right: high-magnification view of dendritic sections (squared on the left with red dash line) of the same CA1-PV.

Fig.2S2. **Comparison of axonal and dendritic arbors of two LC/NE morphological cell types**. (**A**) Morphology of multipolar cells (MPs) including all axonal trees remained in slices. The dendrite and soma are shown in gray and the axon is shown in red. (**B**) Morphology of fusiform cells (FFs) including all axonal trees remained in slices. The dendrite and soma are shown in dark blue, and the axon is shown in purple. (**C**) Sholl analysis of length of both axonal and dendritic arbors of MP and FF. Left: Dendrite length (FF vs MP: $F_{(1)}=130.6$; $p<0.0001$ with two-way ANOVA, n=32 for FF and n=36 for MP). Right: axonal length (FF vs MP: $F_{(1)}=47.3$; $p<0.0001$ with two-way ANOVA, n=25 for FF and n=31 for MP).

Figure 2-figure supplement 3 (Fig.2S3)

Fig.2S3. Machine learning to distinguish two LC/NE morphological cell types. (**A**) Classification performance with one feature alone (the shape of the cell body, i.e., the ratio of major to the minor axis of a fitted ellipse). (**B**) Classification performance with another feature alone (the fraction of dendrites originated around the major axis of the fitted ellipse of the soma). (**C**) Classification performance with both features.

1240

1241

1242

1243

1244

1245 **Figure 2-figure supplement 4 (Fig.2S4)**

Fig.2S4. The proportion of FF or MP across all individual recording sets (slices). (**A**) Histogram of the proportion of FF across all slices. (**B**) Histogram of the proportion of MPs across all slices.

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

Figure 2-figure supplement 5 (Fig.2S5-Table)

Table 1: Electrophysiological Properties of MPs and FFs

Cell Type	MP (n=41)	FF (n=19)
AP threshold (mV)	-31.2 ± 0.9	-30.9 ± 1.5
AP amplitude (mV)	69.6 ± 1.6	77.4 ± 1.6**
AP half-width (ms)	2.20 ± 0.07	1.84 ± 0.06**
AP upstroke-to-downstroke ratio	3.29 ± 0.09	3.68 ± 0.17*
Resting membrane potential (mV)	-44.8 ±1.5	-44.9 ± 2.1
Input resistance (MOhm)	710.3 ± 60.0	707.5 ± 52.9
Time constant (ms)	69.0 ± 5.4	75.4 ± 5.9
AHP (mV)	23.1 ± 0.8	25.1 ± 1.6

**$p < 0.01$, *$p < 0.05$

Appendix 1

Model 4: dependent pairwise connectivity model

Since simulations of all previous models with independent probabilities of pairwise connections failed to reproduce the experimental data (i.e. the numbers of the chain-like pattern of two connections were consistently under-estimated by our models, whereas the numbers of the non-chain-like pattern of two connections were consistently over-estimated), the above constraint was relaxed in a proof-of-concept new model. Here, the probability for a neuron to connect with another neuron was not independent of other connections in the recording. Instead of using an overall connectivity probability for all neurons (P: the ratio of the observed number of electrical connections to the total number of potential electrical connections tested across all recordings), we manipulated a neuron's connectivity as a function of other pairwise inter-neuronal connections in a given recording in two different ways. First, if a given neuron already had a pairwise electrical connection with another neuron, then the probability for it to form a pairwise electrical connection with yet another neuron would be scaled up by a factor f_1 (i.e. the new probability = P*f_1,), whereas if the recording slice already had a pairwise connection between two neurons, none of which was the given neuron, then its probability of forming a pairwise connection would be scaled down by a factor f_2 (i.e. the new probability = P/f_2). The simulation was performed for every recording slice of n neurons (where n = 2,3, …,or 8) with the total number of simulated slices equivalent to the number of slices in our recordings, so that the total number of possible pairwise connections across all the recording slices in our model is equal to the total number of said connections in the biology. Simulated numbers of double or 2-connections were calculated as before. We found that for several different values of the scaling-up factor f_1 and scaling-down factor f_2, model estimates came close to experimental data. For instance, for f_1 and f_2 equal to 2.9 and 3.9 respectively, the observed number of the chain-like pattern (n = 16, Fig.6D) was no different from the numbers predicted by the new dependent pairwise connectivity model (n = 16.0, 95% C.I = [6, 28]), and the observed number of the non-chain-like pattern (n = 5, Fig.6D) was indistinguishable from the numbers predicted by the new model (n = 4.8, 95% C.I = [0, 12]). In further simulations, we found that one factor alone (f_1 or f_2) cannot reproduce the experimental findings. At present, we have no evidence for mechanisms that could

potentially underlie either the scaling-up factor f_1 or the scaling-down factor f_2, but this simulation

indicates that the probability for a neuron to connect with another neuron is contingent upon whether it

has existing electrical connections, supporting that electrical connections among LC neurons are not

random, but incline to cluster together in LC. We are now revising the new model by adding spatial and

cell type-specific rules into the model to test what f_1 and f_2 would be for the model to predict our

experimental data.

1338

1339

1340

A

DBH-Cre: Ai9

LC

100 μm   20 μm   200 μm

B

-41mV   LC-NE

-61mV   CA1-PV

-64mV   CA1-PC

10s

20mV

RMP (mV)

Firing rate (Hz)

CA1-PC   CA1-PV   LC-NE

C

LC-NE

CA1-PV

CA1-PC

I_{inj}

100ms

50mV
500pA

D

50mV

6ms

CA1-PV
CA1-PC
LC-NE

tSNE-2

tSNE-1

A

0.1mm

B

C

FF
MP

Dendrite Length (μm)

Axon Length (μm)

A LC/NE

B CA1-PC

C CA1-PV

0.1mm

0.01mm