Cyclin CLB2 mRNA localization and protein synthesis link cell cycle progression to bud growth

Anna Maekiniemi[1], Philipp Savakis[2], Jacky L. Snoep[3,4], Markus Seiler[5], Kelly van Rossum[2], David D. van Niekerk[3], Kathi Zarnack[5], Robert H. Singer[1], Evelina Tutucci[1,2,*]

[1] Cell Biology, Albert Einstein College of Medicine, 1300 Morris Park Avenue, 10461 Bronx, NY, USA

[2] Systems Biology Lab, Amsterdam Institute of Molecular and Life Sciences (AIMMS), Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[3] Department of Biochemistry, University of Stellenbosch, Private Bag X1, Matieland, 7602, South Africa

[4] Department of Molecular Cell Biology, VU University Amsterdam, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands

[5] Buchmann Institute for Molecular Life Sciences (BMLS) & Faculty of Biological Sciences, Goethe University Frankfurt, Max-von-Laue-Str. 15, 60438 Frankfurt, Germany

*Corresponding author: Evelina Tutucci evelina.tutucci@vu.nl

Keywords: mRNA localization, localized translation, smFISH, MS2 system, ZIP code, cell cycle, mitosis, *CLB2*, B-type cyclin, fluorescence microscopy

Figures: 6; Extended Data Figures: 8; Movies: 6; Resources table: 1.

Abstract

Clb2 is a conserved mitotic B-type cyclin, the levels of which are finely controlled to drive

progression through the cell cycle. While it is known that *CLB2* transcription and Clb2 protein

degradation are important for precise control of its expression, it remains unclear whether the

synthesis of Clb2 is also regulated. Furthermore, it is unclear whether and how Clb2 expression

levels respond to cell growth changes and adapt cell cycle progression. To address these

questions, we combined single-cell and single-molecule imaging methods to measure *CLB2*

mRNA and protein expression throughout the *Saccharomyces cerevisiae* cell cycle. We found

that the *CLB2* mRNA was efficiently localized to the yeast bud as soon as this compartment was

formed, while the Clb2 protein accumulated in the mother nucleus. The *CLB2* mRNA localization

in the yeast bud by the She2-3 complex did not control protein localization but rather promoted

CLB2 translation. Moreover, *CLB2* mRNA bud localization and protein synthesis were coupled

and dependent on a single secondary structure -a ZIP code- located in the coding sequence. In

a *CLB2* ZIP code mutant, mRNA localization was impaired and Clb2 protein synthesis decreased,

resulting in changes in cell cycle distribution and increased size of daughter cells at birth. Finally,

while in WT cells the Clb2 protein concentration followed bud growth, this relationship was

impaired in the ZIP code mutant. We propose that *S. cerevisiae* couples the control of *CLB2*

mRNA bud localization and protein synthesis to coordinate cell growth and cell cycle progression.

This mechanism extends our knowledge of *CLB2* expression regulation, and constitutes a novel

function for mRNA localization.

1   Introduction

Over the past decades, RNA imaging technologies have revealed that hundreds of mRNAs

localize to various subcellular compartments, in organisms ranging from bacteria to multicellular

eukaryotes[1,2], suggesting that mRNA trafficking is a conserved and integral part of gene

expression regulation. Current studies suggest that the primary role of mRNA trafficking is to

control asymmetric protein distribution to sustain local functions such as cell migration and

polarity[1]. Even in the single-cell organism *S. cerevisiae*, dozens of mRNAs localize to the

endoplasmic reticulum, mitochondria, and the bud[5]. The best-characterized localized mRNA is

ASH1, which is transported to the yeast bud on actin filaments by the She2-She3 complex and

the type V myosin motor Myo4[6–13]. The RNA binding proteins (RBP) Khd1 and Puf6 bind the *ASH1*

mRNA and inhibit its translation until the bud-localized kinases Yck1 and CK2 phosphorylate Khd1

and Puf6, releasing the inhibition and allowing local translation to occur[14–19]. The Ash1 protein is

subsequently asymmetrically segregated into the daughter nucleus, where it controls mating-type

switching[5,9]. An additional kinase-RBP pair, Cbk1-Ssd1, has been shown to localize to the bud [20]

and tune the translation of specific mRNAs[21–23].

Besides *ASH1*, multiple mRNAs have been shown to interact with the She2-She3-Myo4 complex

and to localize to the bud[24]. Among these mRNAs is *CLB2*, which encodes a conserved B-type

cyclin, interacting with and controlling the substrate specificity of the cyclin-dependent kinase

Cdk1[25–30]. The Clb2 protein contains two nuclear localization signals and two nuclear export

signals[31]. Most of the protein is found in the nucleus and at spindle pole bodies[31–33]. However,

when the protein is overexpressed or in nuclear import mutants, the Clb2 protein accumulates in

the cytoplasm or at the bud neck[31–33]. Clb2-Cdk1 regulates entry and progression through mitosis

in a threshold-dependent manner[27,34–38], by phosphorylating transcriptional and post-

transcriptional regulators[39–41]. This triggers a positive feedback loop leading to the transcription of

the *CLB2* cluster, a set of 35 genes including *CLB2*, expressed during the G2/M phase

transition[26,42–44]. Aberrant Clb2 expression -depletion or over-expression- results in abnormal

mitotic progression and cell size alteration[27,29,45]. To achieve accurate periodic Clb2 expression,

cells combine cell-cycle-dependent mRNA synthesis[46,47], controlled mRNA decay[48], and

proteasome-dependent protein degradation[49,50]. The molecular events controlling *CLB2*

transcription and protein degradation, as well as Clb2 function during cell cycle progression, are

well characterized. However, it remains unclear whether and how cells modulate *CLB2* mRNA

translation in response to changes in cell growth, such as bud growth during *S. cerevisiae* mitosis,

and how it may be coupled to cell cycle progression. Previous mathematical modelling suggested

that *CLB2* mRNA localization and local translation could act as a bud sizer during the G2/M phase

checkpoint [51], but thus far experimental evidence supporting this hypothesis is lacking.

To investigate this unexplored layer of *CLB2* expression regulation, we performed single-cell

measurements of *CLB2* mRNA and protein expression throughout the budding yeast cell cycle in

fixed and living cells. We combined single-molecule RNA fluorescence *in situ* hybridization

(smFISH)[52,53] and immunofluorescence (IF)[54,55] to detect *CLB2* mRNA and its protein product

simultaneously in individual cells. Furthermore, to study dynamic gene expression changes in

living cells, we utilized the MS2 system (MBSV6) optimized to endogenously tag unstable mRNAs

in *S. cerevisiae*[56–59], and tagged the Clb2 protein with GFP. Our work shows that *CLB2* mRNAs

are efficiently localized in the bud by the She2-She3 complex during the G2/M phase, while the

Clb2 protein is found in the mother nucleus. We show that the *CLB2* mRNA has a single ZIP code

located in the coding sequence, which is important for both mRNA localization and Clb2 protein

synthesis. We found that bud localization stimulated *CLB2* mRNA translation, as in a *CLB2* ZIP

code mutant we observed impaired mRNA bud localization and reduced Clb2 protein synthesis,

which were partially rescued in a ZIP code rescue strain. Furthermore, we found that the ZIP code

mutant showed an increase in bud size at birth and a small cell cycle distribution defect by flow

cytometry. Finally, unlike in wild-type (WT) cells, we observed that protein accumulation in the

ZIP code mutant showed reduced correlation with bud growth and thus lost the ability to be a

consistent predictor of size changes occurring in this compartment. Altogether, we propose that

yeast cells have evolved a mechanism that couples the control of *CLB2* mRNA bud localization

and protein synthesis to regulate Clb2 protein levels in response to bud growth. We suggest that,

in coordination with other known mechanisms controlling *CLB2* expression, this mechanism helps

cells monitoring bud growth and fine-tune cell cycle progression.

Results

CLB2 mRNAs localize in the bud from S phase to Mitosis

To quantify *CLB2* mRNA expression throughout the *S. cerevisiae* cell cycle, we combined

smFISH and IF. To monitor cell cycle progression, nuclear localization of the transcription factor

Whi5 was used to classify early G1 phase, while G2 and mitotic cells were identified by staining

tubulin (Tub1) and monitoring microtubules stretching between the mother and the daughter

mitotic spindles (**Fig. 1a**). *CLB2* smFISH revealed that mRNAs are detected from late S phase,

when the bud emerges from the mother cell, until the end of anaphase. Quantification of *CLB2*

mRNAs showed that mRNAs are found in 63% of cells in an unsynchronized population (**Fig. 1b**

and Extended Data Fig. 1a). The expression peak occurred during G2 (average 10.2 ± 5.7

mRNAs/cell) when about 50% of the cells showed an active transcription site (**Fig. 1b,c,** yellow

allows) with on average 2.9 ± 1.5 nascent RNAs per transcription site, comparable to previous

studies[48] (**Extended Data Fig. 1b**). Furthermore, in expressing cells, *CLB2* nascent RNA

expression showed Poissonian transcription kinetics typical of constitutive genes [60], suggesting

that this cell-cycle regulated gene is likely transcribed in a single activation event with fixed

initiation rate. From late S phase until anaphase, we observed that *CLB2* mRNAs localize to the

bud from the first stages of bud formation. Throughout the budded phases, we measured up to

65.6% of mRNAs in the bud, as compared to the distribution of the non-localized mRNA *MDN1,*

where only 17.2% of mRNAs are found in this compartment (**Fig. 1b,d and Extended Data Fig.**

1c,d). *CLB2* mRNA bud localization is independent of the *S. cerevisiae* background since we

observed it both in BY4741, used throughout this study, as well as in the W303 background

(**Extended Data Fig. 1e**).

CLB2 mRNAs efficiently localize in the bud of living S. cerevisiae cells

To investigate *CLB2* mRNA localization dynamics in living cells, we used an MS2 system

optimized for yeast mRNA tagging (MS2 binding sites V6, MBSV6)[56–58]. We inserted 24xMBSV6

in the 3'UTR of the endogenous *CLB2* locus (**Extended Data Fig. 2a**). To confirm that mRNA

tagging with MBSV6 did not alter *CLB2* mRNA expression, unlike with previous MS2

variants[57,61,62], we performed two-color smFISH with probes targeting either the coding sequence

(CDS) or the MBSV6 loops, to compare the expression of the endogenous and the tagged *CLB2*

mRNA. This confirmed that MS2-tagged mRNAs are full-length and correctly localized in the bud

(**Extended Data Fig. 2b**). Furthermore, comparable mRNA levels were observed whether the

mRNA was MS2-tagged, with or without GFP-tagged MS2 coat protein (MCP-GFP), which is used

to detect mRNAs in living cells (**Extended Data Fig. 2c-e**).

To monitor cell cycle progression and bud emergence in living cells, we tagged the bud neck

protein Cdc10 endogenously with tdTomato in the *CLB2*-MS2-tagged strain (**Extended Data Fig.**

3a). We performed time-lapse imaging every 2 minutes and measured *CLB2* mRNA expression

throughout the cell cycle by acquiring z-stacks encompassing the cell volume (**Video 1**). To

reduce perturbations in gene expression due to synchronization protocols[63], we quantified *CLB2*

mRNA expression in unsynchronized cells, using the bud neck marker expression to compare

cells. This revealed that up to 62.9% of *CLB2* mRNAs localized in the bud (**Fig. 1e,f**), in agreement

with the smFISH quantifications (**Fig. 1d**). Furthermore, mRNAs were degraded before the end

of mitosis with a half-life of 3.8 ± 1.4 min, consistent with previous measurements[48], suggesting

that the MS2 system does not affect *CLB2* mRNA stability (**Extended Data Fig. 3b**). Interestingly,

imaging of mother-daughter pairs for more than one cell cycle showed that the daughter cell

initiated *CLB2* mRNA expression about 20 minutes after the mother (**Fig. 1g**). This observation

is in line with previous evidence showing that *S. cerevisiae* daughter cells are born significantly

smaller than mothers and that cell size control occurring during G1 regulates the entry into the

next cell cycle[64–66].

High frame-rate imaging every 100 ms revealed that, as the bud grows, the number of *CLB2*

mRNAs localized in the bud rapidly increases (**Extended Data Fig. 3c and Video 2**). The high

concentration of mRNAs in the bud and their rapid movement in and outside of the focal plane

did not allow tracking single mRNA molecules to investigate *CLB2* mRNAs bud localization

dynamics. We therefore used mathematical modelling to predict the behavior of *CLB2* mRNA in

the bud of an average G2 cell. The parameters included the bud and mother volumes based on

our measurements, mRNA counts and decay rates based on smFISH and live imaging

experiments and an apparent mRNA diffusion coefficient based on previous measurements

performed in eukaryotic cells[67] (**Methods**). A fast coefficient of 0.4 $\mu m^2/s$ was previously

measured for non-translating mRNAs, and a slower coefficient of 0.1 $\mu m^2/s$ was measured for

translating mRNAs[67]. Interestingly, our model suggested that if we assumed either a slow or a

fast apparent mRNA diffusion coefficient of 0.1 $\mu m^2/s$ or 0.4 $\mu m^2/s$, respectively, we would not

obtain the expected enrichment of the *CLB2* mRNA in the bud (**Extended Data Fig. 3d,e**). To

predict the accumulation of about 65% of the mRNA in the bud observed during the G2 phase,

the presence of a high-affinity anchoring factor promoting *CLB2* mRNA segregation in the bud

had to be included in the simulation (**Extended Data Fig. 3f**). Altogether, these results showed

that *CLB2* mRNAs were efficiently transported to the bud in a cell cycle dependent manner and

the simulation implied that the mRNA might be anchored via a yet uncharacterized mechanism.

The CLB2 mRNA has a single She2 binding site in the coding sequence and is transported to the

bud via the She2-She3 complex

To elucidate the function of *CLB2* mRNA localization, we first characterized the mechanism of

CLB2 mRNA transport. Previous work showed that *CLB2* mRNA is associated with the She2-

She3-Myo4 complex and that localization of *CLB2* to the bud likely requires She2[24]. To expand

this observation and quantify this phenotype, we performed smFISH-IF throughout the cell cycle

for the *CLB2* mRNA in *SHE2* or *SHE3* gene deletion strains to test whether the She2-She3

complex, required for *ASH1* mRNA transport[6–13], was also involved in *CLB2* mRNA localization.

This revealed that in *Δshe2* and *Δshe3* strains, localization was strongly impaired (**Fig. 2a and**

Extended Data Fig. 4a) and that during mitosis only up to 24.5% and 23.6% of *CLB2* mRNAs

were found in the bud of the *Δshe2* and *Δshe3* strains, respectively (**Extended Data Fig. 4b,c**).

As the She2-She3 complex was required for *CLB2* mRNA localization, we hypothesized that the

CLB2 mRNA might possess a ZIP code akin to the *ASH1* ZIP code. Previous work defined the

sequence and structure of the (E3) *ASH1* mRNA ZIP code bound by She2[10,68–70]. Based on

sequence and structure similarity, a pattern search was performed to predict occurrences within

the *CLB2* mRNA (**Methods**). We identified one high-confidence She2-binding site in the CDS at

position 1111-1145 (**Fig. 2b,c**). To test the predicted site, we generated a *CLB2* synonymized

mutant whereby the CDS was mutagenized at nine bases to destroy the ZIP code structure, while

preserving the protein sequence and the codon usage index (*ZIP mut*, **Fig. 2d and Extended**

Data Fig. 4d,e). A pattern search confirmed that the ZIP code was destroyed upon

synonymization. smFISH revealed that, in the *CLB2* ZIP code mutant, the *CLB2* mRNA was no

longer bud localized (**Fig. 2e,f**), demonstrating that the identified ZIP code is sufficient to control

bud mRNA localization, possibly by recruiting the She proteins. Furthermore, the loss of *CLB2*

mRNA localization was not due to the specific mutations introduced, as alternative ZIP code

mutants also showed *CLB2* mRNA mis-localization (**Extended Data Fig. 4d,f**). To measure *CLB2*

mRNA localization, we quantified the mRNA peripheral distribution index (PDI) in budded cells

using the RNA Distribution Index Calculator[71] (**Methods**). The PDI measures the location of the

mRNA relative to the nucleus and allows for the comparison of the localization of distinct mRNA

species. An index value equals 1 for diffusely distributed mRNAs or >1 if the mRNA has a

polarized pattern, and <1 if the mRNA is distributed closer to the nucleus[71] (**Fig. 2g**). This analysis

revealed a PDI of 1.9 ± 0.42 for the *CLB2* mRNA, similar to the index value of the control mRNA

ASH1 (PDI = 2.2 ± 0.43, **Fig. 2h**). The PDI value was significantly reduced for *CLB2* in the *Δshe2*

(PDI = 0.5 ± 0.2), *Δshe3* (PDI = 0.4 ± 0.13), and *CLB2* ZIP code mutant strain (PDI = 0.5 ± 0.16)

(ANOVA statistical test: $F_{(4, 185)}$ = 15.74, $p < 0.0001$), with PDI values similar to the non-localized

mRNA *MDN1* (PDI = 0.6 ± 0.18, **Fig. 2h and Extended Data Fig. 1c,d**). Thus, the She2-3

complex is required to transport *CLB2* mRNAs to the bud via a conserved ZIP code sequence.

Lack of CLB2 mRNA localization affects Clb2 protein expression but not protein localization

To elucidate whether *CLB2* mRNA localization influences its expression, we measured *CLB2*

mRNA and protein levels in the localization mutants. Using smFISH, we found comparable mature

or nascent RNAs counts in the *Δshe2* (5.3 ±5.1 mRNA/cell and 2.7 ±1.1 nascent

RNA/transcription site(TS)) or *Δshe3* mutants (5.2 ±6.2 mRNA/cell and 2.7 ±1.3 nascent RNA/TS)

compared to WT cells (5.4 ±5.1 mRNA/cell and 2.6 ±1.3 nascent RNA/TS). While we observed a

small but significant increase in the *CLB2* ZIP code mutant strain (6.9 ±6.3 mRNA/cell and 3.4

±1.3 nascent RNA/TS) compared to WT cells (**Fig. 3a,b**). Furthermore, a western blot of the

endogenously myc-tagged Clb2 protein showed that the protein expression in Δ*she2* or Δ*she3*

and in the *CLB2* ZIP code mutants was significantly reduced compared to WT cells (**Fig. 3c,d**).

Interestingly, all ZIP code mutants showed reduced Clb2 protein expression regardless of the ZIP

code mutation variant, indicating that this region of the *CLB2* mRNA is important to control mRNA

localization and protein expression (**Extended Data Fig. 4e and Extended Data Fig. 5a,b**). In

view of the similarity in the mRNA localization and protein expression phenotypes of the three ZIP

mutant strains, we continued the characterization of ZIP code mutant 1 (ZIP[1]), henceforth called

'ZIP-mutant'. To test whether the decrease in protein expression was due to a change in protein

degradation, we performed stability assays by treating WT and localization mutants with the

translation inhibitor cycloheximide and measured the protein abundance over time (**Fig. 3e,f and**

Extended Data Fig. 5c,d). No significant difference was observed in the stability of the

localization mutants compared to WT cells, suggesting that *CLB2* mRNA localization, rather than

mRNA or protein stability, regulated Clb2 protein synthesis.

Finally, we investigated whether mRNA localization affected Clb2 protein localization. To this end,

CLB2 was endogenously tagged with yeast-optimized GFP (yeGFP) in WT and *CLB2* mRNA

localization mutants. We observed that Clb2 is predominantly found in the nucleus in all tested

strains (**Fig. 3g**), as previously reported for WT cells[31–33]. Interestingly, Clb2 was observed in the

mother nucleus already during the G2 phase (**Fig. 3g,** top panels), when the mRNAs were already

localized to the bud (**Fig. 1b,e and 2a**), showing an uncoupling between *CLB2* mRNA and protein

localization. Altogether, these results suggested that *CLB2* mRNA localization did not segregate

the Clb2 protein in the daughter cell, unlike Ash1[5,9,72], but rather influenced protein levels.

CLB2 mRNA and protein co-localization suggest preferential translation in the bud

To simultaneously measure *CLB2* mRNA and protein expression in single cells, we generated a

strain where 25 myc tags were inserted at the N-terminus of the endogenous *CLB2* gene (**Fig.**

4a). This N-terminal tagging and amplification strategy increased the Clb2 protein signal without

affecting growth (**Extended Data Fig. 6a,b**). Next, we combined smFISH and IF to simultaneously

detect *CLB2* mRNAs and proteins in fixed cells. Moreover, we used IF against tubulin to score

the cell cycle phases. This approach showed that the bulk of Clb2 proteins accumulated in the

mother (M) nucleus from G2 to mitosis (**Fig. 4b,c**), at the time when the mRNA was preferentially

found in the bud (B) (**Fig. 4b,d**). This indicated that Clb2 proteins were efficiently localized in the

mother nucleus, as observed by live imaging (**Fig. 3g**) and as shown previously[31,33]. In addition,

Clb2 protein foci were also found in the bud in close proximity of *CLB2* mRNAs from G2 to mitosis,

suggesting that these foci may represent sites of mRNA translation (**Fig. 4b**, yellow arrowheads).

Quantification of co-localized single mRNAs and protein foci (within 250 nm, the resolution of our

system) revealed that in WT cells, more co-localized mRNA-protein foci were found in the bud

than in the mother cell, while in the localization mutant *Δshe2*, mRNA-protein foci were

preferentially found in the mother cell where the bulk of mRNAs was distributed (**Fig. 4e,f**).

Furthermore, we found a significant reduction of the percentage of bud-localized mRNAs co-

localized with protein foci in *Δshe2* cells compared to WT cells (**Fig. 4g**, non-parametric Mann

Whitney test, P<0.0001), indicating that in localization mutants *CLB2* mRNA translation efficiency

may be reduced. It is interesting to note that even in WT cells, only about 25% of the bud-localized

mRNAs were found in close proximity to protein foci, suggesting that *CLB2* mRNAs are translated

unfrequently, as previously reported[73]. Altogether, these data suggest that in WT cells, *CLB2*

mRNAs were preferentially translated in the bud, where the mRNA was actively transported.

Furthermore, in *Δshe2* cells we observed an increase of protein foci in the mother cell relative to

the bud (**Fig. 4f**), despite protein levels being decreased under these conditions (**Fig. 3c,d**),

indicating that translation in the mother cell may be less efficient, resulting in reduced Clb2 protein

levels in the localization mutants.

Clb2 protein expression was not affected by the translation regulators Puf6, Ssd1, and Khd1

To investigate if *CLB2* translation efficiency was higher in the bud as a result of translation

repression prior to localization, we tested if the deletion of the RNA binding proteins Puf6[14,15,19],

Khd1[16–18] and Ssd1[21–23], previously shown to inhibit translation of bud-localized mRNAs,

influenced Clb2 protein levels. A western blot of the endogenously myc-tagged Clb2 protein in

Δssd1, Δkhd1, and *Δpuf6* strains (**Extended Data Fig. 6c,d,e**) did not reveal a significant increase

in Clb2 protein levels when comparing the mutant strains to WT (ANOVA statistical test: $F(3, 8) =$

0.7677, $p = 0.3$), which we would have expected if these proteins were inhibiting *CLB2* mRNA

translation. Additionally, because Khd1 and Puf6 were shown to be important for *ASH1* mRNA

localization[15,16], we tested *CLB2* mRNA localization in the mutants. smFISH for the *CLB2* mRNA

in the *Δssd1, Δkhd1*, and *Δpuf6* strains showed that the mRNA was still localized in the bud at a

similar level as WT cells. Altogether, these results indicated that the factors that control *ASH1*

mRNA expression were not required to control *CLB2* mRNA translation. While other yet

unidentified factors may exist, our data suggested the possibility that *CLB2* mRNA might be

translated also outside of the bud, albeit at a reduced rate.

CLB2 mRNA ZIP-mutant displayed increased cell size at birth

The results so far showed that absence of *CLB2* mRNA bud localization correlated with reduced

protein expression, likely due to decreased mRNA translation. This suggested that *CLB2* mRNA

localization may act as a cellular signal reporting on bud growth to the mother nucleus. To test

whether the lack of *CLB2* mRNA localization and the decrease in Clb2 protein levels observed in

the mutants affected cell size and cell cycle progression, we performed time-lapse fluorescence

microscopy. In all strains, the Clb2 protein was tagged with yeGFP while Cdc10 was fused to

mCherry to label the bud neck and monitor division time and mother-daughter size relationships.

Live imaging revealed that mother cells of the ZIP-mutant strain were larger at budding and at the

division time points (**Fig. 5a-d, Videos 3-6**). Furthermore, daughter cells of the ZIP-mutant strain

were approximately 10.9% larger at birth than WT daughters (**Fig. 5b,e**), which resulted in a

significant increase in combined (bud + mother) size (**Fig. 5f**). This did not hold for the *Δshe*

mutant strains, where there was a slight reduction of mother size at both budding and division in

Δshe2, while *Δshe3* was indistinguishable from WT (**Extended Data Fig. 7a-f)**. The ZIP-mutant

strain added more material during the budded phase than both WT and the *Δshe* mutants (**Fig.**

5g and Extended Data Fig. 7g). This effect remained after correcting for mother size (**Extended**

Data Fig. 7h) indicating a longer duration of the budded phase in the ZIP-mutant. While the

median values for the duration of the budded phase in the ZIP-mutant strain and the WT differed

by 5 min, there was significant overlap in the distributions (**Fig. 5h and Extended Data Fig. 7i**)

and this effect was overall not significant (p=0.29, one-way ANOVA). Interestingly, there was a

small but significant decrease in budded time duration for *Δshe2* (**Extended Data Fig. 7j**). Finally,

to investigate whether nuclear Clb2 levels could report bud size we followed individual cells in

time. Fluorescence levels peaked about 20 min before division in all strains (**Fig. 5i and Extended**

Data Fig. 7k). In accordance with reduced Clb2 protein levels (**Fig. 3**), the ZIP-mutant strain

exhibited overall lower peak fluorescence intensity than WT. Additionally, in WT cells the signal

peaked more sharply (**Fig. 5i**). During the buildup to the peak (between 75 to 50 min before

division) the ZIP-mutant strain exhibited marginally higher fluorescence (**Fig. 5i**) but cumulative

fluorescence over the cell cycle indicated that neither a slight increase in baseline nor more time

spent at intermediate Clb2 levels compensated for the reduction in translation efficiency observed

in the mutant strains (**Fig. 5j and Extended Data Fig. 7l**). To measure if localized translation of

CLB2 mRNA could report bud size, we correlated bud size against nuclear fluorescence of the

corresponding mother cell. While nuclear fluorescence tightly predicted bud size in WT and the

Δshe mutants' (**Fig. 5k and Extended Data Fig. 7m**) this effect was significantly more disperse

in the ZIP-mutant strain, indicating a loss of communication between the two compartments.

Altogether the data suggests that control of *CLB2* mRNA translation in the bud compartment

reports bud size (see Discussion).

Adding the CLB2 WT ZIP code to the 3'UTR of the ZIP mutant promotes translation and rescues

cell cycle progression.

We have shown that the ZIP code sequence located in the coding region of the *CLB2* mRNA is

necessary for mRNA localization to the bud via the She2-3 complex. To test if the mere presence

of the ZIP code in the mRNA is enough for efficient localization we reintroduced the ZIP code into

the 3'UTR of the ZIP-mutant strain (**Fig. 6a**). smFISH uncovered that adding the WT ZIP code to

the 3'UTR resulted in a complete rescue of *CLB2* mRNA localization in the bud, comparable to

WT levels (**Fig. 6b**), demonstrating that the WT ZIP code was both necessary and sufficient for

localizing the mRNA. However, Clb2 protein levels were rescued only partially (**Fig. 6c,d**),

suggesting on one hand that the bud promoted Clb2 translation, but also that the ZIP code

structure in the *CLB2* coding sequence increased *CLB2* mRNA translation via a yet unknown

mechanism. Finally, we investigated the growth and cell cycle distribution of the localization

mutants and the ZIP code rescue strain. To measure the distribution of the three major phases of

the cell cycle (G1, S, G2/M), DNA was labelled with propidium iodide and measured by flow

cytometry, followed by Gaussian mixed modelling to estimate the three cell cycle subpopulations.

This revealed that, while growth rates of the localization mutants were not significantly affected

(**Extended Data Fig. 8a**), we observed a modest but significant (two-way Anova, Dunnett's

multiple comparison test) increase in the G2/M fraction in the ZIP-mutant but not in the *Δshe*

mutants nor the ZIP-mutant rescue strain (**Fig. 6e,f**). Altogether, our experiments point to a dual

function of the *CLB2* mRNA ZIP code motif, promoting both efficient bud localization and

translation. For the latter function to take effect, the ZIP motif needs to be located in the coding

sequence. Consistent with this observation, in the *Δshe* mutants where the ZIP code is still intact,

higher levels of Clb2 protein were observed. In the ZIP-mutant, where both the localization is

impaired and the ZIP code is absent, Clb2 protein levels were lowest in bulk populations (**Fig.

3c,d**) and in individuals across the cell cycle (**Fig. 5i**). We also observed an increased G2/M

fraction in this mutant, suggesting that the duration of these stages is increased at the expense

of G1 (**Fig. 6e,f**). Altogether, the higher G2/M fraction (**Fig. 6e,f**) and the larger size at budding

and division (**Fig. 5b-g**) indicated that control of *CLB2* mRNA localization and translation play a

role in regulating Clb2 protein levels, which in turn report translational maturity as well as size of

the bud compartment, providing a mechanism to coordinate growth and cell cycle progression

during the G2/M phase transition.

Discussion

In this study, we investigated the complete lifecycle of the *CLB2* gene by combining single-cell

and single-molecule imaging methods to measure *CLB2* mRNA and protein expression

throughout the *S. cerevisiae* cell division cycle.

CLB2 mRNA localization: same transport as the ASH1 mRNA but distinct regulation

We experimentally demonstrated that the She2-She3 complex was required to transport the *CLB2*

mRNA to the bud as soon as this compartment was formed (**Fig. 1 and 2**), and, based on

mathematical modeling (**Extended Data Fig. 3**) proposed that active mRNA transport needs to

be combined with an anchoring mechanism to explain the highly efficient (>60%) transcript

localization, although anchoring factors have yet to be identified. Our observations are consistent

with previous work showing that the *CLB2* mRNA can be pulled down using the She2, She3 or

Myo4 proteins as bait, as well as with imaging data suggesting *CLB2* mRNA localization in the

bud[24]. The She2-She3 complex localizes the *ASH1* mRNA to the bud, with the She2-mRNA

interaction mediated via the E3 *ASH1* ZIP code located in the *ASH1* coding region[6–13]. We

identified an element in the coding region of *CLB2* whose secondary structure resembles that of

the *ASH1* ZIP code and demonstrated that this element is both necessary and sufficient for

transport into the bud compartment, indicating that it constitutes a bona fide She2 binding site. In

ASH1 the ZIP code motif promotes localization irrespective of its position in the

transcript[6,9,69,70,74,75]. We could confirm this for *CLB2* where moving the ZIP motif to the 3' UTR

resulted in a localization pattern indistinguishable from WT (**Fig. 6**). While translation of *ASH1*

mRNA is tightly regulated by Puf1 and Khd1 we could find no such inhibition of translation for

CLB2 through either of these factors nor through the bud-localized translation regulator Ssd1

(**Extended Data Fig. 6)**, although we cannot exclude regulation via yet unidentified elements. As

Ash1 protein activity needs to be confined to the bud where it controls mating type switching, the

reverse appears to be true for the Clb2 protein, which accumulates in the nucleus of the mother

cell. Thus, while *ASH1* and *CLB2* localization appears to be controlled by the same cellular

machinery, their translational regulation is different.

The CLB2 ZIP code is important for both mRNA localization and protein synthesis control

Alteration of the secondary structure of the ZIP motif through synonymized mutation abolished

localization and led to reduced Clb2 protein levels. Moving the ZIP motif from the coding sequence

to the 3'UTR rescued *CLB2* localization but the protein concentration did not recover to WT levels.

In the *Δshe* mutants, localization was abolished, but protein levels did not decrease to the levels

of the ZIP mutant strain. This suggested that an intact ZIP structure located within the coding

sequence of *CLB2* is required for optimal translation. We therefore propose that the *CLB2* ZIP

motif fulfills a dual role – localization to the bud via She2-She3-Myo4 as well as promotion of

translation through as yet unidentified factors.

While our data points to a role of *CLB2* mRNA localization to control its translation and not Clb2

protein nuclear localization (**Fig. 3**), we cannot exclude that a transient localization of Clb2 in the

bud may have spatially-defined functions that contribute to the observed cell cycle phenotypes.

Previous work showed that Clb2 can be found also at the bud neck, when overexpressed[31,32] or

when Clb2 nuclear import is blocked[33] and that forcing the Clb2 protein in the nucleus can lead to

detrimental effects on nuclear fission[76]. Future studies may uncover additional roles for Clb2 in

the bud compartment and at the neck.

Lack of CLB2 mRNA localization and reduced protein expression impact coupling between growth

and cell cycle

Our data showed that ZIP-mutant cells are larger at birth and at budding (**Fig. 5**). While *S.*

cerevisiae mainly exerts size control in G1[65,77], experimental observations[78] and mathematical

models[79] suggested a second size control mechanism in G2. In *Schizosaccharomyces pombe*

cell size control is mainly realized in G2, likely through the Clb2 homologue Cdc13[35]. A

mechanism in which localization of the *CLB2* transcript functions as a reporter of bud

compartment maturity was explored previously through mathematical models in which translation

efficiency of *CLB2* mRNA is proportional to bud size volume[51]. In these models, G2/M duration is

more variable in the absence of mRNA localization. While our data show a significant increase in

cell size owing to a greater amount of material added to the bud compartment during G2 (**Fig. 5**

and Extended Data Fig. 7), the effect of mislocalized *CLB2* transcript on the duration of the

budded phase were small (**Fig.5 and 6**).

These overall small effects on budded phase duration are likely due to the fact that in the ZIP

code mutant about 40-50% of the Clb2 protein is still expressed. This might be at the lower edge

of the threshold that suffices to drive mitotic entry and progression[27,34–38] which would help to put

the absence of a clear cell cycle phenotype of the *Δshe* mutants into perspective. In these

mutants, protein levels were reduced, yet there is still significant overlap with WT cell levels

(**Extended Data Fig. 7**). Regressing bud size on nuclear Clb2 concentration yielded significantly

more disperse predictions in the ZIP-mutant strain than in WT or in the *Δshe* mutants. This

suggests that *CLB2* mRNA localization and bud translation may be part of a more complex system

coupling bud size control and cell division by combining yet unidentified sizers and timing

mechanisms[80].

Altogether, our data suggest that by coupling *CLB2* mRNA bud localization and protein synthesis,

cells coordinate cell growth with cell cycle progression by sensing the bud translation capacity via

CLB2 mRNA transport and local translation. We propose that by shuttling back to the mother

nucleus, Clb2 protein levels may signal to the mother cell when the bud is ready for mitosis,

establishing a biochemical-based communication between distinct subcellular compartments.

Acknowledgments

We thank X. Meng for help with cloning; J. Wiemerink for help with flow cytometry, A. Gerber for

discussion and critical reading of the manuscript, W. Li, C. Eliscovich and S. Das for discussion.

Funding

This work was supported by NIH grant GM57071 to R.H.S. E.T. was supported by Swiss National

Science Foundation Fellowships P2GEP3_155692 and P300PA_164717, as well as by the Vrije

Universiteit Amsterdam. We acknowledge financial support from the Department of Science and

Technology/National Research Foundation in South Africa, (grant NRF-SARCHI-82813 to J.L.S.)

and for grant 116298 (to D.D.v.N.). K.Z. was supported by the Deutsche Forschungsgemeinschaft

(DFG) via the Research Unit FOR2333 (ZA 881/3-1).

Author contributions

4 Conceptualization, E.T.; Methodology, E.T., A.M., P.S., J.L.S., D.D.N. and K.Z.; Formal Analysis,

5 E.T., A.M., J.L.S., M.S., D.D.N. and P.S.; Investigation, E.T., A.M., P.S. and K.R.; Writing –

6 Original Draft, A.M., E.T.; Writing –Review & Editing, E.T., A.M., R.H.S., P.S., K.Z, and J.L.S.;

7 Funding Acquisition, E.T., J.L.S., K.Z. and R.H.S.; Supervision, E.T., and R.H.S.

Declaration of interests

9 The material in this manuscript (MS2 system V6) is the subject of a patent application to the US

10 Patent and Trademark Office (no. 18242217). It has not been licensed to any corporation, and

11 the authors (R.H.S., E.T.) are the inventors together with Maria Vera Ugalde (McGill University).

Data Availability

13 The raw data plotted in this manuscript can be found in the Zenodo repository DOI:

14 10.5281/zenodo.11636335. Raw images analyzed in this study can be requested to the lead

15 author. In addition, they can find a Resource table (Excel file), where we provided additional

16 information about the reagents (i.e. plasmids, yeast strains, primers, smFISH probes) used in this

17 study.

Code availability

The code used to process flow cytometry, plate reader and Clb2 protein expression

measurements by time-lapse microscopy can be found in the Zenodo repository DOI:

10.5281/zenodo.11636335.

Methods

Yeast strains construction

All strains described in this work were derived from the *S. cerevisiae* background BY4741 (MATa;

his3Δ1; leu2Δ0; met15Δ0; ura3Δ0). Strains are listed in the Resource Table. Yeast strains (i.e.

deletions and taggings) were constructed as detailed in[58]. *CLB2* was tagged with 24xMBSV6 in

the 3' UTR right after the STOP codon. The MBSV6 insert, followed by the kanamycin resistance

gene flanked by LoxP sequences, was PCR amplified from plasmid pET264 (see plasmids in

Resource Table) with primers OET017 and OET254, each containing about 70 nucleotides of

homology sequence for the *CLB2* gene (see Resource Table). The Kanamycin resistance gene

was then removed by expressing the CRE recombinase under the control of the *GAL1* promoter

(Resource Table, plasmids). Obtained strains were tested by PCR and sequencing as previously

described[58].

Plasmids construction

The synonymized *CLB2* ZIP code plasmids were generated by Gibson cloning using the NEB

Gibson Assembly Cloning kit (#E5510) using oligos and plasmids as indicated in the resource

tables. The plasmids were cut using AatII and EcoRI-HF restriction enzymes and the modified

CLB2 inserts were transformed in BY4741 for homologous recombination. The ZIP-code rescue

plasmids were generated by inserting the predicted WT She2 binding site sequence

(AGCAGATGACTACGATATACAGTCTCGAACTCTTGCC) at the ClaI restriction site 43

nucleotides downstream of the *CLB2* stop codon.

Yeast cell cultures

Agar plates 2% (w/v) made with yeast extract peptone dextrose (YEPD) or drop-out medium

lacking leucine (LEU-) with 2% glucose and 6.7 g/L Yeast Nitrogen Base medium (YNB), were

used to grow strains from the -80°C stock or for transformations. Yeasts were grown in 6.7 g/L

Yeast Nitrogen Base medium (YNB) with 2% glucose and the appropriate amino acids to

complement the strains' auxotrophies. All experiments were performed growing cells in Synthetic

Complete (SC), except the experiments in Figure 1e-g and Extended Data Fig. 3b,c, where cells

were grown in Drop-Out media lacking leucine to maintain the expression of the MCP-2xyeGFP

plasmid (pET296). Cells were grown at the indicated temperature using constant shaking at 210

rpm. For smFISH, live imaging and flow cytometry, the details of the cell cultures are described

below.

smFISH probes design

CLB2 probes were designed using the Stellaris™ Probe Designer by LGC Biosearch

Technologies and purchased from Biosearch Technologies. *ASH1, MDN1* and *MBSV6* probes

design was previously described in[57,58]. Probes sequence and fluorophores are provided in the

Resource Table.

Single molecule fluorescence in situ hybridization (smFISH)

Single-molecule FISH (smFISH) was performed as follows. Yeast strains were grown overnight

at 26°C in synthetic medium with 2% glucose and containing the appropriate amino acids to

complement the strain auxotrophies. In the morning, cells were diluted to OD_{600} 0.1 and allowed

to grow until OD_{600} 0.3-0.4. Cells were fixed by adding paraformaldehyde (32% solution, EM

grade; Electron Microscopy Science #15714) to a final concentration of 4% and gently shaken at

room temperature (RT) for 45 minutes. Cells were then washed three times with buffer B (1.2 M

sorbitol and 100 mM potassium phosphate buffer pH 7.5) and resuspended in 500 µL of

spheroplast buffer (buffer B containing 20 mM VRC (Ribonucleoside–vanadyl complex NEB

#S1402S), and 25 U of Lyticase enzyme (Sigma #L2524) per OD of cells (~10^7 cells) for about 7-

8 minutes at 30°C. Digested cells were washed once with buffer B and resuspended in 1 mL of

buffer B. 150 µL of cells were seeded on 18 mm poly-L-lysine treated coverslips and incubated

at 4°C for 30 minutes. Coverslips were washed once with buffer B, gently covered with ice-cold

70% ethanol and stored at -20°C. For hybridization, coverslips were rehydrated by adding 2xSSC

at RT twice for 5 minutes. Coverslips were pre-hybridized with a mix containing 10% formamide

(ACROS organics #205821000)/2xSSC, at RT for 30 minutes. For each coverslip, the probe mix

(to obtain a final concentration in the hybridization mix of 125 nM) was added to 5 µL of 10 mg/mL

E. coli tRNA/ ssDNA (1:1) mix and dried with a speed-vac. The dried mix was resuspended in 25

µL of hybridization mix (10% formamide, 2×SSC, 1 mg/ml BSA, 10 mM VRC, 5 mM $NaHPO_4$ pH

7.5) and heated at 95°C for 3 minutes. Cells were then hybridized at 37°C for 3 hours in the dark.

Upon hybridization, coverslips were washed twice with pre-hybridization mix for 30 minutes at

37°C, once with 0.1% Triton X-100 in 2xSSC for 10 minutes at RT, once with 1xSSC for 10

minutes at RT and once with PBS 1x for 10 minutes at RT. Coverslips were quickly dipped in

100% ethanol and let dry at RT covered form light. Finally, coverslips were mounted on glass

slides using ProLong Gold antifade (4′,6-diamidino-2-phenylindole) DAPI to counterstain the

nuclei (Thermofisher, #P36935).

smFISH-IF

smFISH-IF was performed as previously described in[54,55]. In brief, smFISH-IF was performed in

a similar way as smFISH, described above. After the last 1xPBS wash of the smFISH, IF was

performed on the same coverslips. The smFISH was fixed in 4% PFA in PBS for 10 minutes at

RT and then washed for 5 min at RT with 1x PBS. The coverslips were blocked with 1xPBS, 0.1%

RNAse-free Bovine Serum Albumin for 30 minutes at RT before being incubated with primary

antibodies (Thermofisher, mouse anti-tubulin, 1:1000; Sigma mouse monoclonal anti-myc clone

9E10, 1:1000, Covance, mouse monoclonal anti-HA, 1:1000) in 1xPBS, 0.1% RNAse-free Bovine

Serum Albumin for 45 minutes. After being washed with 1xPBS for 5 minutes at RT, the coverslips

were incubated with the secondary antibody (goat anti-mouse Alexa 647 1:1500, or goat anti-

mouse Alexa 488 1:1500) in 1xPBS, 0.1% RNAse-free Bovine Serum Albumin for 45 minutes at

RT. Next, the coverslips were washed with 1x PBS three times for 5 minutes to remove excess

antibody. Coverslips were dehydrated by dipping them into 100% ethanol and let dry before

mounting onto glass slides using ProLong Gold antifade solution with DAPI.

smFISH and smFISH-IF image acquisition and analysis

Images were acquired using an Olympus BX63 wide-field epi-fluorescence microscope with a

100X/1.35NA UPlanApo objective. Samples were visualized using an X-Cite 120 PC lamp (EXFO)

and the ORCA-R2 Digital CCD camera (Hamamatsu). Image pixel size: XY, 64.5 nm. Metamorph

software (Molecular Devices) was used for acquisition. Z-sections were acquired at 200 nm

intervals over an optical range of 8 μm. FISH images were analyzed using FISHQUANT[81]. Briefly,

after background subtraction, the FISH spots in the cytoplasm were fit to a three-dimensional (3D)

Gaussian to determine the coordinates of the mRNAs. The intensity and width of the 3D Gaussian

were filtered to exclude nonspecific signal. The average intensity of all the mRNAs was used to

determine the intensity of each transcription site.

Quantification of peripheral distribution index

The peripheral distribution index (PDI) was quantified as described in[71]. Briefly, the Matlab-based

software RDI (RNA dispersion index) calculator was used to calculate the peripheral distribution

index for each cell by identifying cellular RNAs and describing their distribution in relation to the

nucleus. Prior to analysis with the RDI calculator, the RNA channel was processed using a 3D

Laplacian of Gaussian filter of radius=5 and standard deviation=1. The cell and nucleus channels

were processed using the brightness/contrast function in ImageJ to enhance the contrast between

the object and the background, as advised in[71].

Co-localization analysis

For the co-localization of *CLB2* mRNA and protein foci in Figure 4, the FISH-quant data for the

individual molecules were used as x, y, z coordinates and euclidean distances for all protein -

mRNA molecule combinations were calculated in the mother and daughter cells. Protein and

mRNA molecules closer than 250 nm were considered to be in a translation complex. Multiple

protein molecules can be within 250 nm of a single mRNA molecule, and this would still be

considered a single translation complex.

PDE solution for mRNA diffusion

We use a modified diffusion equation at steady state to model the mRNA movement in terms of

diffusion of a concentration *c(x,y,z)* in 3 spatial dimensions, and include binding to ribosomes

(uniformly spread) leading to the formation of complexes *b(x,y,z)*:

7
$$0 = D \cdot \nabla^2 c(x,y,z) - k_d \cdot c(x,y,z) + k_p - k_{\text{on}} \cdot c(x,y,z) + k_{\text{off}} \cdot b(x,y,z)$$

8
$$0 = k_{\text{on}} \cdot c(x,y,z) - k_{\text{off}} \cdot b(x,y,z) - k_d \cdot b(x,y,z)$$

With decay constant k_d = Ln(2)/$t_{0.5}$ = Ln(2)/240 s^{-1}, k_{on} = 0.25 (k_{off} + k_d) = 0.0035 and k_{off} = 1/90 =

0.011 chosen to reflect a half-life of 240 s, a 90 s mean lifetime of ribosome-bound complexes,

and that approximately 20% of mRNAs appear bound at steady state. In the high-binding scenario

k_{on} was increase by a factor of 125. For the numerical implementation, the production constant is

represented by a small non-zero spread around the bud centre using a smooth step-function of

which the volume integral is normalized to 1:

15
$$k_p = \frac{k_{\text{max}} \cdot e^{s \cdot m}}{e^{m \cdot r} + e^{s \cdot m}}$$

For the simulation results shown in Extended Fig. 3d-f, the values chosen are *s* = 5 and *m* = 10,

and the normalization results in k_{max} = 0.0019. Although the exact value of this constant affects

the absolute concentration of mRNA it does not affect the ratio of mother to bud RNA. The PDE

is solved in three Cartesian coordinates using the FEM implementation in Wolfram Mathematica

(Wolfram Research, Inc., Mathematica, Version 12.3.1, Champaign, IL).

Ellipsoid fitting to mother and bud DIC images

Differential interference contrast (DIC) images were analyzed in Mathematica to fit 3D ellipsoids

to mother and bud models. X- and y-axes length were measured for the cells and the short axis

was used as estimation of the z-axis. The z-axis origin value was aligned with the z-stack images

by maximizing the FISH-Quant mRNA and protein point inclusions.

Pattern search to predict ZIP codes and synonymization

To identify potential ZIP codes in the *CLB2* mRNA, we performed a targeted pattern search[82]. In

the first step, we leniently screened for two nested pairs of inverted repeats with a minimal length

of four nucleotides that framed an asymmetric bulge region as found in the E3 ZIP code in the

ASH1 mRNA[68]. We also checked for the presence of a CGA motif and a singular cytosine on the

opposite strand with a defined distance of six nucleotides[75]. The search was performed on the

complete *CLB2* mRNA with 1476 nt of CDS (YPR119W; genomic coordinates: chromosome XVI,

771653-773128, +, genome version S288C; *Saccharomyces* Genome Database,

https://www.yeastgenome.org/locus/S000006323). We further added 366 nt 3' UTR and 346 nt 5'

UTR as previously determined by[48]. In the second step, the minimum free energy (MFE) folds of

all initial instances were analyzed using RNAfold with and without including a constraint on the

nested pairs of inverted repeats[83,84]. Fold prediction was performed at 28°C with 80 nt RNA

sequence fragments centered on each instance. Instances were only kept if (i) at least one of

inverted repeat pairs was present in the MFE structure without constraints, and (ii) the free energy

(ΔG) of the constraint structure did not differ by more than 20% from the MFE structure without

constraint. The latter accounts for energetic benefits from interaction with the She2-She3 proteins.

The pattern search predicted a single ZIP code at nucleotide positions +1111 to +1145 of the

CDS (genomic coordinates: chromosome XVI, 772763-772797, +). Fig. 2c displays the predicted

fold with constraint using RNAfold of ZIP code instance plus ±5 nt flanking sequence. Visualization

of the predicted structure in dot-bracket notation was generated using VARNA [85]. Repeating the

pattern search described above on the synonymized ZIP code mutant did not retrieve any hits.

The complete sequences of the synonymized ZIP code mutant are provided in the Resource

Table.

Sample preparation for live yeast fluorescence imaging

Yeast cells were grown at 26°C in synthetic selective medium. Exponentially growing cells (OD_{600}

0.2-0.4) were plated on coated Delta-T dishes (Bioptech 04200417C). The dishes coating was

done by incubating with Concanavalin A 1mg/ml (Cayman chemical company) for 10 minutes at

RT. Excess liquid was aspirated and dishes were dried at RT. To activate Concanavalin A, dishes

were incubated for 10 minutes at RT with a 50 mM $CaCl_2$ 50 mM $MnCl_2$ solution. Excess was

removed and dishes dried at RT. Finally, dishes were washed once with ultrapure water

(Invitrogen) and completely dried at RT. Cell attachment was performed by gravity for 20 minutes

at RT, excess liquid removed and substitution with fresh media. Cells were diluted to OD_{600} 0.1

and grown until OD_{600} 0.3-0.4. before being plated on Concanavalin A coated dish.

Single molecule mRNA live cell fluorescence imaging and image analysis

The two-color simultaneous imaging of mRNAs and the appropriate cellular marker was

performed on a modified version of the home-built microscope described in[57,58]. Briefly, the

microscope was built around an IX71 stand (Olympus). For excitation, a 491 nm laser

(CalypsoTM, Cobolt) and a 561 nm laser (JiveTM, Cobolt) were combined and controlled by an

acoustic-optic tunable filter (AOTF, AOTFnC-400.650-TN, AA Opto-electronic) before coupled

into a single mode optical fiber (Qioptiq). The output of the fiber was collimated and delivered

through the back port of the microscope and reflected into an Olympus 150x 1.45 N.A. Oil

immersion objective lens with a dichroic mirror (zt405/488/561rpc, 2mm substrate, Chroma). The

tube lens (180 mm focal length) was removed from the microscope and placed outside of the right

port. A triple band notch emission filter (zet405/488/561m) was used to filter the scattered laser

light. A dichroic mirror (T560LPXR, 3mm substrate, Chroma) was used to split the fluorescence

onto two precisely aligned EMCCDs (Andor iXon3, Model DU897) mounted on alignment stages

(x, y, z, θ- and φ- angle). Emission filters FF03-525/50-25 and FF01-607/70-25 (Semrock) were

placed in front of green and red channel cameras, respectively. The two cameras were triggered

for exposure with a TTL pulse generated on a DAQ board (Measurement Computing). The microscope was equipped with a piezo stage (ASI) for fast z-stack and a Delta-T incubation system (Bioptech) for live-cell imaging. The microscope (AOTF, DAQ, Stage and Cameras) was automated with the software Metamorph (Molecular Devices). For two-color live-cell imaging, yeast cells were streamed at 50 ms, Z plane was streamed, and z-stacks acquired every 0.5 μm. Single-molecule analysis was done on maximal projected images using Fiji. Maximally projected images were filtered using the Maxican Hat filter (Radius=2) in Fiji. Spots were identified and counted using the spot detection plugin integrated in TrackMate. LoG detector was used for the spot identification, object diameter= 3 and Quality threshold = 2500. Files were exported as csv files and plotted using GraphPad Prism.

Deconvolution algorithm

To reduce imaging artifacts arising from noise and optics of the microscope, we used the Huygens software v3.6, where a Classic Maximum Likelihood Estimation (CMLE) algorithm was applied as a restoration method to deconvolve the images used for protein-mRNA foci co-localization (Fig. 4). CMLE assumes the photon noise to be governed by Poisson statistics and optimizes the likelihood of an estimate of an object in the input 3D image while taking the point spread function into consideration. The CMLE deconvolution method was chosen since it is suited for images with low signal-to-noise ratio and to restore point-like objects. The result is a more accurate identification of the location of the object, which in our case is the fluorescently labeled mRNA and protein molecules. The restoration parameters used with the CMLE deconvolution algorithm was 99 iterations, a quality stop criterion of 0.01, and a signal-to-noise ratio of 15.

CLB2 mRNA bud localization quantification in living cells

For the analysis reported in Extended Data Fig. 3b, the ImageJ plugin Labkit (https://imagej.net/Labkit) was manually used to segment cells and RNAs. Segmented cells were used as input for training the deep learning program Stardist in 2 dimensions. Stardist was used

to automatically detect and segment cells and single mRNAs from live imaging movie frames (Cell Detection with Star-convex Polygons, https://arxiv.org/pdf/1806.03535.pdf). Cell and RNA segmentation was imported into R using the RImageJROI package. In R, the cell size, number of mRNAs in the bud and the distance of each bud localized mRNA to the periphery was calculated and plotted over time using the R packages Spatial Data and PBSmapping[86]. The Stardist segmentations were used to plot the RNAs and the cell's periphery onto the live imaging movie using the FFmpeg wrapper function for the FFmpeg multimedia framework (https://ffmpeg.org/).

Protein extraction and Western blot

Yeast strains were grown overnight at 26°C in yeast extract peptone dextrose (YEPD) medium with 2% glucose. In the morning, cells were diluted to OD_{600} 0.1 and allowed to grow until OD_{600} 0.5-1. Cell lysis was performed by adding 1 ml H_2O with 150 μL of Yex-lysis buffer (1.85 M NaOH, 7.5% 2-mercaptoethanol) to the pellet of 3-5 ODs of cells (~$3x10^7$) and kept 10 minutes on ice. Proteins were precipitated by the addition of 150 μL of TCA 50% for 10 minutes on ice. Cells were pelleted and resuspended in 100 μL of 1X sample buffer (1 M Tris-HCl pH 6.8, 8 M Urea, 20% SDS, 0.5 M EDTA, 1% 2-mercaptoethanol, 0.05% bromophenol blue). Total protein extracts were fractioned on SDS-PAGE and examined by Western blot with mouse anti-myc (Sigma), mouse anti-Pgk1 (Thermofisher). For quantitative Western blot analyses, fluorescent secondary α-Mouse (IRDye 800CW) and α-Rabbit (IRDye 680RD) antibodies were used. The signals were revealed using the LYCOR® scanner and quantified using LITE® Software.

Clb2 protein imaging in live cells

Cells were precultured in liquid SC medium containing 2% glucose (w/v). Exponentially growing cells were transferred to SC medium containing 2% glucose (w/v), solidified with 1.5% low melt agarose in an ibidi u-Slide 4 well sample holder. Cells were imaged at 30°C using a Nikon Ti-eclipse widefield fluorescence microscope equipped with a SOLA 6-LCR-SB light source (Lumencor), an Andor Zyla 5.5 sCMOS camera, and a Plan Apo lambda 100x immersion Oil

objective (NA 1.45, refractive index 1.515). Brightfield images were collected with 20 ms exposure

at an LED (CooLED) intensity of 7.6. Clb2p-GFP was imaged at 10% sola intensity with 200 ms

exposure and 4x4 hardware binning with a 480/40 excitation filter and a 535/50 emission filter

and a 505 nm longpass dichroic (Semrock). Cdc10-mCherry was imaged at 10% sola intensity

with 100 ms exposure and 4x4 hardware binning with a 570/20 excitation filter and a 610 longpass

emission filter and a 600 nm longpass dichroic (Semrock). Images were taken every 5 min in all

three channels.

Image processing

Bright-field images were segmented using a Cellpose model and association of objects in

subsequent frames (tracking) was achieved with a maximum matching approach. Briefly, a graph

was constructed with cell detections as nodes. Edges were drawn between nodes in subsequent

frames if the corresponding cells were less than a predefined threshold (200 px) apart. Edges

were given weights computed as 1/(node distance) and then the edge set was found that

connected each node to exactly one node in the subsequent frame and that minimized the total

distance. Bud necks were detected with a unet-based multiclass detector which was trained to

recognize background, cytosol, neck outline and neck centers. The training data for the detector

was generated through semi-automatic annotation of Cdc10-mCherry fluorescence images using

a napari-based custom interactive random forest plugin. Neck centers in different frames

pertaining to the same mother-bud pair were associated into tracks using the tracking approach

described above. Lengths of the neck tracks were reported as budded phase duration. Budding

frames were determined as the first time-point at which a fluorescent Cdc10-RFP patch could be

detected. In some cases, the emergence of a bud as judged from the bright-field image preceded

the formation of a detectable fluorescent patch. In these cases, the budding frame was adjusted

manually based on the bright-field information. Pixel areas of mother cells in the budding frame

were reported as size of the mother compartment at budding. Pixel areas of mother and daughter

cell in the first frame after a bona fide neck could be detected were reported as birth sizes of the

respective compartments.

Growth curves setup and analysis

Cells were grown overnight at 30°C in SC with 2% glucose. Cells in mid-log phase were spun

down, the supernatant was removed and cells were resuspended at a final OD_{600} of about 0.1 in

in SC with 2% glucose. In 48-well plates with flat bottom, 400 µL were plated per well. At least 3

well replicates were done per experiment. Cells were grown for the indicated time, at 30°C. OD_{600}

measurements were taken every 5 minutes with 700 rpm orbital shaking between time-points

using a CLARIOstar® plate reader (BMG Labtech). Growth curves analysis was performed using

an adaptation of the R package Growthcurver[87] and plotted using the R package ggplot2[88],

tydiverse[89], RColorBrewer[90], dplyr[91]. Growthcurver fits a basic form of the logistic equation to

experimental growth curve data. The logistic equation gives the number of cells N_t at time t.

13
$$N_t = \frac{K}{1 + \left(\dfrac{K - N_0}{N_0}\right) e^{-rt}}$$

The population size at the beginning of the growth curve is given by N_0. The maximum possible

population size in a particular environment, or the carrying capacity, is given by K. The intrinsic

growth rate of the population, r, is the growth rate that would occur if there were no restrictions

imposed on total population size. The best values of K, r, and N_0 for the growth curve data were

found using the implementation of the non-linear least-squares Levenberg-Marquardt algorithm.

The carrying capacity and growth rate values (K and r) were used to compare the growth

dynamics of the strains.

Flow cytometry sample preparation and analysis

Cells were grown overnight at 30°C in SC medium with 2% glucose. In the morning, cells were

diluted to OD_{600} 0.1 and were grown to mid-log phase (OD_{600} 0.3-0.4) with constant shaking (200

rpm) at 30°C. When cells had reached mid-log phase, 1 mL of culture was transferred to a 1.5

mL Eppendorf tube and centrifuged for 3 minutes at 3000 rpm. The supernatant was removed

and cells were resuspended in 70% ethanol and fixed overnight at 4°C. Cells were washed once

with 1xPBS pH 7.4, resuspended in 500 µL of 1xPBS with 1 µL of RNAse A (1 mg/mL) and

incubated at 37°C for 2 hours. After incubation, cells were washed with 1 mL of 1xPBS and

resuspended in 200 µL of 1xPBS. The resuspended cells were divided into two tubes: 94 µl of

cells were supplemented with 6 µL of a 1 mg/mL solution of propidium iodide (PI), obtaining a

final concentration of 60 µg/mL PI, while another 100 µL of cells remained unstained as negative

control. Both negative control and PI stained cells were incubated in a water bath at 30°C for 1

hour, covered from the light. Cells were then washed 3 times with 1 mL 1xPBS and resuspended

in 500 µl of 1xPBS. The cells were analyzed with the Beckman Coulter CytoFLEX S Flow

Cytometer (B2-R0-V2-Y2). A 561 nm laser was used to excite the PI stained cells and a band

pass filter (610/20 nm) was used to filter the emitted fluorescence. 50'000 cells were collected per

sample. Analysis and plotting was performed using R Studio and the following R packages:

ggplot2[88]; tydiverse[89], RColorBrewer[90], dplyr[91], mixtools[92].

Quantifications and statistical analysis

FISH-quant was used to quantify single mRNA molecules and protein foci in fixed samples. Fiji

was used to quantify single mRNA molecules in living cells. GraphPad Prism was used to

calculate the mean and the standard deviation (SD) of all the data and perform statistical analysis.

Flow cytometry data, growth curves analysis was performed in R Studio, as detailed in previous

paragraphs. For each experiment, the number of biological replicates, the number of cells

analyzed (n), statistical analysis applied and significance is indicated in the figure, figure legend

or in the main text. Symbols meaning: ns, $P > 0.05$; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$, ****,

$P \leq 0.0001$.

Figures and Legends

Figure 1

2 *Fig 1. CLB2 mRNAs localize to the bud in a cell-cycle dependent manner*
3 **(a)** Schematic of *CLB2* mRNA expression during the cell cycle. Green dots represent *CLB2*
4 mRNAs. The Whi5 protein (cyan) accumulates in the nucleus during early G1. Tubulin (magenta)
5 is a component of microtubules and the mitotic spindle. The bud stage starts during S phase and
6 ends with the formation of the daughter cell. During anaphase, microtubules stretch between the

mother and the daughter cell. **(b)** Top panels: MERGE Maximal projections of IF anti-HA (Whi5)
(cyan) and DAPI (blue) merged to a single differential interference contrast (DIC) section (grey).
Bottom panels: MERGE Maximal projections of *CLB2* mRNA smFISH (green), anti-tubulin IF
(magenta) and DAPI (blue) merged to a single DIC section (grey). The corresponding cell cycle
phase is indicated on the panels. Scale bars 3 µm. **(c)** smFISH quantifications of *CLB2* mRNA
expression during the different cell cycle phases determined using the markers shown in (b). Dots
correspond to individual cells (2083 cells, from two replicates). The black bar indicates the
average (G1 early= 1.2±1.9; G1 late= 1.2±1.9; S= 3.8±3.0; G2= 10.3±5.7; M= 8.0±6.5 mRNAs/cell
mean ± standard deviation (SD)). **(d)** Relative distribution (bud vs mother) of *CLB2* mRNA in WT
budded cells based on the smFISH-IF data shown in (b) (S phase= 58.4±5.5, G2 phase=
34.6±0.9, M phase= 34.4±7.2; mean ± SD). **(e)** *CLB2* endogenously tagged with 24xMBSV6 to
enable visualization of the mRNA (black) in live cells. The bud neck protein Cdc10-tdTomato is
shown in magenta. Cell A is the mother of cell B. Time point from start of acquisition is indicated
in the upper left corner of each time frame. Scale bars 3 µm. **(f)** Percentage of *CLB2* mRNAs
localized in the bud over time from bud appearance. Error bars indicates mean ±SD **(g)** Number
of *CLB2* mRNAs per cell over time in cell A and cell B shown in panel (e).

a MERGE smFISH-IF
DIC /DAPI/*CLB2*/Tub1p

b Predicted ZIP-code

c Predicted ZIP-code
ΔG = -17.3 kcal/mol

d Synonymized ZIP-code mutant

1111											1145
CLB2 WT	GCAGATGACTACGATATACAGTCTCGAACTCTTGC
CLB2 ZIP Mut	GCCGATGATTATGATATACAGAGCAGAACCCTAGC
Clb2 protein	A D D Y D I Q S R T L A

e *CLB2* DAPI MERGE

f Relative mRNA distribution (%) — Bud / Mother — WT / *ZIP mut*

g Peripheral Distribution Index (PDI) — PDI=3 PDI=2 PDI=1 PDI=0

h PDI — **** — WT ASH1, WT CLB2, Δshe2 CLB2, Δshe3 CLB2, ZIP mut, WT MDN1

1 ***Fig. 2 The She2-3 complex and an RNA ZIP code in CLB2 mRNA CDS are required for bud***
2 ***localization.***
3 **(a)** smFISH-IF in WT, *Δshe2*, and *Δshe3* strains. MERGE Maximal projections of *CLB2* mRNA
4 smFISH (green), anti-tubulin IF (magenta) and DAPI (blue) merged to a single DIC section (grey).
5 The cell cycle phase is indicated on top of the panel. Scale bars 3 μm. **(b)** Schematic of *CLB2*

mRNA coding sequence. The blue box represents the ZIP-code at nucleotides 1111-1145 (relative to START codon). **(c)** Predicted secondary structure of ZIP code (blue box) with flanking sequence (nt 1089- 1168). The free energy (ΔG) of the mRNA folding is indicated. **(d)** Synonymized ZIP-mutant (*ZIP-mut*). Top sequence is *CLB2* WT. Bottom sequence is the synonymized sequence. Mutated nucleotides are indicated in red. Below is the corresponding Clb2 protein amino acid sequence, which is identical for both the WT and synonymized strains. Rare amino acid codons were avoided to maintain the same codon usage frequency = 0.74 for the *CLB2* mRNA sequence. **(e)** smFISH in WT and ZIP mutant strain. Maximal projections of smFISH with *CLB2* probes (green) and DAPI (blue) and fluorescence images overlapped to a single DIC section (MERGE). Scale bars 3 μm. **(f)** Relative bud vs mother distribution of the *CLB2* mRNA in WT and ZIP mutant strain based on smFISH-IF data shown in (e). **(g)** Schematic representation of mRNA peripheral distribution index (PDI). Black dots represent mRNA. Blue is the nucleus. A PDI close to 0 indicates that the RNA of interest is localized near the nucleus. A PDI of 1 indicates that the RNA is diffusely dispersed throughout the cell. As the PDI value grows > 1, the polarization of the mRNA increases. **(h)** PDI in WT cells for *ASH1*, *CLB2* and *MDN1* mRNAs and in *Δshe2*, *Δshe3* and ZIP mutant strains for the *CLB2* mRNA. Index values are calculated from smFISH-IF experiments shown in (e) and Extended Data Fig. 1c and Extended Data Fig. 4a. Significant difference between WT and localization mutants tested by ANOVA statistical test: $F_{(4, 185)} = 15.74$, $p < 0.0001$).

1 **Fig. 3. CLB2 mRNA mis-localization affects Clb2 protein expression but not mRNA levels,**
2 **nor protein stability or localization.**
3 **(a)** smFISH quantifications of *CLB2* mRNA expression in WT and localization mutants. Dots
4 correspond to individual cells, from at least 3 replicates. The black bar indicates the average
5 number of mRNAs/cell in expressing cells (WT n=3035, 5.4±5.1; *Δshe2* n=1787, 5.3±6.2; *Δshe3*

n=1827, 5.2±6.2; ZIP mutant (ZIP[1]) n=667, 6.9±6.3 mRNAs/cell±SD). Statistical analysis: Anova, Dunnet's multiple comparisons test. **(b)** Quantification of nascent *CLB2* RNAs at transcription sites (TS) from smFISH. Dots correspond to individual cells, from at least 3 replicates (WT n=500, 2.6±1.3; *Δshe2* n=162, 2.7±1.1; *Δshe3* n=186; 2.7±1.3, ZIP mutant (ZIP[1]) n=132, 3.4±1.3 nascent RNAs/TS±SD). Statistical analysis: Anova, Dunnet's multiple comparisons test. **(c)** Western blot analysis using anti-myc antibody against Clb2 protein endogenously tagged with 5 myc tags in WT, *Δshe2*, *Δshe3* and ZIP mutant (ZIP[1]) cells. First lane is the control untagged strain. Endogenous Pgk1 protein was used as loading control. **(d)** Quantification of western blot in (c). Myc signal normalized to Pgk1 loading control. Protein levels relative to WT indicated. Mean ± SD from 3 replicates. **(e)** Clb2 protein stability assay in WT, *Δshe2* and *Δshe3* cells. Western blot was performed using an anti-myc antibody to target Clb2 protein tagged with 5 myc tags in cells treated with 100 µg/ml cycloheximide for 0, 5, 10, 20, 30, 45 or 60 minutes. Pgk1 was used as loading control. Error bars indicates mean ± SD. **(f)** Clb2 protein stability assay in WT and ZIP mutant (ZIP[1]) as done in (e). Western blot was performed in the same way as in (c). Error bars indicate mean ± SD **(g)** Clb2 fused to yeGFP in WT, *Δshe2*, *Δshe3* and ZIP mutant cells. Maximal projections of Clb2-GFP (green) overlapped to a single DIC section. Scale bars 2 µm.

1 ***Fig. 4. CLB2 mRNA and protein colocalization suggests preferential translation in the bud.***
2 **(a)** Schematic of the *CLB2* endogenous gene tagging to simultaneously visualize single mRNA
3 by smFISH and Clb2 proteins by IF against the myc tag in fixed cells. 25 myc tags are inserted at
4 the beginning of the *CLB2* CDS, after the ATG. **(b)** smFISH-IF in WT cells. Top panels: MERGE

Maximal projections of IF anti-tubulin (cyan) and DAPI (blue) merged to a single DIC section (grey). Second panel row from the top: MERGE Maximal projections of IF anti-myc-Clb2 protein (magenta), *CLB2* mRNA smFISH (green) and DAPI (blue) merged to a single DIC section (grey). The third and fourth panels from the top are the *CLB2* mRNA smFISH and Clb2 protein IF, respectively. Scale bars 3 μm. **(c)** Quantification of protein foci in WT bud (B) and mother (M) cells from IF experiments shown in (a). Cell cycle classification was performed using tubulin and DAPI as markers. **(d)** Quantification of mRNA in WT bud (B) and mother (M) cells from smFISH experiments shown in (a). Cell cycle classification was performed as in (b). **(e)** smFISH-IF in *Δshe2* cells. Maximal projections of IF anti-myc-Clb2 protein (magenta), *CLB2* mRNA smFISH (green) and DAPI (blue) merged to a single DIC section (grey). Panels description as in (a). Scale bars 2 μm. **(f)** Quantification of *CLB2* mRNA-protein foci found in close proximity (<250 nm) by smFISH-IF experiments performed in bud (B) and mother (M) cells of WT and *Δshe2* cells shown in panels (a) and (d). **(g)** Quantification of bud-localized mRNAs co-localizing with protein foci (<250 nm) in WT and *Δshe2* cells shown in panels (a) and (d). **** indicate a P value < 0.0001 calculated by using the Mann Whitney significance test.

Fig. 5. CLB2 mRNA ZIP code mutant displays larger daughter size and loss of
communication between mother and bud compartment.
(a) Snapshots from Videos 3 (WT) and 6 (ZIP-mutant, ZIP[1]) at indicated times from live cell
imaging of *CLB2* protein endogenously tagged with yeGFP (green) and Cdc10 tagged with
mCherry (magenta). Scale bars 5 μm. Arrows point to single yeast cells undergoing budding and
birth of the daughter cell. **(b)** ZIP mutant daughters are born at a larger size. Scaled daughter size
at birth vs scaled daughter size at division shown for WT and the ZIP-mutant strain. In both strains
daughter size weakly correlates with mother size (Pearson correlation WT: 0.32 (N=431),
ZIP:0.25, (N=428). Dotted lines indicate median of the data, divided by median of the WT strain

(ZIP-mutant mother size at budding: 1.03, daughter size at birth: 1.109). Data shown were pooled
from three biological replicates per strain. **(c)** Boxplots showing mother size at budding, $p=5.8*10^{-4}$;
(d) mother size at division, $p=3.5*10^{-5}$. **(e)** Daughter size at division, $p=2.2*10^{-16}$. **(f)** Combined
size of mother and daughter at division, $p=1.44*10^{-13}$. **(g)** Size added during budded phase
$p=1.19*10^{-12}$. **(h)** Time spent in budded phase $p=0.293$. **(i)** Average fluorescence of Clb2-yeGFP
of the 10% brightest pixels per cell as a proxy for nuclear fluorescence over time. Traces shown
are recorded between budding and division for cells born in during the experiment and are pooled
from three biological replicates. **(j)** Boxplot showing cumulative fluorescence signal for the
trajectories in (I), $p = 4.3*10^{-12}$, one-way ANOVA. **(k)** Boxplots showing the slopes of linear
regression lines of bud size on nuclear fluorescence intensity for individual trajectories.
Individual boxplots show biological replicates, p values reported are from one-way ANOVA.

1 **Fig. 6. The *CLB2* ZIP mutant shows a small but significant cell cycle defect and the ZIP**
2 **code rescue strain partially recovers protein expression and cell cycle progression.**
3 **(a)** Schematic of *CLB2* ZIP code rescue strain. The blue box represents the WT ZIP code inserted
4 41 nt after the stop codon in the 3'UTR. The Pink box represents the mutated ZIP code at

nucleotides 1111-1145 (nucleotide number relative to START codon). **(b)** smFISH in WT, ZIP code and ZIP code rescue strain. Maximal projections of smFISH with *CLB2* probes (green) and DAPI (blue) and fluorescence images overlapped. Scale bars 3 μm. **(c)** Western blot analysis using anti-myc antibody against Clb2 protein endogenously tagged with 5 myc tags in WT, *Δshe2*, *Δshe3*, ZIP mutant and ZIP code rescue cells. First lane is the control untagged strain. Endogenous Pgk1 protein was used as loading control. **(d)** Quantification of western blot in (c). Myc signal normalized to Pgk1 loading control. Protein levels relative to WT indicated. Mean ± SD from 3-6 replicates. Asterisk represent significance of <0.05 as determined by t-test. **(e)** Cell cycle analysis by DNA content estimation with flow cytometry in in WT, *Δshe2*, *Δshe3*, ZIP mutant and ZIP code rescue cells. DNA was stained with propidium iodide to measure the G1 and G2/M components, respectively. **(f)** Relative distribution (%) of cells in G1, S and G2/M phase obtained by performing mixed Gaussian fitting to estimate the three subpopulations. Mean ± SD from 3 experiments. Statistical significance was calculated using Two-ways ANOVA. **(g)** Working model for *CLB2* mRNA localization translation in the *S. cerevisiae* bud.

Extended Data Fig. 1. CLB2 mRNAs are localized also in a different S. cerevisiae
background unlike a control mRNA.
(a) Quantification of *CLB2* mRNA smFISH shown in Fig. 1b reported as relative frequency
distribution of mature mRNAs per cell. Data from two replicates (n=2083). **(b)** Quantification of
CLB2 nascent RNAs at transcriptions sites (TS) from smFISH shown in Fig. 1b reported as
relative frequency distribution of nascent RNAs per TS. Data from two replicates (n=2083). **(c)**
MDN1 smFISH maximal projection (green), and DAPI merged to a single DIC section (grey).
Scale bar 5 µm. **(d)** Relative bud vs mother distribution of *MDN1* mRNA in budded cells based on
the smFISH data shown in (c). Data from two replicates (n=2011; 82.9±3.7, mean ± SD). **(e)**
smFISH in the *S. cerevisiae* background W303. MERGE Maximal projections of *CLB2* mRNA
smFISH (green) and DAPI (blue) merged to a single DIC section (grey). Scale bar 3 µm.

Extended Data Fig.2. CLB2 tagging with the MBSV6 reporter recapitulates CLB2 mRNA
expression.
(a) Schematic of *CLB2* locus endogenously tagged with 24xMBSV6 inserted in the 3' UTR after
the STOP codon. Dotted lines represent smFISH probe positions targeting the CDS (green) or
MBS sequences (magenta). **(b)** Two-color smFISH for cells expressing tagged *CLB2* mRNAs.
Top panels (MBSV6), cells expressing the control vector (YcpLac111). Bottom panels
(MBSV6+MCP) cells expressing MCP (YcpLac111 CYC1p-MCP-NLS-2xyeGFP). DIC/MERGE
shows the overlap of the DAPI signal in the nucleus (blue), smFISH for the *CLB2* CDS (green)
and the MBS (magenta) with the differential interference contrast (DIC) image. Individual
fluorescence channels are shown in grayscale. Scale bars 3 µm. **(c)** Quantification of smFISH
shown in (b) as well as untagged WT cells, with CDS probes (green plots) or MBS probes
(magenta) reported as distribution of mature mRNAs per cell. Mean of three biological replicates,
(*CLB2* MBSV6: CLB2 probes n= 864, mean ± SD 4.9 ± 5.9 mRNA/cell; MBSV6 probes n=771,
mean ± SD 5.4 ± 6.1 mRNA/cell; *CLB2* MBSV6+MCP: *CLB2* probes n = 723, mean ± SD 4.5±
5.0 mRNA/cell, MBSV6 probes n=977, mean ± SD 4.9±6.1 mRNA/cell; WT cells: *CLB2* probes n
= 791, mean ± SD 5.1±5.1 mRNA/cell). Statistical differences were calculated by non-parametric
Mann-Whitney test. **(d)** and **(e)** Correlation between the number of single CDS and MBSV6
molecules per cell in presence or absence of MCP. Pearson *r* values calculated by combining two
independent experiments (n=1908 and n=2284, respectively).

2 *Extended Data Fig. 3. CLB2 mRNA imaging throughout the cell cycle reveals a rapid mRNA*
3 *accumulation and localization to the bud and mathematical modelling predicts the*
4 *existence of mRNA anchoring factors in the bud.*
5 **(a)** Schematic of *CLB2* localization during the cell cycle. The *CLB2* mRNA (green), the bud neck
6 protein Cdc10 is tagged with tdTomato (magenta). **(b)** Average number of *CLB2* mRNA per cell
7 tagged with 24xMBSV6 monitored over time (purple line), grey represents the SD. **(c) Top.** Snap
8 shots from live cell imaging of *CLB2* mRNA endogenously tagged with 24xMBSV6. Approximate

cell outline identified from fluorescent background (dashed blue line). A single Z plane was
acquired every 100 ms. Scale bars 3 µm. **Middle.** Average cell area monitored over time in live
cells (n=3). A single Z plane was acquired every 100 ms. Pink, purple and green lines represent
time points at which the snapshots on top were taken. **Bottom.** Average number of bud localized
mRNAs per cell over time as monitored from 3 single cells. Pink, purple and green lines represent
the time points at which the snapshots on top were taken. **(d-f)** Simulation of localization of mRNA
particles per nm^3 assuming **(d)** fast diffusion coefficient of 0.4 μm^2/s, **(e)** slow diffusion coefficient
of 0.1 μm^2/s and **(f)** slow diffusion coefficient of 0.1 μm^2/s in combination with a high-affinity
binding factor.

a

Whi5-HA
DAPI
DIC

WT | Δ*she2* | Δ*she3*

CLB2 mRNA
Tub1
DAPI
DIC

b Δshe2

Relative distribution %
Bud / Mother
S, G2, Mitosis

c Δshe3

Relative distribution %
Bud / Mother
S, G2, Mitosis

d Synonymized ZIP-code mutants

1111 1146

CLB2 WT   GCAGATGACTACGATATACAGTCTCGAACTCTTGCC
CLB2 ZIP¹ GCCGATGATTATGATATACAGAGCAGAACCCTAGCC
CLB2 ZIP² GCCGACGACTACGATATCCAGTCTCGAACTCTTGCA
CLB2 ZIP³ GCGGATGATTATGATATACAGAGCCGAACGCTGGCG
Clb2 protein A   D   D   Y   D   I   Q   S   R   T   L   A

e

Strain	Amino acid seq												Average
	A	D	D	Y	D	I	Q	S	R	T	L	A	
WT	0.29	0.65	0.35	0.44	0.65	0.56	0.31	0.26	0.07	0.35	0.13	0.22	0.36
ZIP¹	0.22	0.65	0.65	0.56	0.65	0.56	0.31	0.11	0.48	0.22	0.14	0.22	0.40
ZIP²	0.22	0.35	0.35	0.44	0.65	0.26	0.31	0.26	0.07	0.35	0.13	0.38	0.31
ZIP³	0.11	0.65	0.65	0.56	0.65	0.56	0.31	0.11	0.07	0.13	0.11	0.11	0.34

Codon usage freq

f

CLB2 | DAPI | MERGE

CLB2 ZIP²

CLB2 ZIP³

1 **Extended Data Fig. 4. *CLB2* is not bud localized in the Δshe mutants and alternative ZIP**
2 **code mutants.**
3 **(a)** WT, *Δshe2* and *Δshe3* cells: Top panels, MERGE Maximal projections of IF anti-HA (Whi5)
4 (cyan) and DAPI (blue) merged to a single DIC section (grey). Bottom panels, MERGE Maximal

projections of *CLB2* mRNA smFISH (green), anti-tubulin IF (magenta) and DAPI (blue) merged
to a single DIC section (grey). Scale bars 5 µm. **(b)** Relative bud vs mother distribution of the
CLB2 mRNA in *Δshe2* budded cells based on the smFISH-IF data shown in (a) and Fig. 2a. (S
phase= 92.9±0.2, G2 phase= 82.4±7.9, M phase= 75.5±0.7; mean ± SD). **(c)** Relative bud vs
mother distribution of the *CLB2* mRNA in *Δshe3* budded cells based on the smFISH-IF data
shown in (a) and Fig. 2a. (S phase= 92.44±3.3, G2 phase= 84.0±4.6, M phase= 76.4±1.8; mean
± SD) (S phase= 58.4±5.5, G2 phase= 34.6±0.9, M phase= 34.4±7.2; mean ± SD). **(d)**
Synonymized ZIP-mutant variants (ZIP^1, ZIP^2 and ZIP^3) comparison to *CLB2* WT (top sequence).
Mutated nucleotides are indicated in red. Below is the corresponding Clb2 protein amino acid
sequence, which is identical for both the WT and synonymized strains. **(e)** Table summarizing the
codon usage frequency for the sequences of the three ZIP code variants, ZIP^1, ZIP^2 and ZIP^3
compared to the WT ZIP code. **(f)** smFISH in the ZIP code mutant strains ZIP^2 and ZIP^3. Maximal
projections of smFISH with *CLB2* probes (green) and DAPI (blue) and fluorescence images
overlapped (MERGE). Scale bars 3 µm.

Extended Data Figure 5

Extended data Fig. 5. Multiple ZIP code mutants show a reduction in Clb2 protein
expression and *CLB2* mRNA mislocalization is not affecting protein stability.
(a) Western blot analysis using anti-myc antibody against Clb2 protein endogenously tagged with
5 myc tags in WT, ZIP^1, ZIP^2 and ZIP^3. First lane is the control untagged strain. Endogenous Pgk1
protein was used as loading control. **(b)** Quantification of western blot in (a). Myc signal
normalized to Pgk1 loading control. Protein levels relative to WT indicated. Mean ± SD from 4
replicates. **(c)** Example of western blot of Clb2 protein stability assay in WT, *Δshe2* and *Δshe3*
cells. Western blot was performed using an anti-myc antibody to target Myc-tagged Clb2 protein

tagged in cells treated with 100 µg/ml cycloheximide for 0, 5, 10, 20, 30, 45 or 60 minutes. Pgk1
protein was used as loading control. Quantifications are reported in Fig. 3e. **(d)** Example of
western blot of Clb2 protein stability assay in WT and ZIP mutant cells. Western blot was
performed using an anti-myc antibody to target Clb2 protein tagged with 5 myc tags in cells treated
with 100 µg/ml cycloheximide for 0, 5, 10, 20, 30, 45 or 60 minutes. Pgk1 was used as loading
control. Quantifications are reported in Fig. 3f.

Extended Data Figure 6

Extended Data Fig. 6. Known translation inhibitors are not involved in regulating Clb2
protein expression.
(a) Western blot analysis of *CLB2* Myc tagged strains. Top row is the myc signal. Bottom row is
the Pgk1 loading control. First lane is untagged cells. Second and third lanes are the Clb2 protein
tagged with 5 and 25 myc tags, respectively. **(b)** Growth curves of strains with Myc-tagged Clb2
protein performed in SC 2% glucose at 26°C. Growth curves are fitted to a logistic curve (red
curve). **(c)** Western blot analysis using anti-myc antibody against Clb2 protein tagged with 25 myc
tags in WT, *Δssd1, Δkhd1* and *Δpuf6* cells. Pgk1 protein was used as loading control. First lane
is untagged cells. **(d)** Quantification of (c). Mutant strain signal is normalized to WT signal. Each
color corresponds to one replicate experiment. Black bar indicates mean ± SD (WT= 100; Δssd1=
88.3±15.6; Δkhd1= 98.8±13.5; Δpuf6= 83.9±10.4). **(e)** Merge maximal projections of *CLB2*
smFISH (green), DIC (gray), and DAPI (blue) in WT, *Δssd1, Δkhd1* and *Δpuf6* cells. Scale bars 3
µm.

Extended Data Fig. 7: Δshe mutations have minor effects on cell size.

(a) Normalized scatter of scaled daughter size at division vs scaled mother size at budding for *Δshe2* n=598, vs WT (n=431), and **(b)** for *Δshe3,* (n=475) vs WT. **(c)** Boxplots showing mother size at budding, WT vs *Δshe2*: p=1.2*10⁻⁵, WT vs *Δshe3*: p=0.778. **(d)** Mother size at division, WT vs *Δshe2*: p=2.69*10⁻⁵, WT vs *Δshe3*: p=0.573. **(e)** Daughter size at division, WT vs *Δshe2*: p=0.107, WT vs *Δshe3*: p=0.412. **(f)** Combined size of mother and daughter compartments at

division, WT vs *Δshe2*: p=0.035, WT vs *Δshe3*: p=0.410. **(g)** Size added during budded phase, WT vs *Δshe2*: p=0.042, WT vs *Δshe3*: p=0.109. **(h)** Size added during budded phase and normalized by mother size, WT vs zip: $p=1.96*10^{-4}$, WT vs *Δshe2*: $p=6.42*10^{-5}$, WT vs *Δshe3*: p=0.120. **(i)** Normalized histogram of budded phase duration for WT and ZIP-code mutant. **(j)** Boxplot of time spent in budded phase, WT vs *Δshe2*: $p=1.26*10^{-6}$, WT vs *Δshe3*: p=0.067. p values reported are from one-way ANOVA. **(k)** Average fluorescence of Clb2-yeGFP of the 10% brightest pixels per cell as a proxy for nuclear fluorescence over time. Traces shown are recorded between budding and division for cells born in during the experiment and are pooled from three biological replicates. **(l)** boxplots showing the cumulative signal of the trajectories shown in (k) WT vs *Δshe2*: p=0.507, WT vs *Δshe3*: p=0.324. **(m)** Boxplots showing the slopes of linear regression lines of bud size on nuclear fluorescence intensity for individual trajectories. Individual boxplots show biological replicates, p values reported are from one-way ANOVA.

Extended Data Fig.8. The CLB2 localization mutants do not show significant growth rates changes while the ZIP mutant shows an increase in the G2/M population
(a) Growth curves of WT, *Δshe2*, *Δshe3*, ZIP code mutant and ZIP code rescue cells performed in SC 2% glucose at 30°C. **(b)** Cell cycle analysis by DNA content estimation with flow cytometry in in WT, *Δshe2*, *Δshe3*, ZIP code mutant and ZIP code rescue cells. Representative experiment showing the results of the mixed Gaussian fitting to estimate the three subpopulations, G1 (purple), S (green) and G2/M (yellow).

Supplementary Video Legends

Video 1. CLB2 mRNA imaging in live cells throughout the cell cycle

CLB2 endogenously tagged with 24xMBSV6 to enable visualization of the mRNA (black) in live

cells. The bud neck protein Cdc10-tdTomato is shown in magenta. Two-color imaging was

performed by acquiring 13 Z-stacks every 0.5 μm, at two-minutes intervals and with exposure

time of 50 ms. In video, at each time point, Z-stacks were max-projected. Scale bar 3 μm.

Video 2. CLB2 mRNA imaging in live cells at high frame-rate acquisition

CLB2 endogenously tagged with 24xMBSV6 to enable visualization of the mRNA (black) in live

cells. The bud neck protein Cdc10-tdTomato is shown in magenta. One single Z-stack was

streamed every 50 ms. Scale bar 2 μm.

Video 3. Clb2 protein imaging in WT cells

CLB2 endogenously tagged with yeGFP to enable visualization of the protein (green) in WT live

cells. The bud neck protein mCherry-Cdc10 is shown in magenta. One single Z-stack was

acquired every 5 minutes. Clb2p-GFP was imaged with 200 ms exposure, mCherry-Cdc10 was

imaged with 100 ms exposure and brightfield images were collected with 20 ms exposure. Scale

bar 5 μm.

Video 4. Clb2 protein imaging in ∆she2 cells

CLB2 endogenously tagged with yeGFP to enable visualization of the protein (green) in *∆she2*

live cells. The bud neck protein mCherry-Cdc10 is shown in magenta. One single Z-stack was

acquired every 5 minutes. Clb2p-GFP was imaged with 200 ms exposure, mCherry-Cdc10 was

imaged with 100 ms exposure and brightfield images were collected with 20 ms exposure. Scale

bar 5 μm.

Video 5. Clb2 protein imaging in Δshe3 cells

CLB2 endogenously tagged with yeGFP to enable visualization of the protein (green) in *Δshe3*

live cells. The bud neck protein mCherry-Cdc10 is shown in magenta. One single Z-stack was

acquired every 5 minutes. Clb2p-GFP was imaged with 200 ms exposure, mCherry-Cdc10 was

imaged with 100 ms exposure and bright field images were collected with 20 ms exposure. Scale

bar 5 µm.

Video 6. Clb2 protein imaging in ZIP-code mutant cells

CLB2 endogenously tagged with yeGFP to enable visualization of the protein (green) in ZIP-code

mutant live cells. The bud neck protein mCherry-Cdc10 is shown in magenta. One single Z-stack

was acquired every 5 minutes. Clb2p-GFP was imaged with 200 ms exposure, mCherry-Cdc10

was imaged with 100 ms exposure and brightfield images were collected with 20 ms exposure.

Scale bar 5 µm.

References

1. Das, S., Vera, M., Gandin, V., Singer, R. H. & Tutucci, E. Intracellular mRNA transport and localized translation. *Nature Reviews Molecular Cell Biology* 1–22 (2021) doi:10.1038/s41580-021-00356-8.

2. Kejiou, N. S. & Palazzo, A. F. mRNA localization as a rheostat to regulate subcellular gene expression. *Wiley Interdiscip Rev RNA* **8**, (2017).

3. Lecuyer, E. *et al.* Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* **131**, 174–87 (2007).

4. Nevo-Dinur, K., Nussbaum-Shochat, A., Ben-Yehuda, S. & Amster-Choder, O. Translation-Independent Localization of mRNA in E. coli. *Science* **331**, 1081–1084 (2011).

5. Niessing, D., Jansen, R.-P., Pohlmann, T. & Feldbrügge, M. mRNA transport in fungal top models. *Wiley Interdiscip Rev RNA* **9**, (2018).

6. Bertrand, E. *et al.* Localization of ASH1 mRNA particles in living yeast. *Mol Cell* **2**, 437–45 (1998).

7. Böhl, F., Kruse, C., Frank, A., Ferring, D. & Jansen, R.-P. She2p, a novel RNA-binding protein tethers ASH1 mRNA to the Myo4p myosin motor via She3p. *EMBO J* **19**, 5514–5524 (2000).

8. Long, R. M., Gu, W., Lorimer, E., Singer, R. H. & Chartrand, P. She2p is a novel RNA-binding protein that recruits the Myo4p-She3p complex to ASH1 mRNA. *Embo J* **19**, 6592–601 (2000).

9. Long, R. M. *et al.* Mating type switching in yeast controlled by asymmetric localization of ASH1 mRNA. *Science* **277**, 383–7 (1997).

10. Long, R. M. *et al.* Characterization of transport and localization of ASH1 mRNA in yeast. *Mol Biol Cell* **8**, 2060–2060 (1997).

11. Niessing, D., Huttelmaier, S., Zenklusen, D., Singer, R. H. & Burley, S. K. She2p is a novel RNA binding protein with a basic helical hairpin motif. *Cell* **119**, 491–502 (2004).

1 12. Shen, Z., Paquin, N., Forget, A. & Chartrand, P. Nuclear shuttling of She2p couples ASH1

2  mRNA localization to its translational repression by recruiting Loc1p and Puf6p. *Mol Biol Cell*

3  **20**, 2265–75 (2009).

4 13. Takizawa, P. A. & Vale, R. D. The myosin motor, Myo4p, binds Ash1 mRNA via the adapter

5  protein, She3p. *Proc Natl Acad Sci U S A* **97**, 5273–8 (2000).

6 14. Deng, Y., Singer, R. H. & Gu, W. Translation of ASH1 mRNA is repressed by Puf6p-

7  Fun12p/eIF5B interaction and released by CK2 phosphorylation. *Genes & development* **22**,

8  1037–50 (2008).

9 15. Gu, W., Deng, Y., Zenklusen, D. & Singer, R. H. A new yeast PUF family protein, Puf6p,

10  represses ASH1 mRNA translation and is required for its localization. *Genes & development*

11  **18**, 1452–65 (2004).

12 16. Irie, K. *et al.* The Khd1 protein, which has three KH RNA-binding motifs, is required for proper

13  localization of ASH1 mRNA in yeast. *The EMBO journal* **21**, 1158–67 (2002).

14 17. Paquin, N. *et al.* Local activation of yeast ASH1 mRNA translation through phosphorylation of

15  Khd1p by the casein kinase Yck1p. *Mol Cell* **26**, 795–809 (2007).

16 18. Hasegawa, Y., Irie, K. & Gerber, A. P. Distinct roles for Khd1p in the localization and

17  expression of bud-localized mRNAs in yeast. *RNA* **14**, 2333–2347 (2008).

18 19. Shahbabian, K., Jeronimo, C., Forget, A., Robert, F. & Chartrand, P. Co-transcriptional

19  recruitment of Puf6 by She2 couples translational repression to mRNA localization. *Nucleic*

20  *Acids Res* **42**, 8692–8704 (2014).

21 20. Wanless, A. G., Lin, Y. & Weiss, E. L. Cell Morphogenesis Proteins Are Translationally

22  Controlled through UTRs by the Ndr/LATS Target Ssd1. *PLOS ONE* **9**, e85212 (2014).

23 21. Ballou, E. R., Cook, A. G. & Wallace, E. W. J. Repeated Evolution of Inactive

24  Pseudonucleases in a Fungal Branch of the Dis3/RNase II Family of Nucleases. *Molecular*

25  *Biology and Evolution* **38**, 1837–1846 (2021).

22. Bayne, R. A. *et al.* Yeast Ssd1 is a non-enzymatic member of the RNase II family with an alternative RNA recognition site. *Nucleic Acids Research* **50**, 2923–2937 (2022).

23. Jansen, J. M., Wanless, A. G., Seidel, C. W. & Weiss, E. L. Cbk1 Regulation of the RNA-Binding Protein Ssd1 Integrates Cell Fate with Translational Control. *Current Biology* **19**, 2114–2120 (2009).

24. Shepard, K. A. *et al.* Widespread cytoplasmic mRNA transport in yeast: Identification of 22 bud-localized transcripts using DNA microarray analysis. *PNAS* **100**, 11429–11434 (2003).

25. Amon, A., Irniger, S. & Nasmyth, K. Closing the cell cycle circle in yeast: G2 cyclin proteolysis initiated at mitosis persists until the activation of G1 cyclins in the next cycle. *Cell* **77**, 1037–50 (1994).

26. Amon, A., Tyers, M., Futcher, B. & Nasmyth, K. Mechanisms that help the yeast cell cycle clock tick: G2 cyclins transcriptionally activate G2 cyclins and repress G1 cyclins. *Cell* **74**, 993–1007 (1993).

27. Fitch, I. *et al.* Characterization of four B-type cyclin genes of the budding yeast Saccharomyces cerevisiae. *Molecular biology of the cell* **3**, 805–18 (1992).

28. Ghiara, J. B. *et al.* A cyclin B homolog in S. cerevisiae: chronic activation of the Cdc28 protein kinase by cyclin prevents exit from mitosis. *Cell* **65**, 163–174 (1991).

29. Surana, U. *et al.* Destruction of the CDC28/CLB mitotic kinase is not required for the metaphase to anaphase transition in budding yeast. *The EMBO journal* **12**, 1969–78 (1993).

30. Surana, U. *et al.* The role of CDC28 and cyclins during mitosis in the budding yeast S. cerevisiae. *Cell* **65**, 145–161 (1991).

31. Hood, J. K., Hwang, W. W. & Silver, P. A. The Saccharomyces cerevisiae cyclin Clb2p is targeted to multiple subcellular locations by cis- and trans-acting determinants. *Journal of Cell Science* **114**, 589–597 (2001).

32. Bailly, E., Cabantous, S., Sondaz, D., Bernadac, A. & Simon, M.-N. Differential cellular localization among mitotic cyclins from Saccharomyces cerevisiae: a new role for the axial

budding protein Bud3 in targeting Clb2 to the mother-bud neck. *Journal of Cell Science* **116**, 4119–4130 (2003).

33. Eluère, R. *et al.* Compartmentalization of the functions and regulation of the mitotic cyclin Clb2 in S. cerevisiae. *J Cell Sci* **120**, 702–711 (2007).

34. Coudreuse, D. & Nurse, P. Driving the cell cycle with a minimal CDK control network. *Nature* **468**, 1074–1079 (2010).

35. Curran, S., Dey, G., Rees, P. & Nurse, P. A quantitative and spatial analysis of cell cycle regulators during the fission yeast cycle. *Proc Natl Acad Sci U S A* **119**, e2206172119 (2022).

36. Harvey, S. L. *et al.* A phosphatase threshold sets the level of Cdk1 activity in early mitosis in budding yeast. *MBoC* **22**, 3595–3608 (2011).

37. Novak, B., Tyson, J. J., Gyorffy, B. & Csikasz-Nagy, A. Irreversible cell-cycle transitions are due to systems-level feedback. *Nat Cell Biol* **9**, 724–728 (2007).

38. Stern, B. & Nurse, P. A quantitative model for the cdc2 control of S phase and mitosis in fission yeast. *Trends in Genetics* **12**, 345–350 (1996).

39. Darieva, Z. *et al.* Cell cycle-regulated transcription through the FHA domain of Fkh2p and the coactivator Ndd1p. *Curr Biol* **13**, 1740–1745 (2003).

40. Kõivomägi, M. *et al.* Dynamics of Cdk1 Substrate Specificity during the Cell Cycle. *Mol Cell* **42**, 610–623 (2011).

41. Örd, M. & Loog, M. How the cell cycle clock ticks. *MBoC* **30**, 169–172 (2019).

42. Enserink, J. M. & Kolodner, R. D. An overview of Cdk1-controlled targets and processes. *Cell Div* **5**, 11 (2010).

43. Zhu, G. *et al.* Two yeast forkhead genes regulate the cell cycle and pseudohyphal growth. *Nature* **406**, 90–94 (2000).

44. Spellman, P. T. *et al.* Comprehensive Identification of Cell Cycle–regulated Genes of the Yeast Saccharomyces cerevisiae by Microarray Hybridization. *Mol Biol Cell* **9**, 3273–3297 (1998).

45. Machu, C. *et al.* Spatially distinct functions of Clb2 in the DNA damage response. *Cell Cycle* **13**, 383–398 (2014).

46. Veis, J., Klug, H., Koranda, M. & Ammerer, G. Activation of the G2/M-specific gene CLB2 requires multiple cell cycle signals. *Molecular and cellular biology* **27**, 8364–73 (2007).

47. Maher, M., Cong, F., Kindelberger, D., Nasmyth, K. & Dalton, S. Cell cycle-regulated transcription of the CLB2 gene is dependent on Mcm1 and a ternary complex factor. *Mol Cell Biol* **15**, 3129–3137 (1995).

48. Trcek, T., Larson, D. R., Moldon, A., Query, C. C. & Singer, R. H. Single-Molecule mRNA Decay Measurements Reveal Promoter-Regulated mRNA Stability in Yeast. *Cell* **147**, 1484–1497 (2011).

49. Visintin, R. & Amon, A. Regulation of the mitotic exit protein kinases Cdc15 and Dbf2. *Molecular biology of the cell* **12**, 2961–74 (2001).

50. Hotz, M. & Barral, Y. The Mitotic Exit Network: new turns on old pathways. *Trends in Cell Biology* **24**, 145–152 (2014).

51. Spiesser, T. W., Kühn, C., Krantz, M. & Klipp, E. Bud-Localization of CLB2 mRNA Can Constitute a Growth Rate Dependent Daughter Sizer. *PLOS Computational Biology* **11**, e1004223 (2015).

52. Femino, A. M., Fay, F. S., Fogarty, K. & Singer, R. H. Visualization of single RNA transcripts in situ. *Science* **280**, 585–90 (1998).

53. Raj, A., van den Bogaard, P., Rifkin, S. A., van Oudenaarden, A. & Tyagi, S. Imaging individual mRNA molecules using multiple singly labeled probes. *Nat Methods* **5**, 877–9 (2008).

54. Maekiniemi, A., Singer, R. H. & Tutucci, E. Single molecule mRNA fluorescent in situ hybridization combined with immunofluorescence in S. cerevisiae: Dataset and quantification. *Data in Brief* **30**, 105511 (2020).

55. Tutucci, E. & Singer, R. H. Simultaneous Detection of mRNA and Protein in S. cerevisiae by Single-Molecule FISH and Immunofluorescence. *Methods Mol. Biol.* **2166**, 51–69 (2020).

56. Pichon, X., Robert, M.-C., Bertrand, E., Singer, R. H. & Tutucci, E. New Generations of MS2 Variants and MCP Fusions to Detect Single mRNAs in Living Eukaryotic Cells. *Methods Mol. Biol.* **2166**, 121–144 (2020).

57. Tutucci, E. *et al.* An improved MS2 system for accurate reporting of the mRNA life cycle. *Nat Methods* **15**, 81–89 (2018).

58. Tutucci, E., Vera, M. & Singer, R. H. Single-mRNA detection in living S. cerevisiae using a re-engineered MS2 system. *Nat Protoc* **13**, 2268–2296 (2018).

59. Tutucci, E., Livingston, N. M., Singer, R. H. & Wu, B. Imaging mRNA in vivo, from Birth to Death. *Annu Rev Biophys* **47**, (2018).

60. Zenklusen, D., Larson, D. R. & Singer, R. H. Single-RNA counting reveals alternative modes of gene expression in yeast. *Nat Struct Mol Biol* **15**, 1263–71 (2008).

61. Haimovich, G. *et al.* Use of the MS2 aptamer and coat protein for RNA localization in yeast: A response to 'MS2 coat proteins bound to yeast mRNAs block 5' to 3' degradation and trap mRNA decay products: implications for the localization of mRNAs by MS2-MCP system'. *RNA* **22**, 660–6 (2016).

62. Garcia, J. F. & Parker, R. MS2 coat proteins bound to yeast mRNAs block 5' to 3' degradation and trap mRNA decay products: implications for the localization of mRNAs by MS2-MCP system. *RNA* **21**, 1393–5 (2015).

63. Schlichting, J. K. Modeling synchronization effects in the yeast cell cycle. (2019) doi:http://dx.doi.org/10.18452/19835.

64. Ferrezuelo, F. *et al.* The critical size is set at a single-cell level by growth rate to attain homeostasis and adaptation. *Nat Commun* **3**, 1012 (2012).

65. Johnston, G. C., Pringle, J. R. & Hartwell, L. H. Coordination of growth with cell division in the yeast Saccharomyces cerevisiae. *Experimental cell research* **105**, 79–98 (1977).

66. Leitao, R. M. & Kellogg, D. R. The duration of mitosis and daughter cell size are modulated

by nutrients in budding yeast. *J Cell Biol* **216**, 3463–3470 (2017).

67. Katz, Z. B. *et al.* Mapping translation 'hot-spots' in live cells by tracking single molecules of

mRNA and ribosomes. *Elife* **5**, (2016).

68. Edelmann, F. T. *et al.* Molecular architecture and dynamics of ASH1 mRNA recognition by its

mRNA-transport complex. *Nature Structural & Molecular Biology* **24**, 152–161 (2017).

69. Chartrand, P., Meng, X.-H., Singer, R. H. & Long, R. M. Structural elements required for the

localization of ASH1 mRNA and of a green fluorescent protein reporter particle in vivo. *Current*

Biology **9**, 333–338 (1999).

70. Gonzalez, I., Buonomo, S. B. C., Nasmyth, K. & Ahsen, U. von. ASH1 mRNA localization in

yeast involves multiple secondary structural elementsand Ash1 protein translation. *Current*

Biology **9**, 337–340 (1999).

71. Stueland, M., Wang, T., Park, H. Y. & Mili, S. RDI Calculator: An Analysis Tool to Assess RNA

Distributions in Cells. *Sci Rep* **9**, 1–10 (2019).

72. Heym, R. G. & Niessing, D. Principles of mRNA transport in yeast. *Cell Mol Life Sci* **69**, 1843–

53 (2012).

73. Arava, Y. *et al.* Genome-wide analysis of mRNA translation profiles in Saccharomyces

cerevisiae. *Proc Natl Acad Sci U S A* **100**, 3889–3894 (2003).

74. Niedner, A., Edelmann, F. T. & Niessing, D. Of social molecules: The interactive assembly of

ASH1 mRNA-transport complexes in yeast. *RNA Biol* **11**, 998–1009 (2014).

75. Olivier, C. *et al.* Identification of a conserved RNA motif essential for She2p recognition and

mRNA localization to the yeast bud. *Mol Cell Biol* **25**, 4752–66 (2005).

76. Yang, X., Jost, A. P.-T., Weiner, O. D. & Tang, C. A light-inducible organelle-targeting system

for dynamically activating and inactivating signaling in budding yeast. *MBoC* **24**, 2419–2430

(2013).

77. Talia, S. D., Skotheim, J. M., Bean, J. M., Siggia, E. D. & Cross, F. R. The effects of molecular noise and size control on variability in the budding yeast cell cycle. *Nature* **448**, 947–951 (2007).

78. Soifer, I. & Barkai, N. Systematic identification of cell size regulators in budding yeast. *Mol Syst Biol* **10**, 761 (2014).

79. Spiesser, T. W., Müller, C., Schreiber, G., Krantz, M. & Klipp, E. Size homeostasis can be intrinsic to growing cell populations and explained without size sensing or signalling. *FEBS J* **279**, 4213–4230 (2012).

80. Facchetti, G., Chang, F. & Howard, M. Controlling cell size through sizer mechanisms. *Curr Opin Syst Biol* **5**, 86–92 (2017).

81. Mueller, F. *et al.* FISH-quant: automatic counting of transcripts in 3D FISH images. *Nat Methods* **10**, 277–8 (2013).

82. Seiler, M. *et al.* A consensus motif in ASH1 and further transcripts unifies several RNA motifs required for interaction with the She2p/She3p transport machinery and mRNA localization in yeast. 2024.03.10.584336 Preprint at https://doi.org/10.1101/2024.03.10.584336 (2024).

83. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).

84. Lorenz, R., Hofacker, I. L. & Stadler, P. F. RNA folding with hard and soft constraints. *Algorithms Mol Biol* **11**, 8 (2016).

85. Darty, K., Denise, A. & Ponty, Y. VARNA: Interactive drawing and editing of the RNA secondary structure. *Bioinformatics* **25**, 1974–1975 (2009).

86. Bivand, R. S., Pebesma, E. & Gómez-Rubio, V. *Applied Spatial Data Analysis with R*. (Springer-Verlag, New York, 2013). doi:10.1007/978-1-4614-7618-4.

87. Sprouffske, K. & Wagner, A. Growthcurver: an R package for obtaining interpretable metrics from microbial growth curves. *BMC Bioinformatics* **17**, 172 (2016).

88. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer International Publishing, 2016). doi:10.1007/978-3-319-24277-4.

89. Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).

90. RColorBrewer: ColorBrewer Palettes version 1.1-2 from CRAN. https://rdrr.io/cran/RColorBrewer/.

91. dplyr: A Grammar of Data Manipulation — dplyr-package. https://dplyr.tidyverse.org/reference/dplyr-package.html.

92. Benaglia, T., Chauveau, D., Hunter, D. R. & Young, D. S. mixtools: An R Package for Analyzing Mixture Models. *Journal of Statistical Software* **32**, 1–29 (2010).