Model Selection in Occupancy Models: Inference versus Prediction

Peter S. Stewart[1]*, Philip A. Stephens[1], Russell A. Hill[2], Mark J. Whittingham[3], Wayne

Dawson[1].

[1]Department of Biosciences, Durham University, DH1 3LE, Durham, United Kingdom

[2]Department of Anthropology, Durham University, DH1 3LE, Durham, United Kingdom

[3]School of Natural and Environmental Sciences, Newcastle University, NE1 7RU, Newcastle-

Upon-Tyne, United Kingdom

Keywords:

Akaike's information criterion, Causal inference, Collider bias, Information-theoretic approach,

Model selection, Occupancy models

Running headline:

Model selection in occupancy models

Word count:

7944

<u>Abstract</u>

1. Occupancy models are a vital tool for understanding the patterns and drivers of species

occurrence, but their use requires a method for model selection – choosing between models

with different sets of occupancy and detection covariates. The information-theoretic

approach, which employs information criteria such as Akaike's Information Criterion (AIC)

is arguably the most popular approach for model selection in ecology and is often used for

selecting occupancy models. However, the information-theoretic approach risks selecting

models which produce inaccurate parameter estimates, due to a phenomenon called collider

bias.

2. We aimed to investigate the consequences of collider bias (using an illustrative example

called M-bias) in the occupancy and detection processes of an occupancy model, and to

explore the implications for model selection using AIC and a common alternative, the

Schwarz Criterion (or Bayesian Information Criterion, BIC).

3. We simulated datasets using known parameter values and fitted models with different

combinations of covariates. We then compared the models on how accurately they estimated

the effect of a focal covariate on the occupancy probability, the accuracy of their site-level

occupancy predictions, and their level of support from AIC and BIC.

4. When M-bias was present in the occupancy process, AIC and BIC selected models which

inaccurately estimated the effect of the focal occupancy covariate, while simultaneously

producing more accurate predictions of site-level occupancy probability. In contrast, M-bias

in the detection process did not impact the estimate; all models made accurate inferences,

while the site-level predictions of the AIC/BIC-best model were slightly more accurate.

5.   Our results emphasise the importance of distinguishing between parameter inference and

prediction as separate tasks in ecological modelling. In the context of occupancy models, our

findings suggest that information criteria can be used to select occupancy covariates if the

purpose of the model is prediction, but not if the purpose is to understand how environmental

variables affect occupancy. In contrast, detection covariates can usually be selected using

information criteria regardless of the model's purpose. More generally, our results support the

view that information criteria should not be used to compare different biological hypotheses

in observational studies.

Introduction

The patterns and drivers of species occurrence are of fundamental interest to ecologists. Predicting where species occur enables ecologists to tackle key problems such as understanding the spread of invasive species (Gormley *et al.* 2011), assessing the efficacy of protected areas (Midlane *et al.* 2014), and evaluating the extinction risk (Breiner & Bergamini 2018) and recovery of populations and species (Akçakaya *et al.* 2018). Understanding the drivers of occurrence is also important; we must diagnose factors which threaten species to mitigate them through informed interventions (Caughley 1994). Many studies have aimed to determine how occurrence is driven by factors including forest degradation (Zimbres *et al.* 2018), agricultural expansion (Semper-Pascual *et al.* 2020), wildfires (Hossack *et al.* 2013), and anthropogenic noise pollution (Chen & Koprowski 2015).

A key challenge in studying the patterns and drivers of occurrence is that ecologists are often constrained to the use of observational data; experimental manipulations of ecological systems may be physically impossible, logistically unfeasible, or unethical. Consequently, one approach is to use a model which relates observed variation in species occurrence to one or more environmental covariates. The model can then be used to predict species occurrence at new sites, or to examine the effect of each covariate on occurrence (Shmueli 2010). Occupancy models are often used because they deal with imperfect detection (MacKenzie *et al.* 2002, 2006). They do so by modelling the probability that a species occupying the site is detected, often including environmental covariates to explain variation in detectability among sites (MacKenzie *et al.* 2002, 2006). Occupancy models therefore contain one set of covariates for occupancy probability, and a second set for detection probability; the challenge is to select suitable sets of covariates to include in the model. This challenge can also be framed as a problem of model selection (Robins & Greenland 1986; Buckland *et al.* 1997; Forster 2000; Burnham & Anderson 2004; Johnson & Omland 2004).

One approach to model selection is the information-theoretic approach (Anderson *et al.* 2000; Burnham & Anderson 2001, 2004; Lukacs *et al.* 2007; Burnham *et al.* 2011), which compares models in terms of their relative Kullback-Leibler (KL) divergence – the relative distance between each model and "full reality", in units of information entropy (Forster 2000; Burnham & Anderson 2001;

McElreath 2021). Information criteria, of which Akaike's information criterion (AIC; Akaike 1973) is

the most commonly used, estimate the relative KL divergence of each model using the sample data

(McElreath 2021). AIC is calculated for each model by taking the in-sample deviance (a measure of

how well the model fits the data), and adding an overfitting penalty term proportional to the number

of parameters in the model (Akaike 1973; Burnham *et al.* 2011). Consequently, AIC favours

parsimonious models which balance underfitting and overfitting, with the aim of producing better out-

of-sample predictions (McElreath 2021).

Some proponents of the information-theoretic approach argue that each model in the candidate set

should represent a different biological hypothesis, and that the models' relative AIC scores indicate

the strength of evidence for each hypothesis (Burnham *et al.* 2011). However, insights from the field

of causal inference reveal a potential problem with this approach – collider bias. Collider bias is a

form of confounding, where the estimate of an effect is biased due to some feature of the data-

generating process (Greenland 2003; Greenland & Mansournia 2015). However, unlike the classical

notion of confounding, collider bias arises not from omitting an important variable from the statistical

model (omitted variable bias; Clarke 2005), but is instead a form of included variable bias (or "bad

control"; Cinelli *et al.* 2020; McElreath 2021). In the simplest case, collider bias arises from including

a variable which is a function of both the explanatory and response variable of interest (Greenland

2003). For example, in spotted hyenas (*Crocuta crocuta*), reproductive state is likely a function of

both social connectedness and immune function – therefore, including reproductive state as a

covariate will induce a spurious association between social connectedness and immune function

(Laubach *et al.* 2021). In more complex examples with more than three variables, collider bias can

arise from the relationships between variables without the response variable directly affecting any

other variable in the system (Fig. 1).

As AIC and other information criteria select models based on their expected predictive performance,

they will not necessarily select models which return accurate parameter estimates (McElreath 2021).

For example, Luque-Fernandez *et al.* (2019) demonstrated that for a simple collider example,

including the collider covariate improved the model's AIC score, while simultaneously resulting in an

estimated effect size which was far from the true value. It is straightforward to extend their simulation

to show that this result is consistent across a broad range of effect sizes (Fig. 2). However, the

implications for model selection in occupancy modelling, in which covariates must be selected for

both the occupancy and detection sub-models, have yet to be investigated.

To address this topic, we investigated the consequences of a form of collider bias known as "M-bias"

(Fig. 3 ; Greenland 2003; Cinelli *et al.* 2020) in an occupancy modelling framework, and explored the

implications for model selection using the information-theoretic approach (using AIC). We also

examined the performance of a common alternative to AIC, the Schwarz criterion (or Bayesian

Information Criterion, BIC; Schwarz 1978), which can be used for selecting the "true" model from the

candidate set according to some authors (Aho *et al.* 2014). We focused on M-bias because it is a

common illustrative example in the causal inference literature (e.g. Greenland 2003; Cinelli *et al.*

2020), which demonstrates several important points: i) conditioning on a variable may cause more

harm than omitting it from the model (Greenland *et al.* 1999), ii) collider bias can occur even when no

variables are caused by the response variable, and iii) the importance of considering potential latent

(unobserved) variables when drawing inferences. In our simulation-based approach, we generated

datasets where M-bias was present in the occupancy process, the detection process, or both. We then

fitted occupancy models with different sets of covariates to these datasets, and evaluated them on the

accuracy of their parameter inferences, the accuracy of their site-level occupancy predictions, and

their level of support from AIC and BIC.

Figure 1. Directed acyclic graph showing a hypothetical example in which we are interested in

estimating the direct effect of the density of an invasive plant on animal occupancy (red arrow).

Invasive plant density also influences the density of a native food plant, which in turn influences the

animal's occupancy. Soil properties influence the density of both the native and invasive plant – these

properties are latent (unobserved), and so are shown in a circle. Urbanisation influences the densities

of both plant species, as well as animal occupancy – here, urbanisation is also latent. In this example,

native food plant density is a collider which sits on the backdoor path from invasive plant density to

animal occupancy, *Invasive plant ← (Soil properties) → Native food plant ← (Urbanisation) →*

Animal occupancy. Including native food plant density in the model opens the backdoor path, biasing

the estimated direct effect of the invasive plant. Notably, in this example the effect of interest cannot

be estimated by including any of the available covariates – measuring and including urbanisation is

necessary.

Figure 2. Luque-Fernandez *et al.* (2019) presented simulations illustrating classical confounding and collider bias in a simple linear model. We extended their example by conducting 10000 simulation iterations for each example, with each iteration using true effect sizes drawn from a uniform distribution between -2 and 2 to ensure that the results are not specific to the choice of parameter values. In the classical confounding example (**A**), including the variable Z reduces the absolute bias (the absolute difference between the estimated and true effect: $|\hat{\beta}_{XY} - \beta_{XY}|$) when estimating the effect of X on Y. Conversely, in the collider example (**B**), including Z increases the absolute bias. However, in both cases Akaike's Information Criterion (AIC) favours the model which includes Z (**C**), illustrating that AIC does not always favour models which produce accurate parameter estimates. R code to reproduce the simulations is available at: https://zenodo.org/badge/latestdoi/462801230

Methods

Causal inference, directed acyclic graphs and M-bias: a primer

Causal inference is concerned with predicting the consequences of intervening in a system, as well as

inferring counterfactual outcomes – events which might have happened, under hypothetical unrealised

conditions (Pearl 2009; Pearl *et al.* 2016). Importantly, causal inference is not about 'inferring

causation from correlation' – conclusions about causality cannot be made from the data alone, but

require causal assumptions about the process which generated the data (Pearl 2009; Pearl *et al.* 2016).

These assumptions, rather than remaining implicit, should be communicated clearly so that they are

open to scrutiny, debate, sensitivity analysis, and verification (Pearl 2009).

Causal assumptions are typically expressed in a graphical model, usually a directed acyclic graph

(DAG) where the nodes represent variables and the edges represent the relationships between them

(Pearl 1995). The nodes on a DAG can represent both observed and latent variables – the latter are

usually shown surrounded by a circle (e.g., A and C in Fig. 3a), and we show them surrounded by

brackets in the main text. The edges linking each node are directional, and are thus represented as

arrows; these arrows represent the mechanistic links between the variables (Greenland *et al.* 1999).

The succession of edges linking one variable to another, regardless of which direction these edges are

pointing in, is known as a path (Pearl 1995). For instance, Fig. 3a has two paths connecting X and ψ:

$X \rightarrow \psi$, and $X \leftarrow (A) \rightarrow D \leftarrow (C) \rightarrow \psi$.

A DAG corresponds to a non-parametric structural equations model (Pearl 1995), where each node is

represented as a function of the variables which have edges going into it, as well as a disturbance term

(ε) which represents the effect of omitted variables which are assumed to be independent of one

another (Pearl 1995). For example, the DAGs in Fig. 3c correspond to the following set of equations:

$$A = f_A(\varepsilon_A), \quad C = f_C(\varepsilon_C), \quad D = f_D(A, C, \varepsilon_D), \quad X = f_X(A, \varepsilon_X), \quad \psi = f_\psi(X, \varepsilon_\psi),$$

$$Q = f_Q(\varepsilon_Q), \quad V = f_V(\varepsilon_V), \quad R = f_R(Q, V, \varepsilon_R), \quad U = f_U(A, \varepsilon_U), \quad P = f_P(U, \varepsilon_P)$$

Importantly, the model is non-parametric in the sense that the form of these functions does not have to

be specified (Pearl 1995; Greenland *et al.* 1999), allowing DAGs to accommodate non-linear

functions and interactions. For example, the variable D in the above equation could be modelled as a

nonlinear function of A and C, or a linear interaction between A and C.

Once a DAG has been specified, it is analysed to identify the set of variables necessary to remove

structural confounds from the effect of interest. One strategy is to condition on (i.e., include as

covariates) the variables which satisfy the "back-door criterion", in which the aim is to "close all

back-door paths" linking the explanatory and response variable of interest (Pearl 1995). A back-door

path is defined as any path which has an arrow entering the focal explanatory variable (Pearl 1995) –

for example, in Fig. 3a the path $X \leftarrow (A) \rightarrow D \leftarrow (C) \rightarrow \psi$ is a back-door path.

By default, paths which are a "fork" (e.g., $X \leftarrow Z \rightarrow Y$) or "pipe" (e.g., $X \rightarrow Z \rightarrow Y$) are open, and

conditioning on the middle variable (Z) closes them (Greenland 2003; Pearl *et al.* 2016; McElreath

2021). Paths which are a collider (e.g., $X \rightarrow Z \leftarrow Y$) are closed by default, and conditioning on the

middle variable (Z) opens the path (Greenland *et al.* 1999; Greenland 2003; Pearl *et al.* 2016;

McElreath 2021). A back-door path only needs to be closed in one place; for instance, $X \leftarrow (A) \rightarrow D$

$\leftarrow (C) \rightarrow \psi$ is closed because it is blocked by the collider at D, irrespective of the forks at A and C.

The DAG displayed for ψ in Fig. 3a, for example, contains an "M"-shaped structure, which can

induce a type of confounding known as M-bias (Greenland 2003). By conditioning on the collider (D

in Fig. 3a), the backdoor path is opened and the effect of interest (X on ψ in Fig. 3a) is confounded.

When the back-door path contains latent variables (A and C in Fig. 3a) it is not possible to condition

on them to close the path again because they are unobserved by definition. If the objective is to obtain

an unconfounded estimate of the effect of X on ψ, then the correct approach is to avoid conditioning

on D.

Simulation study

To explore the effects of M-bias in both the occupancy and detection components of an occupancy

model, we conducted simulations of three different scenarios (Fig. 3) in which the target for inference

was the effect of variable X on occupancy probability (ψ). In the first scenario (Fig. 3a), ψ was part of

an M-graph while the detection probability (p) was fixed at 50%. In the second scenario (Fig. 3b), ψ

depended only on X, and p was now part of an M-graph. In the final scenario (Fig. 3c), both the

occupancy and detection probabilities were part of M-graphs.

All three simulations followed the same process: **1)** generate a dataset with known parameter values,

using the relationships between variables embodied in the relevant DAGs (Fig. 3); **2)** fit a number of

occupancy models to the dataset (Table 1); **3)** evaluate each model's accuracy in parameter estimation

and prediction; **4)** evaluate each model's quality under the information-theoretic framework. Each

simulation was repeated 1000 times. We conducted our simulations in R (v 4.0.5; R Core Team,

2021), and provide code to reproduce our simulations and analyses at:

https://zenodo.org/badge/latestdoi/462801230

1) Generating a dataset

Data were simulated for 3000 sites with 40 surveys each. These values were deliberately high to

ensure that any inaccuracy was not primarily driven by an underpowered design.

We first drew effect sizes for each arrow in the relevant DAG from a uniform distribution between -1

and 1. Values for explanatory variables with no ingoing arrows on the DAG were then drawn from a

normal distribution with a mean of 0 and standard deviation of 1. We then generated values for the

other explanatory variables from the appropriate variables and effect sizes (i.e., those from ingoing

arrows on the DAG; Fig. 3), plus a disturbance term drawn from a normal distribution with a mean of

zero and standard deviation of 0.025. The disturbance term was required to avoid perfect collinearity

between the explanatory variables which would result in model failure.

We then generated the true occupancy and detection probabilities. The log-odds of occupancy and

detection were first calculated as a linear function of the explanatory variables with ingoing arrows on

the DAG (Fig. 3), and the inverse-logit was then taken to obtain the probability. The true occupancy

state of each site was then simulated as a Bernoulli trial with probability of success equal to the

occupancy probability. We assumed that the true occupancy state remained constant for each site,

which is a key assumption of single-season occupancy models (MacKenzie *et al.* 2002).

Detection histories were then generated for each site, as 40 Bernoulli trials with probability of success

equal to the true occupancy state multiplied by the detection probability. This means that the species

was never observed at sites in which it was absent (occupancy state = 0), while when the species was

present (occupancy state = 1) it had a chance of being observed on each visit, with probability equal to

the detection probability. The detection histories for each site represent the observed data that would

be collected by an ecologist in the field, alongside the covariate values for the non-latent variables.

Figure 3. Directed acyclic graphs (DAGs) showing the relationship between variables in the

occupancy and detection processes of the three simulation scenarios: **a)** scenario 1 (M-bias in the

occupancy process), **b)** scenario 2 (M-bias in the detection process), **c)** scenario 3 (M-bias in both the

occupancy and detection processes). ψ represents the probability that a site is occupied by a species

(occupancy probability), while p represents the probability that a species is detected given that it is

present at a site (detection probability). X represents an explanatory variable for which we are

interested in estimating the effect on the occupancy probability. A, C, D, Q, U, R and V represent

additional variables which are thought to be important in the system. Circled variables are latent

(unobserved).

2)Fitting models

Occupancy models were fitted to each dataset using the *occu* function in the R package *unmarked*

(v.1.0.0; Fiske & Chandler, 2011), which implements the single-season occupancy model developed

by MacKenzie and colleagues (2002). The models used the logit link function because we used the

inverse-logit to simulate the data. We fitted models with various combinations of observed variables

(i.e., excluding latent variables) for each scenario (Table 1). All models included the focal covariate

for occupancy (X). In scenario 1, detection probability was modelled as an intercept only, as it did not

depend on any covariates.

Table 1. Occupancy models fitted in each simulation scenario. All models also included an intercept

term for both occupancy and detection. X represents an explanatory variable for which we are

interested in estimating the effect on the occupancy probability. D, U, and R represent additional

variables which are thought to be important in the occupancy or detection processes.

Model	Occupancy covariates	Detection covariates
Scenario 1		
Model 1	X	n/a
Model 2	X + D	n/a
Scenario 2		
Model 1	X	U
Model 2	X	U + R
Scenario 3		
Model 1	X	U
Model 2	X + D	U
Model 3	X	U + R
Model 4	X + D	U + R

307   *3) Evaluating Model Performance*

308   In each scenario, all models were evaluated on the accuracy of their parameter inferences and

309   predictions. To quantify how accurately each model estimated the effect of covariate X on the

310   occupancy probability ψ, we calculated the bias and absolute bias:

311
$$Bias = \hat{\beta}_{X\psi} - \beta_{X\psi}$$

312
$$Absolute\ Bias = |\hat{\beta}_{X\psi} - \beta_{X\psi}|$$

313   Where $\hat{\beta}_{X\psi}$ and $\beta_{X\psi}$ are the estimated and true effects of X on ψ, respectively. Additionally, we

314   constructed 95% confidence intervals for $\hat{\beta}_{X\psi}$ as follows:

315
$$95\%\ CI = \hat{\beta}_{X\psi} \pm 1.96 \times SE$$

where SE is the standard error of the $\hat{\beta}_{X\psi}$ estimate, obtained from the model summary table. We then

checked whether the true value, $\beta_{X\psi}$, was found within this interval. We also checked whether the

sign (positive or negative) of $\hat{\beta}_{X\psi}$ was the same as that of $\beta_{X\psi}$.

To evaluate each model's predictions, we used the *predict* function in R to make predictions for the

occupancy probability value for each site in two datasets. We first made predictions for the data to

which the model was fitted, to examine how the model retrodicted the sample. We then examined the

model's performance in out-of-sample prediction by making predictions for a new dataset (also 3000

sites with 40 surveys each), which was generated using the same true parameter values as the original

dataset. This corresponds to the common task in which occupancy models are fitted to one set of data

(e.g., grid squares sampled from a region) and used to extrapolate to a wider set of data which are

assumed to arise from the same data-generating process (e.g., other grid squares in the region).

To assess the accuracy of the model retrodictions and predictions, we first calculated the error for

each site:

329
$$Error = \hat{\psi}_i - \psi_i$$

330   where $\hat{\psi}_i$ and ψ_i are the estimated and true occupancy probabilities for site i, respectively. To

331   summarise this error over all sites in the dataset, we calculated the mean absolute error:

332
$$\text{Mean absolute error} = \frac{1}{n}\sum_{i=1}^{n}|\hat{\psi}_i - \psi|$$

where n is the number of sites. Additionally, we checked whether the true occupancy probability for

each site was within the 95% confidence interval obtained from the prediction, and calculated the

proportion of sites for which this was the case.

4) Evaluating models under the information-theoretic framework

To examine the degree of support for each model under the information-theoretic framework we

obtained the AIC value for each model m from the model's summary table. We then calculated the

ΔAIC value for each model in the scenario as:

341
$$\Delta AIC_m = AIC_m - minAIC$$

where AIC_i is the AIC value for model m, and *minAIC* is the lowest AIC value for the set of models in

the scenario.

Proponents of the information-theoretic approach have advocated for multimodel inference (e.g.,

Anderson *et al.* 2000; Burnham & Anderson 2004; Burnham *et al.* 2011), in which inferences are

made using the entire candidate set of models, each weighted using Akaike weights derived from

AIC. This approach accounts for uncertainty in the model selection process (Burnham & Anderson

2004). We calculated Akaike weights (w) for each model m as:

349
$$w_m = \frac{\exp\left(-0.5 \times \Delta AIC_i\right)}{\sum_{r=1}^{R}\exp\left(-0.5 \times \Delta AIC_r\right)}$$

350   where R is the number of models in each scenario.

351   We also considered the Schwarz criterion (BIC; Schwarz 1978) as an alternative to AIC. BIC is

352   defined as:

353
$$BIC_m = -2\ln\left(\hat{L}\right) + j\ln(n)$$

where $\hat{L}$ is the maximum likelihood estimate and thus $-2\ln(\hat{L})$ the in-sample deviance, j is the

number of parameters, and n is the number of observations in the dataset. We calculated BIC, ΔBIC,

and BIC weight values for each model using the R package *AICcmodavg* (Mazerolle, 2020). ΔBIC

and BIC weights are defined as for ΔAIC and AIC weights, but using the BIC values for each model.

<u>Results</u>

Scenario 1: M-Bias in the Occupancy Process

When M-bias was present in the occupancy process, model 1 ($\psi \sim X$) estimated the true effect of X on

ψ much more accurately than model 2 ($\psi \sim X + D$) (Table 2, Fig. 4a,b). However, comparing the two

models' predictive accuracy showed the opposite picture; model 1 generally produced worse

predictions than model 2 (Table 2, Fig. 5a,b). The models' retrodictive accuracy was also qualitatively

similar to their predictive accuracy (Fig. S1). AIC and BIC both showed clear support for model 2 in

the majority of simulations (Fig. 6a,b); in 85.1% of simulations model 2 received an Akaike weight of

>0.99, and in 63.0% of simulations it received the entire weight (Fig. 6b). The few simulations in

which model 1 received more Akaike weight were mostly those in which $\beta_{C\psi}$ was small (Fig. S2). A

similar pattern of results was observed for BIC (Fig. S2), although when BIC assigned weight to

model 1 it generally assigned more weight than AIC (Fig. 6a).

Scenario 2: M-Bias in the Detection Process

When M-Bias was present in the detection process, both models 1 ($\psi \sim X, p \sim U$) and 2 ($\psi \sim X, p \sim U$

$+ R$) accurately estimated the effect of X on ψ (Table 2, Fig. 4c,d). Both models also made very

accurate predictions (Table 2, Fig. 5c,d), although model 2 performed slightly better - in 86.4% of

simulations, the true occupancy probability was contained within model 2's 95% confidence interval

for every predicted site, whereas model 1 accomplished this in 85.8% of simulations. Similar results

were observed for the models' retrodictive accuracy (Fig. S1c,d). Both AIC and BIC assigned the

majority of weight to model 2 in most simulations (Fig. 6c,d).

Scenario 3: M-Bias in the Occupancy and Detection Processes

When M-Bias was present in both the occupancy and detection processes, models 1 ($\psi \sim X, p \sim U$)

and 3 ($\psi \sim X, P \sim U + R$) estimated the effect of X on ψ much more accurately than models 2 ($\psi \sim X$

$+ D, P \sim U$) and 4 ($\psi \sim X + D, P \sim U + R$) (Table 2, Fig. 4e-h). In general, the 95% confidence

interval around the estimate in models 2 and 4 only contained the true value when β_{AD} and $\beta_{C\psi}$ (and to

a lesser extent β_{AX}) were relatively small (Figs. S4,5). In contrast, models 2 and 4 made more accurate

predictions than models 1 and 3 (Table 2, Fig. 5e-h), and similar results were obtained for retrodictive

accuracy (Fig. S1e-h). Both AIC and BIC showed clear support for model 4 in the majority of

simulations (Fig. 6h); the model received an Akaike weight of >0.99 in 82.2% of the simulations.

While model 3 did occasionally receive weight, this mostly occurred when $\beta_{C\psi}$ was small (Fig. S6)

and the model still never received the entire weight (Fig. 6g). BIC weights were similar to the Akaike

weights, although BIC assigned more weight to model 3 in some simulations (Fig.4g). Again, these

instances were generally when $\beta_{C\psi}$ was small (Fig. S6).

Table 2. Inferential and predictive accuracy of occupancy models across all simulations (n=1000) for each of the three scenarios: M-bias in the occupancy process only (left), M-bias in the detection process only (centre), and M-bias in both the occupancy and detection processes (right). Each model is shown as a separate column, with the occupancy *(ψ)* and detection *(p)* covariates displayed at the top. Each row indicates a separate metric summarising the models' performance across the simulations. Full definitions of each metric can be found in the main text.

			M-bias in occupancy process		M-bias in detection process		M-bias in occupancy and detection processes			
			$\psi \sim X$	$\psi \sim X+D$	$\psi \sim X$ $p \sim U$	$\psi \sim X$ $p \sim U+R$	$\psi \sim X$ $p \sim U$	$\psi \sim X+D$ $p \sim U$	$\psi \sim X$ $p \sim U+R$	$\psi \sim X+D$ $p \sim U+R$
Inference	Bias	Q1	-0.07	-1.13	-0.03	-0.03	-0.07	0.93	-0.07	0.93
		Mean	-0.001	0.03	<0.001	<0.001	0.04	0.19	0.05	0.19
		Q3	0.06	0.88	0.03	0.03	0.06	0.99	0.06	0.99
	Absolute bias	Q1	0.03	0.29	0.01	0.01	0.03	0.32	0.03	0.32
		Median	0.07	1.00	0.03	0.03	0.06	0.95	0.06	0.95
		Q3	0.13	2.69	0.05	0.05	0.13	2.47	0.13	2.47
	True value in 95% C.I.		89.6%	20.07%	95.4%	95.4%	91.5%	20.2%	91.6%	20.1%
	Sign correct		94.5%	65.1%	98.3%	98.3%	93.6%	63.3%	93.6%	63.3%
Prediction	Mean absolute error	Q1	0.048	0.001	0.006	0.006	0.047	0.010	0.047	0.010
		Median	0.092	0.015	0.009	0.009	0.087	0.014	0.087	0.014
		Q3	0.128	0.022	0.012	0.012	0.126	0.021	0.126	0.021
	Proportion of sites with true ψ in 95% C.I.	Q1	10.6%	65.4%	100%	100%	10.8%	69.0%	10.8%	69.0%
		Median	15.4%	89.0%	100%	100%	16.2%	90.1%	16.2%	90.2%
		Q3	29.4%	99.9%	100%	100%	30.2%	99.9%	30.1%	99.9%

Figure 4. True versus estimated effect of *X* on occupancy probability (ψ), for the following

occupancy models: **A)** scenario 1, model 1; **B)** scenario 1, model 2; **C)** scenario 2, model 1; **D)**

scenario 2, model 2; **E)** scenario 3, model 1; **F)** scenario 3, model 2; **G)** scenario 3, model 3; **H)**

scenario 4, model 4. Each point represents the result from one simulation, with 1000 simulations in

total. Note that the y-axis is truncated at -10 and 10; plots B, F, and H omit 40, 38 and 38 points

respectively which lay outside this range. Blue points indicate that the true value was contained within

the estimate's 95% confidence interval, while unfilled circles indicate that the true value was not

contained within the interval. Dashed black line indicates equality between the true and estimated

effect – increased vertical distance from this line indicates a more biased estimate. Each model's

covariates for ψ and the detection probability (*p*) are shown above their respective plot. For

explanation of the scenarios, see Table 1 and main text.

Figure 5. Kernel density estimate contours showing two measures of predictive accuracy when predicting site-level occupancy probability (ψ), for 1000 simulations. The x-axis shows the proportion of sites (out of 3000 sites) for which the true occupancy probability was contained within the 95% confidence interval around the model's prediction. The y-axis shows the mean absolute error of the predictions. Consequently, the bottom right of each plot indicates higher predictive accuracy, while the top left indicates lower predictive accuracy. The density of simulations within this area is shown by the coloured contours, with lighter colours indicating a higher density of simulations. Results are displayed for the following occupancy models: **A)** scenario 1, model 1; **B)** scenario 1, model 2; **C)** scenario 2, model 1; **D)** scenario 2, model 2; **E)** scenario 3, model 1; **F)** scenario 3, model 2; **G)** scenario 3, model 3; **H)** scenario 4, model 4. Each model's covariates for ψ and the detection probability (p) are shown above their respective plot. For explanation of the scenarios, see Table 1 and main text.

Figure 6. Akaike weight (yellow area) and BIC weight (blue area) for 1000 simulations of eight

occupancy models. Simulations are shown ranked by weight, with higher Akaike and BIC weights

shown on the right. Higher weights correspond to lower AIC/BIC values, indicating more support for

the model. The panels display: **A)** scenario 1, model 1; **B)** scenario 1, model 2; **C)** scenario 2, model

1; **D)** scenario 2, model 2; **E)** scenario 3, model 1; **F)** scenario 3, model 2; **G)** scenario 3, model 3; **H)**

scenario 4, model 4. Each model's covariates for the occupancy probability (ψ) and the detection

probability (p) are shown above their respective plot. For explanation of the scenarios, see Table 1

and main text. Dashed horizontal lines are shown for weights of 0, 0.5, and 1.

Discussion

Occupancy models are essential for studying the patterns and drivers of species occurrence, but their

use requires a method for choosing between models with different covariates for occupancy and

detection. The information-theoretic approach, which uses information criteria like AIC, is a popular

approach for model selection. However, it runs the risk of selecting models which produce inaccurate

parameter estimates because of collider bias. Here, we investigated the consequences of M-bias (a

specific form of collider bias) for occupancy modelling, and explored the implications for model

selection using AIC and BIC. In our simulations, we observed that when M-bias was present in the

occupancy process, AIC and BIC favoured a model which produced a highly inaccurate estimate of

the focal effect but produced more accurate predictions and retrodictions of the site-level occupancy

probability. In contrast, M-bias in the detection process did not result in inaccurate estimates of the

focal effect – both models made accurate inferences. Both models also made accurate predictions and

retrodictions, although those of the AIC/BIC-best models were slightly better. The same results were

observed when M-bias was present in both the occupancy and detection processes: the model

favoured by AIC and BIC produced inaccurate inferences but more accurate predictions, while

models made similarly accurate inferences regardless of M-bias in the detection process. These results

have important implications for model selection in occupancy models, as well as for how the

information-theoretic approach is applied in ecological modelling more generally.

Information criteria select models which produce poor parameter inferences, but good predictions

When M-bias was present in the occupancy process, the model which received the greatest support

from AIC and BIC produced highly inaccurate estimates of the effect of the variable of interest *(X)* on

the occupancy probability *(ψ)*. Such biased estimates are not informative about the biological drivers

underlying the observed pattern, nor do they accurately predict the consequences of intervening in the

system. To illustrate the importance of this result, imagine that the fourth iteration of the first

scenario, in which M-bias is present in the occupancy process only, represents a real ecological study

where the objective is to inform management to increase the occupancy of a species. In this example,

model 2 ($\psi \sim X + D$) received Akaike and BIC weights of one – it is clearly the "best" model,

according to these predictive criteria. The model estimated the effect of X on ψ as -1.22 ($\pm$ 0.12

standard error) – a strong negative effect on occupancy. However, this effect is confounded; in reality,

the effect of X on ψ is strongly positive (0.72). Furthermore, the estimated effect of D on ψ is also

strongly negative (-1.70 $\pm$ 0.09), when D actually has zero effect on occupancy. In this example, the

AIC/BIC "best" model suggests that to increase ψ, we should intervene to reduce X and D. In fact, this

would reduce occupancy, as well as waste resources managing the unrelated variable D. If applied to

a real conservation problem where occupancy models are important (e.g. Hossack *et al.* 2013; Chen &

Koprowski 2015; Zimbres *et al.* 2018; Semper-Pascual *et al.* 2020), the results could be disastrous.

Such poor inferences were the norm in our simulations; confounded models were only able to

estimate the direction of the focal effect correctly in 65.1% of cases at best – little better than the

accuracy we would expect from guessing.

While the models supported by AIC and BIC produced biased parameter estimates, they also

produced more accurate predictions and retrodictions of the occupancy probability at each site. This is

because these models include the variable D which has an open path to ψ; including D provides

additional information about the variation in ψ, improving prediction. However, because D is a

collider, including it opens the back-door path from X to ψ, biasing the estimated effect of X. From the

perspective of AIC and BIC, including D results in a reduced in-sample deviance which typically

outweighs the penalty for adding the additional variable; this reduction must be greater to outweigh

BIC's larger penalty term, which is why BIC was more conservative in its tendency to select

confounded models in our simulations (Fig. 6). This also explains why AIC and BIC tended to prefer

the non-confounded model (omitting D) when $\beta_{C\psi}$ was close to zero (Figs. S2, S6); the near-zero

effect of C on ψ meant that the path from D to ψ through C was almost blocked (the other path from D

to ψ was blocked by conditioning on X), and therefore D explained relatively little variation in ψ.

This mechanism also explains why M-bias in the detection process did not affect inferences about the

effect of X on ψ. In this scenario, including the variable R explains additional variation in the

detection probability, allowing the model to better account for imperfect detection when estimating

the occupancy probability. As the detection probability is generally regarded as a nuisance parameter

(Karavarsamis 2015), it is inconsequential that the effect of the other detection covariate (U) will be

confounded.

Furthermore, this mechanism explains why information criteria prefer confounded models in

scenarios other than collider bias. For example, simulations by McElreath (2021) showed that

information criteria tend to select models which condition on the mediator (M) in a pipe (e.g., $X \rightarrow M$

$\rightarrow \psi$), inducing post-treatment bias (Rosenbaum 1984). Information criteria prefer models which

include M because M has an open path to ψ, so including M explains additional variation in ψ.

However, including M also confounds the estimated effect of X – not by opening a non-causal back-

door path as in the collider example, but instead by closing the causal front-door path which runs

through M. We also expect information criteria to select confounded models in other scenarios, such

as case control bias (Cinelli *et al.* 2020).

Inference and prediction are separate tasks

The key point supported by our results is that inference and prediction are separate tasks which should

not be conflated in model selection (Shmueli 2010; Laubach *et al.* 2021; McElreath 2021). We echo

Gelman and Rubin's (1995) criticism of selecting "a model that is adequate for specific purposes

without consideration of those purposes". In the context of occupancy models, both explanation and

prediction are important objectives, and conflating the two does justice to neither. Furthermore, our

results emphasise the importance of considering not only the model's purpose, but also the purpose of

sub-models within the model; the purpose of the occupancy sub-model depends on whether we are

interested in predicting the occupancy state or inferring its drivers, while the detection sub-model's

purpose is almost always prediction of the detection probability. Consequently, how occupancy

covariates are chosen depends on the purpose of the model – information criteria are suitable if the

purpose of the model is prediction, but not if the purpose is parameter inference – while detection

covariates can generally be selected using information criteria. This advice also applies to other

predictive model selection methods such as cross-validation; the choice of model made by AIC is

asymptotically equivalent to that made by leave-one-out cross validation (Stone 1977).

Information criteria should not be used to compare biological hypotheses in observational studies

The importance of distinguishing between inference and prediction has wider implications for how

information-theoretic model selection is applied in ecology. Proponents of the information-theoretic

approach have argued that it can be used to compare multiple *a priori* specified models, each

representing a different biological hypothesis, with the relative AIC scores indicating the strength of

evidence for each hypothesis (Johnson & Omland 2004; Richards 2005; Burnham *et al.* 2011).

However, using information criteria in this way conflates inference and prediction; information

criteria select models which make better predictions, but these same models can contain spurious

effect sizes which hold no biological meaning, while the effects of biologically important covariates

are confounded. This is not only the case for occupancy models; the occupancy models we employed

are just an extension of logistic regression (Clark & Altwegg 2019), and these points apply to other

forms of linear model as well (Luque-Fernandez *et al.* 2019; McElreath 2021). The implication is that

information-theoretic model selection should not be used to compare biological hypotheses in

observational studies.

The information-theoretic approach and causal inference are complementary

While we argue that the information-theoretic approach should not be used to compare biological

hypotheses in observational studies, we must emphasise that we are not arguing that the information-

theoretic approach is flawed or useless for model selection. As information criteria select models from

the "predictive point of view" (Akaike 1998), while causal inference is concerned with estimating the

effects of covariates, we see the two approaches as complementary.

We also argue that causal inference and the information-theoretic approach share many philosophical

underpinnings. Proponents of the information-theoretic approach emphasise the importance of subject

expertise and *a priori* "hard thinking" to develop hypotheses which are then compared as models

(Lukacs *et al.* 2007; Burnham *et al.* 2011). In causal inference, subject expertise and *a priori* thought

are vital in making the causal assumptions which are embodied in the DAG (Pearl 1995; Greenland *et*

al. 1999). Causal inference therefore provides a framework to support the "hard thinking" in

ecological modelling (Grace & Irvine 2020). Proponents of the information-theoretic approach also

recognise that "a proper analysis must consider the science context and cannot successfully be based

on 'just the numbers'" (Burnham & Anderson 2004). Similarly, proponents of causal inference argue

that conclusions cannot be drawn from the data alone, but require causal assumptions which come

from the scientific context of the model (Pearl 2009; Pearl *et al.* 2016).

Chamberlin's (1890) method of multiple working hypotheses is often emphasised in the information-

theoretic approach (Burnham & Anderson 2004; Elliott & Brook 2007). We argue that causal

inference is very compatible with Chamberlin's method; constructing a causal model forces us to

consider multiple explanations for a phenomenon, guarding against the threat of "parental affection

for a favourite theory" which concerned Chamberlin. Furthermore, the causal approach does not rely

on the comparison of a model with a null model or hypothesis, a feature of the multiple working

hypothesis approach which is regarded as beneficial (Anderson *et al.* 2000; Burnham & Anderson

2004). Due to the relatively static nature of causal models, we argue they are especially suited to the

case of multiple working hypotheses in parallel (Elliott & Brook 2007), in which causation operates

as multiple factors simultaneously. Furthermore, the tools of causal inference allow this parallel case

to be extended to more complex situations with indirect effects, rather than constraining our thinking

to simple additive terms and interactions.

Finally, proponents of the information-theoretic approach often emphasise George Box's classic point

that "all models are wrong, but some of them are useful" – they do not believe in true models

(Burnham & Anderson 2004). We agree, and argue that causal inference is not about making models

that are true, but is about making models which are useful in the sense of inferring the drivers of

patterns and predicting the effect of interventions. Causal inference provides the tools to express the

assumptions underpinning these inferences in a transparent manner, and allows us to examine how

robust our inferences are by using sensitivity analyses to explore the consequences of different causal

assumptions and latent confounding variables.

Summary

We have demonstrated that when a form of collider bias known as M-bias is present in the occupancy

process, occupancy models which are favoured by AIC and BIC produce inaccurate parameter

estimates but accurate predictions. In contrast, M-bias in the detection process does not affect the

accuracy of parameter estimates. The key conclusion supported by these results is that inference and

prediction are separate tasks which should not be conflated during model selection. The correct choice

of model selection procedure depends on the purpose for which the occupancy model will be used;

information-theoretic approaches are suitable for selecting the occupancy covariates only if the model

is to be used for predicting the site-level occupancy probability. If the goal is instead to infer the effect

of environmental covariates on occupancy, we argue the information-theoretic approach should not be

used, and advocate for an approach based on causal inference in this situation. In contrast, our results

support the selection of detection covariates using information-theoretic methods regardless of the

model's purpose, because detection probability is almost always a nuisance parameter. As the single-

season occupancy models we used are in essence a form of logistic regression, our results have wider

implications for the use of information-theoretic model selection in ecology. In particular, we argue

that our results, alongside those of others (Luque-Fernandez *et al.* 2019; McElreath 2021), emphasise

that the information-theoretic approach should not be used to compare biological hypotheses in

observational studies. Finally, we argue that causal inference and the information-theoretic approach

share similar philosophical underpinnings, and should be seen as complementary tools that

accomplish different tasks.

Acknowledgements

The authors would like to thank the members of the Conservation Ecology Group (Durham

University) for their stimulating discussions which improved the content of this manuscript. P.S.S was

supported by the Natural Environment Research Council (NERC) through the Iapetus2 Doctoral

Training Partnership.

Data Availability

Code to fully reproduce our simulations and analyses is available at:

https://zenodo.org/badge/latestdoi/462801230

Author Contributions

P.S.S and W.D. conceived the study, P.S.S. designed and implemented the simulations and analyses, and wrote the first draft of the manuscript. All authors contributed substantially to revisions of the manuscript.

References

Aho, K., Derryberry, D. & Peterson, T. (2014). Model selection for ecologists: the worldviews of AIC and BIC. *Ecology*, **95**, 631–636.

Akaike, H. (1973). *Information Theory and an Extension of the Maximum Likelihood Principle*. In: Proceeding of the Second International Symposium on Information Theory (eds. Petrov, B.N. & Kaski, F.). Akademiai Kiado, Budapest, pp. 267–281.

Akaike, H. (1998). *Prediction and Entropy*. In: Selected Papers of Hirotugu Akaike, Springer Series in Statistics (eds. Parzen, E., Tanabe, K. & Kitagawa, G.). Springer, New York, NY, pp. 387–410.

Akçakaya, H.R., Bennett, E.L., Brooks, T.M., Grace, M.K., Heath, A., Hedges, S., et al. (2018). Quantifying species recovery and conservation success to develop an IUCN Green List of Species. *Conservation Biology*, **32**, 1128–1138.

Anderson, D.R., Burnham, K.P. & Thompson, W.L. (2000). Null Hypothesis Testing: Problems, Prevalence, and an Alternative. *The Journal of Wildlife Management*, **64**, 912–923.

640 Breiner, F.T. & Bergamini, A. (2018). Improving the estimation of area of occupancy for IUCN Red

641   List assessments by using a circular buffer approach. *Biodiversity Conservation*, **27**, 2443–

642   2448.

643 Buckland, S.T., Burnham, K.P. & Augustin, N.H. (1997). Model Selection: An Integral Part of

644   Inference. *Biometrics*, **53**, 603–618.

645 Burnham, K.P. & Anderson, D.R. (2001). Kullback-Leibler information as a basis for strong inference

646   in ecological studies. *Wildlife Research*., **28**, 111.

647 Burnham, K.P. & Anderson, D.R. (2004). Multimodel Inference: Understanding AIC and BIC in

648   Model Selection. *Sociological Methods & Research*, **33**, 261–304.

649 Burnham, K.P., Anderson, D.R. & Huyvaert, K.P. (2011). AIC model selection and multimodel

650   inference in behavioral ecology: some background, observations, and comparisons.

651   *Behavioral Ecology and Sociobiology,* **l**, 65, 23–35.

652 Caughley, G. (1994). Directions in Conservation Biology. Journal of Animal Ecology, 63, 215–244.

653 Chamberlin, T.C. (1890). The Method of Multiple Working Hypotheses. *Science*, **15**, 6.

654 Chen, H.L. & Koprowski, J.L. (2015). Animal occurrence and space use change in the landscape of

655   anthropogenic noise. *Biological Conservation*, **192**, 315–322.

656 Cinelli, C., Forney, A. & Pearl, J. (2020). A Crash Course in Good and Bad Controls. *SSRN Journal*.

657 Clark, A.E. & Altwegg, R. (2019). Efficient Bayesian analysis of occupancy models with logit link

658   functions. *Ecology and Evolution*, **9**, 756–768.

659 Clarke, K.A. (2005). The Phantom Menace: Omitted Variable Bias in Econometric Research. *Conflict*

660   *Management and Peace Science*, **22**, 341–352.

661 Elliott, L.P. & Brook, B.W. (2007). Revisiting Chamberlin: Multiple Working Hypotheses for the

662   21st Century. *BioScience*, **57**, 608–614.

663 Fiske, I., & Chandler, R. (2011). unmarked: An R Package for Fitting Hierarchical Models of

664   Wildlife Occurrence and Abundance. *Journal of Statistical Software*, **43**, 1-23.

665 Forster, M.R. (2000). Key Concepts in Model Selection: Performance and Generalizability. *Journal of*

666   *Mathematical Psychology*, **44**, 205–231.

Gelman, A. & Rubin, D.B. (1995). Avoiding Model Selection in Bayesian Social Research.

Sociological Methodology, **25**, 165–173.

Gormley, A.M., Forsyth, D.M., Griffioen, P., Lindeman, M., Ramsey, D.S.L., Scroggie, M.P., et al.

(2011). Using presence-only and presence–absence data to estimate the current and potential

distributions of established invasive species. *Journal of Applied Ecology*, **48**, 25–34.

Grace, J.B. & Irvine, K.M. (2020). Scientist's guide to developing explanatory statistical models

using causal analysis principles. *Ecology*, **101**, e02962.

Greenland, S. (2003). Quantifying Biases in Causal Models: Classical Confounding vs Collider-

Stratification Bias. *Epidemiology*, **14**, 300–306.

Greenland, S. & Mansournia, M.A. (2015). Limitations of individual causal models, causal graphs,

and ignorability assumptions, as illustrated by random confounding and design unfaithfulness.

European Journal of Epidemiology, **30**, 1101–1110.

Greenland, S., Pearl, J. & Robins, J.M. (1999). Causal diagrams for epidemiologic research.

Epidemiology, **10**, 37–48.

Hossack, B.R., Lowe, W.H. & Corn, P.S. (2013). Rapid Increases and Time-Lagged Declines in

Amphibian Occupancy after Wildfire. *Conservation Biology*, **27**, 219–228.

Johnson, J.B. & Omland, K.S. (2004). Model selection in ecology and evolution. *Trends in Ecology &*

Evolution, **19**, 101–108.

Karavarsamis, N. (2015). Methods for estimating occupancy. *Bulletin of the Australian Mathematical*

Society, **92**, 518–519.

Laubach, Z.M., Murray, E.J., Hoke, K.L., Safran, R.J. & Perng, W. (2021). A biologist's guide to

model selection and causal inference. *Proceedings of the Royal Society B: Biological*

Sciences, **288**, 20202815.

Lukacs, P.M., Thompson, W.L., Kendall, W.L., Gould, W.R., Doherty, P.F., Burnham, K.P., et al.

(2007). Concerns regarding a call for pluralism of information theory and hypothesis testing:

Concerns about pluralism. *Journal of Applied Ecology*, **44**, 456–460.

Luque-Fernandez, M.A., Schomaker, M., Redondo-Sanchez, D., Jose Sanchez Perez, M., Vaidya, A.

& Schnitzer, M.E. (2019). Educational Note: Paradoxical collider effect in the analysis of

non-communicable disease epidemiological data: a reproducible illustration and web

application. *International Journal of Epidemiology*, **48**, 640–653.

MacKenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Royle, J.A. & Langtimm, C.A. (2002).

Estimating Site Occupancy Rates When Detection Probabilities Are Less Than One. *Ecology*,

83, 2248–2255.

MacKenzie, D.I., Nichols, J.D., Royle, J.A., Pollock, K.H., Bailey, L.L. & Hines, J.E. (2006).

Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species

Occurrence. 1st. edn. Elsevier.

Mazerolle, M.J. (2020). *AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c)*.

R package version 2.3-1. https://cran.r-project.org/package=AICcmodavg.

McElreath, R. (2021). *Statistical Rethinking: A Bayesian Course with Examples in R and STAN*. 2nd

edn. Chapman and Hall/CRC.

Midlane, N., O'Riain, M.J., Balme, G.A., Robinson, H.S. & Hunter, L.T.B. (2014). On tracks: A

spoor-based occupancy survey of lion Panthera leo distribution in Kafue National Park,

Zambia. *Biological Conservation*, **172**, 101–108.

Okes, N.C. & O'Riain, M.J. (2017). Otter occupancy in the Cape Peninsula: Estimating the

probability of river habitat use by Cape clawless otters, Aonyx capensis, across a gradient of

human influence. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **27**, 706–716.

Pearl, J. (1995). Causal Diagrams for Empirical Research. *Biometrika*, **82**, 669–688.

Pearl, J. (2009). Causal inference in statistics: An overview. *Statistics Surveys*, **3**, 96–146.

Pearl, J., Glymour, M. & Jewell, N.P. (2016). *Causal Inference in Statistics: A Primer*. John Wiley &

Sons, Incorporated, Chicester, United Kingdom.

R Core Team (2021). *R: A language and environment for statistical computing.* R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Richards, S.A. (2005). Testing Ecological Theory Using the Information-Theoretic Approach:

Examples and Cautionary Results. *Ecology*, **86**, 2805–2814.

Robins, J.M. & Greenland, S. (1986). The role of model selection in causal inference from

nonexperimental data. *American Journal of Epidemiology*, **123**, 392–402.

Rosenbaum, P.R. (1984). The Consquences of Adjustment for a Concomitant Variable That Has Been

Affected by the Treatment. *Journal of the Royal Statistical Society. Series A (General)*, **147**,

656–666.

Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, **6**, 461–464.

Semper-Pascual, A., Decarre, J., Baumann, M., Camino, M., Di Blanco, Y., Gómez-Valencia, B., et

al. (2020). Using occupancy models to assess the direct and indirect impacts of agricultural

expansion on species' populations. *Biodiversity Conservation*, **29**, 3669–3688.

Shmueli, G. (2010). To Explain or to Predict? *Statistical Science*, **25**, 298-310.

Stone, M. (1977). An Asymptotic Equivalence of Choice of Model by Cross-Validation and Akaike's

Criterion. *Journal of the Royal Statistical Society. Series B (Methodological)*, **39**, 44–47.

Zimbres, B., Peres, C.A., Penido, G. & Machado, R.B. (2018). Thresholds of riparian forest use by

terrestrial mammals in a fragmented Amazonian deforestation frontier. *Biodiversity*

Conservation, **27**, 2815–2836.

736