Individually unique, fixed stripe configurations of *Octopus chierchiae* allow for

photoidentification in long-term studies

Benjamin Liu[†,1], *Leo Song[†,1], Saumitra Kelkar[†], Anna Ramji[†], Roy Caldwell[†]

[†]Department of Integrative Biology, University of California Berkeley, Berkeley, California,

United States of America

* Corresponding author:

Email: leosong@berkeley.edu (LS)

[1] These authors contributed equally to this work.

Abstract

The Lesser Pacific Striped Octopus, *Octopus chierchiae*, is a small iteroparous octopus

known to inhabit intertidal regions of the Pacific coast of Central America. Many details about

its life history and ecology remain unknown. For apparently rare and delicate animals such as *O.*

chierchiae, non-extractive sampling methods are necessary to study individuals and populations

over time. After photographically documenting the physical development of 25 octopuses from

hatching, we have concluded that *O. chierchiae* has individually unique stripe configurations that

remain constant throughout their post-hatchling lifetimes. Furthermore, using photographs taken

of animals in captivity on different dates over many months, we show that untrained volunteers

can accurately identify whether or not a pair of images depicts the same individual octopus.

These results demonstrate that laboratory-reared individuals could be identified via photographs

taken at different points in their lifetimes, which suggests wild individuals can also be

recognized and observed for longitudinal field studies. In addition, our results imply potential for

photoidentification and community science to be used as non-extractive, non-intrusive sampling

methods for future studies of wild *O. chierchiae*.

Introduction

Octopuses' remarkable camouflage ability, complex behavior, and intelligence have

attracted vast scientific interest (Di Cosmo et al., 2021) and pop-culture intrigue (Hackett and

Harrington, 2018). However, their soft body forms, elusive behavior, and ever-threatened

habitats make it difficult to track individuals in the wild over time (Huffard et al., 2008). Octopus

body patterns are produced by a combination of pigment-filled chromatophores, color-reflective

iridophores, and passively reflective leucophores, as well as muscular and hydrostatic forces that

produce textural details (Mäthger et al., 2009; How et al., 2017). While these characteristics

allow for rapid, neurally controlled color change, they operate within the fixed anatomical

architecture of an octopus's skin (Packard and Hochlberg, 1977; Messenger, 2001). In some

species, the consistency of body color patterns may allow humans to identify individuals from

photographs, thereby aiding longitudinal studies (Huffard et al. 2008).

Octopus chierchiae (Jatta, 1889) exhibits qualities that may permit, and benefit greatly

from, photoidentification. Also known as the Lesser Pacific Striped Octopus or the Pygmy Zebra

Octopus, this species of dwarf octopus is found in shallow waters on the Pacific coast of the

Americas, from Colombia to Baja California (Kerstitch, 1989). *Octopus chierchiae* –and the yet-

undescribed Larger Pacific Striped Octopus and *Octopus zonatus* (Voss, 1968) – are commonly

referred to as "harlequin" octopuses in reference to the black and white stripes and spots that

cover their entire body [Fig 1A]. Although the animals may take on all pale or all dark

morphologies, *O. chierchiae* most often display the high-contrast stripe-bar-spot morphology

(defined in reference to the Larger Pacific Striped Octopus by Caldwell et al., 2015)[Fig 2]

where the stripe patterns are most clearly visible while the animal is at rest. These stripes are not

initially apparent in hatchlings [Fig 1B], but may be observed with magnification as early as the

fifth day post-hatching [Fig 1C].

Fig 1. Photographs depicting *Octopus chierchiae* of different ages.

(A) One-year-old *O. chierchiae* adult. (B) One-day-old *O. chierchiae* hatchling. (C) Five-day-old

O. chierchiae hatchling already displaying a developing stripe configuration.

Fig 2. Visibility of *O. chierchiae* stripe configuration during different body color morphs.

(A) *O. chierchiae* displaying dark morph (left) and stripe-bar-spot morph (right) illustrating how

chromatophores activate directly over the leucophores to mask their whiteness. (B) A different

O. chierchiae displaying stripe-bar-spot morph (left) and pale morph (right) illustrating how

chromatophores expand adjacent to leucophore white markings to produce high-contrast

markings.

While most octopus species exhibit semelparity and only reproduce near the end of their

lifespan (Rocha et al., 2001), *O. chierchiae* is iteroparous (Rodaniche, 1984) and have been

observed to lay up to eight clutches of eggs over the course of their lifespan in our laboratory.

Overall, *O. chierchiae's* iteroparous reproductive strategy (Rodaniche, 1984), observed lifespan

of over two years in our lab, and lack of a planktonic developmental stage (Sweeney et al., 1992)

make breeding and maintaining the species in captivity a practical and viable pursuit,

contributing to its growing potential as a model organism (Grearson et al., 2021). As its

candidacy and status as a marine model organism develop, the need to locate, monitor, and study

wild individuals of this species grows as well. Specifically, the ability to identify and observe the

octopuses on an individual basis would be crucial for both field studies (Arechavala-Lopez et al.,

2019) as well as controlled laboratory research involving multiple individual animals (Morse et

al., 2017).

Current studies on wild cephalopods most commonly involve tag-recapture studies

(Semmens et al., 2007), as exemplified by efforts to investigate the growth and movement of

Octopus vulgaris in Central Western Sardinia (Mereu et al., 2010) and *Enteroctopus dofleini* in

the Bering Sea (Brewer et al., 2017). Visible Implant Elastomer tags have demonstrated viable

usage in long term studies of larger cephalopods such as *E. dofleini* (Brewer and Norcross,

2012). However, as Semmens et al. (2007) discuss, external tags can harm octopuses' delicate

tissues, cannot be applied to animals that are too small, and can fall off or be deliberately

removed by the tagged animal. Tattooing and branding techniques have had some success in

cephalopod tracking studies as they eliminate the issue of tag removal and may be applied to

small animals, but they are often practically difficult to execute and can cause the animal severe

distress and damage (Semmens et al. 2007). Other rising technological tracking methods for

cephalopods include chemical tagging, chemical analyses of cephalopod hard parts, geographic

differences in parasite fauna, molecular genetics, and satellite data of detected signals (Semmens

et al. 2007). However, *O. chierchiae*'s delicate and diminutive bodies (mantle length up to 4 cm)

make longitudinal studies of wild individuals through the previously mentioned techniques

difficult to complete or potentially dangerous to the animals.

Contrary to the aforementioned methods, photography serves as a largely inexpensive,

non-invasive, non-extractive, and widely accessible technique to produce high-quality data. The

low-impact nature of this approach as well as the virtual ubiquitousness of high-quality cameras

that can record photographs, videos, geospatial data, and temporal data, has enabled community

science efforts to make increasingly significant contributions to scientific research, as showcased

by the US National Parks Service's What's Invasive! project and NSF NEON's Project BudBurst

(Graham et al., 2011; Dickinson et al 2012; Chandler et. al., 2017). *In situ* research in particular

has benefited tremendously from the growing availability of photodata, as many inferences on

the abundance, distribution, phenology, and ecology of wild animal populations can be drawn

from both the content and the context of these photographic images (Graham et al., 2011;

Ballance, 2018), especially if members of the species of interest are individually and uniquely

identifiable. The employment of such photography-analysis techniques has allowed researchers

to estimate population dynamics (Karanth and Nichols, 1998), determine individual movement

and home ranges (Oliveira-Santos et al., 2010) and quantify disease dynamics within populations

(Geers et al., 1997). Besides its exceptional accessibility, the photography-based approach offers

yet another distinctive advantage: it may be applied retrospectively to existing photobanks

(Johnston et al., 2017). All the aforementioned advantages along with the continual development

of new computer-aided photoidentification methods (Kelly, 2001; Schneider, 2019; Tan et al.,

2021; Vidal, 2021) and analytical techniques make photography an ideal approach for

monitoring and studying soft-bodied cephalopods (Roper and Hochberg 1988; Vecchione and

Roper, 1991; Drerup et al., 2021) like *O. chierchiae*.

Longitudinal field studies conducted on the Caribbean reef squid, *Sepiotheuthis*

sepioidea, have demonstrated that individual identification in the wild via body patterning and

markings from injury can be viable in cephalopods (Byrne et al., 2010). However, as far as we

know, *Wunderpus photogenicus* is the only octopus species in which individual cephalopods are

shown to be identifiable by photographs of their individually unique stripe configurations

(Huffard et al., 2008). Here, we test whether *O. chierchiae* also have identifiable, individually

unique stripe configurations, and whether these remain constant throughout their lifetimes,

allowing for photoidentification over long time spans by untrained volunteers.

Materials and Methods

Ethics Statement

This study was performed at the University of California, Berkeley in an AAALAC

accredited facility, overseen by the Office of Laboratory Animal Care (OLAC), and thus follows

the guidelines set forth by the Guide for the Care and Use of Laboratory Animals. Because the

study species is an invertebrate, at the time of the study, neither USDA nor PHS policy covered

this study and an IACUC approved animal use protocol was not needed. An official statement

from the UC Berkeley OLAC is provided in the Supplementary Materials [S1 File]. Human

participants were recruited via social media by reaching out in Slack channels and Facebook

groups. These social media channels ranged from our workplace channels to hobbyist groups that

are not biology or science oriented in order to minimize bias from participant backgrounds. All

data were analyzed anonymously.

Collection and Rearing

Two male and two female adult *O. chierchiae* were obtained from a fully-licensed

commercial collector in Nicaragua during 2017 and 2018. We housed them individually in one

gallon (3.79 liter) glass aquaria lined with gravel, light sand, and bivalve shell fragments [Fig

3A] and provided PVC tubes, glass vials, or hollow rocks as shelter. Hatchlings were housed

individually in glass jars with screen lids (9 cm tall, 8 cm diameter) containing a variety of

substrates including dark sand, light sand, bivalve shell fragments, snail shells, and gravel, all

within the same 40 gallon tank (151.42 liter) [Fig 3B]. Mature animals were housed individually

in larger perforated plastic jars containing a similar mix of substrates, within a separate 20 gallon

(75.71 liter) tank [Fig 3C]. As gonad maturity is difficult to assess in live cephalopods, we

approximated the age indicative of sexual maturity to be 16 weeks old, based on behavioral

observations, i.e. engaging in copulation. All aquaria in our laboratory were filled and

maintained with artificial seawater (32-36 ppt) at 22-24 □. Detailed breeding and rearing

methods can be found in the Supplementary Materials [S2 Supporting Information].

Fig 3. Photographs depicting lab aquaria.

(A) One gallon (3.79 liter) individual mating tank where matings occurred and eggs were laid.

(B) 40 gallon nursery tank where animals were reared in individual glass jars with screen lids

from hatching until sexual maturity. (C) 20 gallon (75.71 liter) general tank where mature

animals were housed in individual perforated plastic jars.

A total of 156 individuals from 10 clutches hatched over the course of this study (3 to 32

hatchlings per clutch) with an average body weight of 17.4 mg (min = 9.5 mg, max = 25.6 mg)

[S3 Dataset]. Comparisons of individuals from different clutches, hatching times, and parents

were not tested in this study. Captive-bred *O. chierchiae* that survived the first few weeks of life

typically lived for at least a year, with our longest-lived captive-bred octopus being a male

hatched in July 2018 who lived for 22 months and fathered at least 4 different clutches of

offspring before his death in June 2020. While the exact age and influence of life experiences on

the development of wild caught individuals cannot be determined, having study animals bred and

raised entirely in captivity allowed us the unique opportunity to follow the species' development

throughout its entire lifecycle, from hatching until death.

<u>Photography</u>

We photographed individual *O. chierchiae* from above water, using a handheld Apple

iPhone 6s smartphone and a handheld 4K Sony camcorder (model HDR-XR260V) from June

2018 to July 2020. Due to high hatchling and juvenile mortality rate (Grearson et al., 2021) and

some individuals' aversion to open environments, only a subset [n = 25] of our lab reared

animals were involved in this study. We only included individuals whose lifespan and activity

patterns allowed stripe configurations to be clearly photographed on multiple occasions. In order

to obtain a photographic record of each individual's development over the course of their

lifetime, we photographed lab-reared *O. chierchiae* weekly, without removing them from their

containers.

Terms used to describe *O. chierchiae* body patterns follow those defined and illustrated

previously (Packard and Sanders, 1969; Packard and Sanders, 1971; Packard and Hochberg,

1977, Caldwell et al., 2015). Morphological appearance of each animal may vary greatly

depending on body position or chromatophore activation. Consequently, this investigation

focused on the stripe configurations of the chronic patterns located on the animals' dorsal mantle

surface [Fig 4A], because these were the most consistently observable identifying characteristics.

Chronic patterns are those that persist for hours or days, or those that the animal returns to after

acute variations that last seconds or minutes (Packard and Sanders, 1969). The stripe-bar-spot

pattern is generally composed of brown and white head bars, eye bars, extended eye bars, and

transverse (chevron) stripes on the head and mantle, and longitudinal stripes on the arm crown.

Occasionally, small brown mantle dark spots can be found within a white mantle stripe, and vice

versa. The precise shape of each of these components was variable between individuals. We

chose to specifically examine [S7 Dataset] the dark transverse mantle stripes and dark mantle

spots in the chronic patterns of *O. chierchiae*. We defined dark transverse mantle stripes as any

dark pattern component that traverses the width of the individual's mantle. We further

categorized these into dark continuous stripes [Fig 4B] and dark discontinuous stripes [Fig 4C],

those that contained no visible breakage and those that contained visible breakage, respectively.

We also defined dark mantle spots [Fig 4D] as any dark pattern component completely encircled

by white pattern components on an individual's mantle. Although we provide above a list of

body pattern component definitions, this study focused on the animal's body patterning as a

whole. These components may come together in many combinations, including overlapping one

another, to produce a wide variety of patterns unique to each individual animal [Fig 4A]. Each

observer may focus on a different identifying characteristic (such as focusing on the pale

components rather than the dark components) within the stripe configuration and yet still come

to the same conclusion.

Figure 4. Specific and polymorphic dorsal patterns on mantle of different individuals of *O.

chierchiae.**

(A) Side-by-side display of dorsal mantle surface of all *O. chierchiae* individuals involved in the

study. (B) Example of a continuous stripe [red]. (C) Example of a discontinuous stripe [blue].

(D) Example of a spot [green].

Per our observations, disturbances could stimulate a change in the animal's color morph

[Fig 2] or flashing between morphs. Over the course of our study, we attempted to avoid

disturbing and distressing the animals during husbandry and photography sessions. If an

individual was hiding or clinging to the jar lid, they were not photographed that session. One

photograph per individual per session was taken from angles relative to the animal's body

position at the time of viewing in order to capture the octopus' dorsal mantle surface as clearly as

possible. For the four individuals with the most extensive photobanks, we arranged a series of

photographs chronologically, one per month if available, and traced their stripe configurations

using Adobe Illustrator. Each isolated pattern component was color coded [Fig 5] to visualize

and establish the consistency of the stripe configurations. Body patterns of *O. chierchiae* became

visible to the naked eye when individuals were around two weeks old and were consistently fully

visible by four weeks, but can be observed with magnification as early as the fifth day after they

hatch [Fig 1C].

Fig 5. Timeline of *O. chierchiae* individuals showing consistent body patternings.

The bottom row of each panel duplicates these photographs, with analogous pattern components

color coded. (A) Timeline of *O. chierchiae* individual (hatched Nov 12, 2018) over a 5 month

time span. (B) Timeline of *O. chierchiae* individual (hatched May 6, 2019) over a 6 month time

span. (C) Timeline of *O. chierchiae* individual (hatched Nov 26, 2018) over a 7 month time span.

(D) Timeline of Octopus individual (hatched May 6, 2019) over a 4 month time span.

Survey Design and Evaluation

We created a survey following the methods of Huffard et. al. (2008) using the photobank

of images taken over the course of our study. The survey [S4 File] consisted of a series of 20

slides, arranged by a random number generator, each depicting two photographs of *O. chierchiae*

individuals side by side, with no repeated measurements. Photograph pairs were rotated and

cropped to display the animals in similar orientations and sizes. Only photographs of animals

over four weeks old were used in the survey as that was the age when all the individuals' stripe

configurations became clearly and consistently discernable by our photography equipment. We

solicited survey participants via social media for two weeks, and collected responses via Google

Forms for a total of four weeks. For each slide, participants were asked to respond whether the

slide appeared as a "match" (two images of the same individual taken at different times) or "no

match" (two images of different individuals). No time limit or built-in zoom function was

implemented and participants were free to go back and edit responses prior to submission. The

survey contained nine slides depicting matching photographs, taken up to 25 weeks apart, and 11

slides depicting non-matching photographs. Each set of survey responses was given a score

$[\frac{correct\ answers}{total\ questions} \times 100]$. Incorrect answers were categorized as false positive (meaning an

erroneous match) and false negative (meaning a missed match). Unanswered questions were

scored as incorrect, but omitted in our false positive-false negative analysis. In our social media

solicitation, we reminded participants to complete the survey on a computer screen, as designed.

However, because our data collection methods were not face-to-face and were analyzed

anonymously, we cannot confirm the device each survey was completed on.

These survey designs were refined based on a preliminary survey detailed in

Supplementary Materials [S5 File; S6 Dataset].

Results

Over the course of the study, we observed that *O. chierchiae* individuals' [n = 25] stripe

configurations were consistent throughout their lifetimes, evidenced by photographic records

[Fig 5]. The *O. chierchiae* bred and raised in our lab displayed [Fig 4] an average of 4.8 (±0.49

S.D.) dark transverse mantle stripes in their chronic patterns. In 20% (n =5) of the animals, this

included one discontinuous stripe [Fig 4C]. Additionally, 24% (n =6) of the animals also

displayed a dark mantle spot [Fig 4D] [S7 Dataset]. Examples of how pattern component counts

were obtained may be found in Supplementary Materials [S8 Figure]. All members of our

research group (the authors of this paper) were able to recognize individual animals based on the

uniqueness [Fig 4] and consistency [Fig 5] of the octopuses' stripe configurations. Next, we

investigated if untrained observers would be able to do the same. Our study focused on untrained

volunteers specifically to control for potential overestimation from trained experts with daily

exposure to the animals.

Survey respondents (n = 38 responses) had a mean score of 84.2% ± 15.4% standard

deviation, with a standard error of the mean of 2.5%. Survey respondents had a median score of

90%, with 44.7% of respondents (n = 17) scoring 95% or higher [Fig 6]. On average,

respondents missed 13.7% of total possible matches (1.9 false negative responses out of 20

questions) and erroneously matched 17.2% of non-matching individuals (1.2 false positive

responses out of 20 questions). Survey data can be found in the Supplementary Materials [S9

Dataset].

Fig 6. Distribution of survey response scores (n = 38 survey responses).

Survey scores in percent are calculated by $[\frac{correct\ answers}{total\ questions} \times 100]$.

To investigate if there was a significant difference between false positive and false

negative rates in the survey response, we performed the Wilcoxon Signed Rank Test with our

null hypothesis being that false positive rates and false negative rates were the same, while our

alternative hypothesis being that false positive rates and false negative rates were different. The

test result revealed that there was not a significant difference between false positive and false

negative rates (Wilcoxon Signed Rank Test: test statistic = 150, α = 0.05, n = 28, critical value =

116), suggesting a potential future study of wild populations would not be biased towards one

error type or another.

Discussion

We hypothesized that untrained survey participants would be able to identify *O.*

chierchiae individuals based on their stripe configurations. This hypothesis was supported. The

full group of participants had an average score of nearly 84.2%, while 52.6% of all participants

scored at least 90% on the survey [Fig 7], showing that *O. chierchiae* individuals were

distinguishable to a majority of untrained observers. The only participant to score 50% (expected

value of random choice) or below responded to every question the same. Due to the anonymous

nature of our methods, we are unable to confirm if this participant truly could not distinguish

between the individuals in each photograph, or if they did not put forth genuine effort in

responding to the survey questions. No individual question was answered incorrectly by a

majority of participants. Based on these results, we conclude that *O. chierchiae* can be

consistently identified throughout their lives via their individually unique stripe configurations

by both trained and untrained observers, when an unobstructed, roughly in-focus view of the

stripe configuration on the mantle's dorsal surface is provided.

Fig 7. Individual survey response scores (n = 38)

Bar chart illustrating total score (Blue), false negative (Red), and false positive (Orange) when

assessing images.

Having hatched, reared and documented the individual *O. chierchiae* used in the surveys

in captivity over extensive time spans – often over the course of their entire life history – we

observed no evidence that stripe configurations changed over the lifetimes of individuals [Fig 5].

Our survey suggests that these permanent stripe configurations are a viable method of identifying

the same individual at different points in time, even by volunteers that may have no experience

with octopuses. To our knowledge, this is the first evidence that individually unique stripe

configurations can be used by volunteers to consistently identify the same individual octopus

throughout its development, from juvenile to adult.

This information could be used to design non-invasive studies monitoring populations of

O. chierchiae over time *in situ*. It is conceivable that environmental disturbances and injuries

may affect the stripe configuration of *O. chierchiae* in the wild. However, because so little is

known about the species in its natural habitat and we worked to avoid injuring animals over the

course of this study, we do not have conclusive evidence whether or not environmental

conditions and injuries would affect an animal's stripe configuration.

Based on our results that participants could consistently identify whether two

photographs depicted the same octopus, we recommend the survey design methods outlined

above be employed in future community science efforts to study *O. chierchiae* via

photoidentification. Furthermore, based on the high median scores recorded, it seems plausible

that researchers may use this survey as a screener to identify a pool of volunteers capable of

achieving highly accurate results. Smartphone apps or websites can be utilized to gather

responses *en masse* and the process may even be automated via computer-assisted

photoidentification software with enough responses and high quality image data. We envision

untrained volunteers engaging in community science efforts by contributing to the establishment

of a photodata repository. This could be done by sending a photograph in which the stripe

configuration on the animal's dorsal mantle surface is visible along with the date, time, and

geographic coordinates saved from a smartphone. Future research teams may make use of such a

database and draw inferences about the species' natural history, including home range, behavior,

population size, population density, and wild lifespan.

While it is known that there can be variation in the founder chromatophores of individual

octopus hatchlings of the same species (Ylitalo et al., 2014) and that *O. chierchiae* can begin to

display the characteristic stripe pattern as soon as five days after hatching [Fig 1C], we did not

begin photographing animals for this survey study until they were at least four weeks old. Thus,

more work is needed to determine whether the individually unique stripe configurations of *O.*

chierchiae are fixed from hatching, and if not, at what point in early development the stripe

configurations become permanent. Identifying the specific anatomical composition of *O.*

chierchiae's stripe configuration was also beyond the scope of our study, and would have

required sacrificing our study subjects. We encourage future research efforts to be directed into

examining the skin histology and ultrastructure of *O. chierchiae*, when deceased individuals are

available, in order to gain further insight into the physiological basis of their stripe configuration

development.

Does *O. chierchiae*'s individually unique, lifelong body patterning play a role in

intraspecific behavior in the wild? Octopus species are known to have the ability to recognize

conspecifics (Walderon et al., 2011). Octopuses generally have high visual acuity, and the ability

to remember (Hochner et al., 2006). Individual recognition has been explored in several other

cephalopods (Shashar 2004; Boal, 2006), and at least one species, *Octopus vulgaris*, is thought to

have the ability of individual conspecific recognition (Tricario et al., 2011). The harlequin

octopuses, with their bold semi-permanent body patterns, also "rely less upon changes in

chromatophores than upon perceptions of postures and positions," (Moynihan, 1985). In our

view, these factors make it worthwhile to explore whether individual recognition based on visual

cues plays a role in the lives of wild *O. chierchiae* and other octopus species.

Acknowledgements

This study would not have been possible without the hard work and support of the Caldwell

Lab's *O. chierchiae* husbandry team over the past few years (2016-2020). Dr. Christine Huffard

imparted much insight and expertise throughout this investigation and critiqued earlier drafts of

the manuscript. Dr. Huffard's 2008 manuscript on *W. photogenicus* inspired this paper. Dr.

Caroline Williams from UC Berkeley and Bret Grasse from the Woods Hole Marine Biological

Laboratory provided correspondence and guidance. Richard Ross from the California Academy

of Sciences contributed to husbandry methods.

References

Arechavala-Lopez P, Minguito-Frutos M, Follana-Berná G, Palmer M. Common octopus settled

in human-altered Mediterranean coastal waters: from individual home range to population

dynamics. ICES J Mar Sci. 2019;76(2):585–97.

Ballance LT. Contributions of photographs to cetacean science. Aquat Mamm. 2018;44(6):668–

82.

Boal J. Social recognition: a top down view of cephalopod behaviour. Vie et Milieu - Life &

Environment. 2006;56(2):69-79.

Brewer RS, Norcross BL. Long-term retention of internal elastomer tags in a wild population of

North Pacific giant octopus (*Enteroctopus dofleini*). Fish Res. 2012;134–136:17–20. Available

from: http://dx.doi.org/10.1016/j.fishres.2012.07.020

Brewer RS, Norcross BL, Chenoweth E. Temperature- and size-dependent growth and

movement of the North Pacific giant octopus (*Enteroctopus dofleini*) in the Bering Sea. Mar Biol

Res. 2017;13(8):909–18.

Byrne RA, Wood JB, Anderson RC, Griebel U, Mather JA. Non-invasive methods of identifying

and tracking wild squid. Ferrantia. 2010;59:22–31.

Caldwell RL, Ross R, Rodaniche A, Huffard CL. Behavior and body patterns of the Larger

Pacific Striped Octopus. PLoS One. 2015;10(8):e0134152.

Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, et al. Contribution of citizen

science towards international biodiversity monitoring. Biol Conserv. 2017;213:280–94.

Dickinson JL, Shirk J, Bonter D, Bonney R, Crain RL, Martin J, et al. The current state of citizen

science as a tool for ecological research and public engagement. Front Ecol Environ.

2012;10(6):291–7.

Di Cosmo A, Pinelli C, Scandurra A, Aria M, D'Aniello B. Research trends in octopus biological

studies. Animals (Basel). 2021;11(6):1808. Available from: https://www.mdpi.com/2076-

2615/11/6/1808

Drerup C, Jackson A, Rickard C, Skea M, Cooke GM. Field observations on the behavioural

ecology of the stout bobtail squid *Rossia macrosoma* (Cephalopoda: Sepiolidae) from Scottish

waters. Mar Biodivers. 2021;51(4). Available from: http://dx.doi.org/10.1007/s12526-021-

01202-y

Geers P, Puers B, Goedseels V, Wouters P. Electronic identification, monitoring and tracking of

animals. Wallingford, England: CABI Publishing; 1997.

Graham EA, Henderson S, Schloss A. Using mobile phones to engage citizen scientists in

research. Eos. 2011;92(38):313-315.

Grearson AG, Dugan A, Sakmar T, Dölen G, Gire DH, Sivitilli DM, et al. The lesser Pacific

striped octopus, *Octopus chierchiae*: an emerging laboratory model for the study of octopuses.

Front Mar Sci. 2021;8:753483. doi: 10.3389/fmars.2021.753483

Hackett J, Harrington S. Beasts of the Deep: Sea Creatures and Popular Culture. Bloomington:

Indiana University Press; 2018.

Hochner B, Shomrat T, Fiorito G. The octopus: a model for a comparative analysis of the

evolution of learning and memory mechanisms. Biol Bull. 2006;210(3):308–17.

How MJ, Norman MD, Finn J, Chung W-S, Marshall NJ. Dynamic skin patterns in cephalopods.

Front Physiol. 2017;8:393. Available from: http://dx.doi.org/10.3389/fphys.2017.00393

Huffard CL, Caldwell RL, DeLoach N, Gentry DW, Humann P, MacDonald B, et al.

Individually unique body color patterns in octopus (*Wunderpus photogenicus*) allow for

photoidentification. PLoS One. 2008;3(11):e3732.

Jatta G, Elenco dei Cefalopodi delle "Vittor Pisani". Bollettino della Societa di Naturalisti in

Napoli. (series 1) 1889;3(1): 63-67.

Johnston DR, Rayment W, Slooten E, Dawson SM. A time-based method for defining

associations using photo-identification. Behaviour. 2017;154(9–10):1029–50.

Karanth KU, Nichols JD. Estimation of tiger densities in India using photographic captures and

recaptures. Ecology. 1998;79(8):2852–62.

Kelly MJ. Computer-aided photograph matching in studies using individual identification: An

example from Serengeti cheetahs. J Mammal. 2001;82(2):440–9.

Kerstitch AN. Sea of Cortez Marine invertebrates: A guide for the Pacific Coast, Mexico to

Ecuador. 1st ed. Monterey, CA: Sea Challengers; 1989.

Mäthger LM, Denton EJ, Marshall NJ, Hanlon RT. Mechanisms and behavioural functions of

structural coloration in cephalopods. J R Soc Interface. 2009;6 Suppl 2(suppl_2):S149-63.

Available from: http://dx.doi.org/10.1098/rsif.2008.0366.focus

Mereu M, Masala P, Maccioni A, Stacca D, Cau A, Cuccu D. Tagging *Octopus vulgaris*

(Octopoda: Octopodidae) in an area of central western Sardinian waters. Biol Mar Mediterr.

2010;17(1):306-307.

Messenger JB. Cephalopod chromatophores: neurobiology and natural history. Biol Rev Camb

Philos Soc. 2001;76(4):473–528.

Morse P, Zenger KR, McCormick MI, Meekan MG, Huffard CL. Chemical cues correlate with

agonistic behaviour and female mate choice in the southern blue-ringed octopus, *Hapalochlaena*

maculosa (Hoyle, 1883) (Cephalopoda: Octopodidae). J Molluscan Stud. 2017;83(1):79–87.

Moynihan M. Communication and Noncommunication by Cephalopods. Bloomington, MN:

Indiana University Press; 1985.

Oliveira-Santos LGR, Machado-Filho LCP, Tortato MA, Brusius L. Influence of extrinsic

variables on activity and habitat selection of lowland tapirs (*Tapirus terrestris*) in the coastal

sand plain shrub, southern Brazil. Mamm Biol. 2010;75(3):219–26.

Packard A, Sanders GD. What the octopus shows to the world. Endeavour. 1969;28(104):92-98.

Packard A, Sanders GD. Body patterns of *Octopus vulgaris* and maturation of the response to

disturbance. Anim. Behav. 1971;19(4):780-790.

Packard A, Hochberg FG. Skin patterning in Octopus and other genera. Symp. Zool. Soc. Lond.

1977;38:191-231.

Rodaniche AF. Iteroparity in the lesser Pacific striped octopus *Octopus chierchiae* (Jatta, 1889).

Bulletin of Marine Science. 1984 Jul 1;35(1):99-104.

Rocha F, Guerra A, González AF. A review of reproductive strategies in cephalopods. Biol. Rev.

Camb. Philos. Soc. 2001;76(3):291–304. Available from:

http://dx.doi.org/10.1017/s1464793101005681

Roper CFE, Hochberg FG. Behaviors and systematics of cephalopods from Lizard Island,

Australia, based on color and body patterns. Malacologia. 1988;29(1):153-93.

Schneider S, Taylor GW, Linquist S, Kremer SC. Past, present and future approaches using

computer vision for animal re□identification from camera trap data. Methods Ecol Evol.

2019;10(4):461–70. Available from: http://dx.doi.org/10.1111/2041-210x.13133

Semmens JM, Pecl GT, Gillanders BM, Waluda CM, Shea EK, Jouffre D, et al. Approaches to

resolving cephalopod movement and migration patterns. Rev Fish Biol Fish. 2007;17(2–3):401–

23.

Shashar N, Vaughan K, Loew E, Boal J, Hanlon R, Grable M. Behavioral evidence for

intraspecific signaling with achromatic and polarized light by cuttlefish (Mollusca:

Cephalopoda). Behaviour. 2004;141(7):837–61. Available from:

http://dx.doi.org/10.1163/1568539042265662

Sweeney MJ, Roper CFE, Mangold KM, Clark MR, Boletzky SV. "Larval" and juvenile

cephalopods: a manual for their identification. Smithson Contrib Zool. 1992;(513):1–182.

Available from: http://dx.doi.org/10.5479/si.00810282.513

Tan HY, Goh ZY, Loh K-H, Then AY-H, Omar H, Chang S-W. Cephalopod species

identification using integrated analysis of machine learning and deep learning approaches. PeerJ.

2021;9(e11825):e11825. Available from: http://dx.doi.org/10.7717/peerj.11825

Tricarico E, Borrelli L, Gherardi F, Fiorito G. I know my neighbour: individual recognition in

Octopus vulgaris. PLoS One. 2011;6(4):e18710.

Vecchione M, Roper CFE. Cephalopods observed from submersibles in the western North

Atlantic. Bull Mar Sci. 1991;49(13):433-45.

Vidal M, Wolf N, Rosenberg B, Harris BP, Mathis A. Perspectives on individual animal

identification from biology and computer vision. Integr Comp Biol. 2021;61(3):900–16.

Available from: http://dx.doi.org/10.1093/icb/icab107

Voss GL. Biological Investigations of the Deep Sea. 39. Octopods from the R/V Pillsbury

Southwestern Caribbean Cruise, 1966, with a description of a new species, *Octopus zonatus*.

Bulletin of Marine Science. 1968 Jul 2;18(3):645-59.

Walderon MD, Nolt KJ, Haas RE, Prosser KN, Holm JB, Nagle GT, et al. Distance

chemoreception and the detection of conspecifics in *Octopus bimaculoides*. J Molluscan Stud.

2011;77(3):309–11.

Ylitalo HA, Watling L, Toonen RJ. First description of hatchlings and eggs of *Octopus oliveri*

(Berry, 1914) (Cephalopoda: Octopodidae). Molluscan Res. 2014;34(2):79–83.

Supporting Information Captions

S1 File. OLAC Ethics Statement

S2 Supporting Information. *O. Chierchiae.* Breeding and Rearing Methods

S3 Dataset. *O. chierchiae* Hatchling Weights

S4 File. Survey

S5 File. Preliminary Survey and Discussion

S6 Dataset. Preliminary Survey Responses

S7 Dataset. Dark Mantle Pattern Component Counts

S8 Figure. Dark Mantle Pattern Component Counts Example

S9 Dataset. Survey Responses

Keep Closed!

Octopus chierchiae

| May 17, 2019 | Jun 8, 2019 | Aug 2, 2019 | Oct 1, 2019 |

| May 17, 2019 | Jun 8, 2019 | Aug 2, 2019 | Oct 1, 2019 |

May 3, 2019   Jun 8, 2019   Jul 1, 2019   Aug 1, 2019   Sep 17, 2019   Nov 17, 2019

May 3, 2019   Jun 8, 2019   Jul 1, 2019   Aug 1, 2019   Sep 17, 2019   Nov 17, 2019

April 8   June 13   July 1   October 1   November 10

April 8   June 13   July 1   October 1   November 10

Jul 17, 2019   Aug 1, 2019   Sep 17, 2019   Oct 1, 2019   Nov 10, 2019

Jul 17, 2019   Aug 1, 2019   Sep 17, 2019   Oct 1, 2019   Nov 10, 2019

Survey Results of Individual Participants

Percent of Total Score

Participant ID