Individually unique, fixed body patterns of *Octopus chierchiae* allow for

photoidentification in long-term studies

Benjamin Liu[1], *Leo Song[1], Saumitra Kelkar[1], Anna Ramji[1], Roy Caldwell[1]

[1]Department of Integrative Biology, University of California Berkeley, Berkeley, California,

United States of America

* Corresponding author:

Email: leosong@berkeley.edu (LS)

¶These authors contributed equally to this work.

Abstract

The Lesser Pacific Striped Octopus, *Octopus chierchiae*, is a small iteroparous octopus

known to inhabit intertidal regions of the Pacific coast of Central America. Many details about

the species' life history in the wild remain unknown. For apparently rare and delicate animals

such as *O. chierchiae*, non-extractive sampling methods are necessary to study individuals and

populations over time. After photographically documenting the physical development of 25

octopuses from hatching, we have concluded that individual *O. chierchiae* have individually

unique body patterns that remain constant throughout their lifetimes. Furthermore, using

photographs taken of animals in captivity on different dates over many months, we show that

untrained volunteers can accurately identify whether or not a pair of images depicts the same

individual octopus. These results demonstrate that wild individuals could be recognized and

observed for in situ longitudinal studies. In addition, our results imply potential for

photoidentification and community science to be used as non-extractive, non-intrusive sampling

methods for future studies of wild *O. chierchiae* and other octopuses.

Introduction

Constituting an estimated 99% of all marine species and spanning over 30 phyla, marine

invertebrates are amongst the most diverse, ecologically-important, and understudied groups of

organisms on the planet (Mather 2013). Cephalopods such as octopuses, squids, and cuttlefishes

represent a peculiar class of marine invertebrates characterized by a bilateral body plan, a

prominent head, and a set of muscular appendages surrounding a beaked mouth (Vendetti 2006);

(Kaestner 1967). Their remarkable camouflage ability, complex behavior, and intelligence have

attracted vast scientific interest and pop-culture intrigue (Hackett and Harrington 2018).

However, their soft body forms, elusive behavior, and ever-threatened habitats make it difficult

to track individual cephalopods in the wild over time (Huffard et al, 2008).

Octopus chierchiae (Jatta, 1889) is no exception. Also known as the Lesser Pacific

Striped Octopus or the pygmy zebra octopus, this species of dwarf octopus is found in shallow

waters on the Pacific coast of the Americas, from Colombia to Baja California (Kerstitch, 1989).

Octopus chierchiae and its relatives - the yet-undescribed Larger Pacific Striped Octopus and

Octopus zonatus (Voss, 1968) - are commonly referred to as "harlequin" octopuses in reference

to the black and white stripes and spots which cover the entire body [Fig 1]. However, unlike

most other octopus species, which generally exhibit semelparity and only reproduce near the end

of their lifespan (Mather, 2006), *O. chierchiae* is iteroparous (Rodaniche, 1984) with female

individuals having been observed to produce up to eight clutches of eggs over the course of their

lifespan in our laboratory. Another trait that sets *O. chierchiae* apart from most other octopus

species is its lack of a planktonic stage in its development, suggesting a limited potential for

dispersal. Hatchlings [Fig 2] have a mantle length of about 3.5mm (Rodaniche, 1984) and

immediately begin their demersal lifestyle upon leaving the egg, making it possible to raise them

in simple, conventional aquaria. Overall, *O. chierchiae's* iteroparous reproductive strategy,

observed lifespan of over two years, and lack of a planktonic developmental stage makes

breeding and maintaining the species in captivity a practical and viable pursuit, contributing to

its growing potential as a model organism (Grearson 2021). As its candidacy and status as the

marine model organism develops, the need to locate, monitor, and study wild individuals of the

species grows as well, especially because of the rapid change in abundance and distribution of

intertidal species caused by anthropogenic climate change (Helmuth et. al., 2006). Specifically,

the ability to identify and monitor the octopuses on an individual basis would be crucial for both

in situ studies (Arechavala-Lopez 2019) as well as controlled laboratory research involving

multiple individual animals (Morse 2017).

Fig 1. One-year-old O. Chierchiae adult

Fig 2. One-day-old O. Chierchiae hatchling

Current studies on wild cephalopods most commonly involve tag-recapture studies

(Semmens et al. 2007). However, as Semmens et al. discuss, external tags can harm octopuses'

delicate tissues, cannot be applied to animals that are too small, and can fall off or be deliberately

removed by the tagged animal. Chemical tagging, chemical analyses of cephalopod hard parts,

geographic differences in parasite fauna, molecular genetics, and satellite data of detected signals

are amongst some of the other rising technological tracking methods (Semmens et al. 2007). *O.*

chierchiae's delicate and diminutive bodies make longitudinal studies of wild *O. chierchiae*

through the aforementioned techniques both difficult to complete and potentially dangerous to

the animals.

Contrary to the techniques mentioned above, photography serves as a largely

inexpensive, innocuous, non-intrusive, and widely accessible method to produce high-quality

data. The harmless nature of this approach as well as the virtual ubiquitousness of smartphones

that can record photos, videos, and geospatial data has enabled community science efforts to

make increasingly significant and effective contributions to scientific research on the abundance,

distribution, phenology, and ecology of multiple species (Ballance, 2018; Chandler et. al., 2017,

Dickinson et al 2012). In situ research in particular has benefited tremendously from the growing

availability of photodata, as many inferences about wild animal populations can be drawn from

both the content and the context of these photographic images, especially if members of the

species of interest are individually and uniquely identifiable. The employment of such

photography-analysis techniques has allowed researchers to estimate population dynamics

(Karanth & Nichols, 1998), determine individual movement and home ranges (Oliveira-Santos et

al., 2010) and quantify disease dynamics within populations (Geers et al., 1997). Besides its

exceptional accessibility, the photography-based approach offers yet another distinctive

advantage in that it may be applied retrospectively to existing photobanks (Johnston et al., 2017).

All the aforementioned advantages along with the continual development of new computer-aided

photo identification methods (Kelly 2001) and analytical techniques makes photography an ideal

approach for monitoring and studying soft-bodied cephalopods like *O. chierchiae*.

As far as we know, *Wunderpus photogenicus* is the only octopus species in which

individuals have been shown to be identifiable by photos of their individually unique body

patterns, making it possible to use photoidentification methods to study this species (Huffard et

al., 2008). Here, we test whether *O. chierchiae* also have identifiable, individually unique body

patterns, and whether these remain constant throughout their lifetimes, allowing for

photoidentification over long timespans by untrained volunteers.

Materials and Methods

<u>Collection and Rearing</u>

Two male and two female adult *Octopus chierchiae* were obtained from a fully-licensed

commercial collector in Nicaragua during 2017 and 2018. We housed them individually in 1-

gallon(3.79 liter) glass aquaria lined with gravel, light sand and bivalve shell fragments [Fig 1,

Fig 3c], provided PVC tubes, glass vials or hollow rocks as shelter, and fed them bait shrimp

(*Crangon franciscorum* and *Palaemonetes sp.*) and yellow shore crabs (*Hemigrapsus*

oregonensis) ad libitum daily. All aquaria in our laboratory were filled and maintained with

artificial seawater (32-36 ppt) at 22-24 ℃.

Beginning in the summer of 2018, we allowed adult *O. chierchiae* (F0 generation) to

mate with other wild caught individuals, assumed to be unrelated, by temporarily moving a male

into a female's tank, waiting for up to one hour for copulation to occur, then moving the male

back into his tank. Females usually laid eggs within two days of mating, which hatched after

about 45 days. Upon hatching, individuals were transferred via a large pipette (turkey baster) into

glass jars with screen lids (9 cm tall, 8 cm diameter) containing a variety of substrates including

dark sand, light sand, bivalve shell fragments, snail shells, and gravel. To avoid issues of

cannibalism, territorial disputes, and other aggressive interactions, the animals were reared in

individual jars, all within the same 40 gallon tank(151.42 liter)[Fig 3a], initially on a diet of

primarily marine amphipods and brine shrimp (*Artemia salina*). As they grew and matured, the

animals were transitioned to a diet primarily consisting of bait shrimp (*Crangon franciscorum*

and *Palaemonetes sp.*) and yellow shore crabs (*Hemigrapsus oregonensis)* and provided empty

snail shells for shelter. Individuals were fed live foods ad libitum when available, and otherwise

were offered a piece of shrimp every other day. We transferred mature animals of this generation

(F1) individually to larger perforated plastic jars in a separate 20 gallon(75.71 liter) tank[Fig

3b]. In order to reduce the risk of genetic strain caused by inbreeding, we selected the least-

related male-female pairs and moved them to 1-gallon(3.79 liter) tanks for mating[Fig 3c]. After

copulation, we transferred the male back to his plastic jar and provided the female with hollow

rocks, PVC pipes, glass vials, and coral fragments in which to lay eggs. Following these methods

we were able to successfully close the lifecycle and obtain a generation (F2) bred and raised in

captivity. Captive-bred *O. chierchiae* that survived the first few weeks of life typically lived for

at least a year, with our longest-lived captive-bred octopus being a male born in July 2018 who

lived for 22 months and fathered at least 4 different clutches of offspring before his death in June

2020. We hatched a total of 156 individuals from 10 clutches over the course of this study (3 to

32 hatchlings per clutch) with an average body weight of 17.4 mg (min = 9.5 mg, max = 25.6

mg)[S1 Dataset].

Fig 3a. Nursery tank where animals were reared in individual glass jars with screen lids from

hatching until sexual maturity

Fig 3b. General tank where mature animals were housed in individual perforated plastic jars

Fig 3c. Individual mating tank where matings occurred and eggs were laid

Photography

We photographed individual *O. chierchiae* with an iPhone 6s smartphone and a 4K Sony

camcorder (model HDR-XR260V) from June 2018 to July 2020. In order to obtain a

photographic record of each individual's development over the course of their lifetime, we

photographed lab-reared *O. chierchiae* (n=25) weekly from a directly overhead angle [Fig 4],

without removing them from their containers. All of the photographic images used in this study

were taken via this described overhead method except for one pair of photographs which was

taken from the side of the transparent glass jar. Body patterns of *O. chierchiae* become visible to

161 the naked eye when individuals are around two weeks old (~ 8 mm mantle length), but can be

162 observed with magnification as early as the fifth day after they hatch (Fig 5).

164 Fig 4a. Timeline of *O. chierchiae* individual (hatched Nov 12, 2018) showing consistent body

165 patternings over a 5 month timespan. The bottom row duplicates these photographs, with

166 analogous patterns color coded.

167 Fig 4b. Timeline of *O. chierchiae* individual (hatched May 6, 2019) showing consistent body

168 patternings over a 6 month timespan. The bottom row duplicates these photographs, with

169 analogous patterns color coded.

170 Fig 4c. Timeline of *O. chierchiae* individual (hatched Nov 26, 2018) showing consistent body

171 patternings over a 7 month timespan. The bottom row duplicates these photographs, with

172 analogous patterns color coded.

173 Fig 4d. Timeline of Octopus individual (hatched May 6, 2019) showing consistent body

174 patternings over a 4 month timespan. The bottom row duplicates these photographs, with

175 analogous patterns color coded.

176 Fig 5. Five-day-old *O. chierchiae* hatchling already displaying a stripe pattern

178 <u>Survey Design and Evaluation</u>

179  We created two surveys following the methods of Huffard et. al. (2008) from this

180 photobank, solicited participants via social media, and collected responses via google forms.

181 Survey A[S2 File] consisted of a series of slides showing two photographs (taken from 0 days to

182 5 months apart) of *O. chierchiae* individuals side by side. Participants of Survey A were asked to

183 mark whether the 2 photographs were of the same (Fig 6) or different (Fig 8) individuals. Survey

B[S3 File] consisted of the same series of slides as Survey A, but included pattern traces created

in Adobe Illustrator below the photographs (Fig 7, Fig 9). We compiled survey responses into

excel spreadsheets and graded them via a python script[S4 Code] using the pandas v1.3.3

package (McKinney 2010). Statistical analysis of our resulting data were calculated using the

ttest_ind and ttest_1samp functions from the scipy.stats v1.7.1 library (Virtanen 2020) with a

predetermined significance level of 0.05.

Fig 6. Question #17 of survey A. Matching pictures of the same *O. chierchiae* individual taken

on Jul 26, 2019 (left) and Aug 2, 2019 (right)

Fig 7 Question #17 of survey B. Matching pictures of the same *O. chierchiae* individual taken on

Jul 26, 2019 (left) and Aug 2, 2019 (right)

Fig 8 Question #1 of survey A. Non-matching pictures of 2 different *O. chierchiae* individuals,

both taken on Aug 29, 2019

Fig 9 Question #1 of survey B. Non-matching pictures of 2 different *O. chierchiae* individuals,

both taken on Aug 29, 2019

Results

Over the course of the study, we observed that *O. chierchiae* individuals' body patterns

were consistent throughout their lifetimes, confirmed by photographic records [Fig 4]. All

members of our research group were able to recognize individual animals based on their body

patterns. Next we investigated if untrained observers would be able to do the same.

Both Survey A (1 sample t-test: t-statistic = 10.90, p-value = 4.17e-10) and Survey B (1

sample t-test: t-statistic=11.14, p-value=2.84e-10) yielded results significantly different from

50% – the expected value if participants were unable to distinguish individual *O. chierchiae*.

Survey A (n = 22 responses [S5 Dataset]) had an average score of 75.54% correct answers,

standard error of the mean = 2.34%. On average, participants missed 25.16% of total possible

matches (false negative) and erroneously matched 22.11% of non-matching individuals (false

positive), whereas Survey B (n = 22 responses [S6 Dataset]) had an average score of 74.27%,

standard error of the mean = 2.18%. On average, participants missed 25.65% of total possible

matches and erroneously matched 25.21% of non-matching individuals.

Discussion

Our results [Fig 10, Fig 11] show that *O. chierchiae* individuals are distinguishable to a

majority of untrained observers, as 63.6% and 54.5% of all participants scored above 75% on

surveys A and B respectively. There were five questions on each survey where over 50% of

participants answered incorrectly and one question on each where it was an even split between

correct and incorrect responses. Upon specific examination of the problematic questions, the

increased difficulty could be explained by a combination of subpar photo quality, the animal

being in dark morph [Fig 12] thus making its stripes harder to discern, body pattern being

obscured in some manner, or the animal being viewed from different angles. Thus we conclude

that *O. chierchiae* can be consistently identified throughout their lives via their individually

unique body patterns by both trained and untrained observers, when a clear view of the body

patterns is provided.

Fig 10. Scores of each participant of survey A (n=22 survey responses), illustrating total score

(Blue), False Negative (Red) and False Positive (Orange) when assessing images.

Fig 11. Scores of each participant of survey B (n=22), illustrating total score (Blue), False

Negative (Red) and False Positive (Orange) when assessing images.

Fig 12. A pair of images depicting the same *O. chierchiae* individual in dark (left) and

light(right) morph

Interestingly, the average score was lower on the survey where body pattern traces were

provided for assistance, perhaps showing variance in how individual participants recognise

identifying features. Like the results for *Wunderpus photogenicu*s, false positives were more

common than false negatives in both survey groups when photo identifying *O. chierchiae*

(Huffard et. al., 2008). Because some participants responded to both surveys, we cannot combine

the results of surveys B and C. Consequently, unlike Huffard et al. (2008), we considered the

results of both surveys separately.

Having hatched, reared and documented the individuals used in the surveys in captivity

over extensive time spans – often over the course of their entire life history – we observed no

evidence that body patterns change over the lifetimes of individuals. Our surveys confirm that

these permanent body patterns are a viable method of identifying the same individual at different

points in time, even by volunteers with little to no experience with octopuses. To our knowledge,

this is the first evidence that individually unique body patterns can be used by volunteers to

consistently identify the same individual octopus throughout its development, from juvenile to

adult.

This information could be used to design non-invasive studies monitoring populations of

O. chierchiae over time in situ. Based on our results that participants could consistently identify

whether or not two photos depicted the same octopus, we recommend that any future community

science effort to study *O. chierchiae* via photo identification should take the form of survey A,

even if it means a large number of survey questions must be conducted to pair images.

Smartphone apps or websites can be utilized to gather responses en masse and the process may

even be automated with enough responses and high quality image data.

In our time working with this species, we have observed clutches of up to 32 hatchlings

successfully leaving the nest, and an individual female who, in our care, laid 8 fertile clutches of

eggs over the course of just over two years. Although they bear the rare distinction among

octopuses of being iteroparous, *O. chierchiae* seems able to produce a maximum of around 240

offspring in its lifetime, as opposed to the many thousands other species might produce. Because

O.chierchiae have so few young, which begin their demersal lifestyle from hatching and thus do

not disperse as plankton, photo identification has potential to be a useful tool for in situ

demographic studies even on very young individuals. While it is known that there can be

variation in the founder chromatophores of individual octopus hatchlings of the same species

(Ylitalo et. al., 2014), and that *Octopus chierchiae* can begin to display the characteristic stripe

pattern as soon as five days after hatching (Fig 5), we did not begin photographing animals for

this study until they were at least four weeks old. Thus, more work is needed to determine

whether the individually unique body patterns of *O. chierchiae* are fixed from birth, and if not, at

what point in early development the body patterns become permanent.

In addition, we hypothesize that the application of photo identification is not limited to *O.*

chierchiae and *W. photogenicus*, or even the octopuses with conspicuously semi-permanent body

patterns, like the blue-ringed octopuses (e.g. Figure 1 in Huffard et al 2008). Since it is known

that octopuses have a fixed skin anatomy (Packard and Hochberg, 1977; Messenger, 2001), and

that noticeable differences in skin anatomy between individuals can exist in octopuses with a

wide repertoire of body patterns (Huffard, 2006), we think it would be worthwhile to investigate

whether individuals of other octopus species can be identified over their lifetimes by their semi-

permanent or recurring body patterns or skin textures.

Lastly, we hypothesize that *O. chierchiae*'s individually unique, lifelong body patterning

may play a role in intraspecific behavior in the wild. Octopus species are known to have the

ability to detect conspecifics (Walderton 2011), high visual acuity, and the ability to remember

(Hochner 2006). At least one species, *Octopus vulgaris*, is thought to have the ability of

individual conspecific recognition (Tricario et. al., 2011). The harlequin octopuses, with their

bold semi-permanent body patterns, also "rely less upon changes in chromatophores than upon

perceptions of postures and positions," (Moynihan, 1985). In our view, these factors make it

worthwhile to explore whether individual recognition based on visual cues plays a role in the

lives of wild *O. chierchiae* and other octopus species.

Acknowledgements

This study would not have been possible without the hard work and support of our laboratory's

O. chierchiae husbandry team over the past few years. We'd like to thank Dr. Christine Huffard

for devoting so much time and effort to help us with this paper. We could not have gotten this far

without your patience and wisdom. Thank you to Dr. Caroline Williams from UC Berkeley and

Bret Grasse from the Woods Hole Marine Biological Laboratory for your correspondence and

guidance, and Richard Ross from the California Academy of Sciences for your contributions to

our husbandry methods.

References

Arechavala-Lopez P, Minguito-Frutos M, Follana-Berná G, Palmer M. Common octopus settled

in human-altered Mediterranean coastal waters: from individual home range to population

dynamics. ICES J Mar Sci. 2019;76(2):585–97.

Auffenberg K, Bank R, Bieler R, Bouchet P, Faber M, Finn J, et al. MolluscaBase. Catalogue of

Life; 2021.

Ballance LT. Contributions of photographs to cetacean science. Aquat Mamm. 2018;44(6):668–

82.

Boal J. Social recognition: a top down view of cephalopod behaviour. Vie et Milieu/Life &

Environment. 2006:69-79.

Brewer RS, Norcross BL, Chenoweth E. Temperature- and size-dependent growth and

movement of the North Pacific giant octopus (Enteroctopus dofleini) in the Bering Sea. Mar Biol

Res. 2017;13(8):909–18.

Caldwell RL, Ross R, Rodaniche A, Huffard CL. Behavior and body patterns of the Larger

Pacific Striped Octopus. PLoS One. 2015;10(8):e0134152.

Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, et al. Contribution of citizen

science towards international biodiversity monitoring. Biol Conserv. 2017;213:280–94.

Dickinson JL, Shirk J, Bonter D, Bonney R, Crain RL, Martin J, et al. The current state of citizen

science as a tool for ecological research and public engagement. Front Ecol Environ.

2012;10(6):291–7.

Elliser CR, MacIver KH, Green M. Group characteristics, site fidelity, and photo-identification

of harbor porpoises,Phocoena phocoena, in Burrows Pass, Fidalgo Island, Washington. Mar

Mamm Sci. 2018;34(2):365–84.

Geers P, Puers B, Goedseels V, Wouters P. Electronic identification, monitoring and tracking of

animals. Wallingford, England: CABI Publishing; 1996.

Grearson AG, Dugan A, Sakmar T, Dölen G, Gire DH, Sivitilli DM, et al. The lesser Pacific

striped octopus, Octopus chierchiae: an emerging laboratory model for the study of octopuses

[Internet]. bioRxiv. 2021. Available from: http://dx.doi.org/10.1101/2021.08.04.455148

Hackett J, Harrington S. Beasts of the Deep: Sea Creatures and Popular Culture. Bloomington:

Indiana University Press; 2018.

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et al. Array

programming with NumPy. Nature. 2020;585(7825):357–62.

Helmuth B, Mieszkowska N, Moore P, Hawkins SJ. Living on the edge of two changing worlds:

Forecasting the responses of rocky intertidal ecosystems to climate change. Annu Rev Ecol Evol

Syst. 2006;37(1):373–404.

Hochberg FG, Camacho-García YE, Wehrtmann IS, Cortés J. Squids and octopuses. Marine

biodiversity of Costa Rica, Central America. Springer; 2009.

Hochner B, Shomrat T, Fiorito G. The octopus: a model for a comparative analysis of the

evolution of learning and memory mechanisms. Biol Bull. 2006;210(3):308–17.

Huffard CL, Caldwell RL, DeLoach N, Gentry DW, Humann P, MacDonald B, et al.

Individually unique body color patterns in octopus (Wunderpus photogenicus) allow for

photoidentification. PLoS One. 2008;3(11):e3732.

Hupman KE, Pawley MDM, Lea C, Grimes C, Voswinkel S, Roe WD, et al. Viability of photo-

identification as a tool to examine the prevalence of lesions on free-ranging common dolphins

(Delphinus sp.). Aquat Mamm. 2017;43(3):264–78.

Jatta G, Elenco dei Cefalopodi delle "Vittor Pisani". Bollettino della Societa di Naturalisti in

Napoli. (series 1) 1889;3(1): 63-67.

Johnston DR, Rayment W, Slooten E, Dawson SM. A time-based method for defining

associations using photo-identification. Behaviour. 2017;154(9–10):1029–50.

Kaestner A. Pages 394-424 in Invertebrate Zoology. Vol. 1. New York: Interscience Publishers;

1967.

Karanth KU, Nichols JD. Estimation of tiger densities in India using photographic captures and

recaptures. Ecology. 1998;79(8):2852–62.

Kelly MJ. Computer-aided photograph matching in studies using individual identification: An

example from Serengeti cheetahs. J Mammal. 2001;82(2):440–9.

Kerstitch AN. Sea of Cortez Marine invertebrates: A guide for the Pacific Coast, Mexico to

Ecuador. 1st ed. Monterey, CA: Sea Challengers; 1989.

Kozloff EN. Pages 447-462 in: Invertebrates. Philadelphia: Saunders College Publishing; 1990.

Mather JA. Behaviour Development: A Cephalopod Perspective. International Journal of

Comparative Psychology. 2006;19(1):98–115.

Mather JA. Marine invertebrates: communities at risk. Biology (Basel). 2013;2(2):832–40.

McKinney W. Data Structures for Statistical Computing in Python. In: Proceedings of the 9th

Python in Science Conference. SciPy; 2010.

Meglitsch PA. Pages 546-563 in Invertebrate Zoology. London: Oxford University Press; 1967.

Messenger JB. Cephalopod chromatophores: neurobiology and natural history. Biol Rev Camb

Philos Soc. 2001;76(4):473–528.

Morse P, Zenger KR, McCormick MI, Meekan MG, Huffard CL. Chemical cues correlate with

agonistic behaviour and female mate choice in the southern blue-ringed octopus,Hapalochlaena

maculosa(Hoyle, 1883) (Cephalopoda: Octopodidae). J Molluscan Stud. 2017;83(1):79–87.

Moynihan M. Communication and Noncommunication by Cephalopods. Bloomington, MN:

Indiana University Press; 1985.

Oliveira-Santos LGR, Machado-Filho LCP, Tortato MA, Brusius L. Influence of extrinsic

variables on activity and habitat selection of lowland tapirs (Tapirus terrestris) in the coastal sand

plain shrub, southern Brazil. Mamm Biol. 2010;75(3):219–26.

Packard A, Hochberg FG. Skin patterning in Octopus and other genera. InSymp. Zool. Soc. Lond

1977 (Vol. 38, pp. 191-231).

Rodaniche AF. Iteroparity in the lesser Pacific striped octopus Octopus chierchiae (Jatta, 1889).

Bulletin of Marine Science. 1984 Jul 1;35(1):99-104.

Ruppert EE, Fox RS, Barnes RD. Page 990 in: Invertebrate Zoology. A functional evolutionary

approach. 7th ed. Brooks/Cole, Thomson Learning learning, Inc; 2004.

Semmens JM, Pecl GT, Gillanders BM, Waluda CM, Shea EK, Jouffre D, et al. Approaches to

resolving cephalopod movement and migration patterns. Rev Fish Biol Fish. 2007;17(2–3):401–

23.

Theimer TC, Ray DT, Bergman DL. Camera angle and photographic identification of individual

striped skunks: Individual Striped Skunk Identification. Wildl Soc Bull. 2017;41(1):146–50.

Tricarico E, Borrelli L, Gherardi F, Fiorito G. I know my neighbour: individual recognition in

Octopus vulgaris. PLoS One. 2011;6(4):e18710.

Vaissi S, Farassat H, Akmali V, Sharifi M. Consistency of coloration pattern and applicability of

photo identification method as a tool to identify individuals of the kaiser's mountain newt,

Neurergus kaiseri. Russ J Herpetol. 2018;25(4):311.

Vendetti J. Squids, octopuses, nautilus, and ammonites. 2006 [cited 18 november 2021]. In: The

Cephalopoda [Internet]. Berkeley: University of California Museum of Paleontology; 2006.

Available from: https://ucmp.berkeley.edu/taxa/inverts/mollusca/cephalopoda.php.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0:

fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17(3):261–72.

Voss GL. Biological Investigations of the Deep Sea. 39. Octopods from the R/V Pillsbury

Southwestern Caribbean Cruise, 1966, with a description of a new species, Octopus zonatus.

Bulletin of Marine Science. 1968 Jul 2;18(3):645-59.

Walderon MD, Nolt KJ, Haas RE, Prosser KN, Holm JB, Nagle GT, et al. Distance

chemoreception and the detection of conspecifics in Octopus bimaculoides. J Molluscan Stud.

2011;77(3):309–11.

Witteveen BH, Wynne KM. Site fidelity and movement of humpback whales (Megaptera

novaeangliae) in the western Gulf of Alaska as revealed by photo-identification. Can J Zool.

2017;95(3):169–75.

Ylitalo HA, Watling L, Toonen RJ. First description of hatchlings and eggs of Octopus

oliveri(Berry, 1914) (Cephalopoda: Octopodidae). Molluscan Res. 2014;34(2):79–83.

Supporting Information Captions

S1 Dataset. Octopus chierchiae Hatchling Weights

S2 File. Survey A

S3 File. Survey B

S4 Code. Survey Grader

S5 Dataset. Survey A Responses

S6 Dataset. Survey B Responses

Figure1

Figure2

Figure3a

Figure3b

Figure3c

May 17, 2019   Jun 8, 2019   Aug 2, 2019   Oct 1, 2019

May 17, 2019   Jun 8, 2019   Aug 2, 2019   Oct 1, 2019

Figure4a

May 3, 2019   Jun 8, 2019   Jul 1, 2019   Aug 1, 2019   Sep 17, 2019   Nov 17, 2019

May 3, 2019   Jun 8, 2019   Jul 1, 2019   Aug 1, 2019   Sep 17, 2019   Nov 17, 2019

Figure4b

April 8 June 13 July 1 October 1 November 10

April 8 June 13 July 1 October 1 November 10

Figure4c

Jul 17, 2019   Aug 1, 2019   Sep 17, 2019   Oct 1, 2019   Nov 10, 2019

Jul 17, 2019   Aug 1, 2019   Sep 17, 2019   Oct 1, 2019   Nov 10, 2019

Figure4d

Figure5

Question

○ match

○ no match

Figure6

○ match

○ no match

Figure7

Question

○ match

○ no match

Figure8

○ match

○ no match

Figure9

Figure10

Figure11

Figure12