The Grainyhead/LSF transcription factor GRH-1 is rhythmically required for molting

Milou W.M. Meeuse[1,2], Smita Nahar[1], Kathrin Braun[1], Helge Großhans[1,2,3]

[1] Friedrich Miescher Institute for Biomedical Research (FMI), Maulbeerstrasse 66, CH-4058 Basel

[2] University of Basel, Petersplatz 1, CH-4001 Basel

[3] Correspondence to: helge.grosshans@fmi.ch

Abstract

Molting, that is, the synthesis and shedding of a cuticular exoskeleton, is a defining characteristic of ecdysozoa. In nematodes such as *C. elegans*, molts rhythmically terminate each of four larval stages. The molting cycle is tightly coupled to the rhythmic expression of thousands of genes. The mechanisms that support the regular molting cycle and oscillatory gene expression have remained largely elusive. Here, we performed an RNAi-based screen for transcription factors required for molting to identify potential components of a molting clock. We find that depletion of GRH-1, BLMP-1, NHR-23, NHR-25, MYRF-1 or BED-3 impairs progression through the molting cycle. We characterize GRH-1, a Grainyhead/LSF transcription factor whose functions in *C. elegans* development have remained largely unexplored, but whose orthologues in other animals are known to be key epithelial cell fate regulators. We show that GRH-1 depletion causes a dose-dependent extension of molt duration, with severe depletion causing defects in cuticle formation and shedding, and larval death. GRH-1 is required repetitively, during a specific time window prior to each larval molt. This rhythmic activity is consistent with the rhythmic accumulation of GRH-1 protein. These features are consistent with GRH-1 functioning as a key component, or high-level output, of the a gene regulatory network that controls molting. As its mammalian orthologues, as well as those of BLMP-1 and NHR-23, have been implicated in rhythmic homeostatic skin regeneration in mouse, the mechanisms underlying rhythmic *C. elegans* molting may be conserved beyond nematodes.

Introduction

C. elegans larval development subdivides into four larval stages, each terminated by a molt. In a first step, termed apolysis, the connections of the existing cuticle to the underlying epidermis are severed. Subsequently, a new cuticle is synthesized, before the old cuticle is shed in a final step termed ecdysis. Traditionally, the time of molting is equated with lethargus, a period of relative behavioral quiescence, when animals stop feeding. For simplicity we will follow this tradition here, but note that additional events required for successful molting precede lethargus (Cohen et al., 2020; Cohen and Sundaram, 2020); (Tsiairis and Großhans, 2021).

Molts occur with a regular period of 7– 8 hr at 25°C, and a clock-type mechanism has been invoked to explain this regularity (Monsalve and Frand, 2012; Tsiairis and Großhans, 2021). Such a clock mechanism may also explain, and be partially based on, the rhythmic expression of thousands of genes that is coupled to the molting cycle (Hendriks et al., 2014; Kim et al., 2013; Meeuse et al., 2020; Turek and Bringmann, 2014). However, the components of this clock, and accordingly their wiring, have remained largely elusive (Tsiairis and Großhans, 2021).

Recently, we showed that rhythmic RNA polymerase II recruitment drives RNA level oscillations during larval development (Hauser et al., 2021). This finding suggests that rhythmically active transcription factors are components of the underlying machinery, or core oscillator, and thus presumably also the molting cycle clock. To identify possible candidates, we screened through a selection of rhythmically expressed transcription factors (TFs), assuming that rhythmic transcription would be a parsimonious (though not necessarily the only) mechanism of achieving rhythmic transcription factor activity. From a set of 92 such transcription factors (Hendriks et al., 2014), we sought to identify those that affected duration of, or progression through, molts. We identified six hits, including the nuclear hormone receptors NHR-23 and NHR-25 that were previously linked to molting (Frand et al., 2005; Gissendanner et al., 2004; Gissendanner and Sluder, 2000; Johnson et al., 2021; Kostrouchova et al., 1998; Kostrouchova et al., 2001). We characterize the function of GRH-1, the sole *C. elegans* member of the phylogenetically conserved LSF/Grainyhead family (Venkatesan et al., 2003).

Grainyhead proteins are key regulators of epithelial cell fates that function in differentiation, maintenance, integrity and repair of different epithelial tissues in animals (Sundararajan et al., 2020). *C. elegans* GRH-1 was previously found to be required for embryonic cuticle formation, but its postembryonic function and regulation have remained unknown. We report that GRH-1 accumulates

rhythmically and promotes molting through its activity in a specific window during each larval stage. Its depletion delays the onset of ecdysis in a dose-dependent manner to the point that animals severely depleted arrest development and die by bursting through a defective cuticle. The results, together with the validation of an additional hit, BLMP-1, in separate work by us and others (Hauser et al., 2021; Stec et al., 2021) provide new insights into the transcriptional mechanisms that support rhythmic molting and identifies potential molting clock components. The fact that proteins orthologous to GRH-1 as well as to additional screen hits also function in rhythmic homeostatic skin regeneration in mouse (Magnusdottir et al., 2007; Steinmayr et al., 1998; Telerman et al., 2017; Wilanowski et al., 2008) suggest mechanistic similarities between this process and the molting process of nematodes.

Results

A targeted screen identifies transcription factors involved in molting

Since *C. elegans* molting cycles are tightly coupled to oscillatory gene expression (Meeuse et al., 2020), which in turn is driven by rhythmic transcription (Hauser et al., 2021), we sought to identify rhythmically active transcription factors involved in molting. Hence, we performed an RNAi screen targeting 92 transcription factors that exhibit transcript level oscillations according to our previous annotation (Hendriks et al., 2014). Specifically, we screened for aberrant developmental progression or molt execution. To obtain such information, we examined luciferase activity in animals that express a luciferase transgene and that are grown in the presence of D-luciferin (Meeuse et al., 2020; Olmedo et al., 2015). This assay detects lethargus by a drop in luminescence at the level of individual animals, allowing us to quantify durations of molts, intermolts and, as a sum of the two, entire larvael stages for several animals per condition. We depleted the transcription factors by feeding animals on RNAi-expressing bacteria. To control for differences in larval growth among RNAi conditions unrelated to target protein depletion, we performed the experiment in parallel on RNAi–deficient *rde-1(ne219)* mutant animals (Tabara et al., 1999) (**Figure 1A**).

We expected mutants with the following phenotypes: i) altered developmental tempo, which could be faster or slower, and specific for a particular oscillation phase or continuously through the cycle, ii) increased variability of developmental tempo among individuals, and iii) cessation of development and/or failed molts.

By plotting the luminescence intensities sorted by entry into the first molt in a heatmap (**Figure 1B,C**), we identified six genes whose depletion caused abnormal progression through development following atypical molts (*nhr-23, myrf-1* and *grh-1*; **Figure 1B,D, Figure S1**), or aberrant duration of molts (*bed-3, blmp-1* and *nhr-25*; **Figure 1C,D, Figure S2**).

The nuclear hormone receptors NHR-23 and NHR-25 have previously been shown to function in molting (Frand et al., 2005; Gissendanner et al., 2004; Gissendanner and Sluder, 2000; Johnson et al., 2021; Kostrouchova et al., 1998; Kostrouchova et al., 2001), and we and others have described functions of BLMP-1 in oscillatory gene expression and cuticle formation (Hauser et al., 2021; Sandhu et al., 2021; Stec et al., 2021). Here, we sought to establish further the function of GRH-1, first, because we wished to understand whether class III phenotypes can indeed be predictive of clock dysfunction, second, because orthologous Grainyhead(-like) proteins function in epithelial cell fate control and remodeling of epithelial tissues in other animals (Sundararajan et al., 2020), suggesting a potentially conserved function and third, because its function in *C. elegans* development has remained largely unexplored (Venkatesan et al., 2003).

Figure 1: RNAi screen for 'oscillating' transcription factors revealed aberrant duration or progression of molting

A, Heatmap showing trend-corrected luminescence (Lum) of wild-type (WT, strain HW1939, top) and RNAi-deficient (*rde-1(ne219)*), strain HW2150, bottom) animals expressing luciferase from the *eft-3* promoter grown on mock

(empty vector, EV) RNAi in a temperature-controlled incubator set to 20°C. Each line represents one animal. Hatch is set to t = 0 hours and traces are sorted by entry into the first molt. Darker color corresponds to low luminescence and is associated with the molt.

B, C Heatmap showing trend-corrected luminescence as in A, for indicated RNAi conditions causing altered numbers (B) or durations (C) of molts, respectively.

D, Quantification of the percentage of animals entering specific molts on indicated RNAi conditions. Shown are the last molts observed for animals in each condition; e.g., 100 % of GRH-1 depleted fail to progress beyond M2.

GRH-1 is repetitively required for normal duration and progression through molts

Injection of double-stranded RNA targeting *grh-1* into the germline of L4 stage larval animals causes embryonic lethality in the next generation (Venkatesan et al., 2003). Consistent with this finding, we observed a penetrant embryonic lethality phenotype in *grh-1(0)* mutant animals (data not shown). To bypass this defect and investigate the role of GRH-1 during larval development, we performed controlled GRH-1 depletion in larvae using the auxin-inducible degradation (AID) system (Zhang et al., 2015). We tagged *grh-1* endogenously with *degron::3xflag* to generate allele *grh-1(xe135)*, and expressed the plant-specific F-box protein TIR1 as a transgene, generating a strain that, for simplicity's sake, we will refer to as *aid::grh-1* in the following. When we hatched *aid::grh-1* embryos in the presence of 250 µM auxin, animals failed to develop beyond the first molt (M1), i.e., none of them entered M2 (**Figure 2A**).

If GRH-1 were a component of the core molting machinery, we would predict that it be required for each molt. Indeed, when we depleted GRH-1 by addition of auxin at the beginning of the L2, L3 and L4 stage, respectively, animals in each case failed to develop beyond the next molt, i.e., M2, M3 and M4, respectively (**Figure 2B-D**). We conclude that GRH-1 is repetitively required during development, for completion of each molt.

Figure 2: GRH-1 is required for ecdysis of every molt.

A-D, Heatmaps showing trend-corrected luminescence (Lum) of *aid::grh-1* animals constitutively expressing luciferase (HW2434). t = 0h corresponds to time of plating embryos, which subsequently hatch at different times. Arrow indicates time point when 250 μM auxin was added, i.e. prior to the first molt (A; M1), M2 (B), M3 (C) or M4 (D) larval stage. Note that for technical convenience in (A), auxin was provided at time of plating. Animals are sorted by entry into M1 (A), M2 (B), M3 (C), M4 (D), respectively. Note that embryos in (C) were younger than those in (D) when plated so that addition of auxin 40h after plating coincides with L3 in (C) but L4 in (D).

We observed an increased duration of the last observed molt that GRH-1-depleted animals entered (**Figure 2**). To examine this phenotype further, we exposed *aid::grh-1* animals to varying auxin concentrations to titrate GRH-1 depletion. Although auxin concentrations of ≥400 nM at hatching yielded a fully penetrant M1 exit phenotype, a dose-dependent decrease in phenotype penetrance occurred below this concentration (**Figure 3A, S3A**). When we quantified the developmental tempo for animals that, at the lowest auxin concentrations tested (53 nM, 79 nM and 119 nM), completed the first three molts, we observed a dose-dependent lengthening in M1, M2 and M3 (**Figure 3B, S3 – S5**). By contrast, no or little change occurred for the duration of the intermolts preceding the lengthened molts (**Figure 3C, S3 – S5**). Hence, we conclude that the activity of GRH-1 is required for normal duration of the recurring molts, with strong depletion additionally causing molting failure.

Figure 3: GRH-1 depletion extends molt duration in a dose-dependent manner.

A, Quantification of the percentage of *aid::grh-1* animals constitutively expressing luciferase (HW2434) that enter indicated molts after hatching into increasing concentrations of auxin as indicated. Molts in single animals grown at 20°C were determined as in **Figure S1**. For concentrations up to 250 µM, similar results as for 400 nM were observed (**Figure S3**).

B, Boxplot showing the duration of M1, M2 and M3 of animals treated with indicated concentrations of auxin. Animals that revealed the phenotype at M1, M2 or M3 in (A) were excluded. Significant differences relative to 0 nM auxin are indicated. P-values were determined by Wilcoxon test. ns: not significant ,* p<0.05, ** p<0.01, *** p<0.001

Loss of GRH-1 results in fragile cuticles prone to rupturing

When we placed *grh-1::aid* animals on auxin-containing plates, they failed to shed the cuticle (ecdyse). To observe the molting process in greater detail, we used time-lapse DIC imaging to observe L1 animals transferred to an agar pad in a drop of M9 buffer, which allowed them to move. We could readily identify loosened cuticles at the tip of the head in wild-type and GRH-1-depleted animal (**Figure 4A,B**). Next, animals made spontaneous back-and-forth movements and the pharynx contracted rapidly (**Supplemental Movies 1 & 2**), after which wild-type animals shed the cuticle (ecdysed) (**Figure 4A, B**). By contrast, in GRH-1-depleted animals, the cuticle became even looser and more inflated in the head region (**Figure 4C**). Vesicles appeared in the cavity underneath the loosened cuticle (**Figure 4C**). Finally, the cuticle broke in the head region and the underlying tissue was extruded (**Figure 4C**).

These observations likely explain the striking increase and later drop in signal following the aberrant molt observed in the luciferase assays for GRH-1-depleted animals (**Fig. S1, Fig. 1**). Indeed, even under conditions where more modest GRH-1 depletion remains compatible with survival, we can observe

evidence of impaired cuticular integrity: Whereas lack of luciferin ingestion generates a low luminescence signal in molting wild-type animals, the signal increased in GRH-1-depleted animals during this time (**Figure 4D**).

We conclude GRH-1 is required for proper cuticle formation and timely ecdysis. In its absence, the duration of the molt becomes extended while a more fragile cuticle than typical of wild-type animals is formed. We note that although rupturing is lethal, animals also fail to complete development when grown on plate, a condition under which most animals do not burst but nonetheless fail to ecdyse (data not shown).

Figure 4: GRH-1 is required for cuticular integrity and normal ecdysis

A, B, Image sequence of N2 wild-type animals during M1. Lethargic animals were transferred to an agar-pad and observed by DIC at 63x magnification and imaged every 20 s (A) or every 4 s (B) respectively. Selected images with time stamps are shown. Dashed lines indicate cuticle boundaries detached from the body. As reported previously (Singh and Sulston, 1978), two different sequences of events were observed: the pharyngeal lining is removed prior

to ecdysis (A), or the pharyngeal lining is expelled after crawling out of the cuticle (B). Arrows indicate specific features of the molt. In (A): 1. Loosened cuticle, 2. Detachment of pharyngeal lining, 3. Crawling out of cuticle, 4. Final crawling out of cuticle. In (B): 1. Loosened cuticle, 2. Back-and-forth movements, 3. Crawling out of cuticle with pharyngeal lining still attached, 4. Pharyngeal lining expelled.

C, Image sequence of an L1 synchronized *aid::grh-1* animal (HW2418) plated on 250 μM auxin-containing plates, grown at 20°C. A lethargic animal was transferred to an agar-pad containing microscopy slide and images were collected every 1 sec, using DIC, 100x magnification. Selected images of Supplemental Movie 1 are shown. Time stamp (min:sec) is indicated. Arrows indicate phenotypic features: loosening of the cuticle (0:00); back-and-forth movements (0:05); inflation of the cuticle (2:46); vesicles underneath loosened cuticle (6:11); rupturing of the cuticle (6:20, 6:40).

D, Luminescence during molt was quantified for *aid::grh-1* animals (HW2418) grown in the presence of the indicated auxin concentrations.

GRH-1 protein levels oscillate and peak shortly before molt entry

Since GRH-1 is required repetitively, for proper execution of each of the four molts, and its mRNA levels oscillate (Hendriks et al., 2014; Meeuse et al., 2020), we wondered whether GRH-1 protein accumulation is also rhythmic. To test this, we examined a GRH-1::GFP fusion protein expressed from the endogenous *grh-1* locus. We observed the first detectable signal in elongating embryos (**Figure S6**), i.e., at the time when oscillatory gene expression initiates (Meeuse et al., 2020). In larvae, GFP signal accumulated in various tissues, including seam cells, vulva precursor cells, non-seam hypodermal cells, rectum cells, socket cells, and pharyngeal cells (**Figure S7**). Finally, in adults, which lack oscillatory gene expression (Meeuse et al., 2020), GFP::GRH-1 levels were greatly diminished or altogether absent in various tissues (**Figure S8**).

To study the larval dynamics of *grh-1* expression in more detail and relate it to developmental progression, we performed time-lapse microscopy of single animals grown in micro-chambers, acquiring fluorescence and bright-field images in parallel and with high temporal resolution as described previously (Meeuse et al., 2020). We observed rhythmic GRH-1 accumulation with a peak before molt entry (**Figure 5**). Thus, the temporal patterns and dynamics suggest a model whereby rhythmic changes in GRH-1 protein levels may generate rhythmic GRH-1 activity.

Figure 5: GRH-1 protein accumulates rhythmically before each molt

Time-lapse imaging of animals expressing endogenously tagged *gfp::grh-1*. Average +/- 95% confidence interval (cyan shading) are shown; gray shading indicates average time of molt.

Molting requires oscillatory GRH-1 activity

To test further whether the activity of GRH-1 was indeed rhythmic, we performed perturbation at different times during a larval stage, to assess whether we could observe a phase-specific response. We initiated GRH-1 degradation at variable times in L2 by adding auxin and monitored developmental progression using the luciferase assay. Plotting luminescence traces by the time when the animals entered molt 2 in a heatmap (**Figure 6A,B**) revealed a striking cutoff on the onset of the phenotype: addition of auxin up to 3 hours before the M2 was sufficient for phenotypic onset at M2 exit (**Figure 6B,C, green**). However, if animals received auxin later than this, i.e., within 3 hours from M2 entry or during M2, they progressed through M2 and instead exhibited the phenotype at M3 exit (**Figure 6B,C, blue**). We observed analogous outcomes when auxin was added in L3 (**Figure S9**). This period of auxin resistance is not explained by slow GRH-1 depletion kinetics since auxin depletes GRH-1 effectively within < 1h (**Figure 6D**). Hence, a period of time exists in each larval stage during which GRH-1 is dispensable for molt completion. In other words, and as we had hypothesized, GRH-1 exhibits rhythmic activity during larval development.

A

B

  C

D

Figure 6: GRH-1 functions prior to the molt

A,B Heatmap showing trend-correct luminescence (Lum) of *grh-1::aid* animals (HW2434) treated with vehicle (0.25% ethanol (A), or 250 µM auxin (B) at 29 hours after plating (white dashed line). Black intensities reflect low luminescence during lethargus (molt). Embryos of various stages were plated to obtain an asynchronously hatching population. Luminescence traces are sorted by entry into molt 2 (M2) such that traces of early hatched animals are at the bottom and those of late hatched animals are at the top.

C, Plot of phenotype onset over time of auxin application relative to entry into molt 2 (M2 entry). Dots represent individual animals from (B), colored according to whether the last observed molt in the luminescence trace was M2 (M2 exit phenotype; green) or M3 (M3 exit phenotype; blue).

D, Western blot revealing rapid GRH-1 depletion in the *grh-1::aid* strain (HW2434) upon addition of 250 µM auxin. A synchronized culture of animals was grown in liquid at 20°C. After 21 h (denoted t=0 h in the figure), the culture was split in two and either auxin or vehicle were added as indicated. Cultures were sampled hourly and protein lysates probed by Western blotting using anti-GRH-1 and anti-actin antibodies as indicated.

Discussion

Although molting is a fundamental feature of nematode development, little is known about the gene regulatory networks that control it. This is particularly true of the clock-type mechanisms thought to facilitate the regular occurrence of molts, and synchronization of the various processes that they encompasse. Considering that molting involves coordinated expression of tens or hundreds of extracellular matrix proteins, notably collagens, and their modifying enzymes, notably proteases and their inhibitors, transcription factors can be assumed to play important roles, but their identity and functions are not well established.

Here, we have identified six transcription factors with roles in molting and, potentially, oscillatory gene expression. Validating our approach, we identified NHR-23/Rora, arguably the best characterized molting transcription factor, whose depletion was previously shown to cause a failure of ecdysis and larval arrest (Frand et al., 2005; Gissendanner et al., 2004; Johnson et al., 2021; Kostrouchova et al., 1998; Kostrouchova et al., 2001; Kouns et al., 2011) and a role in regulating collagens and hedgehog-related genes (Kouns et al., 2011).

NHR-23 is orthologous to DHR3/Nr1f1, an ecdysone-controlled fly transcription factor important for metamorphosis (Kostrouchova 1998). In metamorphosis, DHR3 activates FTZ-F1 (Ou and King-Jones, 2013), the orthologue of a second one of our screen hits, NHR-25/Nr5a. Based on this orthology, and the finding that NHR-25 depletion impairs ecdysis, albeit more weakly and only in later molts than that of NHR-23, it was previously speculated that a regulatory interaction between the two proteins might be conserved in *C. elegans* and contribute to the molting process (Gissendanner et al., 2004; Gissendanner and Sluder, 2000). However, this notion has remained controversial and seems not supported by current evidence (Kostrouchova et al., 2001), suggesting that further effort will be needed to understand NHR-25 function in the molting cycle.

Elsewhere, we have reported a detailed characterization of a third screen hit, BLMP-1, which we found to modulate both molting timing and oscillatory gene expression (Hauser et al., 2021), possibly through its function as a pioneer transcription factor (Stec et al., 2021). Here, we have focused on characterizing another hit from our screen, GRH-1. Previously the least studied factor among our six leads, it is a member of the Grainyhead/LSF1 protein family that controls epithelial cell fates across animals (Sundararajan et al., 2020). Like NHR-23, it is repetively required for molting, but we additionally show that it acts during a specific window of each larval stage, before late intermolt.

This window is consistent with, and presumably determined by, the rhythmic accumulation of GRH-1 protein, whose levels peak shortly before molt entry. We show that the failure to complete development observed in the screen is due to a cuticle and ecdysis defect: the onset of ecdysis is delayed and the newly formed cuticle is prone to rupturing during ecdysis, particularly in the larval head region. Consistent with the view that generation of the new cuticle is defective, rupturing happens specifically after the onset of ecdysis, with tissue extrusion occurring after the old (outer) cuticle has already visibly detached from the worm body. Indeed, the notion of impaired cuticle biogenesis is also consistent with cuticle defects in GRH-1-depleted *C. elegans* embryos, which do not undergo ecdysis (Venkatesan et al., 2003), although it remains possible that inappropriate proteolytic activities during ecdysis might additionally damage the newly formed cuticle.

Soft, thin and granular cuticles, prone to rupturing, are also a hallmark of *Drosophila* Grainyhead mutant animals (Bray and Kafatos, 1991; Nusslein-Volhard et al., 1984). In mammals, *Grhl1* and *Grhl3* are dynamically expressed in the epidermis (Joost et al., 2020; Joost et al., 2016) and control epidermal differentiation (Yu et al., 2006), wound healing, eye lid closure (Boglev et al., 2011). Hence, our findings agree with a fundamental, evolutionarily conserved role of Grainyhead proteins in epidermis and ECM development and remodeling.

Such parallels extend to additional screen hits, and we are particularly intrigued by the apparent similarites between *C. elegans* molting and the mammalian hair follicle cycle. This process of rhythmic homeostatic skin regeneration is controlled by a poorly characterized "hair follicle clock", modulated by circadian rhythms (Paus and Foitzik, 2004; Plikus et al., 2015). Mutations in the mouse orthologues of *blmp-1* and *nhr-23*, *Prdm1* and *Rora*, respectively, impair the proper execution of hair follicle cycles (Magnusdottir et al., 2007; Steinmayr et al., 1998; Telerman et al., 2017), while *grhl1* ablation causes hair coat defects related to hair anchoring and potentially other processes (Wilanowski et al., 2008)

In summary, we propose that in this study we have identified and characterized a number of factors linked to a molting cycle oscillator that may have a functional correspondence in other animals, and that their further dissection provides a path to a molecular-mechanistic and systems understanding of gene expression oscillations in *C. elegans*. Additionally, we propose that *C. elegans* molting could serve as a powerful, genetically accessible, model of animal skin development and regeneration.

Methods

Transgenic animals

HW1939: *EG6699, xeSi296 [eft-3p::luc::gfp::unc-54 3'UTR, unc-119(+)] II* (Meeuse et al., 2020)

HW1949: *EG8080, xeSi301 [eft-3p::luc::gfp::unc-54 3'UTR, unc-119(+)] III* (this study)

rde-1(ne219) V (Tabara et al., 1999)

HW2150: *EG6699, xeSi296 [eft-3p::luc::gfp::unc-54 3'UTR, unc-119(+)] II; rde-1(ne219) V* (this study)

CA1200: *ieSi57 [eft-3p::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II* (Zhang et al., 2015)

HW1984: *ieSi57 [eft-3p::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] II; EG8080, xeSi301 [eft-3p::luc::gfp::unc-54 3'UTR, unc-119(+)] III* (this study)

HW2079: *EG8080, xeSi376 [eft-3p::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] III* (this study)

HW2418: *grh-1(xe135(grh-1::degron::3xFLAG)) I; EG8080, xeSi376 [eft-3p::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] III* (this study)

HW2434: *grh-1(xe135(grh-1::degron::3xFLAG)) I; EG6699, xeSi296 [eft-3p::luc::gfp::unc-54 3'UTR, unc-119(+)] II; EG8080, xeSi376 [eft-3p::TIR1::mRuby::unc-54 3'UTR, cb-unc-119(+)] III* (this study)

HW2493: *grh-1(xe146) I; hT2*

HW2603: *grh-1(syb616(grh-1::GFP::3xFLAG)) I* (PHX616, three times backcrossed) (this study; custom-made by SunyBiotech)

Generation of transgenic animals

Endogenous *aid::3xflag* tagging *grh-1* and generation of *grh-1(0)* mutants by CRISPR/Cas9-mediated editing was performed using the previously published *dpy-10(cn64)* co-conversion protocol (Arribere et al., 2014). For the sgRNA plasmid, we inserted the sgRNA sequence (5' agaggtttactctcatgagt 3') into NotI-digested pIK198 (Katic et al., 2015) by Gibson assembly (Gibson et al., 2009).

For degron-3XFLAG tagging, we used hybridized MM116 (5' AATTGCAAATCTAAATGTTTagaggtttactctcatgagtGTTTAAGAGCTATGCTGGAA 3') and MM117 (5' TTCCAGCATAGCTCTTAAACactcatgagagtaaacctctAAACATTTAGATTTGCAATT 3'). A degron-linker-3XFLAG-linker cassette was synthesized as gBlocks® Gene Fragments (Integrated DNA Technologies) with 50 bp

homology arms to the *grh-1* locus before the stopcodon: 5'
CCACGTTAATCGAGGTGGCTCCCACCAATCCAAACTCGTATTCCAACTCAATGCCTAAAGATCCAGCCAAACCTCCG
GCCAAGGCACAAGTTGTGGGATGGCCACCGGTGAGATCATACCGGAAGAACGTGATGGTTTCCTGCCAAAAATCA
AGCGGTGGCCCGGAGGCGGCGGCGTTCGTGAAGAGTACCTCAGGCGGCTCGGGTGGTACTGGCGGCAGCGACT
ACAAAGACCATGACGGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATGACAAGAGTACTAGCGGTG
GCAGTGGAGGTACCGGCGGAAGCTGAGAGTAAACCTCTTTAGGTTCTTGTCTTAATTCTCTTAAAGGAGGACT 3'.
Wildtype animals were injected with 10 ng/µL gBlock, 100 ng/µL sgRNA plasmid, 20 ng/µL AF-ZF-827
(Arribere et al., 2014), 50 ng/µL pIK155 (Katic et al., 2015) and 100 ng/µL pIK208 (Katic et al., 2015).
Genome editing was confirmed by sequencing.

For *grh-1(0)*, we used hybrized MM116 and MM117, and hybrized MM118 (5'
AATTGCAAATCTAAATGTTTGCGAAAATGCAAACATTAGAGTTTAAGAGCTATGCTGGAA 3') and MM119 (5'
TTCCAGCATAGCTCTTAAACTCTAATGTTTGCATTTTCGCAAACATTTAGATTTGCAATT 3') to delete the coding
region of *grh-1* and generate the *grh-1(xe146)* allele. Wildtype animals were injected with 100 ng/µL of
each sgRNA plasmids, 19.2 ng/µL AF-ZF-827 (Arribere et al., 2014), 47.9 ng/µL pIK155 (Katic et al., 2015)
and 95.8 ng/µL pIK208 (Katic et al., 2015). Genome editing was confirmed by sequencing. *grh-1(xe146)*
homozygous animals could not be maintained and the *grh-1(xe146)* allele was balanced with hT2.

Luciferase assays

Luciferase assays were performed as described before (Meeuse et al., 2020). In short, single embryos,
expressing luciferase from a constitutive and ubiquitous promoter (xeSi296 transgene) were placed in a
384-well plate (Berthold Technologies, 32505) by pipetting, and left to develop until adulthood in 90 uL S-
Basal medium containing *E. coli* OP50 (OD_{600} = 0.9) and 100 µM Firefly D-Luciferin (p.j.k., 102111).
Luminescence was measured using a luminometer (Berthold Technologies, Centro XS3 LB 960) every 10
minutes for 0.5 sec for 72 hours in a temperature controlled incubator set to 20 degrees.

For auxin experiments, a 400x stock solution of 3-indoleacetic acid (auxin, Sigma Aldrich, I2886) in 100%
ethanol was prepared. The stock solution was 400x diluted in the culture medium at the start each
experiment or at specific time points and in concentrations as indicated.

Luminescence data was analyzed using an automated algorithm to detect the hatch and the molts in
MATLAB with the possibility to annotate molts manually, as described before (Meeuse et al., 2020).
Completion of molts was scored by the presence of a drop in luminescence, followed by a period of stable
and low luminescence and subsequent rise in luminescence.

RNAi screen

To knock-down 92 'oscillating' TRANSCRIPTION FACTOR, we used the RNAi feeding method. *E. coli* HT115 bacteria carrying RNAi plasmids were obtained from either of the two libraries (Ahringer library (Fraser et al., 2000; Kamath et al., 2003), Vidal library (Rual et al., 2004)) or cloned (see generation of RNAi vectors).

Luciferase assays were performed as described above with the following adaptations: RNAi bacteria were grown in 5 mL auto-induction medium (2 mM MgSO4, 3.3 g/L (NH4)2SO4, 6.8 g/LKH2PO4, 7.1 g/L Na2HPO4, 5 g/L glycerol, 0.5 g/L glucose, 2 g/L α-lactose, 100 ug/mL Amp in ZY medium (10 g/L tryptone, 5 g/L yeast extract)) at 37 °C. Bacteria were diluted in S-Basal medium (OD600 = 0.45), with 100 uM Firefly D-luciferin (p.j.k., 102111) and 100 ug/mL Ampicillin.

We used HW1939 animals that express the xeSi296 transgene. As a control strain, we used HW2150 animals expressing xeSi296 in an *rde-1(ne219)* (Tabara et al., 1999) background, which are RNAi deficient. For each RNAi condition we used 2 adjacent columns in the 384-wells plate, i.e. 32 wells with 90 uL culture medium each. To avoid plate effects, the first 8 wells of the first column and the last 8 wells of the second column of the same RNAi condition were filled with an HW1939 animal and the remaining wells with an HW2150 animal.

To identify mutants, we inspected the heatmaps with trend-corrected luminescence (Olmedo et al., 2015) for aberrant duration or number of molts and intermolts.

Generation of RNAi vectors

For clones that were not available in the Ahringer or Vidal libraries, cDNA or genomic DNA was PCR amplified using the following primers:

locus	vector transformed	insert	Primer	Primer sequence
nhr-5	pMM012_R	Y73F8A.21a cDNA	MM070	ccaccggttccatggctagcTTCTGGCGGTAACAGTTCAA
			MM071	ttgatatcgaattcctgcagGATGTGAGTATGGAATATTCGG
dmd-8	pMM013_R	T22H9.4 cDNA	MM076	ccaccggttccatggctagcCCCTGTCATCTTCTTCAAATGC
			MM077	ttgatatcgaattcctgcagGTTTCAGCGCAGCTAATTGC
bcl-11	pMM014_R	F13H6.1a cDNA	MM078	ccaccggttccatggctagcAATAGAAACGTCTTCGGCGG
			MM079	ttgatatcgaattcctgcagTTAACGGTTGGTGTGACTGC
fkh-9	pMM015_R	K03C7.2b cDNA	MM080	ccaccggttccatggctagcGATTTGCTACGATCACCCAT
			MM081	ttgatatcgaattcctgcagGGCCTTGATTGGAGAAAGTG
ztf-16	pMM016_R	R08E3.4a cDNA	MM082	ccaccggttccatggctagcCGACTACTGTATTTTCCGAGTT
			MM083	ttgatatcgaattcctgcagCAGTTAACGAAAGTGATGACTC

sem-2	pMM017_R	C32E12.5.1 cDNA	MM084	ccaccggttccatggctagcGATCTCCAAAAACCGCCCAA
			MM085	ttgatatcgaattcctgcagTGCATCGCTCCATGGATAAT
grh-1	pMM018_R	Y48G8AR.1a cDNA	MM086	ccaccggttccatggctagcGAAGAAGTCCGACGGTGAAT
			MM087	ttgatatcgaattcctgcagGAGTTTGGATTGGTGGGAGC
dmd-9	pMM019_R	Y67D8A.3 genomic DNA	MM088	ccaccggttccatggctagcCTTTGTTCCAGTTCAAACCAC
			MM089	ttgatatcgaattcctgcagAGAGGGAAGGAACTGATAGAC
tbx-7	pMM020_R	ZK328.8.1 genomic DNA	MM090	ccaccggttccatggctagcCCTCATGACAGACAACTACT
			MM091	ttgatatcgaattcctgcagCAACAACTCCAAATCCACTT
M03D4.4b	pMM021_R	M03D4.4b cDNA	MM092	ccaccggttccatggctagcTCGGACACAGATTCATCACAAC
			MM093	ttgatatcgaattcctgcagTCCGGTGTTGCTGTATTTGTC
C08G9.2	pMM022_R	C08G9.2 cDNA	MM094	ccaccggttccatggctagcTACCGGCAAGTGTACCAAAT
			MM095	ttgatatcgaattcctgcagACCTTCACATGGATCTACACAA
nhr-112	pMM023_R	Y70C5C.6a cDNA	MM096	ccaccggttccatggctagcTTTTCCGCAGATTCTATCACTC
			MM097	ttgatatcgaattcctgcagTATGATTCATCTCGCACACCA
ets-4	pMM024_R	F22A3.1a cDNA	MM098	ccaccggttccatggctagcATGCAATCTTCCAATCCAACC
			MM099	ttgatatcgaattcctgcagAGGCAGGAATTTGTACACCA
ztf-14	pMM025_R	M163.2 genomic DNA	MM102	ccaccggttccatggctagcGCCGTCCCTGCATAACTACTC
			MM103	ttgatatcgaattcctgcagAGAGAAGTGAGTTGCGGGAG
ztf-29	pMM026_R	Y66D12A.12 cDNA	MM104	ccaccggttccatggctagcCGTCACCGGCTCAACTTCCA
			MM105	ttgatatcgaattcctgcagCATGTTCTCCTCCTTTCGCTCT

PCR fragments were cloned into the RNAi feeding Pml1 and Sma1 digested L4440 vector (L4440 was a gift from Andrew Fire (Addgene plasmid # 1654 ; http://n2t.net/addgene:1654 ; RRID:Addgene_1654)) using Gibson assembly (Gibson et al., 2009)) and transfected into *E. coli* HT115 bacteria.

Phenotype imaging

For imaging phenotypes, HW2418 worms were mounted on a 2% (w/v) agarose pad with a drop of M9 buffer (42 mM Na_2HPO_4, 22 mM KH_2PO_4, 86 mM NaCl, 1 mM $MgSO_4$). *grh-1::aid* animals were imaged on an Axio Imager Z1 (Zeiss) microscope. We acquired Differential Interference Contrast (DIC) images using a 100x/1.4 oil immersion objective and a TL Halogen Lamp (3.00 Volt, 900 ms exposure). Images (1388x1040 pixels, 142.1 µm x 106.48 µm pixel size, 12 Bit) were acquired every second from the moment that the cuticle became loose around the tip of the head until after the worm burst through the head, which took roughly 5 to 10 minutes. N2 animals were imaged on an Axio Imager Z2 (Zeiss) microscope. We acquired Differential Interference Contrast (DIC) images using a 63x/1.4 oil immersion objective) and a TL Vis-LED Lamp (5.74 Volt, 17 ms (**Figure 4A**) or 19 ms (**Figure 4B**) exposure). Images (1388x1040 pixels, 225.56 µm x 169.01 µm pixel size, 12 Bit) were acquired at 20 seconds ((**Figure 4A**) or 4 seconds (**Figure**

4B) intervals from the moment that the cuticle became loose around the tip of the head until the cuticle was shed.

Single worm imaging

To investigate the temporal expression pattern of endogenously tagged *grh-1::gfp*, strain HW2603 (*grh-1(syb616(grh-1::GFP::3xFLAG)) I*) was observed by time lapse imaging using single worm imaging as previously described (Meeuse et al., 2020) with slight modifications. Briefly, an array of microchambers (Bringmann, 2011; Turek et al., 2015) was made of 4.5% agarose in S-Basal medium (Stiernagle, 2005). OP50 bacteria were grown on agar plates, scraped off, and transferred to the chambers. Single eggs were placed on the chambers and flipped into a glass coverslip surrounded by a silicone insulator. Low melting agarose (3.5%) was used to seal the edges of the array, which was subsequently mounted on a glass slide for imaging. We imaged animals using a 2x sCMOS camera model (T2) on an Axio Imager M2 (upright microsocope) CSU_W1 Yokogawa spinning disk microscope with a 20x air objective (NA=0.8). The 488 nm laser was set to 70%, with 10 ms exposure and a binning of 2. The brightfield and fluorescent images were taken in parallel using a motorized z-drive with a 2-µm step size and 23 images per z-stack for a total duration of 60h at 10 min time intervals in a ~21°C room.

Single worm imaging data analysis

Brightfield images were segmented using a Convolutional Neural Network (CNN v2) (to be communicated elsewhere). Using a default threshold of 127 on the segmentation probability, GFP intensities were quantified on the segmented images using a previously published KNIME workflow (Meeuse et al., 2020). In short, worms are straightened and the GFP intensity of the worm is max projected to one pixel line for each time point. Background-subtracted mean GFP intensities are determined from 20%-80% of the anterior-posterior axis for each time point.To annotate molts, each image was visually inspected for molt entry and molt exit by scrolling through z-stack of the individual timepoints. The GFP intensities and lethargus data was plotted together in Python v3.9 using the Seaborn package.

Acknowledgements

We thank Iskra Katic and Lan Xu for help in generating transgenic strains, Anca Neagu for support in analyzing *grh-1* mutant phenotypes, Yannick Hauser, Marit van der Does, Markus Rempfler, Benjamin

Titze, Jan Eglinger and Laurent Gelman for help with imaging and image analysis, and Dimos Gaidatzis for help in analyzing luciferase screening data.

Funding

M.W.M.M. received support from a Boehringer Ingelheim Fonds PhD fellowship. S.N. from a Marie Sklodowska-Curie grant under the EU Horizon 2020 Research and Innovation Program (Grant agreement No. 842386). This work is part of a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 741269, to H.G.). The FMI is core-funded by the Novartis Research Foundation.

Author contributions

K.B. imaged the N2 molt. S.N. performed GRH-1 time-lapse imaging and related data analysis. M.W.M.M performed the remaining experiments and analyzed the associated data. H.G. and M.W.M.M conceived the project and wrote the manuscript.

References

Arribere, J.A., Bell, R.T., Fu, B.X., Artiles, K.L., Hartman, P.S., and Fire, A.Z. (2014). Efficient marker-free recovery of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans. Genetics *198*, 837-846.

Boglev, Y., Wilanowski, T., Caddy, J., Parekh, V., Auden, A., Darido, C., Hislop, N.R., Cangkrama, M., Ting, S.B., and Jane, S.M. (2011). The unique and cooperative roles of the Grainy head-like transcription factors in epidermal development reflect unexpected target gene specificity. Dev Biol *349*, 512-522.

Bray, S.J., and Kafatos, F.C. (1991). Developmental function of Elf-1: an essential transcription factor during embryogenesis in Drosophila. Genes Dev *5*, 1672-1683.

Bringmann, H. (2011). Agarose hydrogel microcompartments for imaging sleep- and wake-like behavior and nervous system development in Caenorhabditis elegans larvae. J Neurosci Methods *201*, 78-88.

Cohen, J.D., Sparacio, A.P., Belfi, A.C., Forman-Rubinsky, R., Hall, D.H., Maul-Newby, H., Frand, A.R., and Sundaram, M.V. (2020). A multi-layered and dynamic apical extracellular matrix shapes the vulva lumen in Caenorhabditis elegans. Elife *9*, e57874.

Cohen, J.D., and Sundaram, M.V. (2020). C. elegans Apical Extracellular Matrices Shape Epithelia. J Dev Biol *8*.

Frand, A.R., Russel, S., and Ruvkun, G. (2005). Functional genomic analysis of C. elegans molting. PLoS Biol *3*, e312.

Fraser, A.G., Kamath, R.S., Zipperlen, P., Martinez-Campos, M., Sohrmann, M., and Ahringer, J. (2000). Functional genomic analysis of C. elegans chromosome I by systematic RNA interference. Nature *408*, 325-330.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., and Smith, H.O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature Methods *6*, 343-345.

Gissendanner, C.R., Crossgrove, K., Kraus, K.A., Maina, C.V., and Sluder, A.E. (2004). Expression and function of conserved nuclear receptor genes in Caenorhabditis elegans. Dev Biol *266*, 399-416.

Gissendanner, C.R., and Sluder, A.E. (2000). nhr-25, the Caenorhabditis elegans ortholog of ftz-f1, is required for epidermal and somatic gonad development. Dev Biol *221*, 259-272.

Hauser, Y.P., Meeuse, M.W.M., Gaidatzis, D., and Großhans, H. (2021). The BLMP-1 transcription factor promotes oscillatory gene expression to achieve timely molting. bioRxiv, 2021.2007.2005.450828.

Hendriks, G.-J., Gaidatzis, D., Aeschimann, F., and Großhans, H. (2014). Extensive oscillatory gene expression during C. elegans larval development. Mol Cell *53*, 380-392.

Johnson, L.C., Aguilera, J., Levenson, M.T., Rechtsteiner, A., Vo, A.A., Ragle, J.M., and Ward, J.D. (2021). Conditional depletion reveals temporal requirements for the oscillating transcription factor NHR-23/NR1F1 in C. elegans larval progression. bioRxiv, 2021.2010.2027.465992.

Joost, S., Annusver, K., Jacob, T., Sun, X., Dalessandri, T., Sivan, U., Sequeira, I., Sandberg, R., and Kasper, M. (2020). The Molecular Anatomy of Mouse Skin during Hair Growth and Rest. Cell Stem Cell *26*, 441-457 e447.

Joost, S., Zeisel, A., Jacob, T., Sun, X., La Manno, G., Lonnerberg, P., Linnarsson, S., and Kasper, M. (2016). Single-Cell Transcriptomics Reveals that Differentiation and Spatial Signatures Shape Epidermal and Hair Follicle Heterogeneity. Cell Syst *3*, 221-237 e229.

Kamath, R.S., Fraser, A.G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le Bot, N., Moreno, S., Sohrmann, M.*, et al.* (2003). Systematic functional analysis of the Caenorhabditis elegans genome using RNAi. Nature *421*, 231-237.

Katic, I., Xu, L., and Ciosk, R. (2015). CRISPR/Cas9 Genome Editing in Caenorhabditis elegans: Evaluation of Templates for Homology-Mediated Repair and Knock-Ins by Homology-Independent DNA Repair. G3 (Bethesda) *5*, 1649-1656.

Kim, D.h., Grün, D., and van Oudenaarden, A. (2013). Dampening of expression oscillations by synchronous regulation of a microRNA and its target. Nat Genet *45*, 1337-1344.

Kostrouchova, M., Krause, M., Kostrouch, Z., and Rall, J.E. (1998). CHR3: a Caenorhabditis elegans orphan nuclear hormone receptor required for proper epidermal development and molting. Development *125*, 1617-1626.

Kostrouchova, M., Krause, M., Kostrouch, Z., and Rall, J.E. (2001). Nuclear hormone receptor CHR3 is a critical regulator of all four larval molts of the nematode Caenorhabditis elegans. Proc Natl Acad Sci U S A *98*, 7360-7365.

Kouns, N.A., Nakielna, J., Behensky, F., Krause, M.W., Kostrouch, Z., and Kostrouchova, M. (2011). NHR-23 dependent collagen and hedgehog-related genes required for molting. Biochem Biophys Res Commun *413*, 515-520.

Magnusdottir, E., Kalachikov, S., Mizukoshi, K., Savitsky, D., Ishida-Yamamoto, A., Panteleyev, A.A., and Calame, K. (2007). Epidermal terminal differentiation depends on B lymphocyte-induced maturation protein-1. Proc Natl Acad Sci U S A *104*, 14988-14993.

Meeuse, M.W.M., Hauser, Y.P., Hendriks, G.-J., Eglinger, J., Bogaarts, G., Tsiairis, C., and Großhans, H. (2020). Developmental function and state transitions of a gene expression oscillator in C. elegans. Mol Syst Biol *16*, e9498.

Monsalve, G.C., and Frand, A.R. (2012). Toward a unified model of developmental timing: A "molting" approach. Worm *1*, 221-230.

Nusslein-Volhard, C., Wieschaus, E., and Kluding, H. (1984). Mutations affecting the pattern of the larval cuticle inDrosophila melanogaster : I. Zygotic loci on the second chromosome. Wilehm Roux Arch Dev Biol *193*, 267-282.

Olmedo, M., Geibel, M., Artal-Sanz, M., and Merrow, M. (2015). A High-Throughput Method for the Analysis of Larval Developmental Phenotypes in Caenorhabditis elegans. Genetics *201*, 443-448.

Ou, Q., and King-Jones, K. (2013). What goes up must come down: transcription factors have their say in making ecdysone pulses. Curr Top Dev Biol *103*, 35-71.

Paus, R., and Foitzik, K. (2004). In search of the "hair cycle clock": a guided tour. Differentiation *72*, 489-511.

Plikus, M.V., Van Spyk, E.N., Pham, K., Geyfman, M., Kumar, V., Takahashi, J.S., and Andersen, B. (2015). The circadian clock in skin: implications for adult stem cells, tissue regeneration, cancer, aging, and immunity. J Biol Rhythms *30*, 163-182.

Rual, J.F., Ceron, J., Koreth, J., Hao, T., Nicot, A.S., Hirozane-Kishikawa, T., Vandenhaute, J., Orkin, S.H., Hill, D.E., van den Heuvel, S.*, et al.* (2004). Toward improving Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library. Genome Res *14*, 2162-2168.

Sandhu, A., Badal, D., Sheokand, R., Tyagi, S., and Singh, V. (2021). Specific collagens maintain the cuticle permeability barrier in Caenorhabditis elegans. Genetics *217*.

Singh, R.N., and Sulston, J.E. (1978). Some observations on molting in Caenorhabditis elegans. Nematologica *24*, 63-&.

Stec, N., Doerfel, K., Hills-Muckey, K., Ettorre, V.M., Ercan, S., Keil, W., and Hammell, C.M. (2021). An Epigenetic Priming Mechanism Mediated by Nutrient Sensing Regulates Transcriptional Output during C. elegans Development. Curr Biol *31*, 809-826 e806.

Steinmayr, M., Andre, E., Conquet, F., Rondi-Reig, L., Delhaye-Bouchaud, N., Auclair, N., Daniel, H., Crepel, F., Mariani, J., Sotelo, C.*, et al.* (1998). staggerer phenotype in retinoid-related orphan receptor alpha-deficient mice. Proc Natl Acad Sci U S A *95*, 3960-3965.

Stiernagle, T. (2005). Maintenance of C. elegans (February 11, 2006), WormBook, ed. The C. elegans Research Community, WormBook, doi/10.1895/wormbook.1.101.1, http://www.wormbook.org.

Sundararajan, V., Pang, Q.Y., Choolani, M., and Huang, R.Y. (2020). Spotlight on the Granules (Grainyhead-Like Proteins) - From an Evolutionary Conserved Controller of Epithelial Trait to Pioneering the Chromatin Landscape. Front Mol Biosci *7*, 213.

Tabara, H., Sarkissian, M., Kelly, W.G., Fleenor, J., Grishok, A., Timmons, L., Fire, A., and Mello, C.C. (1999). The rde-1 gene, RNA interference, and transposon silencing in C. elegans. Cell *99*, 123-132.

Telerman, S.B., Rognoni, E., Sequeira, I., Pisco, A.O., Lichtenberger, B.M., Culley, O.J., Viswanathan, P., Driskell, R.R., and Watt, F.M. (2017). Dermal Blimp1 Acts Downstream of Epidermal TGFbeta and Wnt/beta-Catenin to Regulate Hair Follicle Formation and Growth. J Invest Dermatol *137*, 2270-2281.

Tsiairis, C., and Großhans, H. (2021). Gene expression oscillations in C. elegans underlie a new developmental clock. Curr Top Dev Biol *144*, 19-43.

Turek, M., Besseling, J., and Bringmann, H. (2015). Agarose Microchambers for Long-term Calcium Imaging of Caenorhabditis elegans. J Vis Exp, e52742.

Turek, M., and Bringmann, H. (2014). Gene expression changes of Caenorhabditis elegans larvae during molting and sleep-like lethargus. PLoS One *9*, e113269.

Venkatesan, K., McManus, H.R., Mello, C.C., Smith, T.F., and Hansen, U. (2003). Functional conservation between members of an ancient duplicated transcription factor family, LSF/Grainyhead. Nucleic Acids Res *31*, 4304-4316.

Wilanowski, T., Caddy, J., Ting, S.B., Hislop, N.R., Cerruti, L., Auden, A., Zhao, L.L., Asquith, S., Ellis, S., Sinclair, R.*, et al.* (2008). Perturbed desmosomal cadherin expression in grainy head-like 1-null mice. EMBO J *27*, 886-897.

Zhang, L., Ward, J.D., Cheng, Z., and Dernburg, A.F. (2015). The auxin-inducible degradation (AID) system enables versatile conditional protein depletion in C. elegans. Development *142*, 4374-4384.

Supplementary figures

Figure S1: Related to Figure 1 – *grh-1* RNAi causes aberrant progression through development

Representative raw luminescence trace of single animals grown at 20°C for 72 hours. Left column showing wildtype animals and right column RNAi-deficient animals (*rde-1(ne219)*). Animals in upper panel were grown in the presence of *grh-1* RNAi and animals in the lower panel were grown in the presence of mock (empty vector) RNAi. Pre-hatch is indicated in red and molts in green.

Figure S2: Related to Figure 1 – Quantification of molt and intermolt duration in screen hits that complete development

A, Quantification of molt durations in RNAi deficient (*rde-1(ne219)*) animals (light grey) grown in the presence of mock (n=16), *bed-3* (n=14), *blmp-1* (n=16) or *nhr-25* (n=15) RNAi, and wildtype animals (dark grey) grown in the presence of mock (n=16)*, bed-3* (n=8)*, blmp-1* (n=16) or *nhr-25* (n=13) RNAi. Significant differences relative to *rde-1(ne219)* for each RNAi condition are indicated.

B, Quantification of intermolt durations in RNAi deficient (*rde-1(ne219)*) animals (light grey) grown in the presence of mock (n=16), *bed-3* (n=14), *blmp-1* (n=16) or *nhr-25* (n=15) RNAi, and wildtype animals (dark grey) grown in the presence of mock (n=16)*, bed-3* (n=8)*, blmp-1* (n=16) or *nhr-25* (n=13) RNAi. Significant differences relative to *rde-1(ne219)* for each RNAi condition are indicated.

* p<0.05, ** p<0.01, *** p<0.001, Wilcoxon test.

Figure S3: Related to Figure 3 – Intermolt is lengthened upon GRH-1 depletion at higher auxin concentrations

A, Quantification of the percentage of *grh-1(xe135); eft-3p::luc; eft-3p::TIR1* animals (HW2434) that enter each of four molts molt upon increasing concentrations of auxin. Note that for some concentrations M1 does not start at 100%, i.e. animals initiated I1 but M1 was not observed.

B, Boxplot showing the duration of the first intermolt of animals treated with indicated concentrations of auxin. Animals that failed to progress development beyond M1 (**A**). Significant differences relative to 0 nM auxin are indicated.

C, Boxplot showing the duration of M1 of animals treated with indicated concentrations of auxin. Animals that failed to progress development beyond M1 were excluded (**A**). Significant differences relative to 0 nM auxin are indicated.

P-values were determined by Wilcoxon test. ns: not significant ,* p<0.05, ** p<0.01, *** p<0.001

Figure S4: Related to Figure 3 – Low auxin concentrations strongly affect molt but not intermolt duration of *aid::grh-1* animals

A, Boxplot showing the duration of the first molt of *grh-1(xe135); eft-3p::luc; eft-3p::TIR1* animals (HW2434) treated with indicated concentrations of auxin. Animals that failed to progress development beyond M1 were excluded (**Figure S3A**). Significant differences relative to 0 nM auxin are indicated.

B, Boxplot showing the duration of the first intermolt of animals treated with indicated concentrations of auxin. Animals that failed to progress development beyond M1 were excluded (**Figure S3A**). Significant differences relative to 0 nM auxin are indicated.

C, Boxplot showing duration of first intermolt for vehicle-treated animals and animals treated with 267 nM and 400 nM auxin (**Figure S3A**). For auxin-treated animals, only those that showed a phenotype at M1 were selected. Significant differences relative to 0 nM auxin are indicated.

D, Boxplot showing the duration of I1, I2 and I3 of animals treated with indicated concentrations of auxin. Animals that revealed the phenotype at M1, M2 or M3 were excluded (**Figure S3A**). Significant differences relative to 0 nM auxin are indicated.

P-values were determined by Wilcoxon test. ns: not significant, * p<0.05, ** p<0.01, *** p<0.001

Figure S5: Related to Figure 3 – Effects of auxin in animal lacking a degron

A, Boxplots showing duration of each molt for indicated concentrations of auxin in *eft-3p::TIR1; eft-3p::luc* animals (HW1984), which lack the *aid* tag on *grh-1*. Differences relative to 0.25% ethanol (0 nM auxin) for 3.9 µM – 1mM auxin and relative to 1% ethanol (0 nM auxin) for 4 mM auxin are indicated (ns: not significant, * p<0.05, ** p<0.01, *** P<0.001, Wilcoxon test). Note that concentrations ≤ 250 µM yield at most minor extensions of molt durations (compare Figure 3). At 4 mM auxin, most animals fail to complete M4.

B, Boxplots showing duration of each intermolt for indicated concentrations of auxin in *eft-3p::TIR1; eft-3p::luc* animals (HW1984). Differences relative to 0.25% ethanol (0 nM auxin) for 3.9 µM-1mM auxin and relative to 1% ethanol (0 nM auxin) for 4 mM auxin are indicated (ns: not significant, * p<0.05, ** p<0.01, *** P<0.001, Wilcoxon test). Note that concentrations ≤ 250 µM yield at most minor extensions of intermolt durations.

Figure S6: Related to Figure 5 – GFP-GRH-1 becomes detectable in elongating embryos

Figure S7: Related to Figure 5 – GFP-GRH-1 accumulates in various tissues during larval development

Figure S9: Related to Figure 5 – GFP-GRH-1 accumulation is greatly reduced in adults

Figure S9: Related to Figure 6 – Molt exit phenotype is dependent on the time of GRH-1 depletion

A,B, Heatmap showing trend-correct luminescence (Lum) of *grh-1(xe135); eft-3p::luc; eft-3p::TIR1* animals (HW2434) treated with vehicle (0.25% ethanol, A) or 250 µM auxin (B) at 40 hours (white dashed line). Black intensities correspond to low luminescence occurring during lethargus (molt). Embryos of various stages were left to hatch and develop during the assay. Luminescence traces are sorted by entry into molt 3 (M3) such that traces of early hatched animals are at the bottom and those of late hatched animals are at the top.

C, Quantification of time at which animals enter molt 2 (M2 entry). Animals were assigned to M3 exit phenotype (blue) or M4 exit phenotype (red) when M3 or M4 was the last observed molt in the luminescence trace respectively. Animals sorted according to B.