Global ecological and biogeochemical impacts of pelagic tunicates

Jessica Y. Luo[1], Charles A. Stock[1], Natasha Henschke[2], John P. Dunne[1], Todd D. O'Brien[3]

[1]NOAA Geophysical Fluid Dynamics Laboratory, 201 Forrestal Rd, Princeton NJ 08540
[2]School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052
[3]NOAA NMFS Office of Science and Technology, Silver Spring MD, USA

Manuscript correspondence: Jessica.Luo@noaa.gov

Keywords: gelatinous zooplankton, pelagic tunicates, ocean biogeochemical cycles, carbon export, global model

16   **Abstract**
17
The pelagic tunicates, gelatinous zooplankton that include salps, doliolids, and appendicularians,
are filter feeding grazers thought to produce a significant amount of particulate organic carbon
(POC) detritus. However, traditional sampling methods (i.e., nets), have historically
underestimated their abundance, yielding an overall underappreciation of their global biomass
and contribution to ocean biogeochemical cycles relative to crustacean zooplankton. As climate
change is projected to decrease the average plankton size and POC export from traditional
plankton food webs, the ecological and biogeochemical role of pelagic tunicates may increase;
yet, pelagic tunicates were not resolved in the previous generation of global earth system climate
projections. Here we present a global ocean study using a coupled physical-biogeochemical
model to assess the impact of pelagic tunicates in the pelagic food web and biogeochemical
cycling. We added two tunicate groups, a large salp/doliolid and a small appendicularian to the
NOAA-GFDL Carbon, Ocean Biogeochemistry, and Lower Trophics version 2 (COBALTv2)
model, which was originally formulated to represent carbon flows to crustacean zooplankton.
The new GZ-COBALT simulation was able to simultaneously satisfy new pelagic tunicate
biomass constraints and existing ecosystem constraints, including crustacean zooplankton
observations. The model simulated a global tunicate biomass of 0.10 Pg C, annual production of
0.49 Pg C y^{-1} in the top 100 m, and export flux of 0.7 Pg C y^{-1}, representing 11% of the total
export flux past 100 m. Overall export from the euphotic zone remained largely constant, with
the GZ-COBALT pe-ratio only increasing 5.3% (from 0.112 to 0.118) compared to the
COBALTv2 control. While the bulk of the tunicate-mediated export production resulted from the
rerouting of phytoplankton- and mesozooplankton-mediated export, tunicates also shifted the
overall balance of the upper oceans away from recycling and towards export. Our results suggest
that pelagic tunicates play important trophic roles in both directly competing with
microzooplankton and indirectly shunting carbon export away from the microbial loop.

1. Introduction

In recent decades, there has been a growing recognition of the prevalence and ecological importance of gelatinous zooplankton (GZ), which include the cnidarian jellyfish, ctenophores, and pelagic tunicates (Hays et al., 2018; Henschke et al., 2016). While they have been a natural component of marine ecosystems extending back to the Cambrian (Hagadorn et al., 2002), their abundance and distributions have been largely overlooked among the public and non-GZ specialists (Condon et al., 2012). This may be due to a combination of the erroneous perception of GZ being "trophic dead-ends" within marine food webs (Hays et al., 2018; Lynam et al., 2006; Verity and Smetacek, 1996), as well as systematic biases in sampling leading to overall under-sampling of their biomass. Net-based sampling has been prevalent for over a century, and is very effective for sampling fish and hard-bodied, crustacean zooplankton (Wiebe and Benfield, 2003). Unfortunately, the fragile gelatinous zooplankton are often broken apart in nets, yielding a *ca.* 3 fold underestimation of their abundance and a *ca.* 10 fold (range: 5-15) underestimation of their carbon biomass relative to non-extractive, optical sampling (Remsen et al., 2004). The rise of in-situ plankton imaging systems have resulted in improved estimates of GZ abundance and distribution (e.g., Luo et al., 2014), yielding advances in understanding of their food-web interactions and biogeochemical impacts (Greer et al., 2021; Robison, 2005; Smith Jr et al., 2014). Combined with the increase in ecosystem-level studies of GZ (e.g., Stukel et al., 2021) and technological advances revealing the importance of GZ as food for higher trophic levels (Hays et al., 2018), there has been an overall paradigm shift in our understanding of the importance and role of gelatinous zooplankton within marine ecosystems.

Amongst zooplankton, GZ are notable for their high clearance rates and boom-and-bust population dynamics, which yield mass mortality events ("jelly-falls") that sink rapidly through the water column (Acuña et al., 2011; Billett et al., 2006; Lebrato et al., 2012; Lucas and Dawson, 2014). As a result, models have estimated, based on population densities and allometric scaling of ecological and physiological rates, a large contribution (e.g., 1.6-5.2 Pg C y^{-1} in Luo et al. 2020) of GZ-mediated carbon in the global biological pump, with their relative impact increasing with depth (Lebrato et al., 2019; Luo et al., 2020). However, these studies have been done independently of other biogeochemical and ecological constraints (i.e., "offline" calculations), which may yield unrealistic estimates of GZ contributions to marine ecosystems. Indeed, in a new model that includes cnidarian jellyfish as a plankton functional type (PlankTOM11), global export production exhibited very modest increases (+0.1 Pg C y^{-1}), suggesting that the online inclusion of a jellyfish class does not by itself substantially increase total export production (Wright et al., 2021). Of the three major groups of GZ considered by Luo et al. (2020), cnidarian jellyfish had the largest standing stock biomass but pelagic tunicates, despite their much lower biomass, over two times more sinking detritus than cnidarians and ctenophores combined.

These pelagic tunicates, small filter-feeders including appendicularians, salps, doliolids, and pyrosomes, are less conspicuous than the cnidarians and ctenophores, but are highly significant components of marine ecosystems due to their low trophic position, high clearance rates, and fast sinking fecal pellets (Andersen, 1998; Berline et al., 2011; Henschke et al., 2016; Hopcroft and Roff, 1998). Compared to crustacean mesozooplankton such as copepods which feed at predator to prey size ratios ranging from 5:1 to 100:1 (Hansen et al., 1994), pelagic tunicates can feed at predator to prey size ratios ranging from 10:1 to 10000:1 (Conley et al., 2018). Salps pump water through their fine mucous meshes that can filter submicron particles

such as bacteria and picoplankton; they are able to sustain the entirety of their energetic demands
by grazing on these size classes alone (Sutherland et al., 2010). Using both external and internal
filters to feed, appendicularians have the widest range of predator to prey size ratios (exceeding
2500:1 to lower than 10:1), and thus can feed on organisms ranging from 0.2-20 μm in size
(Conley et al., 2018; Deibel and Lee, 1992; Fernández et al., 2004). The offline Luo et al. (2020)
model estimated that, due to these feeding characteristics, these pelagic tunicates consume
between 3.8-8.3 Pg C y^{-1} in prey. Of this consumption, approximately 12-17% were later
consumed by higher trophic level predators, and 55-60% became detritus.
Global ocean biogeochemical and marine ecosystem models typically represent marine
food webs with roughly linear food chains of phytoplankton to zooplankton to (implicit) fish
(Kearney et al., 2021). The traditional inclusion of multiple zooplankton groups has been to
distinguish between zooplankton size, such as microzooplankton and mesozooplankton, with the
latter parameterized using crustacean zooplankton measurements (Aumont et al., 2015;
Buitenhuis et al., 2006; Stock and Dunne, 2010; Ward et al., 2012). Even in models with
complex food-webs, predator to prey size ratios beyond ~50:1 (c.f. Hansen et al., 1994) are
rarely considered. As such, current ocean biogeochemical models typically represent a marine
ecosystem in which crustaceans dominate zooplankton ecology. While this view represents
certain ecosystems well (Pershing and Stamieszkin, 2020), globally, there is a tension between
the traditional, crustacean-dominated zooplankton view of marine ecosystems and a shifting
paradigm that emphasizes the role of GZ. Unfortunately, GZ-focused offline models are unable
to reconcile this tension, as evidenced by high GZ-mediated global ingestion and production
rates (Luo et al., 2020) that may not support primary and secondary production rates for
crustacean zooplankton consistent with observations. Additionally, offline studies are limited in
the capacity to explore factors underlying observed GZ niches, and how GZ impacts emergent
food web patterns. These challenges are compounded by stubborn limitations in GZ observations
which are patchy and inconsistently sampled.
In this study, we added two explicit zooplankton functional types that represent
thaliaceans (salps, doliolids, pyrosomes) and appendicularians into Carbon, Ocean
Biogeochemistry, and Lower Trophics version 2 (COBALTv2; Stock et al., 2020), a global
model designed to represent a "traditional" marine ecosystem dominated by crustacean
zooplankton. We ask the following four questions:
1) Can simulations capture the magnitude and gradients of observed GZ biomass across
ocean biomes and along productivity gradients, after accounting for approximately an order of
magnitude under-sampling by nets?
2) Can simulations reconcile recent evidence for the importance of GZ with established
evidence for the prominence of crustacean zooplankton in biogeochemical cycles and the
plankton food web?
3) How does a simulation of GZ-modulated export that satisfies multiple food web
constraints compare with offline estimates?
4) What is the net impact of GZ zooplankton on the partitioning of carbon flows between
recycling, carbon export and energy flows to higher trophic levels?
129

2. Methods

As a brief overview of the methods, we begin with a description of plankton food web dynamics within the original COBALTv2 marine ecosystem model (Stock et al., 2014a, 2020), followed by the GZ additions that comprise GZ-COBALT: small and large pelagic tunicates. We first describe the baseline parameterization of GZ-COBALT, and then a few sensitivity experiments that explore parts of the parameter space and particular elements of the tunicate groups that make them distinct from crustacean zooplankton. Next, we detail the physical framework of the model. Finally, we describe the construction of a validation dataset for the two new GZ groups and the identification of an emergent relationship that contrasts gelatinous against crustacean zooplankton. The model is validated against multiple constraints, comprising new and established ecological and biogeochemical datasets.

2.1 COBALT Ecosystem Model

We use the COBALTv2 marine ecosystem model (Stock et al., 2020) as our baseline model configuration, with slight modifications. COBALTv2 is a 33-tracer, intermediate complexity model, representing biogeochemical cycles of carbon, alkalinity, oxygen, nitrogen, phosphorus, iron, silica, calcium carbonate, and lithogenic materials. The food web consists of three phytoplankton and three zooplankton functional types, as well as a free-living heterotrophic bacteria group. Two phytoplankton size classes (small and large) are represented, as well as diazotrophs, parameterized as a large *Trichodesmium* group. The small phytoplankton type includes cyanobacteria and other phytoplankton, up to 10 µm in equivalent spherical diameter (ESD), and the large phytoplankton type represents diatoms and other large phytoplankton from 10-100 µm in ESD. The different sized phytoplankton are parametrized to capture size-based contrasts in nutrient uptake, light harvesting, carbon to chlorophyll ratios, and susceptibility to microzooplankton grazing (Edwards et al., 2015, 2012; Hansen et al., 1994; Munk and Riley, 1952), such that the small phytoplankton are more successful in the low nutrient, seasonally stable subtropical gyres, and large phytoplankton are more competitive in the highly seasonal, high nutrient oceans (Stock et al., 2014a, 2020).

The base configuration of COBALTv2 contains three zooplankton size classes: a microzooplankton and two size classes of crustacean mesozooplankton. Microzooplankton (< 200 µm ESD) include ciliates and heterotrophic nanoflagellates, medium zooplankton (i.e, small mesozooplankton; 200-2000 µm ESD) represent small to medium-bodied copepods, and large zooplankton (2 – 20 mm ESD) represent large copepods and krill (Stock et al., 2014a). Predator-prey relationships are also largely based on size, with microzooplankton predating on bacteria and small phytoplankton, small mesozooplankton predating on diazotrophs, large phytoplankton, and microzooplankton, and large mesozooplankton predating on diazotrophs, large phytoplankton, and small mesozooplankton (Fig. 1a). Grazing is modeled as a Hollings Type II function with density-dependent switching (Stock et al., 2008), with maximum biomass specific grazing rates decreasing with increasing zooplankton size (Hansen et al., 1997). Grazing half-saturation constants do not vary between the zooplankton classes, and are tuned to reproduce observed patterns in phytoplankton biomass and turnover rates (Stock and Dunne, 2010).

Zooplankton grazing was assimilated at an assimilation efficiency (AE) of 0.7, with the non-assimilated grazing partitioned into dissolved and particulate (detritus) matter, depending on size. Detritus partitioning of non-assimilated matter were: 1/6 for microzooplankton, 2/3 for

medium, and 1 for large zooplankton, and the remainder separated between labile (70%), semi-
labile (20%) and semi-refractory (10%) dissolved matter. Assimilated matter was partitioned into
respiration (basal and active) and zooplankton production. Basal respiration is proportional to
biomass, whereas active respiration is proportional to ingestion rates. When respiration rates
exceed assimilated ingestion, production becomes negative and recontributes to zooplankton
mortality. The temperature dependence of biological rates (phytoplankton nutrient uptake and
growth, zooplankton grazing) is determined by a common Eppley (1972) curve with a Q_{10} factor
of 1.88, representing a near doubling of rates for every 10°C temperature increase.
184
185
186 **2.2 Gelatinous Zooplankton in COBALT (GZ-COBALT)**
187
Following Luo et al. (2020), we introduced two new pelagic tunicate groups into
COBALTv2 (Fig. 1b): small tunicates, which represents Appendicularians, and large tunicates,
which represents Thaliaceans (i.e., salps and doliolids). Appendicularians are small, free-
swimming organisms that produce gelatinous houses for filter-feeding, which are discarded when
clogged and re-created multiple times per day. Salps and doliolids are also filter-feeders, but
unlike appendicularians, are colonial (though salps and doliolids have solitary life stages), form
rapidly sinking fecal pellets (Perissinotto and Pakhomov, 1998), and exhibit mass die-offs (jelly-
falls; Henschke et al., 2013).
196

197
Figure 1. Food web structure of (a) COBALTv2 base model, and (b) GZ-COBALT model. Additional functional
groups in GZ-COBALT include a small and large tunicate. Bact = free living bacteria, SP = small phytoplankton,
Diazo = diazotrophs, LP = large phytoplankton, SZ = small zooplankton, MZ = medium zooplankton, LZ = large
zooplankton, and HP = higher trophic-level predators. Note that within typical zooplankton size categories, small
zooplankton would be considered microzooplankton, and medium and large zooplankton would be considered
mesozooplankton.
204
2.2.1 GZ food web structure
The pelagic tunicates were inserted into the COBALTv2 model as summarized in Fig. 1.
Pelagic tunicates have the widest prey-to-predator size ratios among zooplankton (Conley et al.,
2018), with appendicularians predating on organisms ranging from 0.04 – 20% of its size,

equivalent to 0.2 – 20 µm in length (Deibel and Lee, 1992; Fernández et al., 2004; Lombard et
al., 2011). Small tunicates in the baseline model setting are thus able to consume bacteria, small
phytoplankton, large phytoplankton, and microzooplankton. This is consistent with previous
efforts that included appendicularians in a NPZD-type model (Berline et al., 2011) wherein small
tunicates consumed small phytoplankton, which lie near the center of their prey kernel, with high
preference (100% prey availability), and all others with low preference (25% prey availability).
Appendicularians are important prey for many invertebrates and fish, often contributing large
proportions of the diets of medusae jellyfish and many larval fish species (Gorsky and Fenaux,
1998; Llopiz et al., 2010; Purcell et al., 2005). Thus, small tunicates were predated upon by
medium and large zooplankton and higher trophic level predators with high preference (Fig. 1b).
While they have relatively large body sizes (e.g., the solitary form of *Salpa thompsoni*
can exceed 10 cm; Dubischar et al., 2006), salps have fine mucous meshes that can filter
submicron particles, and have a preference for grazing on small algae such as picoplankton
(Sutherland et al., 2010). Salps are able to consume diatoms and bacteria, but with relatively low
efficiency (Dadon-Pilosof et al., 2019) and with many diatoms passing through salp guts
undigested (Harbison et al., 1986). Additionally, salps have been shown to be dominant grazers
on *Trichodesmium* in certain regions (Post, 2002). While, the prey of doliolids is less studied,
Walters et al. (2019) found using genetic approaches that doliolids preferentially consume
diatoms (particularly in their early life stages) and ciliates. Thus, in the present model, the large
tunicates were able to feed on small phytoplankton and diazotrophs with high preference (100%
prey availability), microzooplankton at medium preference (50% prey availability), and bacteria
and large phytoplankton at low preference (25% prey availability).
Large tunicate predators and parasites were traditionally thought to be primarily
sapphirinid copepods and large hyperiid amphipods, including the predatory and parasitic
Phronima spp., which consume salp tissue and live in their cleared-out barrels (Laval, 1980;
Madin and Harbison, 1977; Takahashi et al., 2013). Restricted emphasis on these relatively rare
crustacean predators and parasites has yielded the misconception of gelatinous zooplankton,
particularly salps, as trophic dead-ends. However, increasing evidence has highlighted the role of
thaliaceans as food for fish and other higher trophic levels. Over 200 species of fish, turtles,
corals, and echinoderms consume salps, doliolids, and pyrosomes, with many predators filling
their guts with thaliaceans during bloom periods (Harbison, 1998; Henschke et al., 2016;
Mianzan et al., 2001). Therefore, in our model, large tunicates are predated upon by medium and
large zooplankton at low preference (25% prey availability), reflecting the specialized nature of
the copepod and amphipod predators relative to the broader crustacean zooplankton population,
and by higher trophic level predators at medium preference (50% prey availability).
The diets of small and large tunicates are fairly similar in the model, as they are both
microphagous generalists. Notably, there was no size scaling in the tunicates' diets relative to
their body size (i.e., smaller tunicates did not consume smaller prey), which is in contrast to
crustacean mesozooplankton; large tunicates have larger predator-to-prey size ratios than small
tunicates (Conley et al., 2018). Rather, the key distinction (with respect to food web dynamics)
between the tunicates is the level of predation. Small tunicates have very strong levels of top-
down control, exerted by all mesozooplankton and higher trophic level predators. While large
tunicates experience predation by similar predators as small tunicates in the model, the strength
of that predation is reduced to account for their larger size. Other distinctions between the two
tunicates, including ingestion rates, metabolic scalings, and susceptibility to jelly-falls are
described in the next sections.

2.2.2 Ingestion and assimilation

As filter feeders, the prey consumption rates of pelagic tunicates are typically measured as biomass-specific clearance (or filtration) rates, yielding ingestion rates (I, mg C m^{-3} d^{-1}) as the product of the specific clearance rate (C_b, m^3 mg C^{-1} d^{-1}), prey biomass (P, mg C m^{-3}), and predator biomass (Z, mg C m^{-3})(Acuña et al., 2011):

$$I = C_b PZ \qquad \text{(eq. 1)}$$

In contrast, COBALTv2 and other NPZD-type models utilize saturating functional response (Hollings Type II in the case of one prey) for zooplankton grazing (Fasham et al., 1990; Gentleman et al., 2003; Stock and Dunne, 2010):

$$I = \frac{I_{max}PZ}{K_i + P} \qquad \text{(eq. 2)}$$

Where I_{max} is the maximum specific ingestion rate (d^{-1}), and K_i is the ingestion half-saturation constant (mg C m^{-3}). At $K_i \gg P$, eq. (2) reduces to I = (I_{max}/K_i)*P*Z, and C_b becomes equivalent to I_{max}/K_i (Acuña and Kiefer, 2000).

For small tunicate clearance rates, we used allometric relationships from Lombard et al. (2009), which measured physiological rates for a common appendicularian, *Oikopleura dioica*. Using a characteristic individual body size of 1 mm and associated biomass of 6.68 µg C, specific filtration rates at 0°C (converted using a Q10 of 1.88) range from 0.010-0.017 (mean: 0.013) m^3 mg C^{-1} d^{-1} (Lombard et al., 2009a). Considering that the clearance rates of both small and large tunicates do not significantly change with low to medium food concentrations (Gibson and Paffenhöfer, 2000; Paffenhöfer and Köster, 2011), we opted for a higher tunicate half-saturation constant (K_i = 250 mg C m^{-3}) than that of crustacean zooplankton (102 mg C m^{-3}; Table 1), consistent with the estimated range of K_i for *O. dioica* of 20-500 mg C m^{-3} (Acuña and Kiefer, 2000). The equivalent I_{max} at low prey concentrations would be 2.50-4.25 (mean: 3.25) d^{-1}. Following model tuning, we used an I_{max} value on the low end of the range (1.875 d^{-1}) given a mean K_i, to avoid overconsumption by small tunicates. However, considering the wide variation in K_i, these values were well within the observational bounds. The tradeoffs between tunicates and crustacean zooplankton were visualized in a plot of specific ingestion at 25°C (Fig. 2a): the small tunicate I_{max} and K_i results in a specific ingestion in between small/micro-zooplankton and medium/crustacean zooplankton. Additionally, a sensitivity run was conducted to illustrate the effect of the I_{max} tuning choice (Section 2.3).

For large tunicates, we were able to use an allometric scaling relationship from a prior effort (Luo et al., 2020) that compiled length, carbon biomass, and clearance rate relationships from various studies (see Madin and Deibel, 1998) into a single equation. Using a characteristic biomass of 1.5 mg C, which corresponds to a 20-38 mm length individual, specific clearance rates at 0°C range from 4.2E-4 to 7.4E-3 (mean: 1.8E-3) m^3 mg C^{-1} d^{-1}. Using a K_i of 250 mg C m^{-3}, and the same relationship between clearance rates, K_i, and I_{max} as above, we estimated large tunicate I_{max} values to be between 0.105-1.85 d^{-1}. In the model, we used a value in the lower half of the reported range (0.55 d^{-1}), tuned in conjunction with other variables with wide uncertainty bounds to match observed biomass concentrations. At the lowest prey concentrations, the I_{max}/K_i of large tunicates matched that of the large crustacean zooplankton. As prey concentrations increased, large tunicate ingestion fell between that of the medium and large crustacean zooplankton and reached its maximum ingestion rate much slower than either crustacean group (Fig. 2a).

Assimilation efficiency (AE) for zooplankton is typically set to be a fixed fraction of ingested material in models (between 0.6-0.8; Carlotti et al., 2000), and for crustacean

zooplankton in COBALTv2, it is set to 0.7, allowing for 70% of all food consumed to be
assimilated independent of prey concentration. However, for pelagic tunicates, in particular
appendicularians, there is evidence of AE declines as prey concentration increases, due to
development of tears of their pharyngeal filter and active prey rejection with increasing food
(Acuña and Kiefer, 2000; Lombard et al., 2009a, 2011). Retention and assimilation efficiencies
for salps and doliolids also vary widely, from 28-90%, which may be due to prey selectivity for
optimal sizes and preferred taxa (Andersen, 1986; Dadon-Pilosof et al., 2019; Pakhomov, 2004;
Pakhomov et al., 2006; Vargas and Madin, 2004; Walters et al., 2019). While we have
implemented prey selectivity in the feeding relationships, there is still evidence for feeding
apparatus clogging at high food concentrations due to the formation of a food bolus (Harbison et
al., 1986). Therefore, we implemented varying assimilation efficiencies for pelagic tunicates:

312
$$AE = AE_{max} - \left((AE_{max} - AE_{min}) \left(\frac{P}{K_{AE}+P} \right) \right) \qquad \text{(eq. 3)}$$

where AE_{max} and AE_{min} are maximum and minimum assimilation efficiencies, respectively, and
K_{AE} is the half-saturation constant for AE (all dimensionless).
This AE equation is a desaturating functional form (Hollings Type II subtracted from
AE_{max})(see also Berline et al., 2011). For small tunicates, AE_{max} and AE_{min} were 0.7 and 0.25,
respectively, and K_{AE} was 110 mg C m^{-3}, which is at the low end of the range for the
appendicularian *O. dioica* (145.52 +/- 33.36 std. err.) as measured by Lombard et al. (2009). For
large tunicates, to simulate the clogging response, we also used the same AE bounds as small
tunicates but with a K_{AE} of 215 mg C m^{-3}, a value at which approximately half of the *Pegea*
confoederata salps studied by Harbison et al. (1986) would form boluses. The difference in K_{AE}
between the tunicates results in the AE declining faster for small tunicates compared to their
larger counterparts (Fig. 2b).
Non-assimilated egestion losses for tunicates were parameterized similar to large
zooplankton with 100% of the detritus losses going towards sinking particulate organic matter
and no dissolved organic matter generated through grazing. This assumption is consistent with
the representation of small tunicates in Berline et al. (2011) and the observation that tunicate
detritus, representing the sinking houses of appendicularians and salp fecal pellets, are an
important source of zooplankton detritus (Alldredge and Silver, 1988), and contribute
significantly to carbon export fluxes (Alldredge, 2005; Andersen, 1998; Luo et al., 2020;
Robison, 2005).
In GZ-COBALT, we did not modify the sinking speed of GZ-mediated detritus, which is
pooled together with non-gelatinous detritus and sinks at a rate of 100 m d^{-1} (Alldredge and
Silver, 1988; Stock et al., 2014a; Turner, 2015). While studies have shown that GZ detritus can
sink at rates greatly exceeding that of marine aggregates and crustacean zooplankton fecal pellets
(Lebrato et al., 2013), we opted to focus this study on the impact of gelatinous zooplankton,
specifically tunicates, within the euphotic zone and leave the assessment of GZ-mediated export
on biogeochemical cycles at depth to future work.
339

Figure 2. Specific ingestion (a), assimilation efficiency (b), and specific production (c) as a function of generalized prey biomass at 25°C for zooplankton in GZ-COBALT. The three original COBALTv2 zooplankton types (small, medium, and large zooplankton; dashed lines) are unchanged in all GZ-COBALT simulations from the COBALTv2 control model.

2.2.3 Metabolism and respiration

In the COBALTv2 base model, the total zooplankton respiration rate is a sum of the basal and active respiration rate, with the former the resting metabolic rate that is proportional to biomass and the latter a fixed fraction of the ingestion rate (Flynn, 2005; Stock et al., 2014a). Pelagic tunicates are true filter feeders, and in contrast with crustacean zooplankton, are constantly in motion regardless of the food concentration, pumping water via tail oscillations (appendicularians), muscle contractions (salps), and ciliary action (doliolids and pyrosomes) (Deibel, 1998; Madin and Deibel, 1998). Therefore, a truly "basal" respiration rate only occurs when the tunicate is anesthetized. For *Salpa fusiformis*, swimming accounts for over half of total oxygen demand (Trueman et al., 1984), while for the appendicularian *O. dioica,* swimming accounted for roughly 34% of active respiration (Lombard et al., 2005). Furthermore, Lombard et al. (2005) found that there was no significant difference between the respiration levels of appendicularians at various food levels (no food, low, and high food), implying that the energy required to digest food was small compared to the energetic requirements of swimming. In contrast, the basal respiration rates for crustacean zooplankton are relatively low compared to respiration rates while feeding, which increase linearly with ingestion (Kiørboe et al., 1985). Thus, a key contrast that is built into GZ-COBALT is the difference between crustacean and gelatinous zooplankton respiration tradeoffs: crustaceans have relatively high active respiration rate (30% of ingestion) but low basal respiration, whereas GZ have low active respiration (15% of ingestion) but high basal respiration rates. Consequently, compared to crustacean zooplankton, tunicates incur a higher metabolic cost in low food concentrations, which prevents them from accumulating biomass in large portions of the subtropical gyres. These tradeoffs can be seen on a plot of specific production as a function of food concentration (Fig. 2c), and is most obvious when comparing small tunicates vs. small mesozooplankton, as the tunicate specific production remains negative for a greater portion of the low food concentrations. A full comparison of parameters is found in Table 1.

Biomass-specific basal respiration for small tunicates was calculated using relationships for *O. dioica* (Lombard et al., 2005). A laboratory study in the absence of food found that a 6.68 μg C individual at 0°C has a weight-specific oxygen consumption of 0.068 (0.055-0.085) μmol

O$_2$ (μmol C)$^{-1}$ d^{-1} (Lombard et al. 2005). Assuming a general zooplankton respiratory quotient of
0.87 (Mayzaud et al., 2005), this translates to a basal respiration of 0.047-0.074 d^{-1}. We opted to
use a value at the low end of the range (0.047 d^{-1}), so as to avoid complete elimination of their
biomass in subtropical environments, where they are found in low concentrations (Steinberg et
al., 2008).
For large tunicates, GZ-COBALT takes advantage of the mean allometric respiration
relationship compiled by Luo et al. (2020) from observations in Madin and Deibel (1998; and
references therein). With an average large tunicate of 1.5 mg C in that study respiring 2.5E-3
(8.0E-4 – 8.1E-3) mmol O$_2$ mg C^{-1} d^{-1}, and using a salp-specific respiratory quotient of 1.16
(Mayzaud et al., 2005), basal respiration varied by an order of magnitude (0.011-0.11 d^{-1}) with
mean of 0.035 d^{-1}. However, even with this large uncertainty range, Luo et al. (2020) found that
under average conditions in the pelagic oceans, even the lower bound of these respiration rates
were too high, such that metabolic demands exceeded available food resources, yielding
unfavorable conditions for survival, particularly in the subtropical gyres. Therefore, in the
baseline model configuration, we used a basal respiration rate (0.008 d^{-1}) that was slightly below
the lower bound, tuned to ensure realistic gelatinous zooplankton in oligotrophic ecosystems.
This choice is consistent with the strategy enlisted for calibrating the basal metabolic costs of the
crustacean zooplankton (Stock and Dunne, 2010) wherein highly uncertain basal metabolic rates
were calibrated to ensure that the simulated biomass was consistent with observations in the
oligotrophic gyres where the impact of basal metabolic costs are most prominent. In nature, To
explore the effect of this tuning, a sensitivity case was run where large tunicate basal respiration
was set to the mean value from allometry (Section 2.3).
398
2.2.4 Other sources of mortality
Another key difference between the small and large tunicates is in the additional sources
of mortality. While small tunicates primarily experience mortality through predation, large
tunicates experience cold temperature reproductive failures (Henschke and Pakhomov, 2019) as
well as mass mortality events (jelly-falls) (Henschke et al., 2015; Lebrato and Jones, 2009).
Large tunicate aggregation losses, representing jelly-falls, were parameterized as a quadratic loss
that is suppressed when food is plentiful, following the same functional form as the
phytoplankton aggregation losses as a function of nutrient limitation in COBALTv2 (Stock et al.,
2020; Waite et al., 1992) and other global biogeochemical models (e.g., PISCES; Aumont et al.,
2015). The aggregation losses are controlled by two parameters: f_{agg}, which represents the
fraction of the maximum ingestion rate above which aggregation losses are suppressed, and m_{agg},
or a maximum aggregation loss rate. f_{agg} was set to 0.1 to account for the salp ability to tolerate
low food concentrations, and m_{agg} was set to 1.0E-3 m^3 mg C^{-1} d^{-1} to achieve jelly-falls
representing approximately 35% of total large tunicate mortality following Luo et al. (2020).
413
414
415 **Table 1.** GZ-COBALT zooplankton ingestion, respiration, and aggregation parameters. Plankton functional types
416 (PFT)s: smt = small tunicates, lgt = large tunicates, smz = small zooplankton, mdz = medium zooplankton, and lgz =
417 large zooplankton.

Parameter	Name (Units)	PFT	Values	References
I_{max}	Maximum ingestion rate (day^{-1})	smt	1.875	(Lombard et al., 2009a)
		lgt	0.55	(Luo et al., 2020; Madin and Deibel, 1998)
		smz	1.42	

		mdz	0.57	(Hansen et al., 1997; Stock et al., 2020; Stock and Dunne, 2010)
		lgz	0.23	
K_i	Ingestion half-saturation constant (mg C m^{-3})	smt, lgt	250	(Acuña and Kiefer, 2000; Gibson and Paffenhöfer, 2000)
		smz, mdz, lgz	102	(Stock et al., 2014a, 2020)
$bresp$	Basal respiration rate (day^{-1})	smt	0.06	(Lombard et al., 2005)
		lgt	0.008	(Luo et al., 2020; Madin and Deibel, 1998); tuned to allow for sufficient biomass in subtropical gyres
		smz	0.018	(Hansen et al., 1997; Stock et al., 2020; Stock and Dunne, 2010)
		mdz	0.008	
		lgz	0.0032	
f_{aresp}	Fraction of ingestion rate for active respiration (dimensionless)	smt, lgt	0.15	See text; also tuned to allow for sufficient biomass in subtropical gyres
		smz, mdz, lgz	0.3	(Stock et al., 2014a, 2020)
AE_{min}, AE_{max}	Minimum and maximum assimilation efficiency (dimensionless)	smt, lgt	0.25, 0.8	(Lombard et al., 2009a; Madin and Deibel, 1998; Pakhomov et al., 2006)
		smz, mdz, lgz	0.7, 0.7	(Stock et al., 2014a, 2020)
K_{AE}	Assimilation efficiency half-saturation constant (mg C m^{-3})	smt	110	(Berline et al., 2011; Lombard et al., 2009a)
		lgt	215	(Harbison et al., 1986)
		smz, mdz, lgz	N/A	N/A
f_{agg}	Fraction of maximum ingestion at which aggregation mortality is suppressed (dimensionless)	lgt	0.1	See text
m_{agg}	Maximum aggregation loss rate (m^3 mg C^{-1} d^{-1})	lgt	1.0E-3	Calibrated to achieve jelly-falls representing 35% of total mortality

418
419
420 **2.4 Physical Framework**
421
The GZ-COBALT model with 35 tracers was run in a global ocean-ice configuration
using the GFDL models Modular Ocean Model 6 (MOM6) and Sea Ice Simulator 2 (SIS2) in a
nominal 0.5°, or roughly 50 km, horizontal resolution (OM4p5, Adcroft et al., 2019). The 0.5°
horizontal grid improves the resolution of boundary currents compared to earlier generations of
1° MOM models. The vertical coordinate in MOM6 is a hybrid z*-isopycnal vertical coordinate
system implemented using an Arbitrary Lagrangian-Eulerian (ALE) method, such that isopycnal
coordinates are used in the ocean interior and a z* coordinate is used in the mixed layer. OM4p5
uses 75 vertical layers, which allows for finer resolution at the ocean surface (~2 m) compared to
earlier model configurations with 10 m surface resolution and 50 vertical layers (Adcroft et al.,
2019). The ocean and ice model configurations are also equivalent to those components used
within the fully-coupled ESM4.1 model (Dunne et al., 2020).

Model simulations were forced using Common Ocean-Ice Reference Experiment II
(CORE-II) (Large and Yeager, 2009), a 60-year interannually varying dataset representing
atmospheric forcings from 1948-2007. The model was initialized similar to that of the fully-
coupled model (Stock et al., 2020): from World Ocean Atlas 2013 (WOA13) data for
temperature, salinity, oxygen, and dissolved inorganic nutrients (Garcia et al., 2013a, 2013b;
Locarnini et al., 2013; Zweng et al., 2013), and from Global Ocean Data Analysis Project
(GLODAPv2) for dissolved inorganic carbon and alkalinity (Lauvset et al., 2016). Other tracers
were initialized from outputs of a previous version of COBALT (Stock et al., 2014a), and the
two new gelatinous zooplankton tracers were initialized with biomass concentrations similar to
medium and large zooplankton. Additional sources of nutrients include atmospheric deposition
of NH_4 and NO_3 (Horowitz et al., 2003), dust from Zhao et al. (2018) with soluble Fe calculated
in accordance with Baker and Croot (2010), as well as coastal iron and river nutrients from the
GlobalNEWS dataset (Seitzinger et al., 2005), as described in (Stock et al., 2020)
GZ-COBALT was run for one 60-year interannual forcing cycle. Results are reported
from a climatology computed from the last 20 years of model simulation, representing 1988-
2007. A COBALTv2 control simulation with the same exact model setup, but with tunicates
turned off, was also run for 60 years.
450
451 **2.5 Parameter Sensitivity Runs**
452
To understand the impact of the unique aspects of GZ physiology and ecology as
described above on their emergent distribution and productivity, we considered a number of
perturbations around the baseline settings described above. These sensitivity runs (Table 2)
examine both the effect of our tuning choices (cases 1-2) as well as some key tradeoffs relative
to crustacean zooplankton (cases 3-5). All parameter sensitivity runs were conducted following
the same physical forcing as the GZ-COBALT base simulation.
459
1) For large tunicates, the basal respiration rate was adjusted to be slightly below the
lower bound from the literature. Sensitivity case 1 examines the effects of using the
mean basal respiration rate from allometric relationships from the literature.
2) For the baseline GZ-COBALT model, we used a maximum ingestion rate of small
tunicates at the lower bound of the range. For sensitivity case 2, we test a case where
the small tunicates' I_{max} is higher, set to the mean of the literature-based range.
3) Sensitivity case 3 explores the impact of the unique feeding behavior of GZ relative
to crustaceans. For the calibrated model (base case), we used a mean K_i value from a
wide range measured by Acuña and Kiefer (2000). However, models are known to be
quite sensitive to this relatively unconstrained parameter (Stock and Dunne, 2010).
Thus, we ran a sensitivity test where the tunicate K_i values were the same as that of
the crustacean zooplankton. Since the maximum ingestion rate (I_{max}) was set in
combination with the K_i values to achieve specific filtration rates at low prey
concentrations consistent with observations, I_{max} was also modified accordingly to
preserve the observational constraint.
4) Sensitivity case 4 examines another unique aspect of pelagic tunicates feeding
relative to that of crustaceans. In the baseline GZ-COBALT, we implemented varying
assimilation efficiency (AE) following Berline et al. (2011). To explore the
uncertainty associated with this assumption, we ran a sensitivity test where the

tunicates' AE were constant, but set at a slightly lower value than other zooplankton,
to account for their comparatively lower retention rates.
5) Finally, a case was run to explore the impacts of ignoring the role of large tunicate
aggregation mortality (representing jelly-falls).
483
484 To illustrate the impact of these various parameter modifications, we computed the
485 specific production of all zooplankton groups under an idealized condition of 25°C with a
486 generalized prey biomass (Fig 3), which can be contrasted against the GZ-COBALT baseline
487 (Fig. 2c).
488
489 **Table 2.** Parameters tested in sensitivity tests, showing the values in the base code as well as the permutation for the
490 individual sensitivity case. For detailed names of parameters, see Table 1. Plankton functional types (PFTs): smt =
491 small tunicates; lgt = large tunicates.

Case #	Parameter	PFT	Base code	Permutation
1	$bresp$ (d^{-1})	lgt	0.008	0.035
2	I_{max} (d^{-1})	smt	1.875	3.25
3	I_{max} (d^{-1}), K_i (mg C m^{-3})	smt	1.875, 250	0.765, 102
		lgt	0.55, 250	0.22, 102
4	AE_{min}, AE_{max}	smt & lgt	0.25, 0.8	0.6, 0.6
5	m_{agg} (m^3 mg C^{-1} d^{-1})	lgt	1.0E-3	0

492
493

494
495 **Figure 3.** Specific production as a function of generalized prey biomass at 25°C for sensitivity cases 1-4 (a-d). The
496 three original COBALTv2 zooplankton types (small, medium, and large zooplankton; dashed lines) are unchanged

497   in all GZ-COBALT simulations from the COBALTv2 control. The specific production in sensitivity case 5 is the
498   same as the base case (Fig. 2c).
499
500
501   **2.6 Validation data**
502
As described in the introduction, a central question in our analysis is whether the model can
simultaneously reconcile recent measurements suggesting that GZ carbon biomass is *ca.* 10x
greater than previously thought while also satisfying core observational constraints on crustacean
biomass, nutrients, chlorophyll and net primary production (NPP). The subsections that follow
describe the datasets compiled and/or enlisted for each of these tasks, and the analyses used to
assess model-data consistency across trophic gradients and ocean biomes.
509
2.6.1 Pelagic tunicate dataset and model-data comparison
To generate biomass validation data for pelagic tunicates, we updated the gridded
tunicate data (primarily salps) from Luo et al. (2020) with additional data on both thaliaceans and
appendicularians from the NOAA Coastal and Oceanic Plankton Ecology, Production, and
Observation Database (COPEPOD) (O'Brien, 2014). From the COPEPOD database, raw data on
urochordates were extracted and divided into small tunicates (all appendicularians) and large
tunicates (salps, doliolids, and pyrosomes). With the exception of data from the ECOSAR-II
cruise (Muxagata, 1999), all other data were in numeric density only (# individuals m^{-3}).
Numeric density data were first converted to a common 330 µm mesh size (Moriarty and
O'Brien, 2013; O'Brien, 2005). Second, since the geometric mean cannot handle zeros, zero
numeric density values were modified to be a non-zero value slightly below the minimum value
for both size fractions (small: 0.0008 ind. m^{-3}, large: 0.001 ind. m^{-3}). Next, numeric density was
converted to carbon biomass using the characteristic individual biomass values defined in section
2.2.3 (appendicularians: 6.7 µg C, salps: 1.5 mg C). Characteristic biomass values for pyrosomes
and doliolids were 22.9 mg C ind^{-1} (100 mm individual) and 19.2 µg C ind^{-1} (5 mm individual),
respectively, following Lucas et al. (2014), which used regression conversions from Gibson and
Paffenhöfer (2000) and Mayzaud et al. (2007). Using the geometric mean, 1° gridded values
were averaged by month, then year, for an annual mean biomass. Finally, the Jellyfish Database
Initiative (JeDI; Condon et al., 2015) database was additionally queried for appendicularian data.
Data from 90 additional 1° grid cells, primarily from the North Atlantic and Eastern Equatorial
Pacific, were present in the JeDI dataset but not in the COPEPOD database. These data were
added to our validation dataset. Appendicularian data were present in a total of 3,914 1° grid
cells (Fig. 4a).
Thaliacean data from the COPEPOD database were combined with the Luo et al. (2020)
gridded salp data, which primarily included gridded biomass data from Lucas et al. (2014), with
updates from JeDI the Palmer LTER site at the Western Antarctic Peninsula (Steinberg et al.,
2015), and KRILLBASE (Atkinson et al., 2017). Out of the 5,468 grid cells with data, there were
1,481 cells where COPEPOD data were only present, 1,952 cells where the Luo et al. (2020) data
were only present, and overlap at 2,035 grid cells (Fig. 4b). Because the raw data with assumed
lengths and carbon conversions from Lucas et al. (2014) were not available, we were unable to
examine individual data points for overlap and cross-validation. However, a broad examination
of the two datasets revealed that at the areas of overlap, carbon biomass compiled from
COPEPOD was 1.4x (geometric mean) that of Luo et al. (2020), with variations likely due to the
finer taxonomic detail of the Lucas et al. (2014) effort. This was within the uncertainty bounds

that we considered acceptable (roughly 2x uncertainty). Ultimately, due to discrepancies in
classification and specificity over time (e.g., broad categories such as "Tunicata" and "Salps and
doliolids" were dominant in classifications from the 1950's and 1960's, but not later), we
decided that using a single characteristic carbon biomass conversion for each broad taxonomic
category in the COPEPOD data gave a taxonomic specificity consistent with the coarsest
taxonomic specificity in the data. Further, this single biomass conversion removed a persistent
discontinuity in the Luo et al. (2020) carbon biomass values in the Indian Ocean south of 5°N
that we were previously unable to resolve. Since the vast majority of the JeDI data sources in the
Indian Ocean were from the 1959-1965 International Indian Ocean Expedition (IIOE), which are
also present in COPEPOD (Condon et al., 2015), we opted to replace the Luo et al. (2020) Indian
Ocean data with the COPEPOD data during our merge. For the rest of the oceans, we merged the
two datasets by taking the geometric mean at every grid cell where there was overlap.
Ultimately, the discrepancies between the datasets were quite low compared to the
differences in biomass due to sampling type, particularly when comparing extractive (nets) vs.
non-extractive (imaging) methods. Since traditional, net-based sampling systems break apart
fragile organisms such as pelagic tunicates and other gelatinous zooplankton, a biomass
adjustment is necessary to account for the reduced sampling from nets. Remsen et al. (2004)
used concurrent sampling with an imaging system and a 162 μm mesh plankton net, and found
that for pelagic tunicates, their abundance was undersampled by nets by a factor of 3-4x, and
their carbon biomass undersampled by a factor of 5-15x. Therefore, we considered an additional
"adjusted biomass" from samples with a 10x increase relative to the unadjusted biomass. We
focus mainly on this adjusted biomass, which is indicative of nascent appreciation of the likely
broader importance of gelatinous zooplankton as revealed by optical instruments. This is also
consistent with our intent to assess whether these high values can be reconciled with the overall
high abundance of mesozooplankton in many regions.
We complement our assessment of the simulated magnitude of tunicate biomass with one
of the relationship between tunicate biomass and other ecosystem properties spanning
oceanographic gradients. As the spatial gradients in tunicate biomass span 5 orders of magnitude,
this assessment provides a second metric less sensitive to the adjustments above. To do this, we
considered the GZ biomass as a function of chlorophyll concentration. The resultant, large-scale
relationship allowed for contrasts between large and small tunicates, and between tunicates and
crustaceans. For chlorophyll, we used the GlobColour merged satellite chlorophyll product (from
MERIS, MODIS-Aqua, and SeaWiFS) monthly climatology for case 1 waters using the weighted
averaging method, blended at latitudes south of 50°S with the Southern Ocean algorithm of
Johnson et al. (2013). We computed a growing season mean, define as all months for latitudes
between 30°N and 30°S, and spring and summer only for latitudes poleward of 30°N/S. The
slope of the log-log relationships between chlorophyll and the biomass of small tunicates, large
tunicates, and crustacean mesozooplankton (Moriarty and O'Brien, 2013; more details in Section
2.6.3) were established as emergent relationships for validation purposes.
583
2.6.2 Biome definitions
Finally, we assess gelatinous zooplankton simulation as a function of ocean biome,
adjusting for any systematic biases in the model by referencing biome locations to chlorophyll,
light, and temperature thresholds. We used the three major ocean biomes of Stock et al. (2014),
following Banse (1992): 1) Low Chlorophyll (LC), which encompasses the subtropical gyres, 2)
High Chlorophyll Seasonally Stratified (HCSS), which encompasses the high latitudes, and 3)

High Chlorophyll Permanently Stratified (HCPS), which includes the coastal and equatorial
upwelling regions. Stock et al. (2014) used a threshold of 0.125 mg Chl m^{-3} to separate between
the low vs. high chlorophyll regions in observational chlorophyll datasets. In our biome
definition, we first calculated the total ocean area with observational chlorophyll values lower
than that threshold (approximately 40% of the world's oceans), then found the model chlorophyll
threshold that resulted in a model LC area that most closely matched the LC surface area from
observations. For the COBALTv2 control and GZ-COBALT, this threshold was 0.162 and 0.184
mg Chl m^{-3}, respectively. Next, to distinguish between the seasonally vs. permanently stratified
regions, we used the minimum of the mixed layer irradiance climatology (light averaged over the
mixed layer). HCSS regions were demarcated as those with minimum mixed layer irradiances
lower than 5 W m^{-2}, while the opposite was true of HCPS. Using mixed layer irradiance more
accurately defined the seasonal seas vs. upwelling areas, preventing HCPS areas from occurring
in Arctic regions with shallow maximum mixed layers (Stock et al., 2014a). Biomes for both
GZ-COBALT and the COBALTv2 control are shown in Fig. S1.
604
2.6.3: Crustacean zooplankton dataset and model-data comparison
To assess whether pelagic tunicate biomass magnitude and cross-biome gradients can be
represented while maintaining crustacean zooplankton populations consistent with observations,
we used the 2012 gridded carbon biomass data compilation from the COPEPOD database
(Moriarty and O'Brien, 2013). The entire COPEPOD database (O'Brien, 2014) consists of
multiple types of data products, including the raw, taxonomic data as used above for pelagic
tunicates, as well as the Moriarty and O'Brien (2013) carbon biomass compilation, which is the
more commonly used dataset for mesozooplankton model validation. In total, the COPEPOD
global carbon biomass compilation includes over 150,000 data points that were converted to an
equivalent 333 μm mesh net size, with each gridded value representing multiple data points.
Given the net-based sampling and the historical focus on crustacean zooplankton, the vast
majority of the individual data points consisted of hard-bodied mesozooplankton. Thus, we used
the COPEPOD global carbon biomass compilation as a proxy of crustacean mesozooplankton to
compare against the small and large crustacean mesozooplankton in GZ-COBALT.
Similar to the GZ data, the crustacean observations were also scaled with chlorophyll on
a log-log scale. This enabled us to make comparisons along trophic gradients and across biomes
for crustaceans and gelatinous zooplankton.
Finally, in addition to GZ and crustacean constraints, we include a suite of standard
biogeochemical metrics to ensure that the model solution satisfies large-scale productivity and
nutrient patterns. The data we used were the dissolved inorganic nutrient concentrations (NO$_3$,
PO$_4$, and SiO$_3$) at the ocean surface from the World Ocean Atlas (WOA) 2018 (Garcia et al.,
2019).
627
628

3. Results

3.1 Global distribution and biomass-chlorophyll scaling

The GZ-COBALT simulation produced values consistent with the adjusted biomass of small and large pelagic tunicates, while also reproducing observed crustacean biomass and satisfying ocean biogeochemical constraints (Figs. 4-5, Tables 3-4). Global NPP was 53.7 Pg C y^{-1} and export flux at 100 m was 6.36 Pg C y^{-1} in GZ-COBALT, compared to 55.4 and 6.23 Pg C y^{-1} in COBALTv2. Surface chlorophyll and macronutrient concentrations in GZ-COBALT also compared well with the COBALTv2 control and observational constraints (Fig. 5).

The modeled global mean annual biomass integrated over the top 100 m was 5.8 Tg C for small tunicates and 81.5 Tg C for large tunicates, yielding a total 100 m biomass of 87 Tg C. A small but non-negligible fraction of tunicate biomass was below 100 m, even with the model lacking vertical migration, such that the water column integrated biomass was 102 Tg C. These values are within the adjusted mean and uncertainty of the observations. In comparison, the medium/large crustacean mesozooplankton biomass (representing the size fraction most closely comparable to the values in COPEPOD database) in GZ-COBALT was 205 Tg C in the top 100 m, which was slightly lower than the COBALTv2 value of 220 Tg C. Observational estimates of large mesozooplankton biomass from COPEPOD, using a biome-specific geometric mean and standard deviation to extrapolate globally, was 133 (+/- 209) Tg C over the top 200 m. See Table 3 for additional comparisons by major ocean biome.

Figure 4. Global mean distributions of small and large tunicates and crustacean mesozooplankton, comparing tunicate adjusted observations (a,b), large crustacean biomass from COPEPOD (c) with results from the top 100 m of the model (d,e,f). Medium/large crustacean mesozooplankton model values are given as the large mesozooplankton plus 0.5*small mesozooplankton. Model data show the time-average of the growing season only (fall and winter months excluded poleward of 30°N/S).

Table 3. Global and biome-specific biomass comparison of the observations and the GZ-COBALT model. Model results are the annual area weighted mean of the top 100 m carbon biomass. Observational values are given as the geometric mean and geometric standard deviation of values within the biomes, with global geometric standard deviation calculated following the procedure in Luo et al. (2020). Biomes: LC = Low Chlorophyll; HCPS = High Chlorophyll Permanently Stratified; HCSS = High Chlorophyll Seasonally Stratified. See Fig. S1 for biome maps.

Biome	Large Tunicates (mg C m^{-3})		Small Tunicates (mg C m^{-3})		Med/Large Crustacean mesozooplankton (mg C m^{-3})	
	Adj. Obs. mean (stdev)	Model annual mean	Adj. Obs. mean (stdev)	Model annual mean	Obs. mean (stdev)	Model annual mean
LC	1.4 (10)	1.4	0.05 (16)	0.06	0.98 (2.3)	2.6
HCPS	1.9 (8.0)	3.2	0.11 (15)	0.16	3.6 (2.5)	6.6
HCSS	2.9 (11)	2.4	0.08 (23)	0.28	2.9 (3.2)	8.3
Global	2.0 (9.9)	1.6	0.07 (18)	0.11	2.0 (3.1)	4.1

Figure 5. Surface chlorophyll, log$_{10}$ scale (mg Chl m^{-3}, a-c) and nutrients (mmol NO$_3$ m^{-3}, d-f; mmol PO$_4$ m^{-3}, g-i; mmol SiO$_3$ m^{-3}, j-i) from GZ-COBALT (left column; a, d, g, j), the COBALTv2 Control (center column; b, e, h, k), and observations (right column; c, f, i, l). Model bias, root mean squared error (RMSE), and Pearson's correlation coefficient (r) are also reported.

In the observations, we found that there was a contrast in the slope and intercept of biomass-chlorophyll scaling relationship between small tunicates, large tunicates, and crustacean mesozooplankton (Fig. 6). The small tunicates had significantly less biomass and a steeper log-

log slope (0.63 +/- 0.045 residual std. err.; Fig. 6a) than the large tunicates, which was much
flatter (0.22 +/- 0.036; Fig. 6b). The crustacean mesozooplankton data had much less variability,
a mean biomass similar to that of the large tunicates, and a biomass-chlorophyll scaling slope a
little shallower than the small tunicates (slope: 0.57 +/- 0.009; Fig. 6c). GZ-COBALT
successfully captured the differences in mean biomass across all three groups, as well as the
contrast in slope between the three groups, though admittedly the modeled slopes were all
slightly steeper than the observational slopes. The large tunicates had the shallowest biomass-
chlorophyll scaling slope (0.43, Fig. 6b), followed by the crustacean mesozooplankton (0.71,
Fig. 6c) and the small tunicates (0.89, Fig. 6a).
691

692
Figure 6. Log-log relationship between tunicate biomass and surface chlorophyll from adjusted observations
(blue) and model data (black), sampled at the same locations as the observational dataset, for (a) small
tunicates, (b) large tunicates, and (c) crustacean mesozooplankton. The observations were adjusted from the
data compilation to account for the systematic low sampling bias from nets (10-fold adjustment), with the grey
bars around the observational regression line showing the 5-15x adjustment range. Model values are from the
top 100-m, and the crustacean mesozooplankton biomass was computed as large mesozooplankton + 0.5*small
mesozooplankton. Observational surface chlorophyll, as well as model data, were time means from the
growing season.
701

The sensitivity tests illustrated the impact of various aspects of the base GZ-COBALT
parameter choices, as well as the distinct physiology of tunicates relative to crustacean
mesozooplankton. For large tunicates, the reduced basal respiration rate, relative to the literature-
based mean, in the base GZ-COBALT simulation was key for achieving a mean biomass
consistent with observations (see case 1, Fig. S2a-c). Similarly, for small tunicates, the lower
maximum ingestion rate relative to the literature-based values, was also essential for achieving a
mean biomass consistent with observations, and a biomass-chlorophyll scaling slope that did not
deviate too far from the observational constraints (case 2, Fig. S2d-f). In case 2, total tunicate
production also doubled, despite modest increases in biomass, largely due to the role of the small
tunicates (Table 4). The tunicate biomass-chlorophyll scaling slope was a result of many factors,
including competition between tunicate size classes, as the relative increase in small tunicate
biomass resulted in a shallower scaling slope for large tunicates (case 2, Fig. S2e).
Sensitivity cases 3-5 focused on model formulations distinct to pelagic tunicates relative
to the original crustacean mesozooplankton in COBALTv2. In case 3, where the ingestion half-
saturation constant (with associated adjustments to the maximum ingestion rate) was set to the
same value as that of the crustaceans, the resultant mean biomass and biomass-chlorophyll
scaling for both tunicates matched the observations more closely (Fig. S2g-i). This was an
interesting result and could have been a tuning choice; however, doing so would have negated a

key criterion (of the half-saturation constant, K_i, being much greater than the prey concentration)
in converting measurements of clearance rate to ingestion rate. Setting the assimilation efficiency
to a constant value (case 4) resulted in small tunicates being closer to observations, but large
tunicates dropping significantly in biomass, particularly in the low productivity areas (Fig. S2j-l).
This suggests that the variable assimilation efficiency was one factor in allowing large tunicates
to survive in the subtropical gyres. Finally, in the case where large tunicate aggregation mortality
was removed, this resulted in large tunicate biomass greatly increasing in the high chlorophyll
areas, with associated increases in the biomass scaling slope (Fig. S2m-o).
728
3.2 Seasonal cycle
All zooplankton exhibited a stronger seasonal cycle in the high chlorophyll seasonally
stratified (HCSS) biome compared to the high chlorophyll permanently stratified (HCPS) and
low chlorophyll (LC) biomes, with the biomass peak shifting later in the summer as zooplankton
size increases. GZ exhibited a late summer (August-September) peak for both small and large
tunicates. The large tunicates were also unique amongst zooplankton in that their biomass in the
HCSS biome did not exceed that of the HCPS biome (Fig. 7). Results from the sensitivity cases
showed that this is largely due to the large tunicate aggregation mortality, or jelly-falls (case 5,
Fig. S3k-o), which serves to strongly dampen blooms. Additionally, reductions in the ingestion
half-saturation constant (and associated maximum ingestion rate; case 3, Fig. S3a-e) and the
constant assimilation efficiency (case 4) also reduced the magnitude of the blooms. Additionally,
in case 4, the small tunicates' bloom timing was also shifted to be slightly earlier (Fig. S3f-j). In
the base GZ-COBALT configuration, the biomass of the non-GZ zooplankton were overall
reduced compared to the COBALTv2 control (-7%, -6%, and -7.2% for small, medium, and
large zooplankton, respectively), with the biggest difference seen in the summer
microzooplankton biomass in the HCPS biome. Other substantial differences included the
wintertime biomass of small crustacean mesozooplankton in the LC biome, due to the low
starting biomass in the control (Fig. 7).
747

748
Figure 7. Seasonal cycle of modeled (a) microzooplankton, (b) medium zooplankton, (c) large zooplankton,
(d) small tunicates, and (e) large tunicates, separated by biome. Biome definitions: LC = low chlorophyll,
HCSS = high chlorophyll seasonally stratified, HCPS = high chlorophyll permanently stratified. Southern
Hemisphere values were shifted six months such that Austral summer is represented by months 6-8. Solid lines
indicate the GZ-COBALT simulation, and dashed lines show zooplankton values from the COBALTv2 control
simulation.
755

3.3. Biogeochemical impacts

The overall impact of gelatinous zooplankton on the partitioning of energy between the microbial food web, export to depth, and energy available to higher trophic levels through mesozooplankton was assessed via the difference between GZ-COBALT and the control formulation (Fig. 8). This comparison suggests that the two tunicate classes have a competitive interaction with microzooplankton (Fig. 8c) and a small, but net negative impact of the total combined production of mesozooplankton (i.e., GZ and crustaceans, Fig. 8f). This is in spite of a competitive impact on crustacean zooplankton, which was greater for the small crustaceans, particularly in the upwelling zones, compared to the large crustaceans (Fig. S4).

The differences between the simulations becomes more pronounced when considering plankton functional types that dominate recycling vs. export processes. With the addition of pelagic tunicates, the routing of carbon to the microbial food-web decreased, as indicated by declines in both the heterotrophic bacteria production ratio and the microzooplankton production ratio (Fig. 8c, 7l). Meanwhile, the particle export ratio (pe-ratio, defined as the export flux at 100 m divided by NPP) increased globally (Fig. 8i). This comes as small and large tunicates contributed 0.19 and 0.79 Pg C y^{-1}, respectively, of total export production in the top 100 m (of which 72% sinks past 100 m). This increase in gelatinous-mediated export reflects a redistribution of export production from existing sources, with the largest coming from small mesozooplankton (Fig. 9a,c), as well as a reduction in the dissolved pool.

Figure 8. Differences in annual mean productivity ratios at the top 100 m between GZ-COBALT and the COBALTv2 control, showing ratios of microzooplankton production to NPP (a-c), mesozooplankton

production to NPP (d-f), POC export past 100 m to NPP, or pe-ratio (g-i), and free-living heterotrophic bacteria production to NPP (j-l). The plots show the GZ-COBALT simulation (left column; a,d,g,j), the COBALTv2 control (center column; b,e,h,k), and the difference between the two (right column; c,f,i,l).

Figure 9. Top 100-m production of sinking detritus (a, c) and loss to higher trophic level (HTL) predators (b, d) in the COBALTv2 control (a, b) and GZ-COBALT (c, d) simulations. Of the total export production, approximately 84% of it sinks below 100 m. All values are from the top 100 m, and are in Pg C y^{-1}.

Table 4. Comparison of the major results from the COBALTv2 control, GZ-COBALT base simulation, and the sensitivity experiments. 'z100' refers to the top 100 m of the water column.

Field	COBALT Control	GZ-COBALT Base	GZ-COBALT Sensitivity Experiments				
			Exp. 1	Exp. 2	Exp. 3	Exp. 4	Exp. 5
NPP (Pg C y^{-1})	55.4	53.7	55	51	54.5	54.5	53.6
Export at 100 m (Pg C y^{-1})	6.23	6.36	6.25	6.37	6.23	6.31	6.39
Total Grazing (Pg C y^{-1})	39.9	38.8	39.7	37.5	39.3	39.3	38.8
Zooplankton Ingestion, z100 (Pg C y^{-1})	53.6	51.1	53.1	47.9	52.2	52.4	50.9
HP Ingestion, z100 (Pg C y^{-1})	2.39	2.18	2.33	1.96	2.26	2.25	2.27
Total Phytoplankton Biomass (Pg C)	0.488	0.482	0.486	0.466	0.485	0.486	0.48
Total Zooplankton Biomass (Pg C)	0.659	0.719	0.659	0.676	0.706	0.677	0.766
Total Tunicate Biomass (Tg C)	---	102	8.69	112	73	36.9	161
Sm. Tunicate Biomass, z100 (Tg C)	---	5.78	6.13	37	3.73	3.2	5.53
Lg. Tunicate Biomass, z100 (Tg C)	---	81.5	1.08	61.5	58.1	26.8	132
Crustacean Mesozoop. Biomass (Pg C)	0.378	0.356	0.373	0.331	0.365	0.368	0.352
Tunicate Prod., z100 (Pg C y^{-1})	---	0.49	0.0361	1	0.259	0.0884	0.0724
Crust. Mesozoo. Prod., z100 (Pg C y^{-1})	4.54	3.89	4.39	3.31	4.14	4.21	2.77
Z-ratio	0.082	0.072	0.08	0.065	0.076	0.077	0.07
Sm. Tunicate Detritus Prod., z100 (Pg C y^{-1})	---	0.19	0.21	1.58	0.11	0.14	0.18
Lg. Tunicate Detritus Prod., z100 (Pg C y^{-1})	---	0.79	0.01	0.51	0.5	0.38	0.96
Tunicate Prod./Biomass, z100 (y^{-1})	---	5.61	5.02	10.2	4.19	2.95	5.25
Mesozoop Prod./Biomass, z100 (y^{-1})	14.70	13.4	14.3	12.3	13.9	14	13.1
Lg. Tunicate Prod. lost to HTL Pred.	---	57%	0%	56%	56%	51%	91%
Lg. Tunicate Prod. lost to Jelly-Falls	---	35%	4%	33%	34%	34%	0%
Slope of Sm. Tun/Chlorophyll scaling	---	0.89	0.9	1.1	0.67	0.79	0.88
Slope of Lg. Tun/Chlorophyll scaling	---	0.43	0.26	0.29	0.39	0.62	0.58

4. Discussion

We have added a simple representation of two pelagic tunicate groups, representing appendicularians and thaliaceans, into the GFDL COBALTv2 ocean biogeochemistry model (GZ-COBALT) that captures large-scale patterns in tunicate distribution consistent with the emerging recognition of their importance to marine ecosystems, while maintaining a skillful representation of crustacean mesozooplankton, surface chlorophyll, and macronutrient concentrations. The GZ-COBALT simulation achieved a reasonable match between the modelled mean tunicate biomass and a global observational dataset, compiled from a range of sources including the COPEPOD database (O'Brien, 2014, 2005), Jellyfish Environmental Database Initiative (JeDI) (Condon et al., 2015), and KRILLBASE (Atkinson et al., 2017). Notably, GZ-COBALT captured a contrast between gelatinous and crustacean zooplankton types in their emergent relationship between biomass and surface chlorophyll (Fig. 6). These results confirm that it is possible to reconcile GZ biomass an order of magnitude above previous estimates (Remsen et al., 2004) with prevalent crustacean zooplankton populations: carbon flows through planktonic food-webs are sufficient to support both GZ and crustacean populations.

Observations of tunicate biomass exhibited high variability, even when compared with crustacean zooplankton observations gridded to the same horizontal resolution (Moriarty and O'Brien, 2013), which indicates either large sampling variability (e.g., from inconsistency in sampling effort and/or gear), or unresolved physical or biological dynamics (Andersen, 1998; Boero et al., 2008; d'Ovidio et al., 2010; Greer et al., 2020; Lévy et al., 2018; Luo et al., 2014), or a combination thereof. The modelled variability for tunicates was lower than the observations may suggest, even when daily rather than monthly outputs were sampled (Fig. S5). This was also apparent for crustacean zooplankton, as it is a near-universal outcome when comparing global biogeochemical fields against point measurements or averages of small numbers of point measures (e.g., Krumhardt et al., 2017; Martiny et al., 2019; Saba et al., 2011; Usbeck et al., 2003). The discrepancy is admittedly acute for the tunicates, which was not unexpected given the sparsity and difficulty of measurements. A more complete understanding the drivers of this patchiness and their implications will likely require high resolution physical simulations and GZ models capable of better resolving unique aspects of GZ life cycles and ecology conducive to patch formation (Groeneveld et al., 2020; Henschke et al., 2018a, 2018b)

In our analysis, we found a strong contrast in the biomass-chlorophyll relationship between crustacean zooplankton, small tunicates, and large tunicates, wherein the large tunicates exhibited a flatter scaling relationship compared to the steeper scalings of the small tunicates and crustaceans (Fig. 6). After incorporating an expanded view of GZ biomass considering undersampling by nets, the resultant observational biomass-chlorophyll scaling became one of our primary validation tools, as this emergent relationship can capture the mean biomass responses across productivity gradients. These relationships become important as steep spatial gradients in the contemporary ocean generally translates to amplified trends with climate change (Stock et al., 2017, 2014b). The shallower slope for large tunicates relative to crustacean zooplankton, in contrast, is indicative of less sensitivity to NPP, and suggestive of greater resilience to NPP declines projected by the majority of models under high emissions scenarios (Kwiatkowski et al., 2020) than their crustacean competitors. This would be consistent with current hypotheses for increased prevalence of GZ under climate change (Henschke et al., 2016).

4.1 Marine food web and biogeochemical impacts

Pelagic tunicates have long been identified as a potentially important source of carbon export, due to fecal pellets from salps (Iversen et al., 2017; Madin et al., 2006; Ramaswamy et al., 2005; Smith Jr et al., 2014; Urrère and Knauer, 1981) and appendicularians (Wilson et al., 2013), discarded appendicularian houses (Berline et al., 2011; Lombard and Kiørboe, 2010; Robison, 2005), and salp and pyrosome carcasses from jelly-falls (Henschke et al., 2013; Lebrato et al., 2013; Lebrato and Jones, 2009). Given the boom-and-bust population dynamic of pelagic tunicates, they can often be found to dominate POC export when present (Madin et al., 2006; Smith Jr et al., 2014). Indeed, a recent study from a NASA EXPORTS cruise found that salp fecal pellets comprised up to 80% of the detrital production in the upper 100 m in the NE Pacific when present, though they contributed an average of 28% of fecal pellet carbon production over a month-long sampling period (Stamieszkin et al., 2021). In our 20-year model climatology, large tunicate detritus production comprises 20% of the total detritus production in the top 100 m from July-September in the same region (Fig. S6). These values are a bit lower, but still fairly consistent with the sampled cruise mean, though the high observed variability in Stamieszkin et al. (2021) highlights the challenge in model-observation comparisons with snapshot studies at a single time point. Model comparisons with GZ-COBALT and sediment trap data, which integrates observations over longer time scales, will need to incorporate tunicate-specific POC sinking speeds and is a target for future work.

One common implication of observations of pelagic tunicate-mediated carbon export is that they would add to the existing POC export out of the surface ocean, often attributed to a combination of phyto-detritus and crustacean zooplankton fecal pellets (Buesseler et al., 2008; De La Rocha and Passow, 2007). Here, we found that when considering the upper oceans (top 100 m), the integration of pelagic tunicates with a "traditional" food-web model did not substantially increase total export flux past 100 m, which was 6.36 Pg C y^{-1} compared with 6.23 Pg C y^{-1} in the COBALTv2 control despite GZ accounting for 0.7 Pg C y^{-1}. The integration of GZ thus led primarily to a redistribution of fluxes away from those previously attributed to crustacean zooplankton, rather than a creation of a new additive flux. The modest increase in particle export that did occur is consistent with compensation for reductions in dissolved organic carbon export arising from GZ-induced redirection of carbon flows away from the microbial food-web (Fig. 8, 9).

Compared to the offline estimates of tunicate export (1.3-3.9 Pg C y^{-1} at 100 m; Luo et al., 2020), the online GZ-COBALT model was lower, suggesting that food web and biogeochemical feedbacks decreased the overall export contribution of tunicates. Rather than relying on the direct application of GZ data, GZ-COBALT accomplishes this correspondence by mechanistically representing the primary observational features and satisfying myriad additional physical, biogeochemical, and food-web constraints. Large tunicates contributed about four times more export production than small tunicates in GZ-COBALT (0.79 vs. 0.19 Pg C y^{-1}), with approximately 0.16 Pg C y^{-1} from jelly-falls, which was only slightly lower than the offline estimates of 0.3-0.7 Pg C y^{-1}. Tunicates in GZ-COBALT also contributed 0.1 Pg C y^{-1} to higher trophic level predators (Fig. 9d), which was much lower than the offline estimates of 0.8-1.1 Pg C y^{-1} (Luo et al., 2020). The higher trophic level predation in the offline model was one of the least constrained parameters, as Luo et al. (2020) extracted a total GZ ecotrophic efficiency (fraction of production to predation) from a combination of EcoPath models (e.g., Ruzicka et al., 2020), and tuned this term for individual GZ groups to achieve a global fraction consistent with EcoPath estimates. Future observational and experimental work aimed to increasing our

understanding of GZ predation by higher trophic levels should reduce the uncertainties
associated with these global models.
The GZ-COBALT simulation showed that, compared with the COBALTv2 control, the
largest impact of pelagic tunicates to ocean biogeochemical cycles is in the partitioning between
the biological pump and the microbial food web. In GZ-COBALT, the impact of tunicates served
to reduce microzooplankton and bacterial production as a function of NPP by 14% and 4%,
respectively (Fig. 8). Pelagic tunicates, unlike other gelatinous zooplankton, are notable for
primarily grazing on small particles and their high predator to prey size ratios (Conley et al.,
2018; Sutherland et al., 2010), though some exceptions exist (Post, 2002; Walters et al., 2019).
Recent work from Stukel et al. (2021) showed that in the Southern Ocean, the dominant salp, *S.*
thompsoni, most strongly competed with protistan grazers instead of with krill due to the large
size-based overlap between the salp and protistan diets. This is in contrast to previous
speculation that salps are a dominant competitor of the Antarctic krill, *Euphausia superba*, and
can be implicated as a factor in its long-term decline (Atkinson et al., 2004) and is consistent
with recent evidence that this decline can be attributed to positive anomalies in the Southern
Annular Mode (SAM) and loss of sea ice in the Southern Ocean (Atkinson et al., 2019). Our
results indicate that while tunicates do compete with large crustacean mesozooplankton for prey,
namely through the grazing of large phytoplankton and diatoms by appendicularians and
doliolids, tunicates also serve as a source of food for both small and large crustacean
mesozooplankton. Instead of competing with crustaceans, the magnitude of decline of
microzooplankton and heterotrophic bacteria production in GZ-COBALT compared to the
control and agreement with observations indicates that the microphagous tunicates serve as a
trophic and carbon export shunt away from the microbial loop and towards the mesozooplankton
food web and biological pump.
911
4.2 Model limitations
Amongst the marine zooplankton, thaliaceans are notable for their complex life cycles
which include the ability to reproduce asexually, alternation between sexual and asexual
reproductive phases (salps and doliolids), and hermaphroditism (pyrosomes), all of which can
yield large, transient, blooms under the right conditions (Andersen, 1998). Here, we have opted
against modeling the complex life cycle of pelagic tunicates (Henschke et al., 2018a, 2015;
Lombard et al., 2009b) for a more simple representation (Berline et al., 2011) aimed at capturing
their mean state, seasonal fluctuations, and long-term trends that can be run in an Earth System
Model over O(100) years. As such, there were a number of necessary simplifications, and
associated insights.
The model suggests that the mean turnover time scale, as measured by the ratio of
production over biomass, or P/B, for pelagic tunicates is overall lower (implying slower growth)
than microzooplankton and crustacean mesozooplankton (Table 4, Fig. S7). While some shallow
coastal areas exhibited P/B exceeding 10% d^{-1} in the summer for both small and large tunicates,
the majority of the oceans had turnover rates < 3% d^{-1}, even in the summer months. In contrast,
the turnover timescales for large tunicates as reported in the literature were 15-71% d^{-1} (Deibel,
1982; Gibson and Paffenhöfer, 2000; Madin and Purcell, 1992). While there may be some
averaging due to the model's monthly output, not even daily data captured the range of
variability in the observations (Fig. S5). Future efforts may focus on determining whether the
model's inability to reproduce observed variability is due to its coarse horizontal resolution
relative to the scales of observed variations in tunicate distributions (Greer et al., 2021; Luo et

al., 2014), or due to the representation of the simplified life cycle. For some gelatinous
zooplankton populations, a representation of the complex life cycle may be key for reproducing
interannual and multi-decadal climate fluctuations (e.g., Henschke et al., 2018b).
936
4.3 Future outlook
Gelatinous zooplankton (GZ) are ubiquitous throughout the world's oceans and a key
contributor to marine food webs (Hays et al., 2018). Of the GZ, pelagic tunicates are likely the
most important group in terms of carbon fluxes, due to their low trophic position and
microphagous diet. We demonstrate, through a new model with food-web and biogeochemical
feedbacks incorporated, that it is possible to reconcile an enhanced role of GZ in marine food
webs with the established importance of crustacean mesozooplankton and other ocean
biogeochemical constraints. Simulation results provide GZ flux estimates arising from a self-
consistent physical-biological model satisfying myriad physical, biogeochemical and plankton
food-web constraints, a unique contribution relative to previous "offline" estimates. Climate
change is projected to drive decreases in NPP; coupled Model Intercomparison Project Phase 6
(CMIP6) models under the Shared Socioeconomic Pathway 5 (SSP5; fossil-fueled development)
project a 3-9% decline in NPP by the year 2100 (Kwiatkowski et al. 2020). Associated with
climate-induced NPP decreases, models also project a shift in mean pelagic body size: the
abundance of large autotrophic phytoplankton will likely be reduced relative to their smaller
counterparts due to increased warming, stratification, and subsequent nutrient limitation (Peter
and Sommer 2013). Consequently, as evidenced by the shallow scaling between biomass and
chlorophyll, the role of large pelagic tunicates (thaliaceans) in marine food webs may further
increase under climate change.
In this study, we have focused primarily on the upper ocean impacts of GZ, both to the
food web and to the balance between recycling and export. Omitted in this work are
considerations of the impact of fast sinking GZ export on the remineralization length scale and
transfer efficiency to "sequestration depths", which may have further impacts on benthic fluxes
and air-sea CO_2 exchange (Kwon et al., 2009; Lebrato et al., 2019; Luo et al., 2020; Sweetman et
al., 2014; Titelman et al., 2006). In particular, there may be important feedbacks between
climate-induced stratification and tunicate-mediated increases in export. Our results indicate that
total carbon export was not significantly increased with inclusion of GZ in an ocean
biogeochemical model. However, these tunicate fluxes are globally quite significant and are
associated with a redistribution of export from existing phytoplankton and mesozooplankton
sources. As climate change will have differing impacts by taxonomic group, better understanding
of the sources of carbon export and the mechanisms that drive their variation will improve our
ability to project changes in the future.

Acknowledgements
We thank colleagues who have contributed to the development and integration of various components of GFDL's Earth System Model that make this work possible, as well as the technical support teams that maintain the NOAA/GFDL computing resources. JYL acknowledges support from the NOAA's Marine Ecosystem Tipping Points Initiative. This work also benefited from internal reviews by E. Drenkard and C. Schultz.

Author contributions
JYL and CAS conceived and designed the study, and evaluated the model. JYL wrote the code, carried out simulations, analyzed data, and led the manuscript writing. JYL and TOB compiled data. All authors contributed critical feedback and edits to the final manuscript.

Competing Interests
The authors have no competing interests to declare.

References

Acuña, J.L., Kiefer, M., 2000. Functional response of the appendicularian *Oikopleura dioica*. Limnol. Oceanogr. 45, 608–618. https://doi.org/10.4319/lo.2000.45.3.0608

Acuña, J.L., Lopez-Urrutia, A., Colin, S., 2011. Faking Giants: The Evolution of High Prey Clearance Rates in Jellyfishes. Science 333, 1627–1629. https://doi.org/Doi 10.1126/Science.1205134

Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C.O., Dunne, J.P., Griffies, S.M., Hallberg, R., Harrison, M.J., Held, I.M., Jansen, M.F., John, J.G., Krasting, J.P., Langenhorst, A.R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan, A., Reichl, B.G., Rosati, T., Samuels, B.L., Shao, A., Stouffer, R., Winton, M., Wittenberg, A.T., Xiang, B., Zadeh, N., Zhang, R., 2019. The GFDL Global Ocean and Sea Ice Model OM4.0: Model Description and Simulation Features. J. Adv. Model. Earth Syst. 11, 3167–3211. https://doi.org/10.1029/2019MS001726

Alldredge, A.L., 2005. The contribution of discarded appendicularian houses to the flux of particulate organic carbon from oceanic surface waters, in: Gorsky, G., Youngbluth, M.J., Deibel, D. (Eds.), Response of Marine Ecosystems to Global Change: Ecological Impact of Appendicularians. Éditions Scientifiques, Paris, pp. 309–326.

Alldredge, A.L., Silver, M.W., 1988. Characteristics, dynamics and significance of marine snow. Progress in Oceanography 20, 41–82. https://doi.org/10.1016/0079-6611(88)90053-5

Andersen, V., 1998. Salp and pyrosomid blooms and their importance in biogeochemical cycles, in: Bone, Q. (Ed.), The Biology of Pelagic Tunicates. Oxford University Press, New York, USA, pp. 125–137.

Andersen, V., 1986. Effect of temperature on the filtration rate and percentage of assimilation of Salpa fusiformis Cuvier (Tunicata: Thaliacea). Hydrobiologia 137, 135–140.

Atkinson, A., Hill, S.L., Pakhomov, E.A., Siegel, V., Anadon, R., Chiba, S., Daly, K.L., Downie, R., Fielding, S., Fretwell, P., Gerrish, L., Hosie, G.W., Jessopp, M.J., Kawaguchi, S., Krafft, B.A., Loeb, V., Nishikawa, J., Peat, H.J., Reiss, C.S., Ross, R.M., Quetin, L.B., Schmidt, K., Steinberg, D.K., Subramaniam, R.C., Tarling, G.A., Ward, P., 2017. KRILLBASE: A circumpolar database of Antarctic krill and salp numerical densities, 1926-2016. Earth System Science Data. https://doi.org/10.5194/essd-9-193-2017

Atkinson, A., Hill, S.L., Pakhomov, E.A., Siegel, V., Reiss, C.S., Loeb, V.J., Steinberg, D.K., Schmidt, K., Tarling, G.A., Gerrish, L., Sailley, S.F., 2019. Krill (Euphausia superba) distribution contracts southward during rapid regional warming. Nature Clim Change 9, 142–147. https://doi.org/10.1038/s41558-018-0370-z

Atkinson, A., Siegel, V., Pakhomov, E., Rothery, P., 2004. Long-term decline in krill stock and increase in salps within the Southern Ocean. Nature 432, 100–103. https://doi.org/10.1038/nature02996

Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., Gehlen, M., 2015. PISCES-v2: An ocean biogeochemical model for carbon and ecosystem studies. Geoscientific Model Development. https://doi.org/10.5194/gmd-8-2465-2015

Baker, A.R., Croot, P.L., 2010. Atmospheric and marine controls on aerosol iron solubility in seawater. Marine Chemistry 120, 4–13. https://doi.org/10.1016/j.marchem.2008.09.003

Banse, K., 1992. Grazing, Temporal Changes of Phytoplankton Concentrations, and the Microbial Loop in the Open Sea, in: Primary Productivity and Biogeochemical Cycles in the Sea. https://doi.org/10.1007/978-1-4899-0762-2_22

Berline, L., Stemmann, L., Vichi, M., Lombard, F., Gorsky, G., 2011. Impact of appendicularians on detritus and export fluxes: a model approach at DyFAMed site. Journal of Plankton Research 33, 855–872. https://doi.org/10.1093/plankt/fbq163

Billett, D.S.M., Bett, B.J., Jacobs, C.L., Rouse, I.P., Wigham, B.D., 2006. Mass deposition of jellyfish in the deep Arabian Sea. Limnology and Oceanography 51, 2077–2083. https://doi.org/10.4319/lo.2006.51.5.2077

Boero, F., Bouillon, J., Gravili, C., Miglietta, M.P., Parsons, T., Piraino, S., 2008. Gelatinous plankton: Irregularities rule the world (sometimes). Marine Ecology Progress Series. https://doi.org/10.3354/meps07368

Buesseler, K.O., Trull, T.W., Steinberg, D.K., Silver, M.W., Siegel, D.A., Saitoh, S.-I., Lamborg, C.H., Lam, P.J., Karl, D.M., Jiao, N.Z., Honda, M.C., Elskens, M., Dehairs, F., Brown, S.L., Boyd, P.W., Bishop, J.K.B., Bidigare, R.R., 2008. VERTIGO (VERtical Transport In the Global Ocean): A study of particle sources and flux attenuation in the North Pacific. Deep Sea Research Part II: Topical Studies in Oceanography 55, 1522–1539. https://doi.org/10.1016/j.dsr2.2008.04.024

Buitenhuis, E., Le Quéré, C., Aumont, O., Beaugrand, G., Bunker, A., Hirst, A., Ikeda, T., O'Brien, T., Piontkovski, S., Straile, D., 2006. Biogeochemical fluxes through mesozooplankton. Global Biogeochem. Cycles 20. https://doi.org/10.1029/2005GB002511

Carlotti, F., Giske, J., Werner, F., 2000. Modeling zooplankton dynamics, in: Harris, R., Wiebe, P., Lenz, J., Skjoldal, H.R., Huntley, M. (Eds.), ICES Zooplankton Methodology Manual. Academic Press, London, pp. 571–667. https://doi.org/10.1016/B978-012327645-2/50013-X

Condon, R.H., Graham, W.M., Duarte, C.M., Pitt, K.A., Lucas, C.H., Haddock, S.H.D., Sutherland, K.R., Robinson, K.L., Dawson, M.N., Decker, M.B., Mills, C.E., Purcell, J.E., Malej, A., Mianzan, H., Uye, S.I., Gelcich, S., Madin, L.P., 2012. Questioning the Rise of Gelatinous Zooplankton in the World's Oceans. BioScience 62, 160–169. https://doi.org/Doi 10.1525/Bio.2012.62.2.9

Condon, R.H., Lucas, C.H., Duarte, C.M., Pitt, K.A., D., H.S.H., Madin, L.P., D., B.R., Sutherland K. R., Mianzan H. W., Purcell J. E., Decker M. B., Uye S.-I., Malej A., Bogeberg M., Everett J., Gibbons M., Gonzalez H., Hay S., Henschke, N., Hobson R. J., Kingsford M. J., Kremer P., Lehtiniemi M., Ohman M. D., Rissik D., Sheard K., Suthers I., Coleman N., Costello J.H., Gershwin L.A., Graham W.M., Robinson K.L., Richardson T.M., Giesecke R., Gorsky G., Greve W., Halsband-Lenk C., Hays G., Hobson V., Klein D., Lebrato, M., J., L., P., M., C., M., Perry G., Stemmann L., Sullivan B., Walker T., Schildhauer M., Regetz, J., 2015. Jellyfish Database Initiative (JeDI). https://doi.org/10.1575/1912/7191

Conley, K.R., Lombard, F., Sutherland, K.R., 2018. Mammoth grazers on the ocean's minuteness: A review of selective feeding using mucous meshes, Proceedings of the Royal Society B: Biological Sciences. https://doi.org/10.1098/rspb.2018.0056

d'Ovidio, F., De Monte, S., Alvain, S., Dandonneau, Y., Levy, M., 2010. Fluid dynamical niches of phytoplankton types. Proceedings of the National Academy of Sciences 107, 18366–18370. https://doi.org/10.1073/pnas.1004620107

Dadon-Pilosof, A., Lombard, F., Genin, A., Sutherland, K.R., Yahel, G., 2019. Prey taxonomy rather than size determines salp diets. Limnol Oceanogr 64, 1996–2010. https://doi.org/10.1002/lno.11165

De La Rocha, C.L., Passow, U., 2007. Factors influencing the sinking of POC and the efficiency of the biological carbon pump. Deep Sea Research Part II: Topical Studies in Oceanography 54, 639–658. https://doi.org/10.1016/j.dsr2.2007.01.004

Deibel, D., 1998. Feeding and metabolism of Appendicularia, in: Bone, Q. (Ed.), The Biology of Pelagic Tunicates. Oxford University Press, New York, USA, pp. 139–149.

Deibel, D., 1982. Laboratory determined mortality, fecundity and growth rates of Thalia democratica Forskal and Dolioletta gegenbauri Uljanin (Tunicata, Thaliacea). Journal of Plankton Research 4, 143–153. https://doi.org/10.1093/plankt/4.1.143

Deibel, D., Lee, S., 1992. Retention efficiency of sub-micrometer particles by the pharyngeal filter of the pelagic tunicate Oikopleura vanhoeffeni. Mar. Ecol. Prog. Ser. 81, 25–30. https://doi.org/10.3354/meps081025

Dubischar, C.D., Pakhomov, E.A., Bathmann, U.V., 2006. The tunicate Salpa thompsoni ecology in the Southern Ocean. II. Proximate and elemental composition. Mar Biol 149, 625–632. https://doi.org/10.1007/s00227-005-0226-8

Dunne, J.P., Horowitz, L.W., Adcroft, A.J., Ginoux, P., Held, I.M., John, J.G., Krasting, J.P., Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C.A., Zadeh, N., Balaji, V., Blanton, C., Dunne, K.A., Dupuis, C., Durachta, J., Dussin, R., Gauthier, P.P.G., Griffies, S.M., Guo, H., Hallberg, R.W., Harrison, M., He, J., Hurlin, W., McHugh, C., Menzel, R., Milly, P.C.D., Nikonov, S., Paynter, D.J., Ploshay, J., Radhakrishnan, A., Rand, K., Reichl, B.G., Robinson, T., Schwarzkopf, D.M., Sentman, L.T., Underwood, S., Vahlenkamp, H., Winton, M., Wittenberg, A.T., Wyman, B., Zeng, Y., Zhao, M., 2020. The GFDL Earth System Model version 4.1 (GFDL-ESM 4.1): Overall coupled model description and simulation characteristics. J. Adv. Model. Earth Syst. https://doi.org/10.1029/2019MS002015

Edwards, K.F., Thomas, M.K., Klausmeier, C.A., Litchman, E., 2015. Light and growth in marine phytoplankton: Allometric, taxonomic, and environmental variation. Limnology and Oceanography. https://doi.org/10.1002/lno.10033

Edwards, K.F., Thomas, M.K., Klausmeier, C.A., Litchman, E., 2012. Allometric scaling and taxonomic variation in nutrient utilization traits and maximum growth rate of phytoplankton. Limnology and Oceanography. https://doi.org/10.4319/lo.2012.57.2.0554

Eppley, R., 1972. Temperature and phytoplankton growth in the sea. Fishery Bulletin.

Fasham, M.J.R., Ducklow, H.W., McKelvie, S.M., 1990. A nitrogen-based model of plankton dynamics in the oceanic mixed layer. Journal of Marine Research. https://doi.org/10.1357/002224090784984678

Fernández, D., López-Urrutia, Á., Fernández, A., Acuña, J., Harris, R., 2004. Retention efficiency of 0.2 to 6 μm particles by the appendicularians Oikopleura dioica and Fritillaria borealis. Mar. Ecol. Prog. Ser. 266, 89–101. https://doi.org/10.3354/meps266089

Flynn, K.J., 2005. Incorporating plankton respiration in models of aquatic ecosystem function, in: del Giorgio, P., Williams, P. (Eds.), Respiration in Aquatic Ecosystems. Oxford University Press, pp. 248–266. https://doi.org/10.1093/acprof:oso/9780198527084.003.0013

Garcia, H.E., Boyer, T.P., Locarnini, R.A., Antonov, J.I., Mishonov, A.V., Baranova, O.K., Zweng, M.M., Reagan, J.R., Johnson, D.R., 2013a. World Ocean Atlas 2013. Volume 3: Dissolved oxygen, apparent oxygen utilization, and oxygen saturation.

Garcia, H.E., Locarnini, R.A., Boyer, T.P., Antonov, J.I., Baranova, O.K., Zweng, M.M., Reagan, J.R., Johnson, D.R., 2013b. World Ocean Atlas 2013, Volume 4 : Dissolved Inorganic Nutrients (phosphate, nitrate, silicate).

Garcia, H.E., Weathers, K.W., Paver, C.R., Smolyar, I.V., Boyer, T.P., Locarnini, R.A., Zweng, M.M., Mishonov, A.V., Baranova, O.K., Seidov, D., Reagan, J.R., 2019. World Ocean Atlas 2018. Vol. 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate).

Gentleman, W., Leising, A., Frost, B., Strom, S., Murray, J., 2003. Functional responses for zooplankton feeding on multiple resources: A review of assumptions and biological dynamics. Deep-Sea Research Part II: Topical Studies in Oceanography. https://doi.org/10.1016/j.dsr2.2003.07.001

Gibson, D.M., Paffenhöfer, G.-A., 2000. Feeding and growth rates of the doliolid, Dolioletta gegenbauri Uljanin (Tunicata, Thaliacea). Journal of Plankton Research 22, 1485–1500. https://doi.org/10.1093/plankt/22.8.1485

Gorsky, G., Fenaux, R., 1998. The role of appendicularia in marine food webs, in: Bone, Q. (Ed.), The Biology of Pelagic Tunicates. Oxford University Press, New York, USA, pp. 161–169.

Greer, A.T., Boyette, A.D., Cruz, V.J., Cambazoglu, M.K., Dzwonkowski, B., Chiaverano, L.M., Dykstra, S.L., Briseño-Avena, C., Cowen, R.K., Wiggert, J.D., 2020. Contrasting fine-scale distributional patterns of zooplankton driven by the formation of a diatom-dominated thin layer. Limnol Oceanogr 65, 2236–2258. https://doi.org/10.1002/lno.11450

Greer, A.T., Chiaverano, L.M., Treible, L.M., Briseño-Avena, C., Hernandez, F.J., 2021. From spatial pattern to ecological process through imaging zooplankton interactions. ICES Journal of Marine Science fsab149. https://doi.org/10.1093/icesjms/fsab149

Groeneveld, J., Berger, U., Henschke, N., Pakhomov, E.A., Reiss, C.S., Meyer, B., 2020. Blooms of a key grazer in the Southern Ocean – An individual-based model of Salpa thompsoni. Progress in Oceanography 185, 102339. https://doi.org/10.1016/j.pocean.2020.102339

Hagadorn, J.W., Dott, R.H., Damrow, D., 2002. Stranded on a Late Cambrian shoreline: Medusae from central Wisconsin. Geology 30, 147–150.

Hansen, B., Bjørnsen, P.K., Hansen, P.J., 1994. The size ratio between planktonic predators and their prey. Limnology and Oceanography 39, 395–403. https://doi.org/10.4319/lo.1994.39.2.0395

Hansen, P.J., Bjørnsen, P.K., Hansen, B.W., 1997. Zooplankton grazing and growth: Scaling within the 2-2,000-µm body size range. Limnology and Oceanography. https://doi.org/10.4319/lo.1997.42.4.0687

Harbison, G.R., 1998. The parasites and predators of Thaliacea, in: Bone, Q. (Ed.), The Biology of Pelagic Tunicates. Oxford University Press, New York, USA, pp. 187–214.

Harbison, G.R., McAlister, V.L., Gilmer, R.W., 1986. The Response of the Salp, Pegea confoederata, to High Levels of Particulate Material: Starvation in the Midst of Plenty. Limnology and Oceanography 31, 371–382.

Hays, G.C., Doyle, T.K., Houghton, J.D.R., 2018. A Paradigm Shift in the Trophic Importance of Jellyfish?, Trends in Ecology and Evolution. https://doi.org/10.1016/j.tree.2018.09.001

Henschke, N., Bowden, D.A., Everett, J.D., Holmes, S.P., Kloser, R.J., Lee, R.W., Suthers, I.M., 2013. Salp-falls in the Tasman Sea: a major food input to deep-sea benthos. Marine Ecology Progress Series 491, 165–175. https://doi.org/10.3354/meps10450

Henschke, N., Everett, J.D., Richardson, A.J., Suthers, I.M., 2016. Rethinking the Role of Salps in the Ocean. Trends in Ecology and Evolution 31, 720–733. https://doi.org/10.1016/j.tree.2016.06.007

Henschke, N., Pakhomov, E.A., 2019. Latitudinal variations in *Salpa thompsoni* reproductive fitness. Limnol. Oceanogr. 64, 575–584. https://doi.org/10.1002/lno.11061

Henschke, N., Pakhomov, E.A., Groeneveld, J., Meyer, B., 2018a. Modelling the life cycle of Salpa thompsoni. Ecological Modelling 387, 17–26. https://doi.org/10.1016/j.ecolmodel.2018.08.017

Henschke, N., Smith, J.A., Everett, J.D., Suthers, I.M., 2015. Population drivers of a Thalia democratica swarm: Insights from population modelling. Journal of Plankton Research 37, 1074–1087. https://doi.org/10.1093/plankt/fbv024

Henschke, N., Stock, C.A., Sarmiento, J.L., 2018b. Modeling population dynamics of scyphozoan jellyfish (Aurelia spp.) in the Gulf of Mexico. Marine Ecology Progress Series. https://doi.org/10.3354/meps12255

Hopcroft, R.R., Roff, J.C., 1998. Production of tropical larvaceans in Kingston Harbour, Jamaica: are we ignoring an important secondary producer? J Plankton Res 20, 557–569. https://doi.org/10.1093/plankt/20.3.557

Horowitz, L.W., Walters, S., Mauzerall, D.L., Emmons, L.K., Rasch, P.J., Granier, C., Tie, X., Lamarque, J.-F., Schultz, M.G., Tyndall, G.S., Orlando, J.J., Brasseur, G.P., 2003. A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2: MOZART-2 DESCRIPTION AND EVALUATION. J. Geophys. Res. 108, n/a-n/a. https://doi.org/10.1029/2002JD002853

Iversen, M.H., Pakhomov, E.A., Hunt, B.P.V., van der Jagt, H., Wolf-Gladrow, D., Klaas, C., 2017. Sinkers or floaters? Contribution from salp pellets to the export flux during a large bloom event in the Southern Ocean. Deep-Sea Research Part II: Topical Studies in Oceanography 138, 116–125. https://doi.org/10.1016/j.dsr2.2016.12.004

Johnson, R., Strutton, P.G., Wright, S.W., McMinn, A., Meiners, K.M., 2013. Three improved satellite chlorophyll algorithms for the Southern Ocean: SOUTHERN OCEAN CHLOROPHYLL ALGORITHMS. J. Geophys. Res. Oceans 118, 3694–3703. https://doi.org/10.1002/jgrc.20270

Kearney, K.A., Bograd, S.J., Drenkard, E., Gomez, F.A., Haltuch, M., Hermann, A.J., Jacox, M.G., Kaplan, I.C., Koenigstein, S., Luo, J.Y., Masi, M., Muhling, B., Pozo Buil, M., Woodworth-Jefcoats, P.A., 2021. Using Global-Scale Earth System Models for Regional Fisheries Applications. Front. Mar. Sci. 8, 622206. https://doi.org/10.3389/fmars.2021.622206

Kiørboe, T., Møhlenberg, F., Hamburger, K., 1985. Bioenergetics of the planktonic copepod Acartia tonsa: relation between feeding, egg production and respiration, and composition of specific dynamic action. Mar. Ecol. Prog. Ser. 26, 85–97. https://doi.org/10.3354/meps026085

Krumhardt, K.M., Lovenduski, N.S., Iglesias-Rodriguez, M.D., Kleypas, J.A., 2017. Coccolithophore growth and calcification in a changing ocean. Progress in Oceanography 159, 276–295. https://doi.org/10.1016/j.pocean.2017.10.007

Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., R. Christian, J., P. Dunne, J., Gehlen, M., Ilyina, T., G. John, J., Lenton, A., Li, H., S. Lovenduski, N., C. Orr, J., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., A. Stock, C., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., Ziehn, T., 2020. Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections. Biogeosciences 17, 3439–3470. https://doi.org/10.5194/bg-17-3439-2020

Kwon, E.Y., Primeau, F., Sarmiento, J.L., 2009. The impact of remineralization depth on the air–sea carbon balance. Nature Geoscience 2, 630–635. https://doi.org/10.1038/ngeo612

Large, W.G., Yeager, S.G., 2009. The global climatology of an interannually varying air - Sea flux data set. Climate Dynamics. https://doi.org/10.1007/s00382-008-0441-3

Lauvset, S.K., Key, R.M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F.F., Suzuki, T., Watelet, S., 2016. A new global interior ocean mapped climatology: the 1° × 1° GLODAP version 2. Earth Syst. Sci. Data 8, 325–340. https://doi.org/10.5194/essd-8-325-2016

Laval, P., 1980. Hyperiid Amphipods as Crustacean Parasitoids associated with Gelatinous Zooplankton, in: Barnes, M. (Ed.), Oceanogr. Mar. Biol. Ann. Rev. Aberdeen University Press, pp. 11–56.

Lebrato, M., Jones, D.O.B., 2009. Mass deposition event of Pyrosoma atlanticum carcasses off Ivory Coast (West Africa). Limnology and Oceanography 54, 1197–1209. https://doi.org/10.4319/lo.2009.54.4.1197

Lebrato, M., Molinero, J.C., Cartes, J.E., Lloris, D., Melin, F., Beni-Casadella, L., 2013. Sinking Jelly-Carbon Unveils Potential Environmental Variability along a Continental Margin. PLoS ONE 8, e82070. doi:10.1371/journal.pone.0082070. https://doi.org/10.1371/journal.pone.0082070

Lebrato, M., Pahlow, M., Frost, J.R., Küter, M., Mendes, P.D.J., Molinero, J.C., Oschlies, A., 2019. Sinking of Gelatinous Zooplankton Biomass Increases Deep Carbon Transfer Efficiency Globally. Global Biogeochemical Cycles 33. https://doi.org/10.1029/2019GB006265

Lebrato, M., Pitt, K., Sweetman, A.K., Jones, D., Cartes, J., Oschlies, A., Condon, R., Molinero, J., Adler, L., Gaillard, C., Lloris, D., Billett, D., 2012. Jelly-falls historic and recent observations: a review to drive future research directions. Hydrobiologia 690, 227–245. https://doi.org/10.1007/s10750-012-1046-8

Lévy, M., Franks, P.J.S., Smith, K.S., 2018. The role of submesoscale currents in structuring marine ecosystems. Nat Commun 9, 4758. https://doi.org/10.1038/s41467-018-07059-3

Llopiz, J.K., Richardson, D.E., Shiroza, A., Smith, S.L., Cowen, R.K., 2010. Distinctions in the diets and distributions of larval tunas and the important role of appendicularians. Limnology and Oceanography 55, 983–996. https://doi.org/10.4319/lo.2010.55.3.0983

Locarnini, R.A., Mishonov, A.V., Antonov, J.I., Boyer, T.P., Garcia, H.E., Baranova, O.K., Zweng, M.M., Paver, C.R., Reagan, J.R., Johnson, D.R., Hamilton, M., Seidov, D., 2013. World Ocean Atlas 2013, Volume 1: Temperature.

Lombard, F., Kiørboe, T., 2010. Marine snow originating from appendicularian houses: Age-dependent settling characteristics. Deep Sea Research Part I: Oceanographic Research Papers 57, 1304–1313. https://doi.org/10.1016/j.dsr.2010.06.008

Lombard, F., Renaud, F., Sainsbury, C., Sciandra, A., Gorsky, G., 2009a. Appendicularian ecophysiology I. Food concentration dependent clearance rate, assimilation efficiency, growth and reproduction of Oikopleura dioica. Journal of Marine Systems. https://doi.org/10.1016/j.jmarsys.2009.01.004

Lombard, F., Sciandra, A., Gorsky, G., 2009b. Appendicularian ecophysiology. II. Modeling nutrition, metabolism, growth and reproduction of the appendicularian Oikopleura dioica. Journal of Marine Systems. https://doi.org/10.1016/j.jmarsys.2009.01.005

Lombard, F., Sciandra, A., Gorsky, G., 2005. Influence of body mass, food concentration, temperature and filtering activity on the oxygen uptake of the appendicularian Oikopleura dioica. Marine Ecology Progress Series. https://doi.org/10.3354/meps301149

Lombard, F., Selander, E., Kiørboe, T., 2011. Active prey rejection in the filter-feeding appendicularian *Oikopleura dioica*. Limnol. Oceanogr. 56, 1504–1512. https://doi.org/10.4319/lo.2011.56.4.1504

Lucas, C.H., Dawson, M.N., 2014. What Are Jellyfishes and Thaliaceans and Why Do They Bloom?, in: Pitt, K.A., Lucas, C.H. (Eds.), Jellyfish Blooms. Springer Netherlands, pp. 9–44. https://doi.org/10.1007/978-94-007-7015-7_2

Lucas, C.H., Jones, D.O.B., Hollyhead, C.J., Condon, R.H., Duarte, C.M., Graham, W.M., Robinson, K.L., Pitt, K.A., Schildhauer, M., Regetz, J., 2014. Gelatinous zooplankton biomass in the global oceans: geographic variation and environmental drivers. Global Ecology and Biogeography 23, 701–714. https://doi.org/10.1111/geb.12169

Luo, J.Y., Condon, R.H., Stock, C.A., Duarte, C.M., Lucas, C.H., Pitt, K.A., Cowen, R.K., 2020. Gelatinous zooplankton-mediated carbon flows in the global oceans: A data-driven modeling study. Global Biogeochemical Cycles. https://doi.org/10.1029/2020GB006704

Luo, J.Y., Grassian, B., Tang, D., Irisson, J.-O., Greer, A.T., Guigand, C.M., McClatchie, S., Cowen, R.K., 2014. Environmental drivers of the fine-scale distribution of a gelatinous zooplankton community across a mesoscale front. Marine Ecology Progress Series 510, 129–149. https://doi.org/10.3354./mesp10908

Lynam, C.P., Gibbons, M.J., Axelsen, B.E., Sparks, C.A.J., Coetzee, J., Heywood, B.G., Brierley, A.S., 2006. Jellyfish overtake fish in a heavily fished ecosystem. Current Biology 16, R492–R493. https://doi.org/10.1016/j.cub.2006.06.018

Madin, L.P., Deibel, D., 1998. Feeding and energetics of Thaliacea, in: Bone, Q. (Ed.), The Biology of Pelagic Tunicates. Oxford University Press, New York, USA, pp. 81–103.

Madin, L.P., Harbison, G.R., 1977. The associations of Amphipoda Hyperiidea with gelatinous zooplankton—I. Associations with Salpidae. Deep Sea Research 24, 449–463. https://doi.org/10.1016/0146-6291(77)90483-0

Madin, L.P., Kremer, P., Wiebe, P.H., Purcell, J.E., Horgan, E.H., Nemazie, D.A., 2006. Periodic swarms of the salp Salpa aspera in the Slope Water off the NE United States: Biovolume, vertical migration, grazing, and vertical flux. Deep-Sea Research Part I-Oceanographic Research Papers 53, 804–819. https://doi.org/10.1016/j.dsr.2005.12.018

Madin, L.P., Purcell, J.E., 1992. Feeding, metabolism, and growth of Cyclosalpa bakeri in the subarctic Pacific. Limnology and Oceanography 37, 1236–1251. https://doi.org/10.4319/lo.1992.37.6.1236

Martiny, A.C., Lomas, M.W., Fu, W., Boyd, P.W., Chen, Y.L., Cutter, G.A., Ellwood, M.J., Furuya, K., Hashihama, F., Kanda, J., Karl, D.M., Kodama, T., Li, Q.P., Ma, J., Moutin, T., Woodward, E.M.S., Moore, J.K., 2019. Biogeochemical controls of surface ocean phosphate. Sci. Adv. 5, eaax0341. https://doi.org/10.1126/sciadv.aax0341

Mayzaud, P., Boutoute, M., Gasparini, S., Mousseau, L., Lefevre, D., 2005. Respiration in marine zooplankton - The other side of the coin: CO 2 production. Limnology and Oceanography. https://doi.org/10.4319/lo.2005.50.1.0291

Mayzaud, P., Boutoute, M., Perissinotto, R., Nichols, P., 2007. Polar and Neutral Lipid Composition in the Pelagic Tunicate Pyrosoma atlanticum. Lipids 42, 647–657. https://doi.org/10.1007/s11745-007-3066-0

Mianzan, H., Pájaro, M., Colombo, G.A., Madirolas, A., 2001. Feeding on survival-food: gelatinous plankton as a source of food for anchovies. Hydrobiologia 451, 45–53.

Moriarty, R., O'Brien, T.D., 2013. Distribution of mesozooplankton biomass in the global ocean. Earth System Science Data 5, 45–55. https://doi.org/10.5194/essd-5-45-2013

Munk, W., Riley, G., 1952. Absorption of nutrients by aquatic plants. J. Mar. Res. 11, 215–240.

Muxagata, E., 1999. Avaliação da biomassa e distribuição zooplanctônica na plataforma continental sudeste brasileira durante o inverno de 1995 (Evaluation of the biomass and distribution of the zooplankton in the southeastern continental shelf of Brazil during the winter of 1995) (M.Sc Thesis). Departamento de Oceanografia - Universidade Federal de Rio Grande (FURG), Rio Grande, Brazil.

O'Brien, T.D., 2014. COPEPOD: The Global Plankton Database. An overview of the 2014 database contents, processing methods, and access interface. (NOAA Tech. Memo. No. NMFS-F/ST-37). U.S. Dept. of Commerce.

O'Brien, T.D., 2005. COPEPOD: A Global Plankton Database. A review of the 2005 database contents and creation of new global zooplankton biomass fields. (NOAA Tech. Memo. No. NMFS-F/SPO-73). U.S. Dept. of Commerce.

Paffenhöfer, G.-A., Köster, M., 2011. From one to many: on the life cycle of Dolioletta gegenbauri Uljanin (Tunicata, Thaliacea). Journal of Plankton Research 33, 1139–1145. https://doi.org/10.1093/plankt/fbr001

Pakhomov, E.A., 2004. Salp/krill interactions in the eastern Atlantic sector of the Southern Ocean. Deep Sea Research Part II: Topical Studies in Oceanography, The SWEDARP 1997/98 Expedition 51, 2645–2660. https://doi.org/10.1016/j.dsr2.2001.03.001

Pakhomov, E.A., Dubischar, C.D., Strass, V., Brichta, M., Bathmann, U.V., 2006. The tunicate Salpa thompsoni ecology in the Southern Ocean. I. Distribution, biomass, demography and feeding ecophysiology. Marine Biology 149, 609–623.

Perissinotto, R., Pakhomov, E.A., 1998. Contribution of salps to carbon flux of marginal ice zone of the Lazarev Sea, Southern Ocean. Marine Biology 131, 25–32.

Pershing, A.J., Stamieszkin, K., 2020. The North Atlantic Ecosystem, from Plankton to Whales. Annu. Rev. Mar. Sci. 12, 339–359. https://doi.org/10.1146/annurev-marine-010419-010752

Post, D.M., 2002. Using stable isotopes to estimate trophic position: Models, methods, and assumptions. Ecology 83, 703–718.

Purcell, J.E., Sturdevant, M.V., Galt, C.P., 2005. A review of appendicularians as prey of fish and invertebrate predators, in: Gorsky, G., Youngbluth, M.J., Diebel, D. (Eds.), Response of Marine Ecosystems to Global Change: Ecological Impact of Appendicularians. Contemporary Publishing International, Paris, France, pp. 359–435.

Ramaswamy, V., Sarin, M.M., Rengarajan, R., 2005. Enhanced export of carbon by salps during the northeast monsoon period in the northern Arabian Sea. Deep Sea Research Part II: Topical Studies in Oceanography 52, 1922–1929. http://dx.doi.org/10.1016/j.dsr2.2005.05.005

Remsen, A., Hopkins, T.L., Samson, S., 2004. What you see is not what you catch: a comparison of concurrently collected net, Optical Plankton Counter, and Shadowed Image Particle Profiling Evaluation Recorder data from the northeast Gulf of Mexico. Deep-Sea Research Part I: Oceanographic Research Papers 51, 129–151. https://doi.org/10.1016/j.dsr.2003.09.008

Robison, B.H., 2005. Giant Larvacean Houses: Rapid Carbon Transport to the Deep Sea Floor. Science 308, 1609–1611. https://doi.org/10.1126/science.1109104

Ruzicka, J., Brodeur, R.D., Cieciel, K., Decker, M.B., 2020. Examining the ecological role of jellyfish in the Eastern Bering Sea. ICES Journal of Marine Science 77, 791–802. https://doi.org/10.1093/icesjms/fsz244

Saba, V.S., Friedrichs, M.A.M., Antoine, D., Armstrong, R.A., Asanuma, I., Behrenfeld, M.J., Ciotti, A.M., Dowell, M., Hoepffner, N., Hyde, K.J.W., Ishizaka, J., Kameda, T., Marra, J., Mlin, F., Morel, A., O'Reilly, J., Scardi, M., Smith, W.O., Smyth, T.J., Tang, S., Uitz, J., Waters, K., Westberry, T.K., 2011. An evaluation of ocean color model estimates of marine primary productivity in coastal and pelagic regions across the globe. Biogeosciences. https://doi.org/10.5194/bg-8-489-2011

Seitzinger, S.P., Harrison, J.A., Dumont, E., Beusen, A.H.W., Bouwman, A.F., 2005. Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application: GLOBAL EXPORT OF C, N, AND P TO COASTAL SYSTEMS. Global Biogeochem. Cycles 19, n/a-n/a. https://doi.org/10.1029/2005GB002606

Smith Jr, K.L., Sherman, A.D., Huffard, C.L., McGill, P.R., Henthorn, R., Von Thun, S., Ruhl, H.A., Kahru, M., Ohman, M.D., 2014. Large salp bloom export from the upper ocean and benthic community response in the abyssal northeast Pacific: Day to week resolution. Limnol. Oceanogr 59, 745–757.

Stamieszkin, K., Steinberg, D.K., Maas, A.E., 2021. Fecal pellet production by mesozooplankton in the subarctic Northeast Pacific Ocean. Limnol Oceanogr 66, 2585–2597. https://doi.org/10.1002/lno.11774

Steinberg, D.K., Cope, J.S., Wilson, S.E., Kobari, T., 2008. A comparison of mesopelagic mesozooplankton community structure in the subtropical and subarctic North Pacific Ocean. Deep Sea Research Part II: Topical Studies in Oceanography 55, 1615–1635. https://doi.org/10.1016/j.dsr2.2008.04.025

Steinberg, D.K., Ruck, K.E., Gleiber, M.R., Garzio, L.M., Cope, J.S., Bernard, K.S., Stammerjohn, S.E., Schofield, O.M.E., Quetin, L.B., Ross, R.M., 2015. Long-term (1993–2013) changes in macrozooplankton off the Western Antarctic Peninsula. Deep Sea Research Part I: Oceanographic Research Papers 101, 54–70. http://dx.doi.org/10.1016/j.dsr.2015.02.009

Stock, C.A., Dunne, J.P., 2010. Controls on the ratio of mesozooplankton production to primary production in marine ecosystems. Deep-Sea Research Part I: Oceanographic Research Papers. https://doi.org/10.1016/j.dsr.2009.10.006

Stock, C.A., Dunne, J.P., Fan, S., Ginoux, P., John, J., Krasting, J.P., Laufkötter, C., Paulot, F., Zadeh, N., 2020. Ocean Biogeochemistry in GFDL's Earth System Model 4.1 and Its Response to Increasing Atmospheric CO2. J. Adv. Model. Earth Syst. 12. https://doi.org/10.1029/2019MS002043

Stock, C.A., Dunne, J.P., John, J.G., 2014a. Global-scale carbon and energy flows through the marine planktonic food web: An analysis with a coupled physical–biological model. Progress in Oceanography 120, 1–28. http://dx.doi.org/10.1016/j.pocean.2013.07.001

Stock, C.A., Dunne, J.P., John, J.G., 2014b. Drivers of trophic amplification of ocean productivity trends in a changing climate. Biogeosciences. https://doi.org/10.5194/bg-11-7125-2014

Stock, C.A., John, J.G., Rykaczewski, R.R., Asch, R.G., Cheung, W.W.L., Dunne, J.P., Friedland, K.D., Lam, V.W.Y., Sarmiento, J.L., Watson, R.A., 2017. Reconciling fisheries catch and ocean productivity. Proceedings of the National Academy of Sciences 201610238. https://doi.org/10.1073/pnas.1610238114

Stock, C.A., Powell, T.M., Levin, S.A., 2008. Bottom-up and top-down forcing in a simple size-structured plankton dynamics model. Journal of Marine Systems 74, 134–152. https://doi.org/10.1016/j.jmarsys.2007.12.004

Stukel, M.R., Décima, M., Selph, K.E., Gutiérrez-Rodríguez, A., 2021. Size-specific grazing and competitive interactions between large salps and protistan grazers. Limnol Oceanogr 66, 2521–2534. https://doi.org/10.1002/lno.11770

Sutherland, K.R., Madin, L.P., Stocker, R., 2010. Filtration of submicrometer particles by pelagic tunicates. Proceedings of the National Academy of Sciences 107, 15129–15134. https://doi.org/10.1073/pnas.1003599107

Sweetman, A.K., Smith, C.R., Dale, T., Jones, D.O.B., 2014. Rapid scavenging of jellyfish carcasses reveals the importance of gelatinous material to deep-sea food webs. Proceedings of the Royal Society B-Biological Sciences 281. https://doi.org/Artn 20142210 10.1098/Rspb.2014.2210

Takahashi, K., Ichikawa, T., Saito, H., Kakehi, S., Sugimoto, Y., Hidaka, K., Hamasaki, K., 2013. Sapphirinid copepods as predators of doliolids: Their role in doliolid mortality and sinking flux. Limnology and Oceanography 58, 1972–1984. https://doi.org/10.4319/lo.2013.58.6.1972

Titelman, J., Riemann, L., Sørnes, T.A., Nilsen, T., Griekspoor, P., B\aamstedt, U., 2006. Turnover of dead jellyfish: Stimulation and retardation of microbial activity. Marine Ecology Progress Series 325, 43–58.

Trueman, E., Bone, Q., Braconnot, J., 1984. Oxygen consumption in swimming salps (tunicata: thaliacea). Journal of Experimental Biology.

Turner, J.T., 2015. Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological pump. Progress in Oceanography 130, 205–248. https://doi.org/10.1016/j.pocean.2014.08.005

Urrère, M.A., Knauer, G.A., 1981. Zooplankton fecal pellet fluxes and vertical transport of particulate organic material in the pelagic environment. Journal of Plankton Research 3, 369–387.

Usbeck, R., Schlitzer, R., Fischer, G., Wefer, G., 2003. Particle fluxes in the ocean: comparison of sediment trap data with results from inverse modeling. Journal of Marine Systems 39, 167–183. https://doi.org/10.1016/S0924-7963(03)00029-0

Vargas, C.A., Madin, L.P., 2004. Zooplankton feeding ecology: clearance and ingestion rates of the salps Thalia democratica, Cyclosalpa affinis and Salpa cylindrica on naturally occurring particles in the Mid-Atlantic Bight. Journal of Plankton Research 26, 827–833. https://doi.org/10.1093/plankt/fbh068

Verity, P., Smetacek, V., 1996. Organism life cycles, predation, and the structure of marine pelagic ecosystems. Mar. Ecol. Prog. Ser. 130, 277–293. https://doi.org/10.3354/meps130277

Waite, A., Bienfang, P.K., Harrison, P.J., 1992. Spring bloom sedimentation in a subarctic ecosystem. Marine Biology 114, 131–138. https://doi.org/10.1007/BF00350862

Walters, T.L., Lamboley, L.M., López-Figueroa, N.B., Rodríguez-Santiago, Á.E., Gibson, D.M., Frischer, M.E., 2019. Diet and trophic interactions of a circumglobally significant gelatinous marine zooplankter, *Dolioletta gegenbauri* (Uljanin, 1884). Mol Ecol 28, 176–189. https://doi.org/10.1111/mec.14926

Ward, B.A., Dutkiewicz, S., Jahn, O., Follows, M.J., 2012. A size-structured food-web model for the global ocean. Limnology and Oceanography. https://doi.org/10.4319/lo.2012.57.6.1877

Wiebe, P.H., Benfield, M.C., 2003. From the Hensen net toward four-dimensional biological oceanography. Progress in Oceanography 56, 7–136. https://doi.org/Doi 10.1016/S0079-6611(02)00140-4

Wilson, S.E., Ruhl, H.A., Smith Jr, K.L., 2013. Zooplankton fecal pellet flux in the abyssal northeast Pacific: A 15 year time-series study. Limnology and Oceanography 58, 881–892. https://doi.org/10.4319/lo.2013.58.3.0881

Wright, R.M., Le Quéré, C., Buitenhuis, E., Pitois, S., Gibbons, M.J., 2021. Role of jellyfish in the plankton ecosystem revealed using a global ocean biogeochemical model. Biogeosciences 18, 1291–1320. https://doi.org/10.5194/bg-18-1291-2021

Zhao, M., Golaz, J.-C., Held, I.M., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L.J., Dunne, J.P., Dunne, K., Durachta, J., Fan, S.-M., Freidenreich, S.M., Garner, S.T., Ginoux, P., Harris, L.M., Horowitz, L.W., Krasting, J.P., Langenhorst, A.R., Liang, Z., Lin, P., Lin, S.-J., Malyshev, S.L., Mason, E., Milly, P.C.D., Ming, Y., Naik, V., Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C.J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L.G., Wilson, J.R., Winton, M., Wittenberg, A.T., Wyman, B., Xiang, B., 2018. The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 2. Model Description, Sensitivity Studies, and Tuning Strategies. J. Adv. Model. Earth Syst. 10, 735–769. https://doi.org/10.1002/2017MS001209

Zweng, M.M., Reagan, J.R., Antonov, J.I., Locarnini, R.A., Mishonov, A.V., Boyer, T.P., Garcia, H.E., Baranova, O.K., Johnson, D.R., Siedov, D., Biddle, M.M., 2013. World Ocean Atlas 2013, Volume 2: Salinity.

Supplemental Tables and Figures

Figure S1. Major ocean biomes for the COBALTv2 control (a) and GZ-COBALT (b). The chlorophyll thresholds delineating the low chlorophyll (LC) regions were 0.162 and 0.184 mg Chl m^{-3} for the control and GZ-COBALT, respectively. The annual minimum of the mixed layer depth irradiance climatology (< 5 W m^{-2}) delineated the high chlorophyll seasonally stratified (HCSS) biome from the high chlorophyll permanently stratified (HCPS) biome.

Figure S2. Biomass chlorophyll scaling for small tunicates (left column), large tunicates (center column) and crustacean mesozooplankton (right column) for the five sensitivity cases: case 1 (a-c), where large

tunicate basal respiration is increased; case 2 (d-f), where small tunicate maximum ingestion rate is increased; case 3 (g-i), where the tunicate ingestion half-saturation constant is the same as crustaceans, and maximum ingestion is adjusted accordingly; case 4 (j-l), where tunicate assimilation efficiency is set to be a constant; and case 5 (m-o), where large tunicate aggregation mortality is turned off. See Table 2, and the caption on Fig. 6 for further details.

Figure S3. Zooplankton seasonal cycle differences between the GZ-COBALT base case (dashed) and three of the sensitivity cases: case 3 (a-e); case 4 (f-j); case 5 (k-o). See caption on Fig. 7 for further details.

Figure S4. Differences in annual mean productivity ratios at the top 100 m between GZ-COBALT and the COBALTv2 control, showing ratios of small mesozooplankton (= medium zooplankton) production to NPP (a-c) and large mesozooplankton (= large zooplankton) production to NPP (d-f). The plots show the GZ-COBALT simulation (left column; a, d), the COBALTv2 control (center column; b, e), and the difference between the two (right column; c, f). Colorbar is set to be the same as in Fig. 8 to allow for a direct comparison.

Figure S5. Biomass chlorophyll scaling for small (a) and large (b) tunicates, using daily output from the GZ-COBALT model, showing a randomly selected day within the growing season from the last year of the model simulation (2007). See also the caption in Fig. 6 for more details.

Figure S6. Fraction of late summertime detritus production in the top 100 m from (a) all tunicates, and (b) large tunicates only. Late summer is defined as July-September in the Northern Hemisphere, and January-March in the Southern Hemisphere.

Figure S7. Mean daily Production/Biomass (P/B, d^{-1}) ratios for small zooplankton (a), medium zooplankton (b), large zooplankton (c), small tunicates (d), and large tunicates (e). Months in the Southern Hemisphere are shifted such that Austral summer occurs during months 6-8, and Austral winter occurs during months 12, 1, and 2.